Title: Caloric-restriction mitigates age-associated alteration in CG and non-CG methylation in the old

brain.

Abbreviated title: CR mitigates DNA methylation changes in the brain

Authors: Niran Hadad[1,2,6], Archana Unnikrishnan[2,5,6], Jordan A. Jackson[2,5,6], Dustin R. Masser[2,3], Laura

Otalora[2,3,5,6], David R. Stanford[2,3,6], Arlan Richardson[2,4,5,6], Willard M. Freeman[1,2,3,5,6*]

Affiliations: [1]Oklahoma Center for Neuroscience, [2]Reynolds Oklahoma Center on Aging, [3]Department of

Physiology, [4]Oklahoma City VA Medical Center, [5]Department of Geriatric Medicine, [6]Oklahoma Nathan

Shock Center for Aging, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA, 73104

*Corresponding Author: Willard M. Freeman, Ph.D.; Mail: SLY-BRC 1370, 975 NE 10[th] St, Oklahoma City,

OK 73104; Telephone: (405) 271-8000 ext. 30729; Email: wfreeman@ouhsc.edu

Conflict of interest: The authors declare no competing financial interests

Acknowledgments: The authors would like to acknowledge Laura Blanco-Berdugo for computational

advice and the University of Oklahoma Supercomputing Center for Education & Research (OSCER) for

allocating computational resources used for data analysis. This work was supported by the Donald W.

Reynolds Foundation, the Oklahoma Center for Neuroscience translational seed grant, the Oklahoma

Nathan Shock Center of Excellence in the Biology of Aging Targeted DNA Methylation and Mitochondrial

Heteroplasmy Core (P30AG050911), the National Institute on Aging (R01AG045693, F31AG038285,

T32AG052363), and National Eye Institute (R21EY024520, R01EY021716).

Abstract

Brain aging is marked by cognitive decline and increased susceptibility to neurodegeneration. Epigenetic

mechanisms, including DNA methylation, are vital to CNS cellular function and memory formation, and

are dysregulated with aging and age-related neurodegenerative disease. Caloric-restriction (CR), an

established pro-longevity intervention, increases neurogenesis, improves memory function, and

protects from age-associated pathologies. However, the molecular mechanisms promoting the

neuroprotective effect of CR remain largely unknown. We tested the role of DNA methylation as a

mechanism for CR-induced neuroprotection in the old brain. Hippocampal DNA from young (3M) and

old (24M) male mice fed *ad libitum* and 24M old mice fed 40% calorie-restricted diet from 3M of age

were examined by genome-wide bisulfite sequencing to measure methylation levels at base-specific

resolution. Over 25 million CG and CH (non-CG) sites were examined. Of the ~40,000 differentially

methylated CGs (dmCGs) and ~80,000 CHs (dmCHs) observed with aging, 35% and 38%, respectively,

were prevented by CR. Unique to dmCHs, CR preferentially prevented age-related hypermethylation. A

diet-specific methylation response was observed in both CG and CH contexts. CR-induced dmCHs were

enriched in CG-rich regions such as promoters and CG islands including hypermethylation of DNMT1 and

TET3 promoters corresponding to reduced gene expression. These findings demonstrate for the first

time that caloric-restriction prevents age-induced cytosine methylation changes in the old brain in

combination with CR-specific methylation changes that may function to maintain epigenetic

homeostasis through epigenetic auto-regulation. The prevention of age-dmCGs/CHs by CR emphasizes

the prominent role of DNA methylation as a driver of the aging process.

Significance Statement

DNA methylation is a central epigenetic regulator of genomic structure and function. Aberrant control of

DNA methylation is associated with brain aging and contributes to cognitive decline and age-related

neurodegenerative disease. Unexplored though is whether anti-aging treatments can maintain

'youthful' methylation patterns in the brain. Our findings demonstrate that caloric-restriction attenuates

age-related changes in DNA methylation and induces changes in DNA methylation at specific sites

independent of aging in both neuroprotective pathways and pathways shared by different tissues

responsive to calorie-restriction, potentially through a combination of epigenetic auto-regulation and

uniquely targeted changes in methylation. These findings contribute to our understanding of the aging

process and the molecular mechanisms underlying anti-aging treatments.

Introduction

With increased life-expectancy, incidence of age-related neurodegenerative diseases such as

Alzheimer's, Parkinson's, and other dementias, has increased and is expected to double by 2040 (Reitz

et al., 2011; Reeve et al., 2014; Hickman et al., 2016). Anti-aging interventions offer promise in delaying

age-related impairments and neurological disease development. The most established anti-aging

intervention is calorie-restriction (CR), where caloric intake is reduced by 10-40%. CR consistently

increases lifespan and improves health span across model organisms including invertebrates (Lee et al.,

2006), rodents (Turturro et al., 1999), and primates (Colman et al., 2014). Several conserved

mechanisms have been proposed to underline the beneficial effects observed following CR including

inhibition of mTOR signaling (Kaeberlein et al., 2005), decreased insulin/IGF-1 signaling (Sonntag et al.,

1999) activation of Sirt1/Sirt2 (Bordone and Guarente, 2005) and activation of redox signaling (Hyun et

al., 2006), however CR is also proposed to act through cell-autonomous and tissue specific mechanisms

(Dacks et al., 2013; Schafer et al., 2015; Barger et al., 2017).

Short- and long-term CR have a protective effect against brain aging. CR increases expression of DNA

repair and anti-stress proteins (Kisby et al., 2010), improves glucose metabolism efficiency (Willette et

al., 2012b) and delays onset of inflammatory cytokines (Swindell, 2009). Specifically to the brain, CR

delays neurodegeneration and synaptic dysfunction (Dietrich et al., 2010; Schafer et al., 2015), improves

neuroendocrine function (Redman and Ravussin, 2009; Willette et al., 2012a) and promotes induction of

genes active in neuroprotection, neural growth (Park et al., 2013) and synaptic and neuronal function

(Adams et al., 2008; Fontan-Lozano et al., 2008; Schafer et al., 2015; Wood et al., 2015). At the

functional level, CR improves learning and memory and is beneficial in reducing anxiety and depression

in both rodents and humans (Inoue et al., 2004; Lutter et al., 2008; Willette et al., 2012a). CR is also

beneficial in alleviating pathology of age-related neurodegenerative diseases, improves cognitive

function (Halagappa et al., 2007) and decreases hippocampal beta-amyloid accumulation in Alzheimer's

models (Wang et al., 2005). Despite these positive effects on brain aging and age-related disease, the

molecular mechanisms underlying brain aging processes and those responsive to CR remain elusive.

Epigenetic modifications in the form of DNA methylation and histone modifications are proposed as

central regulators of the aging process (Lopez-Otin et al., 2013). DNA methylation is a malleable

epigenetic modification in which a methyl group is added to the 5-position of a cytosine ring and plays a

complex role in genomic regulation. Proper control of DNA methylation is instrumental to neural

differentiation, neural growth, synaptic function and intact cognitive function (Feng et al., 2010; Lister et

al., 2013; Grigorenko et al., 2016), key processes hampered by the aging process. Altered methylation at

specific genomic locations is associated with the normal aging process (Xekardaki et al., 2015; Wyss-

Coray, 2016). Recent studies have shown a protective epigenetic effect of anti-aging interventions in the

liver and kidney (Kim et al., 2016; Cole et al., 2017; Hahn et al., 2017). However, whether the response

by DNA methylation to aging is common between tissues or demonstrates tissue specific patterns is

unclear.

We previously described extensive age-related DNA methylation changes in the old brain (Masser et al.,

2017). In the current study we investigated whether these changes are prevented by CR by focusing on

the hippocampus, a tissue demonstrating molecular, cellular and functional impairment with old age

(Vanguilder and Freeman, 2011; Gray and Barnes, 2015; Seib and Martin-Villalba, 2015). Using a

genome-wide sequencing approach we examined the response of age-related CG and for the first time

CH (non-CG) methylation to CR. Additionally, we characterize unique CR-specific changes in both CG and

CH methylation, highlighting potential functional similarities and dissimilarities between CG and CH

methylation. Lastly, we show that enzymes regulating DNA methylation and demethylation are

suppressed by CR but not aging, in part through alteration in promoter methylation induced by CR,

indicating a potential mechanism for CR modulation of the neuroepigenome.

Materials and methods

Animals

Male C57BL/6 mice were obtained from the NIA aging and caloric restriction colony at 3 month and 24

month of age. Mice were housed at the University of Oklahoma Health Sciences Center animal facility

and maintained under SPF conditions in a HEPA barrier environment. Mice were fed irradiated NIH-31

mouse/rat diet (Teklad, Envigo). Both CR and AL mice were individually housed. CR was initiated at 14

weeks of age at 10% restriction, increased to 25% restriction at 15 weeks, and to 40% restriction at 16

weeks of age. Young (3 month) and old (24 month) mice were euthanized and both right and left

hippocampus were harvested, snap frozen in liquid nitrogen, and stored at -80°C. All animal experiments

were performed according to protocols approved by the OUHSC Institutional Animal Care and Use

Committee. Tissues from young and old *ad libitum* animals were used from a previous study (Masser et

al., 2017).

Library preparation and sequencing

Hippocampal gDNA was isolated by spin columns (Zymo duet) and genome-wide bisulfite capture

sequencing in accordance to previously described and validated methods (Hing et al., 2015; Masser et

al., 2016). gDNA was quantified by qubit (Invitrogen) and sheared by sonication (Covaris e220) to an

average 200bp fragment size. Fragment size was confirmed by capillary electrophoresis (DNA1000,

Agilent). Bisulfite sequencing library preparation was performed with 3µg DNA using SureSelect Methyl-

seq XT per manufacturer instructions (Agilent). In brief, sheared DNA was end-repaired, adenylated and

adapter ligated prior to bisulfite conversion. Libraries were sized and quantified by capillary

electrophoresis to ensure recovery of at least 350 ng of library. Libraries were hybridized to mouse

SureSelect Methyl-Seq capture library. Hybridized libraries were captured with Dynabeads MyOne

Streptavidin T1 magnetic beads and then bisulfite converted (EZ DNA Methylation-Gold, Zymo

Research), amplified according to manufacturer recommendations and purified with AMPure XP beads.

Following bisulfite conversion libraries were indexed and confirmed for fragment size by capillary chip

(DNA high sensitivity, Agilent). Prior to sequencing, libraries were quantified by PCR (KAPA library

quantification kit, Kapa Biosystems), diluted to 4nM, and pooled in equimolar concentrations. Pooled

libraries were diluted to 12 pM and then sequenced at 100 bp paired end reads on the Illumina

HiSeq2500 platform. Raw fastq files are available through the Sequencing Read Archive (bioproject ID

PRJNA397575).

Informatics

Paired-end reads were trimmed using CLC Genomics Workbench v10.1.1 (RRID:SCR_011853). Reads with

a Q-score <30 or with >1 ambiguous nucleotide were filtered. Reads passing filter were adapter trimmed

and an additional 3 bp were removed from the 3' and 5' ends of each read. All computational analyses

were performed in UNIX and R using custom scripts or published tools (where specified). Trimmed PE

reads were aligned as PE to the mouse GRCm38/mm10 assembly using Bismark Bisulfite Mapper version

0.14.4 (RRID:SCR_005604) (Krueger and Andrews, 2011). Following read mapping, methylation calling

was performed with Bismark methylation extractor and % methylation was determined by calculating

(total C methylated/total C covered) per cytosine in the genome. Only cytosines covered in all groups

with a cumulative coverage of 35 were used for analysis, exceeding the coverage recommendations in

the field for BS-seq studies (Roadmap Epigenomics et al., 2015; Ziller et al., 2015). Greater than 25M CG

and non-CG sites were carried forward for downstream analysis. Differential methylation between

groups was determined using Kruskal-Wallis H test in R, significant sites were tested for pairwise

multiple comparisons using Conover post hoc test using 'lsmeans', p-values were adjusted for false

discovery using the Benjamini-Hochberg procedure. Age-changes were defined as all differences found

between young-AL and old-AL animals. Diet-changes were defined as all differentially methylated

cytosines between Old-AL and Old-CR animals.

Annotation and enrichment

Coordinates for genic features including exons, introns and promoters (defined as ±1kb from TSS) were

downloaded from UCSC genome browser for GRCm38/mm10 reference genome. CG island shores were

defined as 2kb upstream and downstream from the CG island borders and shelves were defined as 2kb

upstream and downstream from shores. Coordinates for annotated gene regulatory regions for mouse

brain were downloaded from ENSEMBL open database (Zerbino et al., 2015). Over- and under-

representation of differentially methylated cytosines was determined by overlapping all analyzed sites

with the genic feature list and gene regulatory regions using 'bedtools' (Quinlan and Hall, 2010).

Statistical significance of over- and under- representation was determined using hypergeometric test.

For pathway analysis, genes were annotated for differentially methylated cytosines, only genes

containing >2 differentially methylated sites were included in the analysis. Analysis of pathways

enriched for genes affected for differential methylation was performed with the 'ReactomePA' R

package. Pathways with broad (>100 genes) or narrow (<7 genes) definitions were excluded.

Additionally, pathways with similar gene lists were consolidated. A full unedited list will be available

upon request.

Real-Time PCR (qPCR)

Expression of DNMTs (pan-DNMT1, Mm01151063_m1; E1-DNMT1, Mm00599763_m1; Alt. E1-DNMT1,

Mm01174085_m1; DNMT3a1, Mm00432870_m1) and TETs (TET1, Mm01169087_m1; TET2,

Mm00524395_m1; TET3, Mm00805756_m1) was determined by real-time qPCR as described previously

(Mangold et al. 2017b). β-actin (Mm02619580_g1) was used as the endogenous control and each

sample was analyzed in technical triplicates.

Experimental Design and Statistical Analysis

Genome-wide methylation analysis was performed with n=4/group. Gene expression experiments were

performed with n=7-8/group. All statistical analyses were performed in R. Box plots represent the 25^{th}

and 75^{th} percentiles. Differential methylation and enrichments were determined as detailed above.

Global methylation differences were compared using linear regression and are written as mean ± SD.

Analysis of gene expression using RT-qPCR was compared using One-Way ANOVA with pairwise

comparisons and Benjamini-Hochberg multiple testing corrections.

Results

Age related CG methylation is attenuated by life-long caloric restriction

To examine the effect of life-long caloric-restriction on CG methylation in the old brain, hippocampal

DNA was isolated from Young (3 month old, Y-AL, n=4) and old (24 month old, O-AL, n=4) male mice that

were fed *ad libitum* (AL) throughout life and old animals fed a 40% caloric restriction (CR) diet (24 month

old, O-CR, n=4) starting at 3 months of age. Using genome-wide bisulfite sequencing, base-specific

methylation levels across 3 million CpGs were compared between the 3 groups. Global DNA

methylation, i.e. the methylation averages across all analyzed CpG sites did not change with age or diet

(Y-AL: 42.3% ± 0.7; O-AL: 42.6% ± 0.5; O-CR: 42.6%±0.6; p=0.45 by linear regression). We defined age-

associated dmCGs (age-dmCGs), as any methylation difference between young-AL and old-AL animals

that met statistical criteria (methylation difference >|5%| and adjusted p-value < 0.05, for more details

see differential methylation in Methods). Comparing young-AL animals to old-AL animals, 41,586 age-

related age-dmCGs were identified. Examination of the methylation profile of all age-dmCGs for

differences between young-AL and old-CR mice reveals a pattern of regression towards that of young-

AL, indicating a "younger" methylation profile in age matched animals fed CR diet (Fig. 1A, B). At single

base resolution, life-long CR prevented 32% hypermethylated and 36% hypomethylated age-dmCGs (Fig.

1C). The distribution of age-dmCGs shows a higher number of hypermethylated dmCGs than

hypomethylated dmCGs. Prevented age-dmCGs followed a similar distribution that was proportional to

that observed for age-dmCGs. (Fig. 1D). Principal component analysis of age-dmCGs unaffected by CR

demonstrated clustering of all old animals as compared to young-AL (Fig. 1E). Parallel analysis of

prevented age-dmCGs separates young-AL and old-CR from old-AL on the first component, while young-

AL and old-CR separate on the second component (Fig. 1F). The partition of young-AL and old-CR may be

attributed to a subset of 416 loci that demonstrate differential methylation between Y-AL and O-CR with

both diet and aging, e.g. gain in methylation with age and loss in methylation with age and diet as

compared to Y-AL.

The localization of differentially methylated sites by genomic context and their interaction with other

epigenomic marks can aid in understanding the role of DNA methylation changes in genomic regulation.

Methylation changes were mapped with known coordinates of gene-centric features to annotate the

enrichment of differentially methylated sites. In genic features, including CG islands, shores and shelves,

introns, exons and promoters there was no observable difference in enrichment pattern between hyper-

and hypo- methylated age-dmCGs or prevented age-dmCGs (Fig. 1G). Age-dmCGs and prevented age-

dmCGs were enriched in introns and CGI shores while under-represented in islands, CGI shelves, exons,

and promoters. Depletion of age-dmCGs in promoters was more pronounced in promoters containing

CGI. Previous reports demonstrated that DNA methylation may alter gene expression through

interaction with gene regulatory regions (Roadmap Epigenomics et al., 2015). To identify enrichment at

specific gene regulatory regions we used the ENSEMBL brain specific regulatory build (Zerbino et al.,

2015) which was constructed using sequencing methods to identify CTCF binding sites, enhancers, open

chromatin regions, promoters, promoter flanks, and transcription factor binding sites and their specific

activation states (active, inactive, poised, repressed). Enrichment of age-dmCGs was predominantly

observed in inactive state gene regulatory regions. Inactive regions are defined as regions not associated

with other epigenetic marks, indicating that the effect of differential methylation at these regions is

independent of interaction with other known epigenetic marks. Age-dmCGs were enriched in active

state regulatory regions and occurred in CTCF binding sites (hypo only), enhancers (hypo and hyper),

open chromatin regions (hyper only), and promoter flank (hyper only). Prevented age-dmCG generally

demonstrated a similar enrichment pattern (Fig. 1H).

The interaction between gene expression and DNA methylation is still an active area of exploration. The

canonical view of DNA methylation regulation of gene expression through promoter or gene body

methylation has been challenged recently both in neurons (Sharma et al., 2016) and following CR (Hahn

et al., 2017). Therefore, we asked which pathways are affected by DNA methylation without

assumptions as to whether methylation is positively or negatively regulating a specific gene. Only genes

containing more than 2 dmCGs in their gene body (inclusive of promoter, introns and exons) were

included in this analysis. Genes were separated by those affected by hyper- and hypo- methylation.

Hypermethylated age-dmCGs were found in 3,266 genes and hypomethylated age-dmCGs were found in

1,925. Of those genes 948 were found to contain prevented hypomethylated age-dmCGs and 548

contained prevented hypermethylated age-dmCGs. Pathways involved in neural cell communication,

neural cell growth, cellular integrity, cell metabolism and inflammatory pathways were affected by both

hyper- and hypomethylation with aging (Fig. 1I). The Rho GTPase pathway, which is involved in neurite

growth and axon guidance (Hall and Lalli, 2010) and downstream of cognitive decline-related myelin

associated inhibitors of synaptic plasticity (Vanguilder et al., 2012) was enriched in prevented

hypomethylated age-dmCGs while no pathways were enriched amongst the prevented hypermethylated

age-dmCGs. These findings are consistent with the negative effect of aging on neuronal integrity and

positive cellular effect of CR on neuronal survival and growth (Seib and Martin-Villalba, 2015; Hornsby et

al., 2016).

Age related non-CG methylation is attenuated by life-long caloric restriction

The effect of calorie restriction on CH methylation has not been previously examined. CH methylation is

higher in the brain relative to other tissues (Lister et al., 2013), is differentially regulated as compared to

CG methylation in the brain (He and Ecker, 2015), and has unique functional impacts on gene regulation

(Lister and Mukamel, 2015). We analyzed >22M CHs sites and asked whether age-dmCHs respond

similarly to diet as age-dmCGs. Differential methylation of CHs was determined in a base-by-base

fashion, similar to CGs. The absolute methylation level of most dmCHs was generally lower than that of

CGs. Average global CH methylation levels did not change with age or with diet (Y-AL: 1.59% ± 0.07; O-

AL: 1.64% ±0.05; O-CR: 1.58%±0.8; $F_{(1,10)}$ = 0.61, p=0.83 by linear regression). More dmCHs (79,058 age-

dmCHs) were identified to change with age compared to CGs. The methylation profile of age-dmCHs in

old animals fed CR diet was closer to that of young-AL animals compared to that of old animals fed *ad*

libitum (Fig. 2A). The number of hypermethylated age-dmCHs was higher than hypomethylated dmCHs

in a manner similar to CGs. Life-long CR prevented 59% of all hypermethylated age-dmCHs and 11% of

hypomethylated age-dmCHs. Prevention of hypermethylated age-dmCHs by CR was 10 times more likely

than diet induced preventions of hypomethylated age-dmCHs (Fig. 2B-C). Principal component analysis

of age-dmCHs unaffected by diet shows a distinct clustering of both AL and CR old animals compared to

young-AL animals (Fig. 2D). Clustering of samples by the methylation profile of prevented age-dmCHs

shows that old-AL animals separate from young-AL and old-CR animals (Fig. 2E).

Differentially methylated CHs were generally under-represented in CG islands and shelves, and in gene-

centric elements. The exception being an enrichment of prevented hypomethylated age-dmCH in CGI

shores (Fig. 2F). Taken together, under-representation of dmCHs in in CG islands and genic regions

indicate that changes in methylation with age, and those prevented by CR, occur primarily in intragenic

regions and are partially excluded from CG rich regions. Enrichment analysis of age-dmCHs in gene-

regulatory regions reveals that age-dmCHs and prevented age-dmCHs, regardless of gain or loss in

methylation, are enriched mostly in active and inactive enhancers. Hypomethylated age-dmCHs and

prevented age-dmCHs were also enriched in poised TF binding sites and repressed promoters (Fig. 2G).

Pathways enriched for genes affected by age-dmCH were functionally similar to those enriched by age-

dmCG, including pathways involved in metabolic regulation, cellular signaling, and regulation of neural

cell structure and growth. Pathways enriched for prevented age-dmCHs were primarily involved in

energetics, metabolism and neurite growth (Fig. 2H). Of all the pathways found to be affected by

differential methylation 18 were common to age-dmCHs and age-dmCGs (Fig. 2I).

Gene regulatory regions are affected differentially by CR induced CG methylation

Caloric-restriction was sufficient to prevent >1/3 of all age-dmCs (CGs and CHs). CR also induced DNA

methylation changes independent of aging as evident when comparing old-AL and old-CR animals.

Methylation of these CR-induced dmCGs (diet-dmCGs) is not different between young-AL and old-AL

animals (colored green in Fig. 3A-left scatterplot) but are differentially methylated between old-AL and

old-CR animals (Fig. 3A-right scatterplot). Of all differentially methylated dmCGs between old-AL and

old-CR, 69.4% were induced by CR and were independent of the aging process (Fig. 3B). A single base

example of a diet-induced change in methylation is depicted in Fig. 3C. The distribution of all diet-

dmCGs shows significant preference towards hypermethylation (Fig. 3D), though the average magnitude

of methylation change is similar between hyper- and hypo- methylation (12.3% vs -12.5%, Fig 3D).

The genomic localization of diet-dmCGs is different from age-dmCGs. Hypermethylated diet-dmCGs

were over-represented in CG islands and shores and under-represented in CG shelves. Hypomethylated

diet-dmCGs were only enriched in shores. With respect to genic regions diet-dmCGs were primarily

depleted with the exception of a small but significant over representation of hypermethylated sites in

exons and hypomethylated sites in introns (Fig.3E). The same distinction between hyper- and hypo- diet-

dmCGs was observed in gene regulatory regions. Enrichment of diet-dmCGs was primarily in poised

regulatory regions while enrichment of hypomethylated diet-dmCGs was observed in active regulatory

features, including CTCF binding sites, enhancers, open chromatin and promoter flanks. Active

promoters were not enriched for hyper- and hypo- methylated diet-dmCGs; however, hypermethylated

diet-dmCGs were enriched in inactive and repressed promoters (Fig. 3F). Genes affected by diet-dmCGs

were enriched for pathways involving energy regulation, inflammatory pathways and phagocytosis (Fig.

3G). Interestingly, we found that genes containing diet-dmCGs were enriched in DNA methylation and

senescence-associated secretory phenotype pathways; both pathways have been proposed as potential

regulators of the aging process (Lopez-Otin, 2013).

Non-CG diet induced changes are clustered

The diet-specific effect on methylation was also observed in non-CG sites (Fig. 4A). Changes in CH

methylation with diet occurred generally at lower methylation levels with the majority of changes

occurring under 50% (Fig. 4A – right panel), but showed no difference between young-AL and old-CR

(Fig. 4A – left panel). Most diet-dmCHs (75% of all differential methylation observed in CHs) were

independent of the aging process (Fig. 4B). Twice as many hypermethylated (64,191 sites) than

hypomethylated diet-dmCHs (26,237 sites) were observed, similar to diet-dmCGs. Hypermethylation of

diet-dmCHs was higher than hypomethylated diet-dmCHs (10% vs -8%, Fig. 4C).

CG dinucleotides are generally depleted from large regions of the genome but occur at specific, high

density regions such as in CG islands and promoters. On the other hand, the distribution of CHs is

relatively uniform across the genome (He and Ecker, 2015). We investigated whether age and diet

induced dmCHs are sporadically distributed across the genome or clustered as a result of targeted

changes at specific genomic loci. We computed the distance between dmCHs, 62% of all dmCHs (age

and diet) occurred within 10kb of one another; therefore we focused our analysis to 10kb segments. The

proximity of diet-dmCHs to the nearest dmCH was higher than that observed for age-dmCHs (67% diet-

dmCHs compared to 56% age-dmCHs, Fig. 4D). This difference was even more pronounced when

calculating the distance between dmCHs within 1kb (49% vs. 29%). Further examination of the clustering

pattern of dmCHs in 10kb and 1 kb segments revealed that the distance to the next proximal dmCH (2nd

dmCH, 3rd dmCH, etc.) decreased gradually for both age and diet, but more diet-dmCHs clustered

together between the 1st and the 6th diet-dmCHs than the 1st and 3nd age-dmCHs in 10kb segments

(Figure 4E). This clustering of diet-dmCHs suggests differential regulation/targeting mechanisms of CH

methylation in response to diet and between diet and age.

Diet dmCHs are over-represented in CGI-promoters and depleted from non-CGIs promoters.

Hypermethylation of diet-dmCHs was over-represented in CG islands, shores and exons, while

hypomethylation was over-represented in introns, a localization pattern similar to diet-dmCGs.

Interestingly, hypermethylated diet-dmCHs were also over-represented in promoters containing CGIs

but were absent from non-CGI promoters (Fig. 5A). This finding is unexpected since CH methylation is

often precluded from high-CG-rich regions and promoters (Lister et al., 2009; Laurent et al., 2010; Xie et

al., 2012; Guo et al., 2014), further supporting a regulated control of methylation by diet. Enrichment of

diet-dmCHs in gene regulatory regions confirms the observation of over-representation of

hypermethylated diet-dmCHs in promoters. Using the ENSEMBL regulatory build we further divided

promoters by their activation state and found that hypermethylated diet-dmCHs are enriched in active

and poised promoters and promoter flanks, and are under-represented in repressed and inactive

promoters. Enrichment of hypomethylated diet-dmCHs was mainly observed in inactive gene regulatory

elements (Fig. 5B).

Higher numbers of genes contained diet-induced CH hypermethylation (5,340 genes) than

hypomethylation (2,981 genes), consistent with the distribution of diet-dmCHs (Fig. 4C); however genes

with hypomethylated diet-dmCHs were enriched in a greater number of pathways (40 pathways by

hypomethylation vs 21 pathways by hypermethylation). Top enrichment pathways included those

regulating cell structure, neural cell communication and growth, and energy metabolism. Inflammatory

pathways were mostly enriched for genes containing hypomethylated dmCHs (Fig. 5C). Of the 69

pathways affected by both dmCHs and dmCGs we found 12 common pathways regulating inflammation

and neuronal integrity (Fig. 5D).

Caloric restriction changes in methylation auto-regulates the DNA methylation machinery.

DNA methylation regulatory enzymes (DNA methyltransferases; DNMTs and Ten-eleven Translocation;

TETs) mRNA and protein expression do not change in the hippocampus with aging (Hadad et al., 2016).

Given that diet-dmCGs were enriched in the DNA methylation pathway, we examined whether any of

the CR-induced changes in methylation observed in the current study mapped to DNMTs and TETs genic

elements. We identified 6 hypermethylated diet-dmCs (2 dmCGs and 4 dmCHs) in the promoter region

of TET3 (Fig. 6A), corresponding to a decrease in TET3 mRNA expression following CR (Fig. 6B). We also

found 9 hypermethylated diet-dmCs (6 dmCHs and 3 dmCGs) in the promoter region of DNMT1 (Fig. 6C),

but an examination of DNMT1 with a pan-isoform assay inclusive of all known isoforms revealed no

changes in expression (Fig. 6D). Promoter methylation of DNMT1 is located near two known start sites;

therefore we examined the expression of each isoform separately by qPCR. Expression of the exon-1

containing mRNAs declined significantly with diet, and minimally with aging, while the alternate exon-1

isoform was unchanged (Fig. 6E).

No other dmCs induced by CR found in our analysis were observed in the remaining DNA methylation

regulatory enzymes, however examination of the expression levels of these enzymes show a small but

significant decrease with age and diet of DNMT3a1 and a decrease in TET2 expression with diet (Fig. 3F).

TET1 expression did not differ with age or diet as well. These findings suggest that while DNMTs and

TETs are generally not regulated with age, the epigenetic effects of CR may occur through regulation of

DNMTs and TETs.

Discussion

This study examined whether DNA methylation is a neuroprotective mechanism of action of CR in

retarding brain aging. Given that calorie-restriction improves cognition and cellular function and that

altered methylation may underline age-related dysfunction in these processes, hippocampal DNA

methylation was quantified at base-specific resolution to identify age-associated changes in DNA

methylation and the effect of CR in preventing these changes, as previously reported for CG methylation

in the kidney and liver (Kim et al., 2016; Cole et al., 2017; Hahn et al., 2017). Our findings demonstrate

for the first time that base-specific prevention of age-related differential methylation by CR is observed

in the brain, not only in CG context but also CH. This is the first study to evaluate the effect of CR on CH

methylation, an observation that may be unique to neural tissues.

DNA methylation changes in the old hippocampus are observed in both CG and CH contexts (Masser et

al., 2017). These changes were enriched in inflammation related pathways, neuronal integrity, neuronal

function and synaptic transmission. Previous studies examining gene expression change with age in the

hippocampus showed enrichment in similar pathways (Keleshian et al., 2013; Stilling et al., 2014; Meng

et al., 2016; Mangold et al., 2017a; 2017b). Studies examining the relationship between gene expression

and methylation with age and CR (Hahn et al., 2017), and in the brain (Sharma et al., 2016) failed to

identify a simple correlation. While demonstrating causality between differential methylation of

thousands of sites and gene expression remains unfeasible, we hypothesize that methylation changes

with age contribute to the dysregulation of genes in these pathways. Based on our data, enrichment of

methylation changes with age commonly occurred in introns and genomic regulatory regions such as

open chromatin, CGI shores and CTCF binding sites and were under-represented in promoters and CG

islands. These findings may support a more complex role to methylation in regulating gene expression

during the aging process than simplistic alterations in promoters. This may also explain the discrepancies

between current studies and the canonical mechanisms by which methylation regulates expression.

Further investigation is required to discover the many intricate mechanisms by which methylation acts

as a genomic regulator with aging.

Over a third of all age-related differentially methylated cytosines were prevented by CR in age-matched

animals. The distribution of prevented sites in CG context was comparable to that of age-dmCGs,

however for CH sites CR preferentially prevented age-related hypermethylation. This may be caused by

the diet-specific reduction in expression of the DNA methylation regulatory enzymes, or via targeted

mechanisms such as recruitment of DNMTs and TETs to specific genomic locations by targeting co-

factors and changes to histone modifications and genome occupancy. In light of our results in which

both the methylation and demethylation machinery changed with diet and previous findings of altered

genomic regulation following CR (Chouliaras et al., 2013; Gong et al., 2015) we consider the later more

likely. It is worth noting that CR effects on age-related dmCs are classified as 'prevented', however the

possibility remains that these changes are delayed or reversed at different points throughout life. To

further elucidate the dynamics by which CR prevents, delays or reverses changes in DNA methylation

with aging, future studies examining time points across the lifespan are necessary.

The majority of CR-induced differentially methylated sites were independent of aging. These diet-

specific changes demonstrated different distributions in promoters, genes, and enhancers compared to

CR prevented age-related sites. Previous studies in the liver (Cole et al., 2017; Hahn et al., 2017)

reported enrichment of diet-induced CG hypermethylation in CG islands and gene bodies while CG

hypomethylation occurred in gene bodies and enhancers. Interestingly, we observed a similar

enrichment pattern in the brain. The commonalities in enrichment of diet-dmCGs in both liver and

hippocampus suggest common regulation of DNA methylation by diet across tissues that may be caused

by changes in regulation of common core proteins required for the targeting and recruitment of the

DNA methylation machinery to specific locations within the genome. Targeting of DNMTs and TETs to

specific genomic loci is a complex process that is not fully understood (Wu and Zhang, 2017). Post

translational histone modifications, have been positively and negatively correlated with DNA

modification (Lehnertz et al., 2003; Cedar and Bergman, 2009; Balasubramanian et al., 2012) and may be

altered through diet, thus changing recruitment of DNMTs and TETs. In the brain, CR attenuates age-

changes of DNMT3a and other epigenetic marks (Bordone and Guarente, 2005; Chouliaras et al., 2013;

Gong et al., 2015). Additionally, insulin signaling, specifically the PI3K/AKT pathway, a conserved

pathway altered by CR, was shown to modify DNA methylation through destabilization of DNMTs

(Popkie et al., 2010)

Genomic CH methylation is higher in the brain compared to other tissues (Lister et al., 2009), and has

been show to regulate transcription of genes involved in synaptic and neural growth (Chen et al., 2015;

Jang et al., 2017). Differential CH methylation has not been examined previously in the context of CR

and aging. Here we demonstrated that changes induced by diet in CH methylation are not sporadic and

are clustered at higher density than age-associated changes. Surprisingly diet-induced gain of

methylation at CH loci is observed in regions where CH methylation is often excluded such as promoters,

CG islands, TF binding sites and active enhancers (Lister et al., 2009; Laurent et al., 2010; Xie et al., 2012;

Guo et al., 2014). These findings provide evidence for a more controlled regulation of DNA methylation

following CR, potentially lost with aging. Genomic regulation of CH methylation is different than CGs but

remains largely unknown due to the lack of CH data from DNA methylation analyses. In the context of

CR, methylation of CHs appears to respond differently than CGs and serve a non-redundant role in

genomic regulation because 1) enrichment of dmCGs and dmCHs was different at promoters, TF binding

sites, enhancers and open chromatin and 2) the affinity of the DNMTs and TETs to CHs is lower than that

of CGs (Ramsahoye et al., 2000; Suetake et al., 2003; He and Ecker, 2015). The differential localization of

dmCHs as compared to dmCGs provides evidence that different, and as yet unknown, targeting

mechanisms are being used to regulate CH and CG methylation with diet.

Several pathways are known to be upregulated by food-restriction and its mimetics (Fusco and Pani,

2013). Prior gene expression studies in the brain have shown that mTOR signaling and insulin signaling

are downregulated by CR, as they are in other tissues, while brain-specific induction of pathways

including calcium signaling, synaptic vesicle regulation, neurotrophic signaling, neural growth,

differentiation and function, and neuronal integrity are also observed (Schafer et al., 2015; Wood et al.,

2015). We observed diet-induced methylation changes in pathways previously reported to change

following CR in the brain, specifically calcium signaling, energy metabolism, NGF signaling, FGFR

signaling, and pathways involved in axon guidance and growth. While a direct effect was not

established, the common findings between our study and previous gene expression studies suggest a

role for DNA methylation as a possible mechanism for the neuroprotective effect of CR against brain

aging.

Altered methylation patterns were also observed in DNA methylation regulatory pathways. Long-term

CR induced hypermethylation of both CGs and CHs in the promoters of TET3 and DNMT1, accompanied

by reduced expression with diet indicating a potential auto-regulation of the DNA methylation

machinery with diet. CR was reported to decrease TET2 and DNMT3b and increase TET3 expression in

the liver (Hahn et al., 2017) and decrease DNMT3a in the hippocampus (Chouliaras et al., 2011). Both

TET3 and DNMT1 were not previously reported to change with diet in the brain. Consistent with

previous results, DNMT3a1 mRNA expression decreased following CR. DNMT3a1 is considered the

primary enzyme for *de novo* CH methylation along with DNMT3b, however DNMT3b is expressed at low

levels in the brain (He and Ecker, 2015). Based on our findings of brain-specific differential expression of

DNMTs and TETs with CR and enrichment of differential methylation in pathways involved in neuronal

function, we propose that DNA methylation underlies brain-specific protective effects by CR, however

the targeting and recruitment of DNMTs and TETs to their genomic locations shares commonalities

across tissues as evident by enrichment of methylation changes at gene regulatory regions. We are the

first to report auto-regulation of DNA methylation by diet which appears to be specific to the brain.

Taken together our data supports a role of calorie-restriction in maintaining homeostasis through DNA

methylation. Changes in methylation were enriched in pathways shown to improve following CR and are

beneficial to brain aging. By including CH methylation analysis, we demonstrate similarities and

dissimilarities in the functions of CG and CH methylation in the protective effect induced by CR. These

findings highlight the importance in investigating both CG and CH methylation and propose a prominent

systemic role for DNA methylation in response to CR. Future studies are required in order to further

parse the specific mechanisms of diet-induced DNA methylation regulation and how these promote

brain health in old age.

References

Adams MM, Shi L, Linville MC, Forbes ME, Long AB, Bennett C, Newton IG, Carter CS, Sonntag WE, Riddle

DR, Brunso-Bechtold JK (2008) Caloric restriction and age affect synaptic proteins in

hippocampal CA3 and spatial learning ability. Exp Neurol 211:141-149.

Balasubramanian D, Akhtar-Zaidi B, Song L, Bartels CF, Veigl M, Beard L, Myeroff L, Guda K, Lutterbaugh

J, Willis J, Crawford GE, Markowitz SD, Scacheri PC (2012) H3K4me3 inversely correlates with

DNA methylation at a large class of non-CpG-island-containing start sites. Genome Med 4:47.

Barger JL, Vann JM, Cray NL, Pugh TD, Mastaloudis A, Hester SN, Wood SM, Newton MA, Weindruch R,

Prolla TA (2017) Identification of tissue-specific transcriptional markers of caloric restriction in

the mouse and their use to evaluate caloric restriction mimetics. Aging Cell 16:750-760.

Bordone L, Guarente L (2005) Calorie restriction, SIRT1 and metabolism: understanding longevity. Nat

Rev Mol Cell Biol 6:298-305.

Cedar H, Bergman Y (2009) Linking DNA methylation and histone modification: patterns and paradigms.

Nat Rev Genet 10:295-304.

Chen L, Chen K, Lavery LA, Baker SA, Shaw CA, Li W, Zoghbi HY (2015) MeCP2 binds to non-CG

methylated DNA as neurons mature, influencing transcription and the timing of onset for Rett

syndrome. Proc Natl Acad Sci U S A 112:5509-5514.

Chouliaras L, van den Hove DL, Kenis G, Dela Cruz J, Lemmens MA, van Os J, Steinbusch HW, Schmitz C,

Rutten BP (2011) Caloric restriction attenuates age-related changes of DNA methyltransferase

3a in mouse hippocampus. Brain Behav Immun 25:616-623.

Chouliaras L, van den Hove DL, Kenis G, Draanen M, Hof PR, van Os J, Steinbusch HW, Schmitz C, Rutten

BP (2013) Histone deacetylase 2 in the mouse hippocampus: attenuation of age-related increase

by caloric restriction. Curr Alzheimer Res 10:868-876.

Cole JJ, Robertson NA, Rather MI, Thomson JP, McBryan T, Sproul D, Wang T, Brock C, Clark W, Ideker T,

Meehan RR, Miller RA, Brown-Borg HM, Adams PD (2017) Diverse interventions that extend

mouse lifespan suppress shared age-associated epigenetic changes at critical gene regulatory

regions. Genome Biol 18:58.

Colman RJ, Beasley TM, Kemnitz JW, Johnson SC, Weindruch R, Anderson RM (2014) Caloric restriction

reduces age-related and all-cause mortality in rhesus monkeys. Nat Commun 5:3557.

Dacks PA, Moreno CL, Kim ES, Marcellino BK, Mobbs CV (2013) Role of the hypothalamus in mediating

protective effects of dietary restriction during aging. Front Neuroendocrinol 34:95-106.

Dietrich MO, Antunes C, Geliang G, Liu ZW, Borok E, Nie Y, Xu AW, Souza DO, Gao Q, Diano S, Gao XB,

Horvath TL (2010) Agrp neurons mediate Sirt1's action on the melanocortin system and energy

balance: roles for Sirt1 in neuronal firing and synaptic plasticity. J Neurosci 30:11815-11825.

Feng J, Zhou Y, Campbell SL, Le T, Li E, Sweatt JD, Silva AJ, Fan G (2010) DNMT1 and DNMT3a maintain

DNA methylation and regulate synaptic function in adult forebrain neurons. Nat Neurosci

13:423-430.

Fontan-Lozano A, Lopez-Lluch G, Delgado-Garcia JM, Navas P, Carrion AM (2008) Molecular bases of

caloric restriction regulation of neuronal synaptic plasticity. Mol Neurobiol 38:167-177.

Fusco S, Pani G (2013) Brain response to calorie restriction. Cell Mol Life Sci 70:3157-3170.

Gong H, Qian H, Ertl R, Astle CM, Wang GG, Harrison DE, Xu X (2015) Histone modifications change with

age, dietary restriction and rapamycin treatment in mouse brain. Oncotarget 6:15882-15890.

Gray DT, Barnes CA (2015) Distinguishing adaptive plasticity from vulnerability in the aging

hippocampus. Neuroscience 309:17-28.

Grigorenko EL, Kornilov SA, Naumova OY (2016) Epigenetic regulation of cognition: A circumscribed

review of the field. Dev Psychopathol 28:1285-1304.

Guo JU, Su Y, Shin JH, Shin J, Li H, Xie B, Zhong C, Hu S, Le T, Fan G, Zhu H, Chang Q, Gao Y, Ming GL, Song

H (2014) Distribution, recognition and regulation of non-CpG methylation in the adult

mammalian brain. Nat Neurosci 17:215-222.

Hadad N, Masser DR, Logan S, Wronowski B, Mangold CA, Clark N, Otalora L, Unnikrishnan A, Ford MM,

Giles CB, Wren JD, Richardson A, Sonntag WE, Stanford DR, Freeman W (2016) Absence of

genomic hypomethylation or regulation of cytosine-modifying enzymes with aging in male and

female mice. Epigenetics Chromatin 9:30.

Hahn O, Gronke S, Stubbs TM, Ficz G, Hendrich O, Krueger F, Andrews S, Zhang Q, Wakelam MJ, Beyer A,

Reik W, Partridge L (2017) Dietary restriction protects from age-associated DNA methylation and

induces epigenetic reprogramming of lipid metabolism. Genome Biol 18:56.

Halagappa VK, Guo Z, Pearson M, Matsuoka Y, Cutler RG, Laferla FM, Mattson MP (2007) Intermittent

fasting and caloric restriction ameliorate age-related behavioral deficits in the triple-transgenic

mouse model of Alzheimer's disease. Neurobiol Dis 26:212-220.

Hall A, Lalli G (2010) Rho and Ras GTPases in axon growth, guidance, and branching. Cold Spring Harb

Perspect Biol 2:a001818.

He Y, Ecker JR (2015) Non-CG Methylation in the Human Genome. Annu Rev Genomics Hum Genet

16:55-77.

Hickman RA, Faustin A, Wisniewski T (2016) Alzheimer Disease and Its Growing Epidemic: Risk Factors,

Biomarkers, and the Urgent Need for Therapeutics. Neurol Clin 34:941-953.

Hing B, Ramos E, Braun P, McKane M, Jancic D, Tamashiro KL, Lee RS, Michaelson JJ, Druley TE, Potash JB

(2015) Adaptation of the targeted capture Methyl-Seq platform for the mouse genome

identifies novel tissue-specific DNA methylation patterns of genes involved in

neurodevelopment. Epigenetics 10:581-596.

Hornsby AK, Redhead YT, Rees DJ, Ratcliff MS, Reichenbach A, Wells T, Francis L, Amstalden K, Andrews

ZB, Davies JS (2016) Short-term calorie restriction enhances adult hippocampal neurogenesis

and remote fear memory in a Ghsr-dependent manner. Psychoneuroendocrinology 63:198-207.

Hyun DH, Emerson SS, Jo DG, Mattson MP, de Cabo R (2006) Calorie restriction up-regulates the plasma

membrane redox system in brain cells and suppresses oxidative stress during aging. Proc Natl

Acad Sci U S A 103:19908-19912.

Inoue K, Zorrilla EP, Tabarin A, Valdez GR, Iwasaki S, Kiriike N, Koob GF (2004) Reduction of anxiety after

restricted feeding in the rat: implication for eating disorders. Biol Psychiatry 55:1075-1081.

Jang HS, Shin WJ, Lee JE, Do JT (2017) CpG and Non-CpG Methylation in Epigenetic Gene Regulation and

Brain Function. Genes (Basel) 8.

Kaeberlein M, Powers RW, 3rd, Steffen KK, Westman EA, Hu D, Dang N, Kerr EO, Kirkland KT, Fields S,

Kennedy BK (2005) Regulation of yeast replicative life span by TOR and Sch9 in response to

nutrients. Science 310:1193-1196.

Keleshian VL, Modi HR, Rapoport SI, Rao JS (2013) Aging is associated with altered inflammatory,

arachidonic acid cascade, and synaptic markers, influenced by epigenetic modifications, in the

human frontal cortex. J Neurochem 125:63-73.

Kim CH, Lee EK, Choi YJ, An HJ, Jeong HO, Park D, Kim BC, Yu BP, Bhak J, Chung HY (2016) Short-term

calorie restriction ameliorates genomewide, age-related alterations in DNA methylation. Aging

Cell.

Kisby GE, Kohama SG, Olivas A, Churchwell M, Doerge D, Spangler E, de Cabo R, Ingram DK, Imhof B, Bao

G, Kow YW (2010) Effect of caloric restriction on base-excision repair (BER) in the aging rat brain.

Exp Gerontol 45:208-216.

Krueger F, Andrews SR (2011) Bismark: a flexible aligner and methylation caller for Bisulfite-Seq

applications. Bioinformatics 27:1571-1572.

Laurent L, Wong E, Li G, Huynh T, Tsirigos A, Ong CT, Low HM, Kin Sung KW, Rigoutsos I, Loring J, Wei CL

(2010) Dynamic changes in the human methylome during differentiation. Genome Res 20:320-

331.

Lee GD, Wilson MA, Zhu M, Wolkow CA, de Cabo R, Ingram DK, Zou S (2006) Dietary deprivation extends

lifespan in Caenorhabditis elegans. Aging Cell 5:515-524.

Lehnertz B, Ueda Y, Derijck AA, Braunschweig U, Perez-Burgos L, Kubicek S, Chen T, Li E, Jenuwein T,

Peters AH (2003) Suv39h-mediated histone H3 lysine 9 methylation directs DNA methylation to

major satellite repeats at pericentric heterochromatin. Curr Biol 13:1192-1200.

Lister R, Mukamel EA (2015) Turning over DNA methylation in the mind. Front Neurosci 9:252.

Lister R, Pelizzola M, Dowen RH, Hawkins RD, Hon G, Tonti-Filippini J, Nery JR, Lee L, Ye Z, Ngo QM,

Edsall L, Antosiewicz-Bourget J, Stewart R, Ruotti V, Millar AH, Thomson JA, Ren B, Ecker JR

(2009) Human DNA methylomes at base resolution show widespread epigenomic differences.

Nature 462:315-322.

Lister R et al. (2013) Global epigenomic reconfiguration during mammalian brain development. Science

341:1237905.

Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G (2013) The hallmarks of aging. Cell

153:1194-1217.

Lutter M, Krishnan V, Russo SJ, Jung S, McClung CA, Nestler EJ (2008) Orexin signaling mediates the

antidepressant-like effect of calorie restriction. J Neurosci 28:3071-3075.

Mangold CA, Wronowski B, Du M, Masser DR, Hadad N, Bixler GV, Brucklacher RM, Ford MM, Sonntag

WE, Freeman WM (2017a) Sexually divergent induction of microglial-associated

neuroinflammation with hippocampal aging. J Neuroinflammation 14:141.

Mangold CA, Masser DR, Stanford DR, Bixler GV, Pisupati A, Giles CB, Wren JD, Ford MM, Sonntag WE,

Freeman WM (2017b) CNS-wide Sexually Dimorphic Induction of the Major Histocompatibility

Complex 1 Pathway With Aging. J Gerontol A Biol Sci Med Sci 72:16-29.

Masser DR, Hadad N, Porter H, Mangold CA, Unnikrishnan A, Ford MM, Giles CB, Georgescu C,

Dozmorov MG, Wren JD, Richardson A, Stanford DR, Freeman WM (2017) Sexually divergent

DNA methylation patterns with brain aging. Aging Cell *in press.*

Masser DR, Stanford DR, Hadad N, Giles CB, Wren JD, Sonntag WE, Richardson A, Freeman WM (2016)

Bisulfite oligonucleotide-capture sequencing for targeted base- and strand-specific absolute 5-

methylcytosine quantitation. Age (Dordr) 38:49.

Meng G, Zhong X, Mei H (2016) A Systematic Investigation into Aging Related Genes in Brain and Their

Relationship with Alzheimer's Disease. PLoS One 11:e0150624.

Park JH, Glass Z, Sayed K, Michurina TV, Lazutkin A, Mineyeva O, Velmeshev D, Ward WF, Richardson A,

Enikolopov G (2013) Calorie restriction alleviates the age-related decrease in neural progenitor

cell division in the aging brain. Eur J Neurosci 37:1987-1993.

Popkie AP, Zeidner LC, Albrecht AM, D'Ippolito A, Eckardt S, Newsom DE, Groden J, Doble BW, Aronow

B, McLaughlin KJ, White P, Phiel CJ (2010) Phosphatidylinositol 3-kinase (PI3K) signaling via

glycogen synthase kinase-3 (Gsk-3) regulates DNA methylation of imprinted loci. J Biol Chem

285:41337-41347.

Quinlan AR, Hall IM (2010) BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics 26:841-842.

Ramsahoye BH, Biniszkiewicz D, Lyko F, Clark V, Bird AP, Jaenisch R (2000) Non-CpG methylation is

prevalent in embryonic stem cells and may be mediated by DNA methyltransferase 3a. Proc Natl

Acad Sci U S A 97:5237-5242.

Redman LM, Ravussin E (2009) Endocrine alterations in response to calorie restriction in humans. Mol

Cell Endocrinol 299:129-136.

Reeve A, Simcox E, Turnbull D (2014) Ageing and Parkinson's disease: why is advancing age the biggest

risk factor? Ageing Res Rev 14:19-30.

Reitz C, Brayne C, Mayeux R (2011) Epidemiology of Alzheimer disease. Nat Rev Neurol 7:137-152.

Roadmap Epigenomics C et al. (2015) Integrative analysis of 111 reference human epigenomes. Nature

518:317-330.

Schafer MJ, Dolgalev I, Alldred MJ, Heguy A, Ginsberg SD (2015) Calorie Restriction Suppresses Age-

Dependent Hippocampal Transcriptional Signatures. PLoS One 10:e0133923.

Seib DR, Martin-Villalba A (2015) Neurogenesis in the Normal Ageing Hippocampus: A Mini-Review.

Gerontology 61:327-335.

Sharma A, Klein SS, Barboza L, Lohdi N, Toth M (2016) Principles Governing DNA Methylation during

Neuronal Lineage and Subtype Specification. J Neurosci 36:1711-1722.

Sonntag WE, Lynch CD, Cefalu WT, Ingram RL, Bennett SA, Thornton PL, Khan AS (1999) Pleiotropic

effects of growth hormone and insulin-like growth factor (IGF)-1 on biological aging: inferences

from moderate caloric-restricted animals. J Gerontol A Biol Sci Med Sci 54:B521-538.

Stilling RM, Benito E, Gertig M, Barth J, Capece V, Burkhardt S, Bonn S, Fischer A (2014) De-regulation of

gene expression and alternative splicing affects distinct cellular pathways in the aging

hippocampus. Front Cell Neurosci 8:373.

Suetake I, Miyazaki J, Murakami C, Takeshima H, Tajima S (2003) Distinct enzymatic properties of

recombinant mouse DNA methyltransferases DNMT3a and DNMT3b. J Biochem 133:737-744.

Swindell WR (2009) Genes and gene expression modules associated with caloric restriction and aging in

the laboratory mouse. BMC Genomics 10:585.

Turturro A, Witt WW, Lewis S, Hass BS, Lipman RD, Hart RW (1999) Growth curves and survival

characteristics of the animals used in the Biomarkers of Aging Program. J Gerontol A Biol Sci

Med Sci 54:B492-501.

Vanguilder HD, Freeman WM (2011) The hippocampal neuroproteome with aging and cognitive decline:

past progress and future directions. Front Aging Neurosci 3:8.

Vanguilder HD, Bixler GV, Sonntag WE, Freeman WM (2012) Hippocampal expression of myelin-

associated inhibitors is induced with age-related cognitive decline and correlates with deficits of

spatial learning and memory. J Neurochem 121:77-98.

Wang J, Ho L, Qin W, Rocher AB, Seror I, Humala N, Maniar K, Dolios G, Wang R, Hof PR, Pasinetti GM

(2005) Caloric restriction attenuates beta-amyloid neuropathology in a mouse model of

Alzheimer's disease. FASEB J 19:659-661.

Willette AA, Coe CL, Colman RJ, Bendlin BB, Kastman EK, Field AS, Alexander AL, Allison DB, Weindruch

RH, Johnson SC (2012a) Calorie restriction reduces psychological stress reactivity and its

association with brain volume and microstructure in aged rhesus monkeys.

Psychoneuroendocrinology 37:903-916.

Willette AA, Bendlin BB, Colman RJ, Kastman EK, Field AS, Alexander AL, Sridharan A, Allison DB,

Anderson R, Voytko ML, Kemnitz JW, Weindruch RH, Johnson SC (2012b) Calorie restriction

reduces the influence of glucoregulatory dysfunction on regional brain volume in aged rhesus

monkeys. Diabetes 61:1036-1042.

Wood SH, van Dam S, Craig T, Tacutu R, O'Toole A, Merry BJ, de Magalhaes JP (2015) Transcriptome

analysis in calorie-restricted rats implicates epigenetic and post-translational mechanisms in

neuroprotection and aging. Genome Biol 16:285.

Wu X, Zhang Y (2017) TET-mediated active DNA demethylation: mechanism, function and beyond. Nat

Rev Genet.

Wyss-Coray T (2016) Ageing, neurodegeneration and brain rejuvenation. Nature 539:180-186.

Xekardaki A, Kovari E, Gold G, Papadimitropoulou A, Giacobini E, Herrmann F, Giannakopoulos P, Bouras

C (2015) Neuropathological changes in aging brain. Adv Exp Med Biol 821:11-17.

Xie W, Barr CL, Kim A, Yue F, Lee AY, Eubanks J, Dempster EL, Ren B (2012) Base-resolution analyses of

sequence and parent-of-origin dependent DNA methylation in the mouse genome. Cell 148:816-

831.

Zerbino DR, Wilder SP, Johnson N, Juettemann T, Flicek PR (2015) The ensembl regulatory build.

Genome Biol 16:56.

Ziller MJ, Hansen KD, Meissner A, Aryee MJ (2015) Coverage recommendations for methylation analysis

by whole-genome bisulfite sequencing. Nat Methods 12:230-232

Figure Legends

Figure 1: Caloric-restriction prevents age-associated differential CG methylation. **A.** Scatterplot depicting

significant age-dmCG (n = 41,586, adjusted p-value < 0.05 & methylation difference between Old-AL and

Young-AL > |5%|, Kruskal-Wallis H test with Conover pairwise comparisons followed by false-discovery

rate multiple testing corrections) methylation levels between young *ad libitum* (Y-AL) and old *ad libitum*

(O-AL) groups (red dots) and young *ad libitum* (Y-AL) and old calorie-restricted (O-CR) groups (blue dots).

Linear regression lines pass through Y-AL ~ Y-AL (black), O-AL ~ Y-AL (blue) and O-CR ~ Y-AL (dotted

green) age-dmCG methylation values separated by hyper- and hypo- age-dmCGs. Calorie-restriction diet

causes a 'younger' methylation profile in age-associated differentially methylated sites. **B.** Single base

representatives of an age-dmCG (left) and a prevented age-dmCG (right). **C.** Quantification of the

number of age-dmCGs by hypermethylation (blue) and hypomethylation (red) in animals fed *ad libitum*

(AL) or calorie-restricted (CR) diet. Grey indicates age-dmCGs prevented by caloric-restriction

(n=13,950). **D.** Distribution plot of age-dmCGs (red) and prevented age-dmCGs (blue). Age-dmCGs and

those prevented by CR follow a similar distribution. **E, F.** PCA plots of age-dmCGs **(E)** and prevented age-

dmCGs (**F**) showing clustering of old animals in age-dmCGs but separation in prevented age-dmCGs. **G.**

Enrichment of age-dmCGs and prevented age-dmCGs in gene-centric regions, only significant

enrichments are shown (p < 0.05, hypergeometric test). **H.** Enrichment of age-dmCGs and prevented

age-dmCGs in gene-regulatory regions. Red indicates under-representation (log(odds ratio) < 0 & p-

value < 0.05, hypergeometric test), blue indicates over-representation (log(odds ratio) > 0 & p-value <

0.05, hypergeometric test) and white shows no significance (p-value > 0.05). **I.** Pathway analysis of genes

containing significant age-dmCGs or prevented age-dmCGs. Top enrichment pathways are shown (FDR

adjusted p-values < 0.01). Circle size indicates the ratio of genes containing age-dmCGs or prevented

age-dmCGs in a pathway, color indicates p-values. No pathways were enriched for genes containing >2

prevented hypomethylated age-dmCGs.

Figure 2: Caloric-restriction prevents age-associated differential CH methylation. **A.** Scatterplot depicting

significant age-dmCH (n = 79,056, adjusted p < 0.05 & methylation difference between Old-AL and

Young-AL > |5%|, Kruskal-Wallis H test with Conover pairwise comparisons followed by false-discovery

rate multiple testing corrections) methylation levels between young *ad libitum* (Y-AL) and old *ad libitum*

(O-AL) groups (red dots) and young *ad lib* (Y-AL) and old calorie-restricted (O-CR) groups (blue dots).

Linear regression lines pass through Y-AL ~ Y-AL (black), O-AL ~ Y-AL (blue) and O-CR ~ Y-AL (dotted

green) age-dmCH methylation values separated by hyper- and hypo- age-dmCHs. **B.** Quantification of

the number of age-dmCHs by hypermethylation (blue) and hypomethylation (red) in animals fed *ad*

libitum (AL) or calorie-restricted (CR) diet. Grey indicates age-dmCHs prevented by caloric-restriction (n

= 29,880). **C.** Distribution plot of age-dmCHs (red) and prevented age-dmCHs (blue). The probability of

hypermethylated age-dmCH to be prevented by diet was higher than that of hypomethylated age-dmCH

(0.59 prevented hyper compared to 0.11 prevented hypos). **D, E.** PCA plots of age-dmCHs (**D**) and

prevented age-dmCHs (**E**). **F.** Enrichment of age-dmCHs and prevented age-dmCHs in gene-centric

regions, only significant enrichments are shown (p < 0.05, hypergeometric test). **G.** Enrichment of age-

dmCHs and prevented age-dmCHs in gene-regulatory regions. Red indicates under-representation (log

(odds ratio) < 0 & p-value < 0.05, hypergeometric test), blue indicates over-representation (log(odds

ratio) > 0 & p-value < 0.05, hypergeometric test) and white blocks did not reach statistical significance **H.**

Pathway analysis of genes containing significant age-dmCHs or prevented age-dmCHs. Top enrichment

pathways are shown (FDR adjusted p-values < 0.01). Circle size indicates the ratio of genes containing >2

age-dmCHs or prevented age-dmCHs in a pathway, color indicates p-values. I. Overlap between

pathways enriched for age-dmCHs and age-dmCGs.

Figure 3: CR induces changes in CG methylation independent of aging. **A.** *Left*: 3D scatterplot showing

the methylation levels of significant diet-dmCGs including unique diet-dmCGs (green) and age-dmCGs

prevented by CR (blue) (adjusted p < 0.05 & methylation difference between Old-AL and Old-CR > |5%|,

Kruskal-Wallis H test with Conover pairwise comparisons followed by false-discovery rate multiple

testing corrections) focusing on young-AL and old-AL. *Right*: 90° rotation on the Y-axis of the 3D plot

showing that unique diet-dmCGs (green) regress away from the midline when comparing old-AL and

Old-CR methylation levels. **B.** Proportion of diet-dmCGs unique to CR (green) and those with an age

interaction (prevented age-dmCGs, blue) (N = 46,052). **C.** Single base example of a unique diet-dmCG. **D.**

Distribution of diet-dmCGs. Dotted lines mark the average methylation differences of hypomethylated

diet-dmCGs (red, -12.5%) and hypermethylated diet-dmCGs (green, 12.3%). **E.** Enrichment of diet-

dmCGs in gene-centric regions (p < 0.05, hypergeometric test). **F.** Enrichment of diet-dmCGs in gene-

regulatory regions. Red indicates under-representation (log odds ratio < 0 & p-value < 0.05,

hypergeometric test), blue indicates over-representation (log odds ratio > 0 & p-value < 0.05,

hypergeometric test) and white show no significance **G.** Pathway analysis of genes containing >2

significant diet-dmCGs. Top enrichment pathways are shown (FDR adjusted p-values < 0.01). Circle size

indicates the ratio of genes containing diet-dmCGs in a pathway, color indicates p-values.

Figure 4: CR induced differential CH methylation is clustered at specific genomic loci**.** A. *Left*: 3D

scatterplot showing the methylation levels of significant diet-dmCHs including unique diet-dmCHs

(green) and age-dmCHs prevented by diet (blue) (adjusted p < 0.05 & methylation difference between

Old-AL and Old-CR > |5%|, Kruskal-Wallis H test with Conover pairwise comparisons followed by false-

discovery rate multiple testing corrections) focusing on young-AL and old-AL. *Right*: 90° rotation on the

Y-axis of the 3D plot showing unique diet-dmCH (green) regress away from the midline when comparing

old-AL and Old-CR methylation levels. **B.** Proportion of diet-dmCHs unique to CR (green) and those with

age interaction (prevented age-dmCHs, blue) (N = 90,428). **C.** Distribution of diet-dmCHs. Dotted lines

mark the average methylation differences of hypomethylated diet-dmCHs (red, 8%) and

hypermethylated diet-dmCHs (green, 10%). **D.** Distribution plot comparing the proximity between

dmCHs by age and diet focusing on 10 kb segments. **E.** Line graph representing % of dmCHs by degree of

proximity within 10kb (solid lines) or 1kb (dotted line) by age or by diet.

Figure 5: CR-induced differential CH methylation is enriched in CG-rich genomic regions. **A.** Enrichment

of diet-dmCHs in gene-centric regions, only significant enrichments are shown (p < 0.05, hypergeometric

test). Diet-dmCHs were found to be enriched in CG islands and shores, promoters and gene bodies. **B.**

Enrichment of diet-dmCHs in gene-regulatory regions. Red indicates under-representation (log(odds

ratio) < 0 & p-value < 0.05, hypergeometric test), blue indicates over-representation (log(odds ratio) > 0

& p-value < 0.05, hypergeometric test) and white show no significance **C.** Pathway analysis of genes

containing >2 significant diet-dmCHs. Top enrichment pathways are shown (FDR adjusted p-values <

0.01). Circle size indicates the ratio of genes in pathway containing diet-dmCHs, color indicates p-values.

D. Overlap between pathways enriched for diet-dmCHs and diet-dmCGs.

Figure 6: Diet induced differential methylation involved in auto-regulation of DNA regulatory enzymes.

A. Calorie-restriction induces hypermethylation of CGs (red) and CHs (green) in the promoter regions of

TET3 **B.** mRNA expression of TET3 decreases with diet but not age (n=7-8/group, $F_{(2,20)}$ = 11.1, ** p <

0.001, One-way ANOVA with Benjamini-Hochberg correction). **C.** Calorie-restriction induces

hypermethylation of CGs (red) and CHs (green) in the promoter regions of DNMT1. **D.** mRNA expression

of DNMT1 using an assay inclusive of all known DNMT1 isoforms show no significant differences with

age or diet (n=7-8/group, $F_{(2,20)}$ = 0.535, p > 0.05, One-way ANOVA with Benjamini-Hochberg correction).

E. Isoform specific assays were used to assess expression of DNMT1 show significant decrease in DNMT1

expression with age and diet in isoforms utilizing exon 1 but no difference in expression when

examining isoforms using the alternative exon 1 (n=7-8/group, $F_{(2,19)}$ = 13.42, * p < 0.05, **p < 0.001,

One-way ANOVA). **F.** mRNA expression of DNMT3a1 decreases with diet and age. TET1 expression did

not change in any condition and TET2 expression decreased with diet but not age (n=7-8/group,

DNMT3a1 $F_{(2,19)}$ = 18.59, Tet2$_{(2,19)}$ F = 10.6, TET1 $F_{(2,20)}$ = 3.2, *p < 0.05, **p < 0.001, One-way ANOVA

with Benjamini-Hochberg correction).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A.

B.

C.

D.

Figure 6