Twenty-five species of frogs in a liter of water: eDNA survey for exploring tropical frog diversity

Miklós Bálint[1,2], Carsten Nowak[1,2], Orsolya Márton[1,3], Steffen U. Pauls[1,2], Claudia Wittwer[1], Jose Luis Aramayo B.[4], Arne Schulze[5], Thierry Chambert[6], Berardino Cocchiararo[1], Martin Jansen[1]

[1]Senckenberg Research Institute and Natural History Museum Frankfurt, Senckenberganlage 25, 60325 Frankfurt, Germany

[2]LOEWE Centre for Translational Biodiversity Genomics (LOEWE-TBG), Senckenberganlage 25, 60325 Frankfurt, Germany

[3]Institute for Soil Sciences and Agricultural Chemistry, Centre for Agricultural Research, Hungarian Academy of Sciences, Herman Otto str. 15, H-1022 Budapest, Hungary

[4]Museo Historia Natural Noel Kempff Mercado (Universidad Autónoma Gabriel Rene Moreno), Av. Irala 565, Santa Cruz de la Sierra, Bolivia

[5]Hessisches Landesmuseum Darmstadt (HLMD), Friedensplatz 1, 64283 Darmstadt, Germany

[6]Pennsylvania State University, Department of Ecosystem Science and Management, University Park, PA 16802, USA

Abstract

Biodiversity is declining at unprecedented rates, with species extinctions being an important component of this process. Currently it is almost impossible to evaluate this loss since biodiversity data is lacking even for emblematic species. Recent advances in eDNA marker gene studies promise to deliver the necessary data, but these approaches have almost never been evaluated in the tropics. Here we evaluate whether 1) eDNA metabarcoding may reliably generate frog faunistic data in species-rich tropical areas, 2) the ecological signal in eDNA data is compatible with data from visual and audio encounter survey (VAES), and 3) eDNA data represent a financially viable solution to biodiversity data acquisition. We applied eDNA metabarcoding to investigate frog species occurrence in five ponds in the Chiquitano dry forest region in Bolivia and compared our data with a simultaneous VAES. We find that taxon lists generated with eDNA and VAES correspond closely, and most deviations are attributable to different species' life histories. The ecological signal in eDNA closely matches the signal from VAES. We find that the cost efficiency of eDNA surveys is mostly influenced by the richness of local fauna and the number of surveyed sites: VAES may be less costly in low-diversity regions, but eDNA quickly becomes more cost-efficient in high-diversity regions, particularly with many sites. The results underline that eDNA is suitable for large-scale biodiversity surveys in high-diversity areas if precautions in data generation and interpretation are taken and this projects spatially extensive standardized biodiversity surveys in the close future.

Keywords

Amphibians, metabarcoding, tropical biodiversity, conservation, community ecology, cost comparison

Introduction

Improvements on most biodiversity loss indicators lag behind the 20 "Aichi Biodiversity Targets" (UNEP 2016) that aim to reduce the decline of biodiversity by 2020 (Tittensor *et al.* 2014). An important component of the biodiversity crisis is the extinction of species: based on current trends in mammals, birds, reptiles and amphibians, it has been projected that the biodiversity crisis may lead to the 6th Mass Extinction over the next three centuries if all threatened species go extinct (Barnosky *et al.* 2011). Current rates of extinction may even be much higher if one considers the extinction that likely occurred during the last decades-centuries, but went unnoticed because the now-extinct species had small ranges, were never described or only described on the eve of their extinct (Pimm *et al.* 2014; Lees & Pimm 2015). It is however difficult to assess which species are endangered and to what extent. First, most taxa remain undescribed: in some highly diverse regions many species will likely go extinct before they are discovered (Costello *et al.* 2013; Lees & Pimm 2015). Second, cryptic genetic diversity is common within morphospecies (Pfenninger & Schwenk 2007; Pauls *et al.* 2013), and global change may impact cryptic diversity more severely than morphospecies (Bálint *et al.* 2011). Third, data on population-level trends is scarce, even for well-known species (Butchart *et al.* 2010). Better population-level biodiversity data is thus urgently needed to 1) understand biodiversity patterns and extinction threats, 2) improve forecasting abilities about future biodiversity, and 3) improve humanity's responses to the challenges of biodiversity loss. This data is crucial in times when conservation action is increasingly demanded by society (Tittensor *et al.* 2014).

The importance of internationally coordinated, standardized biodiversity data collection is long recognized both in science and in conservation (Henry *et al.* 2008). This is particularly true for the most biodiverse areas. The tropics are generally underrepresented in ecological studies (Clarke *et al.* 2017; Stroud & Feeley 2017). However, encounter-based data collection appears

more difficult here and sufficient funds and expertise are often missing for comprehensive

surveys. Indirect species records through environmental DNA are increasingly heralded as an

alternative to encounter-based surveys (Thomsen & Willerslev 2015; Pedersen *et al.* 2015).

eDNA also allows standardized biodiversity surveys on regional and global scales since

community composition data can be obtained by high-throughput-sequencing of standardized,

taxonomically informative marker genes (metabarcoding) (Taberlet *et al.* 2012a; Cristescu

2014). eDNA-based metabarcoding may present one of several tools needed to globally

coordinate initiatives for ecosystem monitoring and sustainable management (Bush *et al.* 2017;

Schmeller *et al.* 2017).

In this study we evaluate whether eDNA metabarcoding is suitable for inventories of

frogs, a group with particular high species-diversity in tropical regions. Frogs and other

amphibians are particularly hard hit by the biodiversity crisis: more than one-third of the

approximately 7500 described species are endangered (Stuart *et al.* 2004; Bishop *et al.* 2012;

Whittaker *et al.* 2013). Frogs are also known for being a highly diverse, but incompletely

described taxon, especially in the tropics (Ferrão *et al.* 2016; Caminer *et al.* 2017). Many

"widespread" morphospecies harbor considerable cryptic genetic diversity and are better

considered complexes of closely related species with much smaller ranges (Fouquet *et al.* 2007;

Gehara *et al.* 2013, 2014; Ortega-Andrade *et al.* 2015). Efficient actions for amphibian

conservation (e.g. the prioritization of areas for conservation, or informing society and

stakeholders about conservation needs) are only possible with geographically fine-scale,

taxonomically well-resolved faunistic data, but our current understanding of present and future

amphibian biodiversity is often based on rare, spatially and temporally scattered observations of

phenotypically defined taxa.

First efforts have been done to test the suitability of eDNA for the survey of tropical frog

biodiversity (Lopes *et al.* 2016), but important practical aspects remain unaddressed. First, it is

not clear which fraction of the local species pool is represented by amphibian eDNA in tropical

water bodies: existing comparisons of encounter-based surveys and eDNA either do not include

non-adult life stages, or they use already compiled fauna lists for the evaluation of eDNA

performance without consideration of life history traits or behavioral aspects at the moment of

sampling. This may confound biodiversity data collection (Petitot *et al.* 2014). Second,

ecological signal in aquatic eDNA data remains largely unevaluated: it is not clear whether it

reflects ecological conditions. Most studies to date only evaluate the correspondence between

encounter-based and metabarcoded taxon lists (Miya *et al.* 2015; Valentini *et al.* 2016),

although the ecological signal itself may be of higher importance for many applications (Ji *et al.*

2013; Elbrecht *et al.* 2017). Third, it is not clear whether, and under what conditions eDNA is

financially efficient since comparisons are lacking (Lopes *et al.* 2016), although these

comparisons are essential for deciding on data collection strategies.

Here we particularly address whether 1) detectability of tropical amphibians with eDNA

is linked to species' life history, and 2) there is sufficient ecological signal in amphibian eDNA

data to allow inferences about biodiversity patterns. Finally, we present a framework for cost

comparisons between encounter- and eDNA-based biodiversity survey that may be adapted to

other systems beyond amphibians. We compare the results of long- and short-term encounter-

based field surveys, and an eDNA survey of tropical amphibians in a high-diversity area (45

species of frogs on ca. 30 km^2). This area is well characterized with respect to amphibians,

including both larvae and adults (Schulze et al. 2015), and cryptic intraspecific diversity (e.g.

Jansen et al. 2011, Jansen *et al.* 2016). The area is located the Chiquitano region of Bolivia, a

forest-savanna ecotone between Amazon, Cerrado and Gran Chaco in a transition zone among

humid and dry forests that are special in regard to their taxonomic and functional diversity

(Castro *et al.* 1999). The region contains the largest intact, old-growth block of seasonally dry

tropical forests in South America (Miles *et al.* 2006; Power *et al.* 2016).

Materials and methods

A detailed description of the Materials and methods is provided in the Supplementary

materials.

The study area is near the Biological Station "Centro de Investigaciones Ecológicas

Chiquitos" on the San Sebastián cattle ranch (S16.3622, W62.00225, 500 m a.s.l.), 24 km south

of the town of Concepción, Province of Ñuflo de Chávez, Santa Cruz Department, Bolivia. A

description of the area is given by Schulze *et al.* (2009). Intensive long-term assessments have

resulted in a list of 45 frog species in the area (e.g. Jansen 2009; Schulze *et al.* 2009, 2015;

Jansen *et al.* 2011). We selected five ponds for this study. eDNA samples were obtained in

three technical replicates from three sampling points on each pond (Fig. 1). All samples and

replicates were filtered with a two-step sequential filtering scheme: first, two glass fiber filters

(GFF) with a pore size of 2 µm (filter method A: preserved in CTAB; B: dried), and second, the

flow-through was filtered with a nylon filter with (pore size 0.2 µm, method C, dried). Combined

visual and audio encounter surveys (VAES, Zimmerman 1994), and a tadpole survey (TS,

Schulze *et al.* 2015) were conducted by experienced observers (MJ and AS). Frogs were

located at night (between 21:00 and 00:30 hours) during 0.5 - 1 hrs transect walks along the

ponds to detect and identify specimens in vegetation, water and on ground around ponds with

flashlights and headlamps, and by listening. Tadpoles were searched once daytime and once at

night in each pond and the riparian vegetation, and caught with dip nets.

eDNA samples were processed in Germany in a low-DNA laboratory. DNA was

extracted from GFF samples (A, B, see above) with a CTAB chloroform extraction method

according to Strand *et al.* (2014) and Wittwer et al. (2017). Dried nylon filters (C) were extracted

with DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) following Thomsen *et al.* (2012).

Samples were fully randomized before extraction to prevent batch effects (Bálint *et al.* 2017a).

The barcode amplification targeted mitochondrial 16S ribosomal RNA. We designed primers for

a 150 bp fragment with the reference database of local species (sequences from 159

specimens, Jansen *et al.* 2011; Schulze *et al.* 2015). The primers were tested on a subset of 12

species (Supplementary Table S3). PCRs were run on 96-well plates in 15-µl reaction volumes

with a touchdown protocol, with four PCR technical replicates per sample. We included four

negative PCR controls (ultra-sterile water), four extraction blanks, and two positive controls on

each plate. The amplification primers contained multiplexing indices. PCR products were

purified, pooled in equal volumes, and paired-end-sequenced (2 x 150 bp) on an Illumina

NextSeq 500 sequencer.

Bioinformatics were done with OBITools v1.2.0 (Boyer *et al.* 2016). We performed two

reference-based taxonomic assignments, first with a custom database of 16S sequences of 159

specimens from the regional amphibian fauna (16S_custom), and second with all 16S

sequences found in the EMBL (release 125, 16S_EMBL). Further filtering was done in R (R

Core Team 2017) with a script supplied on GitHub: https://github.com/MikiBalint/amphibian-

eDNA.

We followed recommendations for the statistical analyses of marker gene community

data (Bálint *et al.* 2016). eDNA faunistic differences among the five ponds were visualized with a

latent variable model-based ordination in R ('boral', Hui *et al.* 2015), and tested with

multispecies generalized linear models ('mvabund', Wang *et al.* 2012). The surveyed ponds

were markedly different in size, vegetation, water depth, etc, thus we considered the pond

identity as a good predictor of community composition. We performed a model-based ordination

also for the VAES presence/absence data and then used a Procrustean superimposition (Peres-

Neto & Jackson 2001) to evaluate how the VAES-based ordination of ponds matches the

ordination of centroids of eDNA samples. We compared the efficiency of filter preservation

(CTAB or dried) on the successful detection of species with a site occupancy model (MacKenzie

et al. 2002; Bailey *et al.* 2014), implemented in the R package 'unmarked' (Fiske *et al.* 2011).

We used the single season false-positive occupancy model developed by Miller *et al.* (2011).

For cost comparisons, we considered a typical eDNA survey scenario: samples are

collected in the field and later processed in a dedicated laboratory. For the encounter-based

survey we considered a scenario with a similar separation of the fieldwork, and species

identification: species records (audio or visual) are collected by a field biologist, and later

identified by an expert in the office/laboratory. The parameters in our cost model are a good

approximation of the variables involved in the present study and involve a learning effect in the

efficiency of the taxonomic expert. The sampling and identification costs of VAES are

dependent both on the number of sampling sites, and the number of species since each species

needs to be recorded (Fig. 4). During the eDNA survey eDNA samples are collected by a field

biologist, analysed in a lab and sequenced by an external provider. The sampling and

identification costs of eDNA sampling costs of eDNA depend on the number of sites, but not on

the number of species (Fig. 4). We kept some cost factors constant: the costs for training the

frog taxonomic expert and the VAES observers, the costs for building up the eDNA

metabarcoding facilities (clean rooms and equipments to perform DNA manipulations, but not

the sequencing), and the databases necessary for the sequence assignment. We assume that

travel costs are the same for the two survey types, and that the time necessary to walk between

frog observations and eDNA sampling points is the same. All model parameters and

calculations are accessible on FigShare (Bálint *et al.* 2017b;

https://figshare.com/articles/_/5099842).

Results

The sequencing resulted in 12 742 273 read pairs (deposited in ENA as PRJEB22113**)**,

of which 9 479 299 were identified as complete 16S amplicons. These reads were de-replicated

into 631 003 unique sequences variants. Only 22 706 of these variants were represented by at

least 10 reads and retained for further processing. The sequence cleanup resulted in 14 442

high quality sequence variants, 13 497 coming from the present experiment. Of these, 4 805

sequence variants were taxonomically assigned with the 16S_custom database, and 8 692 with

the 16S_EMBL database. The assigned sequence variants represented 8 011 631 sequences,

of which 561 sequence variants were kept for further analyses as they were recorded as 'head'

in at least one sample (6 965 866 reads). After the removal of potential contaminant reads 5 898

184 sequences were retained. The final sample - sequence variant abundance matrix contained

5 815 014 sequences. These belonged to several groups: frogs (2 158 534), fish (1 692 613),

insects (304 059), mammals (14 006), birds (967), and bacteria and other groups (1 063 156).

In the five ponds 31 frog species were detected in total with both methods. The VAES

and the eDNA metabarcoding detected 25 species each, and TS detected 4 species; 19 species

were detected by both eDNA survey and VAES (Fig. 2a; Supplementary Table S1). Six species

were detected only by eDNA, and six species were detected only by VAES. Several of the

species detected by VAES but not by eDNA had no aquatic life phase during the time of survey

(i.e. no larvae or adults in the water). Of the species that were present in the water (N=20),

eDNA detected 19 (Fig. 2b, Supplementary Table S1). With a single sampling eDNA detected

more than half of the 45 species known to be present in the area (Fig. 2c, Supplementary Table

S2). Of these 45 species only 35 were ever detected at the surveyed ponds, and eDNA

detected 23 of those (Fig. 2d, Supplementary Table S2)**.** The eDNA results reported two

additional species that were not known during the time of the survey from the area (Fig. 2c,d).

We detected 11 species (of the twelve) from the equimolar DNA concentration positive controls,

and 6 in the non-equimolar DNA concentration controls (Supplementary Fig. S2). Read

numbers in the equimolar PCR controls were highly variable, but strongly correlated among the

controls ($R > 0.7$ for each pair of positive controls, Supplementary Fig. S2A). The read numbers

in the not equimolar positive control were strongly linked to the DNA template concentrations of

the PCRs (Pearson correlation coefficient $R = 0.97$, Supplementary Fig. S2B). Only

Leptodactylus vastus was recovered in the field positive control (water from an aquarium with *L.*

vastus tadpoles).

222   Sample preservation / extraction method C (0.2 µm nylon filter, dried in the field, DNA

extracted with DNeasy Blood & Tissue Kit) appeared to perform better (p_1=0.327, 95% C.I. =

[0.285; 0.372] than methods A (p_1=0.278 [0.242; 0.317]) and B (p_1=0.243 [0.209; 0.281], both 2

µm glass-fiber filters, A - preserved in CTAB and DNA extracted using CTAB protocol, and B -

dried in the field, DNA extracted using CTAB protocol, Supplementary Fig. S3). Method A had

larger detection probability than method B, but confidence intervals were largely overlapping.

The three methods were equivalent with respect to false positive probabilities (Supplementary

Fig. S3).

230   Regarding the ecological signal in the eDNA data, replicate samples of each pond

grouped relatively tightly on the latent variable model ordination (Fig. 3a). The pond identity was

a statistically significant predictor of frog communities in the five ponds (ANOVA, df = 6, dev =

534.99, p < 0.01). This is reflected in the 95% confidence intervals of the group centroids on the

ordination which clearly separates all ponds except T1 and T3 (Fig. 3a). The ordination of the

eDNA pond centroids closely corresponds with the ordination of observations from the five

ponds with VAES (Procrustes permutation test, R = 0.8, p = 0.03, Fig. 3b).

237   The cost model of VAES and eDNA showed that the starting costs (i.e. with few

sampling sites) for VAES are relatively low, but these costs rapidly increase until the taxonomic

expert becomes familiar with the regional frog fauna (Fig. 4). The VAES price is dependent on

the species richness: first, the VAES observer needs to record each species on the field, and

then the taxonomic expert needs to listen to each recording (Fig. 4). eDNA metabarcoding has a

relatively high entry price since consumables and sequencing are costly, regardless of the

number of sites. eDNA survey prices are then a linear function of the number of sampling sites,

and an increase in the site numbers simply adds to sampling and consumable costs, but does

not influence neither the time spent in the laboratory, nor the sequencing costs (Fig. 4).

Discussion

The one-time eDNA survey detected 25 species in the studied ponds, including formerly cryptic species that were only recently described by integrative taxonomy. This is on par with the one-time VAES (Fig. 2a), which confirmed the presence of (i) almost all species that were in contact with water during the survey time (Fig. 2b), (ii) all species that had tadpoles in the ponds (Supplementary Table S1), (iii) more than half of all 45 species ever recorded from the area (Fig. 2c), and (iv) about 65% of the 35 species ever recorded in the surveyed 5 ponds (Fig. 2d). The eDNA dataset recorded clear differences among the frog communities in the surveyed ponds, a result also in accordance with the VAES data (Fig. 3). The comparison of VAES and eDNA cost models show that eDNA biodiversity surveys may be a cost-efficient alternative to VAES in species-rich areas, but not necessarily in areas with low species numbers. Recommendations and technical remarks are presented in the Supplementary Materials.

Sensitivity of eDNA in diverse tropical frog communities

The comparison of the three filtration-preservation approaches shows that filters can be feasibly dried in the field and sent to a lab. We could not evaluate whether detection probability differs between larger pore-size GFF and small pore-size nylon filters since DNA was extracted with different methods from these. Currently only few comparative studies exist regarding the preservation of eDNA filtrates on filters: Hinlo *et al.* (2017) showed that the simple refrigeration of filters may be preferred to frozen storage. Here we show that filters can simply be dried in the field: this simplifies transportation since no special precautions are needed, unlike for liquid handling.

During the one-time surveys, six species were detected only with eDNA, but not with VAES (Fig. 2a). Four of these species (*Dermatonotus muelleri*, *Leptodactylus elenae*, *Osteocephalus taurinus*, *Rhinella schneideri*) are quite common in the area. The detection of

these four species by eDNA but not by VAES may result from the low abundance of adults or

tadpoles in the ponds, or a lack in acoustic activity. In some cases eDNA detections are likely

based on tadpoles only. For example, eDNA detected *L. syphax* from pond T2, although the

species was only heard calling from a crevice of a rocky outcrop, its usual habitat (de Sá *et al.*

2014), approximately 30 m away from T2. The remaining two species detected only with eDNA

are not known to occur in the area, thus they are candidates for false assignments. Both of

these were assigned with sequence data from EMBL (*L. latinasus*: KM091595, *L. laevis*:

AY843696). *Lysapsus laevis* would be the first record of this genus in the study area, but the

identification of *Lysapsus* populations only using short 16S eDNA sequences is questionable,

especially when considering the unclear taxonomy of the group in Bolivia. Nevertheless, all of

the four known species of *Lysapsus* (*L. boliviana*, *L. caraya*, *L. laevis*, *L. limellum*) may occur at

the site (De la Riva *et al.* 2000; Lavilla *et al.* 2004; Reichle 2004a; b; Angulo 2008; Jansen *et al.*

2011; Frost 2016) since they can easily be overlooked due to their small size and rather

inconspicuous advertisement call. *Leptodactylus latinasus* is another species possibly occurring

in the area but was so far not recorded (de Sá *et al.* 2014) and we cannot exclude the possibility

that the corresponding sequence variant is actually an erroneous variant of one of the six local

Leptodactylus species.

The species from this study either have no sufficient distributional data for an IUCN

evaluation, or they are in the "least concern" category (Supplementary Table S1). Twelve of the

31 species are not assessed by IUCN, because (a) the species were described or removed

from synonymy only recently (*Dendropsophus salli*: Jungfer *et al.* 2010; *Scinax madeirae*:

Brusquetti *et al.* 2014; *Dendropsophus arndti*: Caminer *et al.* 2017), or (b) the assessment of

their taxonomic status is still pending (e.g. various lineages of candidate species *sensu* Jansen

et al. 2011). Nine species with uncertain taxonomic status were classified as distinct

genealogical lineages by Jansen *et al.* (2011). These may represent new species with

considerably smaller distributions. Twelve further species lack information on actual distribution,

although only one species is classified as "data deficient" (*Pithecopus azureus*). Nineteen

species are classified as "least concern". These classifications may become obsolete given

rapid changes in taxonomy and updates on distributions: cryptic species-level diversity seems to

be high in frogs of South America (Fouquet *et al.* 2007; Gehara *et al.* 2013). In summary, very

little is known about actual distributions of Neotropical frogs due to sparse sampling and

unreliable taxonomy. eDNA might help to complete distribution data of amphibians (including

cryptic lineages), and this data is essential for reliable IUCN assessments.

eDNA metabarcoding is strongly facilitated by dedicated sequence databases: we had a

database that contained all 45 species that were ever recorded in the area. This database was

essential for both initial primer development, and taxonomic assignments: indeed, both eDNA-

recorded taxa that were not present on the complete local species list of the present study (Fig.

2, Supplementary Table S2) were identified in the EMBL-based assignment that followed the

assignment with the custom local 16S database. The importance of sequence assignment

databases is long recognized, with considerable efforts underway to establish them

(Ratnasingham & Hebert 2007; Coissac *et al.* 2016).

Six species were detected only with VAES, but not by eDNA (Fig. 2a, Supplementary

Table S1). None of these species are strictly bound to ponds in the life stages occurring during

our sampling: two hylids, *Dendropsophus arndti* and *D. salli*, usually call from plants on the pond

shores (Schulze *et al.* 2009) and have only sporadic contact with water. *Leptodactylus fuscus*

and *Pseudopaludicola* sp. also do not enter the water but usually call from nearby muddy

grounds or grasslands. We did not detect tadpoles of these species in the ponds: the tadpoles

of these species develop during the rainy season which triggers reproduction, but our sampling

slightly preceded the rainy season. Some of the other undetected species have terrestrial or

quasi-terrestrial life histories: *L. fuscus* deposits eggs within foam nests in underground burrows

at some distance from ponds and these are washed into nearby water bodies by floods that

follow heavy rains (Heyer 1978; Lucas *et al.* 2008). If we consider the specific life histories, the

one-time eDNA survey only missed a single species (*Sphaenorhynchus lacteus*) that was found

calling from the water during the VAES (Fig. 2b, Supplementary Table S1). The hylid *Hypsiboas*

geographicus was likely missed as a result of PCR bias: we could never recover it from positive

controls when DNA from other species was also present (Supplementary Fig. S2), although

single-template PCR reactions worked. Nonetheless, *H. geographicus* was also not detected by

VAES, so it is possible that the species was not present during the survey. Regarding the whole

local species pool (45 species in 10 years), life history also explains why at least some species

(N=10) may not be detectable by eDNA: these species reproduce outside of water (e.g.

Leptodactylus mystacinus), or are forest dwellers (e.g. *Leptodactylus* cf. *didymus*, Fig. 2d,

Supplementary Table S2). In conclusion, with a single sampling eDNA detected about 65% of

the detectable species from the area (23 out of 35).

Phenology is long considered important in encounter-based surveys of vertebrates

(Wilson & Bart 1985; Møller 2010) and it was recently shown that life history influences insect

detections with eDNA metabarcoding in temperate freshwaters (Bista *et al.* 2017). Our sampling

was done at the beginning of the rainy season when only few species reproduced in the ponds.

Repeating the sampling at during the rainy season may have potentially increased species

detections by eDNA. The results show the importance of life history in the metabarcoding-based

survey of tropical frogs and emphasize that sampling at specific or even multiple timepoints may

be essential for more complete species pools also with eDNA (see Recommendations and

technical remarks in the Supplementary Materials).

Ecological signal

There was considerable variation in the species recorded with the spatial replicates of

the eDNA samples, and this underlines that eDNA sampling should be replicated for a good

representation of community composition. These samples may be pooled before DNA extraction

to optimize costs if site-level variation is not of interest. We recorded similar ecological signal in

the two datasets (Fig 3). The eDNA data did not distinguish the communities from ponds T1 and

T3, and these communities also grouped closely in the VAES results. We randomized the

samples before DNA extraction (Bálint *et al.* 2017a) so the signal is likely of biological origin and

not a result of laboratory batch effects. Although the comparison of the ecological signal may be

confounded by the different sample sizes (with the eDNA ordination based on many spatial

replicates and the VAES ordination on a single observation event per pond), the

correspondence of the results is still striking. Similar results were found in insect (Ji *et al.* 2013;

Elbrecht *et al.* 2017) and plant communities (Niemeyer *et al.* 2017) where eDNA-based and

morphology-based identifications result in similar conclusions about community structure,

despite markedly different species lists. Our results provide further evidence that eDNA-based

biodiversity surveys are highly sensitive to differences among ecological communities. These

inferences are comparable to those derived with encounter-based observations and are

informative about processes that underlay community assembly.

Cost efficiency of biodiversity surveys with eDNA metabarcoding

Although central to deciding on a method for biodiversity surveys, cost comparisons are

not straightforward since they need to be based on expert knowledge both in VAES and eDNA.

Cost comparisons were performed for single species eDNA detection (Huver *et al.* 2015; Davy

et al. 2015; Smart *et al.* 2016), but we are not aware of frameworks suitable for eDNA

metabarcoding. Here the VAES cost estimation is informed by over a decade of field and

integrative taxonomic work with tropical frogs (Jansen *et al.* 2007, 2016; Schulze *et al.* 2009;

Brusquetti *et al.* 2014), and the eDNA part is informed by multiple years of eDNA biodiversity

surveys (Bálint *et al.* 2016, 2017a; Vörös *et al.* 2017). Such comparisons are urgently needed

due to stakeholder demands in eDNA (governmental agencies, conservation NGOs, fisheries,

etc.).

We defined two species richness scenarios for the cost comparisons (Fig. 4) since

experience shows that the time and resources necessary for VAES depend on the local

biodiversity. VAES costs become high in regions with high frog richness since they are a

function of both the number of sites and the number of species. Biodiversity surveys with eDNA

are not necessarily cheaper in low-richness regions since entry costs for the eDNA work are

high: lab consumables and sequencing are costly. However, eDNA survey costs are not

dependent on the local biodiversity since metabarcoding can consider thousands of species

simultaneously in a sample (Taberlet *et al.* 2012b). Consequently, eDNA costs are function of

sample numbers, which influence the collection time spent on the field and consumables. eDNA

metabarcoding operations are easily scaled up in a sense that hundreds of samples can be

simultaneously processed (Ficetola *et al.* 2015).

Several aspects of our cost models are contentious. One issue is whether the relatively

untrained VAES observers, or taxonomic experts perform the fieldwork, since taxonomic experts

may identify many of the species immediately on the field. Currently, most surveys of high-

diversity areas are directly done by experts interested in the local fauna, but we argue that this

will not work for continental - global biodiversity surveys simply because there are not enough

taxonomic experts (Buyck 1999; Haas & Häuser 2003). We also did not consider a scenario

when VAES surveys are performed with automated recording devices (ARDs), and sounds are

automatically identified by algorithms (see Recommendations and technical remarks in the

Supplementary Materials). The sound complexity in tropical environments currently prohibits the

use of automated sound identifications (Campos-Cerqueira & Aide 2016). It is also difficult to

compare fundamental infrastructure and training costs (i.e. the establishment of an eDNA labor

VS the training of taxonomic experts). Discussions about cost models are timely since they will

play major roles in devising continental-global biodiversity surveys, much needed today to

monitor global biodiversity targets (Tittensor *et al.* 2014).

Conclusions

eDNA seems to be suitable to standardized biodiversity surveys of frogs even in

species-rich areas, but it may be overly costly for smaller studies in low-richness regions.

Differences between eDNA and traditional surveys seem to result largely from different

observational biases, but a consideration of life histories promises to improve correspondence.

The eDNA data is certainly suitable to characterize not only community composition, but also

the ecological signals of factors that shape communities: this gives an unprecedented

opportunity to incorporate eDNA as a standard toolkit for community ecology and macroecology.

The quality of the ecological signal in eDNA-based biodiversity data foresees global biodiversity

surveys and monitoring that will support both biodiversity research, and informed decisions on

sustainable use of biological diversity.

Acknowledgements

The authors thank Jenny Wertheimer for technical assistance during eDNA extraction, José Ribeiro for a helpful discussion on passive acoustic monitoring, and Simon Vitecek for helpfuls suggestions on the manuscript. MJ, AS and JLA thank the owners of Hacienda San Sebastián (family Werding) for their invitation to conduct herpetological surveys on their properties and for logistic support. MJ was supported by the Erika and Walter Datz-Stiftung, Bad Homburg v. d. H., equipment was sponsored by Globetrotter, Frankfurt. This research presents an outcome of the Centre for Translational Biodiversity Genomics (LOEWE-TBG) and was supported by the research funding programme "LOEWE – Landes-Offensive zur Entwicklung Wissenschaftlich-ökonomischer Exzellenz" of Hesse's Ministry of Higher Education, Research, and the Arts.

References

Angulo A (2008) *Lysapsus boliviana. The IUCN Red List of Threatened Species.*

Bailey LL, MacKenzie DI, Nichols JD (2014) Advances and applications of occupancy models. *Methods in ecology and evolution / British Ecological Society*, **5**, 1269–1279.

Bálint M, Bahram M, Eren AM *et al.* (2016) Millions of reads, thousands of taxa: microbial community structure and associations analyzed via marker genes. *FEMS microbiology reviews*, **40**, 686–700.

Bálint M, Domisch S, Engelhardt CHM *et al.* (2011) Cryptic biodiversity loss linked to global climate change. *Nature climate change*, **1**, 313–318.

Bálint M, Márton O, Schatz M, Duering R-A, Grossart H-P (2017a) Proper experimental design requires randomization/balancing of molecular ecology experiments. *bioRxiv*. Doi: 10.1101/109280. https://doi.org/10.1101/109280

Bálint M, Nowak C, Márton O *et al.* (2017b) Input files for the comparison of visual-audio encounter surveys and eDNA metabarcoding of species-rich tropical frog communities. https://doi.org/10.6084/m9.figshare.5099842

Barnosky AD, Matzke N, Tomiya S *et al.* (2011) Has the Earth's sixth mass extinction already arrived? *Nature*, **471**, 51–57.

Bishop PJ, Angulo A, Lewis JP (2012) The amphibian extinction crisis-what will it take to put the action into the amphibian conservation action plan? *SAPI EN. S. Surveys.*

Bista I, Carvalho GR, Walsh K *et al.* (2017) Annual time-series analysis of aqueous eDNA reveals ecologically relevant dynamics of lake ecosystem biodiversity. *Nature communications*, **8**, 14087.

Boyer F, Mercier C, Bonin A *et al.* (2016) obitools: a unix-inspired software package for DNA metabarcoding. *Molecular ecology resources*, **16**, 176–182.

Brusquetti F, Jansen M, Barrio-Amorós et al. (2014) Taxonomic review of *Scinax*

449  *fuscomarginatus* (Lutz, 1925) and related species (Anura; Hylidae). *Zoological journal of*

450   *the Linnean Society*.

451  Bush A, Sollmann R, Wilting A *et al.* (2017) Connecting Earth observation to high-throughput

452   biodiversity data. *Nature Ecology & Evolution*.

453  Butchart SHM, Walpole M, Collen B *et al.* (2010) Global biodiversity: indicators of recent

454   declines. *Science*, **328**, 1164–1168.

455  Buyck B (1999) Taxonomists are an endangered species in Europe. *Nature*, **401**, 321.

456  Caminer MA, Milá B, Jansen M *et al.* (2017) Systematics of the *Dendropsophus leucophyllatus*

457   species complex (Anura: Hylidae): Cryptic diversity and the description of two new species.

458   *PloS one*, **12**, e0171785.

459  Campos-Cerqueira M, Aide TM (2016) Improving distribution data of threatened species by

460   combining acoustic monitoring and occupancy modelling. *Methods in ecology and evolution*

461   */ British Ecological Society*, **7**, 1340–1348.

462  Castro AAJF, A A J, Martins FR, Tamashiro JY, Shepherd GJ (1999) How rich is the flora of

463   Brazilian Cerrados? *Annals of the Missouri Botanical Garden. Missouri Botanical Garden*,

464   **86**, 192.

465  Clarke DA, York PH, Rasheed MA, Northfield TD (2017) Does biodiversity–ecosystem function

466   literature neglect tropical ecosystems? *Trends in ecology & evolution*, **32**, 320–323.

467  Coissac E, Hollingsworth PM, Lavergne S, Taberlet P (2016) From barcodes to genomes:

468   extending the concept of DNA barcoding. *Molecular ecology*, **25**, 1423–1428.

469  Costello MJ, May RM, Stork NE (2013) Can we name Earth's species before they go extinct?

470   *Science*, **339**, 413–416.

471  Cristescu ME (2014) From barcoding single individuals to metabarcoding biological

472   communities: towards an integrative approach to the study of global biodiversity. *Trends in*

473   *ecology & evolution*, **29**, 566–571.

474  Davy CM, Kidd AG, Wilson CC (2015) Development and validation of environmental DNA

(eDNA) markers for detection of freshwater turtles. *PloS one*, **10**, e0130965.

De la Riva I, Köhler J, Lötters S, Reichle S (2000) Ten years of research on Bolivian

amphibians: updated checklist, distribution, taxonomic problems, literature and

iconography. *Revista espanola de herpetologia*, **14**, 19–164.

de Sá RO, Grant T, Camargo A *et al.* (2014) Systematics of the neotropical genus

Leptodactylus Fitzinger, 1826 (Anura: Leptodactylidae): phylogeny, the relevance of non-

molecular evidence, and species accounts. *South american journal of herpetology /*

Sociedade Brasileira de Herpetologia, **9**, S1–S100.

Elbrecht V, Vamos EE, Meissner K, Aroviita J, Leese F (2017) Assessing strengths and

weaknesses of DNA metabarcoding-based macroinvertebrate identification for routine

stream monitoring. *Methods in ecology and evolution / British Ecological Society*.

Ferrão M, Colatreli O, de Fraga R *et al.* (2016) High species richness of *Scinax* treefrogs

(Hylidae) in a threatened Amazonian landscape revealed by an integrative approach. *PloS*

one, **11**, e0165679.

Ficetola GF, Pansu J, Bonin A *et al.* (2015) Replication levels, false presences and the

estimation of the presence/absence from eDNA metabarcoding data. *Molecular ecology*

resources, **15**, 543–556.

Fiske I, Chandler R, Others (2011) unmarked: An R package for fitting hierarchical models of

wildlife occurrence and abundance. *Journal of statistical software*, **43**, 1–23.

Fouquet A, Gilles A, Vences M *et al.* (2007) Underestimation of species richness in neotropical

frogs revealed by mtDNA analyses. *PloS one*, **2**, e1109.

Frost DR (2016) Amphibian Species of the World: an Online Reference. Version 6.0.

Gehara M, Canedo C, Haddad CFB, Vences M (2013) From widespread to microendemic:

molecular and acoustic analyses show that *Ischnocnema guentheri* (Amphibia:

Brachycephalidae) is endemic to Rio de Janeiro, Brazil. *Conservation genetics* , **14**, 973–

982.

Gehara M, Crawford AJ, Orrico VGD *et al.* (2014) High levels of diversity uncovered in a

widespread nominal taxon: continental phylogeography of the Neotropical tree frog

Dendropsophus minutus. PloS one, **9**, e103958.

Haas F, Häuser CL (2003) Taxonomists: an endangered species? In: *Success stories in*

implementation of the programmes of work on dry and sub-humid lands and the global

taxonomy initiative, p. 87. Montreal: Secretariat of the Convention on Biological Diversity

Henry P-Y, Lengyel S, Nowicki P *et al.* (2008) Integrating ongoing biodiversity monitoring:

potential benefits and methods. *Biodiversity and conservation*, **17**, 3357–3382.

Heyer WR (1978) *Systematics of the fuscus group of the frog genus* Leptodactylus *(Amphibia,*

Leptodactylidae). Natural History Museum of Los Angeles County.

Hinlo R, Gleeson D, Lintermans M, Furlan E (2017) Methods to maximise recovery of

environmental DNA from water samples. *PloS one*, **12**, e0179251.

Hui FKC, Taskinen S, Pledger S, Foster SD, Warton DI (2015) Model-based approaches to

unconstrained ordination. *Methods in ecology and evolution / British Ecological Society*, **6**,

399–411.

Huver JR, Koprivnikar J, Johnson PTJ, Whyard S (2015) Development and application of an

eDNA method to detect and quantify a pathogenic parasite in aquatic ecosystems.

Ecological applications: a publication of the Ecological Society of America, **25**, 991–1002.

Jansen M (2009) Measuring temporal variation in calling intensity of a frog chorus with a data

logging sound level meter: results from a pilot study in Bolivia. *Herpetology notes*, **2**, 143–

149.

Jansen M, Álvarez LG, Köhler G (2007) New species of *Hydrolaetare* (Anura, Leptodactylidae)

from Bolivia with some notes on Its natural history. *Journal of Herpetology*, **41**, 724–732.

Jansen M, Bloch R, Schulze A, Pfenninger M (2011) Integrative inventory of Bolivia's lowland

anurans reveals hidden diversity. *Zoologica scripta*.

Jansen M, Plath M, Brusquetti F, Ryan MJ (2016) Asymmetric frequency shift in advertisement

calls of sympatric frogs. *Amphibia-reptilia: publication of the Societas Europaea*

Herpetologica, **37**, 137–152.

Ji Y, Ashton L, Pedley SM *et al.* (2013) Reliable, verifiable and efficient monitoring of

biodiversity via metabarcoding. *Ecology letters*, **16**, 1245–1257.

Jungfer K-H, Reichle S, Piskurek O (2010) Description of a new cryptic southwestern

Amazonian species of leaf-gluing treefrog, genus *Dendropsophus* (Amphibia: Anura:

Hylidae). *Salamandra*, **46**, 204–213.

Lavilla E, Reichle S, Lajmanovich R, Faivovich, Julian (2004) *Lysapsus limellum. The IUCN Red*

List of Threatened Species.

Lees AC, Pimm SL (2015) Species, extinct before we know them? *Current biology: CB*, **25**,

R177–80.

Lopes CM, Sasso T, Valentini A *et al.* (2016) eDNA metabarcoding: a promising method for

anuran surveys in highly diverse tropical forests. *Molecular ecology resources.*

Lucas EM, Brasileiro CA, Oyamaguchi HM, Martins M (2008) The reproductive ecology of

Leptodactylus fuscus (Anura, Leptodactylidae): new data from natural temporary ponds in

the Brazilian Cerrado and a review throughout its distribution. *Journal of natural history*, **42**,

2305–2320.

MacKenzie DI, Nichols JD, Lachman GB *et al.* (2002) Estimating site occupancy rates when

detection probabilities are less than one. *Ecology*, **83**, 2248–2255.

Miles L, Newton AC, DeFries RS *et al.* (2006) A global overview of the conservation status of

tropical dry forests. *Journal of biogeography*, **33**, 491–505.

Miller DA, Nichols JD, McClintock BT *et al.* (2011) Improving occupancy estimation when two

types of observational error occur: non-detection and species misidentification. *Ecology*, **92**,

1422–1428.

Miya M, Sato Y, Fukunaga T *et al.* (2015) MiFish, a set of universal PCR primers for

metabarcoding environmental DNA from fishes: detection of more than 230 subtropical

marine species. *Royal Society open science*, **2**, 150088.

Møller AP (2010) When climate change affects where birds sing. *Behavioral ecology: official*

journal of the International Society for Behavioral Ecology, **22**, 212–217.

Niemeyer B, Epp LS, Stoof-Leichsenring KR, Pestryakova LA, Herzschuh U (2017) A

comparison of sedimentary DNA and pollen from lake sediments in recording vegetation

composition at the Siberian treeline. *Molecular ecology resources*.

Ortega-Andrade HM, Rojas-Soto OR, Valencia JH *et al.* (2015) Insights from integrative

systematics reveal cryptic diversity in *Pristimantis* frogs (Anura: Craugastoridae) from the

Upper Amazon Basin. *PloS one*, **10**, e0143392.

Pauls SU, Nowak C, Bálint M, Pfenninger M (2013) The impact of global climate change on

genetic diversity within populations and species. *Molecular ecology*, **22**, 925–946.

Pedersen MW, Overballe-Petersen S, Ermini L *et al.* (2015) Ancient and modern environmental

DNA. *Philosophical transactions of the Royal Society of London. Series B, Biological*

sciences, **370**, 20130383.

Peres-Neto PR, Jackson DA (2001) How well do multivariate data sets match? The advantages

of a pProcrustean superimposition approach over the Mantel test. *Oecologia*, **129**, 169–

178.

Petitot M, Manceau N, Geniez P, Besnard A (2014) Optimizing occupancy surveys by

maximizing detection probability: application to amphibian monitoring in the Mediterranean

region. *Ecology and evolution*, **4**, 3538–3549.

Pfenninger M, Schwenk K (2007) Cryptic animal species are homogeneously distributed among

taxa and biogeographical regions. *BMC evolutionary biology*, **7**, 121.

Pimm SL, Jenkins CN, Abell R *et al.* (2014) The biodiversity of species and their rates of

extinction, distribution, and protection. *Science*, **344**, 1246752.

Power MJ, Whitney BS, Mayle FE *et al.* (2016) Fire, climate and vegetation linkages in the

Bolivian Chiquitano seasonally dry tropical forest. *Philosophical transactions of the Royal*

Society of London. Series B, Biological sciences, **371**.

Ratnasingham S, Hebert PDN (2007) BARCODING: bold: The barcode of life data system

(http://www.barcodinglife.org). *Molecular ecology notes*, **7**, 355–364.

R Core Team (2017) *R: A Language and Environment for Statistical Computing*.

Reichle S (2004a) *Lysapsus laevis. The IUCN Red List of Threatened Species*.

Reichle S (2004b) *Lysapsus caraya. The IUCN Red List of Threatened Species*.

Schmeller DS, Böhm M, Arvanitidis C *et al.* (2017) Building capacity in biodiversity monitoring at

the global scale. *Biodiversity and conservation*, 1–26.

Schulze A, Jansen M, Köhler G (2009) Diversity and ecology of anuran communities in San

Sebastián (Chiquitano region, Bolivia). *Salamandra*, **45**, 75–90.

Schulze A, Jansen M, Köhler G (2015) Tadpole diversity of Bolivia's lowland anuran

communities: molecular identification, morphological characterisation, and ecological

assignment. *Zootaxa*, **4016**, 1–111.

Smart AS, Weeks AR, van Rooyen AR *et al.* (2016) Assessing the cost-efficiency of

environmental DNA sampling. *Methods in ecology and evolution / British Ecological*

Society, **7**, 1291–1298.

Strand DA, Jussila J, Johnsen SI *et al.* (2014) Detection of crayfish plague spores in large

freshwater systems. *The Journal of applied ecology*, **51**, 544–553.

Stroud JT, Feeley KJ (2017) Neglect of the Tropics Is Widespread in Ecology and Evolution: A

Comment on Clarke et al. *Trends in ecology & evolution*.

Stuart SN, Chanson JS, Cox NA *et al.* (2004) Status and trends of amphibian declines and

extinctions worldwide. *Science*, **306**, 1783–1786.

Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH (2012a) Environmental DNA. *Molecular*

ecology, **21**, 1789–1793.

Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E (2012b) Towards next-

generation biodiversity assessment using DNA metabarcoding. *Molecular ecology*, **21**,

2045–2050.

Thomsen PF, Kielgast J, Iversen LL *et al.* (2012) Detection of a diverse marine fish fauna using

environmental DNA from seawater samples. *PloS one*, **7**, e41732.

Thomsen PF, Willerslev E (2015) Environmental DNA – An emerging tool in conservation for

monitoring past and present biodiversity. *Biological conservation*, **183**, 4–18.

Tittensor DP, Walpole M, Hill SLL *et al.* (2014) A mid-term analysis of progress toward

international biodiversity targets. *Science*, **346**, 241–244.

UNEP (2016) The strategic plan for biodiversity 2011-2020, the Aichi biodiversity targets and

NBSAPs. In: *Sourcebook of Opportunities for Enhancing Cooperation among the*

Biodiversity-Related Conventions at National and Regional Levels, pp. 108–135.

Valentini A, Taberlet P, Miaud C *et al.* (2016) Next-generation monitoring of aquatic biodiversity

using environmental DNA metabarcoding. *Molecular ecology*, **25**, 929–942.

Vörös J, Márton O, Schmidt BR, Gál JT, Jelić D (2017) Surveying Europe's only cave-dwelling

chordate species (*Proteus anguinus*) using environmental DNA. *PloS one*, **12**, e0170945.

Wang Y, Naumann U, Wright ST, Warton DI (2012) mvabund- an Rpackage for model-based

analysis of multivariate abundance data. *Methods in ecology and evolution / British*

Ecological Society, **3**, 471–474.

Whittaker K, Koo MS, Wake DB, Vredenburg VT (2013) Global declines of amphibians. In:

Encyclopedia of Biodiversity (Second Edition) (ed Levin SA), pp. 691–699. Academic

Press, Waltham.

Wilson DM, Bart J (1985) Reliability of singing bird surveys: effects of song phenology during

the breeding season. *The Condor*, **87**, 69–73.

Wittwer C, Stoll S, Cocchiararo B, Strand D, Vrålstad T, Nowak C, Thines M (2017) eDNA-

based crayfish plague monitoring is superior to conventional trap-based assessments in

year-round detection probability. *Hydrobiologia* (pending minor revision).

Zimmerman BL (1994) Audio strip transects. In: *Measuring and monitoring biological*

biodiversity. Standard methods for amphibians (eds Heyer WR, Donnelly MA, McDiarmid

RW, Hayek LC, Foster MS), pp. 99–108. Smithsonian Institution Press, Washington DC.

Data accessibility

Raw sequence data was deposited in ENA as PRJEB22113. Cost model parameters

and calculations, and input data for statistical analyses are accessible through FigShare (Bálint

et al. 2017b) (https://figshare.com/articles/_/5099842). R-based bioinformatic filtering and

statistical analyses are accessible through GitHub: https://github.com/MikiBalint/amphibian-

eDNA.

Author contributions

645   M.B., M.J., C.N., C.W. designed research; J.L.A., M.B., M.J., O.M., A.S., C.W.,

646 performed research; B.C., T.C., C.N., S.U.P. contributed reagents or analytic tools; M.B., M.J.

647 analysed data; M.B., T.C., M.J., O.M., C.N., S.U.P., wrote the manuscript.

Figures

Fig. 1.

Sampled ponds, sampling replication, and filtration - filter preservation strategy. GFF -

glass-fiber filter, 2 μm; nylon - nylon filter, 2 μm. Each sampling replicate (three per pond,

marked with small circles) were processed with each three filtration strategies. Replicated

samples of pond T5 were three times processed with each filtration strategy.

Fig. 2.

Comparison of species lists generated by eDNA (blue, n = 25) and A) visual and audio encounter surveys (VAES, yellow, n = 25), B) tadpole survey (TS) plus reduced VAES - only species with high eDNA detectability in ponds (see text for details; grey, n=20), C) complete local species pool (red, n = 45), and D) species pool of studied ponds based on long-term monitoring (green, n = 35).

Fig. 3.

Comparison of ecological signal between visual and audio encounter surveys (VAES) and

eDNA metabarcoding surveys. A - Latent variable model ordination of the pond samples

according to the read numbers of species. Water samples from each pond were taken at three

locations. For ponds T1-T4, three replicate samples at each location were filtered through two

GFF (A,B), and one nylon filter (C), resulting in nine samples per pond. For pond T5, each

filtration was replicated three times. Not all samples contained reads after the bioinformatic

quality filtering and these samples are not shown on the ordination (see e.g. T3 - red). The

ellipses represent 95% confidence intervals for the standard errors of the pond centroids

(marked with +). B - Non-metric multidimensional scaling plot of the five ponds (Jaccard

distance), according to VAES species presence-absences. The arrows represent the Procrustes

rotation of the VAES pond ordination and they target the group centroids of the latent variable

model ordination.

Fig. 4.

Cost comparisons between visual and audio encounter surveys (VAES) and eDNA

metabarcoding surveys. A - Schematic overview of the visual and audio encounter survey

(VAES) and eDNA cost models. Model parameters and calculations are accessible through

FigShare (Balint_et_al_survey_cost_calculations.xlsx, doi: 10.6084/m9.figshare.5099842). B -

Cost comparisons of frog diversity surveys with eDNA, and two VAES scenarios (a low local

species richness scenario - 10 species per site, and a medium-high species richness scenario -

20 species per site). eDNA surveys have high initial costs: laboratory and sequencing costs are

considerable, regardless of the numbers of sites and species. A consequence is that eDNA

surveys are not sensitive to increases in species numbers. This is in contrast with the VAES,

where the costs are a direct function of species richness.