Horizontal gene cluster transfer increased hallucinogenic mushroom diversity

Hannah T. Reynolds[1,2]
Vinod Vijayakumar[1]
Emile Gluck-Thaler[1]
Hailee Brynn Korotkin[3]
Patrick Brandon Matheny[3]
Jason Christopher Slot[1]

[1]Department of Plant Pathology, The Ohio State University
[2]Department of Biological & Environmental Sciences, Western Connecticut State University
[3] Ecology & Evolutionary Biology, University of Tennessee, Knoxville

Corresponding Author: Jason C. Slot, slot.1@osu.edu

Abstract:

Secondary metabolites are heterogeneous natural products that often mediate

interactions between species. The tryptophan-derived secondary metabolite,

psilocin, is a serotonin receptor agonist that induces altered states of consciousness.

A phylogenetically disjunct group of mushroom-forming fungi in the Agaricales

produce the psilocin prodrug, psilocybin. Spotty phylogenetic distributions of fungal

compounds are sometimes explained by horizontal transfer of metabolic gene

clusters among unrelated fungi with overlapping niches. We report the discovery of

a psilocybin gene cluster in three hallucinogenic mushroom genomes, and evidence

for its horizontal transfer between fungal lineages. Patterns of gene distribution and

transmission suggest that psilocybin provides a fitness advantage in the dung and

late wood-decay niches, which may be reservoirs of fungal indole-based metabolites

that alter behavior of mycophagous and wood-eating invertebrates. These

hallucinogenic mushroom genomes will serve as models in neurochemical ecology,

advancing the prospecting and synthetic biology of novel neuropharmaceuticals.

Secondary metabolites (SMs) are small molecules that are widely employed in

defense, competition, and signaling among organisms (Raguso et al. 2015). Due to

their physiological activities, SMs have been adopted by both ancient and modern

human societies as medical, spiritual, or recreational drugs. Psilocin is a

psychoactive agonist of the serotonin (5-hydroxytryptamine, **5-HT**) -2a receptor

(Halberstadt and Geyer 2011) and is produced as the phosphorylated prodrug

psilocybin by a restricted number of distantly related mushroom forming families of

the Agaricales (Bolbitiaceae, Inocybaceae, Hymenogastraceae, Pluteaceae) (Allen

2010 May 19; Dinis-Oliveira 2017). Hallucinogenic mushrooms have a long history

of religious use, particularly in Mesoamerica, and were a catalyst of cultural

revolution in the West in the mid 20th century (Nyberg 1992; Letcher 2006).

Psilocybin was structurally described and synthesized in 1958 by Albert Hoffman

(Hofmann et al. 1958), and a biosynthetic pathway (Fig. 1B) was later proposed

based on the transformation of precursor molecules by *Psilocybe cubensis* (Agurell

and Nilsson 1968). However, prohibition since the 1970s (21 U.S. Code § 812 -

Schedules of controlled substances) has limited advances in psilocybin genetics,

ecology, and evolution. There has been a recent resurgence of the hallucinogen

research in the clinical setting. Brain state imaging studies of psilocin exposure have

identified changes in neural activity and interconnectivity that underlie subjective

experiences, and therapeutic trials have investigated psilocybin's potential for

treating major depression and addictive disorders (Griffiths et al. 2011; Carhart-

Harris et al. 2012; Petri et al. 2014; Carhart-Harris et al. 2016; Johnson et al. 2017).

While the ecological roles of psilocybin, like most SMs, remain unknown, psilocin's

mechanism of action suggests metazoans may be its principal targets.

A common feature of fungal SM biosynthesis is the organization of most or all

required anabolic, transport, and regulatory elements in gene clusters (GCs). GCs

are often discontinuously distributed among fungal taxa, partly due to horizontal

transfer (HT) among species with overlapping ecological niches (Gluck-Thaler and

Slot 2015). The sparse phylogenetic distribution of psilocybin, coupled with the

requirement for multiple enzymatic steps for its biosynthesis (tryptophan-

decarboxylation, N-methylation, indole-4-hydroxylation, and O-phosphorylation),

suggest the psilocybin pathway might have dispersed via horizontal GC transfer, and

therefore the genetic mechanism for psilocybin biosynthesis might be identified in

searches for GCs with a common phylogenetic history and distribution among

psilocybin producing (PS+) mushrooms. The pattern of HT events may further

suggest ecological pressures that have driven the pathway's persistence (Baquero

2004).

We identified candidate psilocybin genes by sequencing three diverse PS+

mushroom monokaryon genomes -- *Psilocybe cyanescens*, *Panaeolus* (=*Copelandia*)

cyanescens, and *Gymnopilus dilepis* (Table 1), and comparing them to three related

mushrooms not known to produce psilocybin (PS-): *Galerina marginata*, *Gymnopilus*

chrysopellus, and *Hypholoma sublateritium*. Of 37 gene homolog groups (HGs)

consistent with a PS+ distribution among these taxa, only five were clustered, all in

PS+ genomes (Fig. S1-pipeline and HGs). We retroactively designated *Gy.*

chrysopellus, potentially PS+ because it possesses a cluster identical to *Gy. dilepis*,

which is not a surprising oversight given inconsistent identifications, and

geographical variation among *Gymnopilus* spp. phenotypes. Predicted functions of

these five genes were also consistent with psilocybin biosynthesis and metabolite

transport, and were putatively designated tryptophan decarboxylase (TDC),

tryptamine N-methyltransferase (TMT), dimethyltryptamine-4-hydroxylase (D4H),

psilocin phosphotransferase (PPT), and psilocybin transporter (PST). As SM GCs are

infrequently identified in Basidiomycota compared with Ascomycota, this is a

notable discovery (Quin et al. 2014).

To confirm GC function, we profiled the enzymology of heterologously expressed

TDC and PPT, and assayed by LC-MS/MS analyses. We determined that TDC, the first

committed step in the reaction and the only one not producing a drug-scheduled

compound, has specific decarboxylase activity on tryptophan. TDC reactions

produced tryptamine, identified at the characteristic m/z 144.1 [M+H]+, (Fig. S2,

Supplemental data1). TDC did not decarboxylate phenylalanine, tyrosine or 5-

hydroxy-L-tryptophan (5-HTP) under the same conditions. We note that TDC is

similar to type II phosphatidylserine decarboxylases (PSDs), but has no significant

sequence similarity with a pyridoxal-5'-phosphate-dependent decarboxylase

recently characterized in *Ceriporiopsis subvermispora* as specific for L-tryptophan

and 5-HTP (Kalb et al. 2016). A unique GGSS sequence in a conserved C-terminal

motif (Fig. S3), suggests tryptophan decarboxylation is a previously unknown

derived function among PSDs (Wriessnegger et al. 2009; Choi et al. 2015). We

detected no activity of PPT on 5-HT or 4-Hydroxyindole (4-HI) as alternatives to the

psilocin substrate, possibly due to requirements for the 4-hydroxyl and the

methylated amine groups of psilocin, but further characterization of PPT and other

enzymes was prevented by the regulatory status of substrates and products.

Phylogenetic analyses of PS homologs from a local database of 618 fungal

proteomes yielded congruent gene tree topologies with respect to PS[+] taxa, and

clades of clustered PS genes from all gene trees excluded the PS[-] taxa in the

database, suggesting the clustered genes are coordinately-inherited (Fig. 1, Fig. S4

A-F).The gene trees also suggest HT of the cluster from *Psilocybe* to *Panaeolus* and

HT of most PS genes between Atheliaceae and Agaricaceae when compared to a

phylogenomic tree of related Agaricales (Fig. 1). The direction of the latter HT is

ambiguous, and not strongly supported by all five genes. Analyses with TDC and PPT

amplicon sequences retrieved by degenerate PCR of unsequenced *Psilocybe* and

Conocybe genomes (Supplemental data1) suggest the dung fungus *Ps. cubensis*

vertically inherited the cluster, and *Pa. cyanescens* acquired the cluster from

Psilocybe sp., and possibly from a dung-associated lineage. Alternative hypotheses of

vertical inheritance in these lineages were rejected; exclusion of *Pa. cyanescens* and

C. cyanopus (AU test, p = 0.004) or *Pa. cyanescens* alone (p = 0.036) were

significantly worse constrained topologies (Supplemental data1). Furthermore, a

TDC gene tree-species tree reconciliation model allowing duplication, HT, and loss

(6 events: D=1, HT=3, L=2) is more parsimonious than a model that only allows

duplication and loss (28 events: D=3, L=25)(Fig. S5). PS gene orthologs were not

detected in *Ps. fuscofulva*, a PS[-] species representing an early branch in *Psilocybe*

diversification (Borovička et al. 2015). Conservation of synteny flanking the *Ps.*

cyanescens PS cluster (Fig. 1) suggests it may have been recently acquired in

Psilocybe as well, or lost as a unit in close relatives. A genome wide scan did not

identify additional HT genes or clusters between *Psilocybe* and *Panaeolus* (Fig. S6).

HT is comparatively rare in the Basidiomycetes, suggesting the transfer of the PS

cluster may have special significance (Wisecaver et al. 2014), and is to our

knowledge, the first report of HT of a SM GC between lineages of mushroom-forming

fungi (Agaricomycotina).

Recent studies suggest that ecology can select for both genome content (Ma et al.

2010; de Jonge et al. 2013) and organization in eukaryotes through both vertical and

horizontal patterns of inheritance (Holliday et al. 2015; Kakioka et al. 2015).

Ordination of 10,998 HGs identified two principal components (PCs) that describe

22% of the variation in gene content among 16 Agaricales genomes (Fig. 2B).

Discrimination of genome composition along PC1 appears to reflect phylogenetic

differences, while discrimination along PC2 parallels ecological differences between

plant mutualists and other fungi. However, PC2 does not discriminate between dung

and wood-decay fungi. The functions of HGs most associated with each PC are

consistent with this interpretation. All eight metabolism-related processes in the

COG classification system are overrepresented in PC2, but only one is

overrepresented in PC1 HGs (Supplemental data1). The grouping of several

divergent lineages of wood and dung decay fungi to the exclusion of close

ectomycorrhizal relatives along PC2 may reflect similar selective pressures in the

decayed wood and dung environments, from recalcitrant plant polymers like lignin,

and invertebrate predation (Rouland-Lefèvre 2000). However, a small number of

HGs exclusive to either wood or dung associated fungi (Fig. S7, Supplemental data1)

are consistent with ecological specialization within each guild. Wood-specific genes

include functions in lignin degradation (e.g., peroxidase, isoamyl alcohol oxidase)

and carbohydrate transport, while dung-specific genes have functions in bacterial

cell wall degradation (e.g., lysozyme), hemicellulose degradation (e.g., Endo-1,4-

beta-xylanase, Alpha-L-arabinofuranosidase), and inorganic phosphate transport.

Niche-specific genes are largely consistent with vertical inheritance; however,

analyses support HT of a single ferric-reductase-like gene (pfam01794, pfam00175)

likely involved in iron uptake, to Coprinopsis and Panaeolus from dung-associated

Ascomycota (Supplemental data).

In addition to similar ecological pressures, similar genome content among wood and

dung-decaying fungi may also reflect the ecological diversification of

Agaricomycetes that accompanied major geological transformations (Fig. 2B). For

example, the emergence of true wood opened a massive saprotrophy niche space in

the upper Devonian (380 Mya), in which the Agaricomycetes diversified with the aid

of key enzymatic innovations (Floudas et al. 2012). The subsequent radiation of

herbivorous megafauna during the Eocene approximately 50 MYA (MacFadden

2000) and the spread of grasslands 40 MYA (Retallack 2001) expanded the

mammalian dung niche space in which invertebrates and fungi competed. These

changes parallel the repeated emergence of dung-specialization from plant-decay

ancestors in the radiation of *Psilocybe* and other Agaricales lineages (Ramirez-Cruz

et al. 2013; Tóth et al. 2013). Late stage wood decay fungi like *Psilocybe* spp. likely

harbor genetic exaptations for lignin tolerance/degradation, and competition with

invertebrates and prokaryotes, and acquisition of particularly adaptive functions by

other fungi (e.g. *Panaeolus*) through HT may have facilitated additional transitions

to dung saprotrophy.

The evolution of PS genes suggest they originally served roles in the wood-decay

niche, and more recently emerged through both vertical and horizontal transfer in

dung-decay fungi (Fig. 2). HT and retention of PS clusters are evidence of selection

on the PS pathway in the recipient lineage, as SM clusters are inherently unstable in

fungal genomes (Reynolds et al. 2017 Apr 28). Psilocybin neurological activity,

coupled with HT and retention in lineages that colonize dung and/or decayed wood,

which are rich in both mycophagous and competitor invertebrates (Rouland-Lefevre

2000), suggest that psilocybin may be a modulator of insect behavior. Psilocybin

and/or the related aeruginascin have also been identified in the lichenized agaric,

Dictyonema huaorani, and in the ectomycorrhizal genus *Inocybe* (Kosentka et al.

2013; Schmull et al. 2014). PS distribution in *Inocybe* is complementary to that of

the acetylcholine mimic, muscarine, which could suggest alternative strategies and

pressures to manipulate animal behavior beyond the dung and wood decay niches.

Neurotransmitter mimics may provide advantages to fungi by interfering with the

behavior of invertebrate competitors for woody resources (Hunt et al. 2007),

especially social insects, like termites, which emerged ~ 137 Mya, because they rely

on the coordinated activities of multiple castes (Genise 2017). It is thus intriguing

that D4H and PST have experienced massive gene family expansion through

duplication in *Fibulorhizoctonia* sp., which produce termite egg-mimicking sclerotia

in an ancient mutualistic relationship with *Reticulitermes* termites (Matsuura 2005).

While neurotransmitter agonists are not known to mediate this symbiosis, insect

predatory fungi (i.e. *Cordyceps* spp.) use neurotransmitter analogs to influence the

behavior of infected insects (de Bekker et al. 2014), and a number of repellents and

toxins in wood-decay fungi inhibit xylophagy and mycophagy by termites (Rouland-

Lefèvre 2000).

The identification of genes underlying PS biosynthesis is an important advance in

the field of neurochemical ecology, with both social and medical applications. The

sequences of the first *Psilocybe* and *Panaeolus* genomes presented here will be

important resources for the prospecting of novel neurotropic natural products

(Rutledge and Challis 2015). The discovery that a psilocybin cluster has been

horizontally transferred and subsequently maintained among the invertebrate-

challenged environments of dung and late wood-decay suggests these niches may be

reservoirs not only of new antibiotics (Bills et al. 2013 Aug 23), but also novel

neuroactive pharmaceuticals.

Acknowledgements:

The authors thank Jan Borovička and Paul Stamets for providing additional cultures

used in this study. We also thank Michael Kelly, Ma Elena Hernandez, Stephen Opiyo,

206   **Table 1.** Genome assembly and annotation of psilocybin-producing mushrooms.

	length (nt)	scaffolds	contigs	av. depth of coverage (x)	N50 (nt)	Complete BUSCOs (%)	total proteins	decarboxylases / PSD [a]	P450's[a]	methyltransferases / DUF890 domain-proteins[a]	kinases / phosphotransferases / OG term 0PNAW[a]	MFS[a]
G. dilepis[b] TENN071165	47,177,497	8,423	10,681	16.5	33,540	73.43	16,257	28 / 9	151	89 / 4	275 / 29 / 1	37
Pa. cyanescens	44,965,162	9,521	11,850	25.7	32,751	75.66	13,420	28 / 8	148	91 / 2	267 / 16 / 2	34
Ps. cyanescens	53,483,841	18,721	38,006	44.7	46,250	72.18	15,973	38 /17	178	102 / 2	298 / 23 / 1	44

207
208
209   [a]Functional category of PS genes as annotated in Eggnog. [b]*G. dilepis* (= *G. aeruginosus* sensu L.R. Hesler) was isolated from oak
210   sawdust in Knoxville, Tennessee 4-Oct-2013. *Pa. cyanescens* and *Ps. cyanescens* basidiospores were supplied by The Spore Works,
211   Knoxville.

Figure Legends:

Figure 1. Psilocybin evolution. A. The PS cluster consists of tryptophan decarboxylase (TDC), 1-2 P450 monooxygenases (D4H), methyltransferase (TMT), phosphotransferase (PPT), and 1-2 MFS transporters (PST). B. Phylogenomic tree of Agaricales (tan) with Atheliales (blue) outgroup. Support values = (internode certainty, tree certainty). Clades 1-5 are as in Fig. 2B. C. PS locus synteny relative to *Ps. cyanescens* scaffold 5617 and *G. marginata* scaffold 9. D. RAxML phylogeny of TDC indicating putative HT branches; *Eutypa lata* is in Xylariales (Ascomycota, lavendar), an order correlated with absence of termites in coarse woody debris (Kirker et al. 2012), with members that produce a white rot of wood. *Basidiobolus meristosporus* (Zoopagomycota, rose) is commonly associated with amphibian dung and arthropods. Grey taxon names = PCR sequences, black = whole genome. Support is percent of 100 ML bootstraps. **54 similar D4H homologs not shown.

Figure 2. Patterns of ecological diversification of PS genes and Agaricales genomes. A. Ultrametric representation of PPT phylogeny, with root age hypothetically set to align ecological transitions with Earth history events. HT = horizontal transfer, VT = vertical transmission. B. Ordination of Agaricales genome content. Numerals correspond to clades in Fig. 1A.

References Cited:

Agurell S, Nilsson JL. 1968. Biosynthesis of psilocybin. II. Incorporation of labelled tryptamine derivatives. Acta Chem Scand 22:1210–1218.

Allen JW. 2010. A Chemical Referral and Reference Guide to the Known Species of Psilocin and/or Psilocybin-containing Mushrooms and their Published Analysis and Bluing Reactions.

Baquero F. 2004. From pieces to patterns: evolutionary engineering in bacterial pathogens. Nat Rev Micro 2:510–518.

Bills GF, Gloer JB, An Z. 2013 Aug 23. Coprophilous fungi: antibiotic discovery and functions in an underexplored arena of microbial defensive mutualism. Curr. Opin. Microbiol.

Borovička J, Oborník M, Stříbrný J, Noordeloos ME, Sánchez LAP, Gryndler M. 2015. Phylogenetic and chemical studies in the potential psychotropic species complex of *Psilocybe atrobrunnea* with taxonomic and nomenclatural notes. Persoonia 34:1–9.

Carhart-Harris RL, Bolstridge M, Rucker J, Day CMJ, Erritzoe D, Kaelen M, Bloomfield M, Rickard JA, Forbes B, Feilding A, et al. 2016. Psilocybin with psychological support for treatment-resistant depression: an open-label feasibility study. Lancet Psychiatry 3:619–627.

Carhart-Harris RL, Erritzoe D, Williams T, Stone JM, Reed LJ, Colasanti A, Tyacke RJ, Leech R, Malizia AL, Murphy K, et al. 2012. Neural correlates of the psychedelic state as determined by fMRI studies with psilocybin. Proceedings of the National Academy of Sciences 109:2138–2143.

Choi J-Y, Duraisingh MT, Marti M, Ben Mamoun C, Voelker DR. 2015. From Protease to Decarboxylase. Journal of Biological Chemistry 290:10972–10980.

de Bekker C, Quevillon LE, Smith PB, Fleming KR, Ghosh D, Patterson AD, Hughes DP. 2014. Species-specific ant brain manipulation by a specialized fungal parasite. BMC Evol Biol 14:185.

de Jonge R, Bolton MD, Kombrink A, van den Berg GCM, Yadeta KA, Thomma BPHJ. 2013. Extensive chromosomal reshuffling drives evolution of virulence in an asexual pathogen. Genome Res 23:1271–1282.

Dinis-Oliveira RJ. 2017. Metabolism of psilocybin and psilocin: clinical and forensic toxicological relevance. Drug Metab. Rev. 49:84–91.

Floudas D, Binder M, Riley R, Barry K, Blanchette RA, Henrissat B, Martínez AT, Otillar R, Spatafora JW, Yadav JS, et al. 2012. The Paleozoic origin of enzymatic lignin decomposition reconstructed from 31 fungal genomes. Science

336:1715–1719.

Genise JF. 2017. The Trace Fossil Record of Eusociality in Ants and Termites.
Ichnoentomology.

Gluck-Thaler E, Slot JC. 2015. Dimensions of Horizontal Gene Transfer in Eukaryotic
Microbial Pathogens. PLoS Pathog 11:e1005156.

Griffiths RR, Johnson MW, Richards WA, Richards BD, McCann U, Jesse R. 2011.
Psilocybin occasioned mystical-type experiences: immediate and persisting
dose-related effects. Psychopharmacology (Berl.) 218:649–665.

Halberstadt AL, Geyer MA. 2011. Multiple receptors contribute to the behavioral
effects of indoleamine hallucinogens. Neuropharmacology 61:364–381.

Hofmann A, Frey A, Ott H, Petr Zilka T, Troxler F. 1958. [Elucidation of the structure
and the synthesis of psilocybin]. Experientia 14:397–399.

Holliday JA, Zhou L, Bawa R, Zhang M, Oubida RW. 2015. Evidence for extensive
parallelism but divergent genomic architecture of adaptation along altitudinal
and latitudinal gradients in Populus trichocarpa. New Phytologist 209:1240–
1251.

Hunt T, Bergsten J, Levkanicova Z, Papadopoulou A, John OS, Wild R, Hammond PM,
Ahrens D, Balke M, Caterino MS, et al. 2007. A Comprehensive Phylogeny of
Beetles Reveals the Evolutionary Origins of a Superradiation. Science
318:1913–1916.

Johnson MW, Garcia-Romeu A, Griffiths RR. 2017. Long-term follow-up of
psilocybin-facilitated smoking cessation. Am J Drug Alcohol Abuse 43:55–60.

Kakioka R, Kokita T, Kumada H, Watanabe K, Okuda N. 2015. Genomic architecture
of habitat-related divergence and signature of directional selection in the body
shapes of Gnathopogonfishes. Mol Ecol 24:4159–4174.

Kalb D, Gressler J, Hoffmeister D. 2016. Active-Site Engineering Expands the
Substrate Profile of the Basidiomycete L-Tryptophan Decarboxylase CsTDC.
ChemBioChem 17:132–136.

Kirker GT, Wagner TL, Diehl SV. 2012. Relationship between wood-inhabiting fungi
and *Reticulitermes* spp. in four forest habitats of northeastern Mississippi.
International Biodeterioration & Biodegradation 72: 18-25.

Kosentka P, Sprague SL, Ryberg M, Gartz J, May AL. 2013. Evolution of the toxins
muscarine and psilocybin in a family of mushroom-forming fungi. PLoS ONE.
DOI:10.1371/journal.pone.0064646

Letcher A. 2006. Shroom: a cultural history of the magic mushroom. London: Faber
and Faber.

Ma L-J, van der Does HC, Borkovich KA, Coleman JJ, Daboussi M-J, Di Pietro A,
Dufresne M, Freitag M, Grabherr M, Henrissat B, et al. 2010. Comparative
genomics reveals mobile pathogenicity chromosomes in Fusarium. Nature
464:367–373.

MacFadden BJ. 2000. Cenozoic Mammalian Herbivores From the Americas:
Reconstructing Ancient Diets and Terrestrial Communities. Annu. Rev. Ecol.
Syst. 31:33–59.

Matsuura K. 2005. Distribution of termite egg-mimicking fungi ("termite balls") in
Reticulitermes spp. (Isoptera : Rhinotermitidae) nests in Japan and the United
States. Applied Entomology and Zoology 40:53–61.

Nyberg H. 1992. Religious use of hallucinogenic fungi: A comparison between
Siberian and Mesoamerican Cultures. Karstenia 32:71-80.

Petri G, Expert P, Turkheimer F, Carhart-Harris R, Nutt D, Hellyer PJ, Vaccarino F.
2014. Homological scaffolds of brain functional networks. Journal of The Royal
Society Interface 11:20140873.

Quin MB, Flynn CM, Schmidt-Dannert C. 2014. Traversing the fungal terpenome. Nat
Prod Rep 31:1449–1473.

Raguso RA, Agrawal AA, Douglas AE, Jander G. 2015. The raison d'être of chemical
ecology. Ecology 96:617-630.

Ramirez-Cruz V, Guzman G, Rosa Villalobos-Arambula A, Rodriguez A, Matheny PB,
Sanchez-Garcia M, Guzmán-Dávalos L. 2013. Phylogenetic inference and trait
evolution of the psychedelic mushroom genus *Psilocybe* sensu lato
(Agaricales). Botany-Botanique 91:573–591.

Retallack GJ. 2001. Cenozoic Expansion of Grasslands and Climatic Cooling. The
Journal of Geology 109:407–426.

Reynolds H, Slot JC, Divon HH, Lysøe E, Proctor RH, Brown DW. 2017 Apr 28.
Differential retention of gene functions in a secondary metabolite cluster. Mol
Biol Evol. DOI: 10.1093/molbev/msx145

Rouland-Lefèvre C. 2000. Symbiosis With Fungi. In: Termites: evolution. Dordrecht:
Springer Netherlands. pp. 289–306; 18 p.

Rutledge PJ, Challis GL. 2015. Discovery of microbial natural products by activation
of silent biosynthetic gene clusters. Nat Rev Micro 13:509–523.

Schmull M, Dal-Forno M, Lücking R, Cao S, Clardy J, Lawrey JD. 2014. *Dictyonema*
huaorani (Agaricales: Hygrophoraceae), a new lichenized basidiomycete from
Amazonian Ecuador with presumed hallucinogenic properties. The Bryologist
117:386–394.

Tóth A, Hausknecht A, Krisai-Greilhuber I, Papp T, Vágvölgyi C, Nagy LG. 2013.
Iteratively refined guide trees help iImproving alignment and phylogenetic
inference in the mushroom family Bolbitiaceae. PLoS ONE 8:e56143.

Wisecaver JH, Slot JC, Rokas A. 2014. The evolution of fungal metabolic pathways.
PLoS Genet 10:e1004816.

Wriessnegger T, Sunga AJ, Cregg JM, Daum G. 2009. Identification of
phosphatidylserine decarboxylases 1 and 2 from *Pichia pastoris*. FEMS Yeast
Res 9:911–922.

A

L-Tryptophan → **TDC** → Tryptamine → **TMT** → N,N-DMT → **D4H** → Psilocin → **PPT** → Psilocybin → **PST**

Shared synteny   Specific synteny

B Species phylogeny
0.04

C

D TDC phylogeny
80.0

PS cluster

HGT?