Stochastic processes dominate the within and between host evolution of influenza virus

Running Title: Influenza dynamics within and between hosts

John T. McCrone[1], Robert J. Woods[2], Emily T. Martin[3], Ryan E. Malosh[3], Arnold S. Monto[3], and Adam S. Lauring[1,2] [*]

[1] Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI 48109

[2] Division of Infectious Diseases, Department of Internal Medicine, University of Michigan, Ann Arbor, MI 48109

[3] Department of Epidemiology, University of Michigan, Ann Arbor, MI 48109

* Corresponding author

Adam S. Lauring

1150 W. Medical Center Dr.

MSRB1 Room 5510B

Ann Arbor, MI 48109-5680

(734) 764-7731

alauring@med.umich.edu

Abstract Word Count:

Main Text Word Count:

Key Words – influenza virus, bottleneck, transmission, diversity, evolution

Abstract

The global evolutionary dynamics of influenza virus ultimately derive from processes that take

place within and between infected individuals. Here we define the dynamics of influenza A virus

populations in human hosts through next generation sequencing of 249 specimens from 200

individuals collected over 6290 person-seasons of observation. Because these viruses were

collected over 5 seasons from individuals in a prospective community-based cohort, they are

broadly representative of natural human infections with seasonal viruses. We used viral

sequence data from 35 serially sampled individuals to estimate a within host effective population

size of 30-70 and an in vivo mutation rate of 4×10^{-5} per nucleotide per cellular infectious cycle.

These estimates are consistent across several models and robust to the models' underlying

assumptions. We also identified 43 epidemiologically linked and genetically validated

transmission pairs. Maximum likelihood optimization of multiple transmission models estimates

an effective transmission bottleneck of 1-2 distinct genomes. Our data suggest that positive

selection of novel viral variants is inefficient at the level of the individual host and that genetic

drift and other stochastic processes dominate the within and between host evolution of influenza

A viruses.

Introduction

The rapid evolution of influenza viruses has led to reduced vaccine efficacy, widespread drug

resistance, and the continuing emergence of novel strains. Broadly speaking, evolution is the

product of deterministic processes, such as selection, and stochastic processes, such as

genetic drift (Kouyos et al. 2006). The relative contribution of each is greatly affected by the

effective population size, or size of an idealized population whose dynamics are similar to that of

the population in question (Rouzine et al. 2001). If the effective population size of a virus is large,

as in quasispecies models, evolution is largely deterministic and the frequency of a mutation

can be predicted based on its starting frequency and selection coefficient. In small populations,

selection is inefficient, and changes in mutation frequency are strongly influenced by migration

or genetic drift.

Viral dynamics may differ across spatial and temporal scales, and a complete understanding of

influenza evolution requires studies at all levels (Nelson & Holmes 2007; Holmes 2009) .The

global evolution of influenza A virus (IAV) is dominated by the positive selection of novel

antigenic variants that circulate in the tropics and subsequently seed annual epidemics in the

Northern and Southern hemisphere (Rambaut et al. 2008). Whole genome sequencing has also

demonstrated the importance of intrasubtype reassortment to the emergence of diverse strains

that differ in their antigenicity. While continual positive selection of antigenically drifted variants

drives global patterns, whole genome sequencing of viruses on more local scales suggests the

importance of stochastic processes such as strain migration and within-clade reassortment

(Nelson et al. 2006).

It is now feasible to efficiently sequence patient-derived isolates at sufficient depth of coverage

to define the diversity and dynamics of virus evolution within individual hosts (Kao et al. 2014).

Studies of IAV populations in animal and human systems suggest that most intrahost single

nucleotide variants (iSNV) are rare and that intrahost populations are subject to strong purifying

selection (Rogers et al. 2015; Murcia et al. 2010; Iqbal et al. 2009; Poon et al. 2016; Dinis et al.

2016; Debbink et al. 2017). While positive selection of adaptive variants is commonly observed

in cell culture (Doud et al. 2017; ARCHETTI & HORSFALL 1950; Foll et al. 2014), it has only

been documented within human hosts in the extreme cases of drug resistance (Gubareva et al.

2001; Ghedin et al. 2011; Rogers et al. 2015), long-term infection of immunocompromised hosts

(Xue et al. 2017) or experimental infections with attenuated viruses (Sobel Leonard et al. 2016).

Indeed, we and others have been unable to identify evidence for positive selection in natural

human infections (Debbink et al. 2017; Dinis et al. 2016), and its relevance to within host

processes is unclear.

Despite limited evidence for positive selection, it is clear that novel mutations do arise within

hosts. Their potential for subsequent spread through host populations is determined by the size

of the transmission bottleneck (Alizon et al. 2011; Zwart & Elena 2015). If the transmission

bottleneck is sufficiently wide, low frequency variants can plausibly be transmitted and spread

through host populations (Geoghegan et al. 2016). Because the transmission bottleneck is

conceptually similar to the effective population size between hosts, its size will also inform the

relative importance of selection and genetic drift in determining which variants are transmitted.

While experimental infections of ferrets suggest a very narrow transmission bottleneck (Varble

et al. 2014; Wilker et al. 2013), studies of equine influenza support a bottleneck wide enough to

allow transmission of rare iSNV (Hughes et al. 2012; Murcia et al. 2010). The only available

genetic study of influenza virus transmission in humans estimated a remarkably large

transmission bottleneck, allowing for transmission of 100-200 genomes (Poon et al. 2016; Sobel

Leonard et al. 2017).

Here, we use next generation sequencing of within host influenza virus populations to define the

evolutionary dynamics of influenza A viruses (IAV) within and between human hosts. We apply

a benchmarked analysis pipeline to identify iSNV and to characterize the genetic diversity of

H3N2 and H1N1 populations collected over five post-pandemic seasons from individuals

enrolled in a prospective household study of influenza. We use these data to estimate the *in*

vivo mutation rate and the within and between host effective population size. We find that

intrahost populations are characterized by purifying selection, a small effective population size,

and limited positive selection. Contrary to what has been previously reported for human

influenza transmission (Poon et al. 2016), but consistent with what has been observed in other

viruses (Zwart & Elena 2015), we identify a very tight effective transmission bottleneck that

limits the transmission of rare variants.

Results

We used next generation sequencing to characterize influenza virus populations collected from

individuals enrolled in the Household Influenza Vaccine Effectiveness (HIVE) study (Monto et al.

2014; Ohmit et al. 2013; Ohmit et al. 2015; Ohmit et al. 2016; Petrie et al. 2013), a community-

based cohort that enrolls 213-340 households of 3 or more individuals in Southeastern Michigan

each year (Table 1). These households are followed prospectively from October to April, with

symptom-triggered collection of nasal and throat swab specimens for identification of respiratory

viruses by RT-PCR (see Methods). In contrast to case-ascertained studies, which identify

households based on an index case who seeks medical care, the HIVE study identifies

symptomatic individuals regardless of illness severity. In the first four seasons of the study

(2010-2011 through 2013-2014), respiratory specimens were collected 0-7 days after illness

onset. Beginning in the 2014-2015 season, each individual provided two samples, a self-

collected specimen at the time of symptom onset and a clinic-collected specimen obtained 0-7

days later. Each year, 59-69% of individuals had self-reported or confirmed receipt of that

season's vaccine prior to local circulation of influenza virus.

Over five seasons and nearly 6,290 person-seasons of observation, we identified 77 cases of

influenza A/H1N1pdm09 infection and 313 cases of influenza A/H3N2 infection (Table 1).

Approximately half of the cases (n=166) were identified in the 2014-2015 season, in which there

was an antigenic mismatch between the vaccine and circulating strains (Flannery et al. 2016).

All other seasons were antigenically matched. Individuals within a household were considered

an epidemiologically linked transmission pair if they were both positive for the same subtype of

influenza virus within 7 days of each other. Several households had 3 or 4 symptomatic cases

within this one-week window, suggestive of longer chains of transmission (Table 1).

Within host populations have low genetic diversity

We processed all specimens for viral load quantification and next generation sequencing. Viral

load measurements (genome copies per μl) were used for quality control in variant calling,

which we have shown is highly sensitive to input titer (McCrone & Lauring 2016) (Figure 1A).

Accordingly, we report data on 249 high quality specimens from 200 individuals, which had a

viral load of $>10^3$ copies per microliter of transport media, adequate RT-PCR amplification of all

eight genomic segments, and an average read coverage of $>10^3$ across the genome (Table 1,

Supplementary Figure 1).

We identified intrahost single nucleotide variants (iSNV) using our empirically validated analysis

pipeline (McCrone & Lauring 2016). Our approach relies heavily on the variant caller DeepSNV,

which uses a clonal plasmid control to distinguish between true iSNV and errors introduced

during sample preparation and/or sequencing (Gerstung et al. 2012). Given the diversity of

influenza viruses that circulate locally each season, there were a number of instances in which

our patient-derived samples had mutations that were essentially fixed (>0.95 frequency) relative

to the clonal control. DeepSNV is unable to estimate an error rate for the control or reference

base at these positions. We therefore performed an additional benchmarking experiment to

identify a threshold for majority iSNV at which we could correctly infer whether or not the

corresponding minor allele was also present (see Methods). We found that we could correctly

identify a minor allele at a frequency of ≥2% when the frequency of the major allele was ≤98%.

We therefore report data on iSNV present at frequencies between 2 and 98%. As expected, this

threshold improved the specificity of our iSNV identification and decreased our sensitivity to

detect variants below 5% compared to our initial validation experiment (McCrone & Lauring

2016), which did not employ a frequency threshold (Supplementary Table 1).

Consistent with our previous studies and those of others, we found that the within host diversity

of human influenza A virus (IAV) populations is low (Poon et al. 2016; Dinis et al. 2016; Debbink

et al. 2017; Sobel Leonard et al. 2016; McCrone & Lauring 2016). Two hundred forty-three out

of the 249 samples had fewer than 10 minority iSNV (median 2, IQR 1-3). There were 6

samples with greater than 10 minority iSNV. In 3 of these cases, the frequency of iSNVs were

tightly distributed about a mean suggesting that the iSNV were linked and that the samples

represented mixed infections. Consistent with this hypothesis, putative genomic haplotypes

based on these minority iSNV clustered with distinct isolates on phylogenetic trees

(Supplementary Figures 2 and 3). While viral shedding was well correlated with days post

symptom onset (Figure 1A) the number of minority iSNV identified was not affected by the day

of infection, viral load, subtype, or vaccination status (Figure 1B and Supplementary Figure 4).

The vast majority of minority variants were rare (frequency 0.02-0.07), and iSNV were

distributed evenly across the genome (Figure 1C and 1D). The ratio of nonsynonymous to

synonymous variants was 0.64 and was never greater than 1 in any 5% bin, which suggests

that within host populations were under purifying selection. We also found that minority variants

were rarely shared among multiple individuals. Ninety-five percent of minority iSNV were only

found once, 4.7% were found in 2 individuals, and no minority iSNV were found in more than 3

individuals. The low level of shared diversity suggests that within host populations were

exploring distinct regions of sequence space with little evidence for parallel evolution. Of the 31

minority iSNV that were found in multiple individuals (triangles in Figure 1D), 4 were

nonsynonymous.

Although the full range of the H3 antigenic sites have not been functionally defined, it is

estimated that 131 of the 329 amino acids in HA1 lie in or near these sites (Lee & Chen 2004).

We identified 17 minority nonsynonymous iSNV in these regions (Supplementary Table 2). Six

of these were in positions that differ among antigenically drifted viruses (Smith et al. 2004; Wiley

et al. 1981), and two (193S and 189N) lie in the "antigenic ridge" that is a major contributor to

drift (Koel et al. 2013). Three of these have been detected at the global level as consensus

variants since the time of isolation (128A, 193S and 262N) with two (193S and 262N) seemingly

increasing in global frequency (Neher & Bedford 2015) (Supplementary Figure 5). Additionally,

we identified 1 putative H1N1 antigenic variant (208K in C_a) (Caton et al. 1982; Xu et al. 2010).

In total, putative antigenic variants account for 1.0-2.5% of minority iSNV identified and were

found in 3.5-8.0% of infections. None of these iSNV were shared among multiple individuals.

Estimation of effective population size

Given the above observations, we hypothesized that within host populations of IAV are under

purifying selection and that variants that rise to detectable levels do so by a neutral process as

opposed to positive selection. Consistent with this hypothesis, we found that nonsynonymous

and synonymous iSNV exhibited similar changes in frequency over time in the 35 individuals

who provided serial specimens that contained iSNV (Figure 2A and 2B). We used a maximum

likelihood approach to estimate the within host effective population size (N_e) of IAV by fitting a

diffusion approximation of the Wright-Fisher model (Kimura 1955). This model assumes that

changes in iSNV frequency are due solely to random genetic drift and not selection, that iSNV

are independent of one another, and that the effective population is sufficiently large to justify a

continuous approximation to changes in allele frequency. The diffusion approximation of the

Wright-Fisher model assigns probabilities to frequency changes given an N_e and the number of

generations between sample times. In our model we fixed the within host generation time as

either 6 or 12 hours (Geoghegan et al. 2016) and report the findings for the 6 hour generation

time below. We then asked what population size makes the observed changes in frequency

most likely (Figure 2B). We restricted this analysis to samples taken at least 1 day apart (n = 29),

as there was very little change in iSNV frequency in populations sampled twice on the same day

(R^2 = 0.986, Figure 2B and Supplementary Figure 6). The concordance of same day samples

suggests that our sampling procedure is reproducible and that less than a generation had

passed between samplings. Maximum likelihood optimization of this diffusion model revealed a

within host effective population size of 35 (95% CI 26-46, Table 2).

The diffusion approximation makes several simplifying assumptions, which if violated could

influence our findings. To ensure our results were robust to the assumption of a large population,

we employed a discrete interpretation of the Wright-Fisher model which makes no assumptions

about population size (Williamson & Slatkin 1999). In this case we found an effective population

size of 32 (95% CI 28-41), very close to our original estimate (Table 2). Both models assume

complete independence of iSNV. To ensure this assumption did not affect our results, we fit the

discrete model 1000 times, each time randomly subsetting our data such that only one iSNV per

individual was included. This simulates a situation in which all modeled iSNV are independent

and our assumption is met. Under these conditions we found a median effective population size

of 33 (IQR 32-40), demonstrating negligible bias in the initial analysis due to correlation between

iSNV.

As above, most iSNV in the longitudinal samples were rare (< 10%) and many became extinct

between samplings. To ensure that our models were capable of accurately estimating the

effective population size from such data, we simulated 1000 Wright-Fisher populations with

iSNV present at approximately the same starting frequencies as in our data set and an N_e of 30,

50, or 100. In these simulations, we found mean N_e of 34, 56 and 117 (Figure 2C). These

simulations suggest that although this method may slightly overestimate the N_e, our results are

not constrained by the data structure.

To this point, we have assumed that neutral processes are responsible for the observed

changes in iSNV frequency within hosts. Although this assumption seems justified at least in

part by the analysis above, we tested the robustness of our models by fitting the

nonsynonymous (n = 27) and synonymous iSNV (n = 36) separately. Here, we estimated an

effective population size of 30 using the nonsynonymous iSNV and an effective population size

of 37 using the synonymous iSNV (Table 2). These estimates are very close to those derived

from the whole dataset and suggest that nonsynonymous and synonymous mutations are

influenced by similar within host processes. To further ensure that our results were not driven by

a few outliers subject to strong selection, we ranked iSNV by their change in frequency over

time and consecutively removed iSNV with the most extreme changes. We estimated the

effective population size at each iteration and found that removing the top 50% most extreme

iSNV only increased the effective population size to 161 (Figure 2D). Therefore, our estimates

are robust to a reasonable number of non-neutral sites. Finally, we also applied a separate

Approximate Bayesian Computational (ABC) method, which uses a non-biased moment

estimator in conjunction with ABC to estimate the effective population size of a population as

well as selection coefficients for the iSNV present (Foll et al. 2014). This distinct approach

relaxes the previous assumption regarding neutrality. We applied this analysis to the 16

longitudinal pairs that were sampled 1 day apart and estimated an effective population of 69.

We were unable to reject neutrality for just 7 of the 35 iSNV in this data set (Figure 2E). These

seven mutations consisted of 3 nonsynonymous and 4 synonymous mutations and were split

between two individuals. None were putative antigenic variants.

Identification of forty-three transmission pairs

We analyzed virus populations from 85 households with concurrent infections to quantify the

level of shared viral diversity and to estimate the size of the IAV transmission bottleneck (Table

1). Because epidemiological linkage does not guarantee that concurrent cases constitute a

transmission pair (Petrie et al. 2017), we used a stringent rubric to eliminate individuals in a

household with co-incident community acquisition of distinct viruses. We considered all

individuals in a household with symptom onset within a 7-day window to be epidemiologically

linked. The donor in each putative pair was defined as the individual with the earlier onset of

symptoms. We discarded a transmission event if there were multiple possible donors with the

same day of symptom onset. Donor and recipients were not allowed to have symptom onset on

the same day, unless the individuals were both index cases for the household. In these 6

instances, we analyzed the data for both possible donor-recipient directionalities. Based on

these criteria, our cohort had 124 putative household transmission events over 5 seasons

(Table 1). Of these, 52 pairs had samples of sufficient quality for reliable identification of iSNV

from both individuals.

We next used sequence data to determine which of these 52 epidemiologically linked pairs

represented true household transmission events as opposed to coincident community-acquired

infections. We measured the genetic distance between influenza populations from each

household pair by L1-norm and compared these distances to those of randomly assigned

community pairs within each season (Figure 3A, see also trees in Supplementary Figures 2 and

3). While the L1-norm of a pair captures differences between the populations at all levels, in our

cohort, it was largely driven by differences at the consensus level. We only considered

individuals to be a true transmission pair if they had a genetic distance below the 5th percentile

of the community distribution of randomly assigned pairs (Figure 3A). Forty-seven household

transmission events met this criterion (Figure 3B). Among these 47 sequence-validated

transmission pairs, 3 had no iSNV in the donor and 1 additional donor appeared to have a

mixed infection. These four transmission events were removed from our bottleneck analysis as

donors without iSNV are uninformative and mixed infections violate model assumptions of site

independence (see Methods). We estimated the transmission bottleneck in the remaining 43

high-quality pairs (37 H3N2, 6 H1N1, Figure 3B).

A transmission bottleneck restricts the amount of genetic diversity that is shared by both

members of a pair. We found that few minority iSNV where polymorphic in both the donor and

recipient populations (Figure 3C). Minority iSNV in the donor were either absent or fixed in the

recipient (top and bottom of plot). The lack of shared polymorphic sites (which would lie in the

middle of the plot in Figure 3C) suggests a stringent effective bottleneck in which only one allele

is passed from donor to recipient.

Estimation of the transmission bottleneck

We applied a simple presence-absence model to quantify the effective transmission bottleneck

in our cohort. The presence-absence model considers only whether or not a donor allele is

present or absent in the recipient sample. Under this model, transmission is a neutral, random

sampling process, and the probability of transmission is simply the probability that the iSNV will

be included at least once in the sample given its frequency in the donor and the sample size, or

bottleneck. We estimated a distinct bottleneck for each transmission pair and assumed these

bottlenecks followed a zero-truncated Poisson distribution. This model also assumes that the

sensitivity for detection of transmitted iSNVs is perfect and that each genomic site is

independent of all others. We then used maximum likelihood optimization to determine the

distribution of bottleneck sizes that best fit the data. We found a zero-truncated Poisson

distribution with a mean of 1.66 (lambda = 1.12; 0.51-1.99, 95% CI) best described the data.

This distribution indicates that the majority of bottlenecks are 1, and that very few are greater

than 5 (probability 0.2%). There were no apparent differences between H3N2 and H1N1 pairs.

The model fit was evaluated by simulating each transmission event 1,000 times. The presence

or absence of each iSNV in the recipient was noted and the probability of transmission given

donor frequency determined. The range of simulated outcomes matched the data well, which

suggests that transmission is a selectively neutral event characterized by a stringent bottleneck

(Figure 3D).

The majority of transmitted iSNV were fixed in the recipients. Although this trend matches the

expectation given a small bottleneck, these data could also be consistent with a model in which

the probability of transmission is determined by the frequency at which iSNV are found at the

community level. To ensure our bottleneck estimates were an outcome of neutral transmission

and not an artifact of the larger community population structure or selection for the community

consensus, we created a null model by randomly assigning community "recipient-donor"

pairings. Each community "recipient" was drawn from the pool of individuals that were infected

after the "donor" but in the same season and with the same subtype as the donor. We then

identified whether or not each donor iSNV was found in the community recipient and determined

the relationship between "donor" frequency and probability of "transmission" for 1,000 such

simulations. Given the low level of diversity in our cohort, we predicted that rare iSNV would be

unlikely to be found in a random sample, while the major alleles should be fixed in most random

pairs. This trend is clearly demonstrated in Figure 3E. It is also clear that this null model fit the

data much more poorly than the presence/absence model, suggesting that the observed data in

our bona fide transmission pairs were not a product of community metapopulation structure, but

rather an outcome of neutral sampling events.

Because our bottleneck estimates were much lower than what has previously been reported for

human influenza (Poon et al. 2016), we investigated the impact that our simplifying assumptions

could have on our results. In particular, the presence-absence model assumes perfect detection

of variants in donor and recipient, and it can therefore underestimate the size of a bottleneck in

the setting of donor-derived variants that are transmitted but not detected in the recipient. These

"false negative" variants can occur when the frequency of an iSNV drifts below the level of

detection (e.g. 2% frequency) or when the sensitivity of sequencing is less than perfect for

variants at that threshold (e.g. 15% sensitivity for variants at a frequency 2-5%). To determine

the impact of sequencing sensitivity and specificity on our bottleneck estimates, we re-called

variants using our original pipeline without the 2% frequency cut-off. As shown in

Supplementary Table 1, this increases the sensitivity of iSNV detection in the 1-5% frequency

range, and also the number of false positive variant calls (McCrone & Lauring 2016). This

analysis only slightly increased average transmission bottleneck to 2.10 (lambda = 1.67; 0.91-

2.71, 95% CI), and indicates that our results are not biased by the added stringency used in the

initial analysis (Supplementary Figures 7A and 7B).

To further investigate the impact of sequencing accuracy on our estimates, we inferred minor

variants in our current pipeline (see above and methods) without a frequency cutoff. Ultimately,

this reduced variant calling to a count method at a number of positions and greatly increased

the number of shared minority iSNV in our samples (Supplementary Figure 7C). Many of these

presumed false positive variant calls were at similar frequencies (0.1-2%) in donor and recipient.

As such, the "apparent transmission" of rare variants drives an inflated estimate of the

transmission bottleneck (118, see Supplementary Figure 7D). Simulation showed that this

inflated bottleneck no longer fit the trend in the data, likely because the model is now forced to

accommodate shared iSNV that are biased toward sequencing error as opposed to the actual

transmission process.

We also estimated bottleneck size using a beta binomial model, which Leonard *et al.* have used

to account for the stochastic loss of transmitted variants. This model allows for a limited amount

of time-independent genetic drift within the recipient (Sobel Leonard et al. 2017), and we

modified it to also account for our benchmarked sensitivity for rare variants (Supplementary

Table 1, Current Pipeline). For all donor-derived iSNV that were absent in the recipient, we

estimated the likelihood that these variants were transmitted but either drifted below our level of

detection or drifted below 10% and were missed by our variant identification. Despite the

relaxed assumptions provided by this modified beta binomial model, maximum likelihood

estimation only marginally increased the average bottleneck size (mean 1.73: lambda 1.22;

0.57-2.17, 95%CI) relative to the simpler presence-absence model. We simulated transmission

and subsequent random drift using the beta binomial model and the estimated bottleneck

distribution as above (Figure 3F). Although the model matched the data well, the fit was not

better than that of the presence-absence model (AIC 83.0 for beta-binomial compared to 76.7

for the presence-absence model).

The mutation rate of influenza A virus within human hosts

The stringent influenza transmission bottleneck suggests that most infections are founded by

one lineage and develop under essentially clonal processes. The diffusion approximation to the

Wright-Fisher model (see above and Figure 2) can be used to predict the rate at which

homogenous populations diversify from a clonal ancestor as a function of mutation rate and

effective population size (Rouzine et al. 2001). Maximum likelihood optimization of this model

suggested an *in vivo* neutral mutation rate of 4×10^{-6} mutations per nucleotide per replication

cycle and a within host effective population size of 36 (given a generation time of 6 hours).

These estimates are consistent with those above (Table 2). As we have recently estimated that

13% of mutations in influenza A virus are neutral (Visher et al. 2016), we estimated that the true

in vivo mutation rate would be approximately 8 fold higher than our neutral rate – on the order of

$3\text{-}4 \times 10^{-5}$. This in vivo mutation rate is close to our recently published estimate of influenza A

mutation rates in epithelial cells by fluctuation test (Pauly et al. 2017) and within the range of

other estimates for IAV (Sanjuán et al. 2010).

Discussion

We find that seasonal influenza A viruses replicate within and spread among human hosts with

very small effective population sizes. Because we used viruses collected over five influenza

seasons from individuals enrolled in a prospective household cohort, these dynamics are likely

to be broadly representative of many seasonal influenza infections in their natural transmission

context. Our results are further strengthened by the use of a validated sequence analysis

pipeline and models that are robust to the underlying assumptions. The small effective size of

intrahost populations and the tight effective transmission bottleneck suggest that stochastic

processes, such as genetic drift, dominate influenza virus evolution at the level of individual

hosts. This stands in contrast to prominent role of positive selection in the global evolution of

seasonal influenza.

While influenza virus populations are subject to continuous natural selection, selection is an

inefficient driver of evolution in small populations (Rouzine et al. 2001). Despite a large viral

copy number, our findings demonstrate that intrahost populations of influenza behave like much

smaller populations. We therefore expect stochastic fluctuations to be the major force driving

the fixation of novel variants within human hosts. This finding contradicts previous studies,

which have found signatures of adaptive evolution in infected hosts (Gubareva et al. 2001;

Rogers et al. 2015; Ghedin et al. 2011; Sobel Leonard et al. 2016). However, these studies rely

on data from infections in which selective pressures are likely to be particularly strong (e.g. due

to drug treatment or infection with a poorly adapted virus), or in which the virus has been

allowed to propagate for extended periods of time (Xue et al. 2017). Under these conditions,

one can identify the action of positive selection on within host populations. We suggest that

these are important and informative exceptions to the drift regime defined here.

We used both a simple presence-absence model and a more complex beta binomial model to

estimate an extremely tight transmission bottleneck. The estimation of a small bottleneck size is

driven by low within-host diversity and very few minority iSNV shared among individuals in a

transmission pair. While our methods for variant calling may be more conservative than those

used in similar studies, we found that relaxing our variant calling criteria led to the inclusion of

false positive variants that inflated our estimates. Furthermore, the beta binomial model

accounts for false negative iSNV (i.e. variants that are transmitted but not detected in the donor),

which can lead to underestimated transmission bottlenecks (Sobel Leonard et al. 2017). Our

formulation of this model incorporates empirically determined sensitivity and specificity metrics

to account for both false negative iSNV and false positive iSNV (McCrone & Lauring 2016).

Finally, if rare, undetected, iSNV were shared between linked individuals, we would expect to

see transmission of more common iSNV (frequency 5-10%), which we can detect with high

sensitivity. In our data, the transmission probability iSNVs > 5% frequency in the donor were

also well predicted by small bottleneck size (Figure 3D).

Although the size of our transmission bottleneck is consistent with estimates obtained for other

viruses and in experimental animal models of influenza (Zwart & Elena 2015; Varble et al. 2014),

it differs substantially from the only other study of bottlenecks in natural human infection (Poon

et al. 2016; Sobel Leonard et al. 2017). While there are significant differences in the design and

demographics of the cohorts, the influenza seasons under study, and sequencing methodology

(Kugelman et al. 2017), the bottleneck size estimates are fundamentally driven by the amount of

viral diversity shared among individuals in a household. Importantly, we used both

epidemiologic linkage and the genetic relatedness of viruses in households to define

transmission pairs and to exclude confounding from the observed background diversity in the

community. We find that household transmission pairs and randomly assigned community pairs

had distinct patterns of shared consensus and minority variant diversity. The comparison to

random community pairs is important, as an unexplained aspect of the work of Poon et al. is

that rare iSNV were frequently shared by randomly selected individuals, and more common

ones were not (Poon et al. 2016).

Our estimates of IAV population dynamics are consistent across three separate models and

partitions of the data. The measurements of shared diversity are influenced by both between

and within host processes, and the transmission bottleneck is entirely consistent with the small

within host population size derived from the longitudinal samples. We also jointly estimated the

in vivo mutation rate and effective population size based on the frequency distribution of minor

alleles observed in the entire cohort. This model assumed a small transmission bottleneck,

produced a mutation rate that this consistent with previous estimates, and independently

reproduced the within host population size estimate. Given the concordance among these

distinct approaches, it is unlikely that our findings are biased by hidden assumptions or model

limitations.

Accurately modeling and predicting influenza virus evolution requires a thorough understanding

of the virus' population structure. Some models have assumed a large intrahost population and

a relatively loose transmission bottleneck (Geoghegan et al. 2016; Russell et al. 2012; Peck et

al. 2015). Here, adaptive iSNV can rapidly rise in frequency and low frequency variants can

have a high probability of transmission. In such a model, it would be possible for the highly

pathogenic H5N1 virus to develop the requisite 4-5 mutations to become transmissible through

aerosols during a single acute infection of a human host (Herfst et al. 2012; Russell et al. 2012).

Although the dynamics of emergent avian influenza and human adapted seasonal viruses likely

differ (Petrova & Russell 2017), our work suggests that fixation of multiple mutations over the

course of a single acute infection is unlikely.

While it may seem counterintuitive that influenza evolution is dominated by drift on local scales

and positive selection on global scales, these models are certainly not in conflict. Within

individuals we have shown that the effective population is quite small, which suggests that

selection is inefficient. Indeed, we have deeply sequenced 332 intrahost populations from 283

individuals collected over more than 11,000 person-seasons of observation and only identified a

handful of minority antigenic variants with little evidence for positive selection (this work and

(Debbink et al. 2017)). However, with several million infected individuals each year, even

inefficient processes and rare events are likely to happen at a reasonable frequency on a global

scale.

Methods

Description of the cohort

The HIVE cohort (Monto et al. 2014; Ohmit et al. 2013; Ohmit et al. 2015; Ohmit et al. 2016;

Petrie et al. 2013), established at the UM School of Public Health in 2010, enrolled and followed

households of at least 3 individuals with at least two children <18 years of age; households

were then followed prospectively throughout the year for ascertainment of acute respiratory

illnesses. Study participants were queried weekly about the onset of illnesses meeting our

standard case definition (two or more of: cough, fever/feverishness, nasal congestion, sore

throat, body aches, chills, headache if ≥3 yrs old; cough, fever/feverishness, nasal

congestion/runny nose, trouble breathing, fussiness/irritability, decreased appetite, fatigue in <3

yrs old), and the symptomatic participants then attended a study visit at the research clinic on

site at UM School of Public Health for sample collection. For the 2010-2011 through 2013-2014

seasons, a combined nasal and throat swab (or nasal swab only in children < 3 years of age)

was collected at the onsite research clinic by the study team. Beginning with the 2014-2015

seasons, respiratory samples were collected at two time points in each participant meeting the

case definition; the first collection was a self- or parent-collected nasal swab collected at illness

onset. Subsequently, a combined nasal and throat swab (or nasal swab only in children < 3

years of age) was collected at the onsite research clinic by the study team. Families with very

young children (< 3 years of age) were followed using home visits by a trained medical assistant.

Active illness surveillance and sample collection for cases were conducted October through

May and fully captured the influenza season in Southeast Michigan in each of the study years.

Data on participant, family and household characteristics, and on high-risk conditions were

additionally collected by annual interview and review of each participant's electronic medical

record. In the current cohort, serum specimens were also collected twice yearly during fall

(November-December) and spring (May-June) for serologic testing for antibodies against

influenza.

This study was approved by the Institutional Review Board of the University of Michigan Medical

School, and all human subjects provided informed consent.

Identification of influenza virus

Respiratory specimens were processed daily to determine laboratory-confirmed influenza

infection. Viral RNA was extracted (Qiagen QIAamp Viral RNA Mini Kit) and tested by RT-PCR

for universal detection of influenza A and B. Samples with positive results by the universal

assay were then subtyped to determine A(H3N2), A(H1N1), A(pH1N1) subtypes and

B(Yamagata) and B(Victoria) lineages. We used primers, probes and amplification parameters

developed by the Centers for Disease Control and Prevention Influenza Division for use on the

ABI 7500 Fast Real-Time PCR System platform. An RNAseP detection step was run for each

specimen to confirm specimen quality and successful RNA extraction.

Quantification of viral load

Quantitative reverse transcription polymerase chain reaction (RT-qPCR) was performed on 5μl

RNA from each sample using CDC RT-PCR primers InfA Forward, InfA Reverse, and InfA

probe, which bind to a portion of the influenza M gene (CDC protocol, 28 April 2009). Each

reaction contained 5.4μl nuclease-free water, 0.5μl each primer/probe, 0.5μl SuperScript III

RT/Platinum Taq mix (Invitrogen 111732) 12.5μl PCR Master Mix, 0.1μl ROX, 5μl RNA. The

PCR master mix was thawed and stored at 4°C, 24 hours before reaction set-up. A standard

curve relating copy number to Ct value was generated based on 10-fold dilutions of a control

plasmid run in duplicate.

Illumina library preparation and sequencing

We amplified cDNA corresponding to all 8 genomic segments from 5μl of viral RNA using the

SuperScript III One-Step RT-PCR Platinum Taq HiFi Kit (Invitrogen 12574). Reactions consisted

of 0.5μl Superscript III Platinum Taq Mix, 12.5μl 2x reaction buffer, 6μl DEPC water, and 0.2μl

of 10μM Uni12/Inf1, 0.3μl of 10μM Uni12/Inf3, and 0.5μl of 10μM Uni13/Inf1 universal influenza

A primers (Zhou et al. 2009). The thermocycler protocol was: 42˚C for 60 min then 94˚C for 2

min then 5 cycles of 94˚C for 30 sec, 44˚C for 30 sec, 68˚C for 3 min, then 28 cycles of 94˚C for

30 sec, 57˚C for 30 sec, 68˚C for 3 min. Amplification of all 8 segments was confirmed by gel

electrophoresis, and 750ng of each cDNA mixture were sheared to an average size of 300 to

400bp using a Covaris S220 focused ultrasonicator. Sequencing libraries were prepared using

the NEBNext Ultra DNA library prep kit (NEB E7370L), Agencourt AMPure XP beads (Beckman

Coulter A63881), and NEBNext multiplex oligonucleotides for Illumina (NEB E7600S). The final

concentration of each barcoded library was determined by Quanti PicoGreen dsDNA

quantification (ThermoFisher Scientific), and equal nanomolar concentrations were pooled.

Residual primer dimers were removed by gel isolation of a 300-500bp band, which was purified

using a GeneJet Gel Extraction Kit (ThermoFisher Scientific). Purified library pools were

sequenced on an Illumina HiSeq 2500 with 2x125 nucleotide paired end reads. All raw

sequence data have been deposited at the NCBI sequence read archive (BioProject submission

ID: SUB2951236). PCR amplicons derived from an equimolar mixture of eight clonal plasmids,

each containing a genomic segment of the circulating strain were processed in similar fashion

and sequenced on the same HiSeq flow cell as the appropriate patient derived samples. These

clonally derived samples served as internal controls to improve the accuracy of variant

identification and control for batch effects that confound sequencing experiments.

Identification of iSNV

Intrahost single nucleotide variants were identified in samples that had greater than 10^3

genomes/µl and an average coverage >1000x across the genome. Variants were identified

using DeepSNV and scripts available at https://github.com/lauringlab/variant_pipeline as

described previously (McCrone & Lauring 2016) with a few minor and necessary modifications.

Briefly, reads were aligned to the reference sequence (H3N2 2010-2011 & 2011-2012 :

GenBank CY121496-503, H3N2 2012-2013:GenBank KJ942680-8, H3N2 2014-2015 : Genbank

CY207731-8, H1N1 GenBank : CY121680-8) using Bowtie2 (35). Duplicate reads were then

marked and removed using Picard (http://broadinstitute.github.io/picard/). We identified putative

iSNV using DeepSNV. Bases with phred <30 were masked. Minority iSNV (frequency <50%)

were then filtered for quality using our empirically determined quality thresholds (p-value <0.01

DeepSNV, average mapping quality >30, average Phred >35, average read position between

31 and 94). To control for PCR errors in samples with lower input titers, all isolates with titers

between 10^3 and 10^5 genomes/µl were processed and sequenced in duplicate. Only iSNV that

were found in both replicates were included in down stream analysis. The frequency of the

variant in the replicate with higher coverage at the iSNV location was assigned as the frequency

of the iSNV. Finally, any SNV with a frequency below 2% was discarded.

Given the diversity of the circulating strain in a given season, there were a number of cases in

which isolates contained mutations that were essentially fixed (>95%) relative to the plasmid

control. Often in these cases, the minor allele in the sample matched the major allele in the

plasmid control. We were, therefore, unable to use DeepSNV in estimating the base specific

error rate at this site for these minor alleles and required an alternative means of eliminating

true and false minority iSNV. To this end we applied stringent quality thresholds to these

putative iSNV and implemented a 2% frequency threshold. In order to ensure we were not

introducing a large number of false positive iSNV into our analysis, we performed the following

experiment. Perth (H3N2) samples were sequenced on the same flow cell as both the Perth and

Victoria (H3N2) plasmid controls. Minority iSNV were identified using both plasmid controls. This

allowed us to identify rare iSNV at positions in which the plasmid controls differed both with and

without the error rates provided by DeepSNV. We found that at a frequency threshold of 2% the

methods were nearly identical (NPV of 1, and PPV of 0.94 compared to DeepSNV).

Overview of models for effective population size

We estimated the effective population size using two separate interpretations of a Wright-Fisher

population (Ewens 2004). At its base, the Wright-Fisher model describes the expected changes

in allele frequency of an ideal population, which is characterized by non-overlapping generations,

no migration, no novel mutation, and no population structure. We then asked what size effective

population would make the changes in frequency observed in our dataset most likely. We

calculated these values using two applications of the Wright-Fisher model (i) a diffusion

approximation (Kimura 1955)and (ii) a maximum likelihood approach based on the discrete

interpretation (Williamson & Slatkin 1999).

For these estimates we restricted our analysis to longitudinal samples from a single individual

that were separated by at least 1 day and only used sites that were polymorphic in the initial

sample (29 of the 49 total serial sample pairs). We modeled only the iSNV that were the minor

allele at the first time point, and we assumed a within host generation time of either 6 or 12

hours as proposed by Geoghegan *et al.* (Geoghegan et al. 2016).

Diffusion approximation

The diffusion approximation was first solved by Kimura in 1955 (Kimura 1955). This

approximation to the discrete Wright-Fisher model has enjoyed widespread use in population

genetics as it allows one to treat the random time dependent probability distribution of final allele

frequencies as a continuous function (e.g. (Zanini et al. 2017; Kimura & Ohta 1969; Kimura

1971; Myers et al. 2008)). Here, we also included the limitations in our sensitivity to detect rare

iSNV by integrating over regions of this probability density that were either below our limit of

detection or within ranges where we expect less than perfect sensitivity as follows.

Let $P(p_0, p_t, t \mid N_e)$ be the time dependent probability of a variant drifting from an initial

frequency of p_0 at time 0 p_t at time t generations given an effective population size of N_e where

$0 < p_t < 1$.

The time dependent derivative of this probability has been defined using the Kolmogorov

forward equation (Kimura 1955) and for haploid populations is:

$$P(p_0, p_t, t \mid N_e) = \sum_{i=1}^{\infty} p_0\, q_0\, i(i+1)(2i+1) F(1-i, i+2, 2, p) \times F(1-i, i+2, 2, p_t) e^{-\left[\frac{i(i+1)}{2N_e}\right]t}$$

590   (1)

Where $q = 1 - p$ and F is the hypergeometric function. We approximated the infinite sum by

summing over the first 50 terms. When we added an additional 50 terms (100 in total) we found

no appreciable change in the final log likelihoods.

We denote the frequency of allele that is not observed at the second time point as $p_t \approx 0$ and

the probability of such an event as $P(p_0, p_t \approx 0, t \mid N_e)$. This probability is given in equation 2 as

the sum of the probability that the variant is truly lost by generation t (i.e. the other allele is fixed

$P(q_0, q_t = 1, t \mid N_e)$), the probability that it is present but below the limit of detection (i.e.

$P(p, p_t \approx 0, t \mid 0 < p_t < 0.02, N_e)$) and the probability the variant is not detected due to low

sensitivity for rare variant detection (i.e. $P(p_0, p_t \approx 0, t \mid 0.02 < p_t < 0.1, N_e)$). The probability of

not observing an allele at the second time is then

$$P(p_0, p_t \approx 0, t \mid N_e) = P(q_0, 1, t \mid N_e) + P(p, p_t \approx 0, t \mid 0 < p_t < 0.02, N_e) + \\ P(p_0, p_t \approx 0, t \mid 0.02 < p_t < 0.1, N_e)$$

604
605   (2)

607   The first term in equation 2 is adapted from Kimura, 1955 as

$$P(q_0, 1, t \mid N_e) = q_0 + \sum_{i=1}^{\infty} (2i + 1)p_0 q_0 (-1)^i F(1 - i, i + 2, 2, q_0) e^{-[i(i+1)/2N_e]t}$$

(3)

Where q is defined as above. (Note that this is simply the probability of fixation for a variant at

initial frequency q). As in equation 1 the infinite sum was approximated with a partial sum of 50

terms.

The probability of the allele drifting below our limit of detection can be found by integrating

equation 1 between 0 and our limit of detection, 0.02. This was done numerically using the

python package scipy.

$$P(p, p_t \approx 0, t \mid 0 < p_t < 0.02, N_e) = \int_0^{0.02} P\,(p_0, p_t, t | N_e) dp_t$$

618   (4)

619   Finally, the probability of an iSNV being present at the second time point, but escaping detection,

620   is given by the integral of equation 1 between our benchmarked frequencies (0.02,0.05) times

621   the false negative rate for that range. Here, we assumed the entire range had the same

622   sensitivity as the benchmarked frequency at the lower bound and rounded recipient titers down

623   to the nearest $\log_{10}$ titer (e.g. $10^3, 10^4, 10^5$). We also assumed perfect sensitivity above 10%.

$$P(p_0, p_t \approx 0, t \mid 0.02 < p_t < 0.1, N_e)) = \sum_{f_i}^{[0.02,0.05,0.10)} (\,\text{FNR} \mid \text{Titer}_r, f_i) \int_{f_i}^{f_{i+1}} P\,(p_0, p_t, t \mid N_e) dp_t$$

625   (5)

Where $(\text{FNR} \mid \text{Titer}_r, f_i)$ is the false negative rate given the frequency and the sample titer (See

Supplemental Table 1) and $P(p_0, p_t, t \mid N_e)$ is defined in equation 1.

The log likelihood of an effective population size is the sum of the log of $P(p_0, p_t, t \mid N_e)$ for each

minor allele in the data set, where either the position is polymorphic at time t (i.e. equation 1) or

the allele is not observed at time t (i.e. equation 2).

Discrete Wright-Fisher estimation of N_e

The diffusion approximation treats changes in frequency as a continuous process because it

assumes sufficiently large N_e. That assumption can be relaxed, and the effective population size

can be determined, by applying a maximum likelihood method developed by Williamsom and

Slatkin 1999 (Williamson & Slatkin 1999). In this model, the true allele frequencies move

between discrete states (i.e. the frequency must be of the form i/N_e where i is a whole number

in the range $[0, N_e]$. In the original application, allele counts were used, and sampling error was

added to the model as a binomial distribution with n determined by the sample size. Here, we

use the frequencies available from next generation sequencing and estimate sampling error as

a normal distribution with mean equal to the observed frequency and a standard deviation equal

to that observed in our benchmarking study for the 10^4 genomes/μl samples ($\sigma = 0.014$)

(McCrone & Lauring 2016).

In this model, the probability of observing an allele frequency shift from $\hat{p}_0$ to $\hat{p}_t$ in t generations

provided an effective population of N_e is the probability of observing $\hat{p}_0$ given some initial state

p_0 and the probability of the population having that state, times the probability of observing $\hat{p}_t$

given some final state p_t and the probability of moving from the initial to the final state summed

across all possible states.

$$P(p_0, p_t \mid N_e) = \sum_{}^{p_0, p_t} P\left(\hat{p}_0 \mid p_0\right) P(p_0 \mid N_e) P(\hat{p}_t \mid p_t) P(p_t \mid p_0, N_e)$$

651   (6)

653   Where $\hat{p}_x$ are the observed probabilities and p_x are the real ones (of the form i/N_e discussed

654   above). The likelihood of observing a given frequency $\hat{p}_x$ given a defined state p_x is given by the

655   likelihood of drawing $\hat{p}_x$ from a normal distribution with mean p_x and standard deviation 0.014.

$$P(\hat{p}_x \mid p_x) = \text{Norm}(p_x, 0.014)$$

657   (7)

658   As in Williamson and Slatkin 1999, we assume a uniform prior on the initial state. Because we

659   know that our specificity is near perfect (above 2%, Supplemental Table 1) and we restrict our

660   analysis to only polymorphic sites, the probability of any initial state is given by

$$P(p_0 \mid N_e) = \frac{1}{N_e - 1}$$

662   (8)

664   and finally the probability of moving from one state to another in t generations is given by

$$P(p_t, p_0 \mid N_e) = v_0 M^t v_t$$

(9)

Where M is a square transmission matrix with $N_e + 1$ rows and columns. Where $m_{i,j}$ is the

probability of going from the ith configuration to the jth or the probability of drawing $j - 1$ out of

binomial distribution with mean $(i - 1)/N_e$ and a sample size N_e. v_0 is a row vector of initial

frequencies p_0 with 100% chance of initial state p_0, and v_t is column vector of the frequencies at

time point t with 100% chance of the final state. In other words v_0 is a row vector of $N_e + 1$

states with 0 everywhere except in the ith position where $\frac{i-1}{N_e} = p_0$, and v_t is a column vector of

$N_e + 1$ states with 0 everywhere except the jth position where $\frac{j-1}{N_e} = p_t$

Using the scalar and cumulative properties of matrix multiplication equation 6 reduces to

$$P(p_0, p_t \mid N_e) = [0, P(p_0 \mid q_{0_2})P(q_{0_2} \mid N_e), \ldots, P(p_0 \mid q_{0_{N_e-1}})P(q_{0_{N_e-1}} \mid N_e), 0]M^t \begin{bmatrix} P(p_t \mid q_{t_1}) \\ \vdots \\ P(p_t \mid q_{t_{N_e}}) \end{bmatrix}$$

(10)

The first and last entries in v_0 are 0 because we assume all measured sites represent

polymorphisms at the first time of sampling. As above, the log likelihood of a given population

size is then simply the sum of the log of $P(\hat{p}_0, \hat{p}_t, t \mid N_e)$ for each minor allele in the data set.

Simulations

To simulate within host evolution we set N_e in equation 10 to either 30, 50 or 100. For each

minor allele we used the closest available non-zero state given the effective population size as

the starting state. We then calculated the probability of moving to any other state and selected a

final state from this distribution. We then drew a final measured frequency from the normal

distribution to account for measurement errors.

ABC model

We estimated both the effective population size and selection coefficients using the approximate

Bayesian computation (ABC) described in (Foll et al. 2014) with the WFACB_v1.1 software

provided in (Foll et al. 2015). In its current implementation, this analysis requires the same time

points for each sample, and we restricted this analysis to longitudinal samples taken 1 day apart.

This subset constitutes 16 of the 29 modeled longitudinal samples. Briefly, we subsampled

polymorphic sites to 1,000x coverage to estimate allele counts from frequency data as in (Foll et

al. 2014). We then estimated the prior distribution of the effective population size using 10,000

bootstrap replicates. We selected a uniform distribution on the range [-0.5,0.5] as the prior

distribution for the selection coefficients. The posterior distributions were determined from

accepting the top 0.01% of 100,000 simulations.

Overview of models used for estimating the transmission bottleneck

We model transmission as a simple binomial sampling process (Sobel Leonard et al. 2017). In

our first model, we assume any transmitted iSNV, no matter the frequency, will be detected in

the recipient. In the second, we relax this assumption and account for false negative iSNV in the

recipient. To include the variance in the transmission bottlenecks between pairs we use

maximum likelihood optimization to fit the average bottleneck size assuming the distribution

follows a zero-truncated Poisson distribution.

Presence/Absence model

The presence/absence model makes several assumptions. We assume perfect detection of all

transmitted iSNV in the recipient. For each donor iSNV, we measure only whether or not the

variant is present in the recipient. Any iSNV that is not found in the recipient is assumed to have

not been transmitted. We also assume the probability of transmission is determined only by the

frequency of the iSNV in the donor at the time of sampling (regardless of how much time passes

between sampling and transmission). The probability of transmission is simply the probability

that the iSNV is included at least once in a sample size equal to the bottleneck. Finally, we

assume all genomic sites are independent of one another. For this reason, we discarded the

one case where the donor was likely infected by two strains, as the iSNV were certainly linked.

In our within host models, we only tracked minor alleles as in our data set we only ever find 2

alleles at each polymorphic site. In this case, the frequency of the major allele is simply one

minus the frequency of the minor allele. Because the presence/absence model is unaware of

the frequency of alleles in the recipient we must track both alleles at each donor polymorphic

site.

Let A_1 and A_2 be alleles in donor j at genomic site i. Let $P(A_1)$ be the probability that A_1 is the

only transmitted allele. There are three possible outcomes for each site. Either only A_1 is

transmitted, only A_2 is transmitted, or both A_1 and A_2 are transmitted. The probability of only A_1

being transmitted given a bottleneck size of N_b is

$$P_{i,j}(A_1 \mid N_b) = p_1^{N_b}$$

730
(11)

731   where p_1 is the frequency of A_1 in the donor. In other words, this is simply the probability of only

732   drawing A_1 in N_b draws. The probability that only A_2 is transmitted is similarly defined.

734   The probability of both alleles being transmitted is given by

$$P_{i,j}(A_1, A_2 \mid N_b) = 1 - (p_1^{N_b} + p_2^{N_b})$$

736
(12)

where p_1 and p_2 are the frequencies of the alleles respectively. This is simply the probability of

not picking only A_1 or only A_2 in N_b draws.

This system could easily be extended to cases where there are more than 2 alleles present at a

site; however, that never occurs in our data set.

For ease we will denote the likelihood of observing the data at a polymorphic site i in each

donor j given the bottleneck size N_b as $P_{i,j}(N_b)$ where $P_{i,j}(N_b)$ is defined by equation 11 if only

one allele is transmitted and equation 12 if two alleles are transmitted.

The log likelihood of a bottleneck of size N_b is given by

$$LL(N_b) = \sum_j \sum_i \text{Ln}(P_{i,j})$$

749
(13)

where i,j refers to the ith polymorphic site in the jth donor. This is the log of the probability of

observing the data summed over all polymorphic sites across all donors.

Because the bottleneck size is likely to vary across transmission events, we used maximum

likelihood to fit the bottleneck distribution as oppose to fitting a single bottleneck value. Under

this model we assumed the bottlenecks were distributed according to a zero-truncated Poisson

distribution parameterized by λ. The likelihood of observing the data given a polymorphic site i

in donor j and λ is

$$P_{i,j}(\lambda) = \sum_{N_b=1}^{\infty} P_{i,j}(N_b)P(N_b \mid \lambda)$$

759   (14)

where $P_{i,j}(N_b)$ is defined as above, $P(N_b \mid \lambda)$ is the probability of drawing a bottleneck of size N_b

from a zero-truncated Poisson distribution with a mean of $\frac{\lambda}{1-e^{-\lambda}}$. The sum is across all possible

N_b defined on $[1, \infty]$. For practical purposes, we only investigated bottleneck sizes up to 100, as

initial analyses suggested λ is quite small and the probability of drawing a bottleneck size of 100

from a zero-truncated Poisson distribution with $\lambda = 10$ is negligible. We follow this convention

whenever this sum appears.

The log likelihood of λ for the data set is given by

$$LL(\lambda) = \sum_{j}\sum_{i} \text{Ln}\left(\sum_{N_b=1}^{\infty} P_{i,j}(N_b)P(N_b \mid \lambda)\right)$$

770   (15)

Beta Binomial model

The Beta binomial model is explained in detail in Leonard *et al.* (Sobel Leonard et al. 2017). It is

similar to the presence/absence model in that transmission is modeled as a simple sampling

process; however, it relaxes the following assumptions. In this model, the frequencies of

transmitted variants are allowed to change between transmission and sampling according a

beta distribution. The distribution is not dependent on the amount of time that passes between

transmission and sampling, but rather depends on the size of the founding population (here

assumed to equal to N_b) and the number of variant genomes present in founding population k.

Note the frequency in the donor is assumed to be the same between sampling and transmission.

The equations below are very similar to those presented by Leonard *et al.* with one exception.

Because we know the sensitivity of our method to detect rare variants based on the expected

frequency and the titer, we can include the possibility that iSNV are transmitted but are missed

due to poor sensitivity. Because the beta binomial model is aware of the frequency of the iSNV

in the recipient, no information is added by tracking both alleles at a genomic site i.

Let p_{i,j_d} represent the frequency of the minor allele frequency at position i in the donor of some

transmission pair j. Similarly, let p_{i,j_r} be the frequency of that same allele in the recipient of the

jth transmission pair. Then, as in Leonard *et al.*, the likelihood of some bottleneck N_b for the

data at site i in pair j where the minor allele is transmitted is given by

$$L(N_b)_{i,j} = \sum_{k=1}^{N_b} \mathrm{p\_beta}\left(p_{i,j_r} \mid k, N_b - k\right) \mathrm{p\_bin}(k \mid N_b, p_{i,j_d})$$

792   (16)

Where p_beta is the probability density function for the beta distribution and p_bin is the

probability mass function for the binomial distribution.

This is the probability density that the transmitted allele is found in the recipient at a frequency

of p_{i,j_r} given that the variant was in k genomes in a founding population of size N_b times the

probability of drawing k variant genomes in a sample size of N_b and a variant frequency of p_{i,j_d}.

This is then summed for all possible k where $1 \leq k \leq N_b$.

As in equation 14 the likelihood of a zero truncated Poisson with a mean of $\frac{\lambda}{1-e^{-\lambda}}$ given this

transmitted variants is then given by

$$L(\lambda)_{i,j}^{\text{transmitted}} = \sum_{N_b=1}^{\infty} L\,(N_b)_{i,j}\,P(N_b \mid \lambda)$$

(17)

This is simply the likelihood of each N_b weighted by the probability of drawing a bottleneck size

of N_b from bottleneck distribution.

In this model, there are three possible mechanisms for a donor iSNV to not be detected in the

recipient. (i) The variant was not transmitted. (ii) The variant was transmitted but is present

below our level of detection (2%). (iii) The variant was transmitted and present above our level

of detection but represents a false negative in iSNV identification.

As in Leonard *et al.,* the likelihood of scenarios (i) and (ii) for a given N_b are expressed as

$$L(N_b)_{i,j}^{\text{lost}} = \sum_{k=0}^{N_b} \text{p\_beta\_cdf}(p_{i,j_r} < 0.02 \mid k, N_b - k)\,\text{p\_bin}(k \mid N_b, p_{i,j_d})$$

(18)

Where p_beta_cdf is the cumulative distribution function for the beta distribution. Note that if

$k = 0$ (i.e. the iSNV was not transmitted) then the term reduces to the probability of not drawing

the variant in N_b draws.

The likelihood of the variant being transmitted but not detected in the recipient given a

bottleneck of N_b is described by

$$L(N_b)_{i,j}^{\text{missed}} = \sum_{k=0}^{N_b} \sum_{f_e}^{[0.02,0.05,0.1)} \text{p\_beta\_cdf}(f_e < p_{i,j_r} < f_{e+1} \mid k, N_b - k) \times$$

826
$$\text{p\_bin}(k \mid N_b, p_{i,j_d})(\text{FNR} \mid \text{Titer}_r, f_e)$$
(19)

This is the likelihood of the variant existing in the ranges [0.0.2,0.05] or [0.05,0.1] given an initial

frequency of k/N_b and a bottleneck size of N_b multiplied by the expected false negative rate

(FNR) given the titer of the recipient and the lower frequency bound. As in our diffusion model,

we assumed perfect sensitivity for detection of iSNV present above 10%, rounded recipient

titers down to the nearest $\log_{10}$ titer (e.g. $10^3, 10^4, 10^5$) and assumed the entire range $[f_e, f_{e+1}]$

has the same sensitivity as the lower bound.

The likelihood of λ for iSNV that are not observed in the recipient is then given by summing

equations 18 and 19 across all possible N_b.

$$L(\lambda)_{i,j}^{\text{nontransmitted}} = \sum_{N_b=1}^{\infty} (L(N_b)_{i,j}^{\text{lost}} + L(N_b)_{i,j}^{\text{missed}})P(N_b \mid \lambda)$$

(20)

The log likelihood of the total dataset is then determined by summing log of equations 17 and 20

(as applicable) across all polymorphic sites in each donor. (As before here we sum of N_b within

the range $[1,100]$.)

Simulation

In order evaluate the fits of the two transmission models, we simulated whether or not each

donor iSNV was transmitted or not. This involved converting each model to a presence absence

model. In each simulation, we assigned a bottleneck from the bottleneck distribution for each

transmission pair. We then determined the probability of only transmitting one allele (A_x where

$x \in [1,2]$ as in the presence/absence model above) and the probability of transmitted both

alleles (A_1, A_2 above) for each polymorphic site.

For the presence/absence model, the probabilities for each possible outcome are given by

equations 11 and 12. For the beta binomial model, the probability of only observing A_x at site i

is given by

$$P(A_x \mid N_b) = L(N_b)_{i,j}^{\mathrm{lost}} + L(N_b)_{i,j}^{\mathrm{missed}}$$

856   (21)

858   where $L(N_b)_{i,j}^{\mathrm{lost}}$ and $L(N_b)_{i,j}^{\mathrm{missed}}$ are defined as in equations 18 and 19 respectively, but with

859   p_{i,j_d} replaced by $1 - p_{i,j_d}$. This is simply the probability of not observing the other allele in the

860   recipient.

862   Again, the probability of observing both alleles is

$$P(A_1, A_2 \mid N_b) = 1 - (P(A_1) + P(A_2))$$

864
<div style="text-align: right;">(22)</div>

where $P(A_1)$ and $P(A_2)$ are defined as in equation 21.

Fitting mutation rate and N_e

The diffusion approximation to the Wright - Fischer model allows us to make predictions on the

allele frequency spectrum of a population given a mutation rate and an effective population size.

The probability of observing a mutation at frequency p_t given an initial frequency of 0 can be

approximated as in (Rouzine et al. 2001)

$$P(0, p_t, t, \mid \mu, N_e) = \frac{2\mu N_e}{p_t} e^{-\frac{2N_e p_t}{t}}$$

873
<div style="text-align: right;">(23)</div>

Where μ is the mutation rate. In this model mutation increases an allele's frequency from 0 but

after that initial jump, drift is responsible for allowing the mutation to reach it's observed

frequency. Because the limit of equation 23 approaches infinity as p_t approaches 0 and for ease

in numerical integration, we assumed that any variant present at less than 0.1% was essentially

at 0%.

We then assumed each infection began as a clonal infection matching the consensus sequence

observed at the time of sampling. The likelihood of observing minor alleles at the observed

frequency is the given by equation 23.

As in the other within host models, we can account for nonpolymorphic sites by adding the

likelihood that no mutation is present $P(0, p_t \approx 0, t \mid p_t < 0.001, \mu, N_e)$, that a mutation is present

but below our level of detection $P(0, p_t \approx 0, t \mid p_t < 0.02, \mu, N_e)$, and that a mutation is present

but missed due to low sensitivity at low frequencies $P(0, p_t \approx 0, t \mid 0.02 < p_t < 0.1, \mu, N_e)$. In this

model we assumed 13133 mutagenic targets in each sample (the number of coding sites

present in the reference strain from 2014-2015).

The probability of not observing a mutation is given by

$$P(\,0, p_t \approx 0, t, \mid \mu, N_e\,) = P(\,0, p_t \approx 0, t \mid p_t < 0.001, \mu, N_e\,) +$$
$$\text{893} \qquad P(\,0, p_t \approx 0, t \mid p_t < 0.02, \mu, N_e\,) +$$
$$\text{894} \qquad P(0, p_t \approx 0, t \mid 0.02 < p_t < 0.1, \mu, N_e)$$

895   (24)

896   Where

$$P(0, p_t \approx 0, t \mid p_t < 0.001, \mu, N_e) = 1 - \int_{0.001}^{1} P(0, p_t, t, \mid \mu, N_e) dp_t$$

897   (25)

898   and

$$P(0, p_t \approx 0, t \mid p_t < 0.02, \mu, N_e) = \int_{0.001}^{0.02} P(0, p_t, t, \mid \mu, N_e) dp_t$$

899   (26)

900   and

$$P(0, p_t \approx 0, t \mid 0.02 < p_t < 0.1, \mu, N_e) = \sum_{f_i}^{[0.02, 0.05, 0.10)} (\text{FNR} \mid \text{Titer}_r, f_i) \int_{f_i}^{f_{i+1}} P(p_0, p_t, t \mid \mu, N_e) dp_t$$

901   (27)

Where we follow the same convention as in equation 5 for determining the false negative rate.

The log likelihood of a given μ and N_e pair is then the sum of the log of equations 23 and 24 for

all possible sites in the data set.

Annotated computer code for all analyses and for generating the figures can be accessed at

https://github.com/lauringlab/Host_level_IAV_evolution

Acknowledgments

This work was supported by a Clinician Scientist Development Award from the Doris Duke

Charitable Foundation (CSDA 2013105), a University of Michigan Discovery Grant, and R01

AI118886, all to ASL. The HIVE cohort was supported by NIH R01 AI097150 and CDC U01

IP00474 to ASM. JTM was supported by the Michigan Predoctoral Training Program in Genetics

(T32GM007544). RJW was supported by K08AI119182. We thank Alexey Kondrashov and

Aaron King for helpful discussion.

References

Alizon, S., Luciani, F. & Regoes, R.R., 2011. Epidemiological and clinical consequences of
within-host evolution. *Trends in microbiology*, 19(1), pp.24–32.

ARCHETTI, I. & HORSFALL, F.L., 1950. Persistent antigenic variation of influenza A viruses
after incomplete neutralization in ovo with heterologous immune serum. *The Journal of
experimental medicine*, 92(5), pp.441–462.

Caton, A.J. et al., 1982. The antigenic structure of the influenza virus A/PR/8/34 hemagglutinin
(H1 subtype). *Cell*, 31(2 Pt 1), pp.417–427.

Debbink, K. et al., 2017. Vaccination has minimal impact on the intrahost diversity of H3N2
influenza viruses. *PLoS pathogens*, 13(1), p.e1006194.

Dinis, J.M. et al., 2016. Deep Sequencing Reveals Potential Antigenic Variants at Low
Frequencies in Influenza A Virus-Infected Humans S. Schultz-Cherry, ed. *Journal of
Virology*, 90(7), pp.3355–3365.

Doud, M.B., Hensley, S.E. & Bloom, J.D., 2017. Complete mapping of viral escape from
neutralizing antibodies A. S. Lauring, ed. *PLoS pathogens*, 13(3), pp.e1006271–20.

Ewens, W.J., 2004. *Mathematical Population Genetics 1*, Springer Science & Business Media.

Flannery, B. et al., 2016. Enhanced Genetic Characterization of Influenza A(H3N2) Viruses and
Vaccine Effectiveness by Genetic Group, 2014-2015. *The Journal of Infectious Diseases*,
214(7), pp.1010–1019.

Foll, M. et al., 2014. Influenza virus drug resistance: a time-sampled population genetics
perspective. *PLoS genetics*, 10(2), p.e1004185.

Foll, M., Shim, H. & Jensen, J.D., 2015. WFABC: a Wright-Fisher ABC-based approach for
inferring effective population sizes and selection coefficients from time-sampled data.
Molecular ecology resources, 15(1), pp.87–98.

Geoghegan, J.L., Senior, A.M. & Holmes, E.C., 2016. Pathogen population bottlenecks and
adaptive landscapes: overcoming the barriers to disease emergence. *Proceedings.*
Biological sciences / The Royal Society, 283(1837), pp.20160727–9.

Gerstung, M. et al., 2012. Reliable detection of subclonal single-nucleotide variants in tumour
cell populations. *Nature Communications*, 3, p.811.

Ghedin, E. et al., 2011. Deep sequencing reveals mixed infection with 2009 pandemic influenza
A (H1N1) virus strains and the emergence of oseltamivir resistance. *The Journal of*
Infectious Diseases, 203(2), pp.168–174.

Gubareva, L.V. et al., 2001. Selection of influenza virus mutants in experimentally infected
volunteers treated with oseltamivir. *The Journal of Infectious Diseases*, 183(4), pp.523–531.

Herfst, S. et al., 2012. Airborne transmission of influenza A/H5N1 virus between ferrets. *Science*
(New York, NY), 336(6088), pp.1534–1541.

Holmes, E.C., 2009. RNA virus genomics: a world of possibilities. *The Journal of clinical*
investigation, 119(9), pp.2488–2495.

Hughes, J. et al., 2012. Transmission of equine influenza virus during an outbreak is
characterized by frequent mixed infections and loose transmission bottlenecks. *PLoS*
pathogens, 8(12), p.e1003081.

Iqbal, M. et al., 2009. Within-host variation of avian influenza viruses. *Philosophical transactions*
of the Royal Society of London Series B, Biological sciences, 364(1530), pp.2739–2747.

Kao, R.R. et al., 2014. Supersize me: how whole-genomesequencing and big data
aretransforming epidemiology. *Trends in microbiology*, pp.1–10.

Kimura, M., 1955. SOLUTION OF A PROCESS OF RANDOM GENETIC DRIFT WITH A
CONTINUOUS MODEL. *Proceedings of the National Academy of Sciences of the United*
States of America, 41(3), pp.144–150.

Kimura, M., 1971. Theoretical foundation of population genetics at the molecular level.
Theoretical population biology, 2(2), pp.174–208.

Kimura, M. & Ohta, T., 1969. The Average Number of Generations until Fixation of a Mutant
Gene in a Finite Population. *Genetics*, 61(3), pp.763–771.

Koel, B.F. et al., 2013. Substitutions near the receptor binding site determine major antigenic
change during influenza virus evolution. *Science (New York, NY)*, 342(6161), pp.976–979.

Kouyos, R.D., Althaus, C.L. & Bonhoeffer, S., 2006. Stochastic or deterministic: what is the

effective population size of HIV-1? *Trends in microbiology*, 14(12), pp.507–511.

Kugelman, J.R. et al., 2017. Error baseline rates of five sample preparation methods used to
characterize RNA virus populations K. K. Tee, ed., 12(2), pp.e0171333–13. Available at:
http://dx.plos.org/10.1371/journal.pone.0171333.

Lee, M.-S. & Chen, J.S.-E., 2004. Predicting antigenic variants of influenza A/H3N2 viruses.
Emerging infectious diseases, 10(8), pp.1385–1390.

McCrone, J.T. & Lauring, A.S., 2016. Measurements of Intrahost Viral Diversity Are Extremely
Sensitive to Systematic Errors in Variant Calling. *Journal of Virology*, 90(15), pp.6884–6895.

Monto, A.S. et al., 2014. Frequency of acute respiratory illnesses and circulation of respiratory
viruses in households with children over 3 surveillance seasons. *The Journal of Infectious
Diseases*, 210(11), pp.1792–1799.

Murcia, P.R. et al., 2010. Intra- and interhost evolutionary dynamics of equine influenza virus.
Journal of Virology, 84(14), pp.6943–6954.

Myers, S., Fefferman, C. & Patterson, N., 2008. Can one learn history from the allelic spectrum?
Theoretical population biology, 73(3), pp.342–348.

Neher, R.A. & Bedford, T., 2015. nextflu: real-time tracking of seasonal influenza virus evolution
in humans. *Bioinformatics (Oxford, England)*, p.btv381.

Nelson, M.I. & Holmes, E.C., 2007. The evolution of epidemic influenza. *Nature Reviews
Genetics*, 8(3), pp.196–205.

Nelson, M.I. et al., 2006. Stochastic processes are key determinants of short-term evolution in
influenza a virus. *PLoS pathogens*, 2(12), p.e125.

Ohmit, S.E. et al., 2015. Influenza vaccine effectiveness in households with children during the
2012-2013 season: assessments of prior vaccination and serologic susceptibility. *The
Journal of Infectious Diseases*, 211(10), pp.1519–1528.

Ohmit, S.E. et al., 2013. Influenza Vaccine Effectiveness in the Community and the Household.
*Clinical infectious diseases : an official publication of the Infectious Diseases Society of
America*, 56(10), pp.1363–1369.

Ohmit, S.E. et al., 2016. Substantial Influenza Vaccine Effectiveness in Households With
Children During the 2013-2014 Influenza Season, When 2009 Pandemic Influenza A(H1N1)
Virus Predominated. *The Journal of Infectious Diseases*, 213(8), pp.1229–1236.

Pauly, M.D., Procario, M.C. & Lauring, A.S., 2017. A novel twelve class fluctuation test reveals
higher than expected mutation rates for influenza A viruses. *eLife*, 6, p.e26437.

Peck, K.M., Chan, C.H.S. & Tanaka, M.M., 2015. Connecting within-host dynamics to the rate of
viral molecular evolution. *Virus Evolution*, 1(1), p.vev013.

Petrie, J.G. et al., 2017. Application of an Individual-Based Transmission Hazard Model for
Estimation of Influenza Vaccine Effectiveness in a Household Cohort. *American journal of*

epidemiology.

Petrie, J.G. et al., 2013. Influenza Transmission in a Cohort of Households with Children: 2010-
2011 J. McVernon, ed. *PLoS ONE,* 8(9), p.e75339.

Petrova, V.N. & Russell, C.A., 2017. The evolution of seasonal influenza viruses. *Nature*
Reviews Microbiology, 2, p.517.

Poon, L.L.M. et al., 2016. Quantifying influenza virus diversity and transmission in humans.
Nature Genetics, 48(2), pp.195–200.

Rambaut, A. et al., 2008. The genomic and epidemiological dynamics of human influenza A
virus. *Nature*, 453(7195), pp.615–619.

Rogers, M.B. et al., 2015. Intrahost dynamics of antiviral resistance in influenza A virus reflect
complex patterns of segment linkage, reassortment, and natural selection. *mBio*, 6(2).

Rouzine, I.M., Rodrigo, A. & Coffin, J.M., 2001. Transition between stochastic evolution and
deterministic evolution in the presence of selection: general theory and application to
virology. *Microbiology and molecular biology reviews : MMBR*, 65(1), pp.151–185.

Russell, C.A. et al., 2012. The potential for respiratory droplet-transmissible A/H5N1 influenza
virus to evolve in a mammalian host. *Science (New York, NY)*, 336(6088), pp.1541–1547.

Sanjuán, R. et al., 2010. Viral mutation rates. *Journal of Virology*, 84(19), pp.9733–9748.

Smith, D.J. et al., 2004. Mapping the antigenic and genetic evolution of influenza virus. *Science*
(New York, NY), 305(5682), pp.371–376.

Sobel Leonard, A. et al., 2016. Deep Sequencing of Influenza A Virus from a Human Challenge
Study Reveals a Selective Bottleneck and Only Limited Intrahost Genetic Diversification D.
S. Lyles, ed. *Journal of Virology*, 90(24), pp.11247–11258.

Sobel Leonard, A. et al., 2017. Transmission Bottleneck Size Estimation from Pathogen Deep-
Sequencing Data, with an Application to Human Influenza A Virus. *Journal of Virology*,
91(14).

Varble, A. et al., 2014. Influenza A virus transmission bottlenecks are defined by infection route
and recipient host. *Cell Host and Microbe*, 16(5), pp.691–700.

Visher, E. et al., 2016. The Mutational Robustness of Influenza A Virus N. M. Ferguson, ed.
PLoS pathogens, 12(8), pp.e1005856–25.

Wiley, D.C., Wilson, I.A. & Skehel, J.J., 1981. Structural identification of the antibody-binding
sites of Hong Kong influenza haemagglutinin and their involvement in antigenic variation.
Nature, 289(5796), pp.373–378.

Wilker, P.R. et al., 2013. Selection on haemagglutinin imposes a bottleneck during mammalian
transmission of reassortant H5N1 influenza viruses. *Nature Communications*, 4, p.2636.

Williamson, E.G. & Slatkin, M., 1999. Using maximum likelihood to estimate population size
from temporal changes in allele frequencies. *Genetics*, 152(2), pp.755–761.

Xu, R. et al., 2010. Structural basis of preexisting immunity to the 2009 H1N1 pandemic
influenza virus. *Science (New York, NY)*, 328(5976), pp.357–360.

Xue, K.S. et al., 2017. Parallel evolution of influenza across multiple spatiotemporal scales.
eLife, 6, p.e26875.

Zanini, F. et al., 2017. In vivo mutation rates and the landscape of fitness costs of HIV-1. *Virus*
Evolution, 3(1), p.vex003.

Zhou, B. et al., 2009. Single-reaction genomic amplification accelerates sequencing and vaccine
production for classical and Swine origin human influenza a viruses. *Journal of Virology*,
83(19), pp.10309–10313.

Zwart, M.P. & Elena, S.F., 2015. Matters of Size: Genetic Bottlenecks in Virus Infection and
Their Potential Impact on Evolution. *Annual Review of Virology*, 2(1), pp.161–179.

1056

1057

 Figure Legends

 Figure 1. Within host diversity of IAV populations. (A) Boxplots (median, 25th and 75th

 percentiles, whiskers extend to most extreme point within median ± 1.5 x IQR) of the number of

 viral genomes per microliter transport media stratified by day post symptom onset. Notches

 represent the approximate 95% confidence interval of the median. (B) Boxplots (median, 25th

 and 75th percentiles, whiskers extend to most extreme point within median ± 1.5 x IQR) of the

 number of iSNV in 249 high quality samples stratified by day post symptom onset. (C)

 Histogram of within host iSNV frequency in 249 high quality samples. Bin width is 0.05

 beginning at 0.02. Mutations are colored nonsynonymous (blue) and synonymous (gold) (D)

 Location of all identified iSNV in the influenza A genome. Mutations are colored

 nonsynonymous (blue) and synonymous (gold) relative to that sample's consensus sequence.

 Triangles signify mutations that were found in more than one individual in a given season.

 Figure 2. Within host dynamics of IAV. (A) Timing of sample collection for 35 paired longitudinal

 samples relative to day of symptom onset. Of the 49 total, 35 pairs had minor iSNV present in

 the first sample. (B) The change in frequency over time for minority nonsynonymous (blue) and

 synonymous (gold) iSNV identified for the paired samples in (A). (C) The distribution of effective

 population sizes estimated from 1,000 simulated populations. Simulations were run on

 populations with characteristics similar to the actual patient-derived populations and with the

 specified effective population size (x-axis). (D) The effect of iteratively removing iSNV with the

 most extreme change in frequency (fraction of iSNV removed, x-axis) on the estimated effective

 population size. The point represents the estimate when all iSNV are included. (E) The posterior

 distributions of selection coefficients estimated for the 35 iSNV present in isolates sampled one

 day apart. Distributions are colored according to class relative to the sample consensus

 sequence, nonsynonymous (blue) synonymous (gold). Variants for which the 95% highest

1083   posterior density intervals exclude 0.0 are noted in the margin.

1084

Figure 3. Between host dynamics of IAV. (A) The distribution of pairwise L1-norm distances for

household (blue) and randomly-assigned community (gold) pairs. The bar heights are

normalized to the height of the highest bar for each given subset (47 for household, 1,592 for

community). The red line represents the 5^{th} percentile of the community distribution. (B) Timing

of symptom onset for 52 epidemiologically linked transmission pairs. Day of symptom onset for

both donor and recipient individuals is indicated by black dots. Dashed lines represent pairs that

were removed due to abnormally high genetic distance between isolates, see (A). (C) The

frequency of donor iSNV in both donor and recipient samples. Frequencies below 2% and

above 98% were set to 0% and 100% respectively. (D) The presence-absence model fit

compared with the observed data. The x-axis represents the frequency of donor iSNV with

transmitted iSNV plotted along the top and nontransmitted iSNV plotted along the bottom. The

black line indicates the probability of transmission for a given iSNV frequency as determined by

logistic regression. Similar fits were calculated for 1,000 simulations with a mean bottleneck size

of 1.66. Fifty percent of simulated outcomes lie in the darkly shaded region and 95% lie in the

lightly shaded regions. (E) The outcome from 1,000 simulated "transmission" events with

randomly assigned pairings. The black line represents the observed data, as in (D) the shaded

regions represent the middle 50% and 95% of simulated outcomes. The results from the

simulated logit models were smoothed by plotting the predicted probability of transmission at

0.02 intervals. (F) The beta-binomial model fit. Similar to (D) except the simulated outcomes are

the based on a beta-binomial model using a mean bottleneck of 1.73.

1105

1106   **Figure 4.** Combined estimates of within host mutation rate and effective population size.

1107   Contour plot shows the log likelihood surface for estimates of the effective population size and

1108   neutral mutation rate. The point represents the peak ($\mu = 4\times10^{-6}$, $N_e = 36$, log likelihood = -

1109   4,687). Log likelihoods for each contour are indicated.

1110

Supplementary Figure 1. Sequence coverage for all samples. For each sample, the sliding

window mean coverage was calculated using a window size of 200 and a step of 100. The

distributions of these means are plotted as box plots (median, 25^{th} and 75^{th} percentiles,

whiskers extend to most extreme point within median ± 1.5 x IQR) where the y-axis represents

the read depth and the x-axis indicates the position of the window in a concatenated IAV

genome.

1117

1118   **Supplementary Figure 2.** Approximate maximum likelihood trees of the concatenated coding

1119   sequences for high quality H1N1 samples. The branches are colored by season; the tip

1120   identifiers are colored by household. Arrows with numbers indicate consensus and putative

1121   minor haplotypes for samples with greater than 10 iSNV. Trees were made using FastTree.

1122

1123   **Supplementary Figure 3.** Approximate maximum likelihood trees of the concatenated coding

1124   sequences for high quality H3N2 samples. The branches are colored by season; the tip

1125   identifiers are colored by household. Arrows with numbers indicate consensus and putative

1126   minor haplotypes for samples with greater than 10 iSNV. Trees were made using FastTree.

1127

1128   **Supplementary Figure 4.** The effect of titer and vaccination on the number of iSNV identified.

1129   (A) The number of iSNV identified in an isolate (y-axis) plotted against the titer (x-axis,

1130   genomes/µl transport media). (B) The number of iSNV identified in each isolate stratified by

1131   whether that individual was vaccinated or not. Red bars indicate the median of each distribution.

1132

1133   **Supplementary Figure 5.** Minority nonsynonymous iSNV in global circulation.

1134   The global frequencies of the amino acids that were found as minority variants in sample

1135   isolates (x-axis) plotted overtime (y-axis). Each amino acid trace is labeled according to the H3

1136   number scheme. All samples were isolated in December of 2014 (gray line).

1137

1138   **Supplementary Figure 6.** Reproducibility of iSNV identification for paired samples acquired on

1139   the same day. The x-axis represents iSNV frequencies found in the home-acquired nasal swab.

1140   The y-axis represents iSNV frequencies found the clinic-acquired combined throat and nasal

1141   swab.

1142

Supplementary Figure 7. Estimate of effective bottleneck size with relaxed variant calling

criteria. (A) The frequency of iSNV in both recipient and donor isolates. iSNV were identified

using the original variant calling pipeline. (B) The presence-absence model fit compared to the

observed data for iSNV identified using the original variant calling pipeline. The x-axis

represents the frequency of donor iSNV with transmitted iSNV plotted along the top and

nontransmitted iSNV plotted along the bottom. The black line indicates the probability of

transmission for a given iSNV frequency as determined by logistic regression. Similar fits were

calculated for 1,000 simulations with a mean bottleneck size of 2.10. Fifty percent of simulated

outcomes lie in the darkly shaded region and 95% lie in the lightly shaded regions. (C) Similar to

(A) but with minority iSNV identified using the current analytical framework without a frequency

threshold. (D) Similar to B but with minority iSNV identified using the current analytical

framework without a frequency threshold.

1155

1156 **Table 1. Influenza viruses over five seasons in a household cohort**

	2010-2011	2011-2012	2012-2013	2013-2014	2014-2015
Households	328	213	321	232	340
Participants	1441	943	1426	1049	1431
Vaccinated, n (%)[a]	934 (65)	554 (59)	942 (66)	722 (69)	992 (69)
IAV Positive Individuals[b]	86	23	69	48	166
H1N1	26	1	3	47	0
H3N2	58	22	66	1	166
IAV Positive Households[c]					
Two individuals	13	2	9	7	23
Three individuals	5	2	3	3	11
Four individuals	-	-	1	2	4
High Quality NGS Pairs[d]	4	1	2	6	39

1157
[a] Self reported or confirmed receipt of vaccine prior to the specified season.
[b] RT-PCR confirmed infection.
[c] Households in which two individuals were positive within 7 days of each other. In cases of trios and quartets, the
putative chains could have no pair with onset >7 days apart.
[d] Samples with >10^3 genome copies per µl of transport medium, adequate amplification of all 8 genomic segments,
and average sequencing coverage >10^3 per nucleotide.
1164

Table 2. Within host effective population size of IAV

Model	SNV Used	Generation Time (h)	Effective Population Size (95% CI)
Diffusion approximation	All	6	35 (26-46)
	All	12	17 (13-23)
Discrete model	All	6	32 (28-41)
	Nonsynonymous	6	30 (21-40)
	Synonymous	6	37 (27-54)
	All	12	23 (23-29)
	Nonsynonymous	12	19 (19-21)
	Synonymous	12	27 (22-33)

 Supplementary Table 1. Sensitivity and specificity of variant detection

Copy Number[a]	Variant Frequency	Original Pipeline[b]		Current Pipeline[c]	
		Sensitivity	Specificity	Sensitivity	Specificity
$>10^5$	0.05	1	>0.9999	0.85	1.000
	0.02	0.85	0.9999	0.15	1.000
	0.01	0.95	0.9995	-	-
	0.005	0.35	0.9999	-	-
10^4-10^5	0.05	0.95	0.9999	0.85	1.000
	0.02	0.9	0.9999	0.15	1.000
	0.01	0.8	0.9998	-	-
	0.005	0.4	0.9999	-	-
10^3-10^4	0.05	0.8	>0.9999	0.70	1.000
	0.02	0.45	0.9999	0.15	1.000
	0.01	0.2	0.9997	-	-
	0.005	0.1	0.9999	-	-

[a] Per µl transport media
[b] As described in McCrone JT and Lauring AS, J. Virol. 90(15):6884, 2016.
[c] As described in Methods, benchmarked for frequencies 0.02-0.98 only

Supplementary Table 2. Nonsynonymous substitutions in HA antigenic sites

House ID	Enrolment ID	Symptom Onset	Subtype	Frequency	Amino Acid Change	Antigenic Site	Vaccinated	Day of Symptoms
1111	300481	3-30-2011	H3N2	0.071	E62G	E*	No	0
2166	320661	2-13-2012	H3N2	0.071	V297A	C	Yes	1
1302	301355	3-20-2011	H3N2	0.088	L86I	E	Yes	1
3075	331045	12-10-2012	H3N2	0.066	I214T	D	Yes	1
5219	50935	12-5-2014	H3N2	0.175	F193S	B*†	No	3
5263	51106	12-6-2014	H3N2	0.111	T128A	B	Yes	3
5290	51225	12-15-2014	H3N2	0.405	I260V	E*	Yes	1
5302	51273	12-13-2014	H3N2	0.030	S262N	E*	Yes	0
5098	50419	12-22-2014	H3N2	0.364	G208R	D	Yes	4
5033	50141	12-3-2014	H3N2	0.032	A163T	B	Yes	2
5034	50143	1-11-2015	H3N2	0.119	I307R	C	Yes	1
5289	51220	12-13-2014	H3N2	0.038	K189N	B*†	Yes	-1
5033	50141	12-3-2014	H3N2	0.025	D53E	C*	Yes	1
5033	50141	12-3-2014	H3N2	0.023	S312G	C	Yes	1
5269	51132	12-6-2014	H3N2	0.028	I242T	D	Yes	2
5147	50630	11-18-2014	H3N2	0.164	I242L	D	Yes	1
5034	50143	1-11-2015	H3N2	0.161	I307R	C	Yes	2
4185	UM40738	12-14-2013	H1N1	0.021	R208K	Ca	No	2

* Sites observed to vary between antigenically distinct strains in Wiley et al., 1981 and Smith DJ et al., 2004.
† Sites located in the "antigenic ridge" identified in Koel et al., 2013.

Figure 1

Figure 1. Within host diversity of IAV populations. (A) Boxplots (median, 25th and 75th percentiles, whiskers extend to most extreme point within median ± 1.5 x IQR) of the number of viral genomes per microliter transport media stratified by day post symptom onset. Notches represent the approximate 95% confidence interval of the median. (B) Boxplots (median, 25th and 75th percentiles, whiskers extend to most extreme point within median ± 1.5 x IQR) of the number of iSNV in 249 high quality samples stratified by day post symptom onset. (C) Histogram of within host iSNV frequency in 249 high quality samples. Bin width is 0.05 beginning at 0.02. Mutations are colored nonsynonymous (blue) and synonymous (gold) (D) Location of all identified iSNV in the influenza A genome. Mutations are colored nonsynonymous (blue) and synonymous (gold) relative to that sample's consensus sequence. Triangles signify mutations that were found in more than one individual in a given season.

Figure 2

Figure 2. Within host dynamics of IAV. (A) Timing of sample collection for 35 paired longitudinal samples relative to day of symptom onset. Of the 49 total, 35 pairs had minor iSNV present in the first sample. (B) The change in frequency over time for minority nonsynonymous (blue) and synonymous (gold) iSNV identified for the paired samples in (A). (C) The distribution of effective population sizes estimated from 1,000 simulated populations. Simulations were run on populations with characteristics similar to the actual patient-derived populations and with the specified effective population size (x-axis). (D) The effect of iteratively removing iSNV with the most extreme change in frequency (fraction of iSNV removed, x-axis) on the estimated effective population size. The point represents the estimate when all iSNV are included. (E) The posterior distributions of selection coefficients estimated for the 35 iSNV present in isolates sampled one day apart. Distributions are colored according to class relative to the sample consensus sequence, nonsynonymous (blue) synonymous (gold). Variants for which the 95% highest posterior density intervals exclude 0.0 are noted in the margin.

Figure 3. Between host dynamics of IAV. (A) The distribution of pairwise L1-norm distances for household (blue) and randomly-assigned community (gold) pairs. The bar heights are normalized to the height of the highest bar for each given subset (47 for household, 1,592 for community). The red line represents the 5th percentile of the community distribution. (B) Timing of symptom onset for 52 epidemiologically linked transmission pairs. Day of symptom onset for both donor and recipient individuals is indicated by black dots. Dashed lines represent pairs that were removed due to abnormally high genetic distance between isolates, see (A). (C) The frequency of donor iSNV in both donor and recipient samples. Frequencies below 2% and above 98% were set to 0% and 100% respectively. (D) The presence-absence model fit compared with the observed data. The x-axis represents the frequency of donor iSNV with transmitted iSNV plotted along the top and nontransmitted iSNV plotted along the bottom. The black line indicates the probability of transmission for a given iSNV frequency as determined by logistic regression. Similar fits were calculated for 1,000 simulations with a mean bottleneck size of 1.66. Fifty percent of simulated outcomes lie in the darkly shaded region and 95% lie in the lightly shaded regions. (E) The outcome from 1,000 simulated "transmission" events with randomly assigned pairings. The black line represents the observed data, as in (D) the shaded regions represent the middle 50% and 95% of simulated outcomes. The results from the simulated logit models were smoothed by plotting the predicted probability of transmission at 0.02 intervals. (F) The beta-binomial model fit. Similar to (D) except the simulated outcomes are the based on a beta-binomial model using a mean bottleneck of 1.73.

Figure 4

Figure 4. Combined estimates of within host mutation rate and effective population size. Contour plot shows the log likelihood surface for estimates of the effective population size and neutral mutation rate. The point represents the peak ($\mu = 4\times10\text{-}6$, Ne = 36, log likelihood = -4,687). Log likelihoods for each contour are indicated.

Supplementary Figure 1

Supplementary Figure 1. Sequence coverage for all samples. For each sample, the sliding window mean coverage was calculated using a window size of 200 and a step of 100. The distributions of these means are plotted as box plots (median, 25th and 75th percentiles, whiskers extend to most extreme point within median ± 1.5 x IQR) where the y-axis represents the read depth and the x-axis indicates the position of the window in a concatenated IAV genome.

Supplementary Figure 2

Supplementary Figure 2. Approximate maximum likelihood trees of the concatenated coding
sequences for high quality H1N1 samples. The branches are colored by season; the tip
identifiers are colored by household. Arrows with numbers indicate consensus and putative
minor haplotypes for samples with greater than 10 iSNV. Trees were made using FastTree.

Supplementary Figure 3

Supplementary Figure 3. Approximate maximum likelihood trees of the concatenated coding sequences for high quality H3N2 samples. The branches are colored by season; the tip identifiers are colored by household. Arrows with numbers indicate consensus and putative minor haplotypes for samples with greater than 10 iSNV. Trees were made using FastTree.

0.003

Supplementary Figure 4

Supplementary Figure 4. The effect of titer and vaccination on the number of iSNV identified. (A) The number of iSNV identified in an isolate (y-axis) plotted against the titer (x-axis, genomes/μl transport media). (B) The number of iSNV identified in each isolate stratified by whether that individual was vaccinated or not. Red bars indicate the median of each distribution.

Supplementary Figure 5

Supplementary Figure 5. Minority nonsynonymous iSNV in global circulation. The global frequencies of the amino acids that were found as minority variants in sample isolates (x-axis) plotted overtime (y-axis). Each amino acid trace is labeled according to the H3 number scheme. All samples were isolated in December of 2014 (gray line).

Supplementary Figure 6

Supplementary Figure 6. Reproducibility of iSNV identification for paired samples acquired on the same day. The x-axis represents iSNV frequencies found in the home-acquired nasal swab. The y-axis represents iSNV frequencies found the clinic-acquired combined throat and nasal swab.

Supplementary Figure 7

Supplementary Figure 7. Estimate of effective bottleneck size with relaxed variant calling criteria. (A) The frequency of iSNV in both recipient and donor isolates. iSNV were identified using the original variant calling pipeline. (B) The presence-absence model fit compared to the observed data for iSNV identified using the original variant calling pipeline. The x-axis represents the frequency of donor iSNV with transmitted iSNV plotted along the top and nontransmitted iSNV plotted along the bottom. The black line indicates the probability of transmission for a given iSNV frequency as determined by logistic regression. Similar fits were calculated for 1,000 simulations with a mean bottleneck size of 2.10. Fifty percent of simulated outcomes lie in the darkly shaded region and 95% lie in the lightly shaded regions. (C) Similar to (A) but with minority iSNV identified using the current analytical framework without a frequency threshold. (D) Similar to B but with minority iSNV identified using the current analytical framework without a frequency threshold.