The Complex Ecosystem in Non Small Cell Lung Cancer Invasion

Seth Haney[1], Jessica Konen[2,3], Adam I. Marcus[3,4], Maxim Bazhenov[1]

[1]Department of Medicine, Division of Pulmonary, Critical Care & Sleep Medicine, University of California, San Diego, La Jolla, CA

[2]Department of Thoracic/Head and Neck Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, Texas, USA.

[3]Winship Cancer Institute, Emory University, Atlanta, GA

[4]Department of Hematology and Medical Oncology, Emory University, Atlanta, GA

Supported by: R01DC012943 (MB), R01CA194027 (AIM), R21CA201744 (AIM), U54CA209992 (AIM). SH is Postdoctoral Kirschstein-NRSA fellow supported by training award T32HL134632.

Abstract

Many tumors are characterized by genetic instability, producing an assortment of genetic variants of tumor cells called subclones. These tumors and their surrounding environments form complex multi-cellular ecosystems, where subclones compete for resources and cooperate to perform multiple tasks, including cancer invasion. Our recent empirical studies revealed existence of such distinct phenotypes of cancer cells, leaders and followers, in lung cancer. These two cellular subclones exchange a complex array of extracellular signals demonstrating a symbiotic relationship at the cellular level. Here, we develop a computational model of the microenvironment of the lung cancer ecosystem to explore how the interactions between subclones can advance or inhibit invasion. We found that, due to the complexity of the ecosystem, invasion may have very different dynamics characterized by the different levels of aggressiveness. By altering the signaling environment, we could alter the ecological relationship between the cell types and the overall ecosystem development. Competition between leader and follower cell populations (defined by the limited amount of resources), positive feedback within the leader cell population (controlled by the focal adhesion kinase and fibronectin signaling), and impact of the follower cells to the leaders (represented by yet undetermined proliferation signal) all had major effects on the outcome of the collective dynamics. Specifically, our analysis revealed a class of tumors (defined by the strengths of fibronectin signaling and competition) that are particularly sensitive to manipulations of the signaling environment. This class can undergo irreversible changes to the tumor ecosystem that outlast these manipulations of feedbacks and have a profound impact on invasive potential. Our study predicts a complex division of labor between cancer cell subclones and suggests new treatment strategies targeting signaling within the tumor ecosystem.

Author Summary

Cancer is an elusive disease due to the wide variety of cancer types and adaptability to treatment. How is this adaptability accomplished? Loss of genetic stability, a hallmark of cancer, leads to the emergence of many different types of cancer cells within a tumor. This creates a complex ecosystem where cancer cell types can cooperate, compete, and exploit each other. We have previously used an image-guided technology to isolate

distinct cancer subclones and identify how they interact. Here, we have employed

mathematical modeling to understand how the dynamic feedbacks between different

cancer cell types can impact the success of invasion in lung cancer. We found that

successful invasion required for feedbacks to support the less viable but more invasive

cell types. These predictions may have implications for novel clinical treatment options

and emphasize the need to visualize and probe cancer as a tumor ecosystem.

Introduction

Lung cancer is the second most prevalent type of cancer causing over 150,000 deaths per year in the United States [1]. Insufficient progress has been made in achieving efficacious treatments. One of the main barriers in developing new treatment strategies is the vast diversity between and within cancers; heterogeneity exists between patients with the same tumor type, between tumor loci within a patient (i.e. metastases and primary tumor), and within the primary tumor itself [2,3]. Cancer is distinguished by loss of normal control over cell processes leading to genetic instability and unregulated growth. Genetic instability creates array of different clonal populations with different cell fitnesses, renewal and invasion potential [4]. Competition between different cancerous subclones and between cancerous and normal cell types sets the stage for classical ecological dynamics in the tumor microenvironment. The outcome of this process determines success of the tumor progression and its understanding may help discover novel treatment strategies [5,6].

Invasion of surrounding tissue, either locally or distally via metastasis, is a hallmark of cancer [7]. Extensive research has detailed that invasion is mediated by interactions between tumor and extracellular matrix [8,9] and cancer-associated fibroblasts [10], but there is a lack of focus on the cooperative interactions between distinct cancer subclones. Indeed, in mouse models of lung cancer, collective invasion of cancer cells was shown to correspond markedly more successful metastasis [3,11–13], confirming the critical role of collective invasion in driving cancer progression.

We recently developed a novel image-guided genomics approach termed SaGA that allowed us to identify at least two distinct phenotypic cell types in lung cancer invasion packs: highly migratory *leader cells* and highly proliferative *follower cells* [14]. Genomic and molecular interrogation of purified leader and follower cultures revealed differential gene expression prompting distinguishing phenotypes. Specifically, leader cells utilized focal adhesion kinase signaling to stimulate fibronectin remodeling and invasion. Leader cells also overexpressed many components of the vascular endothelial growth factor (VEGF) pathway facilitating recruitment of follower cells but not the leader cell motility itself [14]. However, leader cells proliferated approximately 70% slower than follower cells due to a variety of mitotic and doubling rate deficiencies. These deficiencies could be corrected by addition of cell media extracted from the follower only cell cultures, leading to conclusion that follower cells produce an unknown extracellular factor

responsible for correcting mitotic deficiencies in the leader cells. In sum, leader cells

provide an escape mechanism for followers, while follower cells (and follower cell media

only) help leaders with increased growth. Together, these data support a service-

resource mutualism during collective invasion, where at least two phenotypically distinct

cell types cooperate to promote their escape.

In this new study, we developed population-level computational model to explore

impact of the complex interactions between leaders and followers cell types on cancer

progression. The model implemented effects of critical signaling factors controlling the

communication between cell types and the interaction between cells and environment.

We derived analytic boundaries dividing parameter space, representing the major

signaling feedbacks, by the critical changes to invasion dynamics. Our study predicts the

critical role of specific signaling pathways involved in the symbiotic interactions between

cancer subclones for the overall success of cancer progression.

Methods

Our model tracks the cell counts of leader cells, *L*, and follower cells, *F*, the concentrations of extracellular factors VEGF, *V*, an unidentified Proliferation factor, *P*, and Fibronectin, *N*, as well as the size of the domains for leader cells, Ω_L, and for follower cells Ω_F. Based on the available data [14], the following processes have been implemented. Leader cells can expand their domain, Ω_L, by secreting Fibronectin, which in turn relaxes competitive pressure on leader cell growth. Leaders also secrete VEGF, which is taken up by follower cells and causes follower cells to follow them. This was modeled by increasing the domain for follower cells, which in turn relaxes competitive pressure on follower cells. Follower cells secrete an unknown proliferation signal that increases the reproductive capacity of leader cells (initially smaller than follower cells). Leader and follower cells also must compete with each other for resources at rate *c* (see Figure 1).

We modeled cell counts (*L* and *F* species) as standard Lotka-Volterra competition [15]. The carrying capacity of the leader cells was dynamic and dependent on the amount of proliferative signal, *P*, present. This capacity increased in a saturating manner with *P*, with maximum equal to the follower cell carrying capacity, K_{F0}. Intra- and inter-specific competition was driven by concentration, i.e. $[L]=L/\Omega_L$, and birthrate was driven by absolute number, *L*. The extracellular species (*V*,*P*,*N*) and domain sizes all had linear dynamics for simplicity. Below primes denote the time derivative of the variable.

$$\frac{L'}{L} = r_L \left[1 - \frac{(L/\Omega_L) + c(F/\Omega_F)}{K_{L0} + (K_F - K_{L0})\left(\frac{P}{\delta + P}\right)} \right] \tag{1}$$

$$\frac{F'}{F} = r_F \left[1 - \frac{(F/\Omega_F) + c(L/\Omega_L)}{K_F} \right] \tag{2}$$

$$V' = \beta_V L - \gamma_V V \tag{3}$$

$$P' = \beta_P F - \gamma_P P \tag{4}$$

$$N' = \beta_N L - \gamma_N N \tag{5}$$

$$\Omega_L' = \beta_{OL} N - \gamma_{OL}(\Omega_L - \Omega_{L0}) \tag{6}$$

$\Omega_F' = \beta_{OF} V - \gamma_{OF}(\Omega_F - \Omega_{F0})$ (7)

Here r_L and r_F denote the rate of expansion for leaders and followers, respectively. The

parameter *c* denotes the strength of competition between the two cell types. The

capacity of the environment for follower cells is given by the parameter K_F. The capacity

for leaders depended on an initial capacity, K_{L0}, and on the amount of proliferation signal

in a Hill-like manner with EC50, δ. Each extra-cellular species (*V,P,N)* had a production

rate, β, and a degradation rate γ, the domain size variables (Ω_L and Ω_F) also had a

parameter denoting initial capacity (Ω_{L0} and Ω_{F0}).

Reduction and Feedbacks

Previous 3D spheroid experiments show that invasion occurs on a much faster time

scale than reproduction [14]. By assuming that factors (*V,P,N)* and domains (Ω_L, Ω_F)

change much faster than cell counts, one can reduce these equations to a set of two

equations (*L,F)*, where variables in equations (3)-(7) are at their equilibria

$V_{SS} = \frac{\beta_V}{\gamma_V} L; \quad\quad P_{SS} = \frac{\beta_P}{\gamma_P} F; \quad\quad N_{SS} = \frac{\beta_N}{\gamma_N} L; \quad\quad \Omega_L^{SS} = \frac{\beta_{OL}}{\gamma_{OL}} N; \quad \Omega_F^{SS} = \frac{\beta_{OF}}{\gamma_{OF}} V;$ (8)

Using this reduction drastically reduced the complexity of the system. First, we defined

the feedbacks based on the reduced system. The feedback that determines the leaders

impact on their own domain expansion was denoted by $s_L = \frac{\beta_N \beta_{OL}}{\gamma_N \gamma_{OL}}$, for the strength of the

leader only feedback. The feedback that determines the leaders impact on follower cell

growth was denoted by $s_{LF} = \frac{\beta_V \beta_{OF}}{\gamma_V \gamma_{OF}}$, for the strength of the leader to follower feedback.

The feedback that determines the followers impact on leader cell growth was denoted by

$s_{FL} = \frac{\beta_P}{\gamma_P \delta}$, for the strength of the follower to leader feedback. Second, using these

assumptions, we re-wrote the leader-follower system as

$\frac{L'}{L} = r_L \left[1 - \frac{(L/\Omega_L^{SS}(L)) + c(F/\Omega_F^{SS}(L))}{K_L(F)} \right]$ (9)

$\frac{F'}{F} = r_L \left[1 - \frac{(F/\Omega_F^{SS}(L)) + c(L/\Omega_L^{SS}(L))}{K_F} \right]$ (10)

where

$\Omega_L^{SS}(L) = s_L L + \Omega_{L0}; \quad \Omega_F^{SS}(L) = s_{LF} L + \Omega_{F0}; \quad K_L(F) = K_{L0} + (K_F - K_{L0})\frac{s_{FL}F}{1+s_{FL}F}.$

Using this reduction we can derive several critical points in invasion. The reduced

system (9),(10) may have five equilibrium points: extinction of leaders (O_1: L=0, F>0),

followers (O_2: L>0, F=0), both (O_3: L=0, F=0), and two coexistence points (O_4, O_5)

(where both leaders and followers populations are non-zero: L>0, F>0; O_4 is always

stable, wheras O_5 is unstable). Changes in the feedback strengths cause fundamental

shifts in dynamics. In the following we used parameter values $\boldsymbol{\Omega_L = 1, \Omega_F = 1}$. To match

experimental observations that leader cells grow slower and less effeciently, we set

$\boldsymbol{r_L = 0.3}$ and $\boldsymbol{K_{L0} = 0.3}$ while $\boldsymbol{r_F = 1}$ and $\boldsymbol{K_F = 1}$. The strengths of the various

feedbacks, $\boldsymbol{s_L, s_{LF}}$, and $\boldsymbol{s_{FL}}$ are varied systematically below.

Transcritical Bifurcation at Zero

To determine the critical points in the leader-follower system, we calculated the Jacobian

of the reduced system evaluated for the leader extinction equilibrium (O_1 :$L = 0, F =$

$F_{LE} = \Omega_F \cdot K_F$).

$$J|_{L=0;F=F_{LE}} = \begin{pmatrix} r_L(1 - \frac{cK_F}{K_L^{SS}}) & 0 \\ r_F\left(\frac{K_{F0}s_{FL}}{\Omega_{F0}^2} - \frac{c}{\Omega_{L0}}\right) & -r_F \end{pmatrix} \qquad (11)$$

Here $K_L^{SS} = K_{L0} + (K_F - K_{L0})\frac{s_{FL} \cdot F_{LE}}{1 + s_{FL} \cdot F_{LE}}$, the value of K_L when $F = F_{LE}$. The Jacobian has

eigenvalues

$$\lambda = \left[r_L\left(1 - c\frac{K_F}{K_L^{SS}}\right), -r_F\right] \qquad (12)$$

For $c < \frac{K_L^{SS}}{K_F}$, O_1 is unstable and O_4 (steady state where both L>0 and F>0) is stable. At

$c = \frac{K_L^{SS}}{K_F}$ these two equilibria coincide, and for $c > \frac{K_L^{SS}}{K_F}$ equilibrium O_4 moves to the left of

the L=0 axis and becomes unstable while O_1 gains stability. Thus, extinction of leaders

(O_1) is stable as long as $c > \frac{K_L^{SS}}{K_F}$, which determines an upper bound on competition

where leader and followers can coexist and a bifurcation we call the transcritical
bifurcation at zero.

Saddle Node Bifurcation

The system undergoes a saddle node bifurcation when two coexistence equilibria (O_4
and O_5), representing non-zero populations of both leaders and followers, coincide and
disapper. Beyond this bifurcation point the leader/follower populations undergo
unbounded growth. This bifurcation was determined numerically using MatCont [16]. We
found that this bifurcation point depends critically on both the leader feedback strength,
s_L, and on the competition strength, *c*. One of these coexistence points is effected by the
transcritical bifurcation, below.

Transcritical Bifurcation at Infinity

When the leader feedback strength is sufficiently high relative to competition, leaders
and followers may undergo unbounded growth from the initial conditions belonging to the
certain regions of the phase space. We describe this scenario as an attractor basin in
the phase space for the stable infinity attractor. However, if s_L is reduced (or *c* is
increased) beyond a certain threshold, infinity becomes unstable. This corresponds
precisely with the loss of an unstable coexistence equilibrium with non-zero values of
both leaders and followers (O_5). Leaders and followers that are coexisting must satisfy

$$\frac{L}{\Omega_L} + c\frac{F}{\Omega_F} = K_L(F) \tag{13}$$

and $\frac{F}{\Omega_F} + c\frac{L}{\Omega_L} = K_F$ or equivalently,

$$\frac{F}{\Omega_F} = K_F - c\frac{L}{\Omega_L}. \tag{14}$$

In the case that follower populations are large relative to δ, $K_L(F) \rightarrow K_F$, we substituted
(13) into (14) to find

$$L = \frac{K_F\Omega_{L0}}{(1+c)\left(1-\frac{K_F s_L}{1+c}\right)} \tag{15}$$

which has a discontinuity at

$$c = K_F s_L - 1 \tag{16}$$

defining the loss of one of the coexistence equilibrium points (O_5) when it moves to
infinity. We describe this as the transcritical bifurcation at infinity as the stability of infinity
changes at this point.

Results

Leader and Follower Ecosystem

Leader and follower cell types in non-small cell lung cancer spheroids were previously
isolated using a fluorescence technique termed SaGA [14] (Figure 1A,B). We found that
leaders and followers are genotypically and phenotypically distinct populations of cancer
cells that exchange a variety of signaling molecules to coordinate complex behavior
during invasion. In this new work, we focus on four main channels of communication
(see Figure 1C). Leader cells secrete fibronectin in an autocrine manner. This leads to
ECM restructuring and expansion of leader cell domain, Ω_L, (see Methods) which
ultimately increases the leader cell count. The strength of this positive feedback is
characterized in our model by s_L (strength of Leader only feedback). Leader cells also
secrete VEGF. In the leader-follower ecosystem this promotes follower cells to track
expanding leader cells, increases follower domain size (Ω_F), and ultimately, follower cell
count. In our model, the strength of this feedback is given by s_{LF} (strength of Leader to
Follower feedback). Follower cells secrete an undetermined proliferation signal, as
evidenced by the observation that follower-only cell media increases leader cell growth
rate [14]. The strength of this feedback is given by s_{FL} (strength of Follower to Leader
feedback) in the model. Finally, both cell types compete for resources, which is modeled
here by the feedback *c*.

These feedback mechanisms were incorporated into a modified Lotka-Volterra
type competition-cooperation model. We chose a Lotka-Volterra model to focus on the
ecological aspects of competition in the cancer ecosystem. Here, the leader cells could
grow to a total capacity K_L, which is an increasing function of the proliferation signal
secreted by the follower cells. This capacity was reached when a combination of leader
and follower cell densities (cell counts divided by domains) exceeds K_L (see Methods).
Increases in the domain size of each type (by Fibronectin secretion in the leader case
and VEGF in the follower case) limited the overall density of that cell type and mitigated
its impact on the overall capacity of the system. Increasing competition, for example by
limiting resources, increased the impact of either cell type on the conjugate capacity type
(e.g. how leader density, L/Ω_L, impacts follower capacity K_F).

This system of the feedbacks between the leader and follower cells describes a
complex dynamical ecosystem. The impact these feedbacks may have on cancer growth

or invasion is unclear. Leader and follower cells are engaged in competition for
resources but can also be engaged in cooperation and play supportive roles. For
example, invasive leader cells provide new territory for the follower cell population and
are supported by proliferative follower cells. In the following, we analyzed the model to
find critical turning points for the ecosystem dynamics.

Multiple Types of Invasion Dynamics

We found that multiple feedbacks between the leader and follower cell populations could
produce a wide variety of complex dynamics. When competition strength, *c*, was high
and the strength of the leader only feedback, s_L, was moderate, population dynamic was
bounded and resulted in a stable cell count for both leader and follower cell populations
as well as a stable domain size (Fig. 2A). In contrast, when feedback was large and
competition was moderate, population dynamics revealed an unbounded growth (Fig.
2B). Intermediate values of both *c* and s_L led to dynamic regimes that depended on the
initial cell count: ecosystems with large initial cell count underwent unbounded growth,
while small ecosystems attained a stable size (Figure 2C). These types of dynamics are
in a qualitative agreement with experimental studies which revealed (a) rapid expansion
of intact leader-follower ecosystem and (b) that blocking specific feedback mechanisms
in vitro can reduce or block cell population growth. Specifically, blockade of fibronectin
signaling or blockade of VEGF signaling led to significantly reduced invasion [14].

This array of behaviors can be explained by the critical shifts in the cell
population dynamics due to the changes in the feedbacks strength. We found that
depending on the level of competition, *c*, and the strength of invasiveness of leaders, s_L,
the leader-follower ecosystem can operate in one of five different regimes, as described
below (Figure 3).

Leader Extinction: When competition was high and invasive feedback was minimal, the
leader cells (the weaker competitor) were forced to extinction while the follower cells
persisted and its population reached a stable size (Figure 3A,B). There was a critical
level of competition between leaders and followers, given by $c > \frac{K_L^{SS}}{K_F}$ (see Methods,
Transcritical bifurcation at zero for derivation), required for this type of dynamics. This
critical level of competition, the ratio of the capacity of leader cells, K_L^{SS} , to that of the

follower cells, K_F , essentially depends on the fitness differences between leader only

and follower only cell populations. Leader and follower populations with similar fitness

would tolerate a much higher competition threshold without driving one species to

extinction. From the dynamical systems perspective, when $c > \frac{K_L^{SS}}{K_F}$ and s_L is sufficiently

low (see below), the only stable equilibrium in the phase space is O_1 and all system

trajectories converge to this equilibrium point representing the leader cells extinction

state (Figure 3B).

Leader Extinction with Escape: If competition was above the leader extinction

limit, $c > \frac{K_L^{SS}}{K_{F0}}$, but not high enough to balance the impact of the leader only feedback,

$c < K_F s_L - 1$, there were two possible outcomes depending on the initial population size

(see Methods, *Transcritical Bifurcation at Infinity* for derivation) (Figure 3A,C). The

second condition, $c < K_F s_L - 1$, can be interpreted as a balance between positive

feedback, s_L, and negative feedback, c. In this regime, leaders could go extinct if the

initial population of leader cells was sufficiently small. Alternatively, if initial populations

of leaders and followers both were large enough, the ecosystem could grow

unboundedly. Thus, our model predicts, that the ability to undergo successful collective

invasion depends on whether the initial bulk size is larger than a critical amount. These

types of dynamics with divergent outcomes occur when competition is large enough to

be able to drive leaders extinct, but small enough so it can be outbalanced by the strong

invasive effects of the leader cells.

In the phase space of the model, the basins of attraction of the two distinct

dynamical regimes are separated by a critical boundary (separatrix of a saddle

equilibrium O_5) where the cell bulk size determines its ultimate fate (Figure 3C). Both

infinity and the leader extinction equilibrium (O_1) are stable attractors representing two

possible end solutions of the system dynamics.

Non-invasive Dynamics: When competition was smaller than the extinction limit,

$c < \frac{K_L^{SS}}{K_{F0}}$, but large enough to balance leader feedback strength, $c > K_F s_L - 1$, the

ecosystem size remained bounded and both leaders and followers attained a stable and

non-zero population size. In the phase space, this type of dynamics corresponds to

conversion to the stable equilibrium O_4 (Figure 3A,D). We refer to this as non-invasive

dynamics, as the cells cannot grow beyond a defined size. In this case, while

competition was present, it was too weak to lead to extinction, while leader population
was not invasive enough to promote unlimited growth. This scenario represents stable,
non-invasive dynamics.

Multimodal Dynamics: If competition was (a) small enough to allow leader existence,
$c < \frac{K_L^{SS}}{K_{F0}}$, (b) small enough relative to the leader feedback strength, so that escape was
possible, $c < K_F s_L - 1$, but (c) high enough, so that for small initial population of leader it
could balance the positive leader feedback, leader and follower cell dynamics depended
on the initial population size (Figure 3A,E). Ecosystems with a large initial cell count
would grow without bound but those with a small initial cell count would reach a stable
population size, due to the competition as in the non-invasive dynamics case. On the
phase plane the last outcome was represented by a contraction to a stable equilibrium
O_4. This critical boundary was defined by a separatrix of a saddle fixed point O_5 (Fig. 3E).
(This separatrix was determined numerically by reversing time [17].)

Aggressive Dynamics: When leader invasive strength was sufficiently high and
competition was sufficiently low, the only possible outcome was unbounded growth of
both cell populations (Figure 3A, F). In this case, the only stable attractor in the phase
space is infinity where all system trajectories are converged to.

In summary, our analysis revealed that the complex balance of the feedbacks in
the leader-follower ecosystem can lead to the multiple types of population dynamics.
When the leaders' invasiveness was low, the outcome depended on the competition
between two populations – strong enough competition promoted leader extinction, while
weak competition allowed stable coexistence states with bounded size of both leader
and follower cell populations. As leader invasiveness rate increased, the system
revealed a new state with unbounded growth. This aggressive dynamic state coexisted
with a stable attractor representing a bounded size of both populations if competition
between leaders and followers was strong enough. Otherwise, unlimited population
growth was the only outcome. Based on the system dynamics derived above, next we
will show how critical boundaries between parameter regimes could be exploited to lead
to profound changes in the ecosystem dynamics.

Limiting leader feedback Leads to Irreversible Changes in Invasion

In the multimodal dynamics (e.g., Fig. 3C or 3E), leader and follower cell populations can
undergo explosive growth or achieve a stable count depending on the initial size of the
ecosystem. We examined the impact of limiting the invasive leader feedback in
scenarios of this type (Fig. 4). Even when the ecosystem was initially sufficiently large to
support unbounded growth, after reducing invasive leader feedback s_L (Fig. 4A), the
ecosystem was forced into the non-invasive dynamics type and the total bulk of the cell
population reduced reaching a steady-state (Fig. 4E). Importantly, the leader and
follower cell populations remained stable and bounded after restoring invasive leader
feedback to its original strength (Fig 4E, right side).

From the point of view of the dynamical systems analysis, reducing leader
feedback changed the phase space, so the only stable attractor was non-zero
equilibrium (O_4) (Fig. 4C). In this regime, unlimited growth was abandoned and the
system converged to the equilibrium state (O_4) corresponding to the bounded size of
both cell populations. This equilibrium remained stable even after the feedback was
restored to its original level (Fig. 4D).

Our model predictions (Fig. 5A) are consistent with in vitro data (Fig. 5B). Using
siRNA blocking we previously showed that expression of fibronectin (which is
characterized by the strength of leader only feedback, s_L, in the model) led to the low
invasion potential and a stable cell population size [14].

Increasing Competitive Signals Leads to Leader Extinction

We next tested effect of increasing competition between leader and follower cell
populations on the ecosystem dynamics (Fig. 6). Leader cells excrete extracellular
factors that induce the death of the followers and leaders alike [14], which supports
competition. Here, we again started from aggressive unbounded type of dynamics and
then increased competition strength (Fig. 6A). This caused change of the ecosystem
dynamics. Both cell populations reduced the size, with leader cell population going to
extinction state (Fig. 6E). However, upon restoring competition to the original level,
leader and follower cells reemerge and grow unboundedly again. The last can be
avoided if no leader cells remain (complete extinction).

Again, this dynamic can be easily understood using bifurcation analysis. Increasing
competition strength made leader extinction equilibrium state O_1 stable (Fig. 6C).
However, when competition was restored to its original level, O_1 became unstable again
and leader and follower cells returned to escape dynamics (Fig. 6D). Importantly, in the
extreme case of very small cell populations, cells undergo discrete and stochastic
dynamics and complete extinction of a small population of leaders is possible in a finite
time, leading to irreversible changes due to competition increase (similar dynamics was
described in our previous study [18]).

Support For Leaders has Large Impact on Aggressiveness

Changing the strength of the feedbacks that determine the interaction between
leaders and followers (s_{LF} and s_{FL}) could also impact the dynamics. Leader cells secrete
VEGF (denoted here by s_{LF}) that helps follower cells to expand their territory and follower
cells secrete a proliferation signal (denoted here by s_{FL}) that allows leaders to increase
their proliferative capacity. These two feedbacks have distinct impacts on the overall
ecosystem dynamics. Perturbations to s_{LF} (changing the impact that leaders have on
followers) changed the system dynamics (assuming that the cell count was small
enough at the time of the intervention) from unlimited growth to the bounded type. The
size of both leader and follower cell populations decreased reaching non-zero steady-
state (Fig. 7E). This regime persisted as long as the feedback from the leaders to
followers remained low. However, increasing s_{LF} to its original level restored the system
dynamics with unlimited cell population growth (Fig. 7E, right size).

Using bifurcation analysis, we found that reducing impact that leaders have on
followers shifted the location of the saddle node bifurcation boundary that separated
state with unlimited growth only dynamics and a state with coexistence of the unlimited
growth and a stable equilibrium attractor (O_4) regimes (Fig. 7A). Effectively, decreasing
s_{LF} increased the threshold level of the invasive leader feedback (s_L) needed to cause
unbounded growth. Thus, reducing s_{LF} made the system to converge to the stable
equilibrium state O_4 corresponding to the bounded size of both cell populations (Fig. 7C).
However, increasing s_{LF} to its original level changed the phase space again, so infinity
became the only stable attractor (Fig. 7D) and unlimited growth dynamics resumed.

Our model predictions (Fig. 5C) are consistent with in vitro data (Fig. 5D). Using

siRNA to block the VEGF receptor VEGFR2 (siKDR in Fig. 5D), we previously showed

that blocking the leader to follower feedback led to the limited invasion potential and

stable cell population size (Fig. 5D) [14].

Finally, we tested the role of the follower to leader feedback (s_{FL}) and found that

perturbations to s_{FL} have a significant impact on the system dynamics. In contrast to s_{LF},

changes to the s_{FL} changed both the location of the saddle node bifurcation boundary

and the transcritical bifurcation boundary of the leader extinction (Fig. 8A). Therefore,

decreasing s_{FL} both increased the threshold on the leader invasion strength (s_L) needed

to cause unbounded population growth and decreased the threshold of the competition

strength (c) needed to induce leader population extinction. We have exploited this to

show that decreasing s_{FL} can cause irreversible change in the cell population bulk. Again,

starting with unlimited growth dynamics (Fig. 8B), decreasing follower to leader feedback,

s_{LF}, reversed the dynamics and both leader and follower cell population reduced in size

converging to the steady-state (Fig. 8E). This regime with bounded ecosystem size

persisted after the feedback was restored (Fig. 8E, right side).

Using dynamical systems analysis, we found that reducing follower to leader

feedback (s_{FL}) triggered the system convergence to the stable attractor (O_1) representing

the leader extinction state (Fig. 8C). When the feedback was restored, O_1 becomes

unstable but the ecosystem fell to the attraction basin of the stable equilibrium O_4 and

avoided regime of unlimited growth (Fig. 8D). In more general case, the outcome

depended on the balance between the leader to follower, s_{FL}, and follower to leader, s_{LF},

feedbacks, with higher s_{LF} requiring more significant s_{FL} decrease to avoid unbounded

growth (Fig. 8F).

Summary of Perturbations to Cancer Ecosystem

Complex balance of the feedbacks within the cancer cell ecosystem allows for some

alterations of the feedback parameters to have significant impacts on the ecosystem

dynamics. We summarized these different possibilities in Table 1 from the perspective of

achieving the goal to reduce cell population bulk. Hence, manipulating s_L, s_{LF}, s_{FL} should

be interpreted as decreasing these feedbacks, whereas manipulating *c* should be

interpreted as increasing *c*. We also examined the possibility of non-targeted cell death,

such as might occur during non-specific chemotherapy. Manipulations were either

irreversible, so the system dynamics remained altered upon cessation of the

perturbation (e.g. irreversible leader extinction or irreversible stabilization of the cell

count), or caused only temporal and reversible reduction of the cell bulk. In some cases,

such as leader extinction with escape and multimodal dynamics (see Fig. 3), the size of

the initial cell bulk dictated possible outcomes of the feedback perturbations. The

outcomes described in the Table 1 represent the best-case scenario. Thus,

perturbations were started from an appropriate initial state and maintained long enough

to achieve the desired effect.

This analysis revealed that certain parameter regimes are more sensitive to the

perturbations than others. Specifically, in the leader extinction with escape regime (area

(2) in Figure 3A) and the multimodal dynamics regime (area (4) in Figure 3A)

perturbations could have irreversible impacts on the ecosystem. In these cases, any

perturbation (death, reduction in s_L, s_{LF}, s_{FL}, or increase in c) can potentially force the

system to cross the critical boundary (separatrix) and transition from explosive growth to

a steady-state dynamic. These regimes give a unique opportunity to impact the

invasiveness of the ecosystem.

Also, certain perturbations could force the ecosystem into a state where leader

extinction (O_1) is stable. This occurs when applying sufficient increases in the

competition pressure, c, or decreases in the support from followers to leaders, s_{LF}. In

these cases, it is possible for the discrete and stochastic nature of the cell population

dynamics to define the ecosystem fate. Thus, a sufficiently long perturbation could

irreversibly eradicate a sufficiently small discrete number of leader cells [18].

Type of dynamics	Initial State	Manipulation Type	Outcome
1. Leader Extinction	N/A	death, c, s_L, s_{LF}, s_{FL}	Reversible cell bulk reduction.
2. Leader Extinction w/ Escape	Infinity attractor basin	death, s_L, c, s_{LF}, s_{FL}	Irreversible leader extinction.
	Stable attractor basin	death, c, s_L, s_{LF}, s_{FL}	Reversible cell bulk reduction.
3. Non-invasive Dynamics	N/A	c, s_{FL} s_L, s_{LF}, death	Reversible cell bulk reduction.
4. Multimodal	Infinity attractor basin	c, s_{FL}, s_L, s_{LF}, death	Irreversible stabilization of cell bulk.
	Stable attractor basin	c, $s_{FL,}$ s_L, s_{LF}, death	Reversible cell bulk reduction.
5. Aggressive Dynamics	N/A	c, s_{FL} s_L, s_{LF}, death	Reversible cell bulk reduction.

Table 1: Effect of the different feedback alterations on the ecosystem dynamics.
First column - the different types of the ecosystem dynamics in the (s_L, c) parameter
space are as shown in Figure 3. In some cases, initial population size dictated the
outcome. These outcomes are distinguished in the initial state given in the 2d column.
An initial state in the infinity attractor basin denotes that the cell bulk exceeded the
critical amount and could grow unboundedly; initial state in the stable attractor basin
denotes the case when the cell counts were less than the critical value and the system
converged to the stable attractor. Third column - each transient manipulation to the
model parameters had a goal to reduce cell population size (decrease of s_L, s_{LF}, s_{FL} or
increase of c). The "death" indicates a non-targeted "enforced" reduction of the cell
population. The last column indicates the system dynamics after the original values of
the model parameters were restored. The changes were either irreversible upon
cessation of the perturbation (e.g. irreversible leader extinction and irreversible
stabilization of cell bulk) or caused reversible reduction in the cell population bulk.

Discussion

Heterogeneity of tumors, at the genetic, epigenetic, and phenotypic levels, is one
of the main obstacles to developing new effective treatment strategies. Tumor cells
rapidly evolve forming highly efficient symbiotic systems with well-defined labor division
targeted to augment tumor survival and expansion. In lung cancer collective invasion
packs observed *in vitro*, two distinct populations of cancer cells - highly migratory *leader*
cells and highly proliferative *follower cells* – have been recently identified [14]. In this
new study, we used computational models to explore collective dynamics of the leader-
follower ecosystem and to exploit approaches that can effectively disrupt it.

We found that competition between two populations (defined by the limited
amount of resources), the positive feedback within the leader cell population (controlled
by the focal adhesion kinase and fibronectin signaling) and impact of the follower cells to
the leaders (represented by yet undetermined proliferation signal) all had major effects
on the outcome of the collective dynamics. While increase of the positive feedback
within the leader cell population would ultimately lead to the system state with
unbounded growth, manipulating follower to leader feedback or increasing competition
between leader and follower cell populations was able to reverse this dynamic and to
form a stable configuration of the leader and follower cell populations.

Our model highlights the importance of focal adhesion kinase (FAK) and
fibronectin signaling. Our previous empirical work showed that FAK signaling was a key
distinguishing feature between leader and follower cells and critical for invasive leader
behavior [14]. Our model predicts that FAK is the main driver of invasion by leader cells
and disruptions in the FAK driven feedback loop cause critical changes in the leader-
follower population dynamics. Indeed, FAK is a well-known regulator of the tumor micro-
environment: promoting cell motility and invasion [19]. FAK expression is upregulated in
ovarian [20] and breast cancer [21] tumors with expression levels correlating with
survival [22,23]. Many FAK inhibitors, such as defactinib, are currently in clinical trials
with promising results [19,24,24–28]. A key advantage of FAK inhibitors is that they
impact both the tumor itself and the surrounding stroma where tumor associated
fibroblasts also utilize FAK signaling to promote tumor invasiveness [29,30].

While commonly associated with angiogenesis in healthy and cancerous tissue,
our previous work showed that VEGF mediates communication between leader and
follower cells [14]. There is a long history of targeting VEGF to limit tumor invasiveness
[31,32]. While great success has been seen in preclinical models [33,34], only moderate
success was seen in clinical trials with anti-VEGF drugs such as bevacizumab [35,36].
This is largely due to cancers developing resistance to specific VEGF-therapeutics. In
our model, VEGF stimulated followers to shadow leaders and expand their domain.
However, we found that inhibition of VEGF had little impact on the ecosystem dynamics
relative to the perturbations of the other axes (such as FAK or competition for resources).

Competition for resources is one of the principal forces that structures any
ecosystem, including tumor ecosystems [6,37]. Our modeling work predicts that
competition was a critical component in the leader-follower ecosystem. We found that
when the strength of competition exceeded a critical threshold, leaders (the weaker
competitor) were driven to extinction. Further, enhancements of the competition in the
model changed the fundamental cell population dynamics. In some cases this meant
stopping unbounded growth and promoting the extinction of the leader cells. Our
previous in vitro work demonstrated that leaders may inhibit the growth of followers
through an unknown secreted factor in cell media [14]. While still in the early stages,
exploiting this inhibition may also provide similar benefits to those shown here as
increases in competition.

Our previous study also revealed a currently unknown extracellular factor
secreted by followers that corrects mitotic deficiencies and enhances leader proliferation
[14]. Our modeling highlights this factor as having critical impact on the ecosystem
dynamics. We found that blockade of this proliferation factor, modeled here by the
strength of the followed to leader feedback, can cause critical shifts in the population
dynamics. More work needs to be done to identify and understand the mechanism of this
action, but preliminary results suggests that this may be a potential novel treatment axis
that specifically targets the mutualistic interaction between leaders and followers.

Ecological forces shape the exchange of biomaterial between different biotic and
abiotic environmental agents. These forces determine capacity of the ecosystem for
different species (subclones) and the environment ultimately sets the fitness of each of
the competitors. Classic ecological theory dictates that an abundance of many similar

species (such as similar subclonal populations) will lead to a high competition for
resources [38,39]. This competition can force the exclusion of inferior competitors and
ultimately may reduce heterogeneity of the system. However, when symbiotic and
mutualistic interactions occur, otherwise competitive species support each other and
increase the capacity of the ecosystem [40,41]. Symbiosis between different subclonal
populations may be particularly important during critical times when the tumor survival is
in peril (such as hypoxia, metastasis or therapy). One critical moment in tumor
progression occurs when highly proliferative tumor cells saturate the resource potential
of their current environment. In order to obtain more resources, tumors need to invade
new territory.

Previous results to model complex tumor cell population dynamics range from
detailed cellular level models (e.g. [9,42–44]) to continuous models with a different
degree of complexity (e.g. [18,45–48]) similar to that proposed in our new study. While
cellular level model can directly incorporate heterogeneous cell types and intrinsic tumor
properties, including proliferation, metabolism, migration, protease and basement
membrane protein expression, and cell-cell adhesion, they typically have high-
dimensional variables and parameter space that is difficult to explore. Advantages of the
reduced type of models include the low dimensional parameter space, where
parameters have clear biophysical meanings, and which allows for systematic analysis
to rapidly explore and determine the sensitive parameter space. We previously applied
this approach to study cell interactions in chronic cancers and predicted conditions for
explosive tumor growth [18]. Similar approach was applied to model cancer cell
population dynamics in many other types of cancer [45,48,49].

The vast diversity between different cancers and even between different cell
types within a single tumor remains one of the biggest hurdles to overcome to achieve
personalized cancer treatment. This diversity leads to a complex array of interactions
between different tumor cell types and the healthy surrounding tissue: the tumor
ecosystem. Our work has isolated phenotypically unique lung cancer cells and taken a
dynamical approach to understanding the interactions within the tumor ecosystem. We
identified the critical features and interactions composing the leader-follower ecosystem,
to explore vulnerabilities of the lung cancer invasive cell populations.

1.   American Cancer Society. Cancer Facts & Figures 2016.

2.   Burrell RA, McGranahan N, Bartek J, Swanton C. The causes and
consequences of genetic heterogeneity in cancer evolution. Nature.
2013;501: 338–345. doi:10.1038/nature12625

3.   Gerlinger M, Swanton C. How Darwinian models inform therapeutic failure
initiated by clonal heterogeneity in cancer medicine. Br J Cancer. 2010;103:
1139–1143. doi:10.1038/sj.bjc.6605912

4.   Greaves M, Maley CC. Clonal evolution in cancer. Nature. 2012;481: 306–
313. doi:10.1038/nature10762

5.   Korolev KS, Xavier JB, Gore J. Turning ecology and evolution against
cancer. Nat Rev Cancer. 2014;14: 371–380. doi:10.1038/nrc3712

6.   Pienta KJ, McGregor N, Axelrod R, Axelrod DE. Ecological Therapy for
Cancer: Defining Tumors Using an Ecosystem Paradigm Suggests New
Opportunities for Novel Cancer Treatments. Transl Oncol. 2008;1: 158–164.

7.   Hanahan D, Weinberg RA. Hallmarks of Cancer: The Next Generation. Cell.
2011;144: 646–674. doi:10.1016/j.cell.2011.02.013

8.   Ruoslahti E. The Walter Herbert Lecture. Control of cell motility and tumour
invasion by extracellular matrix interactions. Br J Cancer. 1992;66: 239–242.

9.   Bauer AL, Jackson TL, Jiang Y. Topography of Extracellular Matrix Mediates
Vascular Morphogenesis and Migration Speeds in Angiogenesis. PLOS
Comput Biol. 2009;5: e1000445. doi:10.1371/journal.pcbi.1000445

10.   Paulsson J, Micke P. Prognostic relevance of cancer-associated fibroblasts
in human cancer. Semin Cancer Biol. 2014;25: 61–68.
doi:10.1016/j.semcancer.2014.02.006

11.   Ewald AJ, Huebner RJ, Palsdottir H, Lee JK, Perez MJ, Jorgens DM, et al.
Mammary collective cell migration involves transient loss of epithelial
features and individual cell migration within the epithelium. J Cell Sci.
2012;125: 2638–2654. doi:10.1242/jcs.096875

12.   Friedl P, Noble PB, Walton PA, Laird DW, Chauvin PJ, Tabah RJ, et al.
Migration of coordinated cell clusters in mesenchymal and epithelial cancer
explants in vitro. Cancer Res. 1995;55: 4557–4560.

13.   Friedl P. Prespecification and plasticity: shifting mechanisms of cell
migration. Curr Opin Cell Biol. 2004;16: 14–23.
doi:10.1016/j.ceb.2003.11.001

14.   Konen J, Summerbell E, Dwivedi B, Galior K, Hou Y, Rusnak L, et al. Image-
guided genomics reveals a vascular mimicry during symbiotic collective
cancer invasion. Unpublished. 2016;

15.   Smale S. On the differential equations of species in competition. J Math Biol.
1976;3: 5–7. doi:10.1007/BF00307854

16.   Dhooge A, Govaerts W, Kuznetsov YA, Meijer HGE, Sautois B. New
features of the software MatCont for bifurcation analysis of dynamical
systems. Math Comput Model Dyn Syst. 2008;14: 147–175.
doi:10.1080/13873950701742754

17.   Seydel RU. Practical Bifurcation and Stability Analysis. Springer Science &
Business Media; 2009.

18.   Haney S, Reya T, Bazhenov M. Delayed onset of symptoms through
feedback interference in chronic cancers. Converg Sci Phys Oncol. 2016;2:
045002. doi:10.1088/2057-1739/2/4/045002

19.   Sulzmaier FJ, Jean C, Schlaepfer DD. FAK in cancer: mechanistic findings
and clinical applications. Nat Rev Cancer. 2014;14: 598–610.
doi:10.1038/nrc3792

20.   The Cancer Genome Atlas Network. Integrated genomic analyses of ovarian
carcinoma. Nature. 2011;474: 609–615. doi:10.1038/nature10166

21.   The Cancer Genome Atlas Network. Comprehensive molecular portraits of
human breast tumours. Nature. 2012;490: 61–70. doi:10.1038/nature11412

22.   Sood AK, Armaiz-Pena GN, Halder J, Nick AM, Stone RL, Hu W, et al.
Adrenergic modulation of focal adhesion kinase protects human ovarian
cancer cells from anoikis. J Clin Invest. 2010;120: 1515–1523.
doi:10.1172/JCI40802

23.   Ward KK, Tancioni I, Lawson C, Miller NLG, Jean C, Chen XL, et al.
Inhibition of focal adhesion kinase (FAK) activity prevents anchorage-
independent ovarian carcinoma cell growth and tumor progression. Clin Exp
Metastasis. 2013;30: 579–594. doi:10.1007/s10585-012-9562-5

24.   Tanjoni I, Walsh C, Uryu S, Tomar A, Nam J-O, Mielgo A, et al. PND-1186
FAK inhibitor selectively promotes tumor cell apoptosis in three-dimensional
environments. Cancer Biol Ther. 2010;9: 764–777.

25.   Infante JR, Camidge DR, Mileshkin LR, Chen EX, Hicks RJ, Rischin D, et al.
Safety, pharmacokinetic, and pharmacodynamic phase I dose-escalation
trial of PF-00562271, an inhibitor of focal adhesion kinase, in advanced solid
tumors. J Clin Oncol Off J Am Soc Clin Oncol. 2012;30: 1527–1533.
doi:10.1200/JCO.2011.38.9346

26. Roberts WG, Ung E, Whalen P, Cooper B, Hulford C, Autry C, et al.
Antitumor activity and pharmacology of a selective focal adhesion kinase
inhibitor, PF-562,271. Cancer Res. 2008;68: 1935–1944. doi:10.1158/0008-
5472.CAN-07-5155

27. Kang Y, Hu W, Ivan C, Dalton HJ, Miyake T, Pecot CV, et al. Role of focal
adhesion kinase in regulating YB-1-mediated paclitaxel resistance in ovarian
cancer. J Natl Cancer Inst. 2013;105: 1485–1495. doi:10.1093/jnci/djt210

28. Gilbert-Ross M, Konen J, Koo J, Shupe J, Robinson BS, Wiles WG, et al.
Targeting adhesion signaling in KRAS, LKB1 mutant lung adenocarcinoma.
JCI Insight. 2017;2. doi:10.1172/jci.insight.90487

29. Lu C, Bonome T, Li Y, Kamat AA, Han LY, Schmandt R, et al. Gene
alterations identified by expression profiling in tumor-associated endothelial
cells from invasive ovarian carcinoma. Cancer Res. 2007;67: 1757–1768.
doi:10.1158/0008-5472.CAN-06-3700

30. Jean C, Chen XL, Nam J-O, Tancioni I, Uryu S, Lawson C, et al. Inhibition of
endothelial FAK activity prevents tumor metastasis by enhancing barrier
function. J Cell Biol. 2014;204: 247–263. doi:10.1083/jcb.201307067

31. Sitohy B, Nagy JA, Dvorak HF. Anti-VEGF/VEGFR Therapy for Cancer:
Reassessing the Target. Cancer Res. 2012;72: 1909–1914.
doi:10.1158/0008-5472.CAN-11-3406

32. Folkman J. Tumor angiogenesis: therapeutic implications. N Engl J Med.
1971;285: 1182–1186. doi:10.1056/NEJM197111182852108

33. Inai T, Mancuso M, Hashizume H, Baffert F, Haskell A, Baluk P, et al.
Inhibition of vascular endothelial growth factor (VEGF) signaling in cancer
causes loss of endothelial fenestrations, regression of tumor vessels, and
appearance of basement membrane ghosts. Am J Pathol. 2004;165: 35–52.
doi:10.1016/S0002-9440(10)63273-7

34. Kim ES, Serur A, Huang J, Manley CA, McCrudden KW, Frischer JS, et al.
Potent VEGF blockade causes regression of coopted vessels in a model of
neuroblastoma. Proc Natl Acad Sci U S A. 2002;99: 11399–11404.
doi:10.1073/pnas.172398399

35. Hayes DF. Bevacizumab Treatment for Solid Tumors: Boon or Bust? JAMA.
2011;305: 506–508. doi:10.1001/jama.2011.57

36. Hurwitz H, Fehrenbacher L, Novotny W, Cartwright T, Hainsworth J, Heim W,
et al. Bevacizumab plus Irinotecan, Fluorouracil, and Leucovorin for
Metastatic Colorectal Cancer. N Engl J Med. 2004;350: 2335–2342.
doi:10.1056/NEJMoa032691

37. Basanta D, Anderson ARA. Exploiting ecological principles to better
understand cancer progression and treatment. Interface Focus. 2013;3.
doi:10.1098/rsfs.2013.0020

38. Hardin G. The Competitive Exclusion Principle. Science. 1960;131: 1292–
1297. doi:10.1126/science.131.3409.1292

39. Huston M. A General Hypothesis of Species Diversity. Am Nat. 1979;113:
81–101. doi:10.1086/283366

40. D H Boucher, S James, Keeler and KH. The Ecology of Mutualism. Annu
Rev Ecol Syst. 1982;13: 315–347.
doi:10.1146/annurev.es.13.110182.001531

41. Stachowicz JJ. Mutualism, Facilitation, and the Structure of Ecological
CommunitiesPositive interactions play a critical, but underappreciated, role
in ecological communities by reducing physical or biotic stresses in existing
habitats and by creating new habitats on which many species depend.
BioScience. 2001;51: 235–246. doi:10.1641/0006-
3568(2001)051[0235:MFATSO]2.0.CO;2

42. Bauer AL, Jackson TL, Jiang Y. A cell-based model exhibiting branching and
anastomosis during tumor-induced angiogenesis. Biophys J. 2007;92:
3105–3121. doi:10.1529/biophysj.106.101501

43. Schaller G, Meyer-Hermann M. Multicellular tumor spheroid in an off-lattice
Voronoi-Delaunay cell model. Phys Rev E. 2005;71: 051910.
doi:10.1103/PhysRevE.71.051910

44. Shirinifard A, Gens JS, Zaitlen BL, Popławski NJ, Swat M, Glazier JA. 3D
Multi-Cell Simulation of Tumor Growth and Angiogenesis. PLOS ONE.
2009;4: e7190. doi:10.1371/journal.pone.0007190

45. Rodriguez-Brenes IA, Komarova NL, Wodarz D. Evolutionary dynamics of
feedback escape and the development of stem-cell–driven cancers. Proc
Natl Acad Sci. 2011;108: 18983–18988.

46. Lowengrub JS, Frieboes HB, Jin F, Chuang Y-L, Li X, Macklin P, et al.
Nonlinear modelling of cancer: bridging the gap between cells and tumours.
Nonlinearity. 2010;23: R1. doi:10.1088/0951-7715/23/1/R01

47. Anderson ARA, Weaver AM, Cummings PT, Quaranta V. Tumor morphology
and phenotypic evolution driven by selective pressure from the
microenvironment. Cell. 2006;127: 905–915. doi:10.1016/j.cell.2006.09.042

48. Liu X, Johnson S, Liu S, Kanojia D, Yue W, Singh UP, et al. Nonlinear
Growth Kinetics of Breast Cancer Stem Cells: Implications for Cancer Stem
Cell Targeted Therapy. Sci Rep. 2013;3. Available:

http://www.nature.com/srep/2013/130820/srep02473/full/srep02473.html?W
T.ec_id=SREP-631-20130902

49. Wodarz D, Komarova NL. Dynamics of cancer: mathematical foundations of
oncology [Internet]. World Scientific; 2014. Available:
https://books.google.com/books?hl=en&lr=&id=90i7CgAAQBAJ&oi=fnd&pg=
PR7&dq=Dynamics+of+cancer+komarova&ots=OjSC2djQSP&sig=80eRRrs
kJ6xbQu0Cnk46GomlyDQ

Fig 1: Leader and Follower system. A) Pictorial representation of the spatio-temporal genomic
and cellular analysis (SaGA). Laser excitation of Dendra2 drives a change in fluorescence of user-
specified cells. After degradation of cell matrix, fluorescence-based cell sorting is used to separate
cells into leader and follower groups allowing for genetic analysis on specified groups. Printed with
permission from Fairman Studios, LLC. This image is not included in the Creative Commons
licence for the article. Adapted from Figure 1 in [14]. B) Photo-conversion examples using 3-D
spheroids of H1299-Dendra2 cells. L= leader cell, F = follower cell. Adapted from Figure 1 in [14].
C) Stick representation of mathematical model of leader and follower cell interactions and invasion.
Positive feedbacks are given by arrows, while negative feedbacks are given by flat-ended curves.
The strength of leader only feedback (s_L) is mediated by fibroNectin (N). The strength of leader to
follower feedback (s_{LF}) is mediated by VEGF (V). The strength of follower to leader feedback (s_{FL})
is mediated by a proliferation signal secreted by followers (P). The strength of competition is given
by c.

Fig 2: Ecosystem dynamics depend strongly on the feedbacks strength. Characteristic
examples of the cancer cell population dynamics in the model for different strength of the
competition between leader and followed cell populations, c, and leader population intrinsic
feedback, s_L . Cancer cell populations may attain a stable size (A: s_L = 1.2, c = 0.6), grow
unboundedly (B: s_L = 1.2, c = 0.05), or be dependent on the initial tumor size (C: s_L = 2, c = 0.375).

Fig 3: Dynamics of the leader and follower ecosystem. A) Bifurcation diagram sweeping leader population intrinsic feedback strengths, s_L, and competition strength, c. Abbreviations - LE: Leader Extinction, LE w/ Esc.: Leader Extinction with Escape, ND: Non-invasive Dynamics, MD: Multimodal Dynamics, AD: Aggressive Dynamics, TC Bif. at ∞: Transcritical Bifurcation at infinity (see Methods), TC Bif. at 0: Transcritical Bifurcation at zero (see Methods), SN Bif.: Saddle Node Bifurcation (see Methods). B-E) Phase diagrams for each regime identified in A). Dashed blue and green lines are null-clines for the leaders and followers, respectively. Red curves show the trajectories of the leader-follower system from different initial conditions, with arrows denoting tangent vectors at various points. Open circles denote stable equilibrium points, stars denote unstable equilibria. Specific s_L and c parameters used in panels B-E) are given by the circles in A).

Fig 4: Blocking leader cell population feedback in the Multi-modal Dynamics regime can lead to irreversible changes of the population dynamics. A) Bifurcation diagram depicting the direction of the perturbation in the parameter space - increase in feedback, s_L. Black circuit – initial state; red circuit – perturbed state. B-D) Phase plots of dynamics before (B), during (C), and after (D) blockade of s_L. E) Time-course of cell bulk before, during, and after perturbation. Note irreversible change of the ecosystem dynamics.

Fig 5: Model reproduces in vitro experimental results. A) In the model, leader cell count (blue) and domain size (red) are given in both normal (left) and leader population feedback, s_L, blocked (right) conditions. B) In cell culture, invasion of leader cells was significantly reduced during siRNA block of focal adhesion kinase (right), compare to control (left). Scale bar 100μm. Reproduced from [14]. C) In the model, blocking leader to follower feedback, s_{LF}, limited invasive area and cell count. D) Impact on invasion of leader cell cultures during siRNA block of VEGFR2 (siKDR). Scale bar 100μm. Invasive area was significantly reduced after VEGFR2 block (p<0.0001) (right), compare to control (left). Reproduced from [14].

Fig 6: Enhancing competition between leader and follower populations can drive transient
extinction of leaders. A) Bifurcation diagram depicting the direction of the perturbation - increase
in competition, *c*. Black circuit – initial state; red circuit – perturbed state. B-D) Phase plots of
dynamics before (B), during (C), and after (D) enhancement of competition. E) Time-course of cell
count. Here, we assume total extinction of leaders occurs during treatment, i.e. at some point
during treatment *L=0*. Note that ecosystem dynamics is reversed after perturbation is removed.

Fig 7: Disrupting Leader to Follower feedback, s_{LF}, can trigger transient changes in the

population dynamics. A) Bifurcation diagram depicting the direction of the perturbation in

parameter space. Perturbations in s_{LF} change the location of the saddle node bifurcation boundary.

Black line – initial location of the bifurcation boundary; red line – perturbed location. B-D) Phase

plots of dynamics before (B), during (C), and after (D) blockade of s_{LF}. E) Time-course of cell count.

Note that ecosystem dynamics is reversed after perturbation is removed

Fig 8: Disrupting Follower to Leader feedback, s_{FL}, can have irreversible changes in
dynamics leading to stabilization of cell count. A) Bifurcation diagram depicting the direction of
the perturbation in parameter space. Perturbations in s_{FL} change the location of the saddle node
bifurcation and transcritical bifurcation at zero boundaries. Black lines – initial location of the
bifurcation boundaries; red lines – perturbed location. B-D) Phase plots of dynamics before (B),
during (C), and after (D) blockade of s_{FL}. E) Time-course of cell count. Note irreversible change of
the ecosystem dynamics. F) Bifurcation diagram depicting the position of the saddle node
bifurcation point as a function of s_{LF} and s_{FL}. AD: Aggressive Dynamics; MD: Multi-modal Dynamics.
Here, s_L = 1.4 and c = 0.2.

799