Title: The genomic determinants of adaptive evolution in a fungal pathogen

Authors: Jonathan Grandaubert[1,2], Julien Y. Dutheil[3,4,5], Eva H. Stukenbrock[1,2,5]

Affiliations:

[1] Environmental Genomics Group, Max Planck Institute for Evolutionary Biology, August-Thienemann-Str. 2, 24306 Plön, Germany, [2] Christian-Albrechts University of Kiel, Am Botanischen Garten 1-9, 24118 Kiel, Germany, [3] Research group Molecular Systems Evolution, Max Planck Institute for Evolutionary Biology, August-Thienemann-Str. 2, 24306 Plön, Germany, [4] UMR 5554 Institut des Sciences de l'Evolution, CNRS, IRD, EPHE, Université de Montpellier, Place E. Bataillon, 34095, Montpellier, France.

[5] **Corresponding authors:**

Eva H. Stukenbrock, Environmental Genomics, CAU Kiel, Am Botanischen Garten 1-11, 24118 Kiel, Germany, Email: estukenbrock@bot.uni-kiel.de, phone: +49 (0) 431 880 6368

Julien Y. Dutheil, Research group Molecular Systems Evolution, Max Planck Institute for Evolutionary Biology, August-Thienemann-Str. 2, 24306 Plön, Germany, Email: dutheil@evolbio.mpg.de, phone: +49 (0) 4522 763 298

Keywords: Genome evolution, adaptation, evolutionary rates, recombination, plant pathogenic fungi

Runing Title: Adaptive evolution in a fungal plant pathogen

Abstract:

Background: Antagonistic host-pathogen co-evolution is a determining factor in the outcome of infection and shapes genetic diversity at the population level of both partners. Little is known about the overall genomic rate of evolution in pathogens or the molecular bases of rapid adaptation. Here we apply a population genomic approach to infer genome-wide patterns of selection among thirteen isolates of the fungal pathogen *Zymoseptoria tritici*.

Results: Based on whole genome alignments, we report extensive presence-absence polymorphisms of genes resulting from a high extent of within population karyotypic variation. We apply different test statistics based on the distribution of non-synonymous and synonymous polymorphisms (pN/pS) and substitutions (dN/dS) to estimate rates of adaptation and identify specific targets of selection. We report that the pN/pS ratio of sites under purifying selection negatively correlates with the local recombination rate, as expected under a scenario of background selection. Contrasting polymorphism and divergence with a closely related species, we estimate that 44% of substitutions in the proteome of *Z. tritici* are adaptive and we find that the rate of adaptation is positively correlated with recombination rate. The proportion of adaptive amino acid substitutions in genes encoding determinants of pathogenicity is as high as 68% underlining the importance of positive selection in the evolution of virulence-associated traits. Using a maximum likelihood approach and codon models of sequence evolution, we furthermore identify a set of 786 additional genes showing signatures of diversifying selection, which we find are preferentially located in region of high recombination rates.

Conclusions: We report a high rate of adaptive mutations in the genome of *Z. tritici*, in particular in genes involved in host-pathogen interactions. Furthermore, we identify a set of genes with signature of balancing / diversifying selection, which we find are related to secondary metabolism and host infection. Finally, we report a strong impact of recombination on the efficacy of purifying, adaptive and diversifying selection, suggesting pervasive effect of linked selection in the genome and thereby highlighting the importance of recombination in this pathogen species.

Background

Antagonistic host-pathogen interactions drive co-evolutionary dynamics between pathogens and their hosts. Signatures of selection in genome data can inform about mechanisms of evolution and identify targets of selection at interacting loci [1]. In general, genome studies of microbial pathogens have focused on rapidly evolving genes involved in pathogenicity, such as "effector" genes encoding proteins that interfere with host defences and may determine host range of the pathogen [2]. Effector genes of fungal pathogens have frequently been found to associate with repetitive DNA and it is proposed that repeat rich genome compartments provide particularly favourable environments for rapid evolution of new virulence specificities (e.g. [3–6]). Repetitive DNA may locally increase mutation rate and contribute to gene duplications and structural variation among alleles [7]. Yet little is known about the interplay of mutation, natural selection, genetic drift and, for sexually reproducing species, recombination along the genome of fast evolving pathogens.

Rates of evolution and the type of selection acting on a gene can be inferred from the comparison of sequences sampled from distinct individuals in a population. Using statistical population genetics, it is possible to infer the rate at which certain mutations are fixed in comparison to the 'null' expectation of neutral evolution, that is, without selection [8]. In the case of protein-coding sequences, this neutral rate is typically given by the rate of synonymous substitutions, hereby assumed to be selectively neutral. The relative proportion of synonymous and non-synonymous substitutions thus provides a signature of the underlying type of selection acting on a gene. An excess of segregating or fixed non-synonymous mutations is indicative of positive selection while an absence or reduced proportion of non-synonymous mutations shows conservation of the sequence at the amino acid level indicative of purifying selection [9]. Further tests, building on pioneering work by McDonald and Kreitman [10], assess adaptive evolution by comparing the ratio of non-synonymous to synonymous substitutions between species, dN/dS, to the ratio of non-synonymous to synonymous polymorphisms within species, pN/pS. In the absence of selection, the dN/dS ratio is expected to be equal to the pN/pS ratio. However, for a gene that has undergone positive directional selection, the dN/dS ratio is expected to be higher than the pN/pS ratio. This is because rare, positively selected mutations represent only a minor proportion of all polymorphisms in a population sample. However, positively selected mutations have a higher probability of fixation than neutral mutations and will therefore accumulate over time as species diverge, resulting in a higher dN/dS ratio [11,12].

The McDonald-Kreitman table forms the basis for the quantification of the proportion of

adaptive substitutions in a gene fixed by positive selection defined as $\alpha = 1 - (pN / pS) / (dN / dS) = 1 - (dS * pN) / (dN * pS)$ [13]. Additional modeling of the fitness effect of substitutions

allows α to be estimated in the presence of slightly deleterious or slightly advantageous

mutations that otherwise can bias the inference of positively selected substitutions [14].

These methods have been applied to a range of different species and have been used to show

that the rate of adaptive evolution varies largely among different taxa [11,12].

In pathogen genomes, increased rates of adaptation, as occurring by arms-race evolution

where new non-synonymous mutations repeatedly are selected for, will result in higher

dN/dS ratios. Indeed, evolutionary predictions based on dN/dS analyses and McDonald-

Kreitman tests have identified effectors and other pathogenicity-related genes in pathogen

genomes [15,16]. However, genes that evolve under a trench warfare scenario leave different

signatures in the genome sequence. This type of selection can also be referred to as positive

diversifying selection where a diversity of alleles at the selected locus is maintained by natural

selection. The genic signature of positive diversifying selection is an increased proportion of

non-synonymous polymorphisms [17]. Yet, in a functionally important gene, the majority of

non-synonymous sites will be evolving under selective constraints, and only a small number

will be allowed to change without negatively affecting protein function. Therefore, in most

cases, searching for signatures of positive diversifying selection simply by comparing the

average proportion of non-synonymous to synonymous polymorphisms in a gene will show

little statistical power [18]. To circumvent this, evolutionary codon models that allow for

varying selection intensity across a gene sequences can be applied to identify specific codon

positions that have an excess of non-synonymous to synonymous mutations (a higher pN/pS

ratio) [18].

In this study we assess the different types of selection affecting the genome of a fungal wheat

pathogen *Zymoseptoria tritici* (synonym *Mycosphaerella graminicola*). The genome of this

pathogen consists of thirteen core chromosomes and several accessory chromosomes that

are present at different frequencies in different individuals, constituting a particular case of

karyotypic polymorphism [19]. The accessory chromosomes comprise repeat rich,

heterochromatic DNA with a low gene content and a so far unknown functional relevance

[20,21]. We have previously shown that evolutionary rates on these chromosomes are

particularly high suggesting that genes on the accessory chromosomes in general evolve

under less selective constraints [22].

Previous studies based on comparative population genomic analyses of *Z. tritici* and two closely related species, *Z. pseudotritici* and *Z. ardabiliae* used genome-wide estimates of non-synonymous and synonymous divergence to identify past species-specific signatures of selection in the wheat pathogen [16,22]. Functional characterization of some of these genes revealed amino acid substitutions important for *in planta* development and asexual spore formation in the wheat-adapted pathogen, and thereby confirmed the use of evolutionary predictions to identify functionally relevant traits [23].

While past signatures of selection can be related to ecological specialization of *Zymoseptoria* species, ongoing positive selection likely reflects host-pathogen arms race or trench warfare evolution and recurrent adaptations to changing environmental conditions including fungicide treatments and wheat cultivars. Population genomic studies of *Z. tritici* have revealed that the pathogen has a high effective population size, which correlates positively with a strong efficacy of natural selection, i.e. the ability to rapidly fix advantageous mutations and eliminate non-advantageous [16]. Additional parameters that can influence rapid adaptation are recombination and mutation rates, as well as genome architecture. Recombination rate variation has been studied in *Z. tritici* by experimental mating and population genomic sequencing. Independently, these approaches have confirmed exceptionally high recombination rates (~60cM/Mb) ([24,25]). So far the implications of such high recombination rates on gene evolution are not known.

We apply a population genomics approach to infer genome-wide positive and purifying selection among thirteen isolates of *Z. tritici* collected from bread wheat in Europe and the Middle East. Our analyses based on more than 1.4 million single nucleotide polymorphisms (SNPs) and including 700,000 coding sites allow us to identify past and on-going signatures of selection in the genome of *Z. tritici*. Notably, our analyses reveal a strong importance of recombination in gene evolution and a particularly high rate of adaptive substitutions in genes encoding putative effectors. These analyses underline the potential of rapid adaptation of virulence related traits in this important agricultural pathogen.

Results and Discussion

Genetic structure of *Z. tritici* populations

We generated a population genomic dataset of thirteen *Z. tritici* isolates obtained from different field populations in Europe and Iran (Table S1). Given the high extent of structural variation in genomes of *Z. tritici* isolates, we assembled each genome *de novo* to capture structural differences. We used the resulting thirteen genome assemblies to generate a

filtered multiple genome alignment of ~27 Mb (Table S2) comprising a total of 1,489,362 SNPs of which approximately 50% locate in protein coding regions (Table S1). The SNP data was used to compute the overall genetic diversity of the sample showing a mean value of π = 0.022 per site.

Several studies have addressed the population genetic structure of *Z. tritici* at different spatial scales (e.g. [26,27]). These studies have concluded that *Z. tritici* comprises a panmictic population with very little population structure. We here used the genomic data to investigate the relationship of the *Z. tritici* isolates. First, we assessed population genetic structure by analyzing the ancestral recombination graph of the thirteen genomes. To this end, we slid 10 kb windows along the multiple genome alignment, and estimated the genealogy for each window. The resulting 1,850 trees were combined into a super tree (Figure 1A). If the sample of genomes is taken from a panmictic population, the super tree is expected to be a star tree. However, here we observe at least two clusters, one comprising all European isolates and the other comprising two isolates collected in Iran (Figure 1A). We further investigated population structure using the program ADMIXTURE [28] and found the strongest support for a model with two ancestral populations supporting the tree-based clusters of European and Iranian isolates (Figure 1B). Within the European population four German isolates furthermore form an independent group also clustered together by the ADMIXTURE analysis using k=3. While the population sample used here is smaller than the sample sizes used in previous population genetic studies, our analyses build on a considerably larger number of variable sites providing a higher resolution of the genealogical relationships among *Z. tritici* isolates. The thirteen isolates show genetic differentiation that reflects their geographical origin suggesting some extent of subdivision and geographical barriers to gene flow between populations. This geographical pattern is consistent with the global genetic structure of other important wheat pathogens such as the wheat yellow rust fungus *Puccinia striiformis f.sp. tritici* and the wheat scab pathogen *Fusarium graminearum* [29,30], and shows that local adaptation and population divergence can shape the genetic structure of cereal pathogens even in the presence of global agricultural trade and long distance pathogen dispersal [31].

Z. tritici shows presence-absence polymorphism of genes and whole chromosomes

Z. tritici exhibits dramatic patterns of chromosome length polymorphisms and presence-absence variation of accessory chromosomes [32–34]. Based on the multiple genome alignment, we here investigated presence-absence polymorphisms of chromosomes,

chromosome fragments and genes among the thirteen isolates. We focused on non-repetitive

DNA and show, in agreement with a previous study [35], that 96.4% of the core chromosomes

(chromosome 1 to 13) are shared among all isolates (Figure S1). However, as expected, the

mapping of assembled contigs aligning to the accessory chromosomes varied considerably

between isolates (Figure S1).

Overall, 9,499 of the 11,839 genes predicted in the reference genome of IPO323 are present

in all of the thirteen isolates. Of the remaining, 1,584 genes are located on core chromosomes

and 667 on accessory chromosomes. This set of genes comprises genes present at different

frequencies among the thirteen *Z. tritici* isolates including 89 genes present only in IPO323

and of which 77 locate on core chromosomes.

A previous genome comparison of two Swiss *Z. tritici* isolates revealed nearly 300 novel genes

not characterized in the reference genome [36]. We speculated that the different isolates of

Z. tritici also could contain accessory genes and chromosomes not present in the reference

IPO323. To test this we investigated the subset of contigs that were not aligned to the

reference genome. We found a total of 5.2 Mb in the twelve re-sequenced *Z. tritici* isolates

with no homology to sequences in the reference genome. These new *Z. tritici* sequences,

corresponding to accessory fragments, encompass between 602 kb and 995 kb of non-

repetitive DNA in each genomes. We applied an *ab initio* gene predictor to assess whether

these accessory sequences encode proteins and found evidence for 1,041 novel full-length

gene models encoding proteins of at least 30 amino acids. Of these 75% encode proteins

without known function consistent with a general pattern observed on the accessory

chromosomes with a high proportion of gene models without a predicted function [19].

Isolate-specific fragments were previously shown to encode effector candidate genes strongly

up-regulated in *Z. tritici* during host infection [36]. In other fungal pathogens such as

Verticilium dahliae and *Magnaporthe oryzae* isolate-specific genes were experimentally

shown to encode virulence determinants [37,38]. Based on our findings, we hypothesize that

the large number of isolate-specific genes in thirteen *Z. tritici* genomes could play a role in

host-pathogen interactions. Under this scenario, the observed variation in gene content could

reflect diversifying selection imposed by genetic traits in a variety of host genotypes.

Measures of adaptive evolution in *Z. tritici*

We next aimed to obtain a quantitative assessment of adaptive mutations in the genome of *Z.*

tritici. To this end, we first estimated the non-synonymous and synonymous divergence dN

and dS using a genome alignment of *Z. tritici* and its sister species *Z. ardabiliae*. Furthermore,

we used the *Z. tritici* SNP data to compute the (unfolded) site frequency spectra (SFS) of

synonymous and non-synonymous sites using *Z. ardabiliae* as an outgroup. By contrasting

divergence and polymorphism data, we estimated the parameter α as a measure of the

proportion of adaptive non-synonymous substitutions. The SFS is however strongly affected

by demography and the presence of slightly deleterious mutations segregating at low

frequencies (see review by [14]). State-of-the-art statistical methods account for these effects

by modeling the distribution of fitness effects (DFE) of mutations [12,39]. We used the

program Dfem to compute α as well as ω_a, the rate of adaptive substitutions, using four

distinct DFE models accounting for mutations with both slightly deleterious and beneficial

effects [12]. We fitted and compared the DFE models (see Materials and Methods) and found

that the Gamma-Exponential model best fitted our data (Table 1) in agreement with studies

from animals [12]. This suggests the existence of slightly deleterious, as well as slightly

beneficial segregating mutations in the genome of *Z. tritici* (Table 1). The estimates provide an

α value of 44% as a genome average, and an ω_a value of 0.05. Both values are in the range of

what is observed for Mammals (with the exception of Primates) but considerably higher than

estimates from plants [12,39]. In candidate effector genes, however, the rate of mutations

fixed by selection is more than twice as high as the average of all genes (ω_a equal to 0.14

compared to the genome average of 0.05, Table 1), with 68% of substitutions in these genes

inferred to be adaptive. This average estimate is close to the highest values reported in

animals [12]. We further aimed at assessing the impact of recombination on the fixation of

adaptive mutations, by considering genes in distinct recombination environments. We report

a significant positive correlation between α and ρ (the population scaled recombination rate

inferred as ρ = 4Ne x r) (Kendall's tau = 0.25, p-value = 6.816e-6) and ω_a and ρ (Kendall's tau =

0.21, p-value = 1.95e-4, Figure 2), suggesting that a higher recombination rate favors the

fixation of adaptive mutations, as expected by a Hill-Robertson effect [40,41]. These results,

based on population genomic data of *Z. tritici*, represent the first quantitative assessment of

adaptive evolution in the genome of a fungal pathogen. The exceptionally high rate of

adaptation in effector genes likely reflects arms race evolution driven by the antagonistic

interaction of *Z. tritici* and its host.

Strength of purifying selection along the genome of *Z. tritici*

Loci at different positions of the genome evolve at different speed. This is due to variation in

the effect of different evolutionary processes across the genome as well as variation in the

composition of the DNA sequence itself [42]. In sexual organisms, recombination is a strong

determinant of evolutionary rates, and differences in the frequency of recombination can

strongly impact the rate whereby adaptive mutations can be fixed and whereby (slightly)

deleterious mutations are removed [41]. Evolutionary processes at any given site in the

genome are strongly influenced by selection acting on linked sites, a process referred to as

linked selection [40] and background selection in the case of linkage with mutation under

purifying selection [40,43]. Here, we aimed to address the genome-wide strength of purifying

selection in *Z. tritici* in protein coding genes, using the ratio of non-synonymous to

synonymous polymorphisms (pN/pS ratio).

We computed pN and pS for each gene with two methods: (1) by inference of average

pairwise heterozigosity [44,45] and (2) using a maximum likelihood framework and models of

sequence evolution [18]. In this framework, two types of models were compared: a two-rate

model (M1a), with sites evolving under neutrality or purifying selection with a pN/pS ratio

equal or below one (neutral model with purifying selection) and a three-rate model (M2a)

allowing for some sites to evolve under positive selection with a pN/pS ratio above one

(positive selection model) [46]. The best fitting model according to a likelihood ratio test

(False Discovery Rate < 0.01) was kept for further analyses. Such model-based inference

specifically accounts for heterogeneous selective pressures within genes, allowing inference

of pN/pS ratios for sites under purifying selection and for sites under positive selection, if any.

The average heterozygosity computation assumes an infinite site model and does not allow

for recurrent mutations. The model-based approach, while accounting for recurrent

mutations and more accurate mutational models, assumes an underlying tree structure and

no recombination in the region under study [47]. As *Z. tritici* exhibits both high mutation rates

and intragenic recombination, it is a priori unclear which method provides the best estimates

of pN/pS ratios and purifying selection. To assess the sensitivity of model-based estimates in

the presence of recombination, we used coalescent simulations (See Methods). We find that

estimates of purifying selection based on the pN/pS ratio (denoted "ω0" in the M1a and M2a

models) are indeed inflated in the presence of high recombination rates (Figure S2). While the

effect is significant, it is of small size. Furthermore, it goes in the opposite direction of what is

expected under background selection, as selection should be more efficient in highly

recombining regions and lead to the removal of more deleterious mutations reflected by a

lower omega ω0 value. We therefore conclude that the model-based inference of purifying

selection is conservative and we used these estimates in the further analyses. We estimated

the strength of purifying selection for a gene as the ω0 parameter of the best fitting model

between M1a and M2a. This procedure ensures that we only measure the strength of
purifying selection, even in the presence of positions evolving under positive selection.

To investigate which genome parameters determine the strength of purifying selection in *Z.*
tritici, we compared ω0 for each gene to 1) mean gene expression, 2) population
recombination rate in flanking non-coding regions, 3) gene length, 4) local gene density and 5)
GC content at third codon positions. We further considered whether the gene 6) is predicted
to be secreted, 7) is located on an accessory chromosome and 8) is an orphan gene (Figures 3
and 4). We fitted a linear model with ω0 as a response variable and the above-mentioned
observed statistics as explanatory variables. We identify several variables that significantly
explain the strength of purifying selection (Table 2): the length of the coding protein (number
of codons) has a highly significant negative effect, meaning that evolution of larger genes is
more constrained. Other studies that document an impact of gene length on evolutionary
rates have likewise shown that small genes evolve faster than long genes [48]. We also
observe that orphan genes, which in general are shorter than core genes (mean 325 amino
acids vs 516 amino acids on average, Wilcoxon test with p-value < 2.2e-16), are less
conserved, even when protein length is accounted for (Table 2) [48]. Many orphan genes in *Z.*
tritici have likely evolved de novo and may not encode proteins of essential functions. Our
observation of higher ω0 in orphan genes is consistent with findings in other organisms, e.g.
in primates where younger lineages specific genes evolve under less selective constraints than
older "common" primate genes [49].

We used *Z. tritici* gene expression data from early host colonization (four days after spore
inoculation on leaves of seedlings of a susceptible wheat host) and *in vitro* growth to assess
the correlation of transcription level and gene evolution [50]. Studies in other organisms
including yeast and bacteria have documented a strong impact of expression levels on gene
evolution whereby highly expressed genes are more conserved [48,51]. Consistent with these
observations we also find a significant negative correlation between ω0 and gene expression
(Figure 4 and Table 2) showing that highly expressed genes are more conserved in *Z. tritici*.

We furthermore find a highly significant negative effect of recombination rate on ω0,
implying that purifying selection is more efficient in regions with high recombination rate
(Figures 4 and 5). This result is in agreement with a model of background selection, where
purifying selection at linked loci reduces the local effective population size, therefore
reducing the efficacy of selection and allowing slightly deleterious mutations to spread more
frequently than at unlinked loci [43,52]. This finding supports a central role of recombination
in the removal of non-adaptive mutations in the genome of *Z. tritici* consistent with patterns

described in *Drosophila melanogaster* where rates of evolution were compared in regions of

high and low recombination [53].

Lastly, we find that GC content is significantly negatively correlated with ω0, that is, genes

with higher GC tend to be under stronger purifying selection. This may be explained by an

indirect effect of recombination as recombination rate and the GC content at third codon

positions (GC3) are positively correlated (Figure 4, Kendall's tau = 0.097, p-value < 2.2e16). A

similar correlation of recombination and GC3 is found in other organisms [54]. In

Saccharomyces cerevisiae this correlation has been explained by the impact of biased gene

conversion on sequences evolution [55]. However, a thorough search for signatures of GC-

biased gene conversion did not find any pervasive effect of this phenomenon in *Z. tritici* [25].

An alternative explanation is selection for codon usage. In *Z. tritici,* as in several other

organisms, preferred codons often contain a cytosine or a guanine at their third position

(Figure S3) [56]. As highly expressed genes are more conserved and enriched in preferred

codons, the relationship between ω0 and GC might be a by-product of the correlation with

gene expression.

Detection of on-going diversifying selection in *Z. tritici*

Next we aimed at detecting sites in the *Z. tritici* genome showing signatures of positive

diversifying selection. We fitted models of codon sequence evolution as implemented in the

CodeML program of the PAML package to detect genes with significant signatures of positive

diversifying selection using likelihood ratio tests [46]. Two models are typically compared: the

"null" model M1a, where sites evolve only under neutral or purifying selection (ω <= 1), and

an alternative model M2a where sites in coding sequence also can evolve with a ratio of non-

synonymous over synonymous mutations above one (ω > 1). When applied to population

data, sites with ω > 1 can reflect positive diversifying selection [47].

We fitted codon models for genes present in at least three isolates (83% of genes located on

core chromosomes and 31% of genes on the accessory chromosomes) (Table S3). After

correcting for multiple testing, we identified a final set of 786 genes (including 24 on the

accessory chromosomes) having sites evolving under positive diversifying selection (False

Discovery Rate < 0.01) (Table S3). In total, our analyses recognized 4,686 individual codons

evolving under positive selection in the *Z. tritici* genome (0,001% of 4,640,090 analyzed

codons).

In several pathogen genomes, rapidly evolving genes are found clustered in particular

genomic environments often associated with repetitive sequences (e.g [7,15]). To address if

the same pattern is found in *Z. tritici*, we assessed the spatial distribution of genes under

positive diversifying selection along chromosomes (see Materials and Methods). Of the 786

positively selected genes, 240 are located within a distance of 5 kb from each other and

thereby form 108 clusters containing two to four genes with signatures of positive selection.

At the genome scale, clusters containing two or three genes do not show a significant pattern

as the same pattern can be obtained by randomly distributing the positively selected genes

across the genome. However, there are two significant clusters (p-value = 1.8×10^{-3})

containing four and eight genes, respectively. The clusters comprising four genes is located in

a 24kb region of chromosome 5, and the cluster comprising eight genes in a 31 kb region of

chromosome 9. For genes in both clusters no functional relevance can be assigned. Aside

these two clusters, there is no evidence that positively selected genes are physically

associated with each other in *Z. tritici*, nor that they are significantly enriched or depleted in

repeat rich regions.

The molecular battlefield between plants and their pathogens involves proteins secreted from

both host and pathogen cells [2]. Pathogenicity related genes therefore often encode signal

peptides targeting the translated protein for secretion. As shown above, adaptive

substitutions are fixed at a much higher rate in candidate effectors than in non-effector

genes. This observation is in agreement with an antagonistic co-evolution of genes involved in

host-pathogen protein-protein interactions. We also asked whether positive diversifying

selection, consistent with a trench-warfare scenario of co-evolution, drives effector evolution

in *Z. tritici*. To this end we assessed the proportion of effector candidates among the genes

predicted to be under diversifying selection, but found no evidence for a particular

enrichment of these genes (X^2 test, p-value = 0.171).

To jointly analyse the potential genomic variables affecting the distribution of genes under

positive diversifying selections, we fitted a binary logistic regression where the occurrence of

positive selection, as predicted by the likelihood ratio test on codon models, is used as

response variable. Several explanatory variables were included (Table 3), and three of them

appeared to have a highly significant effect. First, we report that highly expressed genes are

less likely to be under positive selection, which we hypothesise is due to a higher level of

selective constraint. Next, we find that genes in regions with higher recombination rates are

more likely to be under positive diversifying selection, which we interpret as evidence for a

Hill-Robertson effect [41,53]. Lastly, the longer the gene the more likely it is to be under

positive selection. We speculate however that this effect reflect a higher statistical power is

large genes.

We further extended our analyses of the genes predicted to be under positive diversifying

selection by characterizing known proteins domains. From the 786 candidate genes, 602 of

the encoded proteins (76.5%) have an *in silico* attributed function or harbour known protein

domains. We conducted a PFAM domain enrichment analyses and identified 21 significantly

enriched domains (FDR <= 0.05) (Table S4). These domains can be grouped into different

categories based on their associated molecular function and include carbohydrate-active

enzymes (CAZymes), polyketide synthases (PKS), non-ribosomal peptide synthetases (NRPS),

and cellular transporters (Table S4). Several of these categories are relevant for the pathogen

to interact with its host. For example, CAZymes are proteins involved in the break down of

glycosidic bonds contained in plant cell walls [57], and PKS and NRPS are multimodular

enzymes involved in the biosynthesis of secondary metabolites of which many also have been

shown to be involved in virulence of plant pathogenic fungi (reviewed in [58]).

Among the positively selected *Z. tritici* genes, we also searched for homologs of known

virulence factors described in other plant pathogens. The PFAM domain PF14856 corresponds

to the mature part of a virulence factor Ecp2 described in the tomato fungal pathogen

Cladosporium fulvum [59]. Ecp2 has been described in several other plant pathogens [60] and

has three homologs in *Z. tritici*. We find that two of these homologs comprise sites under

positive selection supporting a virulence related role of this gene also in this wheat pathogen.

Transposable elements and the origin of new adaptive mutation

In some filamentous Ascomycetes, Repeat Induced Point (RIP) mutations are a genome

defence mechanism that targets repetitive DNA. While experimental evidence for RIP so far

only has been documented in a few species [61,62], sequence signatures of RIP are reported in

several other Ascomycete and Basidiomycete species [63,64]. During meiosis RIP generates

single nucleotide changes mutating cytosine bases to thymine, preferentially at CpA

dinucleotides [61]. The accelerated mutation rate in repetitive DNA or duplicated sequences

mediated by RIP can interfere with the evolution of non-repetitive DNA via "leakage" of RIP

mutations into neighboring genes [6]. To infer the potential impact of RIP-like mutations on

gene evolution in *Z. tritici* we assessed the proportion of CpA → TpA (and GpT → ApT for the

reverse strand) mutations in all dinucleotide substitutions using a substitution mapping

approach for genes located at a maximum distance of 500, 1,000, 1,500 and 2,000 bp from a

TE. We find a significant enrichment of RIP-associated dinucleotide substitutions in genes

located close to TEs suggesting "leakage" of RIP mutations (Table S5). The proportions of RIP

mutations decrease with the distance to TEs supporting that the CpA → TpA mutations indeed

originate from RIP. However, we find no enrichment of positively selected sites among the RIP-

associated mutations, suggesting that in *Z. tritici*, RIP is not a major driver of adaptive

evolution.

Signatures of past selection in *Z. tritici* and related species

In a previous study the evolutionary history of *Z. tritici* was inferred using a whole genome

coalescence analyses [16]. We showed that divergence of *Z. tritici* and its sister species *Z.*

pseudotritici occurred recently and likely coincides with the onset of wheat domestication

and thereby specialization of *Z. tritici* to a new host. We hypothesize that genes important for

the colonization of distinct hosts have been under selection during the divergence of

Zymoseptoria species. To infer signatures of past selection we applied the branch model

implemented in the program package PAML to estimate the branch-specific dN/dS ratio for

core *Zymoseptoria* genes (present in four species *Z. tritici*, *Z. pseudotritici*, *Z. ardabiliae* and *Z.*

brevis[20] [46]. The branch-specific dN/dS ratios reflect the proportion of non-synonymous to

synonymous substitutions accumulated in each branch of the *Zymoseptoria* phylogeny. Our

analyses identified 47 genes with a dN/dS ratio > 1 on the *Z. tritici* branch indicative of an

increased non-synonymous divergence, 54 genes in *Z. pseudotritici*, 60 genes in *Z. brevis* and

15 genes in *Z. ardabiliae* (Figure 5). Based on their putative function in host-pathogen

interactions, we hypothesized that some positively selected genes encode secreted proteins

and putative effectors. Indeed we find a significant enrichment of putative effectors among

the positively selected genes on the *Z. tritici* branch (10 genes encoding effector candidates,

χ^2 test, p = 5.272 x 10^{-05}), and the *Z. pseudotritici* branch (9 genes encoding effector

candidates, χ^2 test, p = 0.00315). For *Z. ardabiliae* and *Z. brevis* the branch-specific positively

selected genes were however not enriched in effector candidates (in *Z. ardabiliae* 1 gene

encoding an effector candidate, χ^2 test, p=0.768, and in *Z. brevis* 8 genes encoding effector

candidates, χ^2 test, p=0.1165). We next performed a PFAM domain enrichment analysis for

the positively selected genes with a predicted function to address if some functional domains

are significantly enriched in this set of genes (FDR < 0.05) (Table S6). The majority of genes

encode proteins of unknown function, however among the functionally characterized

proteins we find a gene encoding a regulator of chromosome condensation that was

previously described to be functionally relevant for virulence in *Z. tritici* (Poppe et al. 2015).

Our analyses reveal an enrichment of genes encoding proteinases, one gene encoding Lysin

motifs (LysM) already described as an effector in *Z. tritici* and other fungal pathogens [66,67],

and one gene encoding a CFEM domain. Cystein-rich CFEM domains have been described in

G-protein-coupled receptors in the rice blast pathogen *Magnaporthe oryzae* playing an

important role in pathogenicity [68]. These genes provide interesting candidates for further

studies of molecular determinants of host specificity in *Z. tritici*.

Conclusions

The analysis of the assembled genomes of thirteen isolates of the fungus *Z. tritici* revealed a

high nucleotide diversity, including structural variability and gene presence-absence

polymorphisms. Based on models of codon sequence polymorphism and divergence, the

analyses presented here provide evidence for signatures of both positive directional and

diversifying selection in protein coding genes, as expected under arms race and trench

warfare scenarios, respectively. The rate of adaptive substitution in the plant pathogen was

found to be similar to estimates in animal species and considerably faster than corresponding

estimates in plants. Notably, rates of evolution in genes encoding effector proteins were

found to be more than twice as fast as the genome average. Furthermore, our results suggest

widespread occurrence of linked selection (both Hill-Robertson interference and background

selection), as both positively and negatively selected loci were found to impact the genetic

diversity at linked sites. These results highlight the fundamental role of recombination and

sexual reproduction in adaptive processes of rapidly evolving organisms.

Materials and Methods

Re-sequencing, assembly and alignment of *Z. tritici* isolates

In this study we used a geographical collection of thirteen field isolates of *Z. tritici* isolated

from infected leaves of bread wheat (*Triticum aestivum*) (Table S1). Genome data of three

isolates, including the reference isolate IPO323, were published previously [16,19]. For the

remaining ten isolates full genomes were sequenced. DNA extraction was performed as

previously described [16]. Library preparation and paired end sequencing using an Illumina

HiSeq2000 platform were conducted at Aros, Skejby, Denmark. Sequence data of the ten

isolates has been deposited under the NCBI BioProject IDs PRJNA312067. We used

SOAPdenovo2 [30] to construct *de novo* genome assemblies for each isolate independently.

For each genome, the k-mer value maximizing the weighted median (N50) of contigs and

scaffolds was selected.

Prior to generating a multiple genome alignment, we pre-processed the individual genomes

of the thirteen *Z. tritici* isolates. First, we masked repetitive sequences using a library of 497

repeat families identified *de novo* in four *Zymoseptoria* species [19]. Repeats were soft-
masked using the program RepeatMasker (option -xsmall) to retained information of repeat
sites in the alignment (A.F.A. Smit, R. Hubley & P. Green RepeatMasker at
http://repeatmasker.org). Second, we filtered the genome assemblies to contain only contigs
with a length >= 1 kb. Multiple genome alignments were generated by the MULTIZ program
using the LASTZ pairwise aligner from the Threaded Blockset Aligner (TBA) package [69]. The
alignment was projected on the IPO323 reference genome using the *maf_project* program
from the TBA package.

Inference of population structure
In order to infer population structure, we generated genealogies of the thirteen isolates using
the multiple genome alignment. We used the MafFilter program [70] to compute pairwise
distance matrices using maximum likelihood under a Kimura 2 parameter model in 10 kb
sliding windows along the chromosomes of the reference genome. For each window, a BioNJ
tree was reconstructed from the distance matrices. The resulting 1,850 genealogies were
used to build a super tree using SDM for the generation of a distance supermatrix [71] and
FastME was used to infer a consensus tree [72]. We used the program ADMIXTURE [28] to
estimate the number of ancestral populations based on single nucleotide polymorphism data.
Input files for ADMIXTURE were generated from the filtered alignment using MafFilter [70].

Presence-absence patterns of chromosomes and genes
Based on the multiple genome alignment, the numbers of aligned bases for each genome
were computed in 10 kb non-overlapping sliding windows along the alignment projected on
the reference sequence. To assess the presence-absence of protein coding genes we used a
recently updated annotation of the reference genome of *Z. tritici* IPO323 [20]. Nucleotide
sequences corresponding to genes were used as queries for BLASTn searches against each of
the re-sequenced genomes. The presence of each gene in the individual genomes was
assessed if the best BLAST hit showed at least 90% (and up to 110%) of coverage of the total
length of the IPO323 gene.

Estimation of rates of adaptation
Based on the coordinates of each predicted gene model in the reference genome IPO323
[19,20], exons were extracted from the multiple genome alignment of *Z. tritici* isolates using
MafFilter [70]. Complete coding sequences (CDS) were concatenated to generate individual

alignments of all orthologous CDS. If one or more exons were not extracted from the

alignment due to missing information, the gene was discarded from further analyses. Each

complete CDS alignment was filtered according to the following criteria: (i) CDS were

discarded if they contained more than 5% gaps in one or more individuals, (ii) CDS with

premature stop codon were likewise deleted, and (iii) only alignments comprising three or

more CDS were kept. In some cases, due to indels in the genome alignment, the codon

phasing of some genes was lost. This issue was overcome by refining the CDS alignment using

the codon-based multiple alignment program MACSE [73]. The final data set contains 7,040

gene alignments. We further created a data set containing an outgroup sequence, taken from

Z. ardabiliae, leading to 6,767 alignments. The data set with outgroup was used to infer the

synonymous and non-synonymous divergence based on the rate of synonymous and non-

synonymous substitutions. The synonymous and non-synonymous unfolded site frequency

spectra (SFS) were also computed, using the outgroup sequence to orient mutations. All

calculations were performed using the BppPopStats program from the Bio++ Program Suite

[74]. We used the Dfem program in order to estimate the distribution of fitness effects from

the SFS and compute a genome wide estimate of α and ω_a, the proportion of mutations fixed

by selection and the rate of adaptive substitutions respectively [12]. The following models

were fitted and compared using Akaike's information criterion: Neutral, Gamma Gamma-

Exponential, Displaced Gamma, Scaled Beta and Bessel K. Analyses were conducted on the

complete set of gene alignments, as well as on a subset of genes encoding effector candidates

only. We further stratified our data set according to the flanking recombination rate of each

gene in 500 bp 5' and 3' flanking regions [25]. We discretized the observed distribution of the

population recombination rate ρ in 15 categories with similar number of genes, each category

comprising between 676 and 749 genes. For each of the 15 gene sets, 10 bootstrap replicates

were generated by sampling genes randomly in each category. Genes in each replicate were

concatenated and the Dfem program run with the GammaExpo distribution of fitness effect

[12]. For each recombination category, the mean estimates of α and ω_a, as well as the

standard error over the 10 replicates, were computed and plotted (Figure 2). Kendall

correlation tests were performed on the 15 times 10 estimates in order to correlate both α

and ω_a with ρ.

Model of codon sequence evolution and candidate genes under positive diversifying

selection

We used all filtered CDS alignments to reconstruct genealogies for the individual genes using

PhyML (model HKY85) [75]. To investigate patterns of selection and infer the role of positive

selection on adaptive gene evolution, the program CodeML from the PAML package was used

[46] with the filtered multiple CDS alignments and the corresponding phylogenetic trees as

inputs. CodeML allows inference of selection and evolutionary rates by calculating the

parameter ω, the ratio of non-synonymous to synonymous rates (dN/dS) for each gene. More

specifically, we compared site models that allow ω to vary among codons in the protein [18].

The models used in this study include the nearly neutral (M1a), positive selection (M2a), beta

(M7), and beta&ω (M8) models. A likelihood ratio test (LRT) was used to compare the fit of

null models and alternative models, and the significance of the LRT statistic was determined

using a χ^2 distribution. The first LRT tests for the occurrence of sites under positive selection

by comparing the M1a and M2a models. In the model M1a sites can be under purifying

selection ($0 < \omega < 1$) and evolve by neutral evolution ($\omega = 0$) while the M2a model allows for

some sites to be under positive selection ($\omega > 1$). The second LRT compares the M7 and M8

models, where in M7 a discretized beta distribution for ω (limited to the interval [0,1]) over

all sites is assumed, while the model M8 allows sites with $\omega > 1$. By allowing for a wider range

of strength of purifying selection, the M7 and M8 models are more biologically realistic. The

M7-M8 LRT, however, was shown to display an increased false discovery rate compared to

the M1a-M2a comparison [47]. For these two tests, p-values were obtained with a χ^2 test (df

= 2). We corrected for multiple testing and a false discovery rate of 1% was used for the

detection of genes under positive selection [76]. Only genes significant for both tests were

considered as genes evolving under positive selection.

To address divergent adaptation, we compared gene evolution among four closely related

Zymoseptoria species. In a previous study we defined the core proteome of *Z. tritici*, *Z.*

ardabiliae, *Z. brevis* and *Z. pseudotritici* comprising 7,786 orthologous genes [20]. We

generated alignments of the corresponding coding sequences using the MACSE sequence

aligner [73] and used CodeML with a branch model that allows ω to vary among branches of

the phylogeny [18]. As input we applied a non-rooted tree of the four *Zymoseptoria* species as

published in [77]. Branch lengths were re-estimated for each gene by CodeML [46].

Functional enrichment analysis and gene cluster analysis

PFAM domains were extracted from Interproscan results from [20]. Only domain hits with e-

values lower than 1e-5 were considered resulting in 10,026 domains present in 7,343 genes.

Enrichment tests were performed based on contingency tables, counting the number of

genes containing the domain and the number of genes which do not contain it, for both the

complete proteome and a given set of candidates to test. A χ^2 test was performed to assess

significance.

To analyse the distribution of genes under positive selection, we considered two genes

separated by less than 5,000 bp to be clustered and assessed the probability of such clusters

under a random distribution of genes along the chromosomes. To do so on a genome-wide

scale, we calculated the probability to obtain clusters encompassing from two to ten genes

under positive selection when these genes are randomly distributed across all gene

coordinates. Based on 10,000 random permutations, it appeared that only clusters containing

more than three genes were significant at the 5% level.

Genome-wide analysis of selection patterns

We compared the selection regime of each gene to several variables. Local recombination

rate was computed using the fine-scale recombination map of Stukenbrock and Dutheil [25]

as the average in the 500 bp 5' and 3' flanking regions. Local gene density was computed as

the proportion of coding sites in 10 kb upstream and downstream regions. Similarly, the

density of TEs was computed as the proportion of sites annotated as TE in a windows starting

10 kb upstream the gene and ending 10 kb downstream the gene, allowing for TE located

within the gene itself. GC content at third codon position (GC3) was also computed.

Expression levels were calculated as in [50]. Mean expression level was computed as the

maximum value observed for the gene in axenic culture or plant infection, each averaged

over three biological replicates. Genes encoding secreted proteins were predicted by the

program SignalP [78]. Genes with no homolog in any other species were considered as

"orphan" genes and genes located on accessory chromosomes were considered as

"dispensable".

A linear model was defined with the strength of purifying selection ($\omega 0$) as a response

variable, together with all individual explanatory variables. We further included an interaction

of gene density and recombination rate, as expected under a background selection model and

an interaction between protein length given that orphan genes are reported to be

significantly shorter than other genes [48,49]. Genes with $\omega 0 < 0.001$ or $\omega 0 > 999.0$ were

discarded for further analyses as these estimates most likely indicate model fitting issues,

possibly due to lack of data. Recombination rate and protein length variables were log

transformed as they showed extreme distribution. Expression values were log(x+1)

transformed. A BoxCox transform was used to normalize the model residues. After

normalization, residues displayed a good fit to a normal distribution (Shapiro-Wilk normality

test, p-value = 0.07534), no significant heteroscedasticity (Harrison-McCabe test, p-value =

0.0837), and model residues were found to be independent (Box-Ljung test, p-value =

0.1982). A A binary logistic regression was fitted in order to predict the occurrence of genes

under positive diversifying selection. The result of the LRT on codon models was used as a

response variable, and gene expression, recombination rate, protein length, gene density,

prediction of secretion, location on dispensable chromosome and whether the gene is an

orphan or not were used as explanatory variable, and transformed where needed. We further

included the effect of density of transposable elements (TE). The TE density variable was

highly skewed with an excess of low values, it was therefore transformed into a binary

variable with TE density > 0.2 (1,881 genes) and TE density <= 0.2 (9958 genes). For all linear

models, we used the rms package in R to compute robust regression estimates and the

fastbw to select the factors that contribute significantly to the variance explained by the

model [79]. We selected the 10% *Z. tritici* most expressed genes and computed the relative

synonymous codon usage of every codons [80]. Analyses were conducted using the 'uco'

function of the seqinr package for R [81].

Simulation with recombination

We used the MS program in order to simulated ancestral recombination graphs (ARGs) for

thirteen individuals and a given recombination rate [82]. The resulting ARGs were used to

simulate codon alignments using the bppSeqGen program [83]. Codon sites were simulated

with dN/dS values of 0.1, 0.3, 0.6, 1.0 or 3.0, with equal probability. In this set-up,

recombination was effectively allowed only between codons and not within codons. The

resulting alignments were given as input to the PhyML program [84] with a HKY85

substitution model and a discretized gamma distribution of site-specific rates with 4 classes,

as used for inference of gene data sets in this study. M1a and M2a models were then fitted

using the estimated tree as input with CodeML. Recombination rate of ρ = 0.0, 0.001, 0.01,

0.05 and 0.1, for gene lengths of 100, 500, 1,000 and 2,000 codons with 30 independent

replicates for each combination of parameters. CodeML output was parsed using BioPython

[85].

Substitution mapping

674   In order to assess the putative occurrence of RIP mutations among mutations observed in a

675   given gene, we inferred the amount of CpA → TpA and GpT → ApT dinucleotide substitutions

using probabilistic substitution mapping with a dinucleotide substitution model previously fitted to the data using maximum likelihood [83]. Given a substitution model and a phylogenetic tree, the probabilistic substitution mapping procedure enabled us to count each type of substitution for a site of an alignment and each branch of a phylogeny, while accounting for the uncertainty of the ancestral sequences at the inner nodes of the phylogeny.

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and material: The datasets generated and analysed during the current study are available at NCBI under the BioProject ID PRJNA312067.

Competing interests: The authors declare that they have no competing interests

Funding: The study was funded by a Max Planck fellowship and a personal grant from the State of Schleswig-Holstein, Germany both to EHS.

Authors' contributions: JD and EHS conceived and planned the experiments. JG and JD performed the computational framework and analysed the data. All authors contributed to the interpretation of data and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements: The authors thank Nicolas Galtier and Thomas Bataillon for helpful discussion.

References

1. Möller M, Stukenbrock EH. Evolution and genome architecture in fungal plant pathogens. Nat. Rev. Microbiol. 2017; 15, 771.

2. Lo Presti L, Lanver D, Schweizer G, Tanaka S, Liang L, Tollot M, et al. Fungal effectors and plant susceptibility. Annu. Rev. Plant Biol. Annual Reviews; 2015;66:513–45.

3. Spanu PD, Abbott JC, Amselem J, Burgis TA, Soanes DM, Stüber K, et al. Genome Expansion and Gene Loss in Powdery Mildew Fungi Reveal Tradeoffs in Extreme Parasitism. Science (80-.). 2010;330:1543–6.

4. Klosterman SJ, Subbarao K V, Kang S, Veronese P, Gold SE, Thomma BPHJ, et al.

Comparative Genomics Yields Insights into Niche Adaptation of Plant Vascular Wilt Pathogens.

PLoS Pathog; 2011;7:e1002137.

5. Ma L-J, van der Does HC, Borkovich KA, Coleman JJ, Daboussi M-JM-J, Di Pietro A, et al.

Comparative genomics reveals mobile pathogenicity chromosomes in Fusarium. Nature;

2010;464:367–73.

6. Daverdin G, Rouxel T, Gout L, Aubertot J-N, Fudal I, Meyer M, et al. Genome Structure and

Reproductive Behaviour Influence the Evolutionary Potential of a Fungal Phytopathogen.

PLOS Pathog.; 2012;8:e1003020.

7. Dutheil JY, Mannhaupt G, Schweizer G, Sieber CMK, Münsterkötter M, Güldener U, et al. A

tale of genome compartmentalization: the evolution of virulence clusters in smut fungi.

Genome Biol. Evol. 2016;8:681–704.

8. Kimura M. Preponderance of synonymous changes as evidence for the neutral theory of

molecular evolution. Nature; 1977;267:275.

9. Nei M, Gojobori T. Simple methods for estimating the numbers of synonymous and

nonsynonymous nucleotide substitutions. Mol. Biol. Evol. SMBE; 1986;3:418–26.

10. McDonald JH, Kreitman M. Adaptive protein evolution at the Adh locus in Drosophila.

Nature; 1991;351:652.

11. Eyre-Walker A. The genomic rate of adaptive evolution. Trends Ecol. Evol. Elsevier;

2006;21:569–75.

12. Galtier N. Adaptive protein evolution in animals and the effective population size

hypothesis. PLoS Genet.; 2016;12:e1005774.

13. Smith NGC, Eyre-Walker A. Adaptive protein evolution in Drosophila. Nature;

2002;415:1022–4.

14. Eyre-Walker A, Keightley PD. The distribution of fitness effects of new mutations. Nat Rev

Genet. 2007;8:610–8.

15. Raffaele S, Farrer RA, Cano LM, Studholme DJ, MacLean D, Thines M, et al. Genome

Evolution Following Host Jumps in the Irish Potato Famine Pathogen Lineage. Science (80-.).

2010;330:1540–3.

16. Stukenbrock EH, Bataillon T, Dutheil JY, Hansen TT, Li R, Zala M, et al. The making of a new

pathogen: Insights from comparative population genomics of the domesticated wheat

pathogen Mycosphaerella graminicola and its wild sister species. Genome Res. 2011;21:2157–

66.

17. Charlesworth D. Balancing Selection and Its Effects on Sequences in Nearby Genome

Regions. PLOS Genet. 2006;2:e64.

18. Nielsen R, Yang Z. Likelihood models for detecting positively selected amino acid sites and

applications to the HIV-1 envelope gene. Genetics. Genetics Soc America; 1998;148:929–36.

19. Goodwin SB, M'barek S Ben, Dhillon B, Wittenberg AHJ, Crane CF, Hane JK, et al. Finished

genome of the fungal wheat pathogen Mycosphaerella graminicola reveals dispensome

structure, chromosome plasticity, and stealth pathogenesis. PLoS Genet. 2011;7:e1002070.

20. Grandaubert J, Bhattacharyya A, Stukenbrock EH. RNA-seq Based Gene Annotation and

Comparative Genomics of Four Fungal Grass Pathogens in the Genus Zymoseptoria Identify

Novel Orphan Genes and Species-Specific Invasions of Transposable Elements. G3

Genes|Genomes|Genetics. 2015;

21. Schotanus K, Soyer JL, Connolly LR, Grandaubert J, Happel P, Smith KM, et al. Histone

modifications rather than the novel regional centromeres of Zymoseptoria tritici distinguish

core and accessory chromosomes. Epigenetics Chromatin. BioMed Central; 2015;8:1.

22. Stukenbrock EH, Jørgensen FG, Zala M, Hansen TT, McDonald BA, Schierup MH. Whole-

Genome and Chromosome Evolution Associated with Host Adaptation and Speciation of the

Wheat Pathogen Mycosphaerella graminicola. PLoS Genet; 2010;6:e1001189.

23. Poppe S, Dorsheimer L, Happel P, Stukenbrock EH. Rapidly Evolving Genes Are Key Players

in Host Specialization and Virulence of the Fungal Wheat Pathogen Zymoseptoria tritici

(Mycosphaerella graminicola). PLoS Pathog; 2015;11:e1005055.

24. Croll D, Lendenmann MH, Stewart E, McDonald BA. The Impact of Recombination

Hotspots on Genome Evolution of a Fungal Plant Pathogen. Genetics 2015;201:1213–28.

25. Stukenbrock EH, Dutheil JY. Comparison of fine-scale recombination maps in fungal plant

pathogens reveals dynamic recombination landscapes and intragenic hotspots. bioRxiv. 2017;

26. Linde CC, Zhan J, McDonald BA. Population Structure of Mycosphaerella graminicola:

From Lesions to Continents. Phytopathology; 2002;92:946–55.

27. Banke S, McDonald BA. Migration patterns among global populations of the pathogenic

fungus Mycosphaerella graminicola. Mol. Ecol. 2005;14:1881–96.

28. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated

individuals. Genome Res. Cold Spring Harbor Lab; 2009;19:1655–64.

29. O'Donnell K, Kistler HC, Tacke BK, Casper HH. Gene genealogies reveal global

phylogeographic structure and reproductive isolation among lineages of Fusarium

graminearum, the fungus causing wheat scab. Proc. Natl. Acad. Sci. 2000;97:7905–10.

30. Ali S, Gladieux P, Leconte M, Gautier A, Justesen AF, Hovmøller MS, et al. Origin,

Migration Routes and Worldwide Population Genetic Structure of the Wheat Yellow Rust

Pathogen Puccinia striiformis f.sp. tritici. PLoS Pathog; 2014;10:e1003903.

31. Brown JKM, Hovmøller MS. Aerial dispersal of pathogens on the global and continental

scales and its impact on plant disease. Science. 2002;297:537–41.

32. Mehrabi R, Taga M, Kema GHJ. Electrophoretic and cytological karyotyping of the foliar

wheat pathogen Mycosphaerella graminicola reveals many chromosomes with a large size

range. Mycologia. 2007;99:868–76.

33. Brunner PC, Torriani SFF, Croll D, Stukenbrock EH, McDonald BA. Coevolution and life

cycle specialization of plant cell wall degrading enzymes in a hemibiotrophic pathogen. Mol.

Biol. Evol. 2013;30:1337–47.

34. McDonald MC, McGinness L, Hane JK, Williams AH, Milgate A, Solomon PS. Utilizing Gene

Tree Variation to Identify Candidate Effector Genes in Zymoseptoria tritici. G3

Genes|Genomes|Genetics. 2016;6:779–91.

35. Croll D, Zala M, McDonald BA. Breakage-fusion-bridge cycles and large insertions

contribute to the rapid evolution of accessory chromosomes in a fungal pathogen. PLoS

Genet.; 2013;9:e1003567.

36. Plissonneau C, Stürchler A, Croll D. The Evolution of Orphan Regions in Genomes of a

Fungal Pathogen of Wheat. MBio. Am Soc Microbiol; 2016;7:e01231-16.

37. de Jonge R, Peter van Esse H, Maruthachalam K, Bolton MD, Santhanam P, Saber MK, et

al. Tomato immune receptor Ve1 recognizes effector of multiple fungal pathogens uncovered

by genome and RNA sequencing. Proc. Natl. Acad. Sci. U. S. A.; 2012;109:5110–5.

38. Xue M, Yang J, Li Z, Hu S, Yao N, Dean RA, et al. Comparative Analysis of the Genomes of

Two Field Isolates of the Rice Blast Fungus Magnaporthe oryzae. PLOS Genet.;

2012;8:e1002869.

39. Gossmann TI, Song B-H, Windsor AJ, Mitchell-Olds T, Dixon CJ, Kapralov M V, et al.

Genome wide analyses reveal little evidence for adaptive evolution in many plant species.

Mol. Biol. Evol.; 2010;27:1822–32.

40. Hill WG, Robertson A. The effect of linkage on limits to artificial selection. Genet. Res.;

1966;8:269–94.

41. Marais G, Charlesworth B. Genome Evolution: Recombination Speeds Up Adaptive

Evolution. Curr. Biol.; 2003;13:R68–70.

42. Ellegren H. Genome sequencing and population genomics in non-model organisms. Trends

Ecol. Evol. 2014;29:51–63.

43. Nordborg M, Charlesworth B, Charlesworth D. The effect of recombination on background

selection. Genet. Res. 1996;67:159–74.

44. Romiguier J, Gayral P, Ballenghien M, Bernard A, Cahais V, Chenuil A, et al. Comparative

population genomics in animals uncovers the determinants of genetic diversity. Nature.

2014;515:261–3.

45. Ellegren H, Galtier N. Determinants of genetic diversity. Nat. Rev. Genet.; 2016;

46. Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol. Evol.

2007;24:1586–91.

47. Anisimova M, Bielawski JP, Yang Z. Accuracy and power of the likelihood ratio test in

detecting adaptive molecular evolution. Mol. Biol. Evol.; 2001;18:1585–92.

48. Liao B-Y, Scott NM, Zhang J. Impacts of Gene Essentiality, Expression Pattern, and Gene

Compactness on the Evolutionary Rate of Mammalian Proteins. Mol. Biol. Evol.

2006;23:2072–80.

49. Cai JJ, Petrov DA. Relaxed purifying selection and possibly high rate of adaptation in

primate lineage-specific genes. Genome Biol. Evol. 2010;2:393–409.

50. Kellner R, Bhattacharyya A, Poppe S, Hsu TY, Brem RB, Stukenbrock EH. Expression

Profiling of the Wheat Pathogen Zymoseptoria tritici Reveals Genomic Patterns of

Transcription and Host-Specific Regulatory Programs. Genome Biol. Evol. 2014;6:1353–65.

51. Drummond DA, Raval A, Wilke CO. A single determinant dominates the rate of yeast

protein evolution. Mol. Biol. Evol. 2006;23:327–37.

52. Charlesworth B, Morgan MT, Charlesworth D. The effect of deleterious mutations on

neutral molecular variation. Genetics. Genetics Soc America; 1993;134:1289–303.

53. Campos JL, Halligan DL, Haddrill PR, Charlesworth B. The relation between recombination

rate and patterns of molecular evolution and variation in Drosophila melanogaster. Mol. Biol.

Evol. 2014;31:1010–28.

54. Duret L. Evolution of synonymous codon usage in metazoans. Curr. Opin. Genet. Dev.

2002;12:640–9.

55. Birdsell JA. Integrating genomics, bioinformatics, and classical genetics to study the effects

of recombination on genome evolution. Mol Biol Evol 2002;19.

56. Duret L, Mouchiroud D. Expression pattern and, surprisingly, gene length shape codon

usage in Caenorhabditis, Drosophila, and Arabidopsis. Proc. Natl. Acad. Sci. 1999;96:4482–7.

57. André I, Potocki-Véronèse G, Barbe S, Moulis C, Remaud-Siméon M. CAZyme discovery

and design for sweet dreams. Curr. Opin. Chem. Biol. 2014;19:17–24.

58. Howlett BJ. Secondary metabolite toxins and nutrition of plant pathogenic fungi. Curr.

Opin. Plant Biol. 2006;9:371–5.

59. Van den Ackerveken GFJM, Vossen P, De Wit J. The AVR9 race-specific elicitor of

Cladosporium fulvum is processed by endogenous and plant proteases. Plant Physiol. Am Soc Plant Biol; 1993;103:91–6.

60. Stergiopoulos I, van den Burg HA, Ökmen B, Beenen HG, van Liere S, Kema GHJ, et al. Tomato Cf resistance proteins mediate recognition of cognate homologous effectors from fungi pathogenic on dicots and monocots. Proc. Natl. Acad. Sci. 2010;107:7610–5.

61. Cambareri EB, Jensen BC, Schabtach E, Selker EU. Repeat-induced GC to AT mutations in Neurospora. Science (80-.). American Association for the Advancement of Science; 1989;244:1571–5.

62. Idnurm A, Howlett BJ. Analysis of loss of pathogenicity mutants reveals that repeat-induced point mutations can occur in the Dothideomycete Leptosphaeria maculans. Fungal Genet. Biol. 2003;39:31–7.

63. Laurie JD, Ali S, Linning R, Mannhaupt G, Wong P, Guldener U, et al. Genome Comparison of Barley and Maize Smut Fungi Reveals Targeted Loss of RNA Silencing Components and Species-Specific Presence of Transposable Elements. Plant Cell. 2012;24:1733–45.

64. Hood ME, Katawczik M, Giraud T. Repeat-induced point mutation and the population structure of transposable elements in Microbotryum violaceum. Genetics. Genetics Soc America; 2005;170:1081–9.

65. Poppe S, Dorsheimer L, Happel P, Stukenbrock EH. Rapidly Evolving Genes Are Key Players in Host Specialization and Virulence of the Fungal Wheat Pathogen Zymoseptoria tritici (Mycosphaerella graminicola). PLoS Pathog. 2015;11:e1005055.

66. Marshall R, Kombrink A, Motteram J, Loza-Reyes E, Lucas J, Hammond-Kosack KE, et al. Analysis of two in planta expressed LysM effector homologs from the fungus Mycosphaerella graminicola reveals novel functional properties and varying contributions to virulence on wheat. Plant Physiol. 2011;156:756–69.

67. de Jonge R, van Esse HP, Kombrink A, Shinya T, Desaki Y, Bours R, et al. Conserved fungal LysM effector Ecp6 prevents chitin-triggered immunity in plants. Science (80-.); 2010;329:953–5.

68. Kulkarni RD, Thon MR, Pan H, Dean RA. Novel G-protein-coupled receptor-like proteins in the plant pathogenic fungus Magnaporthe grisea. Genome Biol. BioMed Central; 2005;6:R24.

69. Blanchette M, Kent WJ, Riemer C, Elnitski L, Smit AFA, Roskin KM, et al. Aligning Multiple Genomic Sequences With the Threaded Blockset Aligner. Genome Res. 2004;14:708–15.

70. Dutheil JY, Gaillard S, Stukenbrock EH. MafFilter: a highly flexible and extensible multiple genome alignment files processor. BMC Genomics. 2014;15:53.

71. Criscuolo A, Berry V, Douzery EJP, Gascuel O. SDM: a fast distance-based approach for

(super) tree building in phylogenomics. Syst. Biol.; 2006;55:740–55.

72. Lefort V, Desper R, Gascuel O. FastME 2.0: a comprehensive, accurate, and fast distance-

based phylogeny inference program. Mol. Biol. Evol.; 2015;32:2798–800.

73. Ranwez V, Harispe S, Delsuc F, Douzery EJP. MACSE: Multiple Alignment of Coding

SEquences accounting for frameshifts and stop codons. PLoS One. 2011;6:e22594.

74. Guéguen L, Gaillard S, Boussau B, Gouy M, Groussin M, Rochette NC, et al. Bio++: efficient

extensible libraries and tools for computational molecular evolution. Mol. Biol. Evol.;

2013;30:1745–50.

75. Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large

phylogenies by maximum likelihood. Syst. Biol.; 2003;52:696–704.

76. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful

approach to multiple testing. J. R. Stat. Soc. Ser. B.; 1995;289–300.

77. Stukenbrock EH, Quaedvlieg W, Javan-Nikhah M, Zala M, Crous PW, McDonald BA.

Zymoseptoria ardabiliae and Z. pseudotritici, two progenitor species of the septoria tritici leaf

blotch fungus Z. tritici (synonym: Mycosphaerella graminicola). Mycologia. 2012;104:1397–

407.

78. Petersen TN, Brunak S, Heijne G, Nielsen H. SignalP 4.0: discriminating signal peptides

from transmembrane regions. Nat Methods. 2011;8.

79. Harrell F. Regression modeling strategies: with applications to linear models, logistic and

ordinal regression, and survival analysis. Springer; 2015.

80. Sharp PM, Tuohy TM, Mosurski KR. Codon usage in yeast: cluster analysis clearly

differentiates highly and lowly expressed genes. Nucleic Acids Res. 1986;14:5125–43.

81. R Core Team. R: A language and environment for statistical computing. Vienna, Austria;

2013.

82. Hudson RR. Generating samples under a Wright–Fisher neutral model of genetic variation.

Bioinformatics. 2002;18:337–8.

83. Dutheil J, Boussau B. Non-homogeneous models of sequence evolution in the Bio++ suite

of libraries and programs. BMC Evol. Biol. 2008;8:1.

84. Guindon S, Lethiec F, Duroux P, Gascuel O. PHYML Online—a web server for fast

maximum likelihood-based phylogenetic inference. Nucleic Acids Res.; 2005;33:W557–9.

85. Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, et al. Biopython: freely

available Python tools for computational molecular biology and bioinformatics.

Bioinformatics. 2009;25:1422–3.

Tables

Table 1: Estimates of the proportion of adaptive mutation (α) under various models of distribution of fitness effects with the DFEM method. CEG: candidate effector genes, α: proportion of adaptive substitutions, ω_a: rate of adaptive substitutions.

Data set	Model	Nb. parameters	Log likelihood	AIC	alpha	omegaA
All genes	Neutral	16	-1673,86	3379,72	0,26	0,03
All genes	Gamma	17	-256,14	546,28	0,5	0,05
All genes	GammaExpo	19	-237,73	**513,46**	**0,44**	**0,05**
All genes	DisplacedGamma	18	-256,14	548,28	0,5	0,05
All genes	ScaledBeta	18	-303,8	643,6	0,52	0,05
All genes	BesselK	19	-294,94	627,88	0,42	0,04
CEG	Neutral	16	-102,64	237,28	0,32	0,07
CEG	Gamma	17	-90,05	214,1	0,52	0,11
CEG	GammaExpo	19	-84,6	**207,2**	**0,68**	**0,14**
CEG	DisplacedGamma	18	-90,41	216,82	0,53	0,11
CEG	ScaledBeta	18	-86,69	209,38	0,6	0,12
CEG	BesselK	19	-86,41	210,82	0,5	0,1

Table 2: Genomic factors affecting the strength of purifying selection.

Effect	Coefficient	S.E.	P-value
Intercept	2,4900	0,3580	<0.0001
Expression	-0,0602	0,0140	<0.0001
Recombination rate	-0,2030	0,0100	<0.0001
Protein size	-0,3749	0,0353	<0.0001
GC 3rd codon position	-5,2429	0,3692	<0.0001
Orphan	0,6175	0,1401	<0.0001

Table 3: Genomic factors and occurrence of positive diversifying selection.

Effect	Coefficient	S.E.	P-value
Intercept	-8,8321	0,6100	<0.0001
Expression	-0,3574	0,0388	<0.0001
Recombination rate	0,1991	0,0284	<0.0001
Protein size	1,3408	0,0954	<0.0001

Figures

Figure 1: Population structure of the thirteen *Z. tritici* isolates. A) Consensus super tree of
the thirteen isolates based on 1,850 genealogies estimated in 10 kb sliding windows along the
multiple genome alignment. This tree suggests the grouping of the isolates into two
populations originating from Europe and Iran. B) Based on SNP data, the program
ADMIXTURE estimated that the best separation of the isolates is also in two populations
(k=2). However, the use of k=3 highlighted a German sub-population within the European
isolates.

Figure 2: Estimates of A) the proportion of adaptive substitution α, and B) the rate of adaptive
substitution, ω_a as a function of the population recombination rate (ρ). Each point and bars
represent the mean estimate and corresponding standard error for one recombination
category over 10 bootstrap replicates. The blue lines show the fitted linear models.

945
946
947

Figure 3: Patterns of selection along the genome of Z. *tritici*. Recombination, intensity of
purifying selection (as estimated from the ω0 parameter, see Methods), density of coding
sites (CDS) and density of transposable elements (TE) are plotted in windows of 150 kb along
the thirteen essential chromosomes.

Figure 4: Correlation of the strength of purifying selection with several genomic factors. The intensity of purifying selection is measured by the ω0 parameter (see Methods). TE = transposable elements. Points represent median values and error bar the first and third quartiles of the distributions. A-E: x-axis were discretized in categories with equal point densities for clarity of visualization. Lines represent first, median and third quantile regression on non-discretized data.

Figure 5: Branch-specific signatures of divergent adaptation in *Zymoseptoria* species.
Phylogeny of four *Zymoseptoria* species based on integrating the mean dN / dS ratio (in red)
and the number of positively selected genes (in green) estimated from the 7,786 genes
shared by the species using the branch model implemented in PAML [46].

Supplementary materials

Table S1: Summary table of isolates used in this study and genome assembly statistics.

Table S2: Summary statistics of the multiple genome alignment of thirteen *Z. tritici* genomes.

Table S3: Output of the PAML analysis using codon site models for the 9,412 filtered CDS of *Z. tritici*.

Table S4: Functional enrichment analysis using PFAM domains for the 786 genes with sites under positive selection in *Z. tritici*.

Table S5: Representation of RIP mutations in genes according to their distance to transposable elements.

Table S6: Functional enrichment analysis using PFAM domains for the genes under positive selection in four *Zymoseptoria* species.

Figure S1: Chromosome polymorphism in the twelve re-sequenced genomes compared to the IPO323 reference genome. Based on the multiple genome alignment, for each re-sequenced genomes the alignment coverage was calculated in 10 kb sliding windows along the total length of each chromosome using the genome of IPO323 as a reference. Chromosomes 1 to 13 (in blue) are well conserved among the different isolates and constitute the "core" chromosomes while the chromosomes 14 to 21 (in red) appear to be absent or incomplete at different frequencies.

Figure S2: Impact of recombination on the estimation of the strength of purifying selection using phylogenetic models of codon sequence evolution. Parameters are the ratio of transitions over transversions (kappa), the ratio of the rates of synonymous over non-synonymous mutations for sites evolving under purifying selection ($\omega 0$) in the two models M1 and M2, and the ratio of the rates of synonymous over non-synonymous mutations for sites evolving under positive selection ($\omega 1$) for model M2 (see methods). Rho: scaled recombination rate used in the simulations. Length: length of the gene used in the simulation. X-axis: average number of distinct topologies per simulated site, increasing with the recombination rate. Y-axis: parameter estimate.

Figure S3: Codon usage in *Z. tritici*. Relative synonymous codon usage (RSCU) in the 10% most expressed genes of *Z. tritici*. Codon usage, according to the base type at the third position.