1 **Positively selected effector genes and their contribution to virulence in the smut fungus**
2 ***Sporisorium reilianum***
3

4 Gabriel Schweizer[1], Karin Münch[1], Gertrud Mannhaupt[1,2], Jan Schirawski[1,3], Regine
5 Kahmann[1*] and Julien Y. Dutheil[1,4,5*]
6
7 [1]Max-Planck-Institute for Terrestrial Microbiology, Department of Organismic Interactions,
8 Karl von Frisch-Straße 10, 35043 Marburg, Germany.
9
10 [2]Helmholtz Zentrum München, Institute for Bioinformatics and Systems Biology, Ingolstädter
11 Landstraße 1, 85764 Neuherberg, Germany.
12
13 [3]Current address: RWTH Aachen, Institute of Applied Microbiology, Microbial Genetics,
14 Worringer Weg 1, 52074 Aachen, Germany.
15
16 [4]University of Montpellier 2, Institute of Evolutionary Sciences, Place Eugène Bataillon, 34095
17 Montpellier, France.
18
19 [5]Current address: Max-Planck-Institute for Evolutionary Biology, Research Group Molecular
20 Systems Evolution, August-Thienemann-Straße 2, 24306 Plön, Germany.
21
22
23 [*]Authors for correspondence:
24
Dr. Julien Y. Dutheil
Max-Planck-Institute for Evolutionary Biology
Research Group Molecular Systems Evolution
August-Thienemann-Straße 2
24306 Plön
Germany
Phone: +49 4522 763 298
Fax: +49 4522 763 281
Email: dutheil@evolbio.mpg.de
34
Prof. Dr. Regine Kahmann
Max-Planck-Institute for Terrestrial Microbiology
Department of Organismic Interactions
Karl-von-Frisch Straße 10
35043 Marburg
Germany
Phone: +49 6421 178 501
Fax: +49 6421 178 509
Email: kahmann@mpi-marburg.mpg.de

44
45
46

Abstract

Plants and fungi display a broad range of interactions within natural and agricultural ecosystems ranging from symbiosis to parasitism. Pathogenic interactions are governed by secreted fungal effector proteins, which are thought to coevolve with their host targets. Biotrophic smut fungi which belong to the division of Basidiomycota are well-suited to investigate the evolution of plant pathogens, because several quality draft genomes and genetic tools are available for these species. Here, we used the genomes of *Sporisorium reilianum* f. sp. *zeae* and *S. reilianum* f. sp. *reilianum,* two closely related formae speciales infecting maize and sorghum, respectively, together with the genomes of *Ustilago hordei*, *U. maydis* and *S. scitamineum* to identify effector genes showing signs of positive selection. The largest numbers of such genes were identified in the two pathovariants of *S. reilianum* and between paralogues in *U. hordei*, where many belong to families showing species-specific expansions. Next, we assessed the contribution to virulence of candidate effector genes in *S. reilianum* f. sp. *zeae* by deletion of individual genes in a solopathogenic strain. While eight of nine deletions mutants were unaffected in virulence, one mutant had lost virulence. This shows that despite the relatively recent divergent time of the two formae speciales, a signature of positive selection in candidate effector genes in *S. reilianum* is a poor indicator for the identification of genes with virulence functions.

Key words

Positive selection; effector evolution; smut fungi; comparative genomics; virulence

Introduction

Plants and fungi have a long history of coevolution since the emergence of pioneering land plants approximately 400 million years ago. The development of early plants was likely supported by associations with symbiotic fungi, as suggested by analyses of ribosomal RNAs and fossil records (Gehrig et al. 1996; Martin et al. 2017; Remy et al. 1994). Different forms of plant-fungus interactions evolved, including mutualistic symbiosis where both plant and fungus benefit (Parniske 2008), and pathogenic interactions where fungal colonization greatly reduces plant fitness (Dean et al. 2012). Pathogenic interactions play critical roles in natural and agricultural ecosystems, and understanding the evolutionary mechanisms shaping such interactions is an active field of research (Bagchi et al. 2014; Fisher et al. 2012). Pathogenic interactions are governed by secreted fungal effector proteins, which can protect and shield growing hyphae, suppress plant defense responses and change plant physiology to support growth of the pathogen. Many effector proteins lack known functional domains, and their expression is linked to plant colonization (Franceschetti et al. 2017; Lanver et al. 2017; Lo Presti et al. 2015; Toruño et al. 2016). Effector proteins and their plant targets are thought to evolve either in an arms race or a trench-warfare scenario. In the former, effectors manipulating the host for the benefit of the pathogen are selected, whereas plant targets evolve to detect effectors or inactivate their function (Rovenich et al. 2014). In the latter scenario, sets of alleles are maintained by balancing selection on both host and pathogen sides (Brown and Tellier 2011; Tellier et al. 2014). Several methods are in place for identifying genomic regions under selection (Aguileta et al. 2010; Aguileta et al. 2009; Nielsen 2005). These approaches detect genes subject to positive selection in host and pathogen genomes, and it is thought that genes with signs of positive selection have important functions during host pathogen interaction or have contributed to host specialization (Aguileta et al. 2009). Depending on the aim of an investigation, studies identifying genes with signatures of positive selection are carried out

within or between species. Whereas studies on the population level focus on recent and ongoing

selective processes, and are therefore instrumental in understanding the process of adaptation,

comparative genomic studies employing different species encompass a broader time span and

can for instance help to unravel the genetic basis of host specialization (Plissonneau et al. 2017).

Indeed, positive selection studies in a number of plant pathogen systems revealed that genes

encoding secreted effector proteins are enriched in signatures of positive selection. Such studies

include investigations in diverse plant pathogens like *Microbotryum* species causing anther-

smut disease of Caryophyllaceae species (Aguileta et al. 2010), the wheat pathogen

Zymoseptoria graminicola (Stukenbrock et al. 2011), the rust fungus *Melampsora larici-*

populina (Hacquard et al. 2012), the rice blast fungus *Magnaporthe oryzae* (Huang et al. 2014),

the Irish potato famine pathogen *Phytophthora infestans* (Dong et al. 2014), and a group of

smut fungi parasitizing different grasses and a dicot host (Sharma et al. 2014). Links between

genes showing signatures of positive selection and the actual function being selected are only

beginning to be understood. For example, the population genomics study in the wheat pathogen

Z. tritici which identified candidate effector genes under positive selection (Stukenbrock et al.

2011) was followed up experimentally, and in this case it was shown that the deletion of three

of four candidate genes reduced virulence (Poppe et al. 2015). In the grey mold fungus *Botrytis*

cinerea four positively selected genes were deleted without affecting virulence and this finding

was attributed to functional redundancy, the limited number of tested host plants or

experimental conditions which are different from natural infections (Aguileta et al. 2012). An

elegant study with the oomycete effector protein EpiC1 showed that a single amino acid

substitution under positive selection leads to targeting different host proteases determining host

specificity (Dong et al. 2014). Many studies stop at the level of scanning for positive selection

or conduct only initial functional experiments like gene expression profiling (Cantu et al. 2013;

Guyon et al. 2014; Hacquard et al. 2012; Joly et al. 2010; Sharma et al. 2014; Sharma et al.

2015; Silva et al. 2015; Sperschneider et al. 2015; Sperschneider et al. 2014). Therefore, it

remains largely open whether genes with signatures of positive selection in general play a role

in virulence and/or host specificity.

Smut fungi belonging to the division of Basidiomycota are an important group of about 2,500

species parasitizing mostly grasses, including important crops like maize, sorghum, oat, barley

and sugarcane (Vánky 2012). In smut fungi, sexual reproduction is linked to pathogenic

development and smut fungi therefore depend on successful plant colonization to complete their

life cycle. As biotrophic pathogens, they require living plant tissue for establishing a successful

interaction (Martinez-Espinoza et al. 2002). With few exceptions like *Ustilago maydis*, smut

fungi usually develop symptoms only in the female or male inflorescence of their respective

host plants. During the last ten years, quality draft genome sequences of prominent species were

obtained, among which are *U. maydis*, the causative agent of smut disease on maize and teosinte

(Kämper et al. 2006), *Ustilago hordei* infecting barley (Laurie et al. 2012), *Sporisorium*

scitamineum parasitizing sugarcane (Dutheil et al. 2016; Que et al. 2014; Taniguti et al. 2015)

and *Sporisorium reilianum* causing head smut (Schirawski et al. 2010). *S. reilianum* occurs in

two formae speciales that infect maize (*S. reilianum* f. sp. *zeae*) and sorghum *(S. reilianum* f.

sp. *reilianum*), respectively (Zuther et al. 2012). The speciation of *U. hordei*, *U. maydis*, *S.*

scitamineum and *S. reilianum* is hypothesized to have occurred between seven and 50 million

years ago (Munkacsi et al. 2007). The availability of genome sequences of several species with

different host ranges, together with established tools for genetic manipulations (Brachmann et

al. 2004; Kämper 2004; Khrunyk et al. 2010; Schuster et al. 2016) make this group of smut

fungi particularly interesting for studying the evolution of effector genes as well as their

contributions to virulence, speciation and host specificity. Here, we employed the genome of

the recently sequenced strain *S. reilianum* f. sp. *reilianum* SRS1_H2-8

(http://www.ebi.ac.uk/ena/data/view/LT795054-LT795076) (Zuther et al. 2012) together with

the genomes of *U. maydis*, *U. hordei*, *S. scitamineum* and *S. reilianum* f. sp. *zeae* to identify

potential effector genes with signatures of positive selection. To determine whether such

candidate effectors play a role in virulence, they were individually deleted in *S. reilianum f. sp.*

zeae and the phenotype of the deletion strain assessed after infection of maize. Only one of nine

tested candidate effectors showed a strong contribution to virulence, suggesting that genes with

signatures of positive selection are rarely selected for traits related to virulence in *S. reilianum*.

Materials and Methods

Construction of protein families

Fungal species used in this study, their number of gene models, number of predicted secreted

proteins and sources of genome data are listed in supplementary table S1. The predicted

proteome of the five smut *Ustilago hordei*, *U. maydis*, *Sporisorium scitamineum*, *S. reilianum*

f. sp. *zeae* and *S. reilianum* f. sp. *reilianum* were used to perform an all-against-all blastp search

(Altschul et al. 1990). To identify settings for coverage and identity resulting in the maximum

number of families comprised of 1:1 orthologues (*i.e.* families that have an equal number of

members in each species), SiLiX (Miele et al. 2011) was run with a range for coverage and

identity between 5 % and 95 % in 5 % steps. An identity of 40 % and coverage between 5 %

and 45 % lead to the maximum number of families with 1:1 orthologues (5,394; supplementary

fig. S1) while settings with 40 % identity and 80 % coverage lead to 5,326 families with 1:1

orthologues (supplementary fig. S1). Since using a higher coverage had only a cost of 68 core

families, the stricter criteria were applied for family clustering. Families with at least two

members were aligned on the codon level using MACSE 1.01b (Ranwez et al. 2011) and on the

protein level using PRANK v.100802 (Löytynoja and Goldman 2008). Only consistently found

alignment sites with a maximum of 30 % gaps were used for further analysis.

Estimation of genome-wide divergence values and divergence times

The five genomes of *U. hordei*, *U. maydis*, *S. scitamineum*, *S. reilianum* f. sp. *zeae* and *S.*

reilianum f. sp. *reilianum* were aligned using the multiz genome aligner from the TBA package

(Blanchette et al. 2004). The resulting multiple genome alignment was further processed to

remove coding and ambiguously aligned regions. More specifically, the following filters were

applied:

- Alignments blocks with paralogous sequences were discarded.

- Aligned regions containing sequences annotated as part of a protein coding gene in any of the

five species were discarded.

This led to an alignment of 2.1 million nucleotides.

- A window of 5 nucleotide long was slid along the alignment by 1 nucleotide. Windows

containing at least 2 indels were discarded.

- A window of 50 nucleotide long was slid along the alignment by 1 nucleotide. Windows

containing more than 150 gaps were discarded.

- A window of 300 nucleotide long was slid along the alignment by 1 nucleotide. Windows

containing more than 500 gaps were discarded.

- Finally only alignment blocks of at least 300 nucleotides were kept.

The final alignment had a total length of 936,984 nucleotides. Given the strict criteria used to

filter the alignment, the divergence estimates obtained for intergenic sequences might represent

an underestimate of the true divergence times.

Gene families with exactly one member in each species were concatenated and pairwise protein

sequence similarities computed using the seqinr package (Charif and Lobry 2007) for R. Protein

alignments were also used to infer dates of divergence, using a relaxed clock model. The

PhyloBayes version 4.1 software (Lartillot et al. 2009) was used with the auto-correlated model

of Thorne et al. (Thorne et al. 1998) under a GTR + CAT model. Two calibration points were

used, based on the speciation times of the host lineages: the split between *U. maydis* and the

Sporisorium species was assumed to have occurred between 11 and 24 million years (Myr) ago,

corresponding to the estimated speciation time of the maize lineage with the sugarcane lineage,

and the split between *U. hordei* and the other species was constrained to have occurred between

24 and 70 Myr ago (Munkacsi et al. 2007). Uniform priors were used on these intervals for the

Monte-Carlo Markov Chain. As convergence issues arise when large alignments (more than

20k positions) are used, we followed the PhyloBayes authors' recommendation and performed

a jackknife procedure and generated three datasets of ca 20,000 amino acids by randomly

sampling families and concatenating the corresponding alignments. Two chains were run in

each case and convergence assessed. Sampling was performed after a burning of 10,000

iterations, and every 10 subsequent iterations. Chains were run to ensure that the minimum

effective sample size was greater than 30 and maximum relative difference lower than 0.3 in at

least one sample. Results are summarized in supplementary table S2 and supplementary fig. S2

shows the six chains for the three samples. In addition to the convergence of the two chains for

each sample, our results reveal extremely consistent results between samples. Figure 1A shows

estimates from one chain of the first data set.

Detection of positive selection

For gene families with at least three members, translated sequences were employed to create

maximum likelihood phylogenetic trees using PhyML 3.0 (Guindon et al. 2010) with a

minimum parsimony starting tree and the LG amino acid substitution model with a four-classes

gamma distribution of site-specific substitution rate (Le and Gascuel 2008). The best tree

topology obtained from nearest neighbor interchange (NNI) and subtree pruning recrafting

(SPR) searches was kept (Guindon et al. 2010). BppML (Dutheil and Boussau 2008) was then

used to re-estimate branch lengths from the codon alignment using the YN98 substitution model

(Nielsen and Yang 1998). Next, MapNH (Romiguier et al. 2012) was used for mapping

substitutions on the resulting tree. PartNH (Dutheil et al. 2012) was subsequently employed to

fit time non-homogeneous models of codon substitutions. PartNH uses the previously inferred

substitution maps in order to perform model comparisons and select a non-homogeneous model

with minimal number of parameters. Both methods 'free' (differing parameters between

neighboring branches of a tree) and 'join' (shared parameters between neighboring branches)

were applied to scan for positive selection. Finally, putative secreted effector proteins were

identified by predicting secretion using SignalP 4.0 (Petersen et al. 2011) and proteins were

considered as secreted if the program indicated the presence of a signal peptide but no

transmembrane domain.

To detect residues under positive selection in homologues of *pit2*, the branch-site model with

Bayes Empirical Bayes (BEB) analysis as implemented in PAML4 (Yang 2007) was applied.

We employed information about family composition, alignment and phylogeny as outlined

above and defined *sr10529* and *srs_10529* as foreground branches. A posterior probability

threshold of $> 95\%$ was used for the BEB analysis.

Association of positively selected genes with repeats in U. hordei

Since *U. hordei* shows the highest content of repetitive elements in the group of smut fungi

investigated here, it was tested whether genes under positive selection are located significantly

closer to repetitive elements than average genes in the genome. For this analysis, only a group

of "uncharacterized interspersed repeats" was investigated, because it was shown previously

that this is the only category showing a strong association with candidate effector genes (Dutheil

et al. 2016). Binary logistic regressions were conducted in R using the rms package. The

'robcov' function of the rms package was used in order to get robust estimates of each effect.

The variable "distance to the closest interspersed repeat" was transformed by log(x+1) because

of its extreme distribution.

Comparing d_N/d_S ratios of genes residing in virulence clusters

Previous work has identified several virulence gene clusters in *U. maydis,* some of them playing

important roles during pathogenic development (Kämper et al. 2006; Schirawski et al. 2010).

In total, these clusters contain 163 genes, where 100 reside in clusters without virulence

phenotype and 63 reside in clusters with virulence phenotype upon deletion. Both types of

clusters contain each 32 genes for which a d_N/d_S ratio could be determined (the missing genes

are part of families that do not have at least three members and were therefore not analyzed).

The d_N/d_S ratios of all genes in clusters were compared between clusters with and without

virulence phenotype (Wilcoxon Rank-Sum Test).

Gene Ontology terms enrichment analysis

All proteins in *S. reilianum* f. sp. *zeae* and in *S. reilianum* f. sp. *reilianum* were considered for

Gene Ontology (GO) term enrichment analysis. GO terms were assigned using iprscan 1.1.0

(http://fgblab.org/runiprscan; developed by Michael R. Thon) which links GO information

provided by Interpro to each protein. In this way, 1,759 unique GO terms could be assigned to

4,130 proteins in *S. reilianum* f. sp. *zeae* and 1,744 unique GO terms could be assigned to 4,124

proteins in *S. reilianum* f. sp. *reilianum* (supplementary table S3). The Bioconductor package

topGO (Alexa et al. 2006) was then used in R to link each GO term to the three major categories

"Cellular Component", "Biological Process" or "Molecular Function". Enrichment analysis

was performed by computing *P* values for each GO term using Fisher's classic test with parent-

child correction (Grossmann et al. 2007). Cytoplasmic proteins with and without signatures of

positive selection were compared for the two species separately and differences were

considered to be significant at the 5 % level.

Strains and growth conditions

The *Escherichia coli* derivative Top10 (Invitrogen, Karlsruhe, Germany) and the

Saccharomyces cerevisiae strain BY4741 (*MAT*a *his*3Δ1 *leu*2Δ *met*15Δ *ura*3Δ; Euroscarf,

Frankfurt, Germany; kindly provided by M. Bölker, Marburg) were used for cloning purposes.

Sporisorium reilianum strains used in this study are listed in supplementary table 4. They are

derivatives of the haploid solopathogenic strain JS161 (Schirawski et al. 2010). *E. coli* was

grown in dYT liquid medium (1.6% (w/v) Trypton, 1.0% (w/v) Yeast Extract (Difco), 0.5%

(w/v) NaCl) or YT solid medium (0.8 % (w/v) Trypton, 0.5% (w/v) Yeast-Extract, 0.5% (w/v)

NaCl, 1.3% (w/v) agar) supplemented with 100 mg/mL Ampicillin when needed. *S. cerevisiae*

was maintained in YPD solid medium (1% (w/v) yeast extract, 2% (w/v) Bacto-Pepton, 2%

(w/v) Bacto-Agar, 2% (w/v) glucose) and grown on SC URA$^-$ medium (1.7% (w/v) Yeast

Nitrogen Base without ammonium sulfate, 0.147% (w/v) dropout-mix without Uracil, 2% (w/v)

glucose) for selecting transformants containing the plasmid pRS426 (Sikorski and Hieter 1989)

(kindly provided by M. Bölker, Marburg) or derivatives of pRS426. *S. reilianum* strains were

grown in liquid YEPS$_{light}$ medium (1.0 % (w/v) yeast extract, 0.4% (w/v) peptone, 0.4% (w/v)

sucrose) at 28°C on a rotary shaker at 200 rpm.

Construction of S. reilianum strains

Polymerase chain reactions were performed using the Phusion High-Fidelity DNA Polymerase

(New England Biolabs). Templates were either JS161 genomic DNA or indicated plasmid

DNAs. Restriction enzymes were obtained from New England Biolabs. *S. reilianum* was

transformed by protoplast-mediated transformation following a method established for *U.*

maydis (Schulz et al. 1990). Transformants were selected on RegAgar plates (1.0% (w/v) yeast

extract, 0.4% (w/v) Bacto-Pepton, 0.4% (w/v) Sucrose, 1 M Sorbitol, 1.5% (w/v) Bactoagar)

supplemented with 200 µg/mL Geneticin and true resistance was tested by growing single

colonies on PD plates (3.9% (w/v) Potato-Dextrose Agar, 1% (v/v) Tris-HCl (1M, pH 8.0))

supplemented with 50 µg/mL Geneticin. Gene replacements with resistance markers were

generated with a PCR-based method employing the previously described *Sfi*I insertion cassette

system (Brachmann et al. 2004; Kämper 2004) and were confirmed by Southern blot analysis.

Genomic regions residing about 1 kb upstream (left border) or downstream (right border)

adjacent to open reading frames of candidate genes were PCR-amplified using the listed primer

pairs (supplementary table S5) and genomic DNA of JS161 as template. The resulting

fragments were used for cloning plasmids containing the respective deletion constructs.

To obtain deletion constructs for the genes *sr10529* and *sr14347*, PCR fragments containing

the left and right borders of each gene were ligated to the hygromycin resistance cassette of

pBS-hhn (Kämper 2004) via *Sfi*I restriction sites and cloned into pCRII-TOPO (Life

Technologies) to generate pTOPO Δsr10529 #1 and pTOPO Δsr14347 #1, respectively. Since

the use of Geneticin as selection marker resulted in much less false positive transformants

compared to the use of Hygromycin B, the hygromycin resistance cassettes in these plasmids

were replaced by the Geneticin resistance cassette of pUMA 1057 (Brachmann et al. 2004) by

ligation via *Sfi*I restriction sites, yielding plasmids pTOPO Δsr10529 G418 and pTOPO

Dsr14347 Gen #1, respectively. Deletion constructs were PCR-amplified from plasmids

pTOPO Δsr10529 G418 and pTOPO Dsr14347 Gen #1 using the listed primers (supplementary

table S5) and used to transform the *S. reilianum* strain JS161 to generate the gene deletion

strains JS161Δsr10529 and JS161Δsr14347, respectively.

The drag and drop cloning method in yeast (Jansen et al. 2005) was used to generate plasmids

pRS426 Dsr12968 Hyg #1, pRS426 Dsr14944 Hyg #2, pRS426 Dsr10059 Hyg #1, pRS426

Dsr10182 Hyg #1, pRS426 Dsr14558 Hyg #1 and pRS426 Dsr12897 Hyg #1 which contain

deletion constructs for deleting the candidate genes *sr12968*, *sr14944*, *sr10059*, *sr10182*,

sr14558 or *sr12897*. These plasmids are a derivate of plasmid pRS426, which can be maintained

in *E. coli* and *S. cerevisiae* (Sikorski and Hieter 1989). PCR-amplified left and right borders of

each candidate gene and the hygromycin resistance cassette were integrated in pRS426 by

homologous recombination in *S. cerevisiae*. Subsequently, the hygromycin resistance cassette

was replaced with the Geneticin resistance cassette by ligation via *Sfi*I restriction sites, yielding

plasmids pRS426 Dsr12968 Gen #1, pRS426 Dsr14944 Gen #3, pRS426 Dsr10059 Gen #1,

pRS426 Dsr10182 Gen #1, pRS426 Dsr14558 Gen #1 and pRS426 Dsr12897 Gen #5,

respectively. Gene deletion constructs were PCR-amplified from the respective plasmid using

listed primers (supplementary table S5). The obtained deletion constructs were transformed into

the *S. reilianum* strain JS161 to generate the gene deletion strains JS161Δsr12968,

JS161Δsr14944, JS161Δsr10059, JS161Δsr10182, JS161Δsr14558 and JS161Δsr12897,

respectively.

The drag and drop cloning method was also used to generate plasmid pRS426 Dsr12084 Gen

#1. PCR-amplified left and right borders of *sr12084* and the Geniticin resistance cassette were

integrated in pRS426 by homologous recombination in *S. cerevisiae*. The gene deletion

construct for deleting the candidate gene *sr12084* was PCR-amplified from plasmid pRS426

Dsr12084 Gen #1 using primers sr12084_lb_fw/sr12084_rb_rv and transformed into the *S.*

reilianum strain JS161 to generate the gene deletion strain JS161Δsr12084.

Virulence assays

The solopathogenic strain JS161 and derivatives thereof were grown in YEPS$_{light}$ liquid medium

to an optical density at 600 nm (OD$_{600}$) of 0.8 - 1.0 and cell cultures were adjusted to an OD$_{600}$

of 1.0 with sterile water prior to injection into one week old maize (*Zea mays*) seedlings of the

dwarf cultivar 'Gaspe Flint' (originally kindly provided by B. Burr, Brookhaven National

Laboratories and maintained by self-pollination). Plants were sowed in T-type soil of

'Fruhstorfer Pikiererde' (HAWITA, Vechta, Germany) and grown in a temperature-controlled

greenhouse (14h-/10h- light/dark cycle, with 28/20°C and 25,000 – 90,000 lux during the light

period). Virulence symptoms were scored nine to ten weeks post infection according to

previously described symptoms (Ghareeb et al. 2011) and the following categories were

distinguished: the plant did not develop ears, the plant developed healthy ears shorter or equal

to 1 cm or the plant developed healthy ears longer than 1 cm, the plant developed spiky ears,

phyllody in ears or phyllody in tassels. Spore formation was only observed occasionally and

rarely, the plant died due to the infection. Three independent infections were carried out per

strain, mock treated plants were infected with water as control and at least three independent

deletion strains were tested for virulence.

Results

S. reilianum f. sp. zeae and S. reilianum f. sp. reilianum diverged around one million years ago

To establish a frame for our comparative analysis we first estimated the divergence of the five

smut fungi *U. hordei*, *U. maydis*, *S. scitamineum*, *S. reilianum* f. sp. *zeae* and *S. reilianum* f. sp.

reilianum, both at the non-coding and protein coding levels. We used the information that the

speciation of *U. hordei*, *U. maydis*, *S. scitamineum* and *S. reilianum* likely followed the

divergence pattern of the respective crop ancestors (Munkacsi et al. 2007) and performed a

molecular dating analysis based on the complete proteomes of the five pathogens. We supposed

that the speciation time of the maize and sorghum / sugarcane lineages occurred between 11

and 24 million years (Myr) ago while constraining the speciation of the barley lineage to have

occurred between 24 and 70 Myr ago to calibrate the molecular dating analysis. Based on these

assumptions, we estimated the oldest split between *U. hordei* and the other four species to have

occurred ca 30 Myr ago with a 95% posterior interval ranging from 24 to 47 Myr ago, (fig. 1A

and supplementary table S2). *U. hordei* shares 58 % identity in alignable intergenic regions

with *S. reilianum f. sp. reilianum* and 77 % identity at the protein level (fig. 1B). The split

between *U. maydis* and the *Sporisorium* species was estimated to have occurred around 20 Myr

ago (95% posterior interval 13 to 24 Myr; fig 1A and supplementary table S2). *S. reilianum* f.

sp. *reilianum* shares 61% nucleotide identity in alignable intergenic regions with *U. maydis*,

and 79% sequence identity at the protein level (fig. 1B). *S. scitamineum* and the two formae

speciales of *S. reilianum* were found to have diverged 12 Myr ago (95% posterior interval 7.15

to 16.61 Myr; fig. 1A and supplementary table S2), which is consistent with the mean

divergence estimated from eight studies between the hosts sorghum and sugarcane (10 Myr

with a posterior interval of eight to 13 Myr, source: timetree.org (Kumar et al. 2017)). *S.*

reilianum f. sp. *reilianum* and *S. scitamineum* share 75% non-coding nucleotide identity and

88% identity at the protein level (fig. 1B). Finally, the two *S. reilianum* strains diverged 1.4

Myr ago (95% posterior interval 0.65 to 3.63 Myr; fig. 1A and supplementary table S2) and

share 98% noncoding nucleotide identity and 99% protein identity (fig. 1B). Some studies

suggested that the speciation of sugarcane and sorghum might have been as recent as 5 Myr

ago. In order to test how this assumption impacts the estimation of the divergence between the

two formae speciales of *S. reilianum*, we constrained the split of *S. scitamineum* and *S.*

reilianum to have occurred in the last 5 Myr. This did not significantly impact the split date of

U. hordei (30 Myr ago with a 95% posterior interval of 24 to 55 Myr; supplementary table S2),

but lead to a more recent split time for *U. maydis*, around 12 Myr ago (with a 95% posterior

interval of 11 to 15 Myr ago), and as expected for *S. scitamineum*, around 4.75 Myr ago (with

a 95% posterior interval of 4 to 5 Myr ago, supplementary table S2). Under this assumption,

the split between the two *S. reilianum* species is predicted to be more recent, ca 0.33 Myr ago

(95% posterior interval 0.17 to 0.55 Myr; supplementary table S2), but still much before the

beginning of agriculture. This suggests that the two variants did not diverge by a host jump

between the domesticated species *Sorghum* and maize, but rather between some of the wild

ancestors of these two crop plants. The comparison of the five smut genomes therefore

encompasses a broad evolutionary time. The speciation times involved all predate 10,000 yrs,

which implies that several "domestication" events of these agricultural pathogens occurred

independently. A consequence is (with the putative exception of the two *S. reilianum* formae

speciales), that adaptation to the agricultural host will be negligible when interpreting the inter-

specific patterns of sequence divergence, as it represents a marginal proportion of the time since

the divergence from the ancestral species.

S. reilianum contains the largest number of positively selected genes

To identify homologous genes with signs of positive selection in the five smut fungi analyzed

here, we first built gene families using the SiLiX clustering algorithm (Miele et al. 2011). To

maximize the occurrence of orthologues and minimize the number of paralogues within each

gene family we optimized the clustering parameters for our data set. A total of 8,761 families

was inferred, among which 5,326 families consisted of 1:1 orthologues with up to six members

in each species (supplementary table S6). In this set of families, 245 families contained

predicted secreted proteins only. Given that each species encodes on average 588 secreted

proteins predicted with SignalP (supplementary table S1), about 40 % of all putatively secreted

proteins were present in all five species. Interestingly, we observed several species-specific

family expansions in *U. hordei*. There were 17 families which encompassed five to 25

members, but no members in other species (supplementary table S6). Moreover, we identified

three families with up to 62 members in *U. hordei*, but only one member in up to three of the

other species (supplementary table S6).

To detect positive selection, families with at least three members (supplementary table S6)

were, regardless of their species composition, aligned on the codon and amino acid level and a

phylogenetic tree was inferred. Obtaining accurate alignments is critical for detecting positive

selection since alignment errors frequently inflate the false discovery rate (Jordan and Goldman

2012; Schneider et al. 2009). We therefore developed a stringent bioinformatics pipeline for the

filtering of sequence alignments by masking ambiguous alignment positions for further analysis

(see Methods). A non-homogeneous model of sequence evolution allowing d_N/d_S ratios to vary

along the phylogeny was employed to scan for positive selection (Dutheil et al. 2012; Nielsen

and Yang 1998). The largest number of genes with signs of positive selection was found in *S.*

reilianum f. sp. *zeae* (84 genes, of which 25 encode predicted secreted proteins) and *S.*

reilianum f. sp. *reilianum* (111 genes of which 27 encode predicted secreted proteins) (fig. 1C).

In addition, a substantial number of positively selected candidate genes was also found in *U.*

hordei (49, and of these 22 genes are predicted to code for secreted proteins), but only very few

in *U. maydis* (2 genes) and *S. scitamineum* (7 genes) (fig. 1C). A list of all proteins with their

associated d_N/d_S ratios in each species is provided in supplementary table S3. Predicted secreted

proteins were significantly enriched in the group of proteins under positive selection in *U.*

hordei and in the two investigated pathovariants of *S. reilianum* (*P* values < 10^{-5}; Fisher's exact

test). This corroborates results of earlier studies that showed that predicted secreted proteins are

often under positive selection, which can be attributed to their direct interaction with host

proteins. Notably, all genes found under positive selection in the two strains of *S. reilianum*, in

S. scitamineum and in *U. maydis* share orthologous genes in the other species (supplementary

tables S2 and S5). In contrast, genes with signs of positive selection in *U. hordei* belong largely

(36 out of 49 genes) to families showing species-specific expansions (supplementary tables S3

and S6).

Genes under positive selection in U. hordei are associated with uncharacterized interspersed

repeats

Among the species compared here, the genome of *U. hordei* shows the highest fraction of repetitive elements (Dutheil et al. 2016). Such elements are known to contribute to gene family expansions (Kazazian 2004), and have been suggested to contribute to adaptation by providing advantageous mutations, for instance by RIP leakage (Dong et al. 2015; Rouxel et al. 2011). Therefore, we tested whether genes under positive selection in *U. hordei* are physically associated with repetitive elements. We performed a binary logistic regression with the prediction of positive selection as a response variable (that is, whether the underlying branch has a dN/dS ratio higher than one) and we considered three putative explanatory variables: (1) whether the gene was predicted to encode a secreted protein, (2) whether the gene is duplicated and (3) the distance of the gene to the closest interspersed repeat. The complete linear model explains of 50% of the observed variance, and the three explanatory variables are all significant at the 0.1% level (supplementary table S7). These results suggest that positively selected genes in *U. hordei* are associated with duplication events, and positive selection is more likely to occur at genes encoding putative effectors. In addition, the proximity of interspersed repeats increases the odds of positive selection, independently of the two other effects, and is confirmed by a stratification approach: the effect still holds when only duplicated genes are considered, or only genes encoding a secreted protein, or the combination of the two (supplementary table S7). This finding corroborates previous results obtained in other microbial plant pathogens where it was described that effector genes tend to localize in repeat rich regions and where it was suggested that such regions contribute to the rapid evolution of effector genes (Dong et al. 2015).

Positively selected genes encoding cytoplasmic proteins in S. reilianum

While we expect effector genes to be under positive selection, we find that the majority of

positively selected genes in *S. reilianum* encodes cytoplasmic proteins (fig. 1C). To assess the

putative functional role of these genes, we performed a Gene Ontology term enrichment

analysis, comparing cytoplasmic proteins under positive selection to cytoplasmic proteins not

under positive selection (table 1). This analysis revealed that genes with a potential role in

metabolic processes, like sulfur compound metabolism, molybdopterin cofactor metabolic

process, RNA metabolic process, organic cyclic compound metabolic process and

oxidoreductase activity, as well as responses to starvation and extracellular stimuli are

significantly overrepresented (table 1). This could indicate that cytoplasmic proteins under

positive selection contribute to metabolic changes which might be needed to survive with the

limited nutrients available on the surface or in the biotrophic interface of different host plants.

Virulence contribution of effector genes showing signs of positive selection in S. reilianum

Candidate effector genes inferred to be under positive selection in a particular species could

play a critical role in pathogenicity. Therefore, we sought to assess the contribution to virulence

of such candidate genes by creating individual deletion mutants. In total nine candidate genes

with high d_N/d_S ratios were chosen which were all predicted to encode secreted proteins: three

with signatures of positive selection only in *S. reilianum* f. sp. *zeae*, three with signatures of

positive selection in *S. reilianum* f. sp. *zeae* as well as in *S. reilianum* f. sp. *reilianum* and three

with signatures of positive selection only in *S. reilianum* f. sp. *reilianum*. All nine chosen

candidate genes together with their selection criteria are summarized in table 2. Deletion

mutants were generated in the haploid solopathogenic strain JS161 of *S. reilianum* f. sp. *zeae*.

This strain is capable to colonize maize plants and cause disease without a compatible mating

partner (Schirawski et al. 2010). Deletion mutants were also generated in strain JS161 in cases

where positive selection was only detected in *S. reilianum* f. sp. *reilianum* (table 2), because no

solopathogenic strain is presently available for *S. reilianum* f. sp. *reilianum*. For each gene at

least three independent deletion mutants were generated and tested for virulence. To determine

virulence, Gaspe Flint, a dwarf variety of corn, was infected and symptoms were scored in male

and female flowers (fig. 2). Only the deletion of *sr10529*, a gene showing positive selection in

both formae speciales of *S. reilianum*, showed a strong reduction in virulence (table 2 and fig.

2). The gene *sr10529* in *S. reilianum* f. sp. *zeae* is orthologous to the previously identified and

characterized gene *pit2* (*UMAG_01375*) in *U. maydis*.

Pit2 plays an essential role in virulence as inhibitor of a group of maize papain-like cysteine

proteases that are secreted to the apoplast (Doehlemann et al. 2011; Mueller et al. 2013).

Previous work identified a conserved domain of 14 amino acids (PID14) in Pit2 as required and

sufficient for the inhibition of maize cysteine proteases (Mueller et al. 2013). When the branch-

site model of PAML 4 (Yang 2007) was used to identify amino acid residues under positive

selection in the Pit2 orthologues of the two *S. reilianum* species, only two residues residing in

the PID14 domain were found under positive selection. However, 24 positively selected

residues were detected outside this domain in the 57 amino acids long C-terminus (fig. 3).

Discussion

We used evolutionary comparative genomics of five related smut fungi infecting four different

host plants to identify positively selected genes encoding predicted secreted proteins, as such

genes were suggested to contribute to virulence in various plant pathogenic microbes (Aguileta

et al. 2010; Dong et al. 2014; Hacquard et al. 2012; Huang et al. 2014; Sharma et al. 2014;

Stukenbrock et al. 2011). Our analysis revealed that positive selection is found between

paralogous genes in *U. hordei*, where they belong to families with species-specific expansions.

In contrast, genes under positive selection in the other four species belong to families of

orthologous sequences. While we find evidence for a large set of genes under positive selection

in the *S. reilianum* species, signatures for positive selection are hardly detectable in the more

distant relatives *U. hordei*, *U. maydis* and *S. scitamineum* that diverged earlier (fig. 1A). Finding

evidence for positive selection over time spans of several millions of years is notoriously

difficult (Gillespie 1994) because of two main reasons: (1) periods where genes are evolving

under positive selection occur episodically and may be followed by long episodes of purifying

selection, leading to an average d_N/d_S below 1 on long periods of time and (2) fast evolving

genes may diverge to an extent where their homology is difficult to infer and where they can

no longer be aligned reliably. To overcome this problem, more genome information of species

with intermediate branching points is needed (Gillespie 1994).

Predicted secreted proteins were about three times overrepresented in the set of positively

selected genes and this illustrates the importance of secreted proteins in adaptation processes

of smut fungi. This corroborates results in other plant pathogenic microbes like *Melampsora*

sp., *Z. tritici* and the wheat powdery mildew *Blumeria graminis* (Joly et al. 2010; Poppe et al.

2015; Wicker et al. 2013). However, the majority of positively selected genes encodes

cytoplasmic proteins (fig. 1C), suggesting that important targets of adaptation like plant

colonization are not only governed by secreted proteins. A Gene Ontology analysis in *S.*

reilianum showed that mainly processes related to metabolism and its regulation as well as

responses to starvation and external stimuli are enriched in cytoplasmic proteins under positive

selection (table 1). This points at a role of these proteins in adaptation to differences in nutrient

availability in the respective host plants maize and sorghum as well as responses to cues

originating from the respective host (Haueisen and Stukenbrock 2016). A study conducted in

U. maydis has shown that the fungus induces major metabolic changes in the host plant upon

infection during establishment of biotrophy and undergoes a series of developmental transitions

during host colonization which are likely influenced by the host environment (Doehlemann et al. 2008). It is thus conceivable that the two *S. reilianum* accessions have adapted to their different hosts that differ significantly for example in their amino acid and vitamin composition (Etuk et al. 2012).

Out of nine deletions of positively selected genes, only one mutant, lacking *sr10529,* was affected in virulence. While six genes deleted are single genes in *S. reilianum* f. sp. *zeae* for which we failed to identify paralogs, *sr12084* has two paralogs, *sr14347* has five paralogs and *sr10182* has ten paralogs (table 2). In our analysis we have restricted ourselves to generating deletion mutants in the genes under positive selection only. This leaves open the possibility that the paralogous genes have redundant functions in virulence. Adapting the CRISPR-Cas9 technology allowing multiplexing (Schuster et al. 2017) to *S. reilianum* will help to address this in future studies. Alternatively, the candidate effectors we investigated may be needed under conditions which differ from those tested here. For example, *S. reilianum* f. sp. *zeae* can systemically colonize maize plants via root infection (Mazaheri-Naeini et al. 2015), a colonization route we have not assessed in our experimental setup. Knowing the expression profile of effector genes might help to look for characteristic differences in development of the mutants compared to wild type strains. Since we lack this information, we scored disease symptoms only in the inflorescences about nine weeks after infection. In three cases positive selection was detected in orthologous genes in *S. reilianum* f. sp. *reilianum* (table 2) while candidate effector genes were for experimental reasons deleted in *S. reilianum* f. sp. *zeae*. Therefore, it cannot be excluded that these effectors might have a virulence function in *S. reilianum* f. sp. *reilianum.* In this case, the positively selected effector genes might have evolved during adaptation to the sorghum host and present host specificity genes. Candidate effector proteins might also be positively selected for traits which are not directly linked to pathogenicity. Such traits could for instance involve competition with large numbers of other

plant colonizing microbes (Rovenich et al. 2014; Zhan and McDonald 2013). Secreted proteins

of *S. reilianum* could act for example as toxin or could efficiently utilize resources from the

environment and thereby limit the growth of other microbes. In these cases, a contribution to

virulence is not expected to be observed in effector gene mutants. Moreover, our molecular

dating analysis showed that the common ancestors of the investigated smut species originated

before the beginning of crop domestication. Therefore, positive selection, whose signs we

detect by our approach, has most likely occurred on ancestral host plants and not on the

domesticated host maize. Consequently, some of the candidate effector genes under positive

selection might not be important for the colonization of crop plants, but for infection of related

wild species.

In *U. maydis*, we note that effector genes residing in clusters whose deletion affected virulence

(Kämper et al. 2006; Schirawski et al. 2010) have similar d_N/d_S ratios as effector genes in

clusters where the deletion had no effect on virulence (median d_N/d_S ratio 0.0619 vs.0.1094;

Wilcoxon rank test with *P* value = 0.1848). Furthermore, orthologues of the effectors Pep1,

Stp1 and Cmu1, which were shown to have important roles in pathogenicity of *U. maydis*

(Djamei et al. 2011; Doehlemann et al. 2009; Schipper 2009) showed no signatures of positive

selection. These observations could suggest that certain fungal effector proteins are for

unknown reasons under evolutionary constraint and are therefore not free to accumulate non-

synonymous mutations. Such effectors are conserved over long time spans (Hemetsberger et al.

2015; Schirawski et al. 2010; Sharma et al. 2015) and this illustrates that they are instrumental

for successful infections in a large group of smut fungi. They probably target molecules shared

by several host plants, for example housekeeping functions that cannot easily evolve in response

to the binding of an effector.

One candidate gene (*sr10529*) under positive selection in both formae speciales of *S. reilianum*

showed a strong contribution to virulence upon deletion. It is orthologous to the previously

described protease inhibitor Pit2 in *U. maydis*, where the deletion also abolished virulence

(Doehlemann et al. 2011; Mueller et al. 2013). Positively selected residues in the PID14 domain

of Pit2 might reflect that different proteases need to be inhibited in maize and sorghum. Pit2

might thus contribute to determining the host range of the respective species. A role of cysteine

protease inhibitors in host specificity was demonstrated in the pathosystem *Phytophthora*

infestans, a pathogen of potato and its sister species *Phytophthora mirabilis*, which infects the

ornamental plant *Mirabilis jalapa*. Positively selected orthologous protease inhibitors where

shown to inhibit proteases specific to the respective host plants and this specificity could be

traced back to a single amino acid substitution (Dong et al. 2014). Surprisingly, 24 positively

selected sites in Pit2 were detected outside the PID14 domain in the 57 amino acid long C-

terminus in both *S. reilianum* f. sp. *zeae* and f. sp. *reilianum*. This finding raises the intriguing

possibility that the C-terminus of Pit2 might possess a second function, which is independent

of protease inhibition. Earlier work has shown that the *pit1* gene encoding a transmembrane

protein is located next to the *pit2* effector gene and both genes contribute similarly to virulence

(Doehlemann et al. 2011). Furthermore, *pit1* and *pit2* are divergently transcribed which makes

it likely that the expression of *pit1* and *pit2* is co-regulated. In addition, this gene arrangement

of *pit1* and *pit2* is conserved in *U. hordei*, *U. maydis*, *S. scitamineum* and *S. reilianum* (Sharma

et al. 2015). This finding has led to the speculation that Pit1 and Pit2 somehow act together to

govern virulence of *U. maydis* and related smut fungi. It was hypothesized that that Pit2 shuttles

apoplastic maize proteins towards Pit1, thereby scavenging damage-associated molecules

(Doehlemann et al. 2011). In this scenario, the positively selected amino acids in the C-terminus

of Pit2 could have been selected for scavenging such molecules as adaptation to the two hosts.

In future studies it will be highly interesting to complement the *pit2* mutant of *S. reilianum* f.

sp. *zeae* with the *pit2* orthologue of *S. reilianum* f. sp. *reilianum* to see if this promotes virulence

on sorghum.

Conclusions

Screens for genes with signs of positive selection are commonly used to identify candidate
effector genes in various plant pathogenic microbes. However, it is currently largely open
whether positively selected effector genes play indeed a role in virulence. Here, we used
comparative genomics of five smut fungi and showed that only one out of nine genes under
positive selection contribute to virulence of *S. reilianum*. Moreover, the majority of positively
selected genes did not encode predicted secreted proteins. Our results therefore suggest that
many genes with signatures of positive selection are involved in traits that are not directly
related to pathogenicity and that positive selection alone is not a good indicator for a virulence
function of a certain gene.

Acknowledgements

Our work was supported by funds from the Max Planck Society. We are grateful to Markus
Rampp (MPI for Plasma Physics) for valuable support in using the computer cluster of the Max
Planck Computing Center Garching. We thank Stefan Schmidt for expert greenhouse service
and all present and former group members for lively discussions. We thank Michael Bölker for
providing the yeast strain BY4741 and the plasmid pRS426 which was used for cloning
purposes. We thank Benoit Nabholz and Nicolas Lartillot for guidance with the dating analysis.

References

Aguileta G, et al. 2012. Genes under positive selection in a model plant pathogenic fungus,
Botrytis. Infect Genet Evol 12: 987-996.

Aguileta G, et al. 2010. Finding candidate genes under positive selection in non-model species:
examples of genes involved in host specialization in pathogens. Mol Ecol 19: 292-306.

Aguileta G, Refregier G, Yockteng R, Fournier E, Giraud T. 2009. Rapidly evolving genes in
pathogens: methods for detecting positive selection and examples among fungi, bacteria,
viruses and protists. Infect Genet Evol 9: 656-670.

Alexa A, Rahnenführer J, Lengauer T. 2006. Improved scoring of functional groups from gene
expression data by decorrelating GO graph structure. Bioinformatics 22: 1600-1607.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search
tool. J Mol Biol 215: 403-410.

Bagchi R, et al. 2014. Pathogens and insect herbivores drive rainforest plant diversity and
composition. Nature 506: 85-88.

Blanchette M, et al. 2004. Aligning multiple genomic sequences with the Threaded Blockset
Aligner. Genome Res 14: 708-715.

Brachmann A, König J, Julius C, Feldbrügge M. 2004. A reverse genetic approach for
generating gene replacement mutants in *Ustilago maydis*. Mol Genet Genomics 272: 216-226.

Brown JKM, Tellier A. 2011. Plant-parasite coevolution: bridging the gap between genetics
and ecology. Annu Rev Phytopathol 49: 345-367.

Cantu D, et al. 2013. Genome analyses of the wheat yellow (stripe) rust pathogen *Puccinia*
striiformis f. sp. *tritici* reveal polymorphic and haustorial expressed secreted proteins as
candidate effectors. BMC Genomics 14: 270.

Charif D, Lobry JR. 2007. SeqinR 1.0-2: a contributed package to the R project for statistical
computing devoted to biological sequences retrieval and analysis. In: Bastolla U, Porto M,
Roman HE, Vendruscolo M, editors. Structural approaches to sequence evolution: Molecules,
networks, populations. New York: Springer Verlag. p. 207-232.

Dean R, et al. 2012. The top 10 fungal pathogens in molecular plant pathology. Mol Plant Pathol
13: 414-430.

Djamei A, et al. 2011. Metabolic priming by a secreted fungal effector. Nature 478: 395-398.

Doehlemann G, Reissmann S, Aßmann D, Fleckenstein M, Kahmann R. 2011. Two linked
genes encoding a secreted effector and a membrane protein are essential for *Ustilago maydis*-
induced tumour formation. Mol Microbiol 81: 751-766.

Doehlemann G, et al. 2009. Pep1, a secreted effector protein of *Ustilago maydis*, is required for
successful invasion of plant cells. PLoS Pathog 5: e1000290.

Doehlemann G, et al. 2008. Reprogramming a maize plant: transcriptional and metabolic
changes induced by the fungal biotroph *Ustilago maydis*. Plant J 56: 181-195.

Dong S, Raffaele S, Kamoun S. 2015. The two-speed genomes of filamentous pathogens: waltz
with plants. Curr Opin Genet Dev 35: 57–65.

Dong S, et al. 2014. Effector specialization in a lineage of the Irish potato famine pathogen.
Science 343: 552-555.

Dutheil JY, Boussau B. 2008. Non-homogeneous models of sequence evolution in the Bio++
suite of libraries and programs. BMC Evol Biol 8: 255.

Dutheil JY, et al. 2012. Efficient selection of branch-specific models of sequence evolution.
Mol Biol Evol 29: 1861-1874.

Dutheil JY, et al. 2016. A tale of genome compartmentalization: the evolution of virulence
clusters in smut fungi. Genome Biol Evol 8: 681-704.

Etuk EB, et al. 2012. Nutrient composition and feeding value of *Sorghum* for livestock and
poultry: a review. J Anim Sci Adv 2: 510-524.

Fisher MC, et al. 2012. Emerging fungal threats to animal, plant and ecosystem health. Nature
484: 186-194.

Franceschetti M, et al. 2017. Effectors of filamentous plant pathogens: commonalities amid
diversity. Microbiol Mol Biol R 81: e00066-00016.

Gehrig H, Schussler A, Kluge M. 1996. *Geosiphon pyriforme*, a fungus forming endocytobiosis
with *Nostoc* (cyanobacteria), is an ancestral member of the *Glomales*: evidence by SSU rRNA
analysis. J Mol Evol 43: 71-81.

Ghareeb HA, Becker A, Iven T, Feussner I, Schirawski J. 2011. *Sporisorium reilianum*
infection changes inflorescence and branching architectures of maize. Plant Physiol 156: 2037-
2052.

Gillespie JH. 1994. The causes of molecular evolution. Oxford: Oxford University Press.

Grossmann S, Bauer S, Robinson PN, Vingron M. 2007. Improved detection of
overrepresentation of Gene-Ontology annotations with parent child analysis. Bioinformatics
23: 3024-3031.

Guindon S, et al. 2010. New algorithms and methods to estimate maximum-likelihood
phylogenies: assessing the performance of PhyML 3.0. Syst Biol 59: 307-321.

Guyon K, Balagué C, Roby D, Raffaele S. 2014. Secretome analysis reveals effector candidates
associated with broad host range necrotrophy in the fungal plant pathogen *Sclerotinia*
sclerotiorum. BMC Genomics 15: 336.

Hacquard S, et al. 2012. A comprehensive analysis of genes encoding small secreted proteins
identifies candidate effectors in *Melampsora larici-populina* (poplar leaf rust). Mol Plant
Microbe Interact 25: 279-293.

Haueisen J, Stukenbrock EH. 2016. Life cycle specialization of filamentous pathogens -
colonization and reproduction in plant tissues. Curr Opin Microbiol 32: 31-37.

Hemetsberger C, et al. 2015. The fungal core effector Pep1 is conserved across smuts of dicots
and monocots. New Phytol 206: 1116-1126.

Huang J, Si W, Deng Q, Li P, Yang S. 2014. Rapid evolution of avirulence genes in rice blast
fungus *Magnaporthe oryzae*. BMC Genetics 15: 45.

Jansen G, Wu C, Schade B, Thomas DY, Whiteway M. 2005. Drag&Drop cloning in yeast.
Gene 344: 43-51.

Joly DL, Feau N, Tanguay P, Hamelin RC. 2010. Comparative analysis of secreted protein
evolution using expressed sequence tags from four poplar leaf rusts (*Melampsora spp.*). BMC
Genomics 11: 422.

Jordan G, Goldman N. 2012. The effects of alignment error and alignment filtering on the
sitewise detection of positive selection. Mol Biol Evol 29: 1125-1139.

Kämper J. 2004. A PCR-based system for highly efficient generation of gene replacement
mutants in *Ustilago maydis*. Mol Genet Genomics 271: 103-110.

Kämper J, et al. 2006. Insights from the genome of the biotrophic fungal plant pathogen
Ustilago maydis. Nature 444: 97-101.

Kazazian HH. 2004. Mobile elements: drivers of genome evolution. Science 303: 1626-1632.

Khrunyk Y, Münch K, Schipper K, Lupas AN, Kahmann R. 2010. The use of FLP-mediated
recombination for the functional analysis of an effector gene family in the biotrophic smut
fungus *Ustilago maydis*. New Phytol 187: 957-968.

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: a resource for timelines,
timetrees, and divergence times. Mol Biol Evol 34: 1812-1819.

Lanver D, et al. 2017. *Ustilago maydis* effectors and their impact on virulence. Nat Rev
Microbiol 15: 409-421.

Lartillot N, Lepage T, Blanquart S. 2009. PhyloBayes 3: a Bayesian software package for
phylogenetic reconstruction and molecular dating. Bioinformatics 25: 2286-2288.

Laurie JD, et al. 2012. Genome comparison of barley and maize smut fungi reveals targeted
loss of RNA silencing components and species-specific presence of transposable elements.
Plant Cell 24: 1733-1745.

Le SQ, Gascuel O. 2008. An improved general amino acid replacement matrix. Mol Biol Evol
25: 1307-1320.

Lo Presti L, et al. 2015. Fungal effectors and plant susceptibility. Annu Rev Plant Biol 66: 513-
545.

Löytynoja A, Goldman N. 2008. Phylogeny-aware gap placement prevents errors in sequence
alignment and evolutionary analysis. Science 320: 1632-1635.

Martin FM, Uroz S, Barker DG. 2017. Ancestral alliances: Plant mutualistic symbioses with
fungi and bacteria. Science 356: eaad4501.

Martinez-Espinoza AD, Garcia-Pedrajas MD, Gold SE. 2002. The Ustilaginales as plant pests
and model systems. Fungal Genet Biol 35: 1-20.

Mazaheri-Naeini M, Sabbagh SK, Martinez Y, Séjalon-Delmas N, Roux C. 2015. Assessment
of *Ustilago maydis* as a fungal model for root infection studies. Fungal Biol 119: 145–153.

Miele V, Penel S, Duret L. 2011. Ultra-fast sequence clustering from similarity networks with
SiLiX. BMC Bioinformatics 12: 116.

Mueller AN, Ziemann S, Treitschke S, Aßmann D, Doehlemann G. 2013. Compatibility in the
Ustilago maydis–maize interaction requires inhibition of host cysteine proteases by the fungal
effector Pit2. PLoS Pathog 9: e1003177.

Munkacsi AB, Stoxen S, May G. 2007. Domestication of maize, sorghum, and sugarcane did
not drive the divergence of their smut pathogens. Evolution 61: 388–403

Nielsen R. 2005. Molecular signatures of natural selection. Annu Rev Genet 39: 197-218.

Nielsen R, Yang Z. 1998. Likelihood models for detecting positively selected amino acid sites
and applications to the HIV-1 envelope gene. Genetics 148: 929-936.

Parniske M. 2008. Arbuscular mycorrhiza: the mother of plant root endosymbioses. Nat Rev
Microbiol 6: 763-775.

Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal
peptides from transmembrane regions. Nat Methods 8: 785-786.

Plissonneau C, et al. 2017. Using population and comparative genomics to understand the
genetic basis of effector-driven fungal pathogen evolution. Front Plant Sci 8: 119.

Poppe S, Dorsheimer L, Happel P, Stukenbrock EH. 2015. Rapidly evolving genes are key
players in host specialization and virulence of the fungal wheat pathogen *Zymoseptoria tritici*
(*Mycosphaerella graminicola*). PLoS Pathog 11: e1005055.

Que Y, et al. 2014. Genome sequencing of *Sporisorium scitamineum* provides insights into the
pathogenic mechanisms of sugarcane smut. BMC Genomics 15: 996.

Ranwez V, Harispe S, Delsuc F, Douzery EJP. 2011. MACSE: Multiple Alignment of Coding
SEquences accounting for frameshifts and stop codons. PLOS One 6: e22594.

Remy W, Taylor TN, Hass H, Kerp H. 1994. Four hundred-million-year-old vesicular
arbuscular mycorrhizae. PNAS 91: 11841-11843.

Romiguier J, et al. 2012. Fast and robust characterization of time-heterogeneous sequence
evolutionary processes using substitution mapping. PLOS One 7: e33852.

Rouxel T, et al. 2011. Effector diversification within compartments of the *Leptosphaeria*
maculans genome affected by Repeat-Induced Point mutations. Nat Commun 2: 202.

Rovenich H, Boshoven JC, Thomma BPHJ. 2014. Filamentous pathogen effector functions: of
pathogens, hosts and microbiomes. Curr Opin Plant Biol 20: 96–103.

Schipper K 2009. Charakterisierung eines *Ustilago maydis* Genclusters, das für drei neuartige
sekretierte Effektoren kodiert. Philipps-Universität Marburg.

Schirawski J, et al. 2010. Pathogenicity determinants in smut fungi revealed by genome
comparison. Science 330: 1546-1548.

Schneider A, et al. 2009. Estimates of positive Darwinian selection are inflated by errors in
sequencing, annotation, and alignment. Genome Biol Evol 1: 114-118.

Schulz B, et al. 1990. The *b* alleles of *U. maydis*, whose combinations program pathogenic
development, code for polypeptides containing a homeodomain-related motif. Cell 60: 295-
306.

Schuster M, Schweizer G, Kahmann R. 2017. Comparative analyses of secreted proteins in
plant pathogenic smut fungi and related basidiomycetes. Fungal Genet Biol in press.

Schuster M, Schweizer G, Reissmann S, Kahmann R. 2016. Genome editing in *Ustilago maydis*
using the CRISPR–Cas system. Fungal Genet Biol 89: 3-9.

Sharma R, Mishra B, Runge F, Thines M. 2014. Gene loss rather than gene gain is associated
with a host jump from monocots to dicots in the smut fungus *Melanopsichium pennsylvanicum*.
Genome Biol Evol 6: 2034-2049.

Sharma R, Xia X, Riess K, Bauer R, Thines M. 2015. Comparative genomics including the
early-diverging smut fungus *Ceraceosorus bombacis* reveals signatures of parallel evolution
within plant and animal pathogens of fungi and oomycetes. Genome Biol Evol 7: 2781-2798.

Sikorski RS, Hieter P. 1989. A system of shuttle vectors and yeast host strains designed for
efficient manipulation of DNA in *Saccharomyces cerevisiae*. Genetics 122: 19-27.

Silva DN, et al. 2015. Genomic patterns of positive selection at the origin of rust fungi. PLOS
One 10: e0143959.

Sperschneider J, et al. 2015. Genome-wide analysis in three *Fusarium* pathogens identifies
rapidly evolving chromosomes and genes associated with pathogenicity. Genome Biol Evol 7:
1613-1627.

Sperschneider J, et al. 2014. Diversifying selection in the wheat stem rust fungus acts
predominantly on pathogen-associated gene families and reveals candidate effectors. Front
Plant Sci 5: 372.

Stukenbrock EH, et al. 2011. The making of a new pathogen: insights from comparative
population genomics of the domesticated wheat pathogen *Mycosphaerella graminicola* and its
wild sister species Genome Res 21: 2157-2166.

Taniguti LM, et al. 2015. Complete genome sequence of *Sporisorium scitamineum* and
biotrophic interaction transcriptome with sugarcane. PLOS One 10: e0129318.

Tellier A, Moreno-Gámez S, Stephan W. 2014. Speed of adaptation and genomic footprints of
host-parasite coevolution under arms race and trench warefare dynamics. Evolution 68: 2211-
2224.

Thorne JL, Kishino H, Painter IS. 1998. Estimating the rate of evolution of the rate of molecular
evolution. Mol Biol Evol 15: 1647-1657.

Toruño TY, Stergiopoulos I, Coaker G. 2016. Plant-pathogen effectors: cellular probes
interfering with plant defenses in spatial and temporal manners. Annu Rev Phytopathol 54: 419-
441.

Vánky K. 2012. Smut Fungi of the World. St Paul, MN, USA: APS Press.

Wicker T, et al. 2013. The wheat powdery mildew genome shows the unique evolution of an
obligate biotroph. Nat Genet 45: 1092-1096.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol 24:
1586-1591.

Zhan J, McDonald BA. 2013. Experimental measures of pathogen competition and relative
fitness. Annu Rev Phytopathol 51: 131-153.

Zuther K, et al. 2012. Host specificity of *Sporisorium reilianum* is tightly linked to generation
of the phytoalexin luteolinidin by *Sorghum bicolor*. Mol Plant Microbe Interact 25: 1230-1237.

Figure legends

FIG. 1. Phylogeny, divergence estimates and number of genes under positive selection in five

related smut fungi species parasitizing different host plants. A) Chronogram of the five fungal

pathogens as estimated under a relaxed molecular clock. Boxes represent 95% posterior

intervals, with corresponding values indicated below. B) Pairwise sequence differences, for

both the non-coding genome and the proteome (non-synonymous differences). C) Number of

positively selected genes on each terminal branch (total number of genes and genes predicted

to encode a secreted protein).

FIG. 2. Virulence phenotypes of single deletion mutants of positively selected genes in *S.*

reilianum f. sp. *zeae*. Plants of the maize variety 'Gaspe Flint' were infected with water, the

solopathogenic strain JS161 or independent deletion mutants of candidate genes as indicated

below each bar. Deletion of *sr10529* lead to a strong reduction in virulence (left panel in first

row). In contrast, deletion of the candidate genes *sr12968* (left panel in second row), *sr14558*

(right panel in second row), *sr14944* (left panel in third row), *sr14347* (right panel in third row),

sr10059 (left panel in forth row), *sr12897* (right panel in forth row), *sr10182* (left panel in fifth

row) and *sr12084* (right panel in fifth row) did not alter virulence. Symptoms were scored about

9 weeks post infection and categorized according to severeness as illustrated in the legend

below the bar plots. Results are shown as mean of three independent experiments in relation to

the total number of infected plants, which is indicated above each bar (n). Note that strains

JS161ΔSr10529 #G4 and #G5 (left panel in first row) were only infected in one replicate.

FIG. 3. Distribution of positively selected amino acids in the cysteine protease inhibitor Pit2.

The alignment shows the protein sequences of orthologues in *U. hordei* (UHOR_02064), *U.*

maydis (UMAG_01375), *S. scitamineum* (SPSC_03677), *S. reilianum* f. sp. *zeae* (sr10529) and

S. reilianum f. sp. *reilianum* (srs_10529). Sites under positive selection detected by a branch-

site model are indicated by colored bold letters. Residues colored in red indicate positive

selection detected in the respective species and purple residues indicate sites found under

positive selection in both species. The yellow shaded area is orthologous to the previously

identified conserved PID14 domain, which is required and sufficient for inhibition of a group

of papain-like cysteine proteases. Green sequences indicate secretion signal peptides and bold

numbers above the alignment indicate positions in UHOR_02064.

Supplementary figure 1. Number of families consisting of 1:1 orthologues in relation to

varying settings for coverage and identity in the clustering program SiLiX. The maximum

number of families containing 1:1 orthologues can be obtained with a coverage between 5 %

and 45 % and an identity of 40 %.

Supplementary figure 2. Trace of the Monte-Carlo Markov chains for 3 gene samples (see

Methods). Vertical lines show the burning phase (10,000 iterations). A) With 2 calibration

points B) With 3 calibration points.

List of tables

Table 1: Gene Ontology terms significantly overrepresented in positively selected genes encoding cytoplasmic proteins in *S. reilianum*

Table 2: Genes under positive selection that were individually deleted in *S. reilianum* f. sp. *zeae* and the resulting virulence phenotype

Supplementary table S1: Sources of genomic data for the five smut fungi species compared

Supplementary table S2: Results of the molecular dating analysis

Supplementary table S3: Omega values of all genes, predictions of secretion and Gene Ontology terms in *S. reilianum*

Supplementary table S4: List of *S. reilianum* strains used in this study

Supplementary table S5: List of all primers

Supplementary table S6: Composition of 8,761 families containing proteins of five smut fungi

Supplementary table S7: Results of the linear model describing the association of gene duplication, positive selection and candidate effector genes in *U. hordei*

Table 1: Gene Ontology terms significantly overrepresented in positively selected genes encoding cytoplasmic proteins in *S. reilianum*

Gene Ontology Id	Gene Ontology Description	Category	*P* values[a]	Species with significantly enriched category
GO:0030532	small nuclear ribonucleoprotein complex	Cellular Component	0.023	*S. reilianum* f. sp. *reilianum*
GO:0045263	proton-transporting ATP synthase complex, coupling factor F(o)	Cellular Component	0.038 and 0.036	*S. reilianum* f. sp. *reilianum* and f. sp. *zeae*
GO:0044425	membrane part	Cellular Component	0.047	*S. reilianum* f. sp. *reilianum*
GO:0031668	cellular response to extracellular stimulus	Biological Process	0.015 and 0.013	*S. reilianum* f. sp. *reilianum* and f. sp. *zeae*
GO:0051186	cofactor metabolic process	Biological Process	0.015	*S. reilianum* f. sp. *reilianum*
GO:0042594	response to starvation	Biological Process	0.017 and 0.012	*S. reilianum* f. sp. *reilianum* and f. sp. *zeae*
GO:0009267	cellular response to starvation	Biological Process	0.020 and 0.014	*S. reilianum* f. sp. *reilianum* and f. sp. *zeae*
GO:0006790	sulfur compound metabolic process	Biological Process	0.022	*S. reilianum* f. sp. *reilianum*
GO:0051301	cell division	Biological Process	0.023	*S. reilianum* f. sp. *reilianum*
GO:0006777	Mo-molybdopterin cofactor biosynthetic process	Biological Process	0.027	*S. reilianum* f. sp. *reilianum*
GO:0009605	response to external stimulus	Biological Process	0.027 and 0.012	*S. reilianum* f. sp. *reilianum* and f. sp. *zeae*
GO:0043545	molybdopterin cofactor metabolic process	Biological Process	0.034	*S. reilianum* f. sp. *reilianum*
GO:0043413	macromolecule glycosylation	Biological Process	0.036	*S. reilianum* f. sp. *reilianum*
GO:0022402	cell cycle process	Biological Process	0.037	*S. reilianum* f. sp. *reilianum*
GO:0051189	prosthetic group metabolic process	Biological Process	0.039	*S. reilianum* f. sp. *reilianum*
GO:0006139	nucleobase-containing compound metabolic process	Biological Process	0.012	*S. reilianum* f. sp. *zeae*
GO:0016070	RNA metabolic process	Biological Process	0.023	*S. reilianum* f. sp. *zeae*
GO:1901360	organic cyclic compound metabolic process	Biological Process	0.016	*S. reilianum* f. sp. *zeae*
GO:0006725	cellular aromatic compound metabolic process	Biological Process	0.024	*S. reilianum* f. sp. *zeae*
GO:0046483	heterocycle metabolic process	Biological Process	0.023	*S. reilianum* f. sp. *zeae*
GO:0035383	thioester metabolic process	Biological Process	0.041	*S. reilianum* f. sp. *reilianum*
GO:1902589	single-organism organelle organization	Biological Process	0.048	*S. reilianum* f. sp. *reilianum*
GO:0010181	FMN binding	Molecular Function	0.047 and 0.022	*S. reilianum* f. sp. *reilianum* and f. sp. *zeae*
GO:0030515	snoRNA binding	Molecular Function	0.021	*S. reilianum* f. sp. *zeae*
GO:0016491	oxidoreductase activity	Molecular Function	0.028	*S. reilianum* f. sp. *zeae*

[a] calculated by Fisher's classic test with Parent-Child correction; only entries with *P* value ≤ 0.05 are shown

Table 2: Positively selected genes that were deleted in *S. reilianum* f. sp. *zeae* and their selection criteria

Family	Gene	Description	d_N/d_S (model free)	d_N/d_S (model join)
FAM001428	*sr10529*	conserved hypothetical protein	31.147	31.147
FAM005472	*sr10059*	conserved hypothetical *Ustilaginaceae*-specific protein	6.539	6.539
FAM000532	*sr10182*	conserved hypothetical protein	1.575	1.575
FAM002067	*sr12968*	conserved hypothetical protein	37.901	37.901
FAM003728	*sr14558*	conserved hypothetical protein	24.355	24.355
FAM004113	*sr14944*	conserved hypothetical *Ustilaginaceae*-specific protein	4.305	4.305
FAM003465	*sr14347*	conserved hypothetical protein	544.340	544.340
FAM001868	*sr12897*	conserved hypothetical protein	infinite[a]	infinite[a]
FAM000842	*sr12084*	conserved hypothetical protein	infinite[b]	infinite[b]

Table 2: Positively selected genes that were deleted in *S. reilianum* f. sp. *zeae* and their selection criteria (continued)

Gene	Species with detected positive selection	Number of paralogs[c]	Virulence phenotype of candidate gene deletion	Closest ortholog in *U. maydis* [d]
sr10529	*S. reilianum* f. sp. *zeae* and f. sp. *reilianum*	0	virulence abolished	*UMAG_01375* (*pit2*)[e]
sr10059	*S. reilianum* f. sp. *zeae* and f. sp. *reilianum*	0	virulence unaffected	*UMAG_05306* (cluster 19A)[f]
sr10182	*S. reilianum* f. sp. *reilianum*	10[1)]	virulence unaffected	*UMAG_00492*
sr12968	*S. reilianum* f. sp. *reilianum*	0	virulence unaffected	*UMAG_02006*
sr14558	*S. reilianum* f. sp. *zeae*	0	virulence unaffected	*UMAG_03564*
sr14944	*S. reilianum* f. sp. *zeae and* f. sp. *reilianum*	0	virulence unaffected	*UMAG_04034* (cluster 11-16)[g]
sr14347	*S. reilianum* f. sp. *zeae*	5[2)]	virulence unaffected	*UMAG_03349*
sr12897	*S. reilianum* f. sp. *reilianum*	0	virulence unaffected	*UMAG_01820*
sr12084	*S. reilianum* f. sp. *zeae*	2[3)]	virulence unaffected	UMAG_00792 (cluster 1-32)[g]

a) infinity due to low value of d_S

b) infinity due to long branch for *sr12085*, a species-specific duplicate

c) based on blastp search with an e-Value cutoff of 0.001

d) based on blastp search

e) as described in Doehlemann *et al*., 2011 and Mueller *et al., 2013*

f) as described in Kämper *et al*., 2006

g) as described in Schirawski *et al*., 2010

[1)] The ten paralogs include: *sr13431*, *sr11876*, *sr16607*, *sr11405*, *sr10621*, *sr16723*, *sr16877*, *13293*, *sr11163.2* and *sr15970*

[2)] The five paralogs include: *sr12257*, *sr11661*, *sr13976*, *sr14607* and *sr11273*

[3)] The two paralogs include: *sr12085* and *sr12086*

A

[46.55,24.19]

U. hordei — — — — Barley

U. maydis — — — — Maize

[23.85,13.59]

S. scitamineum — — — — Sugar cane

[16.61,7.15]

S. reilianum f. sp. *zeae* — — — —

[3.63,0.65]

S. reilianum f. sp. *reilianum* — — — — Sorghum

Divergence time (Myr ago)

−50   −40   −30   −20   −10   0

B

Divergence of *S. reilianum* f. sp. *reilianum* with...

U. hordei

U. maydis

S. scitamineum

S. reilianum f. sp. *zeae*

0%   10%   20%   30%   40%

Sequence differences

Type   Non coding   Non−synonymous

C

U. hordei

U. maydis

S. scitamineum

S. reilianum f. sp. *zeae*

S. reilianum f. sp. *reilianum*

0   30   60   90

Number of genes under positive selection

Type   All genes   Genes encoding a secreted protein (prediction)

□ plant without ears ■ healthy ears ≤ 1 cm ■ healthy ears > 1 cm ■ spiky ears ■ phyllody in ears ≤ 1 cm ■ phyllody in ears > 1 cm ■ phyllody in tassels ■ spore formation ■ dead plants

```
                20                                          40
UHOR_02064   MLHYLGRLFLVAALAVACLR  PATQ---NRPLRRAIV----  -GDNNDNYITKLHRRWYFLW
UMAG_01375   MLFRSAFVLLIVAFASACLV  QHVQAI---PVRRSLSTDAS  M----SSAAGKLNRRWWFGF
SPSC_03677   MLVHSAP-AFIATLVALCLA  QHVQAIQLPAIRRSLTHNDD  A------ANLERRWFWNF
srs_10529    MLVHSAR-AFVAALL-LGLV  LHVHAIQMPAMRRSLSSHAD  AGAAGGSTLGKLARRWFFNE
sr10529      MLVHSAR-AFVAALL-LGLV  LHVHA--MPAMRRSLSSHAD  AGAAGGSTLGKLARRWFFDF

                60                  80                  100
UHOR_02064   PGSLAPKPPREGEEHKIIYA  DWIVHHDPAYNSNVQKEIEL  ARLQNPTFIQVSVGESSSS
UMAG_01375   TGSLGKEPDNGQVQIKIIPD  ALIIKNPPANKDDLNKLIEN  LKRKHPRFKTVVMPTDPNGD
SPSC_03677   GSSLGRSPDNN---ALIVPE  DMIKKHTAALVTEWQTYLNE  MHRQHPNWKRIDWRDDGPAG
srs_10529    GGSLAPLDAVP---IFEIPK  SLIKTHKPAEVTKWEVFLQR  VHRKHPDWTHVHWTDGPVG
sr10529      GGALSRWDVAP---ILRIPE  DVAKAHSRAEVARWEVYLER  VHREHPDWQYVHWTDNGPIG

                120
UHOR_02064   SSSSSKKS-----  120
UMAG_01375   VVIWE-------  118
SPSC_03677   FARWESEKQGRSH  121
srs_10529    YKGH--------  119
sr10529      YKSH--------  117
```