# Delayed activation of the DNA replication licensing system in Lgr5(+) intestinal stem cells

T.D. Carroll[1], I.P. Newton[1], Y. Chen[1], J.J. Blow*[2], I. Näthke*[1]

Affiliations: [1]Cell & Developmental Biology and [2]Centre for Gene Regulation and Expression, University of Dundee, Dundee, Scotland, UK, DD15EH
*Correspondence to: j.j.blow@dundee.ac.uk or i.s.nathke@dundee.ac.uk

**ABSTRACT**

During late mitosis and early $G_1$, replication origins are licensed for replication by binding to double hexamers of MCM2-7. Here, we investigate how licensing and proliferative commitment are coupled in the small-intestinal epithelium. We developed a method for identifying cells in intact tissue containing DNA-bound MCM2-7. Interphase cells above the transit-amplifying compartment had no DNA-bound MCM2-7, but still expressed MCM2-7 protein, suggesting that licensing is inhibited immediately upon differentiation. Strikingly, we found most proliferative Lgr5(+) stem cells are in an unlicensed state. This suggests that the elongated cell-cycle of intestinal stem-cells is caused by an increased $G_1$ length, characterised by dormant periods with unlicensed origins. Significantly, the unlicensed state is lost In *Apc* mutant epithelium, which lacks a functional restriction point, causing licensing immediately upon $G_1$ entry. We propose that the unlicensed $G_1$ of intestinal stem cells creates a temporal window when proliferative fate decisions can be made.

**Delayed activation of the DNA replication licensing system in Lgr5(+) intestinal stem cells**

**INTRODUCTION**

Cell division is necessary for adult tissue homeostasis. It allows for the replacement of aged

or damaged cells and the provision of specialised cells critical for tissue function. The

decision to proliferate is crucial, especially for stem cells, which produce daughter cells that

either maintain a stem cell fate or differentiate to produce specialised cells. The rapidly-

renewing intestinal epithelium replenishes its cellular content every 4-5 days. This high

turnover rate is maintained primarily by Lgr5(+) intestinal stem cells in the crypt base,

thought to be continually proliferative (Basak et al., 2014) as confirmed by proteomic and

transcriptomic analysis (Munoz et al., 2012). There is also a quiescent stem cell-population

that can re-engage with the cell-cycle to repopulate the Lgr5(+) cell population if it becomes

depleted. These quiescent stem cells reside at the +4 position and constitute a subset of

Lgr5(+) cells and are immature secretory lineage precursors (Buczacki et al., 2013). Lgr5(+)

stem cells can divide to form transit-amplifying (TA) cells, which undergo several rounds of

cell division before differentiating and losing proliferative competency (Potten and Loeffler,

1990).

How proliferative fate decisions are governed in stem cells and transit-amplifying cells is not

understood. Lineage tracing studies suggest that in homeostatic intestinal tissue only 5-7

intestinal stem cells are 'active' out of the 12-16 Lgr5(+) cells present in the crypt base

(Baker et al., 2014, Kozar et al., 2013). Interestingly, Lgr5(+) cells have a significantly longer

cell-cycle than transit-amplifying cells (Schepers et al., 2011). The functional significance of

the prolonged cell-cycle time of Lgr5(+) stem cells is currently unknown, but suggests active

regulation of cell-cycle progression and proliferative fate commitment.

Proliferative fate decisions are typically visualised by detecting markers that are present in

all cell-cycle phases, and only distinguish proliferative from quiescent cells. Visualising the

incorporation of labelled nucleosides such as BrdU or EdU marks cells in S-phase. The

limitation of these methods is that they cannot discriminate early proliferative fate

decisions made during the preceding mitosis, or in the early stages of $G_1$. DNA replication in

S phase depends on origin licensing, involving the regulated loading of minichromosome

maintenance 2-7 (MCM2-7) complexes onto origins of DNA replication (reviewed in (Blow

and Hodgson, 2002, Champeris Tsaniras et al., 2014)). During S phase, DNA-bound MCM2-7

hexamers are activated to form the catalytic core of the DNA helicase as part of the CMG

(Cdc45, MCM2-7, GINS) complex (Moyer et al., 2006, Ilves et al., 2010, Makarova et al.,

2012). Replication licensing is thought to occur from late mitosis throughout $G_1$ until

passage through the restriction point (Dimitrova et al., 2002, Haland et al., 2015, Namdar

and Kearsey, 2006, Symeonidou et al., 2013). Correspondingly, insufficient origin licensing

directly limits the ability to progress past the restriction point causing cell cycle arrest (Alver

et al., 2014, Liu et al., 2009, Shreeram et al., 2002). When functional, this licensing-

checkpoint can delay S-phase if an insufficient amount of origins have been licensed.

When cells enter $G_0$, MCM2-7 proteins are downregulated and degraded, primarily via E2F-

mediated transcriptional control of MCM2-7, Cdc6 and Cdt1 (Leone et al., 1998, Ohtani et

al., 1999, Williams et al., 1998). This prevents terminally differentiated cells from re-entering

the cell cycle. In mammalian cells, artificial induction of quiescence through contact

inhibition leads to gradual downregulation of Cdc6 and MCM2-7 over several days

(Kingsbury et al., 2005). These features have led to the suggestion that quiescence can be

defined by an unlicensed state (Blow and Hodgson, 2002). Equally, the licensing status can

define a different restriction point that signals proliferative fate commitment at the end of

mitosis and in early $G_1$, independently of the Rb/E2F restriction point.

The dynamics of replication licensing in the intricate cellular hierarchy of a complex, rapidly

renewing adult tissue, is not understood. Therefore, we investigated the licensing system in

the intestinal epithelium, aiming to understand dynamics of early cell-cycle commitment in

stem and transit-amplifying cells and during terminal differentiation.

**MATERIALS AND METHODS**

**Mice**

All experiments were performed under UK home office guidelines. CL57BL/6 (Wild-type),

*R26-rtTA Col1A1-H2B-GFP* (H2B-GFP), Lgr5-EGFP-IRES-creERT2 (Lgr5$^{GFP/+}$) and Apc$^{Min/+}$ mice

were sacrificed by cervical dislocation or $CO_2$ asphyxiation. Fucci2aR mice (Mort et al., 2014)

were a kind gift of Dr Richard Mort, University of Edinburgh.

**Tissue preparation: Whole small intestine**

Dissected pieces of adult mouse small-intestine were washed briefly in PBS and then fixed in

4% PFA for 3 hours, 4°C. Intestines were cut into 2x2 cm$^2$ pieces and fixed overnight in 4%

PFA, 4°C. Tissue was embedded in 3% low melting temperature agarose and cut into 200 μm

sections using a Vibratome (Leica). Sections were washed in PBS, permeabilised with 2%

Triton-X100 for 2 hours and incubated with Blocking Buffer (1% BSA, 3% Normal Goat

serum, 0.2% Triton-X100 in PBS) for 2 hours, 4°C. Tissue was incubated in Working Buffer

(0.1% BSA, 0.3% Normal Goat Serum, 0.2% Triton-X100 in PBS) containing primary antibody,

Mcm2 (Cell Signalling, 1:500), for 48 hours, 4°C. Sections were washed 5x with Working

Buffer prior to 48 hour incubation with secondary antibodies diluted in Working Buffer:

Alexafluor™ conjugated goat anti-rabbit (1:500, Molecular Probes) plus 5 μg/ml Hoechst

33342 and Alexafluor™ conjugated Phalloidin (1:150, Molecular Probes). Sections were

mounted on coverslips in Prolong Gold between 2x120 μm spacers.

**Tissue preparation:**

**Isolating and staining crypts**

Small intestines were dissected, washed in PBS and opened longitudinally. Villi were

removed by repeated (up to 10 times) scraping of the luminal surface with a coverslip.

Tissue was washed in PBS, incubated in 30 mM EDTA (25 minutes, 4°C) and crypts isolated

by vigorous shaking in PBS. Crypt suspensions were centrifuged (fixed rotor, 88 RCF, 4°C)

and the pellet washed twice in cold PBS. Crypts were fixed in 4% PFA (30min, room

temperature), permeabilized in 1% Triton-X100 (1 hour, room temperature) and blocked in

Blocking Buffer (2 hours, 4°C). Crypts were incubated with primary antibodies diluted in

Working Buffer: Mcm2 (Cell Signalling, 1:500), phospo-HistoneH3 (Abcam, 1:500), Ki67

(Abcam ab15580, 1:250), αGFP (Abcam, 1:500), washed 5x with Working Buffer before

overnight incubation with secondary antibodies diluted in Working buffer: Alexafluor™

conjugated goat anti-mouse or anti-rabbit (1:500, Molecular Probes) or stains: Rhodamine

labelled Ulex Europaeus Agglutinin I (UEA, 1:500), 5 μg/ml Hoechst 33342 or Alexafluor™

conjugated Phalloidin (1:150), at 4°C. Crypts were mounted directly on slides in Prolong

Gold, overnight.

**107**  **CSK extraction of isolated crypts**

**108**  Soluble proteins were extracted from crypts isolated as described above by incubation with

**109**  CSK extraction buffer (10 mM HEPES, 100 mM NaCl, 3 mM $MgCl_2$, 1 mM EGTA, 300 mM

**110**  sucrose, 0.2% TritonX-100, 1 mM DTT, 2% BSA) supplemented with protease inhibitors

**111**  (PMSF, Pepstatin, Leupeptin, Cystatin, $Na_3VO_4$, NaF, aprotinin) for 20 minutes on ice prior to

**112**  fixation. Crypts were then fixed with 4% PFA and processed for imaging as described above.

**113**  **H2B-GFP label retention**

**114**  H2B-GFP expression in transgenic *R26-rtTA Col1A1-H2B-GFP* mice was induced by replacing

**115**  normal drinking water with 5% sucrose water supplemented with 2 mg/ml doxycycline.

**116**  After 7 days, doxycycline water was replaced with normal drinking water. Subsequently,

**117**  mice were sacrificed after 7 days.

**118**  **EdU incorporation and detection**

**119**  Mice were injected intraperitoneally with 100 µg EdU (Invitrogen) prepared in 200 µl sterile

**120**  PBS. Mice were sacrificed 1 hour or 17 hours post induction. For organoids, 10 µM EdU was

**121**  included in crypt media for 1 hour before harvesting. EdU was detected by Click-it

**122**  chemistry, by incubation in EdU working buffer (1.875 µM Alexafluor 488 azide (Invitrogen),

**123**  2 mM $CuSO_4$, 10 mM Ascorbic acid), overnight at 4°C, prior to processing for

**124**  immunofluorescence staining.

**125**  **Organoid Culture**

**126**  Isolated crypts were dissociated to single cells with TripLE express (Life Technologies) at

**127**  37°C, 5 minutes. Dissociated cells were filtered through a 40 µm cell strainer (Greiner) and

suspended in growth factor reduced Matrigel (BD Biosciences). Organoids were grown in

crypt media (Advanced DMEM/F12 (ADF) supplemented with 10 mM HEPES, 2 mM

Glutamax, 1 mM N-Acetylcysteine, N2 (Gemini), B27 (Life technologies),

Penicillin/Streptomycin (Sigma) supplemented with growth factors – ENR media (EGF (50

ng/ml, Invitrogen), Noggin (100 ng/ml, eBioscience) and RSpondin conditioned media

produced from stably transfected L-cells (1:4). Chiron99021 (3 µM), Valproic acid (1 mM,

Invitrogen) and Y27632 (10 µM, eBiosciences) were added to the culture for the first 48

hours. Organoids were passaged every 3-5 days by mechanically disrupting Matrigel and by

washing and pipetting in ADF. Dissociated crypts were re-suspended in fresh Matrigel and

grown in crypt media supplemented with growth factors.

For small molecule treatments, primary intestinal epithelial cells were cultured in ENR-CVY

(ENR plus Chiron99021, Valproic acid and Y27632) for 3 Days, and then organoids were sub-

cultured in ENR for two further days prior to the start of the experiment. Organoids were

then treated with the stated small molecules for the indicated time periods. For induction of

Unlicensed-G$_1$, organoids were treated with Gefitinib (5 µM) coupled with removal of EGF

from the crypt media. For re-activation/chase period, the media was removed and fresh

growth factors added. All growth factors and inhibitors were replenished every 2 days

throughout the experiment.

**Flow cytometry and cell sorting**

Intestinal crypts were isolated and dissociated to single cells as described above. Isolated

cells were filtered through 40 µm cell strainers (Greiner). For organoids, following one PBS

wash, organoids were dissociated to single cells by incubation in TripLE express for 15

minutes at room temperature followed by manual disruption by pippetting. Cells were then

extracted with CSK buffer for 20 minutes on ice, followed by fixation in 0.5% PFA (pH7.40,

15 minutes, room temperature). Cells were then washed once in 1% BSA and permeabilized

with ice-cold 70% EtOH, 10 minutes. Cells were then washed in 1% BSA and re-suspended

with primary antibodies (Mcm2, 1:500; GFP, 1:500; Ki67, 1:200) diluted in Working buffer

(overnight, 4°C). Following two washes in working buffer, cells were re-suspended in

secondary antibodies goat anti-mouse or anti-rabbit (Alexafluor647, 1:500 (Molecular

Probes); Alexafluor488-Ki67, 1:400 (Clone SolA15, BD Biosciences), diluted in working buffer

(1 hour, room temperature). After two washes in 1% BSA, cells were suspended in working

buffer containing 15 µg/ml DAPI. Samples were analysed on a FACS Canto (BD Biosciences).

For cell sorting, cells were isolated from Lgr5-GFP mice as described above by treatment

with TripLE express for 15 minutes, 37°C followed by filtration through 40 µm filters

(Greiner). Cells were sorted in ADF supplemented with 1% FBS and DAPI (15 µg/ml). Sorting

was performed using an Influx$^{TM}$ Cell sorter (BD biosciences). Cells were checked post-sort

to ensure sample purity by re-examining Lgr5 expression in the sorted gates.

**Microscopy and Image analysis**

Samples were imaged using a Zeiss LSM 710 microscope using a 40X LD Pan-Neofluar

objective lens and immersion oil with a refractive index of 1.514. Z-stacks were acquired at

optimal section intervals between 0.3 and 0.8 µm. For quantification, images were acquired

at 16 bit-depth.

Image processing and analysis were performed using Imaris (Bitplane). Images of individual

crypts were manually cropped, ensuring that an individual crypt was the only region of

interest. All nuclei were detected in individual crypts using automated thresholding in Imaris

using the measurement point function, set to detect nuclei at an estimated size of 3.5 μm.

Missed or incorrectly assigned nuclei were manually identified. This function produced

measurement points that segmented the specific region at the corresponding co-ordinate of

the measurement point. Mean intensities for different channels were calculated per spot.

This equates to the intensity at the centre region of each nucleus. A reference nucleus at the

crypt base was used to define the crypt base position. The Euclidean distance to this point

was measured and defined as the distance to the crypt base. Multiple images were analysed

using the same workflow and the analysed files collated. For vibratome sections, a plane

was manually defined running through to the muscle layer beneath the epithelium. The

smallest distance to this surface was defined for segmented nuclei. For nuclear volume

estimation, nuclei were manually segmented in 3D using the manual segmentation tools

within Imaris (**S1 Figure**).

**Flow cytometry analysis**

Flow cytometry data was analysed using FlowJo (Treestar) using a standardised gating

strategy (**S1 Figure**). Briefly, cells were identified using FSC and SSC. Following doublet

discrimination, gates were set using appropriate controls lacking conjugated secondary

antibodies and without primary antibodies. Mcm2 negative gates were set by secondary

only controls in conjunction with the Mcm2 intensity of $G_2$ cells. $G_1$ cells were discriminated

based on the maximal DNA-bound Mcm2 intensity prior S-phase, and by DAPI intensity.

**Computational Modelling**

A deterministic computer model for licensing in G1 was written in the Swift programming

language using the Xcode 9 development environment. The model assumes that there is a

minimum G1 period that is required for cells to grow to a critical size before they can enter S

phase. Licensing can take place during this period at a constant rate. A licensing rate of 1

means that cells will be fully licensed in exactly the minimum G1 period. It is also assumed

that cells have a robust 'licensing checkpoint' (Shreeram et al., 2002, Blow and Gillespie,

2008) so that they cannot enter S phase until origins have been fully licensed. An optional

'unlicensed G1 period' occurs at the start of G1, during which time no licensing takes place.

In the simulation, cells enter G1, wait for any optional 'unlicensed G1 period', then start to

license origins at a fixed rate; cells exit G1 and modelling ceases only when they have

become maximally licensed and also the minimum G1 period has elapsed. The model divides

the minimum G1 period into 10,000 equal steps and records the degree of licensing at the

end of each step in the 'Licensing Array'. The contents of the Licensing Array are distributed

into 101 different frequency bins, ranging from 0% (no licensing) to 100% (maximal

licensing). To model the background signal recorded by flow cytometry, all the licensing

values are increased by 5%. To model flow cytometry measurement error, the frequency

array is smoothed by starting at the smallest bin and pushing 80% of the cell counts into the

next largest bin, and then starting at the largest bin pushing 80% of the cell counts into the

next smallest bin.

```
212   //
213   //  JBLicensingRateModel.swift version 1.0
214   //
215   //  Created by Julian Blow on 05/10/2017.
216
217
218   import Foundation
219
220   class JBLicensingRateModel {
221      let licensingRate: Double    // rate of licensing in proportion to minimum G1 length
222      let proportionUnlicensedG1: Double     // proportion of minimum G1 before licensing occurs
223      let timeStepsForMinimumG1: Int
224      let licensingPerStep: Double
225      let timeStepsBeforeLicensingStarts: Int
226      let binsForFACS: Int
227      let backgroundForFACS: Double
228      let binSmoothingProportion: Double
229      let totalNumberOfCells: Double
230      let precisionCutOffForSmoothing: Double
231      let numberOfTimepointsOutput: Int
232
233      init (licensingRate: Double, proportionUnlicensedG1: Double) {
234         self.licensingRate = licensingRate
235         self.proportionUnlicensedG1 = proportionUnlicensedG1
236         timeStepsForMinimumG1 = 10000
237         licensingPerStep = (licensingRate / Double(timeStepsForMinimumG1)) * 100
238         timeStepsBeforeLicensingStarts = Int(proportionUnlicensedG1 * Double(timeStepsForMinimumG1))
239         binsForFACS = 100
240         backgroundForFACS = 5
241         binSmoothingProportion = 0.8
242         totalNumberOfCells = 10000.0
243         precisionCutOffForSmoothing = totalNumberOfCells / 10000     // cutoff for the smallest number of cells we continue to
244   smooth for the far right of the array
245         numberOfTimepointsOutput = 100
246      }
247
248      func run () -> ([Double], [Double]) {
249         var licensingTimepoints: [Double]
250         if licensingRate > 0 { licensingTimepoints = simulateLicensing() }
251         else { licensingTimepoints = Array(repeating: 0.0, count: timeStepsForMinimumG1) }     // if licensing can never occur
252         let FACSOutputExact = simulateFACSExact(licensingTimepoints: licensingTimepoints)
253         let FACSOutput = smoothFACSOutput(FACSOutputExact: FACSOutputExact)
254         var simplifiedLicensingTimepoints = [Double]()
255         for index in stride(from: 0, to: licensingTimepoints.count, by: licensingTimepoints.count/numberOfTimepointsOutput) {
256            simplifiedLicensingTimepoints.append(licensingTimepoints[index])
257         }
258         return (FACSOutput, simplifiedLicensingTimepoints)
259      }
260
261      func simulateLicensing() -> [Double] {
262      // simulates the degree of MCM loading given the licensingRate and proportionUnlicensed
263      // returns amount of MCM loaded (percent max) at different time points
264         var amountOfLicensing = 0.0
265         var licensingArray = Array(repeating: 0.0, count: timeStepsBeforeLicensingStarts)
266         var timeStep = timeStepsBeforeLicensingStarts
267
268         repeat {
269            amountOfLicensing += licensingPerStep
270            if amountOfLicensing > 100 { amountOfLicensing = 100 }
271            licensingArray.append(amountOfLicensing)
```

```
272        timeStep += 1
273      } while (amountOfLicensing < 100) || (timeStep < timeStepsForMinimumG1)
274
275      return licensingArray
276    }
277
278    func simulateFACSExact (licensingTimepoints: [Double]) -> [Double] {
279      // takes an array amount of MCM loaded at different time points as created by simulateLicensing()
280      // turns this into a simulated output from a FACS (flow cytometry) analysis with totalNumberOfCells
281      var maxMCMValue = licensingTimepoints.max()! // can be >100 because of smoothing
282      if maxMCMValue == 0 { maxMCMValue = 100 }    // if no licensing occurred
283      var FACSOutputExact = Array(repeating: 0.0, count: binsForFACS+1)      // binsForFACS+1 because we 0 and 100 are
284  both possible
285      let numberOfCellsPerTimepoint = totalNumberOfCells / Double(licensingTimepoints.count)         // so we get
286  totalNumberOfCells events
287      for anMCMContent in licensingTimepoints {
288        let binIndex = Int(anMCMContent * Double(binsForFACS) / maxMCMValue)
289        assert(binIndex <= binsForFACS, "Cannot have an MCM content greater than 100, the bin number")
290        if binIndex>=0 { FACSOutputExact[binIndex] += numberOfCellsPerTimepoint }            // add cells to bin
291      }
292
293      return FACSOutputExact
294    }
295
296    func smoothFACSOutput (FACSOutputExact: [Double]) -> [Double] {
297      // simulates the error in FACS data by:
298      // first adding a background of backgroundForFACS to all samples
299      // then pushing binSmoothingProportion of each bin into the higher bin, when the value > cutOffForDither
300      // then pushing binSmoothingProportion of each bin into the lower bin
301      let backgroundBins = Array(repeating: 0.0, count: Int(backgroundForFACS))
302      var FACSOutput = backgroundBins + FACSOutputExact       // add the background of empty pbins
303      var binIndex = 0
304      var amountToShift = 0.0
305    repeat {                    // expand rightwards by dithering
306        amountToShift = FACSOutput[binIndex] * binSmoothingProportion
307        FACSOutput[binIndex] -= amountToShift
308        if binIndex+1 < FACSOutput.count { FACSOutput[binIndex+1] += amountToShift }
309        else { FACSOutput.append(amountToShift) }
310        binIndex += 1
311      } while (amountToShift >= precisionCutOffForSmoothing) || (binIndex < FACSOutputExact.count +
312  Int(backgroundForFACS))
313
314      binIndex = FACSOutput.count-1
315      while binIndex >= 0 {              // expand leftwards by dithering
316        amountToShift = FACSOutput[binIndex] * binSmoothingProportion
317        FACSOutput[binIndex] -= amountToShift
318        if binIndex > 0 { FACSOutput[binIndex-1] += amountToShift }
319        binIndex -= 1
320      }
321
322      return FACSOutput
323    }
324
325  }
```

**RESULTS**

**Mcm2 expression declines along the crypt-villus axis**

Because of their abundance, strong conservation and association with the core DNA

replication process, the presence of MCM2-7 proteins is commonly used to establish

proliferative capacity in tissues, similar to Ki67 or PCNA (Gonzalez et al., 2005, Jurikova et

al., 2016, Stoeber et al., 2001, Williams et al., 1998). Usually, terminally differentiated cells

in mammalian tissues do not contain MCM2-7 (Stoeber et al., 2001, Todorov et al., 1998,

Eward et al., 2004). To establish the overall MCM2-7 protein abundance along intestinal

crypts, we first examined the expression of MCM2-7 proteins in adult murine small-

intestinal epithelium using high-resolution immunofluorescence microscopy. We focused on

Mcm2 as a surrogate for all the members of the MCM2-7 complex based on their similar

function and localisation. However, we repeated a subset of the experiments using an

antibody to Mcm4, which is less effective in detecting endogenous proteins. Nonetheless, in

all cases, the results were identical.

Consistent with previous reports, Mcm2 was highly expressed in both murine and human

intestinal epithelium. Mcm2 was highly expressed in intestinal crypts (**Figure 1A**) and

declined gradually along the crypt-villus axis (**Figure 1B**), but persisted in a few cells in the

villus compartment (**Figure 1D**). Mcm2 was nuclear in interphase cells but cytoplasmic

during mitosis (**Figure 1C**). Although the majority of intestinal crypt cells expressed Mcm2,

at the crypt base, Mcm2(+) and Mcm2(-) cells were interspersed (**Figure 1A, D**), consistent

with previous reports (Pruitt et al., 2010). This pattern is reminiscent of the alternating

arrangement of Lgr5(+) stem cells and Paneth cells at the crypt base (Barker et al., 2007).

Lgr5(+) stem cells express Ki67 and are continually proliferative whereas Paneth cells are

fully differentiated and are Ki67(-) (Basak et al., 2014). As expected, Mcm2 was expressed in

all Lgr5(+) stem cells and there was a strong correlation between Mcm2 and Lgr5 expression

(**Figure 1E**). This is consistent with the idea that Lgr5[Hi] stem cells are the main proliferative

stem cells in the intestinal crypt. Staining with Ulex Europaeus Agglutinin I (UEA),

demonstrated that most Mcm2(-) cells in the crypt base are UEA(+) Paneth cells (**Figure 1F**).

Normally, MCM2-7 expression is lost in terminally differentiated cells (Eward et al., 2004,

Stoeber et al., 2001, Williams et al., 1998, Williams et al., 2004). The loss of expression has

been suggested as a major contributor to the proliferation-differentiated switch *in vivo.* To

test this idea, we measured the Mcm2 content of young and mature secretory cells in

intestinal crypts and villi (**Figure 1G, H**). There was differential expression of Mcm2 in

distinct secretory lineages. Many mature secretory cells including Paneth, Goblet and

enteroendocrine cells were Mcm2(-), consistent with their differentiation status and long

life-span in the epithelium (van der Flier and Clevers, 2009). We detected a small number of

UEA(+) Mcm2(+) cells in intestinal crypts (**Figure 1G**). Assuming that Mcm2 expression

declines slowly after terminal differentiation, the presence of Mcm2 in UEA(+) secretory

cells could reflect their immaturity. Consistently, Mcm2 expression in UEA(+) cells in crypts

was significantly higher than in villi (**Figure 1H**), supporting the idea that MCM2-7 are

gradually lost upon terminal differentiation. Since MCM2-7 are highly abundant and have a

long (>24 hour) half-life (Musahl et al., 1998), it likely that after cells differentiate, their

MCM2-7 content declines at a slow rate, explaining why Mcm2 persists in the villus

compartment.

**Visualisation of DNA replication licensing *In vivo***

MCM2-7 exist in three states: as hexamers free in the nucleoplasm, as double hexamers

bound to DNA during late mitosis, G1 and S phase, or as CMG complexes at replication forks

during S phase (Evrin et al., 2009, Gambus et al., 2011, Remus et al., 2009). To distinguish

between DNA-bound and soluble forms, we developed a protocol involving a brief

extraction of isolated crypts with non-ionic detergent to remove soluble MCM2-7. The

remaining Mcm2 should mark cells whose origins are licensed for replication. Extraction did

not visibly affect intestinal crypt integrity but made them more opaque compared to

unextracted tissue (**Figure 2A**). The majority of cells in unextracted crypts were Mcm2(+)

(**Figure 2B,** 'Total') similar to tissue sections and mirroring the ubiquitous expression of Ki67

along the crypt axis. After extraction, the majority of the Mcm2 content in cells was lost

(**Figure 2B**, 'Licensed'), with only 10-30% of cells maintaining high levels of Mcm2 (**Figure**

**2C**). After extraction, Mcm2(+) was not present in mitotic cells expressing phosphorylated

histone H3, confirming the extraction procedure successfully removed non-DNA bound

MCM2-7 proteins (**Figure 2D**).

We used flow cytometry to measure MCM2-7 content more directly and further confirm the

effectiveness of the extraction procedure. Whereas the majority of isolated epithelial cells

expressed Mcm2 that persisted throughout the cell-cycle, extraction revealed a distinct

profile of Mcm-containing cells in crypts (**Figure 2E**). These profiles are consistent with those

reported for cultured cell lines (Friedrich et al., 2005, Haland et al., 2015, Moreno et al.,

2016, Matson et al., 2017). Mcm2 is present throughout the cell cycle (**Figure 2E**,

'Unextracted'), but extraction shows that it binds to DNA throughout $G_1$, reaching a

maximum level before cells enter S-phase, and is subsequently displaced from DNA during S

phase. This behaviour, which matches the known cell cycle behaviour of MCM2-7, confirms

the efficiency of our extraction protocol. An antibody against Mcm4 produced similar results

(data not shown). We observed that the majority of cells with a G1 DNA content appeared

to be unlicensed, having a DNA-bound Mcm2 content similar to $G_2$/M cells (**Figure 2E, E'**).

This is substantially different from typical profiles observed in cultured cells lines in which

most $G_1$ cells are fully licensed (Friedrich et al., 2005, Haland et al., 2015, Moreno et al.,

2016, Matson et al., 2017). Similar results were observed in cells isolated from intestinal

organoids (**Figure 2F, G**).

**Licensing status and cell-cycle progression along the crypt-villus axis**

Cell-cycle dynamics of intestinal stem and progenitor cells are highly heterogeneous (Pruitt

et al., 2010). The majority of Lgr5(+) stem cells are considered to be continually

proliferative, but with a much longer cell-cycle than transit-amplifying progenitor cells,

which are most commonly found in S-phase (Schepers et al., 2011). To investigate

proliferative fate decisions of intestinal epithelial cells, we used our MCM2-7 extraction in

crypts where S phase cells were labelled in vivo with the nucleoside analogue EdU. We then

used image analysis software to correlate Mcm2 content with cell-cycle stage along the

crypt-villus axis (**S1 Figure A-F**). This allowed quantification of licensing in relation to the

cell-cycle and 3D spatial information.

Figure 3A shows tissue labelled in vivo with a 1 hour EdU pulse followed by extraction of

soluble MCM2-7. As expected, the majority of cells in the transit-amplifying compartment

were labelled with EdU suggesting that most cells were in S-phase, consistent with early

studies using BrdU and [$^3$H]-thymidine labelling (Chwalinski and Potten, 1987). The patterns

of replication foci were consistent with the reported S-phase replication timing programme

(Rhind and Gilbert, 2013). Typically, all licensed cells had intense nuclear Mcm2 staining.

Some cells completely lacked Mcm2 and EdU labelling, suggesting they were in either $G_0$,

very early $G_1$ or in $G_2$. Some cells were labelled with both Mcm2 and EdU. These double-

labelled cells typically showed patterns of EdU labelling consistent with early to mid S phase

and Mcm2 labelling of DNA compartments expected to replicate later in S phase. This

relationship has been observed in tissue culture cells (Krude et al., 1996) and is consistent

with the idea that DNA-bound MCM2-7 are displaced from chromosomal domains as

replication is completed. Cells with late S-phase patterns of EdU labelling had little or no

detectable Mcm2, consistent with the displacement of the majority of MCM2-7 by the end

of S phase. We also measured nuclear volume, which increases during S phase and $G_2$. This

showed that nuclear volume increased up to two-fold in cells classified as S-phase and Late-

$S/G_2$ by Mcm2 and EdU staining (**Figure 3B)**. This confirms our cell-cycle assignment and also

suggests that most Mcm2(-) cells are in $G_0$ or $G_1$, rather than in $G_2$.

The combination of concurrently labelling DNA-bound Mcm2 and EdU showed a clear

correlation between cell position and cell-cycle stage. As noted previously, Mcm2 is

expressed in cells throughout the crypt (**Figure 3C, D**). At the base of the crypt, unlicensed

cells predominate (**Figure 3C, E**). At increasing distances from the crypt base there is a

successive rise in licensed $G_1$ cells, early/mid S phase cells and then late $S/G_2$ cells. Further

up the crypt, at the end of the TA compartment, these cell cycle stages decline in reverse

order, until unlicensed cells again predominate. This suggests that there is a co-ordinated

progression through the cell division cycle as cells enter, then leave, the TA compartment.

This was also observed as a field effect with many neighbouring cells showing similar

replication patterns (**S2 Figure A, B; S1 Movie**).

**Terminal differentiation is associated with a binary licensing decision**

At the terminal boundary of the transit-amplifying compartment, the majority of cells were

unlicensed and had no DNA-bound Mcm2 (**Figure 3C, E**). Similarly, there were no licensed $G_1$

cells beyond the TA compartment as defined by incorporation of EdU (**Figure 3F**). However,

total Mcm2 expression extended significantly beyond the last cells with DNA-bound Mcm2

or incorporated EdU (**Figure 3C, D, F**). The distribution of total Mcm2 expression

corresponded to the zone where cells express Ki67 (**S3 Figure**). Although Mcm2 and Ki67

expression persists beyond the TA compartment, licensing does not occur in this area. This

suggests that differentiation is not governed by a gradual reduction in total MCM2-7 levels,

but is a binary decision and licensing is abolished immediately after the final mitosis

preceding differentiation. To further examine this, we marked the terminally differentiated

zone by a 1 hour EdU pulse, followed by a 16 hour chase (**Figure 3G, H**). After 16 hours, the

majority of the distal end of the TA compartment became labelled with EdU. All labelled

nuclei in this area were significantly smaller than EdU(+) cells at the proximal end of the TA

compartment (data not shown), suggestive of their differentiation status. Importantly the

EdU(+) differentiated cells at the distal end of the TA compartment lacked DNA-bound

Mcm2, supporting our suggestion that licensing is inhibited immediately at terminal

differentiation.

**The majority of Intestinal stem cells spend the majority of G$_1$ in an unlicensed state**

The majority of cells in the crypt base expressed Mcm2, consistent with the finding that all

Lgr5(+) cells express Mcm2 but mature secretory cells, such as Paneth cells, do not (**Figure**

**1D, E**). Surprisingly, extraction revealed that only 7-15% of cells were licensed in the crypt

base (**Figure 3C, D**), with most cells in an unlicensed state despite expressing Mcm2. The

abundance of licensed cells peaked 40-60 µm away from the crypt base, corresponding to

just above the +4/+5 cell position (**Figure 3D, E**). This suggests that the majority of stem cells

remain unlicensed. In contrast, the majority TA cells appear to progress rapidly through the

cell-cycle, as many more actively incorporate EdU (**Figure 4A**).

We next confirmed that the unlicensed cells at the crypt base were Lgr5(+) stem cells. As it

is not possible to identify Lgr5 in these experiments, as it is extracted along with unbound

Mcm2, we instead identified Paneth cells by UEA staining and considered all UEA(-) cells in

the crypt base to stem cells (**Figure 4B**). >50% of the UEA(-) stem cells were in an unlicensed

state and were not incorporating EdU (**Figure 4C**). Approximately 30-40% of all UEA(-) cells

in the stem cell compartment were in an active phase of the cell cycle (licensed $G_1$, S or $G_2$;

**Figure 4C**), corresponding to 5-6 stem cells out of the total 14 present (Snippert et al.,

2010). This number is similar to the small number of proposed 'working' stem cells in the

crypt base (Baker et al., 2014, Kozar et al., 2013). Unlicensed cells not incorporating EdU (i.e.

unlabelled in this experiment) could theoretically be in either $G_0$ or in $G_2$. To distinguish

between these possibilities we first isolated crypt cells from Lgr5-GFP mice and measured

both GFP and DNA content. Both Lgr5(+) and Lgr5(-) cell populations had a similar cell-cycle

profile with the majority of cells having 2N DNA content (**S2 Figure C**). We also examined the

nuclear volume of cells at different positions along the crypt axis, after staining for EdU

incorporation and DNA-bound Mcm2. The majority of unlicensed cells had a similar nuclear

volume to that of fully licensed cells in $G_1$ and not cells in Late-S/$G_2$ phase (**S2 Figure D**).

Together, these results suggest that, although they express abundant Mcm2, the majority of

intestinal stem cells reside in an unlicensed $G_1$ state.

To confirm this conclusion, we flow-sorted unextracted Lgr5-GFP(+) cells, extracted

unbound MCM2-7 and stained them for Mcm2 and Ki67. Consistent with our previous

results, most Lgr5(+) cells with a 2N DNA content had low levels of DNA-bound Mcm2, and

were in an unlicensed state (**Figure 4Di, ii**). Importantly, both the licensed and unlicensed

cells were Ki67(+) indicating that they had not withdrawn from the cell-cycle long-term

(**Figure 4Dii**).

This unlicensed $G_1$ state - 2N DNA content, high Mcm2 expression but with low levels of

DNA-bound Mcm2 - could be explained by two, slightly different, scenarios. 1) MCM2-7 are

loaded on to DNA slowly in Lgr5(+) cells, thereby extending $G_1$ (Schepers et al., 2011)

(Dalton, 2015); or 2) Most Lgr5(+) cells enter $G_1$ and remain in an unlicensed state but do

not load MCM2-7 until an active decision is made to commit to cell cycle progression and

activate the licensing system, at which time MCM2-7 proteins are rapidly loaded. In option

1) where licensing is slow, the presence of unlicensed cells simply reflects the increased

time required to fully license origins, and different levels of Mcm2 loading should be equally

distributed between $G_1$ cells. In option 2), however, where there is an early stage of G1

where licensing does not occur, there should be a discrete peak of unlicensed cells with a $G_1$

DNA content representing cells that have withdrawn from the cell cycle, and a lower

frequency of $G_1$ cells that have loaded different amounts of MCM2-7. To distinguish

between these two possibilities, we utilized ergodic rate analysis. When examining the

frequency distribution of DNA-bound Mcm2 in Lgr5(+) cells with 2N DNA content we found

a discrete peak of unlicensed cells (**Figure 4E**) consistent with the second model. To confirm

these observations, we performed computer modelling in which the licensing rate was

varied in the presence or absence of an initial $G_1$ period when licensing did not occur.

(**Figure 4F**). With no unlicensed G1 period and low licensing rates (**Figure 4F**, left hand

panels), G1 cells were equally distributed across the different degrees of licensing. With

faster licensing rates and no unlicensed G1 period, a distinct peak of fully licensed G1 cells

appeared, similar to what has been observed in cell lines. Only when an unlicensed G1

period was introduced did a distinct peak of unlicensed G1 cells appear (**Figure 4F**, left hand

panels). Since most of these unlicensed Lgr5(+) cells express abundant Mcm2 (**Figure 1D**),

this suggests that their $G_1$ is characterised by a long unlicensed period. This may explain why

the cell-cycle length of intestinal stem cells is significantly longer than transit-amplifying

cells (Schepers et al., 2011).

To further confirm that many intestinal stem cells exist in an unlicensed $G_1$ state, we

examined the expression of Fluorescence ubiquitination cell-cycle indicator (FUCCI)

reporters as an independent indicator of cell-cycle progression. We harvested intestinal

tissue from Fucci2aR mice (Mort et al., 2014), and examined the expression of the $G_1$

specific hCdt1(30/120) and $S/G_2/M$ specific hGeminin(1/110) reporters (**S4 Figure A**). As

expected, many TA cells were hGeminin(1/110)(+). We also noticed that terminally

differentiated cells, such as Paneth cells in the stem cell compartment, and cells at the tips

of villi expressed high levels of hCdt1(30/120) (**S4 Figure B**) reflecting accumulation of the

reporter in differentiated cells (Mort et al., 2014). However, we found that most Mcm2(+)

stem cells in the crypt base expressed very low levels of the hCdt1(30/120) reporter, and

only few cells expressed high levels (**S4 Figure C, D**). In contrast, the majority of TA cells

were either hGeminin(1/110)$^{High}$ or hCdt1(30/120)$^{low}$, consistent with a short $G_1$ phase.

Together, this suggests that stem cells remain in a paused unlicensed $G_1$ state where they

do not rapidly accumulate hCdt1(30/120) unlike rapidly proliferating cells.

It has previously been reported that embryonic stem cells license more replication origins

than neural stem/progenitor cells differentiated from them (Ge et al., 2015). To determine if

adult stem and non-stem cells in intestinal crypts exhibit such differences, we compared the

amount of DNA-bound Mcm2 in $G_1/G_0$ and early S phase Lgr5(+) cells with that in Lgr5(-)

cells (Moreno et al., 2016). Although the majority of Lgr5(+) cells were unlicensed, when

they entered S phase they had approximately twice as much DNA-bound Mcm2 compared

to Lgr5(-) cells (**Figure 4G**). This is consistent with the idea that adult intestinal stem cells

license more origins than TA cells and may represent a mechanism to protect genomic

integrity.

**Intestinal label retaining cells are in a deep $G_0$ state**

Although the intestinal crypt base primarily consists of Lgr5(+) stem cells there is also a

reserved pool of quiescent stem cells, often referred to as '+4 label retaining cells' (LRCs)

reflecting their position in the crypt base and their ability to retain nascent DNA labels

(Potten et al., 2002). These cells are a rare subset of Lgr5(+) cells and are also secretory

precursors (Buczacki et al., 2013). To further define the licensing status of these label-

retaining intestinal stem cells, we identified UEA(-) LRCs by expressing H2B-GFP (which is

incorporated into the chromatin of dividing cells) for 7 days and then chasing for a further 7

days (Buczacki et al., 2013, Roth et al., 2012). Labelled cells that did not divide during the 7-

day chase period contain high levels of H2B-GFP (hence are LRCs), but cells that divided

multiple times only have low levels. LRCs were then distinguished from Paneth cells based

on UEA staining (**Figure 5A**). After induction, most cells in the epithelium expressed H2B-

GFP (**Figure 5Bi**). After the 7D chase, H2B-GFP expression was restricted to cells near the

villus tips, and cells at the crypt base (**Figure 5Bii**). We could successfully distinguish

between Paneth cells and LRCs on the basis of UEA staining (**Figure 5Bii**). Unlike the majority

of the Lgr5(+) cells, LRCs with high GFP-H2B did not express Mcm2 (**Figure 5C**). As expected,

only non-LRC daughter cells with low levels of H2B-GFP had DNA-bound Mcm2 (**Figure 5D**).

This shows that, LRC stem cells are in deep $G_0$, unable to license because they do not

express MCM2-7. In contrast, 'active' intestinal stem cells mostly reside in an unlicensed $G_1$

state.

**Licensing dynamics in intestinal organoids**

Intestinal stem cells reside in a highly specialised niche at the base of crypts. It is therefore

possible that this niche specifically allows stem cells to pause in an unlicensed $G_1$ state

where origin licensing is prevented until a further proliferative fate signal is received. To

understand the contribution of the stem cell niche to the dynamics of entry and exit from

the unlicensed $G_1$ state, we used intestinal organoids, which allowed us to manipulate the

stem cell niche/environment with small-molecules. Although the distribution of licensed

cells was similar between the branches of intestinal organoids and crypts in tissue, there

were considerably more cells with DNA-bound Mcm2 in the former (**Figure 6A, Ciii**).

Importantly, cells in organoids showed a discrete peak of fully licensed $G_1$ cells in addition to

the cells in the unlicensed $G_1$ state (**Figure 6Bii**). This suggests that the epithelium in

organoids represents an accelerated state of self-renewal and may not fully recapitulate

cell-cycle dynamics of intestinal epithelial cells *in vivo*.

We designed an assay to robustly assess licensing dynamics during entry and exit from the

unlicensed $G_1$ state, and transition towards $G_0$. Specifically, we used flow cytometry based

quantification of DNA-bound Mcm2, Ki67 and DNA content to measure licensing dynamics

during entry and exit from the unlicensed $G_1$ state, and transition towards $G_0$ (**S5 figure A**).

Most cells in organoids express Ki67 and it increased during cell-cycle progression (**Figure**

**6Bi**). The DNA-bound Mcm2 profile was similar to that in isolated crypts (**Figure 6Bii**).

Correlating Ki67 and DNA-bound Mcm2 produced a distinctive profile that is similar for

isolated cells from organoids (**Figure 6Ci**) and intestinal crypts (**Figure 6Cii**). This profile

reveals a population of cells that appear to lose Ki67 (dashed arrow in **Figure 6Cii**) and may

represent cells decreasing in proliferative capacity and transitioning towards differentiation

(**S5 Figure A**). Such a loss of proliferative capacity appeared to initiate in cells that express

Ki67 but are unlicensed, i.e. cells in unlicensed $G_1$. These data suggest that different stages

of quiescence can exist that reflected by a spectrum of Ki67 and Mcm levels.

Stem-cell niche maintenance in organoids mainly depends on a combination of EGF, Wnt

and Notch signalling (Sato et al., 2011). To identify the pathway that can modulate the

unlicensed $G_1$ state, we systematically treated organoids with a small molecule inhibitor of

EGFR (Gefitinib), a Wnt agonist (Chir99021), and a Notch activator (Valproic acid). Short-

term treatment with Gefitinib, which reduces MAP kinase activity and blocks DNA

replication and cell division (Lynch et al., 2004), immediately caused cells to accumulate in

the unlicensed $G_1$ state, with a 2N DNA content but continuing to express Mcm2 and Ki67

(**Figure 6D, E**). Only prolonged EGFR inhibition (4 days) caused transition to an intermediate

$G_0$ state, with significantly reduced Ki67 expression but with total Mcm2 levels maintained

(**Figure 6D, E**). Both states were reversed by removal of EGFRi and addition of fresh growth

factors (**Figure 6D, E**). Previously it was shown that EGFR increases Lgr5 expression (Basak et

al., 2017), suggesting that these transitional states (Unlicensed $G_1$ and $G_0$) are associated

with 'stemness'. Additionally, EGFRi appeared to potently kill transit-amplifying cells (**S5**

**Figure B, C**), leaving branches containing only stem cells (Basak et al., 2017). This suggests

that both, the fully quiescent $G_0$ and unlicensed $G_1$ states, may provide protection to stem

cells.

Treatment with the Wnt agonist Chir99021 did not appear to significantly affect licensing

dynamics (**S5 Figure C**). Strikingly, treatment with the Valproic acid (a Notch activator) alone

or in combination with Chir99021 significantly altered licensing profiles (**S5 Figure D, F**). The

combination of Chir99021 and Valproic acid induces Lgr5 expression throughout the

organoid epithelium (**S5 Figure E**) (Yin et al., 2014). Our data showed that this is associated

with the appearance of a population of cells with low levels of Ki67 and intermediate levels

of DNA-bound Mcm2 (**S5 Figure D**) similar to the intermediate, shallow $G_0$ state induced by

EGFRi. Surprisingly, we observed an arc of cells connecting this to the fully licensed state,

suggesting that re-licensing of these cells occurs before they express high levels of Ki67 and

that cells can reactivate licensing from a deeper state of $G_0$ directly (**S5 Figure D, F**).

Inhibiting Notch signalling with DAPT, which induces terminal secretory cell differentiation

(van Es et al., 2005), also significantly altered licensing dynamics and induced deep $G_0$, with

reduced Ki67 and loss of Mcm2 proteins (**S5 Figure G**). Together, these data suggest that

both EGFR signalling and Notch signalling can significantly influence the licensing dynamics

during the transition between quiescence and unlicensed $G_1$.

**The Unlicensed-$G_1$ state is lost in *Apc* mutant organoids**

Many established cell lines appear to lack an unlicensed $G_1$ state, instead licensing all origins

immediately upon mitotic exit (Friedrich et al., 2005, Haland et al., 2015, Moreno et al.,

2016, Matson et al., 2017). In addition, many of these cell-lines lack a functional licensing

checkpoint (Shreeram et al., 2002, Feng et al., 2003, Liu et al., 2009, Nevis et al., 2009) that

has been suggested to arrest cells in $G_1$ by inactivation of the RB-E2F restriction point(Liu et

al., 2009, Machida et al., 2005, Nevis et al., 2009, Shreeram et al., 2002, Teer et al., 2006).

To understand the biological relevance of the unlicensed $G_1$ state, we determined whether

origin licensing dynamics during $G_1$ are altered during the initial stages of tumorigenesis.

The first initiating mutations in colorectal cancer are usually in *Adenomatus polyposis coli*

(Apc). Therefore, we investigated whether licensing dynamics are altered in *Apc* mutant

intestinal epithelium. In Apc$^{Min/+}$ epithelium *in vivo*, Mcm2 expression appeared normal in

areas of normal histology but was greatly increased in polyps (**Figure 7A**). In isolated

Apc$^{Min/+}$ crypts, there was a slight increase in the size of the proliferative compartment

similar to previous reports (Trani et al., 2014), and a slight increase in the number of EdU(+)

cells in the stem cell compartment (**Figure 7Bi, iii**). However, in the majority of crypts there

were no significant differences in the number or distribution of licenced cells in the stem cell

compartment (**Figure 7Bi, iii**). However, we found a small number of crypts that were

considerably larger. Strikingly, within these crypts, we noticed small 'ribbons' of EdU(+) cells

that extended significantly into the transit-amplifying compartment (**Figure 7Bii**). We

suspected that these ribbons reflected a clone of cells that had undergone loss of

heterozygosity (LOH), and continually re-engage with the cell cycle. This in turn suggests

that the LOH event that converts *Apc$^{Min/+}$* to *Apc$^{Min/Min}$* cells significantly alters cell-cycle

dynamics.

To determine whether LOH alters origin licensing dynamics by modifying $G_1$, we compared

wild-type and *Apc$^{Min/Min}$* organoids (**Figure 7C**). *Apc$^{Min/Min}$* organoids contained many more

licensed cells that were distributed randomly throughout the organoid (**Figure 7D**),

reminiscent of the altered distribution of Ki67 (+) cells in these organoids (Fatehullah et al.,

2013). Strikingly, we found that licensing dynamics in the $G_1$ phase of $Apc^{Min/Min}$ cells were

significantly different and there was a significant loss of the unlicensed $G_1$ population.

(**Figure 7E**). Instead, there was a significant population of cells that appear to license

immediately upon $G_1$ entry and progressed into S-phase immediately after minimal licensing

(**Figure 7E, F**). Interestingly, we still detected a population of cells that licensed as many

origins as in WT organoids. These may be $Apc^{Min/+}$ cells that had not undergone LOH yet and

thus maintained near-normal cell cycle dynamics.

A functional licensing checkpoint depends on an intact RB restriction point and we tested

whether there the relative amounts of hypo- and hyper-phosphorylated RB varied between

wild type and $Apc^{Min/Min}$ epithelia. In *wild-type* organoids, we noticed that the majority of RB

appeared hypo-phosphorylated (**Figure 7G**). In contrast, in $Apc^{Min/Min}$ organoids, at least half

of RB appeared hyper-phosphorylated, with at least 3-fold more phosphorylated RB.

Together, this data is consistent with the idea that cells exist in an unlicensed $G_1$ state prior

to passage through the restriction point and activation of E2F-driven transcription. In

contrast, *Apc* mutant cells appear to have lost normal restriction point control, so that they

have constitutively hyper-phosphorylated RB, allowing them to completely bypass the

unlicensed $G_1$ state.

**Discussion**

The cell-cycle of intestinal stem and transit-amplifying cells is poorly understood. By

comparing the total and DNA-bound Mcm2 in intact intestinal crypts we provide new

insights into how licensing and cell-cycle commitment are coupled in this tissue. We provide

evidence that after their final mitosis, transit amplifying cells do not license their replication

origins and immediately exit the cell cycle. We show that many Lgr5(+) stem cells spend the

majority of $G_1$ in an unlicensed state, continually expressing Mcm2 and Ki67. In unlicensed-

$G_1$, stem cells could be poised to respond to cues and progress past this restriction point to

resume the cell division cycle.

Lgr5(+) stem cells have a cell-cycle length greater than transit-amplifying cells (Schepers et

al., 2011). The biological relevance of this is currently unknown. The data presented here

suggest a delay in origin licensing is a key feature of the prolonged cell-cycle of Lgr5(+) cells.

Although ~80% of Lgr5(+) cells are thought to be continually proliferative and express high

levels of both Ki67 (Basak et al., 2014) and Mcm2, we found that most Lgr5(+) cells reside in

an unlicensed state, with 2N DNA content and Mcm2 not bound to DNA. Since the licensed

state defines proliferative fate commitment (Blow and Hodgson, 2002), we suggest that

these cells are temporarily paused in $G_1$, continuing to express proliferative makers such as

Ki67 and Mcm2, but without fully committing to the cell cycle (**Figure 8**). We show that the

number of Lgr5(+) cells with DNA bound-Mcm2 was similar to the number of proposed

'active' stem cells determined in lineage tracing experiments (Baker et al., 2014, Kozar et al.,

2013).

Prolonged arrest may eventually result in degradation of MCM2-7 proteins and lead to

induction of a state of deep quiescence ($G_0$). Consistent with this idea, we observed that

LRCs, thought to provide a reserve of quiescent stem cells, did not express Mcm2. The lack

of Mcm2 expression may reflect that a significant period of time has passed since these cells

divided. The delay in activating the licensing system may create a prolonged time-window

for Lgr5(+) cells to receive and interpret environmental cues before deciding to commit to

duplication, offering a means to control their number. It is likely that the majority of Lgr5(+)

cells regularly resume their cell cycle, given their continual expression of proliferation

markers (Basak et al., 2014). The identity and decisions of Lgr5(+) cells are governed by

stochastic choices and the ability to pause briefly in $G_1$ offers them unique flexibility in

making these choices.

As expected, the transition from unlicensed $G_1$ to licensed $G_1$ seems dependent on EGFR

signalling. However, other pathways responsible for stem-cell maintenance can also

significantly cause the appearance of a unique population of unlicensed cells with distinct

cell-cycle dynamics. Growing evidence suggests that intestinal stem cell fate is not governed

by asymmetric segregation of fate determinants (Lopez-Garcia et al., 2010, Snippert et al.,

2010, Steinhauser et al., 2012). Instead, factors operating in the stem cell niche, such as Wnt

and Notch signalling affect stem cell fate decisions and also reduce the cycle rate of

intestinal stem cells (Hirata et al., 2013). This is consistent with the idea that cell fate

choices are affected by decreasing proliferation rate and increased $G_0/G_1$ length. Indeed,

extending $G_1$ in mouse and human embryonic stem cells can drive differentiation (Calder et

al., 2013, Coronado et al., 2013). Similarly, long $G_1$ phases are associated with the

generation of fate-restricted progenitors during neurogenesis (Arai et al., 2011). An

extended time window in the cell-cycle has been suggested to allow niche factors and/or

fate determinants to accumulate to direct progenitor fate (Calegari and Huttner, 2003). In

the case of adult intestinal stem cells, holding cells in $G_1$ may allow an extended time for

stem cell fate factors to act and maintain stem cell fate. In contrast, many embryonic stem

cells licence rapidly and the cell cycle slows throughout differentiation (Matson et al., 2017).

However, like embryonic stem cells (Ge et al., 2015), intestinal stem cells appear to have

licensed more origins than non-stem cells when they enter S phase. This may help ensure

accurate and complete genome duplication in long-lived stem cells (Moreno et al., 2016).

With a higher demand for licensed origins, intestinal stem cells may therefore more readily

engage the licensing checkpoint that ensures that all origins are licensed before cells enter S

phase (Alver et al., 2014, Liu et al., 2009, Shreeram et al., 2002). This additional demand for

licensed origins in stem cells may also explain why crypts hypomorphic for Mcm2 have a

stem-cell deficiency (Pruitt et al., 2007).

It is unclear how intestinal stem cells enter a significant unlicensed $G_1$ state. The simplest

explanation is that licensing factors such as Cdt1 or Cdc6 are not readily available in new-

born stem cells, and their synthesis has to be stimulated by an upstream signal for fate

commitment via activation of E2F-driven transcription. This is the situation after prolonged

quiescence, which is accompanied by passive downregulation of licensing factors (Coller,

2007). In contrast, in continually dividing cells their levels are maintained. Consistent with

this idea, licensing factors such as Cdc6, along with many cyclin-CDK complexes, are down

regulated beyond the end of the TA zone (Frey et al., 2000) (Smartt et al., 2007). Cells

without a functional restriction point, such as *Apc* mutant cells or most cancer cell lines,

could immediately license their origins upon entering $G_1$ and can progress into S-phase

without sufficient origins being licensed. Interestingly, both transit-amplifying cells and

highly proliferative *Apc* mutant cells are highly sensitive to replication inhibitors, such as

Gefitinib (**S5 Figure H**). Wild-type stem cells survive this treatment, potentially by engaging

the licensing checkpoint to reversibly stall in Unlicensed $G_1$. This suggests that the

unlicensed $G_1$ can protect stem cells from replication inhibitors and offers a potentially

selective means to kill highly proliferative cells, such as *Apc* mutant cells (Shreeram et al.,

2002, Blow and Gillespie, 2008).

In summary, we demonstrate that the dynamics of the DNA replication licensing system

provides a new way for measuring the proliferative fate of intestinal stem cells. We suggest

a model for 'working' intestinal stem cells that spend a significant proportion of $G_1$ in a

unlicensed state until a proliferative fate decision is made. Correspondingly, exit from the

cell-cycle in label retaining '+4' cells leads to loss of proliferative capacity and loss of Mcm2

expression causing cells to enter a deeply $G_0$ quiescent state (**Figure 6**). The unlicensed $G_1$

state is lost in *Apc* mutant epithelia, which lack a functional Rb-restriction point. We suggest

that the unlicensed $G_1$ state serves stem cells in controlling their numbers by regulating the

cell-cycle.

**Author contributions**

T.D.C, J.J.B and I.N conceived and designed the study; T.D.C and I.P.N collected the data; Y.C

assisted with organoid experiments; T.D.C performed the data analysis. J.J.B performed the

modelling simulations.  T.D.C, I.N and J.J.B wrote the manuscript.

**Conflicts of Interest**

The authors report no conflicts of interest.

**Acknowledgements**

We would like to thank members of the Näthke and Blow laboratories for general assistance

and helpful discussions; Dr Paul Appleton, Dr Graeme Ball and the Dundee Imaging and

Tissue Imaging Facility for support with microscopy and image analysis. The imaging facility

is funded by the Welcome Trust Technology Platform award (097945/B/11/Z) and Welcome

Trust award (101468/Z/13/Z). We thank Dr Rosemary Clarke and the Dundee Flow

Cytometry Facility for support with flow cytometry, cell sorting and analysis. We also thank

Dr Richard Mort (University of Edinburgh) for Fucci2aR mouse tissue. This work was

supported by a programme grant from Cancer Research UK to I.N (C430/A11243) and to

J.J.B (C303/A14301), Wellcome Trust grant WT096598MA and an MRC studentship award to

T.D.C.

**REFERENCES**

ALVER, R. C., CHADHA, G. S. & BLOW, J. J. 2014. The contribution of dormant origins to
765            genome stability: from cell biology to human genetics. *DNA repair,* 19**,** 182-9.
766    ARAI, Y., PULVERS, J. N., HAFFNER, C., SCHILLING, B., NUSSLEIN, I., CALEGARI, F. & HUTTNER,
767            W. B. 2011. Neural stem and progenitor cells shorten S-phase on commitment to
768            neuron production. *Nature communications,* 2**,** 154.
769    BAKER, A. M., CERESER, B., MELTON, S., FLETCHER, A. G., RODRIGUEZ-JUSTO, M., TADROUS,
770            P. J., HUMPHRIES, A., ELIA, G., MCDONALD, S. A., WRIGHT, N. A., SIMONS, B. D.,
771            JANSEN, M. & GRAHAM, T. A. 2014. Quantification of crypt and stem cell evolution in
772            the normal and neoplastic human colon. *Cell reports,* 8**,** 940-7.
773    BARKER, N., VAN ES, J. H., KUIPERS, J., KUJALA, P., VAN DEN BORN, M., COZIJNSEN, M.,
774            HAEGEBARTH, A., KORVING, J., BEGTHEL, H., PETERS, P. J. & CLEVERS, H. 2007.
775            Identification of stem cells in small intestine and colon by marker gene Lgr5. *Nature,*
776            449**,** 1003-7.
777    BASAK, O., BEUMER, J., WIEBRANDS, K., SENO, H., VAN OUDENAARDEN, A. & CLEVERS, H.
778            2017. Induced Quiescence of Lgr5+ Stem Cells in Intestinal Organoids Enables
779            Differentiation of Hormone-Producing Enteroendocrine Cells. *Cell stem cell,* 20**,** 177-
780            190 e4.
781    BASAK, O., VAN DE BORN, M., KORVING, J., BEUMER, J., VAN DER ELST, S., VAN ES, J. H. &
782            CLEVERS, H. 2014. Mapping early fate determination in Lgr5+ crypt stem cells using a
783            novel Ki67-RFP allele. *The EMBO journal,* 33**,** 2057-68.
784    BLOW, J. J. & GILLESPIE, P. J. 2008. Replication licensing and cancer--a fatal entanglement?
785            *Nature reviews. Cancer,* 8**,** 799-806.
786    BLOW, J. J. & HODGSON, B. 2002. Replication licensing--defining the proliferative state?
787            *Trends in cell biology,* 12**,** 72-8.

BUCZACKI, S. J., ZECCHINI, H. I., NICHOLSON, A. M., RUSSELL, R., VERMEULEN, L., KEMP, R. &
WINTON, D. J. 2013. Intestinal label-retaining cells are secretory precursors
expressing Lgr5. *Nature,* 495**,** 65-9.
CALDER, A., ROTH-ALBIN, I., BHATIA, S., PILQUIL, C., LEE, J. H., BHATIA, M., LEVADOUX-
MARTIN, M., MCNICOL, J., RUSSELL, J., COLLINS, T. & DRAPER, J. S. 2013. Lengthened
G1 phase indicates differentiation status in human embryonic stem cells. *Stem cells*
*and development,* 22**,** 279-95.
CALEGARI, F. & HUTTNER, W. B. 2003. An inhibition of cyclin-dependent kinases that
lengthens, but does not arrest, neuroepithelial cell cycle induces premature
neurogenesis. *Journal of cell science,* 116**,** 4947-55.
CHAMPERIS TSANIRAS, S., KANELLAKIS, N., SYMEONIDOU, I. E., NIKOLOPOULOU, P.,
LYGEROU, Z. & TARAVIRAS, S. 2014. Licensing of DNA replication, cancer,
pluripotency and differentiation: an interlinked world? *Seminars in cell &*
*developmental biology,* 30**,** 174-80.
CHWALINSKI, S. & POTTEN, C. S. 1987. Influence of irradiation or thymidine (TdR) on the
pattern of 3H-TdR incorporation at each cell position in the crypts of the small
intestine of the mouse. *International journal of radiation biology and related studies*
*in physics, chemistry, and medicine,* 51**,** 243-54.
COLLER, H. A. 2007. What's taking so long? S-phase entry from quiescence versus
proliferation. *Nature reviews. Molecular cell biology,* 8**,** 667-70.
CORONADO, D., GODET, M., BOURILLOT, P. Y., TAPPONNIER, Y., BERNAT, A., PETIT, M.,
AFANASSIEFF, M., MARKOSSIAN, S., MALASHICHEVA, A., IACONE, R., ANASTASSIADIS,
K. & SAVATIER, P. 2013. A short G1 phase is an intrinsic determinant of naive
embryonic stem cell pluripotency. *Stem cell research,* 10**,** 118-31.
DALTON, S. 2015. Linking the Cell Cycle to Cell Fate Decisions. *Trends in cell biology,* 25**,** 592-
600.
DIMITROVA, D. S., PROKHOROVA, T. A., BLOW, J. J., TODOROV, I. T. & GILBERT, D. M. 2002.
Mammalian nuclei become licensed for DNA replication during late telophase.
*Journal of cell science,* 115**,** 51-9.
EVRIN, C., CLARKE, P., ZECH, J., LURZ, R., SUN, J., UHLE, S., LI, H., STILLMAN, B. & SPECK, C.
2009. A double-hexameric MCM2-7 complex is loaded onto origin DNA during
licensing of eukaryotic DNA replication. *Proceedings of the National Academy of*
*Sciences of the United States of America,* 106**,** 20240-5.
EWARD, K. L., OBERMANN, E. C., SHREERAM, S., LODDO, M., FANSHAWE, T., WILLIAMS, C.,
JUNG, H. I., PREVOST, A. T., BLOW, J. J., STOEBER, K. & WILLIAMS, G. H. 2004. DNA
replication licensing in somatic and germ cells. *Journal of cell science,* 117**,** 5875-86.
FATEHULLAH, A., APPLETON, P. L. & NATHKE, I. S. 2013. Cell and tissue polarity in the
intestinal tract during tumourigenesis: cells still know the right way up, but tissue
organization is lost. *Philosophical transactions of the Royal Society of London. Series*
*B, Biological sciences,* 368**,** 20130014.
FENG, D., TU, Z., WU, W. & LIANG, C. 2003. Inhibiting the expression of DNA replication-
initiation proteins induces apoptosis in human cancer cells. *Cancer research,* 63**,**
7356-64.
FREY, M. R., CLARK, J. A., LEONTIEVA, O., URONIS, J. M., BLACK, A. R. & BLACK, J. D. 2000.
Protein kinase C signaling mediates a program of cell cycle withdrawal in the
intestinal epithelium. *The Journal of cell biology,* 151**,** 763-78.

FRIEDRICH, T. D., BEDNER, E., DARZYNKIEWICZ, Z. & LEHMAN, J. M. 2005. Distinct patterns
of MCM protein binding in nuclei of S phase and rereplicating SV40-infected monkey
kidney cells. *Cytometry. Part A : the journal of the International Society for Analytical*
*Cytology,* 68**,** 10-8.
GAMBUS, A., KHOUDOLI, G. A., JONES, R. C. & BLOW, J. J. 2011. MCM2-7 form double
hexamers at licensed origins in Xenopus egg extract. *The Journal of biological*
*chemistry,* 286**,** 11855-64.
GE, X. Q., HAN, J., CHENG, E. C., YAMAGUCHI, S., SHIMA, N., THOMAS, J. L. & LIN, H. 2015.
Embryonic Stem Cells License a High Level of Dormant Origins to Protect the
Genome against Replication Stress. *Stem cell reports,* 5**,** 185-94.
GONZALEZ, M. A., TACHIBANA, K. E., LASKEY, R. A. & COLEMAN, N. 2005. Control of DNA
replication and its potential clinical exploitation. *Nature reviews. Cancer,* 5**,** 135-41.
HALAND, T. W., BOYE, E., STOKKE, T., GRALLERT, B. & SYLJUASEN, R. G. 2015. Simultaneous
measurement of passage through the restriction point and MCM loading in single
cells. *Nucleic acids research,* 43**,** e150.
HIRATA, A., UTIKAL, J., YAMASHITA, S., AOKI, H., WATANABE, A., YAMAMOTO, T., OKANO,
H., BARDEESY, N., KUNISADA, T., USHIJIMA, T., HARA, A., JAENISCH, R.,
HOCHEDLINGER, K. & YAMADA, Y. 2013. Dose-dependent roles for canonical Wnt
signalling in de novo crypt formation and cell cycle properties of the colonic
epithelium. *Development,* 140**,** 66-75.
ILVES, I., PETOJEVIC, T., PESAVENTO, J. J. & BOTCHAN, M. R. 2010. Activation of the MCM2-7
helicase by association with Cdc45 and GINS proteins. *Molecular cell,* 37**,** 247-58.
JURIKOVA, M., DANIHEL, L., POLAK, S. & VARGA, I. 2016. Ki67, PCNA, and MCM proteins:
Markers of proliferation in the diagnosis of breast cancer. *Acta histochemica,* 118**,**
544-52.
KINGSBURY, S. R., LODDO, M., FANSHAWE, T., OBERMANN, E. C., PREVOST, A. T., STOEBER,
K. & WILLIAMS, G. H. 2005. Repression of DNA replication licensing in quiescence is
independent of geminin and may define the cell cycle state of progenitor cells.
*Experimental cell research,* 309**,** 56-67.
KOZAR, S., MORRISSEY, E., NICHOLSON, A. M., VAN DER HEIJDEN, M., ZECCHINI, H. I., KEMP,
R., TAVARE, S., VERMEULEN, L. & WINTON, D. J. 2013. Continuous clonal labeling
reveals small numbers of functional stem cells in intestinal crypts and adenomas. *Cell*
*stem cell,* 13**,** 626-33.
KRUDE, T., MUSAHL, C., LASKEY, R. A. & KNIPPERS, R. 1996. Human replication proteins
hCdc21, hCdc46 and P1Mcm3 bind chromatin uniformly before S-phase and are
displaced locally during DNA replication. *Journal of cell science,* 109 ( Pt 2)**,** 309-18.
LEONE, G., DEGREGORI, J., YAN, Z., JAKOI, L., ISHIDA, S., WILLIAMS, R. S. & NEVINS, J. R.
1998. E2F3 activity is regulated during the cell cycle and is required for the induction
of S phase. *Genes & development,* 12**,** 2120-30.
LIU, P., SLATER, D. M., LENBURG, M., NEVIS, K., COOK, J. G. & VAZIRI, C. 2009. Replication
licensing promotes cyclin D1 expression and G1 progression in untransformed
human cells. *Cell cycle,* 8**,** 125-36.
LOPEZ-GARCIA, C., KLEIN, A. M., SIMONS, B. D. & WINTON, D. J. 2010. Intestinal stem cell
replacement follows a pattern of neutral drift. *Science,* 330**,** 822-5.
LYNCH, T. J., BELL, D. W., SORDELLA, R., GURUBHAGAVATULA, S., OKIMOTO, R. A.,
BRANNIGAN, B. W., HARRIS, P. L., HASERLAT, S. M., SUPKO, J. G., HALUSKA, F. G.,

880            LOUIS, D. N., CHRISTIANI, D. C., SETTLEMAN, J. & HABER, D. A. 2004. Activating
881              mutations in the epidermal growth factor receptor underlying responsiveness of
882              non-small-cell lung cancer to gefitinib. *The New England journal of medicine,* 350**,**
883              2129-39.
884 MACHIDA, Y. J., TEER, J. K. & DUTTA, A. 2005. Acute reduction of an origin recognition
885              complex (ORC) subunit in human cells reveals a requirement of ORC for Cdk2
886              activation. *The Journal of biological chemistry,* 280**,** 27624-30.
887 MAKAROVA, K. S., KOONIN, E. V. & KELMAN, Z. 2012. The CMG (CDC45/RecJ, MCM, GINS)
888              complex is a conserved component of the DNA replication system in all archaea and
889              eukaryotes. *Biology direct,* 7**,** 7.
890 MATSON, J. P., DUMITRU, R., CORYELL, P., BAXLEY, R. M., CHEN, W., TWAROSKI, K.,
891              WEBBER, B. R., TOLAR, J., BIELINSKY, A. K., PURVIS, J. E. & COOK, J. G. 2017. Rapid
892              DNA replication origin licensing protects stem cell pluripotency. *eLife,* 6.
893 MORENO, A., CARRINGTON, J. T., ALBERGANTE, L., AL MAMUN, M., HAAGENSEN, E. J.,
894              KOMSELI, E. S., GORGOULIS, V. G., NEWMAN, T. J. & BLOW, J. J. 2016. Unreplicated
895              DNA remaining from unperturbed S phases passes through mitosis for resolution in
896              daughter cells. *Proceedings of the National Academy of Sciences of the United States*
897              *of America,* 113**,** E5757-64.
898 MORT, R. L., FORD, M. J., SAKAUE-SAWANO, A., LINDSTROM, N. O., CASADIO, A., DOUGLAS,
899              A. T., KEIGHREN, M. A., HOHENSTEIN, P., MIYAWAKI, A. & JACKSON, I. J. 2014.
900              Fucci2a: a bicistronic cell cycle reporter that allows Cre mediated tissue specific
901              expression in mice. *Cell cycle,* 13**,** 2681-96.
902 MOYER, S. E., LEWIS, P. W. & BOTCHAN, M. R. 2006. Isolation of the Cdc45/Mcm2-7/GINS
903              (CMG) complex, a candidate for the eukaryotic DNA replication fork helicase.
904              *Proceedings of the National Academy of Sciences of the United States of America,*
905              103**,** 10236-41.
906 MUNOZ, J., STANGE, D. E., SCHEPERS, A. G., VAN DE WETERING, M., KOO, B. K., ITZKOVITZ,
907              S., VOLCKMANN, R., KUNG, K. S., KOSTER, J., RADULESCU, S., MYANT, K., VERSTEEG,
908              R., SANSOM, O. J., VAN ES, J. H., BARKER, N., VAN OUDENAARDEN, A., MOHAMMED,
909              S., HECK, A. J. & CLEVERS, H. 2012. The Lgr5 intestinal stem cell signature: robust
910              expression of proposed quiescent '+4' cell markers. *The EMBO journal,* 31**,** 3079-91.
911 MUSAHL, C., HOLTHOFF, H. P., LESCH, R. & KNIPPERS, R. 1998. Stability of the replicative
912              Mcm3 protein in proliferating and differentiating human cells. *Experimental cell*
913              *research,* 241**,** 260-4.
914 NAMDAR, M. & KEARSEY, S. E. 2006. Analysis of Mcm2-7 chromatin binding during anaphase
915              and in the transition to quiescence in fission yeast. *Experimental cell research,* 312**,**
916              3360-9.
917 NEVIS, K. R., CORDEIRO-STONE, M. & COOK, J. G. 2009. Origin licensing and p53 status
918              regulate Cdk2 activity during G(1). *Cell cycle,* 8**,** 1952-63.
919 OHTANI, K., IWANAGA, R., NAKAMURA, M., IKEDA, M., YABUTA, N., TSURUGA, H. & NOJIMA,
920              H. 1999. Cell growth-regulated expression of mammalian MCM5 and MCM6 genes
921              mediated by the transcription factor E2F. *Oncogene,* 18**,** 2299-309.
922 POTTEN, C. S. & LOEFFLER, M. 1990. Stem cells: attributes, cycles, spirals, pitfalls and
923              uncertainties. Lessons for and from the crypt. *Development,* 110**,** 1001-20.
924 POTTEN, C. S., OWEN, G. & BOOTH, D. 2002. Intestinal stem cells protect their genome by
925              selective segregation of template DNA strands. *Journal of cell science,* 115**,** 2381-8.

PRUITT, S. C., BAILEY, K. J. & FREELAND, A. 2007. Reduced Mcm2 expression results in severe
stem/progenitor cell deficiency and cancer. *Stem cells,* 25**,** 3121-32.
PRUITT, S. C., FREELAND, A. & KUDLA, A. 2010. Cell cycle heterogeneity in the small
intestinal crypt and maintenance of genome integrity. *Stem cells,* 28**,** 1250-9.
REMUS, D., BEURON, F., TOLUN, G., GRIFFITH, J. D., MORRIS, E. P. & DIFFLEY, J. F. 2009.
Concerted loading of Mcm2-7 double hexamers around DNA during DNA replication
origin licensing. *Cell,* 139**,** 719-30.
RHIND, N. & GILBERT, D. M. 2013. DNA replication timing. *Cold Spring Harbor perspectives in*
*biology,* 5**,** a010132.
ROTH, S., FRANKEN, P., SACCHETTI, A., KREMER, A., ANDERSON, K., SANSOM, O. & FODDE,
R. 2012. Paneth cells in intestinal homeostasis and tissue injury. *Plos One,* 7**,** e38965.
SATO, T., VAN ES, J. H., SNIPPERT, H. J., STANGE, D. E., VRIES, R. G., VAN DEN BORN, M.,
BARKER, N., SHROYER, N. F., VAN DE WETERING, M. & CLEVERS, H. 2011. Paneth cells
constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature,* 469**,** 415-8.
SCHEPERS, A. G., VRIES, R., VAN DEN BORN, M., VAN DE WETERING, M. & CLEVERS, H. 2011.
Lgr5 intestinal stem cells have high telomerase activity and randomly segregate their
chromosomes. *The EMBO journal,* 30**,** 1104-9.
SHREERAM, S., SPARKS, A., LANE, D. P. & BLOW, J. J. 2002. Cell type-specific responses of
human cells to inhibition of replication licensing. *Oncogene,* 21**,** 6624-32.
SMARTT, H. J., GUILMEAU, S., NASSER, S. V., NICHOLAS, C., BANCROFT, L., SIMPSON, S. A.,
YEH, N., YANG, W., MARIADASON, J. M., KOFF, A. & AUGENLICHT, L. H. 2007. p27kip1
Regulates cdk2 activity in the proliferating zone of the mouse intestinal epithelium:
potential role in neoplasia. *Gastroenterology,* 133**,** 232-43.
SNIPPERT, H. J., VAN DER FLIER, L. G., SATO, T., VAN ES, J. H., VAN DEN BORN, M., KROON-
VEENBOER, C., BARKER, N., KLEIN, A. M., VAN RHEENEN, J., SIMONS, B. D. &
CLEVERS, H. 2010. Intestinal crypt homeostasis results from neutral competition
between symmetrically dividing Lgr5 stem cells. *Cell,* 143**,** 134-44.
STEINHAUSER, M. L., BAILEY, A. P., SENYO, S. E., GUILLERMIER, C., PERLSTEIN, T. S., GOULD,
A. P., LEE, R. T. & LECHENE, C. P. 2012. Multi-isotope imaging mass spectrometry
quantifies stem cell division and metabolism. *Nature,* 481**,** 516-9.
STOEBER, K., TLSTY, T. D., HAPPERFIELD, L., THOMAS, G. A., ROMANOV, S., BOBROW, L.,
WILLIAMS, E. D. & WILLIAMS, G. H. 2001. DNA replication licensing and human cell
proliferation. *Journal of cell science,* 114**,** 2027-41.
SYMEONIDOU, I. E., KOTSANTIS, P., ROUKOS, V., RAPSOMANIKI, M. A., GRECCO, H. E.,
BASTIAENS, P., TARAVIRAS, S. & LYGEROU, Z. 2013. Multi-step loading of human
minichromosome maintenance proteins in live human cells. *The Journal of biological*
*chemistry,* 288**,** 35852-67.
TEER, J. K., MACHIDA, Y. J., LABIT, H., NOVAC, O., HYRIEN, O., MARHEINEKE, K., ZANNIS-
HADJOPOULOS, M. & DUTTA, A. 2006. Proliferating human cells hypomorphic for
origin recognition complex 2 and pre-replicative complex formation have a defect in
p53 activation and Cdk2 kinase activation. *The Journal of biological chemistry,* 281**,**
6253-60.
TODOROV, I. T., WERNESS, B. A., WANG, H. Q., BUDDHARAJU, L. N., TODOROVA, P. D.,
SLOCUM, H. K., BROOKS, J. S. & HUBERMAN, J. A. 1998. HsMCM2/BM28: a novel
proliferation marker for human tumors and normal tissues. *Laboratory investigation;*
*a journal of technical methods and pathology,* 78**,** 73-8.

TRANI, D., NELSON, S. A., MOON, B. H., SWEDLOW, J. J., WILLIAMS, E. M., STRAWN, S. J.,
APPLETON, P. L., KALLAKURY, B., NATHKE, I. & FORNACE, A. J., JR. 2014. High-energy
particle-induced tumorigenesis throughout the gastrointestinal tract. *Radiation*
*research,* 181**,** 162-71.
VAN DER FLIER, L. G. & CLEVERS, H. 2009. Stem cells, self-renewal, and differentiation in the
intestinal epithelium. *Annual review of physiology,* 71**,** 241-60.
VAN ES, J. H., VAN GIJN, M. E., RICCIO, O., VAN DEN BORN, M., VOOIJS, M., BEGTHEL, H.,
COZIJNSEN, M., ROBINE, S., WINTON, D. J., RADTKE, F. & CLEVERS, H. 2005.
Notch/gamma-secretase inhibition turns proliferative cells in intestinal crypts and
adenomas into goblet cells. *Nature,* 435**,** 959-63.
WILLIAMS, G. H., ROMANOWSKI, P., MORRIS, L., MADINE, M., MILLS, A. D., STOEBER, K.,
MARR, J., LASKEY, R. A. & COLEMAN, N. 1998. Improved cervical smear assessment
using antibodies against proteins that regulate DNA replication. *Proceedings of the*
*National Academy of Sciences of the United States of America,* 95**,** 14932-7.
WILLIAMS, G. H., SWINN, R., PREVOST, A. T., DE CLIVE-LOWE, P., HALSALL, I., GOING, J. J.,
HALES, C. N., STOEBER, K. & MIDDLETON, S. J. 2004. Diagnosis of oesophageal cancer
by detection of minichromosome maintenance 5 protein in gastric aspirates. *British*
*journal of cancer,* 91**,** 714-9.
YIN, X., FARIN, H. F., VAN ES, J. H., CLEVERS, H., LANGER, R. & KARP, J. M. 2014. Niche-
independent high-purity cultures of Lgr5+ intestinal stem cells and their progeny.
*Nature methods,* 11**,** 106-12.

**Figure Legends**

**Figure 1. Mcm2 is expressed ubiquitously along the crypt-villus axis and declines slowly as**
**cells differentiate**

(**A**) Sections of normal human (top panel) and mouse (bottom panel) small-intestine were
stained with Phalloidin (Green) and an antibody against Mcm2 (Red). Scale bars: 200 μm.

(**B**) Mean Mcm2 intensities for segmented nuclei were plotted along the crypt-villus axis for
mouse and human tissue. Location of the crypt and villus domains is indicated.

(**C**) An intestinal crypt stained with Hoechst (Blue), Phalloidin (Green) and an antibody
against Mcm2 (Red). Individual cells in interphase and mitosis (metaphase and cytokinesis)
are outlined by dashed white lines.

(**D**) Maximum intensity projections of whole-mount intestinal tissue revealing intestinal
crypts and villi (left panels). Individual X-Y sections are also shown to reveal the epithelium
(right panels). Tissue was stained with Phalloidin (Green) and Hoechst (Blue) and an
antibody against Mcm2 (Red). The alternating pattern of Mcm2(+) (Green stars) and Mcm2(-
) (Orange stars) in the crypt base is highlighted)

(**E**) Images of Lgr5-GFP stem cells (Green) (top panel) co-stained with a Mcm2 antibody
(Red). The correlation (Pearson's correlation R=0.81, p<0.0001) between mean Mcm2 and
Lgr5-GFP intensities for 69 Lgr5-GFP(+) cells normalised to the maximum intensity for an
individual crypt is shown.

(**F**) Images of UEA(+) Paneth Cells (top panels) co-stained with an Mcm2 antibody (Red) and
UEA (Green). Mean Mcm2 intensity for segmented nuclei of UEA(+) Paneth cells was
compared with interphase cells (Right panels).

(**G**) Mcm2 (Green) and UEA (Red) expression in subsets of UEA(+) cells in crypt and villus
domains. UEA(+) cells at the crypt base represent Paneth cells.

(**H**) Quantification of mean Mcm2 intensity in individual UEA(+) cell populations. UEA(+) cells
in the crypt base (Paneth cells, N=224), in the upper crypt compartment (Crypt, N = 132) and
in the villus compartment (Villus, N = 225) were identified manually, and the nuclear Mcm2
intensity was determined for individual cells (All cells, N = 33,736). There was a significant
difference between UEA(+) cells in the crypt and villus compartments (T test, p<0.0001)

1024

**Figure 2. Visualising Mcm2 licensing in intestinal crypts**

(**A**) Representative bright-field images of extracted and unextracted isolated intestinal
crypts. Scale bar: 90 µm

(**B**) Representative images of isolated crypts stained with antibodies against Mcm2 (Red) or
Ki67 (Purple). Scale bar: 10 µm

(**C**) The Mcm2 labelling index for unextracted and extracted crypts is significantly different
(Mean +/- SEM, N=10 crypts (T test, p<0.0001).

(**D**) Representative intestinal crypts stained with Hoechst (Blue) and antibodies against
Mcm2 (Red) and phospho-histone H3 (pH-H3) (Green).

(**E**) Representative flow cytometry profiles for extracted and unextracted isolated crypt
epithelial cells showing Mcm2 vs. DNA content.

(**E'**) Suggested model of the licensing profile shown in panel E. Deeply quiescent cells, do not
express Mcm2 and have a no detectable Mcm2 signal. Cells expressing soluble Mcm2
(Unlicensed $G_1$) show a similar Mcm2 signal to $G_2$ cells. After a proliferative fate decision has
been made, origins become licensed and cells commit to S-phase entry. Cells enter S-phase
after maximal origin licensing (Active-$G_1$). During S-phase, Mcms are then displaced from
DNA during replication.

(**F**) Representative images of extracted and unextracted intestinal organoids stained with an
antibody against Mcm2 (Red).

(**G**) The Mcm2 labelling index for unextracted and extracted organoids. Data is displayed as
Mean +/- SEM, N = 3 organoids and shows a significant difference (T test, p<0.0001).

**Figure 3. The licensing state defines distinct proliferative zones in intestinal crypts**

(**A**) Representative image of an extracted intestinal crypt isolated after a 1 hour EdU pulse *in*
*vivo* (EdU, Green) and stained with Hoechst (Blue) and antibodies against Mcm2 (Red) and
pH-H3 (White). Co-staining shows distinct cell-cycle phases (bottom panels): Licensed cells
committed in $G_1$ (Mcm2(+), EdU(-)); Early (Mcm2(+), EdU(+)) to Late (Mcm2(-), EdU(+)) S-
phase), and mitotic cells (pH-H3(+)). Negative cells represent deeply quiescent ($G_0$),
terminally differentiated cells or cells in $G_1$, which have not made a proliferative fate
decision, remaining unlicensed. The crypt base is to the left of the displayed image.

(**B**) Nuclear volume was estimated in cells at the distinct cell-cycle phases identified
previously: Negative ($G_0$/$G_1$/$G_2$: N =115), $G_1$ Licensed (Mcm(+), EdU(-): N=38), S-phase
(Mcm(+), EdU(+): N=24) and Late-S/$G_2$ (Mcm(-), EdU(+): N=26). Top Panels show
representative examples of each cell-cycle phase and the associated 3D rendered nuclei.
There was a significant difference in the size of Licensed $G_1$, S and Late-S/$G_2$ nuclei (T test,
p<0.0001).

(**C**) Representative images of intestinal crypts isolated after a 1 hour EdU pulse (Green) *in*
*vivo*. Displayed are 3D projections of extracted and unextracted crypts stained with Hoechst
(Blue) and an antibody against Mcm2 (Red). The crypt base is to the left of the displayed
image.

(**D**) Comparison between cells expressing Mcm2 protein and DNA-bound Mcm2 along the
crypt-villus axis between unextracted (N=101 crypts) and extracted (N=109 crypts) (taken
from 3 mice). Data is displayed as the mean % of cells per set distance bin.

(**E**) All cells were divided into 4 distinct groups based on Mcm2 and EdU intensities. These
groups represent distinctive cell-cycle phases as defined by their total (unextracted N=101
crypts) or licensed (extracted N=109 crypts) Mcm2 content: **Extracted:** 1) **Unlicensed**
(Mcm2(-), EdU(-)), 2) **$G_1$ licensed** (Mcm2(+), EdU(-)), 3) **Early/Mid S-phase** (Mcm2(+),
EdU(+)) and 4) **Late-S/$G_2$** (Mcm2(-), EdU(+)). The data is represented as the population mean
of the total cells per distance bin, Mean +/- SEM.
(**F**) The distance of the most distal Mcm2(+) and EdU(+) cells to the crypt base was
compared in extracted and unextracted crypts. Data was scored manually for 10
representative crypts per condition. Licensed cells (Mcm2(+)) were significantly closer to the
crypt base than EdU(+) cells (T test, p=0.0015. Cells expressing Mcm2 protein extended
significantly above the last EdU(+) cell (T test, p<0.0003)

(**G**) Representative images of crypts isolated 17 hours after administration of EdU (Green).
3D projections of extracted and unextracted crypts stained with Hoechst (Blue) and an
antibody against Mcm2 (Red) are shown.

(**H**) Cells were divided into 4 distinct groups as in Panel **E** (N=51 Crypts).

1082

**Figure 4. Intestinal stem cells reside in a paused unlicensed G$_1$**

(**A**) Quantification of cell cycle stages for cells in the stem cell (SC) compartment (<40 μm
from the crypt base) and in the early transit-amplifying (TA) compartment (40-80 μm from
the crypt base), as described in Figure 3E (**N=49 crypts**).

(**B**) Representative image of an extracted crypt base isolated 1 hour after a pulse of EdU
(Green) and stained with Hoechst (Blue), UEA (Red) and antibodies against Mcm2 (White).
Nuclear morphology and UEA signal were used to distinguish between UEA(+) Paneth Cells
(outlined by dashed lines and nuclei marked with blue stars) and UEA(-) stem cells (situated
between outlined Paneth cells and with nuclei marked by red stars).

(**C**) The average % of UEA(-) stem cells that fall into the previously defined cell-cycle bins:
(Unlicensed; G$_1$ licensed; S-phase; S/G$_2$ phase; N=68 crypts).

(**D**) Representative flow cytometry profiles of (**i**) isolated Lgr5$^{Hi}$ intestinal stem cells from 3
mice, showing DNA-bound Mcm2 and DNA content. The 2N (G$_1$) cells of the same
population are also shown (ii). The respective gated populations for Unlicensed G$_1$ (Black),
Licensed (Red), S-phase (Yellow) and G$_2$/M (Green) are shown. The Ki67 content of the
Licensed and Unlicensed G$_1$ cells are also shown (iii).

(**E**) The frequency distribution of mean DNA-bound Mcm2 intensities for Mcm2(+) cells in G$_1$
cells.

(**F**) Simulated ergodic rate analysis of origin licensing during G$_1$. We simulated origin
licensing dynamics during G$_1$, varying the licensing rate and incorporating a significant
paused period (Unlicensed G$_1$). The displayed histograms show the frequency distribution of
DNA-bound Mcm2 of G$_1$ cells. Please refer to materials and methods for further
information. However, an analogy to explain the model is as follows: Origin licensing
through G$_1$ can be thought of in terms of cars travelling along a long stretch of motorway: if
cars enter the motorway at a fixed rate (i.e. unsynchronised cell cycles, cells entering G1)
the density of cars at any one point is inversely proportional to the speed they are travelling
at ('speed' in our case means the rate of licensing). We have added the concept of delays at
the start and end, like tollbooths. In our model, there is a minimum drive time (the
minimum length of G$_1$ required for cells to grow to a critical size before they enter S phase
(arbitrarily set at '100%'). If the speed of the cars means they don't reach the end of the
motorway before this time is up, cars maintain a constant speed along the entire road, and
exit the motorway as soon as they reach the end. If cars drive faster and they reach the end
before this time is up (>100%), they have to wait at the end of the motorway till the critical
time is expired, resulting in a peak accumulation at the fully licensed point. Unlicensed G$_1$ is
an enforced time that cars have to wait once they enter the motorway before they are
allowed to drive along it. This creates a peak on the left (unlicensed cells). If the cars then
drive slowly, when they get to the end of the motorway, they exit immediately because the
critical time has expired (centre panels), **but** if they drive fast enough they reach the end of
the motorway before the critical time has expired. This results in a peak accumulation at
both minimally and maximally licensed points.

(**G**) Comparison of DNA-bound Mcm2 content of Lgr5(+) and Lgr5(-) $G_1$ cells and of cells in
very early S-phase.

**Figure 5. Label retaining cells are in a deep-$G_0$ state**

(**A**) Labelling strategy. H2B-GFP expression was induced in all intestinal epithelial cells in
H2B-GFP mice by administration of doxycycline for 7 days. After complete labelling,
doxycycline was removed and mice rested for 7 days. During this chase period, the majority
of H2B-GFP(+) cells are lost by label dilution due to cell division and upward migration. Both
Paneth cells and +4 Label retaining cells are H2B-GFP(+) after the chase period. +4 Label
retaining cells were distinguished from Paneth cells on the lack of UEA staining (Buczacki et
al., 2013).

(**B**) Representative images of whole-mount sections of H2B-GFP expressing small-intestinal
tissue after a 7-day labelling period (i). A vibratome section (ii) of intestinal tissue after a
subsequent 7-day chase period, stained with Hoechst (nuclei), UEA (Grey), and an antibody
against Mcm2 (Red). An enlarged image of the marked crypt is shown. A UEA(-) label
retaining cell (LRC) is marked by the white arrow). Quantification of the mean GFP intensity
for all cells along the crypt-villus axis after the 7 days chase period is shown (iii).

(**C**) A representative image (i) of a label-retaining cell in an intestinal crypts stained with
Hoechst (Blue), UEA (White) and an antibody against Mcm2 (Red). The label retaining cell
(White arrow) does not express Mcm2. The quantification of Mcm2 expression in label
retaining cells (N=12), Paneth cells (N= 116) and all cells (N=543) is shown (iii).

(**D**) Representative images of extracted H2B-GFP crypts after the 7-Day chase period. H2B-
GFP$^{Hi}$ cells (bright Green) represent Paneth cells and +4 cells (i), and H2B-GFP$^{Low}$ cells (Faint
Green) represent daughter cells (white arrow) that have diluted H2B-GFP content as a result
of cell division (ii). The quantification of DNA-bound Mcm2 in GFP$^{Hi}$ label retaining (LR) and
GFP$^{low}$ daughter cells compared with the total cell population (all cells) is shown (iii).

**Figure 6. Licensing dynamics in intestinal organoids**

(**A**) Representative image of an unextracted and extracted intestinal organoid stained with
Hoechst (Blue), and an antibody against Mcm2 (Red) after a 1 hour EdU pulse (green).

(**B**) Representative flow cytometry profiles from isolated organoids cells showing Ki67 (i) or
DNA-bound Mcm2 (ii) content vs DNA content.

(**C**) Representative flow cytometry profile from cells isolated from cultured organoids (i) or
intestinal crypts (ii) showing DNA-bound Mcm2 vs Ki67 (**See also S5 Figure A**). Ki67 loss and
subsequent loss of Mcm2 during differentiation is apparent, starting from Unlicensed $G_1$
(dashed arrow). Quantification of discrete populations (gates shown in i) was performed: 1)
$G_0$, 2) Transition ($G_0 \leftrightarrow$ Unlicensed $G_1$), 3) Unlicensed $G_1$, 4) Licensed $G_1$, 5) S-Phase, 6) Late-
S/$G_2$/M. (N=3).

(**D**) Representative flow cytometry profiles of isolated organoid epithelial cells grown in ENR
(control) and treated with the EGFR inhibitor Gefitinib, for the indicated times (**i-iii**). After 4
Days in Gefitinib, organoids were reactivated by removal of the Gefitinib and re-addition of
fresh growth factors (ENR) for 2 days (**iv**). Displayed are profiles comparing DNA-bound
Mcm2 vs Ki67 (Top panels) or total Mcm2 content (Bottom panels).

(**E**) The $G_1$ cell populations as shown in (D) were quantified. DNA-bound Mcm2 profiles (i)
show 1) Unlicensed $G_1$, 2) Licensed $G_1$, 3) Transition ($G_0 \leftrightarrow$ Unlicensed $G_1$), 4) $G_0$ populations.
Total-Mcm2 profiles (ii) show Mcm2 expressing (Mcm2(+)) and non-expressing (Mcm2(-))
Cells. (N=3 organoids)
1168

**Figure 7. Unlicensed G$_1$ is lost in *Apc* mutant epithelium**

(**A**) A representative vibratome section of *Apc$^{Min/+}$* intestinal epithelium stained with
Hoechst (nuclei), Phalloidin and an antibody against Mcm2. Regions of normal histology and
a region containing a polyp are highlighted.

(**B**) Representative image of an extracted *Apc$^{Min/+}$* isolated crypt (i) stained with Hoechst
(Nuclei) and an antibody against Mcm2 (Red), after a 1 hour EdU pulse (Green). A
representative image of an abnormally elongated crypt is displayed (ii) showing a clonal
ribbon of cells in the distal transit-amplifying compartment which are EdU(+). Quantification
of cell-cycle stages across the crypt axis (Figure 3) is shown (iii).

(**C**) Representative bright field images of organoids cultured from wild-type and organoids
from Apc$^{Min/+}$ mice that have undergone loss of heterozygosity (Apc$^{Min/Min}$).

(**D**) Representative images of unextracted and extracted *Apc$^{Min/Min}$* organoids stained with
Hoechst (nuclei), an antibody against Mcm2 (red), after a 1 hour EdU pulse.

(**E**) Representative flow cytometry profiles from cells from extracted wild-type and *Apc$^{Min/Min}$*
organoids showing DNA-bound Mcm2 vs DNA content (i). The 2N G$_1$ cells of the shown
profile are displayed (ii) showing Unlicensed G$_1$, Early G$_1$ and Peak (fully licensed) G$_1$
populations.

(**F**) Quantification of the populations described in Panel Eii. Data is displayed as mean +/-
SEM, N=3 organoids.

(**G**) Western blot of RB and pRB (low and high exposures) levels between wild-type and
*Apc$^{Min/Min}$* organoids. Two bands corresponding to hypo- and hyper-phospohorylated RB can
be observed.

(**H**) Quantification of the fold increase of pRB:RB for Apc$^{Min/Min}$ organoids (N=3) and polps
isolated from Apc$^{Min/+}$ mice (N=6), compared to *wild-type* organoids and tissue.

1193

**Figure 8. Model of Origin licensing dynamics in intestinal epithelial cells**

In a normal cell-cycle, Mcms are expressed ubiquitously in all stages. The licensing of DNA
with MCM2-7 occurs in late M and throughout $G_1$, when a cell receives a stimulus to commit
to the cell cycle. As DNA is replicated during S-phase, MCM2-7s are displaced from DNA and
are prevented from relicensing in $G_2$. During terminal differentiation, MCM2-7 are not
actively transcribed and the proteins are gradually lost in post-mitotic cells. However, after
the final mitotic division, cells make a binary decision never to license their DNA, even
though the protein is still present. Mcm proteins then degrade slowly, where cells enter a
terminally differentiated state (deep $G_0$). Alternatively, cells can exit mitosis, not relicense
their DNA but maintain proliferative markers and disengage from the cell cycle for some
time (Unlicensed $G_1$). Two major classes of intestinal stem cells exist: 'Active' stem cells,
engaged with the cell-cycle, and reserve, quiescent Label Retaining Cells. Label retaining
cells are in a state of 'deep' quiescence, and do not contain MCM2-7 because they have
disengaged from the cell cycle for some time. In this study, we show that most 'Active'
Lgr5(+) stem cells reside in an unlicensed state, but contain MCM2-7 proteins. These cells
reside in an 'Unlicensed-$G_1$' until they make a proliferative fate decision, enter the cell-cycle
and license. This provides an explanation for the elongated cell-cycle of Intestinal stem cells:
They reside in a partial resting state where they may be able to respond to niche cues to
divide. This therefore may constitute a unique mechanism to control stem cell numbers.

**Supplementary Figure Legends**

**S1 Figure. Image analysis**

(**A**) Image analysis work-flow.

(**B**) Representative image of an extracted isolated crypt 1 hour after an EdU pulse (Green)
stained with Hoechst (Blue) and an antibody against Mcm2 (Red).

(**C**) Detection of nuclei in the crypt in panel A in 3D. Nuclei were detected in 3D using
segmentation tools in Imaris. Detection was validated visually or each individual crypt.

(**D**) Segmentation of the region of interest defined by nuclei detection in panel B.

(**E**) A distance transform was performed in Imaris to measure the distance of each nucleus
to a reference nucleus at the crypt base. Visual representations of distances divided into
different bins are displayed (Green, 0-10μm; Yellow, 10-20 μm; Red, 20-30μm; Blue, 30-
40μm; Magenta, 40-50μm).

(**F**) Representative 3D quantification of the crypt in panel A shows the distance from the
crypt base (X-axis), DNA-bound Mcm2 (Y-axis) and EdU incorporation (Z-axis).

(**G**) An overview of the gating strategy used for flow cytometry experiments. Intestinal
epithelial cells were separated from debris on the basis of forward and side scatter. Single
cells were subsequently distinguished from doublets based on pulse-width gating. Negative
gates were set on control samples stained with secondary antibody alone. Unlicensed $G_1$
(Population 1) and licensed-$G_1$ (Population 2) were distinguished based on DNA-content and
based on the threshold set by the negative control samples. S-phase cells (Population 3) can
be seen as an arc of cells emanating from the fully licensed cell population, increasing in
DNA content whilst losing DNA-bound Mcm2. $G_2$/M cells (Population 4) can be seen as a
population with 4N DNA content and lacking DNA-bound Mcm2.

**S2 Figure. Clonal cell-cycle patterns in the intestinal epithelium**

(**A**) Representative section through an extracted crypt after a 1 hour EdU pulse (Green) and
stained with Hoechst (Blue) and an antibody against Mcm2 (Red). Discrimination of cell-
cycle staging using DNA-bound Mcm2 and EdU incorporation patterns allows visualisation of
clonal cell-cycle field effects revealing many neighbouring cells with similar DNA-bound
Mcm2 and DNA replication patterns. These clones may represent lineages of from single
cells that progress through the cell cycle at the same rate.

(**B**) Representative image of an isolated crypt in which surface rendering was performed on
all nuclei and colour codes applied to reflect cell-cycle stage. Representative cell-cycle
distributions for isolated Ki67(+), Lgr5(+) and Lgr5(-) intestinal epithelial cells are shown.

(**C**) Representative cell-cycle distributions for isolated Ki67(+), Lgr5(+) and Lgr5(-) intestinal
epithelial cells. The average of each cell-cycle phase is displayed for duplicate isolations.

(**D**) Nuclear volumes were rendered for individual nuclei in whole intestinal crypts isolated 1
hour after labelling with EdU. Image shows nuclei (Blue), EdU (Green) and licensed Mcm2
(Red). Maximum intensity projections of the original image are displayed at the top and
corresponding rendered nuclei at the bottom. Nuclear surfaces were colour-coded
according to cell-cycle states: Blue, unlicensed; Red, Licensed $G_1$; Yellow, S-phase; Green,
Late-S/$G_2$. Nuclear volumes were measured for all nuclei in representative crypts (N=3).
Unlicensed (N=368); $G_1$ (N=104); S-phase (N=41); Late-S/$G_2$ (N=70) and the distance of cells
from the crypt base were binned into 10µm intervals. Known parameters of the nuclear
volume for known cell-cycle stages (Figure 3B) are overlaid.

1257

**S3 Figure. Ki67 expression along the crypt-villus axis**

(**A**) A representative isolated crypt 1 hour after labelling with EdU (Green) and stained with
an antibody against Ki67 (magenta)

(**B**) Quantification of the distribution of Ki67(+) cells along the crypt axis. Cells were binned
into four groups: Negative (Ki67(-), EdU(-)); Ki67(+), EdU(-); Ki67(+), EdU(+) and Ki67(-),
(EdU(+) and by their distance from the crypt base. Data is displayed as the average
percentage of a particular cell subtype, per distance bin. Data is displayed as Mean +/- SEM.
Data from 75 crypts is displayed, N = 14,264 cells.

1266

**S4 Figure. Delayed accumulation of Fucci2aR reporter hCdt1(30-120) in intestinal stem**
**cells**

(**A**) Representative image of a vibratome section of intestinal tissue derived from Fucci2aR
mice, stained with Hoechst (Nuclei), an antibody against Mcm2 and showing expression of
the $G_1$ marker hCdt1(30-120) and S/$G_2$/M marker hGeminin(1/110).

(**B**) Representative images hCdt1(30-120) expression in the crypt base (left panels) and villi
(right panels). Most cells in the crypt base that highly express hCdt1(30-120) are UEA(+)
Paneth cells.

(**C**) A representative image of a Fucci2aR intestinal crypt stained with Hoecsht and and
antibody against Mcm2. All Mcm2(-) Paneth cells (PC) in the crypt base express high levels
of hCdt1(30-120). Most Mcm2(+) stem cells in the crypt base express very low levels of
hCdt1(30-120) (hCdt1$^{low}$). Only few Mcm2(+) cells were found to express high levels of
hCdt1(30-120) (hCdt1$^{High}$). Many cells in the transit-amplifying compartment expressed
hGeminin(1/110).

(**D**) Quantification of the expression of Mcm2, hCdt1(30/120) and hGeminin(1/110) in
Mcm2(+) and Mcm2(-) cells in the stem cell and transit-amplifying compartment. Data is
displayed as a ratio of signal to the background signal detected in negative cells (Stem cell
compartment: Mcm2(+) N=220, Mcm2(-) N=112; Transit-amplifying compartment: Mcm2(+)
N=383, Mcm2(-) N=18).

1286

**S5 Figure. Manipulation of the stem cell niche can artificially induce Unlicensed-G$_1$**

(**A**) Representative flow cytometry profile of extracted epithelial cells isolated from
organoids in ENR media. The displayed image is the same profile displayed in Figure 5C. The
populations in boxed regions 1-7 are overlaid onto the individual Mcm2 and Ki67 cell-cycle
profiles plotted against DNA-content (right hand box).

(**B**) Representative bright-field images of organoids in normal (ENR) media, treated with
Gefitinib for 4 days, and after 2 days in fresh ENR media post Gefitinib treatment.

(**C**) Representative flow cytometry profiles from extracted cells isolated from organoids
treated with 10μM Chir99021 for the indicated time intervals, and after 2 days in fresh ENR
media post Chir99021 treatment.

(**D**) Representative flow cytometry profiles from extracted cells isolated from organoids
treated with normal media (ENR) or in normal media containing Chir99021 and valproic acid
(ENR-CV). Displayed is the comparison of DNA-bound Mcm2 vs Ki67 content (i). The gated
populations: 1) Transition (Unlicensed-G$_1$↔ intermediate G$_0$), 2) Ki67 independent origin
licensing pathway,  3) Fully licensed G$_1$ cells, 4) Normal Licensing, 5) Unlicensed G$_1$ have
been quantified (N=3).

(**E**) Representative images of Lgr5-GFP organoids treated with ENR or ENR-CV. In ENR-CV
treated organoids, the majority of cells express Lgr5.

(**F**) Representative flow cytometry profiles from extracted cells isolated from organoids
treated with Valproic acid for indicated time intervals, and after 2 days in fresh ENR media
post Valproic acid treatment.
1308
1309 (**G**) Representative flow cytometry profiles from extracted cells isolated from organoids
1310 treated with DAPT for indicated time intervals, and after 2 days in fresh ENR media post
1311 DAPT treatment.

1312 (**H**) Representative images of *wild-type* and *Apc$^{Min/Min}$* intestinal organoids after treatment
1313 with 5μM Gefitinib for 3 days.
1314
(**I**) Model for the unique cell-cycle characteristics of organoid epithelial cells.
Normal, highly proliferative cells, express Ki67 and Mcm2 protein that is not DNA-bound (**1**).
During a normal cell-cycle, cells are activated from Unlicensed-G$_1$, and rapidly license origins
(**2**). Mcms are subsequently displaced during DNA replication (**3**) and remain unlicensed
through G$_2$ (**3**). Inhibiting EGFR causes highly proliferative cells (Ki67$^{hi}$) to arrest in
Unlicensed-G$_1$ with maintained Mcm2 protein expression. Prolonged EGFRi treatment
causes transition into an intermediate state of G$_0$ accompanied by loss of Ki67 expression
(Ki67$^{lo}$), but maintenance of MCM2-7 protein expression (**4**). Induction of terminal
differentiation by inhibition of Notch signalling is associated with a terminal loss of MCM2-7
proteins, and entry into deep-G$_0$ (**5**). Notch activation (or HDAC inhibition (HDACi)) induces a
unique subset of Ki67$^{lo}$ Unlicensed-G$_1$ cells to license origins independently of Ki67 **6**). We
suggest that the unique cell population observed upon ENR-CV / ENR-V treatment may be a
reserve subset of stem cells that express Lgr5 and start expressing MCM2-7 and enter

Unlicensed-$G_1$ from deep-$G_0$. These cells have unique cell cycle characteristics, and can
immediately license origins independently of Ki67 expression ($6\rightarrow2\rightarrow3$).

1330

1331    **S1 Movie. Cell-cycle clones**

1332

1333    A 3D rotation of a representative intestinal crypt showing nuclei (Blue), DNA-bound Mcm2
1334    (Red) and EdU (Green; 1 hour pulse). The crypt base is at the bottom of the image. An
1335    isosurface rendering of nuclei within the crypt has also been performed, and have been
1336    coloured to match specific cell-cycle stage: Unlicensed (Blue), $G_1$ Licensed (Red), Early S-
1337    phase (Yellow), Late S/$G_2$ (Green).

1338

Carroll et al., Figure 1

A **Un-extracted** **Extracted** Bright-field

B **Total** **Licensed** Mcm2 Ki67

C Labeling Index (%) Total Mcm2 Licensed Mcm2

E' DNA-bound Mcm2 — Active G₁ — Licensing — S-phase — Mcm('+') — Unlicensed G₁ — G₂ — Mcm('-') — DNA Content

D **Un-extracted** Total Mcm2 **Extracted** Licensed Mcm2 Hoechst Mcm2 pH-H3

E **Unextracted** 97.7% 2.3% **Extracted** 36.3% 64.3%

F **Un-extracted** Total Mcm2 **Extracted** Licensed Mcm2 Mcm2

G Organoids Labeling Index (%) Total Mcm2 Licensed Mcm2

Carroll et al., Figure 2

Carroll et al., Figure 3

Carroll et al., Figure 4

Carroll et al., Figure 5

Carroll et al., Figure 6

Carroll et al., Figure 7

Carroll et al., Figure 8

Carroll et al., S1 Figure

Carroll et al., S2 Figure

# 1 Hour pulse

Ki67
EdU

A

B

Ki67(-)EdU(-) — Ki67(+)EdU(-)
Ki67(+)EdU(+) — Ki67(-)EdU(+)

Average % of Cells in Distance Bin

Distance from crypt base (µm)

A    **Nuclei hCdt1(30/120) hGem(1/110) Mcm2**

B   Crypts                       Villi

**Nuclei hCdt1(30/120) Lysozyme**        **hCdt1(30/120)**

C

**Nuclei hCdt1(30/120) hGem(1/110) Mcm2**

D

**Mcm2(+)**                   **Mcm2(−)**

Fold increase over background

Stem Cell    Transit-amplifying       Stem Cell    Transit-amplifying

**Compartment**                 **Compartment**

**Mcm2**
**hCdt1(30/120)**
**hGeminin(1/110)**

Carroll et al., S5 Figure