Decoupling the roles of cell shape and mechanical stress in orienting and cueing epithelial mitosis

Alexander Nestor-Bergmann[1,2,3][*][§], Georgina A. Stooke-Vaughan[1, 4][*], Georgina K. Goddard[1][*], Tobias Starborg[1], Oliver E. Jensen[2], Sarah Woolner[1][§]

Affiliations:

1: Wellcome Trust Centre for Cell-Matrix Research, Division of Developmental Biology and Medicine, School of Medical Sciences, Faculty of Biology, Medicine & Health, Manchester Academic Health Science Centre, University of Manchester, Oxford Road, Manchester M13 9PT UK

2: School of Mathematics, University of Manchester, Manchester M13 9PL, UK

3: Current address: Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge, CB2 3DY

4. Current address: Department of Mechanical Engineering, University of California, Santa Barbara, Santa Barbara, California, United States of America

* These authors contributed equally to the work

§ Corresponding authors: an529@cam.ac.uk; sarah.woolner@manchester.ac.uk (lead contact)

Abstract

Distinct mechanisms involving cell shape and mechanical force are known to influence the rate and orientation of division in cultured cells. However, uncoupling the impact of shape and force in tissues remains challenging. Combining stretching of *Xenopus laevis* tissue with a novel method of inferring relative mechanical stress, we find separate roles for cell shape in orientating division and mechanical stress in cueing division. We demonstrate that division orientation is best predicted by an axis of cell shape defined by the position of tricellular junctions (TCJs), which aligns exactly with the principal axis of local cell stress rather than the tissue-level stress. The alignment of division to cell shape requires functional cadherin and the localisation of the spindle orientation protein, LGN, to TCJs, but is not sensitive to relative cell stress magnitude. In contrast, cell proliferation rate is more directly regulated by mechanical stress, being correlated with relative isotropic stress, and can be decoupled from cell shape when myosin II is depleted.

Keywords: cell division, mitosis, mitotic spindle, cell proliferation, vertex model, mechanical stress, cell shape.

INTRODUCTION

Cell division orientation and timing must be carefully regulated in order to shape tissues and determine cell fate, preventing defective embryonic development and diseases such as cancer (Mishra and Chan, 2014; Pease and Tirnauer, 2011; Quyn et al., 2010). Recent work has shown that mechanical cues from the extracellular environment can influence cell division rate (Benham-Pyle et al., 2015; Streichan et al., 2014a) and orientation (Campinho et al., 2013; Legoff et al., 2013; Mao et al., 2011; Fink et al., 2011). What remains unclear is whether dividing cells are directly sensing mechanical forces or are responding to changes in cell shape induced by these forces. This distinction is crucial as the molecular mechanisms involved in either shape- or force-sensing could be very different (Luo et al., 2013; Nestor-Bergmann et al., 2014).

Several mechanisms of division orientation control have been postulated in single cells, with evidence for both shape- and stress-sensing (Fink et al., 2011; Minc et al., 2011; Minc and Piel, 2012; Théry et al., 2006). There is limited understanding of how these models could apply to tissues, where cells are linked together by adhesions and it is far more difficult to exclusively manipulate either cell shape or mechanical stress. Recent evidence for a shape-sensing mechanism was found in the *Drosophila* pupal notum. The spindle orientation protein, Mud (*Drosophila* orthologue of NuMA), localises at tricellular junctions (TCJs), recruiting force generators to orient astral microtubules in rounding mitotic cells (Bosveld et al., 2016). However, this mechanism has yet to be demonstrated in another system or related to mechanical stress. In contrast, recent work in a stretched monolayer of MDCK cells has indicated that division orientation may be mediated by a tension-sensing mechanism requiring E-cadherin, although an additional role for cell shape sensing could not be excluded (Hart et al., 2017). Indeed, divisions in MDCK cells have also been found to align better with cell shape than a global stretch axis, though local cell stress was not known in this case (Wyatt et al., 2015).

Separating the roles of shape and stress in tissues will inevitably require an understanding of how force is distributed through heterogeneous cell layers. Experimental methods of assessing stress include laser ablation, atomic force microscopy and micro-aspiration (Campinho et al., 2013; Davidson et al., 2009; Hoh and Schoenenberger, 1994; Hutson et al., 2003). Whilst informative, these techniques are invasive, perturbing the stress field through the measurement, and usually require constitutive modelling for the measurement to be interpreted (Stooke-Vaughan et al., 2017; Sugimura et al., 2016). However, mathematical modelling combined with high quality fluorescence imaging now provides the possibility of non-invasively inferring mechanical stress in tissues (Brodland et al., 2014; Chiou et al., 2012; Feroze et al., 2015; Ishihara and Sugimura, 2012; Nestor-Bergmann et al., 2017; Xu et al., 2015).

In this article, we apply a reproducible strain to embryonic *Xenopus laevis* tissue to
investigate the roles of shape and stress in cell division in a multi-layered tissue. We
particularly focus on mathematically characterising local (cell-level) and global (tissue-level)
stress and the relation to cell shape and division. Our data suggest that mechanical stress is
not directly sensed for orienting the mitotic spindle, acting only to deform cell shape, but is
more actively read as a cue for mitosis.
87
88 **RESULTS**
89
Application of tensile force to a multi-layered embryonic tissue
To investigate the relationship between force, cell shape and cell division in a complex tissue,
we developed a novel system to apply reproducible mechanical strain to a multi-layered
embryonic tissue. Animal cap tissue was dissected from Stage 10 *Xenopus laevis* embryos
and cultured on a fibronectin-coated elastomeric PDMS substrate (Figure 1A). A uniaxial
stretch was applied to the PDMS substrate using an automated stretch device (Figure 1A),
and imaged using standard microscopy. The three-dimensional structure of the stretched
tissue (assessed using 3View EM) could be seen to comprise of approximately three cell
layers (Figure 1B), as would be expected in a stage 10 *Xenopus laevis* embryo (Keller, 1980;
Keller and Schoenwolf, 1977), therefore maintaining the multi-layered tissue structure present
in vivo.
101
Stretching elongates cell shape and reorients divisions.
A 35% stretch of the PDMS substrate led to a 19.67 ± 1.91% (95% confidence interval)
elongation of apical cells in the animal cap along the stretch axis (measured change in length
of 1-dimensional lines drawn on opposite sides of the animal cap; displacement field shown in
Figure 1C). The difference in elongation between the substrate and apical cells is presumably
a result of the mechanical stress being dissipated through multiple cell layers. The qualitative
change in cell shape was not as substantial as was previously observed in stretched
monolayers (Wyatt et al., 2015) (Figure 1D).
110
We mathematically characterised shape using two parameters: orientation of the principal
axis of cell shape relative to the stretch axis (0º), θ_A, and cell circularity, C_A (derived in Section
1 of the Supplementary Document). C_A describes the degree of elongation of a cell (ranging
from 0 being a straight line to 1 being a perfect circle) and θ_A indicates the principal direction
in which this occurs. Stretching oriented the majority of cells with the direction of stretch
(Figure 1E) and caused a highly reproducible elongation of cell shape (Figure 1F). However,
when the substrate was held fixed following stretch, cell elongation reduced over time and
returned close to the unstretched shape profile after 90 minutes (95% confidence intervals of
stretched animal caps at t = 90 minutes overlap with unstretched caps; Figure 1F). Therefore,

cells in this tissue adapt to the elongation caused by stretching and do not behave like a
purely elastic material.

In unstretched tissue, division orientation, θ_D, was not significantly different from a uniform
distribution (p = 0.36, Kolmogorov-Smirnov test; Figure 1G). In contrast, divisions in the
stretched tissue were significantly oriented along the axis of stretch, (p < 1.43×10^{-9},
Kolmogorov-Smirnov test; Figure 1G), with 52% of divisions oriented within 30º of the stretch
axis (compared to 36% in unstretched).

Shape-based models of division differ significantly depending on the cellular
characteristics used to define shape
A shape-based 'long axis' division rule may explain why stretching reorients divisions.
However, the precise molecular mechanism behind shape-based models remains unclear
and may vary across cell type and tissue context (Campinho et al., 2013; Fink et al., 2011;
Minc et al., 2011). Past models have used different characteristics to determine the shape of
a cell, usually selecting one of the following: cell area, cell perimeter and tricellular junction
location (TCJ, which we define here as the meeting point of three or more cells). Though
often used interchangeably, these shape characteristics model different biological functions.
We investigated their differences and determined if one characteristic predicts division
orientation better than the others.

We modelled cell shape by area, perimeter and TCJs to derive three respective measures of
cell shape orientation, θ_A, θ_P, and θ_J, and circularity, C_A, C_P, and C_J (Supplementary
Document Section 1). Cells tend to have $C_P > C_A > C_J$ i.e. shape generally appears less
anisotropic using the perimeter-based measure. C_A and C_P (and correspondingly θ_A and θ_P)
are reasonably well correlated, while C_J (and θ_J) tends to coincide less well with the others
(Figure 2A&B and S1A). Thus a cell that appears round by area and perimeter can have clear
elongation as measured by TCJs. This is intuitive for rounding mitotic cells, where TCJs can
be distributed non-uniformly around the circular periphery (Bosveld et al., 2016). However, it
is surprising that this can also be the case in cells with relatively straight edges (Figure 2A'',
note how θ_J, – yellow line – differs from θ_A and θ_P – blue and red lines – in the central dark
green cells). Notably, cells in the *Xenopus* animal cap do not undergo the dramatic mitotic
cell rounding seen in some other systems (Bosveld et al., 2016) (Figure S1B&C).

TCJ placement is a better predictor of division orientation than cell area, cell perimeter
or microtubule length.
Given that θ_A, θ_P, and θ_J are often highly correlated, division orientation is generally well
predicted by all three. We therefore focused on cases in which the orientations of shape
differed by at least 15°. In a pooled sample of 600 cells from stretched and unstretched
tissue, only 7 cells were found to have $| \theta_A - \theta_P | \geq 15°$. 58 satisfied $\left| \theta_A - \theta_J \right| \geq 15°$ and 60

satisfied $\left|\theta_P - \theta_J\right| \geq 15°$. In the latter two cases, θ_J was a significantly better predictor of

division angle than random (p < 0.0162 when $\left|\theta_A - \theta_J\right| \geq 15°$; p < 0.0042 when $\left|\theta_P - \theta_J\right| \geq$

15°; Mann-Whitney U test), but θ_A and θ_P were not (Figure 2C&D). Furthermore, C_A, C_P, and

C_J were all significantly higher in these subpopulations (Figure S1D&E; 95% confidence

intervals do not overlap), indicating that these cells are rounder, yet can still effectively orient

their spindle in-line with their TCJs. This result is strengthened considering that TCJs provide

fewer data points than area or perimeter, thus junctional data may be more susceptible to

geometric error than area and perimeter. For all of our data comparing cell shape with division

orientation, we use shape determined just prior to nuclear envelope breakdown (NEB),

avoiding any possible shape changes due to mitosis (e.g. cell rounding on entry into

mitosis/elongation at anaphase). However, to test whether the fidelity of division alignment to

TCJ shape changes depending on when shape is measured we compared $\left|\theta_D - \theta_J\right|$ at time-

points through mitosis, finding no significant difference (Figure S1F). It is important to note

that we do not see significant cell rounding in the *Xenopus* animal cap upon entry into mitosis

(Figure S1C), so static fidelity is likely a reflection of relatively static cell shape in this system,

a feature which helps simplify our analysis.

In unstretched tissue, cells which we classed as "rounded" (C_A > 0.65; Figure 2E) showed no

significant correlation between θ_A and θ_D or θ_P and θ_D, as could be expected from previous

work (Minc et al., 2011). However, θ_J was significantly aligned with division angle in these

round cells, when compared to random (p = 0.025, Mann-Whitney U test) (Figure 2F&G). This

degree of sensitivity is striking and further demonstrates that TCJ-sensing could function

effectively in round cells, which may have previously been thought to divide at random.

Our analysis is based purely on predictions arising from the data and thereby has the

advantage of being independent of unknown model parameters and assumptions. However,

to test how our division predictions compare with previous models of division orientation we

turned to a well-known shape-based model of division in isolated cells (Minc et al., 2011). The

"Minc" model hypothesises that astral microtubules exert length-dependent pulling forces on

the spindle, thereby exerting a torque and rotating the spindle, with division predicted to occur

along the axis of minimum torque. In this shape-based model, the shape of the cell

determines the distribution of torque on the spindle and thereby the division axis (see the

Supplementary Document for further details of this model and its implementation). As with our

purely geometric measures of shape, we found that the Minc model predicts division

orientation significantly better than a random distribution (Figure 2H; p < 4.1×10^{-40} for TCJs

and p < 1.2×10^{-39}, Mann-Whitney U test). However, for cells where the predicted division axes

according to TCJs (θ_J) and the Minc model (θ_{Minc}) differed by more than 15º, TCJs (θ_J)

provided a prediction of division angle that was significantly better than random (p < 0.028,

Mann-Whitney U test), while division predicted by microtubule pulling forces (θ_{Minc}) did not

(Figure 2I & J), indicating that TCJs provide a better prediction of division orientation. This

result held for multiple scaling laws between microtubule length and force (Figure S1G&H).

Local cell shape aligns with local stress and predicts division orientation better than

global stretch and stress

Contrary to observations in monolayers (Hart et al., 2017), we found that cells in stretched

tissue divide according to cell shape both when θ_J is oriented with (Figure 3A) and against

(Figure 3B&C) the direction of stretch. Moreover, in the case of cells that are relatively round

in shape (C$_J$>0.65), there is no preference for aligning with the global stretch direction and

indeed alignment with TCJ shape still appears more accurate than with the stretch axis

(Figure S2A&B; p < 0.005 for TCJS, not significant for stretch direction, Mann-Whitney U

test). These data indicate that global stretch direction is a poor predictor of division angle

when compared to cell shape. However, little is known about the local stress distribution

around individual cells in a tissue subjected to a stretch, which may not coincide with global

stress in such a geometrically heterogeneous material.

We extended a popular vertex-based model to mathematically characterise cell stress

(Brodland et al., 2014; Chiou et al., 2012; Ishihara and Sugimura, 2012; Nestor-Bergmann et

al., 2017; Nestor-Bergmann et al., 2018). Predicted orientations of forces from the model

have been found to be in accordance with laser ablation experiments (Farhadifar et al., 2007;

Landsberg et al., 2009), indicating that the model can provide a physically relevant description

of cellular stresses. Our methodology allows relative cell stress to be inferred solely from the

positions of cell vertices, without invasively altering the mechanical environment

(Supplementary Document, Section 2). The model predicts that the orientation of cell shape

based on TCJs, θ_J, aligns exactly with the principal axis of local stress (Nestor-Bergmann et

al., 2017) (Figure 3D). We demonstrated this computationally in stretched tissue by simulating

a uniaxial stretch (Figure 3E-F). Following stretch, we see that local cell stress remains

aligned with θ_J, rather than the global stress along the x-axis. Much previous work assumes

that the local axis of stress coincides with the global stress. Significantly, the model predicts

that a stress-sensing mechanism would align divisions in the same direction as a shape-

based mechanism (as in Figure 3B).

The magnitude of cell stress does not correlate with the alignment of division angle

and TCJ positioning

If a stress-sensing mechanism were contributing to orienting division, we hypothesised that

cells under higher net tension or compression might orient division more accurately with the

principal axis of stress (θ_J). We infer relative tension/compression using the isotropic

component of stress, effective pressure (P^{eff}) (Nestor-Bergmann et al., 2017):

$$P^{\text{eff}} = \frac{\tilde{A}}{\tilde{A}_0} - 1 + \frac{\Gamma \tilde{L}^2}{2\tilde{A}} + \frac{\Lambda \tilde{L}\sqrt{\tilde{A}_0}}{4\tilde{A}}$$

where $\tilde{A}$ is cell area, $\tilde{L}$ is perimeter, $\tilde{A}_0$ is the preferred area and (Λ, Γ) are model parameters,

defined in Section 2 of the Supplementary Document and inferred from data (Nestor-

Bergmann et al., 2017). Cells under net tension have $P^{\text{eff}} > 0$, whereas $P^{\text{eff}} < 0$ indicates net

compression. We provide a novel method for estimating $\tilde{A}_0$ in Section 3 of the Supplementary

Document. A representative segmentation, showing cells predicted to be under net tension

and compression, from an unstretched experiment is given in Figure 3G. Interestingly, we

found no correlation between the value of P^{eff} (relative isotropic stress) and the alignment of

division orientation to θ_J $\left(\left|\theta_D - \theta_J\right|\right)$ (Figure S2C). The mechanical state of a cell may also be

characterised by shear stress, ξ (defined as the eigenvalue of the deviatoric component of the

stress tensor, see Section 2 of the Supplementary Document). Larger values of $|\xi|$ indicate

increased cellular shear stress. Again, we found no correlation between ξ and the alignment

of division to θ_J (Figure S2D).

Despite the lack of correlation with stress magnitude, cell shape anisotropy, measured by C_J,

correlates significantly with $\left|\theta_D - \theta_J\right|$ (p < 3.04x10^{-10}, Spearman rank correlation coefficient;

Figure 3H), with elongated cells having θ_D aligned with θ_J significantly better than round cells

(p < 1.64 x 10^{-8}; Figure 3I).

Cadherin is required for positioning the mitotic spindle relative to cell shape

Immunofluorescence staining of β-catenin confirmed that adherens junctions were distributed

along the apical cell cortex, but particularly concentrated at the meeting points of three or

more cells (Figure 4E). To test a functional requirement for adherens junctions in orienting the

spindle, we focused on maternal C-cadherin (cadherin 3), which is expressed at the highest

level in Stage 10-11 *Xenopus* embryos (Heasman et al., 1994; Lee and Gumbiner, 1995). We

used two constructs to manipulate C-cadherin in the tissue: C-cadherin FL -6xmyc (CdhFL:

Full length C-cadherin with 6xmyc tags at the intracellular c-terminus) and C-cadherin ΔC -

6xmyc (CdhΔC: C-cadherin with extracellular and transmembrane domains, but lacking the

cytosolic domain) (Figure 4A) (Kurth et al., 1999). CdhFL- and CdhΔC-injected embryos

developed normally up to Stage 10/11 (Figure S3A), but the majority of embryos failed to

complete gastrulation (Lee and Gumbiner, 1995) (and data not shown). We observed no

change in the cumulative distribution of cell circularities in CdhFL- and CdhΔC-injected

tissues compared to control tissue (Figure S3B). We also saw no difference in the rate of cell

divisions (data not shown).

CdhΔC-injected tissue was elongated by application of stretch (Figure S3C), but showed a

worse alignment of divisions to stretch direction compared to uninjected control and CdhFL-

injected tissue (Figure 4B; Mann-Whitney U test p < 0.0162 for CdhΔC less than CdhFL).
Moreover, unstretched CdhΔC-injected tissue showed a significant decrease in the alignment
of division angle to θ_J, when compared to uninjected controls (Figure 4C; p < 0.016
Kolmogorov-Smirnov test on distributions differing), though both were significantly different to
random (control: p < 3.6x10^{-11}; CdhΔC: p < 4.3x10^{-11}; Kolmogorov Smirnov test). To further
investigate a requirement for adherens junctions in division orientation, we overexpressed C-
cadherin in the cell cortex by injecting CdhFL. Focussing on cells which satisfied $|\theta_P - \theta_J| \geq$
15°, we found the striking result that division orientation was now significantly well predicted
by cell perimeter, but no longer by TCJs (Figure 4D; p < 0.0027 for alignment θ_D to θ_P but not
significant for θ_D to θ_J; Mann-Whitney U test). Therefore, overexpression of CdhFL was
sufficient to switch division orientation from alignment with TCJs to alignment with the shape
of the entire cortex.
285
To investigate the mechanism behind the observed switch in division orientation, we explored
how overexpression of CdhFL alters the localisation of spindle orientation machinery at the
cell cortex. We found that overexpression of CdhFL led to a loss of the "hotspots" of β-catenin
localisation at TCJs seen in control tissue, in both interphase and mitotic cells (Figure 4E and
F). When CdhFL is overexpressed, β-catenin is more equally spread around the entire apical
perimeter of the cell (Figure 4E and F). The "hotspots" of β-catenin localisation in controls are
not purely a result of more cells contributing to this focal point, but are also seen when
fluorescence intensity is measured in single β-catenin-GFP expressing cells in the animal cap
(Figure 4F). To determine how this observed change in adherens junction localisation might
alter spindle orientation we investigated how the localisation of the spindle orientation protein,
LGN, was altered by overexpression of CdhFL. Mosaic expression of GFP-LGN allowed us to
analyse at the single-cell level in stretched and unstretched animal caps. In control tissue
LGN, like β-catenin, shows a more concentrated localisation at TCJs (Figure 4G and H). We
observed no significant difference in LGN localisation between unstretched and stretched
tissue (data not shown). However, we saw a loss of concentrated "hotspots" of LGN
localisation when CdhFL is overexpressed, with LGN instead spread more equally around the
whole perimeter (Figure 4G and H). We conclude that overexpression of CdhFL switches
division orientation from alignment with TCJs to alignment with the shape of the whole cortex
by altering the localisation of LGN. Moreover, our data indicate that, in normal circumstances,
the concentration of LGN at TCJs plays an important role in aligning divisions with cell shape.
306
Cell division rate is temporarily increased following change in global stress
Stretch elicited a reproducible and significant increase in cell division rate, with 6.47 ± 1.12 %
of cells dividing per hour in the stretched tissue compared to 3.22 ± 0.55 % in unstretched
tissue (Figure 5A, 95% confidence intervals do not overlap), as reported for cultured cells and
monolayers (Fink et al., 2011; Streichan et al., 2014b; Wyatt et al., 2015). We roughly classify
two distinct periods of division after stretch; there is an initial period of high proliferation (8.1%

cells undergoing division per hour; Figure 5B), which drops, after 40-60 minutes, to near-
unstretched control levels (4.2% cells undergoing division per hour). Stretching increases
apical tissue area by 6 ± 2.69% (95% confidence interval), and is predicted to increase global
stress by increasing individual values of P^{eff}. We sought to determine whether the increase in
division rate is a response to these changes.

In both stretched and unstretched experiments, dividing cells had a larger area than the
population, being about 22.7% and 25.7% larger on average respectively (Figure 5C).
Similarly, the mean perimeter was significantly larger in the dividing cells by about 14.1% in
unstretched and 13.8% in stretched (Figure 5D). However, there was no significant difference
in the level of cell elongation in dividing cells (Figure S2E). Crucially, we found that dividing
cells were more likely to be under predicted net tension than compression (Figure 5E, more
cells in red region). However, P^{eff} is correlated with cell area (though the two are not always
equivalent), thus a further perturbation was required to separate their effects.

Loss of myosin II reduces cell contractility
We perturbed the mechanical properties of the tissue with targeted knockdown of non-muscle
myosin II using a previously published morpholino (Skoglund et al., 2008). As expected,
myosin II knockdown disrupted cytokinesis, seen by the formation of 'butterfly' shaped nuclei,
where daughter cells had not fully separated (Figure 6A&B). However, division rate and
orientation could still be assessed using the same methods described for control tissue.
Myosin II is known to generate contractility within a tissue (Clark et al., 2014; Effler et al.,
2006; Gutzman et al., 2015). Accordingly, we found evidence for reduced contractility in the
myosin II MO tissue by observing that cells were much slower at adapting to stretch,
remaining elongated for longer (compare Figure 6C to Figure 1F).

Myosin II is required for mitotic entry in unstretched tissue
Somewhat surprisingly, considering suggestions that myosin II may play a stress-sensing role
in orienting the spindle (Campinho et al., 2013), we found that alignment of division angle to
stretch and θ_J was unaffected in global myosin II knockdown experiments (Figure 6D&E). In
contrast, proliferation rate was significantly affected, with divisions virtually ceasing in
unstretched myosin II MO tissue. Strikingly, stretching the myosin II MO tissue increased the
division rate to significantly higher levels (Figure 6F). Thus myosin II is required to cue cells
into division in the unstretched tissue, but this can be partially overridden by applying an
external loading. Unlike in control experiments, dividing cells in myosin II knockdown stretch
experiments were not significantly larger than the population in area (Figure 6G) or perimeter
(Figure 6H), thus cell area has been uncoupled as a cue to divide in the myosin II
knockdowns. This finding, along with our observation that dividing cells were more likely to be
under relative net tension than relative compression (Figure 5E), indicates that in a tissue the
cue to divide, in contrast to division orientation, is directly sensitive to mechanical force.

DISCUSSION

DISCUSSION

Previous models of cell division have demonstrated that specific features of cell shape, such
as the cell cortex or TCJs, may be important in orienting the spindle (Bosveld et al., 2016;
Hertwig, 1893; Luxenburg et al., 2011; Minc et al., 2011). We have presented a framework for
characterising cell shape in terms of its area, perimeter or TCJs (Supplementary Document).
We find that the principal axis of shape defined by TCJs is the best predictor of division angle,
better than cell shape as determined by area, perimeter or by a previous shape-sensing
model based on microtubule length (Minc et al., 2011). Moreover, the principal axis of shape
defined by TCJs aligns exactly with the principal axis of local stress (Nestor-Bergmann et al.,
2017), providing a non-invasive way to infer mechanical stress in individual cells in the
epithelium. However, division angle is not better predicted in cells with higher/lower relative
isotropic or shear stress, suggesting that cell-level mechanical stress is not a direct cue to
orient the spindle. Our findings share similarities with observations in the *Drosophila* pupal
notum, where TCJs have been hypothesised to localise force generators to orient the spindle
(Bosveld et al., 2016). Notably, however, *Xenopus* animal cap cells do not undergo the
dramatic mitotic rounding exhibited by cells in the notum.

Cell-cell adhesion has been linked to spindle orientation in MDCK cells, where E-cadherin
instructs LGN/NuMA assembly at cell-cell contacts to orient divisions (Gloerich et al., 2017).
E-cadherin polarises along a stretch axis, reorienting divisions along this axis rather than
according to cell shape (Hart et al., 2017). In accordance, we find division is less well
predicted by shape in embryos injected with C-cadherin ΔC -6xmyc, lacking the cytosolic
domain. Interestingly, over-expression of C-cadherin around the entire cell cortex leads to a
switch in division orientation, from TCJs to division best predicted by a perimeter-based
shape axis. As β -catenin is increased around the cell cortex when C-cadherin is
overexpressed, we hypothesized that this may lead to altered recruitment of spindle
orientation proteins, such as LGN and NuMA (Gloerich et al., 2017). Indeed we find that while
LGN is normally most highly localised to TCJs, overexpression of C-cadherin leads to a loss
of these "hotspots" and instead a more even spread of LGN around the entire cell perimeter.
We suggest that in the wild type situation, the hotspots of LGN localisation at TCJs will recruit
more NuMA and dynein providing localised force generation to orient the spindle according to
TCJ shape. When C-cadherin is overexpressed, and the LGN hotspots are no longer present,
we suggest that a perimeter-based shape sensing mechanism, similar to that proposed by
Minc et al (Minc et al., 2011) predominates. Our TCJ-based system of spindle orientation is
similar to the Mud-dependent TCJ-sensing mechanism in the *Drosophila* pupal notum
(Bosveld et al., 2016). However, it is important to note a key difference: NuMA, the vertebrate
homologue of *Drosophila* Mud, localises to the nucleus during interphase, only localising to
the cortex after NEB (Bowman et al., 2006; Kiyomitsu et al., 2012; Seldin et al., 2013;
Gloerich et al., 2017). Of future interest will be to determine in vertebrate tissue how the TCJ

localisation of LGN influences the highly dynamic recruitment of NuMA. Indeed, recent work
in MDCK cells has shown that on mitotic entry, when NuMA is released from the nucleus, it
competes LGN away from E-cadherin at the cortex to locally form the LGN/NuMA complex
(Gloerich et al., 2017); it will be important to determine if this is happening specifically at
TCJs.

Stretching increases proliferation rate, which correlates with cell area, perimeter and effective
pressure. We see almost no proliferation in unstretched myosin II MO experiments, although,
rather strikingly, the division rate is significantly increased following stretch. Dividing myosin II
MO cells are not significantly larger in area or perimeter than the population as a whole,
indicating that cell area has been decoupled as a division cue. Considering the established
role of myosin II as a force generator (Clark et al., 2014; Gutzman et al., 2015; Vicente-
Manzanares et al., 2009), it is possible that the myosin II MO cells cannot generate enough
internal contractility in neighbouring cells to engage the mechanical cues required for mitotic
entry. Myosin II has also been shown to function in stress-sensing pathways (Hirata et al.,
2015; Priya et al., 2015), which may explain why the proliferation rate in stretched myosin II
MO cells does not reach the levels of stretched controls. Contrary to findings in other systems
(Campinho et al., 2013), a global loss of myosin II does not alter division orientation relative to
cell shape. However, future work should look to explore whether anisotropic biases in
junctional myosin II affect division orientation, as was recently seen in the Drosophila
germband (Scarpa et al., 2018).

In conclusion, we have combined whole-tissue stretching with a biomechanical model to
propose separate roles for cell shape and mechanical stress in orienting the spindle and
cueing mitosis. The mechanism involved in orienting the mitotic spindle does not appear to
sense relative cell stress directly. Instead, division is best predicted by an axis of shape
defined by TCJs and is dependent on functional cadherin and the recruitment of LGN. In
contrast to this shape-based mechanism, we find that cells may directly sense mechanical
stress as a cue for mitotic entry, in a myosin II-dependent manner.

Author Contributions

Conceptualization, S.W. and O.E.J.; **Methodology**, A.N-B., S.W., and O.E.J.; **Software**,
A.N.B. **Investigation**, G.A.S-V., G.K.G, T.S., A.N-B. and S.W.; **Formal Analysis**, G.A.S-V.,
G.K.G. and A.N-B. **Writing – Original Draft**, S.W. and A.N-B. , G.A.S-V. and G.K.G;
Writing – Review & Editing, G.A.S-V., G.K.G., S.W., A.N-B and O.E.J.; **Funding**
Acquisition, S.W. and O.E.J.; **Supervision**, S.W. and O.E.J.

Acknowledgements
ANB was supported by a BBSRC studentship. SW, GSV and GG were supported by a

Wellcome Trust/Royal Society Sir Henry Dale Fellowship to SW [098390/Z/12/Z] with
additional funding from the Wellcome Trust ISSF [105610/Z/14/Z]. The Bioimaging Facility
microscopes used in this study were purchased with grants from BBSRC, Wellcome and the
University of Manchester Strategic Fund. We are grateful to Egor Zindy for his assistance in
developing in-house nuclei tracking scripts. Thanks to Peter March and Roger Meadows for
their help with the microscopy and to Lance Davidson for sharing Cadherin constructs. Also,
special thanks to Viki Allan, Tom Millard and Nancy Papalopulu for their critical reading of the
manuscript.

Bibliography

Benham-Pyle, B.W., Pruitt, B.L., Nelson, W.J., 2015. Mechanical strain induces E-cadherin-
dependent Yap1 and beta-catenin activation to drive cell cycle entry. Science 348, 1024–
1027. doi:10.1126/science.aaa4559

Bosveld, F., Markova, O., Guirao, B., Martin, C., Wang, Z., Pierre, A., Balakireva, M.,
Gaugue, I., Ainslie, A., Christophorou, N., Lubensky, D.K., Minc, N., Bellaïche, Y., 2016.
Epithelial tricellular junctions act as interphase cell shape sensors to orient mitosis.
Nature 530, 495–. doi:10.1038/nature16970

Bowman, S. K., Neumüller, R. A., Novatchkova, M., Du, Q., & Knoblich, J. A. (2006). The
Drosophila NuMA Homolog Mud Regulates Spindle Orientation in Asymmetric Cell
Division. Developmental Cell, 10(6), 731–
742. http://doi.org/10.1016/j.devcel.2006.05.005

Brodland, G.W., Veldhuis, J.H., Kim, S., Perrone, M., Mashburn, D., Hutson, M.S., 2014.
CellFIT: A Cellular Force-Inference Toolkit Using Curvilinear Cell Boundaries. PLoS ONE
9, e99116. doi:10.1371/journal.pone.0099116

Campinho, P., Behrndt, M., Ranft, J., Risler, T., Minc, N., Heisenberg, C.-P., 2013. Tension-
oriented cell divisions limit anisotropic tissue tension in epithelial spreading during
zebrafish epiboly. Nature cell biology 15, 1405–1414. doi:10.1038/ncb2869

Chiou, K.K., Hufnagel, L., Shraiman, B.I., 2012. Mechanical stress inference for two
dimensional cell arrays. PLoS Comput. Biol. 8, e1002512.
doi:10.1371/journal.pcbi.1002512

Clark, A.G., Wartlick, O., Salbreux, G., Paluch, E.K., 2014. Stresses at the Cell Surface
during Animal Cell Morphogenesis. Current Biology 24, R484–R494.
doi:10.1016/j.cub.2014.03.059

Davidson, L., Dassow, von, M., Zhou, J., 2009. Multi-scale mechanics from molecules to
morphogenesis. Int. J. Biochem. Cell Biol. 41, 2147–2162.
doi:10.1016/j.biocel.2009.04.015

Effler, J.C., Kee, Y.-S., Berk, J.M., Tran, M.N., Iglesias, P.A., Robinson, D.N., 2006. Mitosis-
Specific Mechanosensing and Contractile-Protein Redistribution Control Cell Shape.
Current Biology 16, 1962–1967. doi:10.1016/j.cub.2006.08.027

Farhadifar, R., Röper, J.C., Aigouy, B., Eaton, S., Jülicher, F., 2007. The influence of cell
mechanics, cell-cell interactions, and proliferation on epithelial packing. Current Biology
17, 2095–2104.

Feroze, R., Shawky, J.H., Dassow, von, M., Davidson, L.A., 2015. Mechanics of blastopore
closure during amphibian gastrulation. Dev. Biol. 398, 57–67.
doi:10.1016/j.ydbio.2014.11.011

Fink, J., Carpi, N., Betz, T., Bétard, A., Chebah, M., Azioune, A., Bornens, M., Sykes, C.,
Fetler, L., Cuvelier, D., Piel, M., 2011. External Forces Control Mitotic Spindle
Positioning. Nature cell biology 13, 771–778. doi:10.1038/ncb2269

Gloerich, M., Bianchini, J.M., Siemers, K.A., Cohen, D.J., Nelson, W.J., 2017. Cell division
orientation is coupled to cell-cell adhesion by the E-cadherin/LGN complex. Nat Commun
8. doi:10.1038/ncomms13996

Gutzman, J.H., Sahu, S.U., Kwas, C., 2015. Non-muscle myosin IIA and IIB differentially
regulate cell shape changes during zebrafish brain morphogenesis. Dev. Biol. 397, 103–
115. doi:10.1016/j.ydbio.2014.10.017

Hart, K.C., Tan, J., Siemers, K.A., Sim, J.Y., Pruitt, B.L., Nelson, W.J., Gloerich, M., 2017. E-

cadherin and LGN align epithelial cell divisions with tissue tension independently of cell
shape. Proc. Natl. Acad. Sci. U.S.A. 114, E5845–E5853. doi:10.1073/pnas.1701703114
Heasman, J., Ginsberg, D., Geiger, B., Goldstone, K., Pratt, T., Yoshindanoro, C., Wylie, C.,
1994. A Functional Test for Maternally Inherited Cadherin in Xenopus Shows Its
Importance in Cell-Adhesion at the Blastula Stage. Development 120, 49–57.
Hertwig, O., 1893. Ueber den Werth der ersten Furchungszellen für die Organbildung des
Embryo Experimentelle Studien am Frosch-und Tritonei. Archiv f. mikrosk. Anat. 42,
662–807. doi:10.1007/BF02976796
Hirata, H., Gupta, M., Vedula, S.R.K., Lim, C.T., Ladoux, B., Sokabe, M., 2015. Actomyosin
bundles serve as a tension sensor and a platform for ERK activation. EMBO Rep. 16,
250–257. doi:10.15252/embr.201439140
Hoh, J.H., Schoenenberger, C.A., 1994. Surface morphology and mechanical properties of
MDCK monolayers by atomic force microscopy. J Cell Sci 107 (Pt 5), 1105–1114.
Hutson, M.S., Tokutake, Y., Chang, M.S., Bloor, J.W., Venakides, S., Kiehart, D.P., Edwards,
G.S., 2003. Forces for morphogenesis investigated with laser microsurgery and
quantitative modeling. Science 300, 145–149. doi:10.1126/science.1079552
Ishihara, S., Sugimura, K., 2012. Bayesian inference of force dynamics during
morphogenesis. J. Theoret. Biol. 313, 201–211. doi:10.1016/j.jtbi.2012.08.017
Jones, E., Oliphant E, P., 2001. SciPy: Open Source Scientific Tools for Python [WWW
Document]. httpwww.scipy.org. URL http://www.scipy.org/ (accessed 6.8.17).
Jones, L.A., Villemant, C., Starborg, T., Salter, A., Goddard, G., Ruane, P., Woodman, P.G.,
Papalopulu, N., Woolner, S., Allan, V.J., 2014. Dynein light intermediate chains maintain
spindle bipolarity by functioning in centriole cohesion. J. Cell Biol. 207, 499–516.
doi:10.1083/jcb.201408025
Joshi, S.D., Davidson, L.A., 2010. Live-cell imaging and quantitative analysis of embryonic
epithelial cells in Xenopus laevis. J Vis Exp. doi:10.3791/1949
Kanda, T., Sullivan, K.F., Wahl, G.M., 1998. Histone-GFP fusion protein enables sensitive
analysis of chromosome dynamics in living mammalian cells. Current Biology 8, 377–
385. doi:https://doi.org/10.1016/S0960-9822(98)70156-3
Keller, R.E., 1980. The cellular basis of epiboly: an SEM study of deep-cell rearrangement
during gastrulation in Xenopus laevis. J Embryol Exp Morphol 60, 201–234.
Keller, R.E., Schoenwolf, G.C., 1977. An SEM study of cellular morphology, contact, and
arrangement, as related to gastrulation inXenopus laevis. Development Genes and
Evolution.
Kiyomitsu, T., & Cheeseman, I. M. (2012). Chromosome- and spindle-pole-derived
signalsgenerate an intrinsic code for spindle positionand orientation. Nature Cell
Biology, 14(3), 311–317. http://doi.org/10.1038/ncb2440
Klymkowsky, M.W., Hanken, J., 1991. Whole-Mount Staining of Xenopus and Other
Vertebrates. Methods in Cell Biology 36, 419–&. doi:10.1016/S0091-679X(08)60290-3
Kremer, J.R., Mastronarde, D.N., McIntosh, J.R., 1996. Computer visualization of three-
dimensional image data using IMOD. J. Struct. Biol. 116, 71–76.
doi:10.1006/jsbi.1996.0013
Kurth, T., Fesenko, I.V., Schneider, S., Münchberg, F.E., Joos, T.O., Spieker, T.P., Hausen,
P., 1999. Immunocytochemical studies of the interactions of cadherins and catenins in
the early Xenopus embryo. Dev. Dyn. 215, 155–169. doi:10.1002/(SICI)1097-
0177(199906)215:2<155::AID-DVDY8>3.0.CO;2-S
Landsberg, K.P., Farhadifar, R., Ranft, J., Umetsu, D., Widmann, T.J., Bittig, T., Said, A.,
Juelicher, F., Dahmann, C., 2009. Increased Cell Bond Tension Governs Cell Sorting at
the Drosophila Anteroposterior Compartment Boundary. Current Biology 19, 1950–1955.
doi:10.1016/j.cub.2009.10.021
Lee, C.H., Gumbiner, B.M., 1995. Disruption of Gastrulation Movements in Xenopus by a
Dominant-Negative Mutant for C-Cadherin. Dev. Biol. 171, 363–373.
doi:10.1006/dbio.1995.1288
Legoff, L., Rouault, H., Lecuit, T., 2013. A global pattern of mechanical stress polarizes cell
divisions and cell shape in the growing Drosophila wing disc. Development 140, 4051–
4059. doi:10.1242/dev.090878
Luo, T., Mohan, K., Iglesias, P.A., Robinson, D.N., 2013. Molecular mechanisms of cellular
mechanosensing. Nature Materials 12, 1063–1070. doi:10.1038/NMAT3772
Luxenburg, C., Pasolli, H.A., Williams, S.E., Fuchs, E., 2011. Developmental roles for Srf,
cortical cytoskeleton and cell shape in epidermal spindle orientation. Nature cell biology

13, 203–U51. doi:10.1038/ncb2163
Mao, Y., Tournier, A.L., Bates, P.A., Gale, J.E., Tapon, N., Thompson, B.J., 2011. Planar
polarization of the atypical myosin Dachs orients cell divisions in Drosophila. Genes Dev.
25, 131–136. doi:10.1101/gad.610511
Minc, N., Burgess, D., Chang, F., 2011. Influence of cell geometry on division-plane
positioning. Cell 144, 414–426. doi:10.1016/j.cell.2011.01.016
Minc, N., Piel, M., 2012. Predicting division plane position and orientation. Trends in Cell
Biology 22, 193–200. doi:DOI: http://dx.doi.org/10.1016/j.tcb.2012.01.003
Mishra, P., Chan, D.C., 2014. Mitochondrial dynamics and inheritance during cell division,
development and disease. Nature Reviews Molecular Cell Biology 15, 634–646.
doi:10.1038/nrm3877
Nakagawa, S., Cuthill, I.C., 2007. Effect size, confidence interval and statistical significance: a
practical guide for biologists. Biol Rev Camb Philos Soc 82, 591–605. doi:10.1111/j.1469-
185X.2007.00027.x
Nestor-Bergmann, A., Goddard, G., Woolner, S., 2014. Force and the spindle: Mechanical
cues in mitotic spindle orientation. Semin. Cell Dev. Biol. 34, 133–139.
doi:10.1016/j.semcdb.2014.07.008
Nestor-Bergmann, A., Goddard, G., Woolner, S. and Jensen, O.E., 2017. Relating cell
shape and mechanical stress in a spatially disordered epithelium using a vertex-based
model. Mathematical medicine and biology: a journal of the IMA, 35(suppl_1), pp.1-27.
doi:10.1093/imammb/dqx008
Nestor-Bergmann, A., Johns, E., Woolner, S. and Jensen, O.E., 2018. Mechanical
characterization of disordered and anisotropic cellular monolayers. Physical Review
E, 97(5), p.052409. doi: 10.1103/PhysRevE.97.052409
Pease, J.C., Tirnauer, J.S., 2011. Mitotic spindle misorientation in cancer – out of alignment
and into the fire. J Cell Sci 124, 1007–1016. doi:10.1242/jcs.081406
Petridou, N.I., Stylianou, P., Skourides, P.A., 2013. A dominant-negative provides new
insights into FAK regulation and function in early embryonic morphogenesis.
Development 140, 4266–4276. doi:10.1242/dev.096073
Priya, R., Gomez, G.A., Budnar, S., Verma, S., Cox, H.L., Hamilton, N.A., Yap, A.S., 2015.
Feedback regulation through myosin II confers robustness on RhoA signalling at E-
cadherin junctions. Nature cell biology 17, 1282–1293. doi:10.1038/ncb3239
Quyn, A.J., Appleton, P.L., Carey, F.A., Steele, R.J.C., Barker, N., Clevers, H., Ridgway,
R.A., Sansom, O.J., Naethke, I.S., 2010. Spindle Orientation Bias in Gut Epithelial Stem
Cell Compartments Is Lost in Precancerous Tissue. Cell Stem Cell 6, 175–181.
doi:10.1016/j.stem.2009.12.007
Scarpa, E., Finet, C., Blanchard, G. and Sanson, B., 2018. Actomyosin-driven tension at
compartmental boundaries orients cell division independently of cell geometry in
vivo. bioRxiv, p.397893. doi: 10.1101/397893
Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of
image analysis. Nat. Methods 9, 671–675. doi:10.1038/nmeth.2089
Seldin, L., Poulson, N. D., Foote, H. P., & Lechler, T. (2013). NuMA localization, stability, and
function in spindle orientation involve 4.1 and Cdk1 interactions. Mol Biol Cell, 24(23),
3651–3662. http://doi.org/10.1091/mbc.E13-05-0277
Skoglund, P., Rolo, A., Chen, X., Gumbiner, B.M., Keller, R., 2008. Convergence and
extension at gastrulation require a myosin IIB-dependent cortical actin network.
Development 135, 2435–2444. doi:10.1242/dev.014704
Sokac, A.M., Co, C., Taunton, J., Bement, W., 2003. Cdc42-dependent actin polymerization
during compensatory endocytosis in Xenopus eggs. Nature cell biology 5, 727–732.
doi:10.1038/ncb1025
Starborg, T., Kalson, N.S., Lu, Y., Mironov, A., Cootes, T.F., Holmes, D.F., Kadler, K.E.,
2013. Using transmission electron microscopy and 3View to determine collagen fibril size
and three-dimensional organization. Nat Protoc 8, 1433–1448.
doi:10.1038/nprot.2013.086
Stooke-Vaughan, G.A., Davidson, L.A., Woolner, S., 2017. Xenopus as a model for studies in
mechanical stress and cell division. Genesis 55, e23004. doi:10.1002/dvg.23004
Streichan, S.J., Hoerner, C.R., Schneidt, T., Holzer, D., Hufnagel, L., 2014a. Spatial
constraints control cell proliferation in tissues. PNAS 111, 5586–5591.
doi:10.1073/pnas.1323016111
Streichan, S.J., Hoerner, C.R., Schneidt, T., Holzer, D., Hufnagel, L., 2014b. Spatial

constraints control cell proliferation in tissues. PNAS 111, 5586–5591.

Sugimura, K., Lenne, P.-F., Graner, F., 2016. Measuring forces and stresses in situ in living
tissues. Development 143, 186–196. doi:10.1242/dev.119776

Théry, M., Pépin, A., Dressaire, E., Chen, Y., Bornens, M., 2006. Cell distribution of stress
fibres in response to the geometry of the adhesive environment. Cell Motility and the
Cytoskeleton 63, 341–355. doi:10.1002/cm.20126

Vicente-Manzanares, M., Ma, X., Adelstein, R.S., Horwitz, A.R., 2009. Non-muscle myosin II
takes centre stage in cell adhesion and migration. Nature Reviews Molecular Cell Biology
10, 778–790. doi:10.1038/nrm2786

Williams, M.E., Wilke, S.A., Daggett, A., Davis, E., Otto, S., Ravi, D., Ripley, B., Bushong,
E.A., Ellisman, M.H., Klein, G., Ghosh, A., 2011. Cadherin-9 Regulates Synapse-Specific
Differentiation in the Developing Hippocampus. Neuron 71, 640–655.
doi:10.1016/j.neuron.2011.06.019

Woolner, S., O'Brien, L.L., Wiese, C., Bement, W.M., 2008. Myosin-10 and actin filaments are
essential for mitotic spindle function. J Cell Biol 182, 77–88. doi:10.1083/jcb.200804062

Woolner, S., Papalopulu, N., 2012. Spindle Position in Symmetric Cell Divisions during
Epiboly Is Controlled by Opposing and Dynamic Apicobasal Forces. Developmental cell
22, 775–787. doi:10.1016/j.devcel.2012.01.002

Wyatt, T.P.J., Harris, A.R., Lam, M., Cheng, Q., Bellis, J., Dimitracopoulos, A., Kabla, A.J.,
Charras, G.T., Baum, B., 2015. Emergence of homeostatic epithelial packing and stress
dissipation through divisions oriented along the long cell axis. PNAS 112, 5726–5731.
doi:10.1073/pnas.1420585112

Xu, G.-K., Liu, Y., Li, B., 2015. How do changes at the cell level affect the mechanical
properties of epithelial monolayers? Soft Matter 11, 8782–8788.
doi:10.1039/c5sm01966d

Figure Legends

Figure 1: Application of tensile force to a multi-layered tissue. **A.** Animal cap tissue was dissected from Stage 10 *Xenopus laevis* embryos and adhered to fibronectin-coated PDMS membranes and a 35% uniaxial stretch of the membrane was applied. **B**. The animal cap tissue is 2-3 cells thick; cell shape and divisions were assessed in the apical cell layer. **C**. Displacement of nuclei was tracked in a stretched animal cap. **D**. Confocal images of the apical cells in unstretched and stretched animal caps (green: GFP-alpha-tubulin; magenta: cherry-histone2B), taken 0 and 90 minutes after stretch. Representative cells outlined by dashed lines. **E**. Rose plot showing orientation of cell shape relative to direction of stretch in unstretched (blue) and stretched (red; measured immediately following stretch) experiments. **F**. Cumulative plots of cell circularity in unstretched (blue) and stretched (red; at 0, 30, 60 and 90 mins after stretch) animal caps (0=straight line; 1=circle). 100% of cells have circularity ≤ 1. Markers slightly off-set for clarity. **G**. Rose plot of division angle relative to direction of stretch for unstretched (red) and stretched (blue) experiments. Kolmogorov-Smirnov test indicates that the unstretched distribution is not significantly different from a uniform distribution, n = 343 divisions, 15 animal caps; Kolmogorov-Smirnov test indicates that stretched distribution is significantly different from uniform, p < 1.4×10^{-9}, n = 552 divisions, 17 animal caps. Scale bar: 10μm in **B**, 500μm in **C**, 50μm in **D**.

Figure 2: Cell division orientation is best predicted by an axis of shape defined by TCJs. A. Representative image of control cells from an unstretched experiment. Scale bar: 20μm **A'.** Overlay of segmentation of cells given in **A**, with the principal axis of shape characterised by area, perimeter and junctions drawn in red, blue and yellow respectively. **A''.** Enlargement of segmented cells from white box drawn in **A'**, cells analysed are outlined by dashed white line **B.** Circularities of 2035 cells from unstretched experiments, with shape characterised by area, perimeter and junctions plotted in red, blue and yellow respectively. Cells have been ordered in descending order of perimeter-based circularity (C_P), with the corresponding values of C_A and C_J plotted alongside. **C**. Rose plot of difference between division angle, θ_D, and orientation of shape based on perimeter (blue; $\theta_{shape} = \theta_P$) and junctions (yellow; $\theta_{shape} = \theta_J$), for cells which satisfy $|\theta_P - \theta_J| \geq 15°$. **D**. Rose plot of difference between division angle, θ_D, and orientation of shape based on area (red; $\theta_{shape} = \theta_A$) and junctions (yellow; $\theta_{shape} = \theta_J$), for cells which satisfy $|\theta_A - \theta_J| \geq 15°$. **E**. Examples of round (top) and elongated (bottom) cells where division angle (black arrows) is well predicted by the principal axis of shape defined by area (yellow arrows). **F.** Rose plot of difference between division angle, θ_D, and orientation of shape based on perimeter (blue; $\theta_{shape} = \theta_P$) and junctions (yellow; $\theta_{shape} = \theta_J$), for round cells which satisfy C_A > 0.65. **G**. Rose plot of difference between division angle, θ_D, and orientation of shape based on area (red; $\theta_{shape} = \theta_A$) and junctions (yellow; $\theta_{shape} = \theta_J$), for round cells which satisfy C_A> 0.65. See also Figure S1. **H** Rose plot of difference between division angle, θ_D, and orientation of shape based on

Minc model when β = 3 (magenta; $\theta_{shape} = \theta_{Minc}$) and junctions (yellow; $\theta_{shape} = \theta_J$) for all

cells in stretched and unstretched experiments (n = 599 cells). **I.** Rose plot of difference

between division angle, θ_D, and orientation of shape based on Minc model when β = 3

(magenta; $\theta_{shape} = \theta_{Minc}$) and junctions (yellow; $\theta_{shape} = \theta_J$), for cells which satisfy $|\theta_{Minc} -$

$\theta_J| \geq 15°$ (n = 65 cells). **J.** Cumulative plot of difference between division angle, θ_D , and

orientation of shape for data shown in I.

Figure 3: Division orientation is better predicted by shape rather than high relative

isotropic or shear stress. A. Images taken from a confocal timelapse movie of a division in

a cell in stretched tissue whose interphase shape (dashed line, 0:00) is oriented with the

stretch (horizontal) axis. Cell division aligns with both cell shape and stretch axis. **B.**

Timelapse images of an unusual cell in a stretched tissue, whose interphase shape (dashed

line, 0:00) is oriented against the stretch axis. Cell division aligns with cell shape but against

the stretch axis. **C.** Rose plot of difference between division angle, θ_D, and orientation of

shape based on junctions, θ_J, for cells from stretched experiments, where θ_J was at least $60°$

divergent to the direction of stretch. 29 cells satisfied this condition. Kolmogorov-Smirnov test

found a significant difference from a uniform distribution (p=0.022). **D.** Representative cells

showing classification of cell stress configurations. Red (blue) cells are under net tension

(compression), where P^{eff} is positive (negative). Larger (smaller) black arrows indicate the

orientation of the principal (secondary) axis of stress, with inward- (outward)-pointing arrows

indicating the tension (compression) generated by the cell. Yellow arrows indicate the

principal axis of shape defined by cell junctions, which aligns exactly with a principal axis of

stress. **E.** 50 simulated cells randomly generated in a periodic box, relaxed to equilibrium with

parameters (Λ,Γ) = (-0.259, 0.172), under conditions of zero global stress (Nestor-Bergmann

et al., 2017). Red (blue) cells are under net tension (compression). Principal axis of stress

(shape) indicated in black (yellow). **F.** Cells from **E** following a 13% area-preserving uniaxial

stretch along the x-axis. **G.** Example segmented cells from an unstretched experiment. Cells

in red (blue) are predicted to be under net tension (compression). **H.** Cell circularity defined by

junctions, C_J, vs $|\theta_D - \theta_J|$. Spearman rank correlation coefficient found a significant

correlation (p < 3.04 x 10⁻¹⁰). Elongated cells ($C_J \leq 0.65$) cluster in blue box, whereas rounded

cells ($C_J > 0.65$) have a more uniform distribution. **I.** Rose plot of difference between division

angle, θ_D, and orientation of shape based on junctions, θ_J for round ($C_J > 0.65$; right) and

elongated ($C_J \leq 0.65$; left) cells shown in **H**. Mann-Whitney U test indicated that elongated

cells have θ_J aligned significantly more with θ_D than rounded cells (p < 1.64 x 10⁻⁸). Scale bar

in **A&B:** 20μm. All rose plots show percentage of cells. See also Figure S2.

Figure 4: C-Cadherin is required to orient the mitotic spindle according to cell shape.

A. Schematic of Cadherin contructs CdhFL and CdhΔC **B.** Rose plot of division angles, θ_D,

relative to direction of stretch for cells from stretched CdhΔC-injected (411 cells; cyan) and

stretched CdhFL-injected experiments (552 cells; orange). CdhFL-injected cells align significantly better with direction of stretch (p < 0.0162, Mann-Whitney U test). **C.** Rose plot of difference between division angle, θ_D, and orientation of shape based on junctions, θ_J, for cells from CdhΔC-injected experiments (390 cells; cyan) and control experiments (239 cells; blue). Distributions are significantly different (p < 0.016 Kolmogorov-Smirnov test). **D.** Rose plot of difference between division angle, θ_D, and orientation of shape based on perimeter, θ_P, (blue) and junctions, θ_J, (yellow) for 96 cells from CdhFL-injected experiments which satisfied $|\theta_P - \theta_J| \geq 15°$. θ_D aligns significantly better to θ_P than a random distribution (p < 0.004; Kolmogorov-Smirnov test), but not to θ_J. **E.** Single confocal slices from immunofluorescent staining for β-catenin (green) and myc-tag (magenta) in uninjected and CdhFL-injected St 12 embryos (stage matched to time that animal caps are stretched and imaged). Hotspots of β-catenin localization (arrows) are seen at TCJs in controls but are lost when CdhFL is overexpressed. **F.** Quantification of β-catenin-GFP localization at TCJs compared to cell edges in single mitotic cells in animal caps. β-catenin-GFP is more strongly localised at TCJs compared to cell edges in controls but this bias is lost in CdhFL-injected tissue (*p < 0.01 Komogorov-Smirnov test; n = 14, 16 mitotic cells from 6, 4 unrelated animal caps for control and CdhFL, respectively). **G.** Time-lapse movies of control and CdhFL-injected animal cap tissue expressing GFP-LGN in a mosaic fashion. In control cells, GFP-LGN is enriched at TCJs during interphase (arrows) and this localization persists through mitosis. The enrichment of GFP-LGN at TCJs is lost when CdhFL is expressed, with localisation spread throughout the cell edge. **H.** Quantification of GFP-LGN localisation at TCJs compared to cell edges in single mitotic cells in animal caps. GFP-LGN is more strongly localised at TCJs compared to cells edges in controls but this bias is lost in CdhFL-injected tissue (*p < 0.05 Komogorov-Smirnov test; n = 21, 23 mitotic cells from 7, 6 unrelated animal caps for control and CdhFL, respectively). Red points show quantification for mitotic cells highlighted in G. Rose plots show percentage of cells. Scale bars represent 20μm. See also Figure S3.

Figure 5: Stretching increases division rate. Dividing cells have large area, perimeter and relative effective pressure. A. Division rate (percentage of cells entering mitosis per hour) increases in stretched tissue compared to unstretched. 95% confidence intervals do not overlap, indicating significant difference. Each point represents the mean division rate from an animal cap. **B.** Percentage of cells that have undergone nuclear envelope breakdown (NEB) with respect to time in control stretched (red) and unstretched (blue) experiments from **A**. Dashed lines indicate linear lines of best fit; control unstretched experiments have gradient 4.2% cells undergoing division per hour. Stretched experiments have initial gradient 8.1% and then 4.35% cells undergoing division per hour. **C.** Comparison of mean area of population of all cells vs dividing cells from unstretched and stretched control experiments. Error bars represent mean and 95% confidence intervals, which do not overlap between the population and dividing cells, indicating a significant difference. **D.** Comparison of mean perimeter of population of all cells vs dividing cells from unstretched and stretched control experiments.

Error bars represent mean and 95% confidence intervals, which do not overlap between the

population and dividing cells, indicating a significant difference. **E.** Heat map showing

predicted relative isotropic stress (effective pressure, P^{eff}) of dividing cells from control

unstretched experiments. Areas and perimeters have been nondimensionalised using the

preferred areas, $\tilde{A}_0$, fitted to each experiment in Supplemental Figure 4C. Polygonal class

(number of neighbours) indicated by marker colour and style, with (4,5,6,7,8+) sided cells

given in (blue, green, red, purple, yellow). Dashed vertical line represents mean area of all

cells. Cells lying in red (blue) regions are under predicted net tension (compression).

Figure 6: Myosin II MO cells maintain alignment of division to TCJ shape, but have

perturbed proliferation rate. A. Images taken from a confocal timelapse movie of stretched

myosin II morpholino injected animal cap explants at 0 and 90 minute intervals. Butterfly

nuclei seen prominently at 90 minutes, where nuclei are in contact. **B.** Timelapse images of

control morpholino-injected stretched animal cap explants at 0 and 90 minute intervals. **C.**

Cumulative distribution of cell circularity defined by area, C_A, in myosin II MO knockdown

stretched animal caps (shaded green) at t=0, 30, 60 and 90 minutes after stretch. Cumulative

distribution for unstretched t=0 control MO knockdown experiments shown in blue. Error bars

represent 95% confidence intervals. Error bars for myosin II MO t=90 minutes distribution

does not overlap with control MO, indicating a significant difference from unstretched shape.

Markers are slightly off-set for clarity. **D.** Rose plot of difference between division angle, θ_D,

and orientation of shape based on junctions, θ_J, for 216 cells from myosin II knockdown

stretched experiments. Mann-Whitney U test found significant alignment compared to random

(p < 5.72x10^{-15}), but no significant difference from equivalent dataset in control stretched

experiments. Percentages of cells shown. **E**. Rose plot of division angle relative to direction

for stretch for control MO (532 cells; blue) and myosin II MO (301 cells; green) experiments.

Mann-Whitney U and Kolmogorov Smirnov test found no significant difference between the

two. **F.** Division rate (percentage of total cells entering mitosis per hour) in unstretched and

stretched tissue from myosin II MO (green; n=10 for unstretched and n=12 for stretched) and

control MO (blue; n=13 for unstretched and n=10 for stretched) experiments. Error bars

represent mean and 95% confidence intervals. **G.** Comparison of mean area of population of

all cells vs dividing cells from stretched myosin II knockdown experiments. Error bars

represent mean and 95% confidence intervals, which overlap, indicating no significant

difference. **H.** Comparison of mean perimeter of population of all cells vs dividing cells from

stretched myosin II knockdown experiments. Error bars represent mean and 95% confidence

intervals, which overlap, indicating no significant difference. Scale bar in **A** and **B**: 100µm.

Materials and Methods

Xenopus laevis embryos and microinjection

Xenopus laevis embryos were obtained and injected as described previously (Woolner and Papalopulu, 2012). RNA was synthesised as described previously (Sokac et al., 2003) and microinjected at the following needle concentrations: 0.5 mg/ml GFP-α-tubulin; 0.1 mg/ml cherry-histone2B(Kanda et al., 1998); 0.125 mg/ml cadherin 3a full length:6x myc-tag; 0.125 mg/ml cadherin 3a deleted cytosolic domain:6x myc-tag (Kurth et al., 1999). For mosaic expression of β-catenin-GFP (Addgene plasmid #16839, Randall Moon) and GFP-LGN (sub-cloned into pCS2+ from Addgene plasmid #37360, Iain Cheeseman), RNA was injected into a single cell at the 4-cell stage at 0.25 mg/ml (needle concentration). Morpholinos prepared as 1mM stocks (diluted in water) were heated at 65°C for 5 minutes and microinjected at a needle concentration of 1mM and needle volume of 2.5nl into all cells of four-cell stage embryos. The morpholinos used were MHC-B (Myosin Heavy Chain-B, myosin II) MO (5'-CTTCCTGCCCTGGTCTCTGTGACAT-3'; (Skoglund et al., 2008) and standard control MO (5'-CCTCTTACCTCAGTTACAATTTATA-3'; Gene Tools LLC). All embryos were incubated at 16°C for approximately 20 hours prior to animal cap dissection.

Animal cap dissection and culture

Animal cap tissue was dissected from the embryo at stage 10 of development (early gastrula stage) following a previously described protocol (Joshi and Davidson, 2010), and cultured in Danilchik's for Amy explant culture media (DFA; 53mM $NaCl_2$, 5mM Na_2CO_3, 4.5mM Potassium gluconate, 32mM Sodium gluconate, 1mM $CaCl_2$, 1mM $MgSO_4$) on a 20mm × 20mm elastomeric PDMS (Sylgard 184, SLS) membrane made in a custom mold and coated with fibronectin (fibronectin from bovine plasma, Sigma). Explants were held in place by a coverslip fragment. Each membrane was then incubated at 18°C for at least 2 hours prior to imaging.

Animal cap stretch manipulation and confocal imaging

Each PDMS membrane was attached to a stretch apparatus (custom made by Deben UK Limited) fixed securely to the stage of a Leica TCS SP5 AOBS upright confocal and a 0.5mm (to remove sag on the membrane) or 8.6mm uniaxial stretch was applied for unstretched and stretched samples respectively. Images were collected on a Leica TCS SP5 AOBS upright confocal using a 20x/0.50 HCX Apo U-V-I (W (Dipping Lens)) objective and 2x confocal zoom. The distance between optical sections was maintained at 5μm and the time interval between each frame was 20 seconds, with each sample being imaged for up to 2.5 hours. For quantification of β-catenin-GFP and GFP-LGN localisation, animal caps were prepared as described but timelapse movies were collected with 2μm optical sections and a time interval of 1 min between frames. Maximum intensity projections of these 3D stacks are shown in the

829   results; except for the GFP-LGN timelapse (Figure 4G), which is an average intensity
830   projection.
831
Scanning EM
Uninjected embryos were allowed to develop to stage 10 at 16°C and then animal cap tissue
was dissected and allowed to adhere to a fibronectin PDMS membrane as described
previously. After 2 hours the animal caps were fixed following a protocol previously detailed
by (L. A. Jones et al., 2014). Briefly, the animal caps were fixed in 3.7% PFA and 2.5%
Glutaraldehyde in BRB80 buffer (80mM PIPES, 1mM $MgCl_2$, 1mM EGTA, pH 6.8) overnight
at 4°C. Samples were processed using a high density staining method detailed in full by
(Williams et al., 2011) (supplementary protocol), but briefly comprising a 1 hour fix in 2%
(wt/vol) osmium tetroxide and 1.5% (wt/vol) potassium ferrocyanide in 0.1M cacodylate buffer.
This was followed by a 20 minute incubation in 1% (wt/vol) thiocarbohydrazide and a 30
minute incubation in 2% (wt/vol) osmium teroxide, followed by a final incubation in 1% (wt/vol)
uranyl acetate overnight at 4°C. Samples were then stained with freshly prepared Walton's
lead aspartate (0.02M lead nitrate and 0.03M in aspartic acid, adjusted to pH 5.5) for 30
minutes at 60°C prior to dehydration, embedding in Epon 812 (hard formulation), and
trimming on a standard microtome. Samples were visualised using a microtome (3View;
Gatan) within a Quanta 250 FEG; FEI scanning electron microscope using the following
imaging conditions: indicated quadrant magnification of 1600x, accelerating voltage of 3.8kV,
pressure at 0.33 Torr. Images were collected at 4000 x 5000 pixels with a dwell time of 10 µs.
Raw data was converted to an MRC file stack using IMOD (Kremer et al., 1996; Starborg et
al., 2013) and further processed using Imaris software (Bitplane).
852
Image analysis
Image analysis was performed using ImageJ (Schneider et al., 2012). Cell division orientation
was quantified using the straight-line tool to draw a line between the dividing nuclei of a cell in
late anaphase (a stage in mitosis where division orientation is set and the spindle undergoes
no further rotation (Woolner et al., 2008; Woolner and Papalopulu, 2012)). Using the ROI
manager the angle of division relative to stretch (horizontal axis) was recorded along with the
frame and location of the division. Single cell edges and junctions were manually traced 40s
before NEB using the freehand paintbrush tool. The whole population of cells in the apical
layer of the animal cap was manually traced, along with peripheral junctions and cell centres,
using the freehand paintbrush tool. Segmentation of the cell boundaries was performed using
in-house Python scripts implementing a watershed algorithm. Geometric features of the cells,
such as area and perimeter, were extracted and analysed in Python; for further details on how
cell shape was characterised using the segmented images, please see the Supplementary
Document. To quantify β-catenin-GFP and GFP-LGN localisation at TCJs in mitotic cells,
movies of unstretched and stretched animal caps were analysed as follows: mitotic cells
which had non-expressing neighbours were selected at early metaphase. A single optical

slice which was level with the centre of metaphase nuclei (visualised by mCherry-H2B) was
selected and ROI's were drawn around TCJs and the corresponding cell edges in Image J.
Mean grey values were measured for each ROI and TCJ and cell edge grey values were
averaged (mean) for each cell. A ratio between average TCJ intensity and average cell edge
intensity (Mean TCJ intensity/Mean cell edge intensity) for each mitotic cell was then
calculated.
875

Data analysis

The data analysis and plotting was carried out using in-house Python scripts. Statistical tests
were performed using the SciPy library (E. Jones and Oliphant E, 2001) and Prism 7
(GraphPad Software, Inc). Mann-Whitney U tests were used to assess if rose histograms
were distributed closer to zero. Kolmogorov-Smirnov tests were used to assess if two
distributions were significantly different. Otherwise, bootstrapping with 95% confidence
intervals, which allow the precision of the estimate to be seen (Nakagawa and Cuthill, 2007),
were used to assess significance.
884

Immunofluorescence

Embryos were fixed at stage 12 following the protocol previously detailed by Jones et al.,
(2014)(L. A. Jones et al., 2014). Embryos were incubated in primary and secondary
antibodies in TBSN/BSA (Tris- buffered saline: 155mM NaCl, 10mM Tris-Cl [pH 7.4]; 0.1%
Nonidet P-40; 10 mg/ml BSA) overnight at 4°C, with five 1 hour washes with TBSN/BSA
following each incubation. Primary antibodies were: anti-β-catenin at 1:200 dilution, raised in
rabbit (Abcam) and anti c-myc 9E10 at 1:1000 dilution, raised in mouse (Santa-cruz). Alexa
Fluor secondary antibodies, anti-rabbit 488 and anti-mouse 568 (Invitrogen) were used at a
dilution of 1:400. After staining, embryos were methanol dehydrated, then cleared and
mounted in Murray's Clear (2:1, benzyl benzoate:benzyl alcohol; (Klymkowsky and Hanken,
1991)). Images were collected on a Leica TCS SP5 AOBS inverted confocal using a 63x HCX
PL APO (Oil λ BL) objective and 1024 x 1024 format. Single confocal slices are shown in the
results.
898

Implementation of microtubule length division model by Minc et al. (2011)

900 Details of the model and implementation are given in the Supplementary Document. The
901 predicted torques and corresponding division axes were calculated using in-house Python
902 scripts that are available upon request.
903

Implementation of the vertex-based model

905 The numerical simulations of the vertex-based model were carried out using the same scripts
906 outlined in section 3.8 of (Nestor-Bergmann et al., 2017). Model parameters used for all
907 simulations were $(\Lambda, \Gamma) = (-0.259, 0.172)$, determined using a fitting procedure described in
908 (Nestor-Bergmann et al., 2017).

Figure 1

Figure 2

Figure 3: Local stress aligns with shape. Division orientation is better predicted by shape in elongated cells, rather than those with higher relative isotropic or shear stress. A. Images taken from a confocal timelapse movie of a division in a cell in stretched tissue whose interphase shape (dashed line, 0:00) is oriented with the stretch (horizontal) axis. Cell division aligns with both cell shape and stretch axis. **B.** Timelapse images of an unusual cell in a stretched tissue, whose interphase shape (dashed line, 0:00) is oriented against the stretch axis. Cell division aligns with cell shape but against the stretch axis. **C.** Rose plot of difference between division angle, θ_D, and orientation of shape based on junctions, θ_J, for cells from stretched experiments, where θ_J was at least $60°$ divergent to the direction of stretch. 29 cells satisfied this condition. Kolmogorov-Smirnov test found a significant difference from a uniform distribution (p=0.022). **D.** Representative cells showing classification of cell stress configurations. Red (blue) cells are under net tension (compression), where P^{eff} is positive (negative). Larger (smaller) black arrows indicate the orientation of the principal (secondary) axis of stress, with inward- (outward)-pointing arrows indicating the tension (compression) generated by the cell. Yellow arrows indicate the principal axis of shape defined by cell junctions, which aligns exactly with a principal axis of stress. **E.** 50 simulated cells randomly generated in a periodic box, relaxed to equilibrium with parameters (Λ, Γ) = (-0.259, 0.172), under conditions of zero global stress (Nestor-Bergmann et al., 2017). Red (blue) cells are under net tension (compression). Principal axis of stress (shape) indicated in black (yellow). **F.** Cells from **E** following a 13% area-preserving uniaxial stretch along the x-axis. **G.** Example segmented cells from an unstretched experiment. Cells in red (blue) are predicted to be under net tension (compression). **H.** Cell circularity defined by junctions, C_J, vs $|\theta_D - \theta_J|$. Spearman rank correlation coefficient found a significant correlation (p < 3.04 x 10^{-10}). Elongated cells ($C_J \leq 0.65$) cluster in blue box, whereas rounded cells ($C_J > 0.65$) have a more uniform distribution. **I.** Rose plot of difference between division angle, θ_D, and orientation of shape based on junctions, θ_J for round ($C_J > 0.65$; right) and elongated ($C_J \leq 0.65$; left) cells shown in **H**. Mann-Whitney U test indicated that elongated cells have θ_J aligned significantly more with θ_D than rounded cells (p < 1.64 x 10^{-8}). Scale bar in **A&B**: 20µm. All rose plots show percentage of cells.

Figure 4

Figure 5: Stretching increases division rate. Dividing cells have large area, perimeter and relative effective pressure. A. Division rate (percentage of cells entering mitosis per hour) increases in stretched tissue compared to unstretched. 95% confidence intervals do not overlap, indicating significant difference. Each point represents the mean division rate from an animal cap. **B.** Percentage of cells that have undergone nuclear envelope breakdown (NEB) with respect to time in control stretched (red) and unstretched (blue) experiments from **A**. Dashed lines indicate linear lines of best fit; control unstretched experiments have gradient 4.2% cells undergoing division per hour. Stretched experiments have initial gradient 8.1% and then 4.35% cells undergoing division per hour. **C.** Comparison of mean area of population of all cells vs dividing cells from unstretched and stretched control experiments. Error bars represent mean and 95% confidence intervals, which do not overlap between the population and dividing cells, indicating a significant difference. **D.** Comparison of mean perimeter of population of all cells vs dividing cells from unstretched and stretched control experiments. Error bars represent mean and 95% confidence intervals, which do not overlap between the population and dividing cells, indicating a significant difference. **E.** Heat map showing predicted relative isotropic stress (effective pressure, P^{eff}) of dividing cells from control unstretched experiments. Areas and perimeters have been nondimensionalised using the preferred areas, $\tilde{A}_0$, fitted to each experiment in Supplemental Figure 4C. Polygonal class (number of neighbours) indicated by marker colour and style, with (4,5,6,7,8+) sided cells given in (blue, green, red, purple, yellow). Dashed vertical line represents mean area of all cells. Cells lying in red (blue) regions are under predicted net tension (compression).

Figure 6: Myosin II MO cells maintain alignment of division to tricellular junctional shape, but have perturbed proliferation rate. A. Images taken from a confocal timelapse movie of stretched myosin II morpholino injected animal cap explants at 0 and 90 minute intervals. Butterfly nuclei seen prominently at 90 minutes, where nuclei are in contact. **B.** Timelapse images of control morpholino-injected stretched animal cap explants at 0 and 90 minute intervals. **C.** Cumulative distribution of cell circularity defined by area, C_A, in myosin II MO knockdown stretched animal caps (shaded green) at t=0, 30, 60 and 90 minutes after stretch. Cumulative distribution for unstretched t=0 control MO knockdown experiments shown in blue. Error bars represent 95% confidence intervals. Error bars for myosin II MO t=90 minutes distribution does not overlap with control MO, indicating a significant difference from unstretched shape. Markers are slightly off-set for clarity. **D.** Rose plot of difference between division angle, θ_D, and orientation of shape based on junctions, θ_J, for 216 cells from myosin II knockdown stretched experiments. Mann-Whitney U test found significant alignment compared to random ($p < 5.72 \times 10^{-15}$), but no significant difference from equivalent dataset in control stretched experiments. Percentages of cells shown. **E.** Rose plot of division angle relative to direction for stretch for control MO (532 cells; blue) and myosin II MO (301 cells; green) experiments. Mann-Whitney U and Kolmogorov Smirnov test found no significant difference between the two. **F.** Division rate (percentage of total cells entering mitosis per hour) in unstretched and stretched tissue from myosin II MO (green; n=10 for unstretched and n=12 for stretched) and control MO (blue; n=13 for unstretched and n=10 for stretched) experiments. Error bars represent mean and 95% confidence intervals. **G.** Comparison of mean area of population of all cells vs dividing cells from stretched myosin II knockdown experiments. Error bars represent mean and 95% confidence intervals, which overlap, indicating no significant difference. **H.** Comparison of mean perimeter of population of all cells vs dividing cells from stretched myosin II knockdown experiments. Error bars represent mean and 95% confidence intervals, which overlap, indicating no significant difference. Scale bar in **A** and **B**: 100μm.