Stereotyped p53 binding tuned by chromatin accessibility

Antonina Hafner[1,*,2]**, Galit Lahav**[1,3] **and Jacob Stewart-Ornstein**[1*]

[1]Department of Systems Biology, Harvard Medical School, Boston, MA 02115, USA

*These authors contributed equally to this work

Corresponding author: jacob_stewart-ornstein@hms.harvard.edu

[2]Antonina Hafner: antonina@hms.harvard.edu

[3]Galit Lahav: galit_lahav@hms.harvard.edu

Keywords: p53, ChIPseq, Chromatin, DNA damage, Gene Expression

Abstract

<u>Background</u>

Transcription factors orchestrate the cellular response to stimuli by binding specifically to DNA and

activating associated genes. In the DNA damage response, the tumor suppressor p53 regulates much of

the transcriptional response and has been suggested to selectively regulate gene expression in different

contexts. However, comparison between genome-wide studies shows a large overlap between p53 bound

loci and thus questions this selectivity.

<u>Results</u>

To systematically assess the cell type specificity of p53, we directly measured its association with DNA in

12 p53 wild-type cell lines in response to ionizing radiation. We found that the vast majority of bound

sites are occupied across all cells lines uniformly. Gene expression, on the other hand showed substantial

variability between cell lines. The coherence of our dataset, allowed us to identify a small subset of

binding sites that appeared in just one or a few cell lines. We found that intrinsic chromatin accessibility

of a cell line explained these differential p53 binding preferences. Moreover, we were able to manipulate

p53 binding by altering chromatin state.

<u>Conclusion</u>

Our results show the limited contribution of genomic sequence to p53 binding and suggest that in vivo

factors including chromatin accessibility largely regulate its binding.

Background

Transcription factors (TF) are a large family of proteins that can bind to DNA and induce or repress transcription of genes [1]. Substantial effort, notably through the ENCODE project, has gone into generating genome wide maps of TF-DNA associations, gene expression and chromatin states. These genome wide datasets have been used to identify gene regulatory networks that allow cell type and stimulus-specific gene expression and led to characterization of a handful of networks, as for example in the embryonic stem cells system [2, 3]. However, these studies have also revealed the complexity of gene regulation and the large space of combinatorial interactions between TFs [4, 5]. Identifying the mechanisms that allow tissue and stimulus-specific gene expression remains a challenge and an active field of research.

In this work, we explored cell type and stimulus specificity of the tumor suppressing transcription factor p53 at the level of DNA binding and regulation of gene expression. The major DNA damage response (DDR) regulator in mammals, p53 is necessary and sufficient to impose terminal cell fates on irradiated cells. Though ubiquitously expressed across human tissues, it remains unclear if p53 DNA binding is universal, which could be advantageous for a critical stress response system, or shaped by the chromatin and cofactor environment of different cell types and lineages. Context specific regulation of gene expression by p53 has been a long standing hypothesis in the p53 field, and implies that p53 can integrate information about cellular context and the type of stress to selectively activate some target genes versus others [6-9]. However, studies querying p53 binding genome wide in different human cell lines and upon different treatments have variously found strong agreement in p53 binding locations [10, 11] and activation of a core set of target genes [12]. Other studies have argued for unique cell type or transformation status specific p53 activity [13-15]. These studies compared pairs of cell lines or supplemented single cell line data with meta-analysis of published datasets, an approach that is powerful for identifying universal p53

binding sites, but has limits for detection of cell line specific binding patterns due to divergent

experimental conditions across datasets.

To study how p53 binding varies across cell lines, we measured p53 DNA binding in 12 cell lines from

different tissue types in response to a single treatment. Indeed, we took advantage of our previous

characterization of these cell lines which showed a comparable acute response of p53 [16] in response to

ionizing radiation (IR). By treating this panel of epithelial cell lines with a dose of IR sufficient to induce

uniform p53 activation across cell lines, and measuring p53 binding at an early (2 hours) time-point we

minimized secondary effects and focused on measuring the rapid and direct binding of p53. Our approach

differs from the majority of p53 datasets in the literature, which use chemotherapy agents such as

doxorubicin or the p53 activator Nutlin3A at later time-points of 6-12 hours. This coherent set of samples

allowed us to rigorously explore the heterogeneity of p53 binding and identify the influence of universal

genomic and cell line specific chromatin factors on p53 binding and gene expression.

Results

p53 binding across the genome is stereotyped between cell lines and treatments

To study how p53 binding varies across cell lines we treated a set of 12 cell lines expressing wild type p53

with ionizing radiation (IR; X-Ray 4Gy) to induce p53 activity and 2 hours later cross-linked and harvested

each cell line for ChIP-Seq analysis. Well established p53 target genes showed robust binding in all 12 cell

lines (Fig. 1A). Overall, by pooling data from all cell lines we confidently called ~9000 p53 ChIP peaks. *De*

novo motif analysis identified a motif bearing strong similarity to previously identified p53 binding motifs

as the strongest single motif and also found it to be centrally enriched within peaks (Fig. 1B).

The quantitative strength of p53 binding at each genomic locus was highly conserved across the 12 cell

lines (Fig. 1C). Though no strong groups of cell lines appeared by eye, hierarchical clustering correctly

sorted the cell lines by tissue of origin, with pairs of lung and kidney lines, and all five melanoma lines

clustering together (Fig. 1C). These p53 bound regions were also similar to other published datasets

(average within dataset correlation 0.53, average correlation to external datasets 0.49; Fig. S1) and

consistent with previous work on p53 binding locations [10]. It was previously suggested that cancer cell

lines show a different p53 binding profile from non-cancerous cells [13]. We therefore compared the 12

cancer cell lines to an identically treated non-transformed line, RPE1. We found that p53 binding at

identified sites in RPE1 cells in response to IR was highly correlated with our dataset in transformed cell

lines (Fig. 1D; average R= 0.48 for correlation (RPE, Cancer Lines) vs 0.53 for correlation (Cancer, Cancer)).

To further explore if the apparent uniformity of p53 binding is specific to IR, we compared p53 ChIP peaks

in MCF7 and UACC257 cell lines treated with either IR or the p53 activator Nutlin3A. We observed

condition-condition correlations within each line that were stronger than any line- line correlations (Fig.

1E; R=0.87 or 0.88 for MCF7 and UACC257, respectively, vs R=0.73 for the maximum line-line). Thus in

addition to the uniformity we observe between cell lines, IR induced and pharmacologically induced p53

do not lead to distinct p53 function as measured by acute p53 DNA binding.

Genomic DNA sequence has limited predictive power for p53 binding

Given the strong conservation of p53 binding across cell lines we hypothesized that p53 binding might be

largely determined by genomic DNA sequences. We tested this by comparing motif scores (calculated

from the position weight matrix) with p53 ChIP-Seq signal intensity. The extent of the correlation between

p53 binding and PWM score was highly cell line dependent (Fig. 2A), ranging from no correlation to

correlation of 0.22 in a single cell line. Averaging p53 binding over increasing numbers of cell lines resulted

in better agreement between genomic motif score and p53 binding, with the highest correlation of 0.26

averaged across all datasets (Fig. 2 A,B). Therefore, although the motif score significantly correlates with

p53 DNA binding (p=1.98e-132), it only allows to account for slightly more than 5% of the variance.

One possible explanation for why the p53 motif only had modest predictive power was the presence of

binding elements in the DNA that were not well accounted for by a PWM. It has been proposed, for

example, that flanking nucleotide identity and also general 'DNA shape' features of the binding site might

influence binding site choice [17, 18]. To explore if our motif analysis was simply a poor model of p53

binding we performed an *in vitro* ChIP experiment where recombinant p53 was incubated with

fragmented genomic, followed by immunoprecipitation and deep sequencing similarly to a recently

published protocol [19]. As this assay uses sheared genomic DNA with a size of ~300-600bp factors such

as flanking nucleotides and DNA shape should be appropriately measured. Preforming this assay, we

obtained a strong signal of p53 binding, recovering a consensus p53 motif (1e-2422, Fig 2C). We observed

p53 binding sites, such as the one proximal to the CDKN1A/p21 promoter, that showed strong *in vivo*

binding, a strong motif, and substantial *in vitro* p53 binding signal (Fig. 2D). Whereas, other binding sites

such as the one contained in the first intron of MDM2 showed substantial *in vivo* binding, but little *in vitro*

binding (and no strong motif). Other sites such as one in the MDM4 gene showed strong *in vitro* binding

(and a strong motif), but little *in vivo* binding. Overall, the *in vitro* p53 binding signal did not show a better

correlation (R=0.252, P= 3.10e-127, Fig. 2E) with *in vivo* p53 binding than the motif score These results

suggest that factors other than just DNA sequence determine p53 binding *in vivo*.

A subset of p53 binding sites is highly variable

To see if we could identify what factors regulate p53 binding *in vivo* we looked for binding sites that were

variably occupied across cell lines, with the reasoning that perhaps by understanding cell line-cell line

variation in binding would give a clue to *in vivo* regulators of p53 binding. We compared the cell line to

cell line variability in p53 peak height, correcting for the average peak height (which contributes shot noise

to our analysis), and identified about 5% of binding sites (494 peaks) that showed high variation between

cell lines relative to their average peak strength (Fig. 3A, B). For example, p53 peaks nearby the

inflammatory associated genes IL1A and CXCL1 showed clear p53 binding in the LOXIMVI line, weaker

association in the UO31 and H460 lines, and no binding in other cell lines (Fig. 3B). We also found

variability in p53 binding at the promoters of previously reported p53 target genes, ALDH3A1and EPHA2,

ranging from no binding in some cell lines to strong peaks in others (Fig. 3B). *De novo* motif search on this

set of variable peaks identified the p53 binding site as the most significantly enriched motif (HOMER,

p=1.0e-46), suggesting that these sites represent direct p53 binding events.

To determine if these highly variable binding sites had novel cell line specific functions, we assigned each

peak to its closest gene (with a 10 kb cutoff) and clustered the resulting 218 peaks on their p53 binding.

We found that most cell lines showed a few unique p53 binding sites, but without strong clustering which

would sort the lines into groups (Fig. 3C) as in Fig. 1C.   Enrichment analysis identified

inflammatory/chemotaxis associated genes as being enriched in these highly variable p53 bound genes.

The cell line LOXIMVI showed particularly strong enrichment for p53 binding to inflammatory genes

including IL1A, IL1B, CLL20, and CXCL1 genes, although UO31 also showed substantial binding for many of

these targets. We also observed, that in the estrogen receptor (ER) positive MCF7 breast cancer cell line,

several unique binding sites overlapped with ESR1 (estrogen receptor) binding sites, including TFF1,

IGFBP4, and PRLH. These results suggest that the non-uniformities in p53 binding we observed may be

linked to cell line specific regulatory programs.

p53 binding correlates with basal gene expression

To explore the connection between cell line specific p53 binding and gene expression, we collected RNA

sequencing data from each cell line in the basal, untreated state and 3 hours after IR. In contrast to the

uniformity of p53 binding, we saw substantially different gene expression programs in response to IR

across cell lines.   Focusing on established p53 target genes [12] (list of target genes provided in

Supplement), we found that while some canonical p53 target genes were universally activated, others

showed more variable activation after IR. Notably, while CDKN1A/p21 showed fairly uniform induction in

all cell lines, MDM2 and BBC3/Puma showed variable activation despite having conserved p53 binding

(Fig. 4A; Fig. 1A). More generally, there was substantial variation in the activation of p53 targets after IR

(Fig. 4B, 77 genes).However we found that, cell lines originating from the same tissue were highly

correlated in their gene expression response to IR. For example, the two lung cancer cell lines (A549 and

H460) showed a correlation of R=0.90 in their p53 target gene response to IR (Fig. 4C) compared to low

or even negative correlation, between other cell line pairs as for example UO31 and H460 (R=-0.49, Fig.

4B,C).

We next asked if differential p53 binding could explain this variation in IR induced gene expression in

different cell lines. Quantitative variation in p53 binding was essentially uncorrelated (median R=0.01) to

the induction of p53 target genes after IR (Fig. 4D). However, for the highly variable p53 bound genes

(Fig. 2C), p53 binding was correlated with basal gene expression (p=1.9e-31, t-test; Fig. 4D, E). Taken

together, these results suggest that while the transcriptional response of cell lines to IR is diverse,

relatively little of this diversity is explained by p53 binding site occupancy. Conversely, basal gene

expression appears to correlate with the ability of p53 to bind to a subset of genes.

Cell line specific chromatin accessibility accounts for variability in p53 binding sites

Given our observation that basal gene expression correlated with cell line specific p53 binding events, we

wondered if the p53 peaks specific to some cell lines can be attributed to increased chromatin accessibility

in these lines. It has been suggested that p53 can act as a pioneer factor with a high affinity for histone

occupied regions [20-22], whereas others have shown that p53 binds readily in open regions [23, 24]. Our

results linking basal expression of nearby genes to p53 binding suggest that the 'openness' of the genomic

region might influence p53 binding. To study this we focused on the LOXIMVI cell line, which showed

strong, and unique binding of p53 nearby inflammatory related genes and the MCF7 cell line, which

showed p53 binding at estrogen receptor associated genes, and for which DNase accessibility data was

available. We performed a modified ATAC-Seq protocol using the MuA transposase to generate genome

wide maps of accessible regions in the MCF7 and LOXIMVI cell lines. We note that in this context the MuA

transposition occurs in a 5min 30C step, a potential advantage over the more conventional TN5 approach

that requires a 30min 37C incubation. Our ATAC-Seq data and ENCODE produced DNase sensitivity data

from MCF7 showed strong overlap with greater than 90% of ATAC-Seq peaks being DNase accessible [25].

We compared our ATAC-Seq data to the p53 ChIP-Seq signal for the inflammatory genes that showed p53

binding in LOXIMVI but not in MCF7, and observed strong ATAC-Seq signal only in the LOXIMVI cell line

(Fig. 5A), consistent with increased accessibility at these loci leading to stronger p53 binding. Conversely,

GREB1, a breast cancer associated gene showed only p53 binding and ATAC-Seq sensitivity in the MCF7

cell line (Fig. 5A). Genome wide the difference in ATAC-Seq signal between the two lines accounted for

just over 22% of the variance in p53 binding between the two datasets (R^2=0.225; Fig 5B). More generally,

as has been observed for other transcription factors [26], combining accessibility and motif scoring allows

for improved prediction of DNA binding. Indeed, accessibility and motif score accounted for 13.8% and

20.9% of the variance in the log2(p53 ChIP-Seq peak signal) for MCF7 and LOXIMVI respectively, compared

to ~5% with the motif alone. We therefore conclude that chromatin accessibility favors p53 binding and

accounts for a substantial fraction of the cell line specific gain of p53 DNA binding sites between MCF7

and LOXIMVI cells. Interestingly, we also found that genome wide chromatin accessibility was negatively

correlated with *in vitro* p53 binding (R=-0.2, p=2.1e-80, MCF7 ATAC-Seq vs. *in vitro* binding), suggesting

that many strong p53 binding sites are obscured by local chromatin context.

To explore if, within a single cell line, perturbations to accessibility could alter p53 binding we treated

MCF7 cells with decitabine, a demethylase inhibitor that has been shown to broadly alter chromatin

structure [27]. We then treated these cells with IR and preformed p53 ChIP-Seq and ATAC-Seq on the

samples.   Comparing p53 binding between the decitabine treated and untreated cells, we observed

minimal alterations in p53 binding (Fig. 5C). Indeed, we found only one binding site, adjacent to the

DLGAP5 gene, that showed a substantial change in p53 binding (Fig. 5D). This increase in p53 binding was

accompanied by increased accessibility (Fig. 5D). The DLGAP5 binding site has a consensus p53 motif and

showed occupancy in other cell lines such as UACC62 (Fig. 5D), consistent with this binding site being

obscured by chromatin in wild type MCF7 cells. Computing across p53 binding sites genome-wide, there

was a modest but significant correlation between change in chromatin accessibility and change in p53

DNA binding between decitabine treated and untreated samples (R=0.16, p=3.99e-13).

Discussion

The transcription factor p53 regulates the cellular response to DNA damage, including up regulating

repair, cell cycle arrest and apoptotic proteins. The nature, strength, and balance between the DNA repair

and cell death arms of p53 signaling varies across tissues in the body [6, 16, 28], and can be modified by

drug treatment [29, 30] and genetic perturbation [31]. The role of the p53 itself in this decision making is

controversial, with arguments for p53 behaving as a smart 'signal integrator' (reviewed in [8]) or a simple

effector [10]. We sought to understand the role of p53 in diverse cell lines by focusing on p53 DNA binding

and gene expression in response to ionizing radiation.

As other studies have suggested, p53 DNA binding does not greatly vary across cell lines or treatments.

However, we did find that p53 binding could group cell lines by their tissue of origin, suggesting some

degree of tissue specificity. Further, we noted a modest, but significant correlation between the strength

of p53 binding (measured by ChIP-Seq) and the predicted strength of p53 association (using the PWM

model). This correlation varied across cell lines and was strongest in the pooled dataset containing all cell

lines. More strikingly, we observed a similar correlation when comparing genome wide *in vitro* association

of p53 with *in vivo* p53 binding. These results suggest that the PWM motif is a relatively accurate measure

of p53 binding affinity, and that flanking regions or other DNA features do not greatly improve p53 binding

prediction. In general, p53 binding at any given location in the genome was relatively poorly predicted by

either *in vitro* binding or motif analysis suggesting that *in vivo* factors greatly contribute to p53 binding

specificity.

Taking advantage of the coherence of our dataset we identified p53 binding sites that were variably

occupied across cell lines. This subset of peaks were nearby genes enriched for specific cellular programs,

most notably the inflammatory response in the melanoma LOXIMVI cell line and ER specific response in

the MCF7 cell line. Our ATAC-Seq data showed that this differential p53 binding could be substantially

explained by chromatin accessibility in the MCF7 and LOXIMVI cell lines we examined. Globally, our data

showed that a higher degree of chromatin accessibility favored p53 binding and is consistent with a

previous report showing that open chromatin can provide a permissive environment for p53 [21] as well

as other transcription factors such as GR [32] for example. We tested this prediction by using a

pharmacological agent to modify p53 chromatin state and observed increased accessibility around the

DLGAP5 gene that correlated with increased p53 binding.

Looking at gene expression of p53 bound genes, we observed no direct correlation between variation in

p53 binding and induction of gene expression. This suggests that although p53 binding is required for the

DNA damage response, the extent and mode of the IR gene expression response, is dictated by other

factors than p53 binding itself. We recently showed that mRNA stability of direct p53 target genes can

lead do differences in the timing and level of expression [33]. Further studies coupling chromatin

accessibility, p53 binding, post-translational modifications, and measurements of RNA synthesis and

degradation rates will be required to reconcile these observations and identify what features tune the

cellular response to DNA damage in different cellular backgrounds.

Conclusions

We found that the vast majority of p53 binding events as universal across cancerous and non-cancerous

cell lines, with strong quantitative agreement in binding magnitude. We further found that Nutlin3A

treatment resulted in a nearly identical set of p53 binding events as IR, emphasizing the universality of

these binding sites. These binding sites were not, however, well predicted by the local genomic sequence.

We identified a set of variable p53 binding events (~5%) present in only one or a handful of cell lines.

These binding events were often in the vicinity of transcriptionally active genes and correlated strongly

with cell line specific chromatin accessibility. Consistent with this, pharmacological modification of

chromatin state could modify p53 binding. Interestingly, we did not find a strong correlation of p53 DNA

binding to the transcriptional response to IR, suggesting additional layers of transcriptional control.

Overall we found that the acute activation and binding of p53 to DNA in response to damage is highly

stereotyped, with modest tuning by the chromatin environment.

Methods

<u>General genomic analysis</u>

All DNA reads were single end Illumina reads and were aligned to HG19 genome build using bowtie [34].

Aligned reads were further processed using HOMER (V4.6, [35]) to assemble tag files and call peaks. Peak

locations and tag files were read by custom Matlab code (Mathworks) which was used to integrate the

different datasets/types. For each ChIP-Seq dataset the number of reads in p53 peaks were normalized

to the average of all cell lines, and for subsequent analyses and comparisons, peaks with less than 2

normalized counts were discarded. Clustering and comparison was based on Pearson correlations

between p53 binding signals. RNA data was aligned to hg19 and the Refseq HG19 transcriptome using

Tophat, CuffQuant, and CuffMerg [36]. Most computation was done on the Harvard Medical School

cluster (Orchestra). Genomic binding and signals were visualized using the UCSC genome browser [37].

Motif analysis was performed in Matlab on the Hg19 genome using a ChIP-Seq derived PWM adjusted to

have a minimum probability of each nucleotides occurrence.

<u>ATAC-Seq</u>

ATAC seq was performed as described [38], with the major exception of the use of a MuA transposase

(Thermo) rather than the TN5 transposase. Briefly, MCF7 or LOXIMVI cells were trypsinized and 50K cells

spun down, washed once with PBS, and lysed with a hypotonic buffer containing 0.1% NP-40, and spun

down to generate a crude nuclei pellet. This pellet was transposed in a 30µl volume using MuA (0.7µl),

MuA buffer (10µl), and H2O (19µl) for 5min at 30C. The sample was treated with 3µl stop solution, and

incubated at 30C for a further minute. The sample was then collected and purified by addition of 45µl of

SPRI beads (Aline Biosciences). The purified sample was PCR amplified in two steps to add barcoded

adaptors suitable for Illumina sequencing. Samples were sequenced with single end 75bp reads on an

Illumina NextSeq. Reads (>30M) were trimmed to remove adaptors with cutadapt [39], aligned to the

genome with Bowtie, and analyzed with Matlab. Genomic DNA (50ng) from MCF7 and LOXIMVI was

transposed, amplified and sequenced in parallel to estimate background.

<u>ChIP-Seq</u>

Briefly, 10M cells were treated with 4Gy IR (RS-2000, RadSource) and 2hrs later were fixed by addition of

1% paraformeldahyde (Alfa Aesar) at room temperature for 10 minutes with agitation. Fixation was

stopped by addition of 250mM glycine. Cells were scraped and flash frozen. Cell pellets were thawed in

hypotonic lysis buffer and spun to generate a crude nuclei prep. These nuclei were lysed in a SDS buffer

and sonicated (Bioruptor) to fragment DNA. Fragmented DNA was diluted in IP buffer and agitated

overnight with 2mg/ml DO-1 (anti-p53, Santa Cruz). 20µl of protein A magnetic beads (Invitrogen) were

used to isolate the p53 associated fragments and samples were washed with low salt, high salt, and LiCl

buffers. DNA was eluted from beads with an SDS/NaCO3 buffer and was decross-linked at 65C for 6hrs in

a high salt buffer. ChIP libraries were constructed with the commercial NEBnext kit (NEB) and associated

protocols, although reaction volumes were reduced by 4 fold and custom adaptors and barcodes were

employed. Libraries were sequenced with single end 75bp reads on Illumina NextSeq.

Reads were aligned to the genome with Bowtie1.1 [34], and analyzed with HOMER [35], MACS2 [40] and

custom Matlab scripts. Peak calling was done after pooling reads from ChIP-Seq experiments in all cell

lines. The final set of peaks represented the consensus of HOMER (default settings) and MACS2 (using the

q 0.01 threshold) identified peaks, and was filtered to remove ENCODE black-list locations. The number

of reads within each peak region was computed from HOMER tag files using custom Matlab scripts.

Background regions around each peak were subtracted from peak scores to correct for high background

regions. The HOMER package [35] was used for *de novo* motif discovery. WebLogo was used to generate

the motif plot [41] in (Figs. 1B, 2C) for the top enriched motif. The top enriched motif (Fig. 1B) was then

used to re-scan and score all peaks and background regions. Background regions were generated by

selecting 500bp regions adjacent to either side of the peak and excluding regions that overlap with p53

peak regions. Clustering of peaks was accomplished using a correlation distance metric and average

linkage in Matlab.

In vitro ChIP-Seq

To generate recombinant p53 we in vitro transcribed/translated human p53 with a c-terminal HA tag using

a rabbit reticulocyte system (Promega). To generate fragmented genomic DNA we tagmented 50ng of

human genomic DNA from MCF7 cells using the MuSeq kit (Thermo) and amplified it using PCR and custom

adaptor primers for 8 cycles. DNA was cleaned up on SPRI beads (Aline Biosciences) and quantified. At

room temperature 20ng of DNA and recombinant p53 (0.1uM final) were combined in a binding buffer

(10mM TRIS, 5mM MgCl2, 10% glycerol, 1mM DTT) and incubated at room temperature for 30 min. The

mixture was diluted 2 fold (to 20ul) and 1.5ul of anti-HA antibody was added (Rockland) and the sample

incubated at 4C overnight with shaking. A 1:1 mixture of magnetic proteinA/proteinG beads was added

(sigma) and incubated at 4C for 1hr with shaking. The beads were then washed 3x with washing buffer

(10mM Tris, 5mM HCL, 0.1% triton, 150mM NaCl) and DNA eluted with elution buffer (1%SDS, 100mM

Na2CO3) at 37C for 15 minutes. Samples were cleaned up, and adaptors and barcodes added by PCR.

Reads (>30M) were trimmed to remove adaptors with cutadapt [39], aligned to the genome with Bowtie,

and analyzed with Matlab.

RNA-Seq

For each cell line 50,000 cells were plated in 35 mm dishes, 24hrs later cells were treated (or not) with

4Gy IR (RS-2000, RadSource), 3hrs after that cells were lysed with Trizol (Ambion). RNA was purified on

affinity columns and DNAse treated (Zymo). Purified RNA (500ng) was polyA purified using magnetic

beads (NEB), fragmented and reverse transcribed using protoscript RT (NEB), second strand synthesized

(NEB), and then assembled into libraries with the commercial NEBnext kit (NEB) and associated protocols,

although reaction volumes were reduced by 4 fold and custom adaptors and barcodes were employed.

Libraries were sequenced with single end 75bp reads on a NextSeq. Reads were aligned to the

genome/transcriptome with Bowtie/Tophat and analyzed with custom Matlab scripts.

Cell Culture and Cell Treatment

Parental cell lines were obtained from ATTC, with the exception of the RPE cells which were a gift from

Prof. Steve Elledge (Harvard Medical School), and were thawed and propagated in RPMI (GIBCO) with 5%

FBS. All experiments were performed in this media. All media was supplemented with 1% antibiotic and

antimycotic (Corning). Treatment with Nutlin3A (sigma) was at 5uM. X-ray induced DNA damage was

generated with a RS-2000 source (RadSource, 160KeV). MCF7 cells were treated with 2uM (5-AZA-2'-

deoxcytidine, MP Biomedicals) for 5 days, cells were split on day 2, replated in decitabine containing

media. Treated and untreated cells were then further treated with IR or not as with other samples.

Public datasets

Raw fastq datasets were downloaded from the Sequence Read Archive (see supplement). These datasets

were all single end Illumina reads and were aligned to the HG19 genome with using the same pipeline as

described above for our ChIP-Seq samples, and further analyzed with HOMER to generate tag files.

Custom Matlab code was used to compare these datasets to our ChIP-seq data.

Declarations

Ethics approval and consent to participate

Not Applicable.

Consent for publication

Not Applicable

Availability of data and material

All sequencing data has been deposited in NCBI's Gene Expression Omnibus under accession number

GSE100292. Data is also available as UCSC tracks as a custom session accessible at:

https://genome.ucsc.edu/cgi-

bin/hgTracks?hgS_doOtherUser=submit&hgS_otherUserName=jacobso&hgS_otherUserSessionName=H

afner_et_al_hg19

Competing interests

The authors declare no conflicts of interest.

Funding

AH was supported by a Boehringer Ingelheim Fonds PhD fellowship. This study was supported by NIH

grants GM083303 and CA207727 (J.S-O).

Authors' contributions

JSO and AH initiated and conceived the study; JSO and AH collected and analyzed the data; AH, JSO and

GL wrote the paper, JSO and GL supervised the study.

Acknowledgements

We thank Megha Padi and all members of the Lahav lab for feedback. We thank the Bauer sequencing

core and Christian Daly for sequencing support and HMS research computing for maintaining the

Orchestra cluster.

References

1.   Vaquerizas, J.M., et al., *A census of human transcription factors: function, expression and*

evolution. Nat Rev Genet, 2009. **10**(4): p. 252-63.

2.   Chen, X., et al., *Integration of external signaling pathways with the core transcriptional network*

in embryonic stem cells. Cell, 2008. **133**(6): p. 1106-17.

3. Boyer, L.A., et al., *Core transcriptional regulatory circuitry in human embryonic stem cells.* Cell,

2005. **122**(6): p. 947-56.

4. Gerstein, M.B., et al., *Architecture of the human regulatory network derived from ENCODE data.*

Nature, 2012. **489**(7414): p. 91-100.

5. Garber, M., et al., *A high-throughput chromatin immunoprecipitation approach reveals principles*

of dynamic gene regulation in mammals. Mol Cell, 2012. **47**(5): p. 810-22.

6. Fei, P., E.J. Bernhard, and W.S. El-Deiry, *Tissue-specific induction of p53 targets in vivo.* Cancer Res,

2002. **62**(24): p. 7316-27.

7. Brady, C.A. and L.D. Attardi, *p53 at a glance.* J Cell Sci, 2010. **123**(Pt 15): p. 2527-32.

8. Kruiswijk, F., C.F. Labuschagne, and K.H. Vousden, *p53 in survival, death and metabolic health: a*

lifeguard with a licence to kill. Nat Rev Mol Cell Biol, 2015. **16**(7): p. 393-405.

9. Aylon, Y. and M. Oren, *The Paradox of p53: What, How, and Why?* Cold Spring Harb Perspect Med,

2016. **6**(10).

10. Verfaillie, A., et al., *Multiplex enhancer-reporter assays uncover unsophisticated TP53 enhancer*

logic. Genome Res, 2016. **26**(7): p. 882-95.

11. Nikulenkov, F., et al., *Insights into p53 transcriptional function via genome-wide chromatin*

occupancy and gene expression analysis. Cell Death Differ, 2012. **19**(12): p. 1992-2002.

12. Fischer, M., *Census and evaluation of p53 target genes.* Oncogene, 2017.

13. Botcheva, K., et al., *Distinct p53 genomic binding patterns in normal and cancer-derived human*

cells. Cell Cycle, 2011. **10**(24): p. 4237-49.

14. Smeenk, L., et al., *Role of p53 serine 46 in p53 target gene regulation.* PLoS One, 2011. **6**(3): p.

e17574.

15.   Akdemir, K.C., et al., *Genome-wide profiling reveals stimulus-specific functions of p53 during*

differentiation and DNA damage of human embryonic stem cells. Nucleic Acids Res, 2014. **42**(1):

p. 205-23.

16.   Stewart-Ornstein, J. and G. Lahav, *p53 dynamics in response to DNA damage vary across cell lines*

and are shaped by efficiency of DNA repair and activity of the kinase ATM. Sci Signal, 2017.

10(476).

17.   Gordan, R., et al., *Genomic regions flanking E-box binding sites influence DNA binding specificity*

of bHLH transcription factors through DNA shape. Cell Rep, 2013. **3**(4): p. 1093-104.

18.   Rohs, R., et al., *The role of DNA shape in protein-DNA recognition.* Nature, 2009. **461**(7268): p.

1248-53.

19.   Bartlett, A., et al., *Mapping genome-wide transcription-factor binding sites using DAP-seq.* Nat

Protoc, 2017. **12**(8): p. 1659-1672.

20.   Lidor Nili, E., et al., *p53 binds preferentially to genomic regions with high DNA-encoded*

nucleosome occupancy. Genome Res, 2010. **20**(10): p. 1361-8.

21.   Su, D., et al., *Interactions of chromatin context, binding site sequence content, and sequence*

evolution in stress-induced p53 occupancy and transactivation. PLoS Genet, 2015. **11**(1): p.

e1004885.

22.   Sammons, M.A., et al., *TP53 engagement with the genome occurs in distinct local chromatin*

environments via pioneer factor activity. Genome Res, 2015. **25**(2): p. 179-88.

23.   Tebaldi, T., et al., *Whole-genome cartography of p53 response elements ranked on transactivation*

potential. BMC Genomics, 2015. **16**: p. 464.

24.   Allen, M.A., et al., *Global analysis of p53-regulated transcription identifies its direct targets and*

unexpected regulatory mechanisms. Elife, 2014. **3**: p. e02200.

25.   ENCODE, *An integrated encyclopedia of DNA elements in the human genome.* Nature, 2012.

426   **489**(7414): p. 57-74.

427   26.   Zhong, S., X. He, and Z. Bar-Joseph, *Predicting tissue specific transcription factor binding sites.*

428   BMC Genomics, 2013. **14**: p. 796.

429   27.   Ponnaluri, V.K.C., et al., *NicE-seq: high resolution open chromatin profiling.* Genome Biol, 2017.

430   **18**(1): p. 122.

431   28.   Gudkov, A.V. and E.A. Komarova, *The role of p53 in determining sensitivity to radiotherapy.* Nat

432   Rev Cancer, 2003. **3**(2): p. 117-29.

433   29.   Purvis, J.E., et al., *p53 dynamics control cell fate.* Science, 2012. **336**(6087): p. 1440-4.

434   30.   Paek, A.L., et al., *Cell-to-Cell Variation in p53 Dynamics Leads to Fractional Killing.* Cell, 2016.

435   **165**(3): p. 631-42.

436   31.   Sullivan, K.D., et al., *ATM and MET kinases are synthetic lethal with nongenotoxic activation of*

437   *p53.* Nat Chem Biol, 2012. **8**(7): p. 646-54.

438   32.   John, S., et al., *Chromatin accessibility pre-determines glucocorticoid receptor binding patterns.*

439   Nat Genet, 2011. **43**(3): p. 264-8.

440   33.   Hafner, A., Stewart-Ornstein, J., Purvis, J.E., Forrester, W.C., Bulyk, M.L., Lahav, G, *p53 pulses lead*

441   *to distinct patterns of gene expression albeit similar DNA binding dynamics.* Nature Structural &

442   Molecular biology, 2017. **In press**.

443   34.   Langmead, B., et al., *Ultrafast and memory-efficient alignment of short DNA sequences to the*

444   *human genome.* Genome Biol, 2009. **10**(3): p. R25.

445   35.   Heinz, S., et al., *Simple combinations of lineage-determining transcription factors prime cis-*

446   *regulatory elements required for macrophage and B cell identities.* Mol Cell, 2010. **38**(4): p. 576-

447   89.

36.   Trapnell, C., et al., *Differential gene and transcript expression analysis of RNA-seq experiments*

with TopHat and Cufflinks. Nat Protoc, 2012. **7**(3): p. 562-78.

37.   Kent, W.J., et al., *The human genome browser at UCSC.* Genome Res, 2002. **12**(6): p. 996-1006.

38.   Buenrostro, J.D., et al., *Transposition of native chromatin for fast and sensitive epigenomic*

profiling of open chromatin, DNA-binding proteins and nucleosome position. Nat Methods, 2013.

10(12): p. 1213-8.

39.   Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing reads.*

EMBnet.journal, 2011. **17**(1): p. 10-12.

40.   Zhang, Y., et al., *Model-based analysis of ChIP-Seq (MACS).* Genome Biol, 2008. **9**(9): p. R137.

41.   Crooks, G.E., et al., *WebLogo: a sequence logo generator.* Genome Res, 2004. **14**(6): p. 1188-90.

42.   Zeron-Medina, J., et al., *A polymorphic p53 response element in KIT ligand influences cancer risk*

and has undergone natural selection. Cell, 2013. **155**(2): p. 410-22.

467   **Figure Legends**

Fig. 1: Stereotyped p53 binding across twelve cell lines. (A) ChIP-Seq for p53 in 12 p53 wild-type cell lines. UCSC screen shots of p53 binding sites for three canonical p53 target genes are shown. (B) Motif analysis recovered a p53 motif as the single most enriched sequence. Centrality analysis showed that this motif was also centrally enriched within peaks. (C) Heatmap showing p53 binding intensity in 8742 locations in the genome. Cell lines were clustered on p53 binding and resulted in lineages clustering together. (D) Comparison of p53 binding in two cancer cell lines (UACC62 and UACC257) as well as between one cancer (UACC257) and one non-cancerous cell line (RPE1). (E) Comparison of p53 binding between Nutlin3A and IR treated samples in MCF7 or UACC257 cells.

Fig. 2: Genomic sequence is weakly predictive of p53 binding. (A) The correlation between motif strength and p53 binding is shown as a function of the number of cell lines across which the peak height was averaged, box plot represent the distribution of correlations across all possible cell line combinations. (B) The degree to which p53 binding motif predicts the strength of p53 binding is shown in a box plot, with p53 binding sites binned by their motif strength. (C) The top enriched motif identified by *in vitro* ChIP. (D) UCSC screen shots of p53 binding sites, motif score, and *in vitro* p53 binding signal are shown for CDKN1A/p21, MDM2, and MDM4. (E) I*n vivo* p53 binding strength is shown in a box plot, binned by *in vitro* p53 binding signal at each genomic site.

Fig. 3: Variable p53 binding sites show cell type specific functional enrichment. (A) Scatterplot of all 8742 p53 binding sites by their average ChIP signal and coefficient of variation (CV). Highlighted in red are 'highly variable' peaks defined as having higher than expected CV relative to the peak height. Example binding sites are labeled with the associated gene names. (B) UCSC screen shots of four example 'variable' peaks. (C) Heatmap of 'variable' p53 peaks that are also nearby (<10kb) transcription start sites of genes. The intensity of each peak is normalized to the average across 12 cell lines. Cell lines and peaks were hierarchically clustered, with no grouping by lineage observed for cell lines. Groups of inflammatory and the ER associated are highlighted.

Fig. 4: Basal gene expression, but not DNA damage induced gene expression, correlates with p53

binding. (A) Box plots of the fold change of three canonical p53 target genes 3h after IR. Each dot

represents a cell line. CDKN1A/p21 is induced in all cell lines, while MDM2 and BBC3/Puma is cell line

dependent. (B) Correlation of 12 cell lines by p53 target gene (77 genes) induction after DNA damage. (C)

Scatter plots of p53 target genes comparing the highly correlated lung cancer lines (A549 and H460, upper)

and negatively correlated lung and kidney lines (H460 and UO31, lower). (D) Boxplots showing the

correlation of either basal or DNA damage induced fold change of gene expression with p53 binding at

the p53 target or variable p53 binding gene sets (Fig. 3). (E) Scatterplots illustrating the relationship

between basal gene expression and p53 binding across the 12 cell lines for four p53 peaks/genes (note

that in many cases multiple cell lines show little gene expression or p53 binding and therefore cluster near

the origin).

Fig. 5 Chromatin accessibility contributes to variable p53 binding. (A) UCSC screen shots of two p53

binding sites; p53 binding in the proximity of GREB1 is found in MCF7 treated with IR, while IL1A was

bound in IR treated LOXIMVI cells. ATAC-Seq data and published DNase hypersensitivity data (for MCF7,

untreated) showing that IR induced p53 binding correlates with basal DNA accessibility in each cell line.

(B) Scatter plot of p53 binding post IR in MCF7 compared to LOXIMVI, colored by the difference in ATAC-

Seq signal log2(LOXIMVI-MCF7) between the two cell lines. (C) Scatter plot of p53 binding in IR treated

MCF7 cells compared to MCF7 treated with IR and decitabine, colored by the difference in ATAC-Seq signal

between decitabine treated and untreated cells: log2(decitabine-not treated). (D) UCSC screen shot of

region around the gene DLGAP5, showing changes in p53 binding and accessibility in the decitabine

treated MCF7 cells (the new peak is indicated by an arrow). Binding of p53 in IR treated UACC62 cells at

the DLGAP5 locus without decitabine treatment is also shown.

Supplement

<u>Public p53 ChIP-Seq datasets</u>

SRR048928, SRR048929 – U2OS cells: Actinomycin D or Etoposide treated [14]

SRR287798, SRR287799, SRR287800 – MCF7: Nutlin3A, RITA, 5-FU [11]

SRR575903, SRR575904, SRR575905 – hESC : RA or Doxorubicin [15]

SRR851807, SRR851807, SRR851807 -- lymphoblastoid cell lines: doxorubicin [42]

<u>p53 Target Genes used in this analysis</u>

This gene list is based on the list of [12] and removing genes with low read coverage in our dataset.

CDKN1A	HSPA4L	ZNF337
RRM2B	ISCU	ZNF79
MDM2	PHLDA3	CERS5
GDF15	SLC12A4	CSF1
SUSD6	TRAF4	DUSP14
BTG2	CCDC90B	EPS8L2
DDB2	CES2	FAM210B
GADD45A	DYRK3	FUCA1
PLK3	KITLG	IER5
RPS27L	NADSYN1	IKBIP
TNFRSF10B	NTPCR	MICALL1
TRIAP1	ORAI3	PMAIP1
BAX	SESN2	RAP2B
BLOC1S2	SLC30A1	RNF19B
POLH	TM7SF3	SAC3D1

PPM1D	TMEM68	TNFRSF10D
SULF2	ANXA4	
XPC	APOBEC3C	
AEN	ASCC3	
ANKRA2	BBC3	
NINJ1	DCP1B	
PLK2	EPHA2	
SERTAD1	FDXR	
SESN1	FOSL1	
TP53I3	LIF	
CCNG1	PGPEP1	
CMBL	PRKAB1	
CYFIP2	PTP4A1	
DRAM1	TGFA	
FBXO22	ZNF219	

522 **Supplementary Figure Legends**

523 **Fig. S1: Comparison of p53 DNA binding with published p53 ChIP-Seq datasets.** Heatmap showing p53

524 binding intensity in 8742 locations in the genome in 12 IR treated cell lines (same as Fig. 1C) as well as

525 published datasets.

Figure 1

A

p53 ChIPseq signal

CDKN1A/p21   MDM2   BBC3/Puma

A498
A549
H460
HCT116
LOXIMVI
MALM3E
MCF7
SKMEL5
U2OS
UACC257
UACC62
UO31

B

bits

p53 peaks
background

# of peaks

[bp] from peak center

C

Sorted Peaks

Peak Height

UO31
A498   Kidney
A549   Lung
H460
HCT116
LOXIMVI
UACC62
UACC257   Melanoma
MALM3E
SKMEL5
U2OS
MCF7

Correlation

D

UACC62 p53 ChIP signal
R=0.643
UACC257 p53 ChIP signal

UACC257 p53 ChIP signal
R=0.46
RPE1 p53 ChIP signal

E

p53 Nut ChIP signal
MCF7
R=0.866
p53 IR ChIP signal

p53 Nut ChIP signal
UACC257
R=0.878
p53 IR ChIP signal

Figure 2

A

correlation of ChIP signal and motif

number of cell lines

B

log2 average ChIP signal (AU)

log10 motif strength (PWM)

C

bits

D

CDKN1A/p21   MDM2   MDM4

A549 p53
UO31 p53
UACC257 p53
Motif Score
In vitro p53

10 kb   10 kb   10 kb

E

ChIP signal (AU)

log10 *in vitro* p53 signal

Figure 3

A

B

C

Figure 4

Figure 5

Figure S1

H460 -2hr 4Gy
HCT116 -2hr 4Gy
A549 -2hr 4Gy
MCF7 -2hr 4Gy
UO31 -2hr 4Gy
UACC62 -2hr 4Gy
U2OS-2hr 4Gy
A498-2hr 4Gy
UACC257 -2hr 4Gy
SKMEL5 -2hr 4Gy
LOXIMVI -2hr 4Gy
MALM3E -2hr 4Gy
U2OS -ActD
U2OS -Etop
HCT116 -5FU
MCF7 -RITA
MCF7 -5FU
MCF7 -Nutlin
hESC -Doxo
hESC -RA
LCL -Doxo
LCL -IR

Our data   Published data