microRNA tissue atlas of the malaria mosquito *Anopheles gambiae*

Lena Lampe[1] and Elena A. Levashina[1]

[1] Max Planck Institute for Infection Biology, Berlin

Corresponding author: levashina@mpiib-berlin.mpg.de

Keywords: *Anopheles gambiae*, *Plasmodium falciparum*, microRNAs, reproductive cycle

ABSTRACT

Anopheles gambiae mosquitoes transmit the human malaria parasite *Plasmodium falciparum*, which causes the majority of fatal malaria cases worldwide. The hematophagous life style defines the mosquito reproductive biology and is exploited by *P. falciparum* for its own sexual reproduction and transmission. The two main phases of the mosquito reproductive cycle, pre-vitellogenic (PV) and post-blood meal (PBM) shape its capacity to transmit malaria. Transition between these phases is tightly coordinated to ensure homeostasis between mosquito tissues and successful reproduction. One layer of control is provided by microRNAs, well-known regulators of blood meal digestion and egg development in mosquitoes. Here, we report a global overview of tissue-specific miRNA expression during the PV and PBM phases and identify miRNAs regulated during PV to PBM transition. The observed coordinated changes in the expression levels of a set of miRNAs in the energy-storing tissues suggest a role in the regulation of blood meal-induced metabolic changes.

INTRODUCTION

Anopheles gambiae s.l. females are major vectors of the human malaria parasite *Plasmodium falciparum* in sub-Saharan Africa. The parasite exploits the mosquito's reproductive demand for blood feeding as a route of transmission, which tightly links the mosquito infection and its reproductive (also called gonotrophic) cycle. As reproduction is central to every organism, the hematophagic life style directs mosquito behavior, physiology, and metabolism. The gonotrophic cycle is divided into two phases – the pre-vitellogenic (PV) and post blood meal (PBM) phase. During the PV phase, egg maturation (vitellogenesis) is arrested at a premature stage until the blood meal-mediated uptake of amino acids and lipids (1–3). The PV phase, controlled by the juvenile hormone (JH), prepares the carbohydrate-consuming female mosquitoes for blood feeding by accumulating energy resources in the fat body, the major storage tissue in insects, and by activating their host-seeking behavior (4–9). Successful blood feeding initiates the PBM phase, which is driven by the steroid hormone ecdysone (10,11). Digestion of the blood nutrients in the midgut generates lipids, sugars, and amino acids that allow ovary development and egg maturation (12,13). Within 48 h, a blood meal is fully digested, nutrients absorbed, and oogenesis is completed. However, a blood meal may also bring *Plasmodium* infective stages or gametocytes that fuse in the mosquito midgut to produce motile ookinetes (14). The ookinetes traverse the midgut epithelium at 18-24 h post BM and establish an infection at the basal side of the midgut wall (14). Here within the next ten days, the parasites undergo replication and maturation of the infective forms (15). Therefore, the PBM phase shapes both mosquito reproduction and *P. falciparum* development, two crucial components of malaria transmission. Repeated blood feedings increase female reproductive fitness by boosting its nutritional resources

and promote *P. falciparum* development by providing key nutrients and a route for

transmission to the next host.

The metabolic transition from the PV to PBM phase entails a complex regulatory

network of hormonal, transcriptional and post-transcriptional changes in multiple

tissues. However, the mechanisms of tissue homeostasis during this transition

remain incompletely understood. Recently, microRNAs (miRNAs) emerged as

powerful regulators of developmental and metabolic switches. miRNAs fine tune JH

and ecdysone pulses and, thereby, shape developmental transitions in the fruit flies

(16–18). In mosquito adults, miR-8, miR-309, miR-275, miR-1174 and miR-1890

have been implicated in the regulation of PBM (19–24), and miR-305 in parasite

development (25). However, only three of ~180 miRNAs identified by bioinformatics

and experimental approaches have been functionally characterized in *A. gambiae*

(24,26–28). The functional diversity of miRNAs is further amplified by the capacity of

each miRNA locus to generate two miRNAs arms, which differ in their seed sequence

and target distinct sets of mRNAs. Several studies observed a bias in arm selection

for different species and tissues, suggesting that it might represent an additional layer

of miRNA regulation (26,29–31). In addition, the same miRNA can regulate distinct

processes in a tissue-specific manner by binding different targets. For example, the

Drosophila miR-8 regulates the production of myogenic peptide hormone in the fat

body, whereas it controls synapse structure in the brain (32,33). So far, all studies on

A. gambiae miRNAs focused either on whole mosquitoes or on few selected tissues

(19,24–28). However, a comprehensive overview of tissue-specific miRNA

expression would be beneficial for further characterization of miRNA function during

PV to PBM transition.

Here, we report a global overview of miRNA expression in the head, midgut,

abdominal carcass (the fat body) and ovaries of *A. gambiae* females during the

transition from the PV to PBM phase, and after *P. falciparum*-infection. We show that

the majority of mosquito miRNAs are tissue-specific, except for a small cluster of

ubiquitously expressed miRNAs. Further, we identify four miRNAs, whose expression

levels are regulated by blood feeding in the fat body, ovaries, and midgut. Our study

provides the first comprehensive miRNA tissue atlas of the female *A. gambiae*

mosquito during the gonotrophic cycle and identifies miRNAs with potential roles in

this critical process.

METHODS
83

Mosquitoes
84

85 *Anopheles coluzzii* Ngousso (*TEP1*S1*) mosquitoes were reared at 30°C 80%

86 humidity at 12/12 h day/night cycle with a half-hour long dawn/dusk periods. All

87 mosquitoes were fed *ad libitum* with 10% sugar solution.

Blood meal and *P. falciparum* infections
89

Blood meals and *P. falciparum* infections were performed using membrane feeders

(34,35). To establish gametocyte cultures, *Pf* NF54 asexual cultures (parasitaemia

>2%) were harvested by centrifugation for 5 min at 1,500 rpm, washed with fresh red

blood cells and diluted to 1% total parasitaemia in complete gametocyte medium at

4% hematocrit. Gametocyte cultures were incubated at 37°C with 3% O_2 and 4%

CO_2. Medium was changed daily for 15-16 days on heated plates to reduce

temperature drops. On day 14 after establishment, (i) gametocytaemia was checked

by Giemsa stain of gametocyte cultures-smears, and (ii) parasite exflagellation rates

were estimated by microscopy.

Mosquitoes were fed with either uninfected or *P. falciparum*-infected blood using an

artificial feeder system as described in Lensen *et al.* (34,35). The feeding system was

prepared by covering the bottom of the midi-feeders with a stretched parafilm. The

midi-feeder was then attached to a 37°C water bath system, allowing water flow

through the feeder. The blood (Haema, Berlin, Germany) was introduced to the

feeder and mosquitoes were fed for 15 min. Unfed mosquitoes were removed and

only fully engorged females were kept for further analyses.

Sample collection

Microarray and qPCR

Three-day-old virgin females were divided into three groups and fed with: (i) 10%

sugar (sugar-fed; PV phase); (ii) human blood (blood-fed; PBM phase); (iii) *P.*

falciparum-infected blood (infected; PBM phase). After feeding, all groups were kept

at 26°C 80% humidity. At 18 h post feeding, 15 females per group were dissected on

ice and the heads, midguts, ovaries and abdominal carcasses were pooled according

to the tissue. The samples were homogenized in TRIzol using a beat-beater (Qiagen)

at 50 rpm. Homogenized samples were kept at -80°C until further usage.

Time course

For miRNA expression kinetics, 15 unfed and 15 blood-fed females were dissected

on ice at 20, 24, 28, 32 and 48 h after blood meal. The heads, midguts, ovaries and

abdominal carcasses were pooled according to the tissue, immediately homogenized

in TRIzol and kept at -80°C for RNA isolation.

RNA isolation

Microarray

Total RNA was isolated using the miRNeasy Mini Kit (Qiagen) according to the manufacturer's recommendations. Briefly, after phase separation by centrifugation total RNA was purified by silica membrane, eluted with water and stored at -80°C. The total RNA yield was measured with a NanoDropND-1000 Spectrophotometer. The integrity of total RNA was assessed with a 2100 Bioanalyzer and a RNA 6000 Nano LabChip kit (Agilent). Furthermore, the ratio of miRNA/small RNAs from isolated total RNA was monitored by the Agilent Small RNA kit.

qPCR

Total RNA was isolated by TRIzol according to the manufacturer's recommendations. The total RNA yield was measured with a NanoDropND-1000 spectrophotometer.

Microarray analyses

Custom 8-plex 60K mosquito miRNA microarrays (Design Name: Agilent-049943, ID Name: Custom_Mosquito_miRNA, Design Format: IS-62976-8-V2, AMADID 016436) (Agilent) were used for one-color hybridizations. The microarray included miRNAs identified in *Apis mellifera*, *Anopheles gambiae*, *Tribolium castaneum*, *Culex quinquefasciatus*, *Aedes aegypti*, *Bombyx mori* and *Drosophila melanogaster.* Total RNA (100 ng) was processed with the miRNA Complete Labeling and Hyb Kit (Agilent) according the supplier's recommendations. In brief, samples were dephosphorylated with Calf Intestine Alkaline Phosphatase and labeled with Cy3 in a

T4 RNA ligase-mediated reaction with 3', 5'-cytidine bisphosphate (Cy3-pCp). The

labeling reaction was column-purified, vacuum-dried using a speed-vac at 45°C and

resuspended in a blocking and hybridization buffer. After hybridization, the

microarrays were washed, scanned at 5 µm resolution with a G2565CA high-

resolution laser microarray scanner, and features were extracted. Results were

analyzed by the R limma package. Briefly, signal intensities were corrected for the

background using the normexp function and quantile normalized. All spot replicates

were pooled using the avereps function producing an Elist containing log2 relative

expression data. Values of the negative control spots were subtracted from the mean

intensities and all negative probes were removed. Expression values were calculated

as mean values of the dipteran miRNA expression of the four biological replicates. As

the microarray contained miRNAs of multiple insect species including miRNA

orthologues, the probe sequences were blasted against the annotated *A. gambiae*

miRNAs and all probes with at least 14 nucleotides complementarity were included in

further analysis. In total, signals were detected for 86 miRNAs (Table S1).

Differentially regulated miRNAs were identified by fitting a linear regression model to

the data and empirical Bayes statistics in R.

cDNA synthesis and qPCR

Total RNA was reverse-transcribed with the miScript Kit (Qiagen). Expression levels

of mature miRNAs and of mRNAs were measured using the Quantitect SYBR Green

PCR Kit (Qiagen). Relative quantities of miRNA expression were normalized to the

gene encoding ribosomal protein S7 (*RPS7*). The miRNA primers were obtained as

miRNA Primer Assay (Qiagen).

RESULTS

We set out to develop a global overview of tissue-specific miRNA expression in the head, midgut, ovary and fat body, during the gonotrophic cycle of *A. gambiae* and after *P. falciparum* infections. To this end, we determined miRNA expression levels using the Agilent custom-designed miRNA microarray in the three groups of females fed with: (1) sugar; (2) human blood or (3) *P. falciparum*-infected blood (Fig. 1).

MicroRNA expression in the pre-vitellogenic phase

We first examined expression levels of miRNAs in the pre-vitellogenic (PV) phase. Microarray analysis detected expression of 74 out of 86 miRNA probes in at least one tissue. The microarray probes matched almost all *A. gambiae* miRNAs annotated in the miRBase, as well as the highly conserved insect miRNAs (Table S1). High levels of expression across all tissues were detected only for miR-8, whereas nine miRNAs (miR-1, miR-2, let-7, miR-34, miR-125, miR-184, miR-277, miR-306, and miR-957) had moderate to high levels of expression in at least two tissues (Fig. 2). A second cluster of 18 miRNAs included tissue-specific miRNAs with intermediate expression levels. The largest number of tissue-specific miRNAs (cut-off 1.5-fold) was observed in the head (17 miRNAs), including the well-known brain-enriched miR-7 and miR-124. Three miRNAs, miR-281, miR-1174 and miR-1175, were enriched in the midgut, whereas miR-10, miR-92b, miR-279, miR-989, and miR-998 were predominantly detected in the ovaries. Surprisingly, not a single miRNA was exclusively expressed in the fat body. Microarray results were further validated by independent quantitative PCR (Fig. 3A-D).

The mature miRNA can be generated from the 3p or 5p arm of the pre-miRNA. We

used arm-specific probes to quantify tissue-specific expression of miRNA arms and

identified 21 miRNAs expressing both arms (Table S2). Among those, six miRNAs

showed tissue-dependent arm bias (Fig. 3E). Head-specific enrichment in the 3p arm

was observed for miR-124 and miR-210, whereas the 5p arm bias was detected for

miR-1175 in the midgut and for miR-263 in the head, fat body and ovaries.

Furthermore, higher levels of the 5p arm were detected in the head and midgut for

miR-10. A complex pattern of arm usage was observed for miR-281. While the 5p

arm was enriched in the head and midgut, the 3p arm was predominantly expressed

in the fat body and ovary. We concluded that during the PV phase, expression of the

majority of the miRNAs is tissue-specific and, in some cases, is regulated at the arm

level.

Tissue-specific miRNA expression after blood feeding and after P.

falciparum *infection*

We next examined miRNA expression during the PBM phase and after *P. falciparum*

infection at 18 h post feeding (hpb) (Fig. S1). At this time point, all midgut samples

from blood-fed females had high background levels, probably due to unspecific

binding of human short RNAs and DNAs present in the blood meal. Therefore, we

excluded these samples from further analyses. Interestingly, blood feeding impacted

expression levels of very few miRNAs in other tissues. Major changes were observed

in the fat body, where blood feeding increased levels of miR-275-3p, miR-276-5p and

miR-305-5p, and reduced levels of miR-989-3p (Fig. 4A). Higher levels of miR-305-

5p were also detected in the ovaries (Fig. 4B). No changes in miRNA levels after

blood meal were observed in the head (Fig. 4C). Surprisingly, infections with *P.*

falciparum had no effect on miRNA expression in any tested tissue (Fig. S2A-C). The

expressional changes detected by microarray analysis were further confirmed by

qPCR, except for miR-989-3p with very low expression levels in the fat body (Fig.

4D). Taken together, our results identified three miRNAs whose expression was

modulated during PV to PBM transition in the fat body, whereas expression of miR-

305-5p was also upregulated in the ovaries.

Expression of blood meal-induced miRNAs during the PBM phase

The kinetics of tissue-specific expression of the blood feeding-induced miRNAs was

further analyzed by qPCR. The highest expression levels of miR-305-5p, miR-275-3p

and miR-276-5p were observed in the head before blood feeding, while expression of

miR-989-3p was only detected in the ovaries (Fig. 5A). Expression levels of miR-275-

3p, miR-276-5p and miR-305-5p transiently declined in the head at 24 and 28 hpb

and regained the initial levels by 32 hpb (Fig. 5A). In contrast, high levels of these

miRNAs were detected in the fat body and midgut at the same time points (Fig. 5B

and 5C). In the fat body, miR-275-3p and miR-276-5p levels peaked at 28 and 32

hpb and remained high at 48 hpb. High inter-replicate variation was observed for

miR-305-5p whose levels increased between 20-48 hpb (Fig. 5B). In the midgut, miR-

275-3p, miR-276-5p and miR-305-5p peaked at 24 to 32 hpb, but in contrast to the

fat body, their levels declined by 48h post BM (Fig. 5C). High expression levels of

miR-989-3p detected in the ovaries before blood feeding, gradually declined during

48 hpb. In contrast, low transcript levels of miR-305-5p, miR-275-3p and miRNA-276-

5p in the ovaries slightly increased at 28 and 32 hpb (Fig. 5D). We concluded that PV

to PBM transition regulates miRNA expression levels in a tissue- and time-specific

manner.

DISCUSSION

We report a global overview of miRNA expression in the *A. gambiae* tissues during the late PV and PBM phases. We found that the majority of *A. gambiae* miRNAs are expressed in a tissue-specific manner with the exception of the ubiquitously highly expressed miR-8. While six miRNAs showed tissue-specific bias in arm expression, one third of the miRNAs expressed both arms. Previous RNA sequencing studies of whole mosquitoes identified miR-10, miR-184, miR-263, miR-281 and miR-306 as the most abundant miRNAs in adult females (26,27). We show that expression of these abundant miRNAs is restricted to specific tissues (Fig. 6A). These results are in agreement with functional specialization of tissues, which is in part regulated by miRNAs (16,33,36).

Reproduction in many mosquito species is tightly coupled to blood feeding. During the PV phase, females feed on sugars to increase energy reserves for efficient egg development (4,5). Previous reports showed that miR-1174 regulates sugar absorption in the midgut of *Aedes* female mosquitoes (19). We demonstrate conservation of the midgut-specific expression of miR-1174/1175 and miR-281 between *A. gambiae* and *Ae. aegypti* (19,28,37). In addition, we also observed high levels of miR-277 in the midgut. *Drosophila* miR-277 regulates branched-chain amino acid (BCAA) catabolism and activates TOR signaling in the thoracic muscles (38). BCAAs serve as signaling metabolites in tissue communication, systemically promote anabolic metabolism and release hormones from the intestinal tract in mice (39–42). It is possible, that BCAA catabolism in the midgut signals food uptake to distant tissues. Further investigations of miR-277 function in the midgut are needed to explore an interesting link between food digestion, TOR signaling and systemic metabolic changes. TOR signaling is tightly connected to synthesis of JH, the major

driver of the PV phase (43). During insect development, systemic JH levels serve as

a checkpoint for sufficient energy reserves and body size before transition to the next

developmental stage. *Drosophila* miR-2 generates a threshold for developmental

transitions by regulating downstream JH signaling (44). The rise of JH levels with

nutrient availability during the PV phase suggests that the JH checkpoint function

may be also conserved in mosquitoes (8,9,45). Furthermore, high expression levels

of miR-2 in the PV phase indicate that expression of this miRNA may also regulate

JH signaling in *A. gambiae*.

Brain tissues, especially the central nervous system, play essential roles in the PV

phase by regulating mosquito olfactory and circadian behaviors (46,47). Efficient

response to blood feeding depends on timely accumulation of the ovary

ecdysteroidogenic hormone in the granules of neurosecretory cells (48). We found

that the majority of the tissue-specific miRNAs is expressed in the head, including the

well-known miRNAs that regulate olfactory sensing and circadian rhythms (49–53). In

mosquitoes, these processes shape such important vector competence traits as host

seeking and biting behavior, therefore, further investigation of the head-specific

miRNAs should identify new factors and mechanisms that regulate vector

competence.

Although feeding induces massive physiological changes in the mosquito, we

identified only modest changes in expression levels of miRNAs during the PBM

phase (Fig. 6B). Interestingly, similar expressional patterns of the blood-feeding

induced miRNAs in the fat body and midgut indicate potentially important roles of

these miRNAs in the regulation of mosquito metabolism. In line with this hypothesis,

miR-275 and miR-305 modulate metabolic processes in other insects. In *Ae. aegypti*,

miR-275 ensures successful blood meal digestion, fluid excretion and, consequently,

egg development (20). In agreement with miR-275 expression patterns in *A.*

gambiae, TOR and ecdysone control expression of this miRNA in *Ae. aegypti* (20).

Similarly, TOR and insulin pathways in *Drosophila* regulate expression of miR-305 in

the fat body and the gut, respectively, where miR-305 fosters adaptation to starvation

by balancing stem cell renewal and differentiation in a nutrient-dependent context

(54,55). Interestingly, increased levels of miR-305 in the midgut 24 h post *P.*

falciparum infection negatively impact parasite development in *A. gambiae* (25).

Although the mechanisms of parasite inhibition are currently unknown, it is plausible

that miR-305 affects *Plasmodium* development by regulating midgut homeostasis

and/or nutrient availability. Further studies should examine whether the effect of miR-

305 on *Plasmodium* development is midgut-specific. We did not detect any miRNAs

regulated by *P. falciparum* infection in the head, fat body and ovaries during the early

PBM phase. Whether parasite replication regulates miRNA expression during later

stages of the PBM phase remains to be investigated.

Ovary development and maturation is initiated in the second half of the PBM phase.

The only minor changes in the expression levels of ovarian miRNAs detected in this

study are probably due to the choice of an early time point. Nevertheless, the

observed down regulation of miR-989 coincides in time with initiation of border cell

migration, which in *Drosophila* is regulated by miR-989 (56). Therefore, *A. gambaie*

miR-989 may have a conserved role in induction of oogenesis.

Collectively, our results show tissue-specific expression patterns of 74 *A. gambiae*

miRNAs during the PV and PBM phases. Tissue-specific regulation of miRNA

expression by blood feeding is primarily observed at time points of high metabolic

activity suggesting a role in the regulation of metabolic changes throughout the

mosquito gonotrophic cycle.

Acknowledgments

The authors thank Liane Spohr and Sandrina Koppitz for mosquito breeding as well

as Dana Tschierske and Daniel Eyermann for cultivation of the *P. falciparum*

parasite. We further acknowledge the technical support of Dr. Hans-Joachim

Mollenkopf and Ina Wagner from the microarray core facility at the Max Planck

Institute for Infection Biology. Finally, we are also grateful to Dr. Maiara Severo for

comments on the manuscript.

REFERENCES

1.   Smykal V, Raikhel AS. Nutritional Control of Insect Reproduction. *Curr. Opin.*
insect Sci. 2015;11:31–38.

2.   Hansen IA, Attardo GM, Rodriguez SD, et al. Four-way regulation of mosquito
yolk protein precursor genes by juvenile hormone-, ecdysone-, nutrient-, and
insulin-like peptide signaling pathways. *Front. Physiol.* 2014;5:103.

3.   Hagedorn HH, Turner S, Hagedorn E a, et al. Postemergence growth of the
ovarian follicles of Aedes aegypti. *J. Insect Physiol.* 1977;23(2):203–206.

4.   Hou Y, Wang X-L, Saha TT, et al. Temporal Coordination of Carbohydrate
Metabolism during Mosquito Reproduction. *PLoS Genet.* 2015;11(7):e1005309.

5.   Wang X, Hou Y, Saha TT, et al. Hormone and receptor interplay in the
regulation of mosquito lipid metabolism. *Proc. Natl. Acad. Sci. U. S. A.*
2017;201619326.

6.   Lu T, Qiu YT, Wang G, et al. Odor coding in the maxillary palp of the malaria
vector mosquito Anopheles gambiae. *Curr. Biol.* 2007;17(18):1533–44.

7.   Shapiro JP, Hagedorn HH. Juvenile hormone and the development of ovarian
responsiveness to a brain hormone in the mosquito, Aedes aegypti. *Gen.*
Comp. Endocrinol. 1982;46(2):176–183.

8.   Clifton ME, Noriega FG. Nutrient limitation results in juvenile hormone-
mediated resorption of previtellogenic ovarian follicles in mosquitoes. *J. Insect*
Physiol. 2011;57(9):1274–81.

9.   Perez-Hedo M, Rivera-Perez C, Noriega FG. Starvation Increases Insulin

Sensitivity and Reduces Juvenile Hormone Synthesis in Mosquitoes. *PLoS*

One. 2014;9(1):e86183.

10. Spielman A, Gwadz RW, Anderson WA. Ecdysone-initiated ovarian

development in mosquitoes. *J. Insect Physiol.* 1971;17(10):1807–1814.

11. Hagedorn HH, O'Connor JD, Fuchs MS, et al. The ovary as a source of alpha-

ecdysone in an adult mosquito. *Proc. Natl. Acad. Sci. U. S. A.*

1975;72(8):3255–3259.

12. Zhou G, Flowers M, Friedrich K, et al. Metabolic fate of [14C]-labeled meal

protein amino acids in Aedes aegypti mosquitoes. *J. Insect Physiol.*

2004;50(4):337–349.

13. Ziegler R, Ibrahim MM. Formation of lipid reserves in fat body and eggs of the

yellow fever mosquito, Aedes aegypti. *J. Insect Physiol.* 2001;47(6):623–627.

14. Bennink S, Kiesow MJ, Pradel G. The development of malaria parasites in the

mosquito midgut. *Cell. Microbiol.* 2016;18(7):905–918.

15. Beier JC. MALARIA PARASITE DEVELOPMENT IN MOSQUITOES. *Annu.*

Rev. Entomol. 1998;43(1):519–543.

16. Jin H, Kim VN, Hyun S. Conserved microRNA miR-8 controls body size in

response to steroid signaling in Drosophila. *Genes Dev.* 2012;26(13):1427–32.

17. Verma P, Cohen SM. miR-965 controls cell proliferation and migration during

tissue morphogenesis in the Drosophila abdomen. *Elife.* 2015;4:1–18.

18. Boulan L, Martín D, Milán M. bantam miRNA promotes systemic growth by

connecting insulin signaling and ecdysone production. *Curr. Biol.*

2013;23(6):473–8.

372 19. Liu S, Lucas KJ, Roy S, et al. Mosquito-specific microRNA-1174 targets serine

373   hydroxymethyltransferase to control key functions in the gut. *Proc. Natl. Acad.*

374   *Sci. U. S. A.* 2014;111(40):14460–14465.

375 20. Bryant B, Macdonald W, Raikhel AS. microRNA miR-275 is indispensable for

376   blood digestion and egg development in the mosquito Aedes aegypti. *Proc.*

377   *Natl. Acad. Sci. U. S. A.* 2010;107(52):22391–8.

378 21. Lucas KJ, Zhao B, Roy S, et al. Mosquito-specific microRNA-1890 targets the

379   juvenile hormone-regulated serine protease JHA15 in the female mosquito gut.

380   *RNA Biol.* 2015;12(12):1383–90.

381 22. Lucas KJ, Roy S, Ha J, et al. MicroRNA-8 targets the Wingless signaling

382   pathway in the female mosquito fat body to regulate reproductive processes.

383   *Proc. Natl. Acad. Sci.* 2015;112(5):1440–1445.

384 23. Zhang Y, Zhao B, Roy S, et al. microRNA-309 targets the Homeobox gene

385   SIX4 and controls ovarian development in the mosquito Aedes aegypti. *Proc.*

386   *Natl. Acad. Sci. U. S. A.* 2016;113(33):E4828-36.

387 24. Fu X, Dimopoulos G, Zhu J. Association of microRNAs with Argonaute proteins

388   in the malaria mosquito Anopheles gambiae after blood ingestion. *Sci. Rep.*

389   2017;(June):1–13.

390 25. Dennison NJ, BenMarzouk-Hidalgo OJ, Dimopoulos G. MicroRNA-regulation of

391   Anopheles gambiae immunity to Plasmodium falciparum infection and midgut

392   microbiota. *Dev. Comp. Immunol.* 2015;49(1):170–8.

393 26. Biryukova I, Ye T, Levashina E, et al. Transcriptome-wide analysis of

394   microRNA expression in the malaria mosquito Anopheles gambiae. *BMC*

395   *Genomics.* 2014;15(1):557.

396   27.   Castellano L, Rizzi E, Krell J, et al. The germline of the malaria mosquito

397   produces abundant miRNAs, endo-siRNAs, piRNAs and 29-nt small RNAs.

398   *BMC Genomics.* 2015;16(1):100.

399   28.   Winter F, Edaye S, Hüttenhofer A, et al. Anopheles gambiae miRNAs as actors

400   of defence reaction against Plasmodium invasion. *Nucleic Acids Res.*

401   2007;35(20):6953–62.

402   29.   Zhou H, Arcila ML, Li Z, et al. Deep annotation of mouse iso-miR and iso-moR

403   variation. *Nucleic Acids Res.* 2012;40(13):5864–75.

404   30.   Chiang HR, Schoenfeld LW, Ruby JG, et al. Mammalian microRNAs:

405   experimental evaluation of novel and previously annotated genes. *Genes Dev.*

406   2010;24(10):992–1009.

407   31.   Marco A, Hui JHL, Ronshaugen M, et al. Functional shifts in insect microRNA

408   evolution. *Genome Biol. Evol.* 2010;2:686–96.

409   32.   Loya CM, McNeill EM, Bao H, et al. miR-8 controls synapse structure by

410   repression of the actin regulator enabled. *Development.* 2014;141(9):1864–74.

411   33.   Lee GJ, Jun JW, Hyun S. MicroRNA miR-8 regulates multiple growth factor

412   hormones produced from *Drosophila* fat cells. *Insect Mol. Biol.* 2015;24(3):311–

413   318.

414   34.   Lensen A, van Druten J, Bolmer M, et al. Measurement by membrane feeding

415   of reduction in Plasmodium falciparum transmission induced by endemic sera.

416   *Trans. R. Soc. Trop. Med. Hyg.* 1996;90(1):20–22.

35.   A. Lensen, Bril A, Van de Vegte M, et al. Infectivity of cultured Plasmodium

falciparum gametocytes to mosquitoes. *Exp. Parasitol.* 1999;91(1989):101–
103.

36. Loya CM, McNeill EM, Bao H, et al. miR-8 controls synapse structure by
repression of the actin regulator enabled. *Development.* 2014;141(9):1864–74.

37. Zhou Y, Liu Y, Yan H, et al. miR-281, an abundant midgut-specific miRNA of
the vector mosquito Aedes albopictus enhances dengue virus replication.
Parasit. Vectors. 2014;7(1):488.

38. Esslinger SM, Schwalb B, Helfer S, et al. Drosophila miR-277 controls
branched-chain amino acid catabolism and affects lifespan. *RNA Biol.*
2013;10(6):1042–56.

39. Nishitani S, Matsumura T, Fujitani S, et al. Leucine promotes glucose uptake in
skeletal muscles of rats. *Biochem. Biophys. Res. Commun.* 2002;299(5):693–
6.

40. Ijichi C, Matsumura T, Tsuji T, et al. Branched-chain amino acids promote
albumin synthesis in rat primary hepatocytes through the mTOR signal
transduction system. *Biochem. Biophys. Res. Commun.* 2003;303(1):59–64.

41. She P, Reid TM, Bronson SK, et al. Disruption of BCATm in Mice Leads to
Increased Energy Expenditure Associated with the Activation of a Futile Protein
Turnover Cycle. *Cell Metab.* 2007;6(3):181–194.

42. Chen Q, Reimer RA. Dairy protein and leucine alter GLP-1 release and mRNA
of genes involved in intestinal lipid metabolism in vitro. *Nutrition.*
2009;25(3):340–349.

43. Pérez-Hedo M, Rivera-Perez C, Noriega FG. The insulin/TOR signal

transduction pathway is involved in the nutritional regulation of juvenile hormone synthesis in Aedes aegypti. *Insect Biochem. Mol. Biol.* 2013;43(6):495–500.

44. Lozano J, Montañez R, Belles X. MiR-2 family regulates insect metamorphosis by controlling the juvenile hormone signaling pathway. *Proc. Natl. Acad. Sci. U. S. A.* 2015;112(12):3740–5.

45. Shiao S-H, Hansen IA, Zhu J, et al. Juvenile hormone connects larval nutrition with target of rapamycin signaling in the mosquito Aedes aegypti. *J. Insect Physiol.* 2008;54(1):231–9.

46. Kessler S, Vlimant M, Guerin PM. Sugar-sensitive neurone responses and sugar feeding preferences influence lifespan and biting behaviours of the Afrotropical malaria mosquito, Anopheles gambiae. *J. Comp. Physiol. A.* 2015;201(3):317–329.

47. Carey AF, Wang G, Su C-Y, et al. Odorant reception in the malaria mosquito Anopheles gambiae. *Nature.* 2010;464(7285):66–71.

48. Brown MR, Cao C. Distribution of Ovary Ecdysteroidogenic Hormone I in the nervous system and gut of mosquitoes. *J. Insect Sci.* 2001;11(133).

49. Chen X, Rosbash M. mir-276a strengthens Drosophila circadian rhythms by regulating timeless expression. *Proc. Natl. Acad. Sci. U. S. A.* 2016;113(21):E2965-72.

50. Li W, Cressy M, Qin H, et al. MicroRNA-276a functions in ellipsoid body and mushroom body neurons for naive and conditioned olfactory avoidance in Drosophila. *J. Neurosci.* 2013;33(13):5821–33.

464   51.   Zhang Y, Lamba P, Guo P, et al. miR-124 Regulates the Phase of Drosophila
465   Circadian Locomotor Behavior. *J. Neurosci.* 2016;36(6).

466   52.   Wang C, Feng T, Wan Q, et al. miR-124 controls Drosophila behavior and is
467   required for neural development. *Int. J. Dev. Neurosci.* 2014;38:105–112.

468   53.   Garaulet DL, Sun K, Li W, et al. miR-124 Regulates Diverse Aspects of
469   Rhythmic Behavior in Drosophila. *J. Neurosci.* 2016;36(12).

470   54.   Foronda D, Weng R, Verma P, et al. Coordination of insulin and Notch pathway
471   activities by microRNA miR-305 mediates adaptive homeostasis in the
472   intestinal stem cells of the Drosophila gut. *Genes Dev.* 2014;28(21):2421–31.

473   55.   Barrio L, Dekanty A, Milán M. MicroRNA-Mediated Regulation of Dp53 in the
474   Drosophila Fat Body Contributes to Metabolic Adaptation to Nutrient
475   Deprivation. *Cell Rep.* 2014;8(2):528–541.

476   56.   Kugler J-M, Chen Y-W, Weng R, et al. miR-989 Is Required for Border Cell
477   Migration in the Drosophila Ovary. *PLoS One.* 2013;8(7):215–233.

FIGURES

Figure 1*: Experimental set up for tissue-specific miRNA expression analysis.*
(A) Mosquitoes were fed with: (1) 10% sugar, (2) blood or (3) *P. falciparum*-infected blood. At 18 h post blood feeding, mosquitoes from each group were dissected. (B) The head, midgut, fat body and ovaries were examined for tissue-specific miRNA expression.

Figure 2: Tissue-specific miRNA expression in the 3-day-old *A. gambiae* female. Heatmap of miRNAs expression levels in the head, midgut, fat body and ovaries of sugar-fed mosquitoes in the PV phase. Colour gradient from light blue to dark brown represents an increase in miRNA expression.

**Figure 3: Tissue-*specific expression of miRNAs quantified by qPCR* of (A) miR-276-5p, (B) miR-281-5p, (C) miR-989-3p and (D) miR-305-5p. miRNA expression values are plotted as dots with the line indicating the mean. Results of six independent experiments are shown. miRNA expression levels were normalized using the ribosomal protein *RPS7* gene. (E) Tissue-specific differences in miRNA arm ratios.

Figure 4: Differentially expressed miRNAs at 18 h post blood feeding (hpb) in: (A) the fat body, (B) the ovaries and (C) the head. Cut-off log10 p-value of +/-1.3 (p<0.05) and log2 fold change of +/-0.58 (FC>1.5) (n=4) were used to detect significant changes in miRNA expression. (D) Blood feeding-induced changes (log2 FC) in the fat body and ovary of miRNA expression levels at 18 hpb measured by qPCR. Results of three independent experiments are shown, horizontal lines indicate mean values.

Figure 5: Tissue-specific expression of miR-276-5p, miR-305-5p, miR-275-3p and miR-989-3p during the PBM phase in (A) the head, (B) the fat body, (C) the midgut and (D) the ovary. Results of three independent experiments are shown as mean +/- SEM. For statistical analyses all time points were compared to 0h (sugarfed) by t-test and significant differences are shown by asterisk (* = p<0.05).

Figure 6: Schematic summary of tissue-specific miRNA expression during the gonotrophic cycle of the female *A. gambiae* mosquito (A) Tissue-enriched miRNAs in sugar-fed mosquitoes. Enrichment cut-off for tissue-specific miRNAs was set at 1.5-fold. (B) miRNAs regulated during the PBM phase. The coloured boxes show tissue-specific miRNAs (blue = head, yellow = fat body, green = ovary, red = midgut).

Figure S1: Tissue-specific miRNA expression in blood-fed *A. gambiae.* Heatmap of miRNA expression levels in the head, midgut, fat body and ovaries at 18 h post blood meal. Colour gradient from light blue to dark brown represents the increase in miRNA expression.

Figure S2: Differentially expressed miRNAs at 18 h post *P. falciparum*-infected blood feeding in *A. gambiae* tissues. (A) fat body, (B) ovaries and (C) head. Cut-off log10 p-value of +/-1.3 (p<0.05) and log2 fold change of +/-0.58 (fold-change>1.5) (n=4) were used to detect significant differences in miRNA expression.

TABLE

497   Table S1: The microarray repertoire of *A. gambiae* miRNA probes

aga-miR-212	aga-miR-79-5p	aga-miR-278-3p
aga-miR-929-5p	aga-miR-87-3p	aga-miR-9c-5p
aga-miR-100-5p	aga-mir-2796-5p	aga-miR-9a-5p
aga-miR-281-5p	aga-mir-2940-5p	aga-miR-957-3p
aga-miR-1175-5p	aga-mir-33-3p	aga-miR-1890-3p
aga-miR-308-3p	aga-miR-263b-3p	aga-miR-317-3p
aga-miR-282-5p	aga-miR-1174-5p	aga-mir-286b-5p
aga-miR-92b-3p	aga-miR-988-5p	aga-mir-2945-5p
aga-miR-125-3p	aga-miR-79-3p	aga-miR-279-3p
aga-miR-1889-3p	aga-miR-9c-3p	aga-miR-996-3p
aga-mir-10368-5p	aga-miR-283-5p	aga-bantam-3p
aga-miR-10-5p	aga-miR-277-3p	aga-miR-1175-3p
aga-miR-iab-4-5p	aga-mir-10375b-3p	aga-miR-1891-5p
aga-miR-307-5p	aga-miR-965	aga-let-7-5p
aga-miR-9b-5p	aga-miR-965-3p	aga-miR-12-5p
aga-miR-190-5p	aga-miR-124-3p	aga-miR-219-5p
aga-miR-276-5p	aga-miR-8-3p	aga-mir-932-3p
aga-miR-210-5p	aga-miR-993-5p	aga-mir-2c-3p
miR-iab-8-3p	aga-mir-998-5p	aga-miR-7-5p
aga-mir-10355-3p	aga-mir-285-5p	aga-miR-1-3p
aga-miR-1000-5p	aga-mir-31-3p	aga-miR-184-3p
aga-miR-305-5p	aga-mir-980-3p	aga-mir-31-5p
aga-miR-927-3p	chr2R_3959-miR*	aga-miR-72-1
aga-miR-10-3p	aga-miR-276-3p	aga-miR-34-5p
aga-miR-1889-5p	chr2L_3426-miR	aga-miR-263a-5p
aga-miR-11-3p	aga-mir-2a-1-3p	aga-miR-281-3p
aga-miR-8-5p	aga-miR-2-1-3p	chrX_40848
aga-miR-1-5p	aga-miR-13b-3p	aga-chrX_353716
aga-miR-190-3p	aga-mir-2944a-1-3p	aga-miR-989-3p
aga-bantam-5p	aga-miR-92a-3p	aga-mir-999-5p
aga-miR-305-3p	aga-miR-137-3p	chrUNKN_30285
aga-miR-iab-4-3p	aga-mir-932-5p	chr2R_12239-miR
aga-mir-980-5p	aga-miR-307-3p	aga-miR-981-3p
aga-miR-263a-3p	aga-miR-309-3p	aga-miR-133-3p
aga-miR-929-3p	aga-miR-3-1-miR	aga-miR-210-3p
aga-mir-252-3p	aga-miR-1174-3p	aga-miR-927-5p
aga-miR-263b-5p	aga-miR-275-3p	aga-miR-375-1-3p
aga-miR-315-3p	aga-miR-306-5p	aga-miR-315-5p
aga-miR-993-3p	aga-miR-14-3p	
aga-mir-2944b-5p	aga-miR-970-3p	
aga-mir-2944a-5p	aga-miR-125-5p	

498   Table S2: Proportion of miR-3p arm expression of miRNAs expressing both miRNA

499   arms.

miRNA	Fat body	Head	Midgut	Ovary	SD
bantam-3p	0,590378	0,539231	0,50896	0,546386	0,029099
miR-10-3p	0,425485	0,362922	0,297304	0,546466	0,091875
miR-1175-3p	0,491238	0,524269	0,310941	0,510286	0,086387
miR-124-3p	0,418944	0,74533	0,470842	0,494804	0,125911
miR-125-3p	0,22676	0,183561	0,170767	0,208014	0,021653
miR-13b-3p	0,568517	0,59165	0,656206	0,614877	0,032393
miR-1-3p	0,733715	0,744	0,764682	0,775197	0,016388
miR-1889-3p	0,534361	0,553319	0,547071	0,649744	0,045902
miR-190-3p	0,503467	0,533528	0,510703	0,46622	0,024204
miR-210-3p	0,514505	0,714925	0,475985	0,502771	0,095069
miR-263-3p	0,288323	0,267782	0,439021	0,225327	0,08058
miR-276-3p	0,501213	0,57535	0,515418	0,594905	0,039345
miR-281-3p	0,431388	0,517835	0,215577	0,292874	0,117573
miR-305-3p	0,472667	0,401627	0,451947	0,460314	0,027013
miR-307-3p	0,597067	0,572176	0,465073	0,511524	0,05164
miR-31-3p	0,520295	0,399548	0,564907	0,54837	0,06477
miR-79-3p	0,57127	0,671821	0,698043	0,740639	0,062303
miR-8-3p	0,550598	0,595151	0,556974	0,595009	0,02077
miR-927-3p	0,468846	0,445864	0,467125	0,481053	0,01266
miR-929-3p	0,458689	0,414248	0,496434	0,503308	0,035465
miR-993-3p	0,529512	0,506608	0,507545	0,494083	0,012751