Awake fMRI Reveals Brain Regions for Novel Word Detection in Dogs

Ashley Prichard[1], Peter F. Cook[2], Mark Spivak[3], Raveena Chhibber[1], & Gregory S. Berns[1*]

[1]Psychology Department, Emory University, Atlanta, GA 30322; [2]Psychology Department, New College of Florida, Sarasota, FL 34243; and [3]Comprehensive Pet Therapy, Atlanta, GA, 30328

[*] E-mail: gberns@emory.edu

Abstract

How do dogs understand human words? At a basic level, understanding would require the discrimination of words from non-words. To determine the mechanisms of such a discrimination, we trained 12 dogs to retrieve two objects based on object names, then probed the neural basis for these auditory discriminations using awake-fMRI. We compared the neural response to these trained words relative to "oddball" pseudowords the dogs had not heard before. Consistent with novelty detection, we found greater activation for pseudowords relative to trained words bilaterally in the parietotemporal cortex. To probe the neural basis for representations of trained words, searchlight multivoxel pattern analysis (MVPA) revealed that a subset of dogs had clusters of informative voxels that discriminated between the two trained words. These clusters included the left temporal cortex and amygdala, left caudate nucleus, and thalamus. These results demonstrate that dogs' processing of human words utilizes basic processes like novelty detection, and for some dogs, may also include auditory and hedonic representations.

Introduction

Because dogs can learn basic verbal commands, it is obvious that they have the capacity for discriminative processing of some aspects of human language. For humans, words represent symbolic placeholders for a multitude of people, objects, actions, and other attributes. However, just because a dog can match a word with an action, like 'fetch,' does not mean that the dog understands the word has meaning in the same way humans do. For example, dogs may rely on other cues to follow verbal commands such as gaze, gestures and emotional expressions, as well as intonation (Fukuzawa et al., 2005;Mills, 2015;Müller et al., 2015;Persson et al., 2015;D'Aniello et al., 2016). This raises the question of what cognitive mechanisms dogs use to differentiate between words, or even what constitutes a word to a dog.

Part of the problem in studying word comprehension in dogs is the necessity of a behavioral response to demonstrate understanding. Some dogs can retrieve a named object based on a command combined with the name of the object, but this often requires months of training. Examples include Chaser, the border collie who learned over one thousand object-word pairings, and the border collie Rico, who demonstrated the ability to select a novel object among familiar objects based on a novel label (Kaminski et al., 2004;Pilley and Reid, 2011;Zaine et al., 2014).

But these dogs may have been exceptional. Few other dogs have been documented to have this
level of expertise. It may be that most dogs rely on simple mechanisms of discrimination – like
novelty detection – coupled with other cues from the human to figure out an appropriate
behavioral response.

The auditory oddball task, where subjects behaviorally discriminate between target and novel
acoustic stimuli, is a well-established task used to measure the processing of target detection and
decision-making in humans and nonhumans. The neural regions responsible for target detection
and novelty processing not only include primary sensory areas associated with the stimulus
modality, but also recruit broader areas such as the posterior cingulate, inferior and middle
frontal gyri, superior and middle temporal gyri, amygdala, thalamus, and lateral occipital cortex
(Linden et al., 1999;Kiehl et al., 2001;Brazdil et al., 2005;Goldman et al., 2009;Cacciaglia et al.,
2015). This suggests that differentiating between target versus novel sounds requires primary
auditory cortex as well as an additional attentional network to discriminate between competing
sensory stimuli. At least one event-related potential study in dogs suggested similar mechanisms
might be at work, finding mismatch negativity to deviant tones (Howell et al., 2012).

Recent advances in awake neuroimaging in dogs have provided a means to investigate many
aspects of canine cognition using approaches similar to those in humans. Since 2012, pet dogs
have been trained using positive reinforcement to lie still during fMRI scans in order to explore a
variety of aspects of canine cognition (Berns et al., 2012;Berns et al., 2013). These studies have
furthered our understanding of the dog's neural response to expected reward, identified
specialized areas in the dog brain for processing faces, observed olfactory responses to human
and dog odors, and linked prefrontal function to inhibitory control (Cook et al., 2014;Berns et al.,
2015;Dilks et al., 2015;Cook et al., 2016a;Cuaya et al., 2016). In one fMRI study, dogs listened
to human and dog vocalizations through headphones and showed differential activation within
regions of the temporal and parietal cortex (Andics et al., 2014). A follow-up study suggested a
hemispheric bias for praise words versus neutral words, a finding that was interpreted as proof of
semantic processing in dogs. However, a subsequent correction in which left and right were
reversed raised questions about the interpretability of this finding (Andics et al., 2016).

To examine auditory processing in dogs, we used fMRI to measure activity in dogs' brains in
response to both trained words and novel pseudowords. Over several months prior to scanning,
owners trained their dogs to select two objects based on the objects' names. During the fMRI
session, the owner spoke the names of the trained objects as well as novel pseudowords the dog
had never heard before. If dogs discriminate target words from novel words as humans do, they
should show differential activity in the parietal and temporal cortex in response to trained words
relative to pseudowords (Friederici et al., 2000;Humphries et al., 2006;Raettig and Kotz, 2008).
In addition, if dogs use hedonic mechanisms to associate reward value with trained words, then
differential activity should also be observed in the caudate.

Methods

Ethics Statement

This study was performed in accordance with the recommendations in the Guide for the Care and
Use of Laboratory Animals of the National Institutes of Health. The study was approved by the
Emory University IACUC (Protocol DAR-2002879-091817BA), and all owners gave written
consent for their dog's participation in the study.

Participants

Participants were pet dogs from the Atlanta community volunteered by their owners for fMRI
training and experiments (Table 1). All dogs had previously completed one or more scans for the
project and had demonstrated the ability to remain still during training and scanning (Berns et al.,
2012).

Table 1. Dogs and their object names.

Dog	Breed	Age	Sex	Years with fMRI project	Object 1	Object 2
Caylin	Border Collie	8	Spayed F	4	Monkey	Blue
Eddie	Golden Retriever-Lab mix	6	Neutered M	2	Piggy	Monkey
Kady	Golden Retriever–Lab mix	7	Spayed F	4	Taffy	Yellow
Libby	Pit mix	11	Spayed F	4	Duck	Hedge Hog
Ninja	Australian Cattle dog- mix	2	Spayed F	1	Block	Monkey
Ohana	Golden Retriever	7	Spayed F	3	Blue	Star
Pearl	Golden Retriever	7	Spayed F	3	Duck	Elephant
Stella	Bouvier	6	Spayed F	3	Stick	Tuxy
Truffles	Pointer mix	12	Spayed F	2	Pig	Blue
Velcro	Viszla	8	Intact M	3	Rhino	Beach Ball
Zen	Golden Retriever– Lab mix	8	Neutered M	4	Teddy	Duck
Zula	Lab-Mastiff mix	4	Spayed F	1	Goldie	Bluebell

85   Dog's names, breed, age in years when undergoing scanning, sex, years participating in fMRI experiments, and
86   training objects (S+) are listed
87
Word-Object Training

In the current experiment, dogs were trained to reliably fetch or select a trained object given the
matching verbal name for the object. The dogs were trained by implementing the "Chaser
Protocol" in which object names were used as verbal referents to retrieve a specific object (Pilley
and Reid, 2011). To keep the task simple, each dog had a set of two objects, selected by the
owner from home or from dog toys provided by the experimenters. One object had a soft texture,
such as a stuffed animal, whereas the other was of a different texture such as rubber or squeaked,
to facilitate discrimination (Fig. 1).

Each dog was trained by his or her owner at home, approximately 10 minutes per day, over 2 to
6 months, as well as at biweekly practices located at a dog training facility. Initial shaping

involved the owner playing "tug" or "fetch" with her dog and one object while verbally
reinforcing the name of the object. Later, the objects were placed at a distance (four feet on
average) and the owner instructed the dog to "go get [object]" or "where is [object]?" or
"[object]!" The dog was reinforced with food or praise (varied per dog) for retrieving or nosing
the object. Next, the object was placed beside a novel object roughly two feet apart, at least 4
feet from the dog, and the command repeated. The dog was reinforced only for correctly
selecting the trained object if it was her first selection. Otherwise, if the dog selected the wrong
object, the owner made no remark and a new trial began. Regardless of the selection, objects
were rearranged before each trial to limit learning by position. If the dog failed to approach an
object, the trial was repeated. This training was repeated for each dog's second object against a
different comparison object, to limit the possibility of learning by exclusion. Owners were
instructed to train one object per day, alternating between objects every other day until they
showed the ability to discriminate between the trained and novel object, at which point they
progressed to discrimination training between the 2 trained objects.

All dogs in the current study participated in training for previous fMRI experiments. As
described in previous experiments (Berns et al., 2012;Berns et al., 2013;Cook et al., 2014;Cook
et al., 2016b), each dog had participated in a training program involving behavior shaping,
desensitization, habituation and behavior chaining to prepare for the loud noise and physical
confines of the MRI bore inherent in fMRI studies.

Word-Object Discrimination Tests

Two weeks after progressing to two-object discrimination training, and every two weeks
thereafter, each dog was tested on her ability to discriminate between the two trained objects.
Discrimination between the two named objects was chosen as the measure of performance, as
both objects had a similar history of reinforcement, and this precluded the possibility that
performance was based on familiarity. Discrimination testing consisted of the observer placing
both trained objects 2-3 feet apart, and at least 4 feet from the dog (Ann Young, 1991), though
the number of distractor objects was sometimes increased during training to maximize
discriminatory performance. With the dog positioned next to the owner in the heel position, the
owner gave the dog the command to "go get [object]" or "[object]!" The dog was reinforced only
for correctly selecting the trained object if it was her first selection. If the dog selected the
incorrect object, the owner made no remark. After each trial, the objects were rearranged, and the
test progressed to the next trial. A performance criterion to move forward to the MRI scan was
set at 80% correct for at least one of the objects, with the other object at or above 50%.

During training, owners were asked to report if their dog showed a preference for one object over
the other. For the majority of the dogs, the preference was for the softer object of the two, and
both the preferred word and the object were consistently labeled as word 1 and object 1. Though
Zula passed the discrimination test, she was unable to complete the MRI scan and was excluded

from the remainder of the study. Individuals varied on the amount of time needed to train both
objects.

Scan Day Discrimination Test

Scan day tests were conducted in a neighboring room to the MRI room, and were typically
conducted prior to the MRI scan. Test procedure was identical to the word-object discrimination
test as described above, although the number of trials was increased from 10 to 12 trials if the
dog failed to make a response during one or more trials.

fMRI Stimuli

The stimuli consisted of the two trained words and the corresponding objects. Pseudowords were
included as a control condition. Pseudowords were matched to the group of trained words based
on the number of syllables and bigram frequency where possible using a pseudoword generator
(Keuleers and Brysbaert, 2010) (Table 2). Phoneme substitution was necessary in some cases to
ensure that trained words and pseudowords did not overlap at onset or coda. During the scan,
pseudowords were followed by the presentation of novel objects with which the dogs had no
previous experience. The novel objects included a bubble wand, Barbie doll, stuffed caterpillar,
wooden train whistle, plastic gumball dispenser, yellow hat, watermelon seat cushion, Nerf ball
launcher, etc.

Table 2. List of pseudowords per run.

Run 1	Run 2	Run 3
prang	cal	Cloft
risnu	o gri	Sowt
doba	ropp	bodmick
bobbu	prel	Fons
zelve	thozz	Stru

fMRI Experimental Design

As in previous studies, dogs were stationed in the magnet bore using custom chin rests. All
words were spoken by the dog's primary owner, who stood directly in front of the dog at the
opening of the magnet bore. Both owners and dogs wore ear plugs, which reduced scanner noise
by approximately 30 decibels, but allowed for intelligible human speech over the sound of the
scanner. The spoken words were intelligible to the experimenters, who also wore ear plugs while
next to the MRI during scanning, as well as human operators in the control room via the
intercom. At the onset of each trial, a word was projected onto the surface of the scanner, directly
above the owner's head. An experimenter stood next to the owner, out of view of the dog. The
experimenter controlled the timing and presentation of the words to the owner via a four-button
MRI-compatible button box (Fig. 2A). Onset of words and objects were controlled by the

simultaneous presentation and press of the button box by the experimenter marking the onset and
duration of presentation. This was controlled manually by the experimenter during each dog's
scan, as opposed to a scripted presentation as in human fMRI studies, because dogs may leave
the MRI at any time and data for absentee trials would be lost.

An event-based design was used, consisting of four trial types presented semi-randomly:
expected, unexpected, pseudoword, and reward. On expected trials, the owner repeated a trained
object's name five times, once per second. Words were repeated to ensure a robust hemodynamic
response on each trial and spoken loudly to be heard above the scanner noise. After a variable 3
to 8 s delay, the dog was shown the corresponding object for 5 s and was subsequently allowed
to interact with the object. During unexpected trials, the owner repeated the name for a trained
object as above, but following the delay period a novel object was presented instead of the
corresponding object. In pseudoword trials, the owner repeated a pseudoword, and the delay was
followed by a novel object. Reward trials were interspersed throughout each run, during which
the owner rewarded the dog's continued down-stay with food. Trials were separated by a 6 s
inter-trial interval, and each dog received the same trial sequence (Fig. 2B). Each of three runs
consisted of 26 trials, for a total of 78 trials. The trial types included: 30 expected (15 each of
word1 and word2), 15 unexpected (7 or 8 of word1 and word2), 15 pseudowords, and 18 food
rewards.

Imaging

Scanning for the current experiment was conducted with a Siemens 3 T Trio whole-body scanner
using procedures described previously (Berns et al., 2012;Berns et al., 2013). During previous
experiments, a T2-weighted structural image of the whole brain was acquired using a turbo spin-
echo sequence (25-36 2mm slices, TR = 3940 ms, TE = 8.9 ms, flip angle = 131˚, 26 echo trains,
128 x 128 matrix, FOV = 192 mm). The functional scans used a single-shot echo-planar imaging
(EPI) sequence to acquire volumes of 22 sequential 2.5 mm slices with a 20% gap (TE = 25 ms,
TR = 1200 ms, flip angle = 70˚, 64 x 64 matrix, 3 mm in-plane voxel size, FOV = 192 mm).
Slices were oriented dorsally to the dog's brain (coronal to the magnet, as in the sphinx position
the dogs' heads were positioned 90 degrees from the prone human orientation) with the phase-
encoding direction right-to-left. Sequential slices were used to minimize between-plane offsets
from participant movement, while the 20% slice gap minimized the "crosstalk" that can occur
with sequential scan sequences. Three runs of up to 700 functional volumes were acquired for
each participant, with each run lasting 10 to 14 minutes.

Analysis

Preprocessing

Data preprocessing included motion correction, censoring and normalization using AFNI (NIH)
and its associated functions. Two-pass, six-parameter affine motion correction was used with a
hand-selected reference volume for each dog that best reflected their average position within the

scanner. All volumes were aligned to the reference volume. Aggressive censoring (i.e., removing
bad volumes from the fMRI time sequence) was used because dogs can move between trials,
when interacting with the object, and when consuming rewards. Data were censored when
estimated motion was greater than 1 mm displacement scan-to-scan and based on outlier voxel
signal intensities. Smoothing, normalization, and motion correction parameters were identical to
those described previously (Cook et al., 2016b). A high-resolution canine brain atlas was used as
the template space for individual spatial transformations (Datta et al., 2012). The atlas resolution
was 1 mm x 1 mm x 1 mm. Thus voxel volumes are in mm^3.

General Linear Model

For *a priori* hypotheses, each participant's motion-corrected, censored, smoothed images were
analyzed with a general linear model (GLM) for each voxel in the brain using 3dDeconvolve
(part of the AFNI suite). Nuisance regressors included motion time courses generated through
motion correction, constant, linear, quadratic, and cubic drift terms. The drift terms were
included for each run to account for baseline shifts between runs as well as slow drifts unrelated
to the experiment. Task related regressors included: (1) spoken word1; (2) spoken word2; (3)
spoken pseudowords; (4) presentation of object1; (5) presentation of object2; (6) presentation of
unexpected objects (novel object following either word1 or word2); and (7) presentation of novel
objects following a pseudoword. The object on which each dog performed best during the day of
the MRI scan as well as the object owners reported as being the preferred of the two was labeled
as word1 and object1 when creating the GLM regressors. Stimulus onset and duration were
modeled using the dmUBLOCK function, with the 5 utterances treated as a block.

Whole Brain Analysis

Contrasts focused on the dogs' response to words and pseudowords. Auditory novelty detection
was probed with the contrast: [pseudowords – (word1 + word2)/2]. Low-level aspects of
language processing (including acoustic and hedonic representations) were probed with the
contrast [word1 – word2] and expectation violation with [novel objects – unexpected objects].

Each participant's individual-level contrast from the GLM was normalized to template space as
described in (Berns et al., 2012;Cook et al., 2014) via the Advanced Normalization Tools
(ANTs) software (Avants et al., 2011). Spatial transformations included a rigid-body mean EPI
to structural image, affine structural to template, and diffeomorphic structural to template. These
spatial transformations were concatenated and applied to individual contrasts from the GLM to
compute group level statistics. 3dttest++, part of the AFNI suite, was used to compute a t-test
across dogs against the null hypothesis that each voxel had a mean value of zero. All contrasts
mentioned above as part of the GLM were included.

As there is spatial heterogeneity within fMRI data, the average smoothness of the residuals from
each dog's time series regression model was calculated using AFNI's non-Gaussian spatial
autocorrelation function 3dFWHMx –acf. The acf option leads to greatly reduced FPRs clustered

around 5 percent across all voxelwise thresholds (Cox et al., 2017). AFNI's 3dClustsim was then
used to estimate the significance of cluster sizes across the whole brain after correcting for
familywise error (FWE). Similar to human fMRI studies, a voxel threshold of $p \leq 0.005$ was
used, and a cluster was considered significant if it exceeded the critical size estimated by
3dClustsim for a FWER ≤ 0.01, using two-sided thresholding and a nearest-neighbor of 1.

Multivoxel Pattern Analysis (MVPA)

In previous fMRI studies of the oddball task, it was noted that attentional differences occurring
trial-by-trial may go undetected in the univariate analysis (Goldman et al., 2009). As an
exploratory analysis, we used searchlight MVPA to identify regions potentially involved in the
representation of words that were not captured in the univariate analysis. We were primarily
interested in the representation of word1 vs. word2.

We used a linear support vector machine (SVM) for a classifier because of its previously
demonstrated robust performance (Misaki et al., 2010;Mahmoudi et al., 2012). Unsmoothed
volumes were censored for motion and outlier count as in the univariate GLM. We then made a
model for the unsmoothed data using AFNI's 3dDeconvolve stim_times_IM function. This
model yielded trial-by-trial estimates (betas) for each repetition of word1 and word2, regardless
of which object followed. Although it is common in the human literature to use each scan
volume as a data point in MVPA (for training and testing), we have found this approach to be
problematic with dogs, who move more than humans, resulting in spurious volumes that should
be censored. Estimating the beta for each trial affords an additional level of robustness with less
sensitivity to potential outlier volumes due to motion. As an additional check for outliers, masks
were drawn of the left and right caudate on each dogs' T2-weighted structural image. Average
beta values were extracted from both the left and right caudate for each trial of word1 and word2.
Trials with beta values greater than |3%| were assumed to be non-physiological and were
removed prior to MVPA. Finally, these trial-dependent estimates were then used as inputs to a
whole-brain searchlight MVPA for each individual dog using PyMVPA2 (Hanke et al., 2009).
The classifier was trained on the fMRI dataset for each dog by training on 2 runs and testing on
the third using the NFoldPartitioner. We used the Balancer function to retain the same number of
trials for word1 and word2 across training and testing for 100 repetitions. For the searchlight, we
used a 3-voxel radius sphere. This yielded a map of classification accuracies throughout each
dog's brain.

Given the difficulty in finding significant effects in small datasets using cross-validation and
parametric methods, we used a permutation approach outlined by Stelzer et al. (2013) to
determine the significance of any cluster of common voxels across dogs (Stelzer et al.,
2013;Varoquaux, 2017). Briefly, we permuted the order of attributes—but not their
corresponding data—and ran the searchlight in individual space for all dogs. This created a null
distribution of accuracies. The mean of these distributions was noted to be very close to 0.5,
confirming that the classifiers wasn't biased or skewed. The cumulative distribution of that an

accuracy ≥ 0.63 corresponded to the top 5% of voxels, and this was used as a cut-off threshold
for the individual maps. These binarized maps were transformed into template space and the
average computed across dogs. The resultant group map represented the locations of potentially
informative voxels and served as qualitative representation of the relative consistency versus
heterogeneity of word-processing in the dogs' brains. Somewhat arbitrarily, we only considered
locations in which at least two dogs had informative voxels.

283
284   **Results**
285
Scan Day Discrimination Tests

Scans were scheduled as close as possible to the day on which object identification criterion was
met ($M = 9.33$ days, $SD = 4.92$ days) based on owner availability. On the day of the scheduled
MRI scan, each dog was tested on her ability to behaviorally differentiate between the two
trained objects out of 5 trials each. With the exception of Eddie, each dog correctly selected
object 1 on 80 to 100 percent of the trials [$M=85.73\%$, $SE = 3.87\%$], and object 2 on 60 to 100
percent of the trials [$M=64.27\%$, $SE=5.91\%$] (Fig. 3). The percent correct performance
(subtracting 50 percent for chance levels of responding) on scan days for each object was
compared in a mixed-effect linear model and showed that performance was significantly greater
than chance [$T(17.1) = 3.00$, $P = 0.008$] and that there was a significant difference in
performance between word1 and word2 [$T(11) = 4.67$, $P < 0.001$].

Primary Auditory and Visual Activation

To confirm that the dogs clearly heard the words during scanning, a simple contrast subtracting
activation to objects (trained and novel) from activation to words (trained and pseudowords) was
performed. In human fMRI, the MRI operator may ask the participant whether they can hear
auditory stimuli, which is not necessarily possible in dog fMRI, so this was included as a quality
check. We opted for an unthresholded image not only to highlight the activation in bilateral
auditory cortex but, just as important, to show what was not activated. Notably in the contrast
[Words—Objects] positive activation was localized to the auditory cortex for words and negative
activation for presentation objects in parietal cortex (Fig. 4), confirming that the dogs heard the
words and saw the objects.

Whole Brain Analyses

Whole brain analysis of the contrasts of interest revealed significant activation only within the
right parietotemporal cortex for the contrast [pseudowords – trained words]. With a voxel-level
significance threshold of $P \leq 0.005$, the cluster size in the right hemisphere (839 voxels) was
statistically significant at $P \leq 0.005$ after correction for whole-brain FWE (although activation
appeared bilaterally) (Fig. 5). Whole brain analysis of the contrasts of [word1– word2] and

[novel – unexpected] were not significant as no cluster survived thresholding at the voxel
significance mentioned above.

315
316 *MVPA*
317
Because the univariate analysis of word1 vs. word2 did not reveal any region with a significant
difference, we used MVPA to explore potential regions that may code for different
representations of the words. The searchlight map of word1 vs. word2, which identified regions
involved in the discrimination of the trained words, showed four clusters of informative voxels
(Fig. 6): posterior thalamus/brainstem; amygdala; left temporoparietal junction (TPJ); and left
dorsal caudate nucleus. Seven dogs shared informative voxels in or near the left temporal cortex
that passed the 0.63 accuracy threshold (Fig. 7).

Discussion

Using awake-fMRI in dogs, we found neural evidence for auditory novelty detection in the
domain of human speech. The hallmark of this finding was greater activation in parietotemporal
cortex to novel pseudowords relative to trained words. Thus, even in the absence of a behavioral
response, we demonstrate that dogs process human speech at least to the extent of differentiating
words they have heard before from those they have not. The mechanism of such novelty
detection may be rooted in either the relatively less frequent presentation of the pseudowords
(oddball detection) or the lack of meaning associated with them (lexical processing).

The activation observed in the parietotemporal cortex to pseudowords relative to trained words
meets current standards of human fMRI analyses concerning up-to-date methods for cluster
thresholds. Specifically, to address concerns raised by Eklund et al. (2016), present analyses for
cluster inferences address the former Gaussian-shaped assumption about spatial structure in the
residuals of fMRI data and provide more accurate false positive rates compared to previous
methods (Eklund et al., 2016;Cox et al., 2017;Slotnick, 2017). As the identified cluster was
significant at $P \leq 0.005$, corrected for whole-brain FWE, the result does not appear to be a false
positive. However, as the study was limited to 11 participants, future studies with an increased
number of participants could produce a more robust finding.

In humans, greater BOLD response to pseudowords versus real words has been noted in the
superior temporal gyrus – an area potentially analogous to the one we identified in dogs (Kotz,
2002;Raettig and Kotz, 2008). In humans, stronger activation to pseudowords depends on
whether the pseudoword strongly resembles a known word or is so unlike known words as to
prevent any semantic retrieval. When the pseudoword is similar to a known word, more
processing has been observed in the superior temporal gyri, presumably to disambiguate it from
known words (Raettig and Kotz, 2008). Thus, in dogs, the greater activation to the pseudowords
could be due to the acoustic similarity between pseudowords and words that the dogs "knew"
and their attempt to resolve the ambiguity. This would be a form of low-level lexical processing.

However, previous research has shown that dogs can discriminate between altered phonemes of
well-known commands (Fukuzawa et al., 2005), suggesting that it is unlikely that the dogs in our
study were confused by acoustic similarity of words and pseudowords.

More likely, a novel word resulted in increased auditory processing to facilitate learning the
association with the novel object that followed. A dog's behavioral bias for novelty is often
described as an explanation for performance otherwise labeled as learning by exclusion (Bloom,
2004;Markman and Abelev, 2004;Zaine et al., 2014). As such, a dog may select a novel item
because it is novel among other stimuli, but not because she has learned all other stimuli and
associated a new word with the novel item. A bias for novelty would therefore be reflected in the
dog's brain as with her behavior.

Auditory stimuli can be difficult to discriminate in the scanner. We used a continuous scanning
protocol because that is what the dogs were accustomed to. The simple contrast of all words vs.
all objects showed bilateral activation of the superior temporal lobe, indicating that the dogs
heard something. However, the main effect of pseudowords vs. trained words showed that the
majority of dogs discriminated well enough to tell the difference. The predominant location in
the auditory pathway also suggests that the effect wasn't based on non-verbal cues from the
handler (i.e. Clever Hans effect).

The manner in which dogs learn words is different than humans do, and this undoubtedly affects
their performance on behavioral tests and the patterns of brain activation we observed. Humans
acquire nouns as early as six months of age and differentiate between nouns prior to their ability
to use verbs (Bergelson and Swingley, 2012;Waxman et al., 2013). In contrast, dogs do not
typically have much experience with nouns because humans tend to train them on actions/verbs
(e.g. sit and fetch). Consequently, even the trained words in our study were novel for the dogs in
comparison to years of experience with verbs as commands. Prior studies have shown only three
dogs that consistently retrieved objects given a verbal referent (Kaminski et al., 2004;Pilley and
Reid, 2011). Additionally, those dogs had been trained to retrieve from a young age (<11
months), and in most cases rarely attained 100 percent accuracy. Object retrieval training for the
current experiment was modeled from these studies; however, because the dogs' owners
conducted training at home on a voluntary basis, training rigor could not be enforced.

Although humans readily generalize the meaning of words to a variety of contexts, this may not
be the case for dogs. The environment in which the dogs learned the words was different than
both the testing and scanning environments (Mills, 2015). In addition, although human fMRI
language studies do not typically repeat the spoken word each trial, as is common in oddball
paradigms, it was necessary for the dogs to make sure that they heard each word. Trials also did
not include a condition in which a spoken pseudoword was followed by a trained object, or trials
in which a trained object was mismatched to a trained word. These types of trials would have
provided additional evidence for violation of expected semantic content; however, these types of
trials have the potential to confuse the dogs and result in extinction of the words already learned.

Lastly, dogs might have habituated to the continued presentation of trained words followed by
trained objects, as opposed to the single trial presentations of pseudowords and the
accompanying novel objects.

So what do words mean to dogs? Even though our findings suggest a prominent role for novelty
in dogs' processing of human words, this leaves the question of what the words represent. One
possibility is that the words had no further representation than the relative hedonic value of the
objects. While some dogs showed a behavioral preference for one object over the other, this
preference was not reflected in whole brain analyses. Admittedly, the somewhat arbitrary
designation of word1 / word2 and object1 / object2 could explain the nonsignificant results in the
univariate analysis. Indeed, the MVPA of word1 vs. word2, which identified regions that
classified the words above chance regardless of directionality, showed one cluster in the left
caudate. However, the MVPA also identified clusters in the left TPJ, amygdala, and posterior
thalamus. The TPJ was located just posterior to the region in the univariate analysis, which
would take it out of the area of cortex associated with low-level acoustic processing. Its location
appears similar to human angular gyrus – aka Wernicke's area. If so, this could be a potential site
for receptive word processing in dogs (e.g. the Dog Wernicke's Area), but future work would
need to verify this.

Evaluating classifier performance for MVPA remains a complex task. We used MVPA as an
exploratory analysis to identify brain regions that potentially discriminate between trained words
across dogs. But classification using the whole brain may result in a high classification accuracy
that is not generalizable across subjects. Indeed, the regions identified using MVPA were of
marginal statistical significance, especially given the small sample size. Further, it should be
noted that only a subset of dogs contained informative voxels in the TPJ region. Although all
dogs had informative voxels somewhere in the brain, only seven dogs had informative voxels in
the TPJ area. Thus, even though all the dogs were cleared for scanning by reaching performance
criterion, they may have used different mechanisms to process the words. Like our previous
fMRI studies, heterogeneity seems to be the rule (Cook et al., 2016a;Cook et al., 2016b). Even
so, the accuracy of the classifier was not correlated with a dog's performance. This suggests that
performance on such tasks may be influenced by factors other than word discrimination alone.

These results highlight potential mechanisms by which dogs process words. Word novelty
appears to play an important role. The strong response of the parietotemporal region to
pseudowords suggests that dogs have some basic ability to differentiate words with associations
from those that do not. Future studies may reveal whether these representations remain in the
auditory domain or whether such representations are invariant to modality.

Acknowledgments

Thanks to Dr. Kate Revill for advice about language processing and generating pseudowords.
Thank you to all of the owners who trained their dogs over the course of 6 months for this study:

Lorrie Backer, Darlene Coyne, Vicki D'Amico, Diana Delatour, Marianne Ferraro, Patricia
King, Cecilia Kurland, Claire Mancebo, Cathy Siler, Lisa Tallant, Nicole & Sairina Merino Tsui,
and Nicole Zitron.

Funding:

This work was supported by the Office of Naval Research (N00014-16-1-2276). M.S. is the
owner of Comprehensive Pet Therapy (CPT). ONR provided support in the form of salaries for
authors [PC, MS, & GSB], scan time, and volunteer payment, but did not have any additional
role in the study design, data collection and analysis, decision to publish, or preparation of the
manuscript.

Competing Interests:

G.B. and M.S. own equity in Dog Star Technologies and developed technology used in some of
the research described in this paper. The terms of this arrangement have been reviewed and
approved by Emory University in accordance with its conflict of interest policies. MS is the
owner of Comprehensive Pet Therapy (CPT) but no CPT technology or IP was used in this
research.

Author Contributions

A.P., P.C., and G.B. designed and performed research; A.P, R.C., and G.B. analyzed data; A.P.,
P.C., G.B., & M. S. trained dogs; and A.P., G.B., P.C., R. C., and M.S. wrote the paper.

References

Andics, A., Gabor, A., Gacsi, M., Farago, T., Szabo, D., and Miklosi, A. (2016). Neural
mechanisms for lexical processing in dogs. *Science* 353**,** 1030-1032.
Andics, A., Gacsi, M., Farago, T., Kis, A., and Miklosi, A. (2014). Voice-sensitive regions in the
dog and human brain are revealed by comparative fMRI. *Curr Biol* 24**,** 574-578.
Ann Young, C. (1991). Verbal commands as discriminative stimuli in domestic dogs (Canis
familiaris). *Applied Animal Behaviour Science* 32**,** 75-89.
Avants, B.B., Tustison, N.J., Song, G., Cook, P.A., Klein, A., and Gee, J.C. (2011). A
reproducible evaluation of ANTs similarity metric performance in brain image
registration. *Neuroimage* 54**,** 2033-2044.
Bergelson, E., and Swingley, D. (2012). At 6-9 months, human infants know the meanings of
many common nouns. *Proc Natl Acad Sci U S A* 109**,** 3253-3258.
Berns, G.S., Brooks, A., and Spivak, M. (2013). Replicability and heterogeneity of awake
unrestrained canine FMRI responses. *PLoS One* 8**,** e81698.
Berns, G.S., Brooks, A.M., and Spivak, M. (2012). Functional MRI in awake unrestrained dogs.
PLoS One 7**,** e38027.
Berns, G.S., Brooks, A.M., and Spivak, M. (2015). Scent of the familiar: An fMRI study of
canine brain responses to familiar and unfamiliar human and dog odors. *Behav Processes*
110**,** 37-46.
Bloom, P. (2004). Behavior. Can a dog learn a word? *Science* 304**,** 1605-1606.

Brazdil, M., Dobsik, M., Mikl, M., Hlustik, P., Daniel, P., Pazourkova, M., Krupa, P., and
Rektor, I. (2005). Combined event-related fMRI and intracerebral ERP study of an
auditory oddball task. *Neuroimage* 26**,** 285-293.
Cacciaglia, R., Escera, C., Slabu, L., Grimm, S., Sanjuan, A., Ventura-Campos, N., and Avila, C.
(2015). Involvement of the human midbrain and thalamus in auditory deviance detection.
Neuropsychologia 68**,** 51-58.
Cook, P.F., Brooks, A., Spivak, M., and Berns, G.S. (2016a). Regional brain activations in
awake unrestrained dogs. *Journal of Veterinary Behavior-Clinical Applications and*
Research 16**,** 104-112.
Cook, P.F., Prichard, A., Spivak, M., and Berns, G.S. (2016b). Awake canine fMRI predicts
dogs' preference for praise vs food. *Soc Cogn Affect Neurosci* 11**,** 1853-1862.
Cook, P.F., Spivak, M., and Berns, G.S. (2014). One pair of hands is not like another: caudate
BOLD response in dogs depends on signal source and canine temperament. *PeerJ* 2**,**
e596.
Cox, R.W., Chen, G., Glen, D.R., Reynolds, R.C., and Taylor, P.A. (2017). FMRI Clustering in
AFNI: False-Positive Rates Redux. *Brain Connect* 7**,** 152-171.
Cuaya, L.V., Hernandez-Perez, R., and Concha, L. (2016). Our Faces in the dog's brain:
Functional imaging reveals temporal cortex activation during perception of human faces.
PLoS One 11**,** e0149431.
D'aniello, B., Scandurra, A., Alterisio, A., Valsecchi, P., and Prato-Previde, E. (2016). The
importance of gestural communication: a study of human-dog communication using
incongruent information. *Anim Cogn* 19**,** 1231-1235.
Datta, R., Lee, J., Duda, J., Avants, B.B., Vite, C.H., Tseng, B., Gee, J.C., Aguirre, G.D., and
Aguirre, G.K. (2012). A digital atlas of the dog brain. *PLoS One* 7**,** e52140.
Dilks, D.D., Cook, P., Weiller, S.K., Berns, H.P., Spivak, M., and Berns, G.S. (2015). Awake
fMRI reveals a specialized region in dog temporal cortex for face processing. *PeerJ* 3**,**
e1115.
Eklund, A., Nichols, T.E., and Knutsson, H. (2016). Cluster failure: Why fMRI inferences for
spatial extent have inflated false-positive rates. *Proc Natl Acad Sci U S A* 113**,** 7900-
7905.
Friederici, A.D., Meyer, M., and Von Cramon, D.Y. (2000). Auditory language comprehension:
An event-related fMRI study on the processing of syntactic and lexical information.
Brain Lang 74**,** 289-300.
Fukuzawa, M., Mills, D.S., and Cooper, J.J. (2005). The effect of human command phonetic
characteristics on auditory cognition in dogs (Canis familiaris). *J Comp Psychol* 119**,**
117-120.
Goldman, R.I., Wei, C.Y., Philiastides, M.G., Gerson, A.D., Friedman, D., Brown, T.R., and
Sajda, P. (2009). Single-trial discrimination for integrating simultaneous EEG and fMRI:
identifying cortical areas contributing to trial-to-trial variability in the auditory oddball
task. *Neuroimage* 47**,** 136-147.
Hanke, M., Halchenko, Y.O., Sederberg, P.B., Hanson, S.J., Haxby, J.V., and Pollmann, S.
(2009). PyMVPA: A python toolbox for multivariate pattern analysis of fMRI data.
Neuroinformatics 7**,** 37-53.
Howell, T.J., Conduit, R., Toukhsati, S., and Bennett, P. (2012). Auditory stimulus
discrimination recorded in dogs, as indicated by mismatch negativity (MMN). *Behav*
Processes 89**,** 8-13.

Humphries, C., Binder, J.R., Medler, D.A., and Liebenthal, E. (2006). Syntactic and semantic
modulation of neural activity during auditory sentence comprehension. *J Cogn Neurosci*
18**,** 665-679.
Kaminski, J., Call, J., and Fischer, J. (2004). Word learning in a domestic dog: evidence for "fast
mapping". *Science* 304**,** 1682-1683.
Keuleers, E., and Brysbaert, M. (2010). Wuggy: A multilingual pseudoword generator. *Behav*
Res Methods 42**,** 627-633.
Kiehl, K.A., Laurens, K.R., Duty, T.L., Forster, B.B., and Liddle, P.F. (2001). An event-related
fMRI study of visual and auditory oddball tasks. *Journal of Psychophysiology* 15**,** 221-
240.
Kotz, S. (2002). Modulation of the lexical–semantic network by auditory semantic priming: An
event-related functional MRI study. *NeuroImage* 17**,** 1761-1772.
Linden, D.E., Prvulovic, D., Formisano, E., Vollinger, M., Zanella, F.E., Goebel, R., and Dierks,
T. (1999). The functional neuroanatomy of target detection: an fMRI study of visual and
auditory oddball tasks. *Cereb Cortex* 9**,** 815-823.
Mahmoudi, A., Takerkart, S., Regragui, F., Boussaoud, D., and Brovelli, A. (2012). Multivoxel
pattern analysis for FMRI data: A review. *Comput Math Methods Med* 2012**,** 961257.
Markman, E.M., and Abelev, M. (2004). Word learning in dogs? *Trends Cogn Sci* 8**,** 479-481;
discussion 481.
Mills, D.S. (2015). What's in a word? A review of the attributes of a command affecting the
performance of pet dogs. *Anthrozoös* 18**,** 208-221.
Misaki, M., Kim, Y., Bandettini, P.A., and Kriegeskorte, N. (2010). Comparison of multivariate
classifiers and response normalizations for pattern-information fMRI. *Neuroimage* 53**,**
103-118.
Müller, C.A., Schmitt, K., Barber, A.L.A., and Huber, L. (2015). Dogs can discriminate
emotional expressions of human faces. *Current Biology* 25**,** 601-605.
Persson, M.E., Roth, L.S., Johnsson, M., Wright, D., and Jensen, P. (2015). Human-directed
social behaviour in dogs shows significant heritability. *Genes Brain Behav* 14**,** 337-344.
Pilley, J.W., and Reid, A.K. (2011). Border collie comprehends object names as verbal referents.
Behav Processes 86**,** 184-195.
Raettig, T., and Kotz, S.A. (2008). Auditory processing of different types of pseudo-words: an
event-related fMRI study. *Neuroimage* 39**,** 1420-1428.
Slotnick, S.D. (2017). Cluster success: fMRI inferences for spatial extent have acceptable false-
positive rates. *Cogn Neurosci* 8**,** 150-155.
Stelzer, J., Chen, Y., and Turner, R. (2013). Statistical inference and multiple testing correction
in classification-based multi-voxel pattern analysis (MVPA): random permutations and
cluster size control. *Neuroimage* 65**,** 69-82.
Varoquaux, G. (2017). Cross-validation failure: Small sample sizes lead to large error bars.
Neuroimage.
Waxman, S., Fu, X., Arunachalam, S., Leddon, E., Geraghty, K., and Song, H.J. (2013). Are
nouns learned before verbs? Infants provide insight into a longstanding debate. *Child Dev*
Perspect 7**,** 155-159.
Zaine, I., Domeniconi, C., and Costa, A.R.A. (2014). Exclusion performance in visual simple
discrimination in dogs (Canis familiaris). *Psychology & Neuroscience* 7**,** 199-206.

FIGURES

556
557 **Fig. 1. Individual dogs and their trained objects.** All 12 dogs successfully trained to retrieve
558 two objects using object names as verbal referents.

Fig. 2. Experimental design. A) Experimental setup with mirror relay projected words onto MRI surface. Owner is facing the projected word and her dog while the experimenter controls the presentation of words and objects to the owner. B) Trial timeline indicating spoken word over 5 s, 3-8 s delay, 5 s presentation of object, 3 s for the dog to interact with the object, followed by a 6 s intertrial interval.

Fig. 3. Individual performance on two object discrimination tests. Tests were conducted on the day of the fMRI scan. Each dog's Object 1 is in black, object 2 is in grey. All dogs performed significantly greater than chance, with the dog's greater performance or owner's report of their preference for one object over the other designating object 1.

570
Fig 4 Whole brain group map showing unthresholded activation to all words versus all
objects. **A)** Location of crosshairs in superior temporal lobe on average image of all dogs. **B)**
Sagittal view of left hemisphere. Colors represent T-statistics. The primary auditory region
extending into the parietotemporal area showed greater activation to words (*red*), whereas
parietal and occipital areas showed greater activation to objects (*blue*). **C)** Dorsal view. **D)**
Transverse view.

577
Fig. 5. Whole brain response to [pseudowords – words] contrast. Whole brain analysis
revealed significant activation within a parietotemporal region including primary auditory cortex
and neighboring regions. **A)** Location of crosshairs on average image of all dogs without overlay.
B) Sagittal view of right hemisphere. Colors represent T-statistics. With a single voxel
significance of 0.005, the clusterwise significance (Right: 839 voxels; Left: 43 voxels) corrected
across the whole brain was $P = 0.005$ for the right hemisphere, though activation seemed
bilateral. **C)** Dorsal view. **D)** Transverse view.

585
Fig. 6. Aggregate performance of searchlight MVPA classifier for word1 and word2 across
dogs. Color intensity indicates fraction of dogs with informative voxels at each location. The
image is thresholded such that only voxels that were informative for more than one dog are
shown. This map showed four clusters: posterior thalamus/brainstem; amygdala; left
temporoparietal junction; and left dorsal caudate nucleus. The temporoparietal junction appears
similar to human angular gyrus and could be a potential site for receptive language processing in
dogs.

Fig. 7. **Dogs with informative voxels for word1 and word2 in the left temporal and parietal lobes.** Color intensity indicates classification accuracy at each location, thresholded ≥ 0.63. Seven dogs displayed clusters in the left temporal and parietal lobes, suggesting some heterogeneity in the location underlying word discrimination.