1 **Enhancer connectome in primary human cells reveals target genes of disease-**
2 **associated DNA elements**
3

Maxwell R. Mumbach[1,2,12], Ansuman T. Satpathy[1,3,12], Evan A. Boyle[1,2,12], Chao Dai[1,12],
Benjamin G. Gowen[4,5], Seung Woo Cho[1], Michelle L. Nguyen[6], Adam J. Rubin[7], Jeffrey
M. Granja[1,2], Katelynn R. Kazane[4,5], Yuning Wei[1], Trieu Nguyen[8], Peyton G.
Greenside[2], M. Ryan Corces[1], Josh Tycko[2], Dimitre R. Simeonov[6,10], Nabeela Suliman[2],
Rui Li[1], Jin Xu[1], Ryan A. Flynn[1], Anshul Kundaje[2], Paul A. Khavari[7], Alexander
Marson[5,6,8], Jacob E. Corn[4,5], Thomas Quertermous[9], William J. Greenleaf[1,2,8,11,13],
Howard Y. Chang[1,7,13]
11
12 **AFFILIATIONS**
13
[1] Center for Personal Dynamic Regulomes, Stanford University School of Medicine,
Stanford, CA 94305, USA
[2] Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305,
USA
[3] Department of Pathology, Stanford University School of Medicine, Stanford, CA 94305,
USA
[4] Department of Molecular and Cell Biology, University of California, Berkeley, CA
94720, USA
[5] Innovative Genomics Institute, University of California, Berkeley, CA, 94720, USA
[6] Department of Microbiology and Immunology, University of California, San Francisco,
CA 94143, USA
[7] Program in Epithelial Biology, Stanford University School of Medicine, Stanford, CA,
94305, USA.
[8] Chan Zuckerberg Biohub, San Francisco, CA 94158, USA
[9] Department of Medicine, Division of Cardiovascular Medicine, Stanford University
School of Medicine, Stanford, California 94305, USA.
[10] Biomedical Sciences Graduate Program, University of California, San Francisco, CA
94143, USA
[11] Department of Applied Physics, Stanford University, Stanford, CA, 94305, USA
[12] These authors contributed equally
[13] These authors jointly directed this work
35
36 *To whom correspondence should be addressed. E-mail: howchang@stanford.edu and
37 wjg@stanford.edu

ABSTRACT

The challenge of linking intergenic mutations to target genes has limited molecular
understanding of human diseases. Here, we show that H3K27ac HiChIP generates
high-resolution contact maps of active enhancers and target genes in rare primary
human T cell subtypes and coronary artery smooth muscle cells. Differentiation of naïve
T cells to T helper 17 cells or regulatory T cells creates subtype-specific enhancer-
promoter interactions, specifically at regions of shared DNA accessibility. These data
provide a principled means of assigning molecular functions to autoimmune and
cardiovascular disease risk variants, linking hundreds of noncoding variants to putative
gene targets. Target genes identified with HiChIP are further supported by CRISPR
interference and activation at linked enhancers, by the presence of expression
quantitative trait loci, and by allele-specific enhancer loops in patient-derived primary
cells. The majority of disease-associated enhancers contact genes beyond the nearest
gene in the linear genome, leading to a four-fold increase of potential target genes for
autoimmune and cardiovascular diseases.

Gene expression programs are intimately linked to the hierarchical organization
of the genome. In mammalian cells, each chromosome is organized into hundreds of
megabase-sized topologically associated domains (TADs), which are conserved from
early stem cells to differentiated cell types[1]. Within this invariant TAD scaffold, cell type-
specific enhancer-promoter (E-P) interactions establish regulatory gene expression
programs[2]. Standard methods require tens of millions of cells to obtain high-resolution
interaction maps and confidently assign E-P contacts[3–5]. Thus, the principles that
govern E-P conformation in disease-relevant patient samples are incompletely
understood. This gap in understanding is particularly problematic for interpreting the
molecular functions of inherited risk factors for common human diseases, which reside
in intergenic enhancers or other non-coding DNA features in up to 90% of cases[6–9].
Such disease-relevant enhancers may not influence the expression of the nearest gene
(often reported as the default target in the literature), and instead act in a cell-type
specific manner on distant target genes residing up to hundreds of kilobases (kb)
away[2,10–14]. Recently, systematic perturbations of regulatory elements in select gene loci
have shown that effects of individual regulatory elements on gene activity can be
predicted from the combination of (i) enhancer activity [marked by histone H3 lysine 27
acetylation (H3K27ac) level] and (ii) enhancer-target looping[5,15]. Here we leverage this
insight to capture the combination of these two types of information genome-wide in a
single assay, mapping the enhancer connectome in disease-relevant primary human
cells.
110
111   **RESULTS**
112
113   **H3K27ac HiChIP identifies functional enhancer interactions**
114
We recently developed HiChIP, a method for sensitive and efficient analysis of
protein-centric chromosome conformation[16]. Cohesin HiChIP in GM12878 cells
identified similar numbers of loops as *in situ* Hi-C (~10,000) with high correlation (R =
0.83), demonstrating that HiChIP captures loops with high sensitivity and specificity.
Here, we evaluated the enhancer and promoter-associated mark H3K27ac[17–19] as a
candidate factor to selectively interrogate E-P interactions genome-wide. We performed
H3K27ac HiChIP in mouse embryonic stem (mES) cells to compare to cohesin HiChIP
(**Supplementary Fig. 1a, Supplementary Table 1**)[16]. 3,552 of 4,191 H3K27ac HiChIP
loops in mES cells were also identified by cohesin HiChIP. The H3K27ac-biased loops
($\log_2$ fold-change > 1) spanned shorter distances than cohesin-biased loops, and were
enriched for H3K27ac ChIP-seq peaks (78.9%; **Supplementary Fig. 1b-f,**
Supplementary Table 2). Moreover, systematic titration of input material showed
H3K27ac HiChIP retained high signal fidelity and reproducibility from 25 million to
50,000 cells as input material (loop signal correlation = 0.918; **Supplementary Figs. 2**
and 3). Therefore, H3K27ac HiChIP identifies high-confidence chromatin loops focused
around enhancer interactions from limited cell numbers.
In order to capture (i) conformational change during T cell differentiation and (ii)
cell type-specific chromatin contacts of autoimmune risk variants in protective and

pathogenic T cell types, we performed H3K27ac HiChIP on primary human Naïve T
cells (CD4$^+$CD45RA$^+$CD25$^-$CD127^{hi}), regulatory T cells (T$_{reg}$; CD4$^+$CD25$^+$CD127^{low}) and
T helper 17 cells (T$_H$17; CD4$^+$CD45RA$^-$CD25$^-$CD127^{hi}CCR6$^+$CXCR5$^-$) directly isolated
from donors (**Fig. 1a,b and Supplementary Fig. 4a**)[20,21]. T$_H$17 cells were sorted to
include autoimmune disease-relevant pathogenic T$_H$17 cells and to exclude follicular
helper T cells with a distinct surface phenotype and immune function (**Supplementary**
Fig. 4a)[22–24]. Peripheral blood CD4$^+$ T cells were isolated from three healthy subjects,
isolated by FACS, and subjected to H3K27ac HiChIP. HiChIP libraries from each subset
were high quality; greater than 40% of the reads represented unique paired-end tags
(PETs) (**Supplementary Fig. 4b-d and Supplementary Table 1**). Furthermore,
libraries exhibited high 1D signal enrichment at enhancers and promoters, and globally
recapitulated publically available H3K27ac ChIP-seq datasets (74.7% overlap of ChIP-
seq and 1D HiChIP peaks; **Fig. 1c**)[25]. Inspection of the interaction matrix at
progressively higher resolution revealed chromatin compartments, TADs, and focal
loops, as previously reported in high-resolution Hi-C and HiChIP analyses from cell lines
(**Fig. 1b**)[4,16]. Importantly, H3K27ac HiChIP maps were capable of identifying focal
interactions at 1 kb resolution, which is comparable to *in situ* Hi-C maps generated from
100-fold more cells and sequenced to 13-fold greater depth[4] (**Fig. 1b**).
Previous saturation perturbation screens demonstrated that functional enhancers
can be identified by integrating H3K27ac ChIP-seq signal with chromosome
conformation contact strength (Hi-C)[5]. Since H3K27ac HiChIP combines these two
components into one assay, we reasoned that HiChIP signal, which we term Enhancer
Interaction Signal (EIS), should identify functional regulatory elements. To validate this
prediction, we first generated H3K27ac HiChIP maps in a chronic myelogenous
leukemia cell line (K562) as a direct comparison to published high-resolution CRISPR
interference (CRISPRi) screens[5]. We then examined the 3D enhancer landscape of the
MYC and *GATA1* loci using virtual 4C (v4C) analysis, where a specific genomic position
is set as an anchor viewpoint, and all interactions occurring with that anchor are
visualized in 2D[16]. v4C analysis of the *MYC* promoter demonstrated that EIS in K562
cells captured all functional enhancers identified in the CRISPRi screen (**Fig. 2a**).
Analysis of the *GATA1* locus demonstrated a similar agreement between both methods
(**Fig. 2b**). Quantitatively, EIS in K562 cells was significantly correlated with CRISPRi
score in the same cell type, whereas EIS in GM12878 (GM; B cell lymphoblast) cells
was not correlated with K562 CRISPRi (Spearman's rho = 0.332 and 0.145; p-value =
9.25 x 10^{-5} and 0.1246; **Fig. 2c**).
We found the enhancer landscapes of the *MYC* promoter to be highly cell-type
specific. v4C analysis of the *MYC* promoter in GM and My-La (CD4$^+$ T cell leukemia)
cells showed dramatically different regulatory interactions with the promoter compared
to K562 cells (**Fig. 2d**). To validate EIS specificity, we performed CRISPRi experiments
in GM cells using sgRNAs targeting enhancers identified in either GM or My-La HiChIP
maps as well as a positive control sgRNA targeting the *MYC* promoter and a negative
control sgRNA targeting lambda phage sequence (**Fig. 2e**). As expected, we found that
CRISPRi of GM, but not My-La, enhancers impacted *MYC* expression and cell growth in
GM cells (**Fig. 2e**).

Finally, we focused on the *CD69* locus, where a high-resolution CRISPRa screen
identified three enhancers upstream of the transcription start site[26]. These sites were
also identified by Naïve T cell H3K27ac HiChIP. Moreover, HiChIP identified four
additional distal enhancers that were outside the region spanned by the sgRNA tiling
array (**Fig. 2f and Supplementary Fig. 5**). To functionally validate these novel
enhancers, we performed CRISPRa experiments in Jurkat cells with sgRNAs targeting
these enhancers, the *CD69* promoter, the *KLRF2* promoter as a locus negative control,
and a non-human genome-targeting negative control. We observed a significant
increase in *CD69* RNA and protein levels in the four HiChIP enhancers compared to
negative controls (**Fig. 2g and Supplementary Fig. 5**). Interestingly, two of the four
identified novel enhancers were within promoter regions of distant genes. These
findings are in line with previous reports that identified widespread distal gene regulatory
functions of promoters genome-wide[27,28]. Altogether, these results suggest that
H3K27ac HiChIP EIS identifies functional regulatory elements, and that enhancers that
regulate a gene of interest can differ significantly between cell-types.
192
193 **Landscape of enhancer interactions in primary T cells**
194
We examined global features of the enhancer connectome associated with
cellular differentiation from Naïve T cells to either T_H17 cells or T_{reg} cells. We identified a
total of 10,706 high confidence loops in the union set of the three cell types
(**Supplementary Table 2**). Analysis of loop read support between biological replicates
demonstrated high reproducibility (**Supplementary Fig. 4c**), and ~91% of loop anchors
were associated with either a promoter or enhancer[29], as expected, with a median
distance of 130 kb (**Supplementary Fig. 6a,b**). Importantly, high-resolution E-P
connectivity maps revealed several features that could not be discerned from 1D
epigenomic data (i.e. H3K27ac ChIP-seq or ATAC-seq; **Fig. 3a**). These features
included: (i) 'enhancer skipping': enhancers that have stronger EIS with a more distal
target promoter, (ii) higher order structures such as 'enhancer cliques' (related to loop
cliques[30]): multiple regulatory elements that have strong EIS with a single target
promoter, (iii) promoter to promoter interactions[13,31], and (iv) 'enhancer switching':
enhancers that exhibit differential EIS with a target promoter in a cell type-specific
manner (**Fig. 3a**).
We found that EIS contacts were very cell type-specific. After quantile-quantile
normalization of contact reads at high-confidence loops (correcting for false positives
caused by 1D fragment visibility; **Methods**), we focused on the top and bottom 5% of
EIS ranked by cell-type bias for each pair-wise comparison (**Supplementary Figs. 6c-g
and 7, Supplementary Tables 3-4**). Cell type-specific enhancer loop anchors revealed
genes encoding canonical T cell subtype TFs and effector molecules (**Fig. 3b,
Supplementary Figs. 8 and 9**). Deeper v4C analysis of shared and cell type-specific
loci pinpointed regulatory elements interacting with each gene promoter of interest as
well as local conformational landscape changes (**Supplementary Figs. 8 and 9**). TF
motifs located within cell type-specific loop anchors were enriched for TFs known to
drive T cell subtype differentiation and nominated novel TFs involved in regulation (**Fig.**

3c). Furthermore, cell type EIS bias was associated with differential expression of genes
located within corresponding EIS anchors for the same cell type (Spearman's rho =
0.242 and 0.207; p-value = 4 x 10^{-15} and 2 x 10^{-11}; **Fig. 3d**).
Cell type-specific EIS may be driven by cell type-specific enhancer activation
(based on H3K27ac ChIP-seq) or stable enhancer activation with cell type-specific
looping (Hi-C) in a gene specific manner. We first examined H3K27ac ChIP-seq at
differential EIS anchors and found that many biased H3K27ac HiChIP interactions also
exhibited biased ChIP-seq signal, as expected. 58.5% of Naïve-biased loops contain at
least one Naïve-biased ChIP-seq peak (log_2 fold change > 1) located on the anchors.
Similarly, 66.7% of T_H17-biased and 67.8% of T_{reg}-biased interaction anchors were cell
type-specific in 1D (**Supplementary Fig. 10a**). Therefore, while on average ~64% of
the differential EIS corresponded to change in 1D data, ~36% were likely also driven by
change in 3D chromatin loop strength. To further assess the contribution of cell type-
specific 3D signal to EIS, we examined HiChIP 1D signal at differential EIS anchors. We
found that HiChIP 1D signal correlated better with ChIP-seq signal than EIS, with a
higher likelihood of differential ChIP-seq signal overlapping differential HiChIP 1D signal
compared to 3D, suggesting EIS bias is in part driven by 3D changes (**Supplementary
Fig. 10b**).
We asked whether the integration of reference cell line Hi-C data with primary T
cell H3K27ac ChIP-seq could recapitulate HiChIP EIS in primary T cells. We binned GM
Hi-C loops with increasing primary T cell ChIP-seq signal at loop anchors and then
determined the overlap of loops in each bin with loops derived from H3K27ac HiChIP.
As expected, increased ChIP-seq signal at the Hi-C anchors led to increased overlap
with the HiChIP loops. However, the overlap was lower in all T cell subtypes compared
to the same analysis performed using GM HiChIP data. These observations
demonstrate that cell-type specific 3D interactions can impact EIS independent of
differences in 1D ChIP-seq signal (**Supplementary Fig. 10c**). Similarly, previously
generated enhancer-promoter maps obtained from bulk T cells did not identify T cell
subtype-specific interactions obtained using H3K27ac HiChIP. To assess the unique
information obtained through cell type-specific interaction maps, we compared promoter
Capture Hi-C maps in bulk $CD4^+$ T cells to H3K27ac HiChIP maps in Naïve, T_H17, and
T_{reg} cells[14]. Strikingly, the most cell type-specific loops in T_H17 and T_{reg} (16-fold
enriched) demonstrated a low discovery rate in promoter Capture Hi-C T cells (11.83%
in 415 loops and 13.83% in 373 loops, respectively; **Supplementary Fig. 10d**). Many of
these subset-specific interactions included genomic loci encoding functionally important
effector genes, such as *LRRC32*. The *LRRC32* locus contains T_{reg}-specific loops that
are neither visualized in HiChIP maps from Naïve or T_H17 cells nor in bulk $CD4^+$
promoter Capture Hi-C maps (**Supplementary Fig. 10e**). Since primary human T_H17
and T_{reg} cells are present in human blood with low frequency, it would also be
challenging to generate subset-specific promoter Capture Hi-C maps with published
promoter Capture Hi-C protocols. In summary, EIS is derived from a combination of 1D
ChIP-seq and 3D interaction signal and cannot be accurately predicted from 3D maps in
reference cell lines or unsorted primary cell datasets.

Cell type-specific EIS can occur at sites of shared chromatin accessibility. Paired
chromatin accessibility profiles by Assay of Transposase-Accessible Chromatin by
sequencing (ATAC-seq)[32] from each T cell subset revealed most cell type-specific loop
anchors had equivalent chromatin accessibility across all three cell types (**Fig. 3e-g**). To
illustrate this finding, we examined the *BACH2* promoter, which exhibits shared
chromatin accessibility at enhancers, but increased EIS in Naïve cells (**Fig. 3e**).
Globally, only 14.2%, 27.8%, and 16.5% of Naïve-, T_H17-, and T_{reg}-biased loops,
respectively, contained at least one biased ATAC-seq peak ($\log_2$ fold change > 1)
located on the anchors. Furthermore, the majority of cell type-specific TF motifs were
observed in shared ATAC-seq peaks within differential interactions, highlighting that
these regions are functioning in T cell differentiation (**Fig. 3f-g**). Altogether, these results
suggest that in highly related – yet functionally distinct – cell types, a portion of
transcriptional control is achieved through differential chromosome looping, rather than
differential chromatin accessibility. This finding is consistent with previous studies which
demonstrated that T cell subset-specific TFs, such as Foxp3, act predominantly at pre-
accessible chromatin sites to establish subset-specific gene expression[33].
280
281　**Enhancer interactions link disease variants to target genes**
282
The high specificity of EIS enabled us to identify putative target genes of
autoimmune disease risk loci in functionally relevant T cell subsets. To achieve this, we
used a previously described list of putatively causal variants associated with 21
autoimmune diseases, known as PICS SNPs, which were fine-mapped based on dense
genotyping data[25]. We determined that PICS autoimmune SNPs were significantly
enriched in T cell loop anchors, with specific autoimmune diseases showing greater
than 5-fold enrichment compared to a shuffled control loop set (**Supplementary Fig.
11**). Next, we constructed a set of all possible connections between autoimmune risk
SNPs and TSS within 1 Mb and measured the EIS for each SNP-TSS pair (**Fig. 4a**). We
aggregated these signals to determine the overall interaction activity in each T cell
subtype in each disease (**Fig. 4b**). We observed high interaction strength enrichments
and cell type specificity in autoimmune disease SNPs, but low enrichment and cell
specificity in non-immune traits (**Fig. 4b**). To further visualize HiChIP bias in shared or
differential enhancers, we analyzed SNP-TSS interactions grouped by their presence
near H3K27ac ChIP-seq peaks (**Supplementary Fig. 12a,b**). We observed a large
number of active SNP-TSS pairs that were present in regulatory regions that were
shared between T effector cell types (T_{reg} and T_H17), while relatively less EIS signal was
observed in SNPs located in cell-type specific enhancers, supporting the concept that
many autoimmune disease variants impact common T cell effector/activation
pathways[25,34]. Notably, SNPs present in enhancers shared across all three cell types
could still be distinguished by HiChIP bias (**Supplementary Fig. 12a,b**). For example,
although we could not detect cell type bias at risk loci for Alopecia Areata using
H3K27ac ChIP-seq (**Supplementary Fig. 12a,b** and ref. 3), H3K27ac HiChIP identified
increased SNP-TSS activity in T_{reg} cells among shared T cell enhancers, consistent with
several studies identifying the crucial role of this cell type in disease pathogenesis[35].

Importantly, autoimmune signal enrichments were not readily apparent from 1D
H3K27ac ChIP-seq peaks, aggregated ChIP-seq signal within the TAD containing the
SNP, nor cell line H3K27ac HiChIP datasets (**Fig. 4b and Supplementary Fig. 12c**).
Therefore, examining 3D disease variant interactions may capture cell type biases more
robustly than 1D epigenomic data. Finally, to validate our findings with an orthogonal
dataset, we performed SNP-TSS EIS analysis on an overlapping set of autoimmune
disease-associated SNPs obtained from the NHLBI GRASP catalog and observed
similar enrichments of specific T cell subsets (**Supplementary Fig. 12d**).
We leveraged HiChIP to identify potential gene targets of intergenic SNPs, which
have classically been paired to the nearest neighboring gene. We overlapped the SNP-
TSS pairs with loops to call a discrete set of target pairs. We then performed differential
analysis on the SNP-TSS loops to ascertain bias for specific T cell subsets (**Fig. 4c and
Supplementary Table 5**). Examples of biased SNP-TSS pairs included *FOXO1* in
Naïve T cells (rs9603754), *BATF* (rs2300604) in Memory T cells, *CTLA4* (rs10186048)
in T_{reg} cells, and *IL2* (rs7664452) in T_H17 cells (**Fig. 4c and Supplementary Table 5**).
Next, we sought to characterize the connectivity landscape of the SNP-TSS loops. We
identified an average of 1.75 gene targets per autoimmune SNP (ranging from 0 to over
10 target genes), while non-immune traits did not demonstrate an increase in targets
(0.33 genes per SNP; **Supplementary Fig. 12e**). For 684 autoimmune intergenic SNPs,
we identified a total of 2,597 HiChIP target genes, representing a four-fold increase in
target genes for known disease SNPs (**Fig. 4d**). Only 367 (~14%) of all targets were the
nearest gene to the SNP, while approximately ~86% of SNPs skipped at least one gene
to reach a predicted target TSS (**Supplementary Fig. 12e**). Furthermore, approximately
~45% of SNP to HiChIP target interactions had increased signal compared to the same
SNP to nearest gene, despite distance biases.
333
334 **Target gene validation by eQTL and CRISPRi**
335
HiChIP enhancer-target gene interactions can be validated using previously
identified point mutations that alter expression at distantly located genes in T cells—i.e.
expression quantitative trait loci (eQTL)[36]. For example, the celiac disease-associated
SNP rs2058660 impacts the expression of the inflammatory cytokine receptor genes
IL18RAP, *IL18R1*, *IL1RL1*, and *IL1RL2*, which are known regulators of intestinal T cell
differentiation and response[37]. HiChIP EIS revealed contacts between rs2058660 and
each of these predicted gene promoters (**Supplementary Fig. 13a**). Similarly, the
Crohn's disease risk variant rs6890268 and the multiple sclerosis (MS) risk variant
rs12946510 impact the expression of *PTGER4* and *IKZF3*, respectively, and H3K27ac
HiChIP also demonstrated clear contacts between these SNPs and their predicted
promoter (**Supplementary Fig. 13a**). Globally, HiChIP contact signal was increased in
eQTLs in T cells compared to a distance-matched background loop set (p-value < 2.2 x
10^{-16}; **Fig. 4e)** or to eQTLs identified in an unrelated cell type (liver; p-value < 2.2 x 10^{-}
^{16}). The overlap of HiChIP and eQTL loci provides support for chromosome interactions
as a physical basis for distal eQTLs[10–12] and further validates the HiChIP approach to
assign enhancer-target gene relationships.

We next sought to directly validate HiChIP SNP-gene targets using CRISPRi in
My-La cells. First, we focused on three loci of interest in primary T cells and then
confirmed that the SNP-TSS loops were also present in My-La cells (**Fig. 4f and**
Supplementary Fig. 13b). We then targeted sgRNAs to these SNP-containing
enhancers, as well as positive control sgRNAs to the HiChIP target gene promoters and
a negative control sgRNA targeting lambda phage sequence. As expected, we observed
a significant reduction of RNA levels in the HiChIP target genes upon CRISPRi of its
SNP-containing enhancer (**Fig. 4f**).

Fine-mapping of disease-associated DNA variants

Since SNP-TSS HiChIP signal is capable of identifying target genes of candidate
SNPs, we asked whether TSS-SNP HiChIP signals could also be used to nominate
functional causal variants within haplotype blocks in a reciprocal manner. We first
performed a proof-of-principle analysis using fine-mapped SNPs associated with
inflammatory bowel disease (IBD)[38] or Type 1 Diabetes (T1D)[39] as well as high
confidence PICS SNPs and examined EIS from putatively causal SNPs to all gene
promoters within 300 kb. EIS from putatively causal SNPs to gene promoters was
significantly higher than EIS from a distance-matched set of SNPs within the same LD
block to gene promoters (p-value = 2.4×10^{-15}, 8.7×10^{-8}, 3.9×10^{-3} for IBD fine-mapped
SNPs, T1D fine-mapped SNPs, and high confidence PICS, respectively; **Fig. 5a** and
Supplementary Fig. 14a). Next, we assessed the fine-mapping ability of HiChIP EIS at
individual loci of interest. We focused on IBD- and MS-associated SNPs neighboring the
PTGER4 and *SATB1* loci and performed v4C analysis anchored at the gene promoters.
We calculated EIS signal at 1 kb resolution and identified specific regions within the
linkage disequilibrium (LD) blocks that contained the highest EIS to the target
promoters, positioning the likely causal SNPs within these regions (**Fig. 5b** and
Supplementary Fig. 14b). For example, at the *PTGER4* locus (**Fig. 5b**), the ~160 kb
genomic interval spanned by LD SNPs in association with Crohn's disease is refined to
two bins of 3kb and 4kb, which both contain PICS SNPs.
We asked whether complex disease-associated loci containing more than one
gene could be fine-mapped using HiChIP. We focused on two disease-associated
enhancers in between the *STAT1* and *STAT4* gene promoters (**Fig. 5c**). These two
genes encode transcription factors with distinct roles in immune regulation. Signal
transducer and activator of transcription 1 (STAT1) is critical for type I IFN and IFNγ
signaling, whereas STAT4 induces T_H1 differentiation and IFNγ expression[40]. We
investigated bias of these enhancers to *STAT1* and *STAT4* and found that, despite
comparable linear distance and 1D signal at the promoters, the enhancers were
significantly biased to interact with *STAT4*. Next, we fine-mapped the disease
associated SNPs within this locus using 1 kb resolution EIS from the *STAT4* promoter,
and narrowed down candidate functional variants within the two enhancers (**Fig. 5c**). In
summary, HiChIP EIS can nominate functional causal variants within haplotype blocks,
and two-way analysis of target gene identification from an enhancer of interest and high-
resolution interaction maps of that enhancer with its target gene can be used to fine-

map disease-associated loci containing several candidate genes.

Allelic target gene bias of cardiovascular disease variants

Finally, we asked whether this approach could be applied broadly to other
categories of human disease, and whether we could directly test SNP-TSS associations
using allele-specific HiChIP. We generated high-resolution E-P maps from primary
human coronary artery smooth muscle cells (HCASMC), which can be used to inform
variants linked to cardiovascular diseases[41]. First, to validate cell type specificity, we
examined the *TCF21* gene promoter, a transcription factor required for the
differentiation of HCASMC[42] and observed enrichment in HCASMC EIS relative to Naïve
T cells (**Fig. 6a**). We next examined the 9p21.3 locus, which harbors risk associations
with several cardiovascular disorders[43–45]. We found that the promoters of all three
genes in the locus interact with one another and with CAD variant-containing enhancers
located approximately 100 kb upstream of the *CDKN2B* promoter (**Supplementary Fig.**
15). We then generated SNP-TSS target lists using CAD SNPs identified in the
CARDIoGRAMplusC4D study[46]. We again performed differential analysis on the SNP-
TSS loops to ascertain bias for HCASMC versus Naïve T cells (**Fig. 6b**). Overall, 75.1%
of biased HCASMC SNP-TSS pairs were CAD SNPs, while only 5.5% of Naïve T cell
biased SNP-TSS pairs were CAD SNP-TSS loops. Next, we examined the connectivity
of the HCASMC SNP-TSS contacts and identified 1,062 gene targets, of which only 120
(~11%) mapped to the nearest gene. Furthermore, approximately 89% skipped at least
one gene to reach a predicted target TSS, and 64% of SNPs were mapped to more than
a single gene target.
We took advantage of genome phasing information in HCASMC to measure E-P
interactions at allele-specific CAD SNPs, allowing us to examine the functional
consequence of a risk variant compared to its alternative allele in the same nucleus.
First, 4.2% of high confidence loops in HCASMC with no observed mapping bias in the
anchors exhibited significant allelic bias (FDR < 0.05, **Fig. 6c**), consistent with
frequency of allelic imbalance of RNA expression and prior evidence of allele-specific
regulation of specific E-P interactions[47,48]. We leveraged this global E-P allelic bias to
examine the effect of a risk variant compared to its control alternative allele for a set of
CAD-associated SNP-target gene pairs (**Fig. 6d**)[49]. We found that many risk alleles
disrupt enhancer-target gene interactions, but a subset of pathogenic SNPs increased
enhancer-target gene interaction. At CAD risk variant rs1537373 in the 9p21.3 locus,
the risk allele (T) showed increased EIS to the *CDKN2A* promoter as well as an
additional enhancer within the lncRNA *ANRIL* relative to the reference allele (G) (**Fig.**
6e). We further observed increased EIS of the CAD risk variant rs4562997 to an
additional *SMAD3* enhancer 10 kb downstream of the TSS (**Fig. 6e**). The ability to
resolve enhancer connectomes of the risk and reference alleles in the same nucleus
demonstrates that the mutated base in the risk allele suffices to alter enhancer looping
in *cis* in disease-relevant primary cells.

DISCUSSION

440
Here, we developed an approach to define the high-resolution landscape of E-P
regulation in primary human cells. We find that E-P contacts are highly dynamic in
related cell types and often involve genomic elements with shared accessibility.
Accordingly, many complex features of the 3D enhancer connectome cannot simply be
predicted from 1D, which demonstrates that mapping conformation in primary cells can
identify novel regulatory connections underlying gene function in human disease. We
take advantage of this principle to chart the connectivity of autoimmune and
cardiovascular GWAS SNPs and link SNPs to hundreds of potential target genes.
Although non-genic SNPs have previously been paired with their closest neighboring
gene, we find that the majority of these variants can engage in long-distance
interactions, including skipping several promoters to predicted target genes, connecting
to multiple genes, or acting in concert with enhancer cliques to contact a single gene.
Further use of this approach will help to clarify hidden mechanisms of human disease
that are driven by genetic perturbations in non protein-coding DNA elements, which can
now be linked to their cognate gene targets in primary cells.

ACKNOWLEDGEMENTS

458
We thank members of the Chang and Greenleaf laboratories for helpful discussions and
Justin Tumey for artwork. This work was supported by National Institutes of Health
(NIH) P50HG007735 (H.Y.C. and W.J.G.) and U19AI057266 (W.J.G), Human Frontier
Science Program (W.J.G.), Rita Allen Foundation (W.J.G.), and the Scleroderma
Research Foundation (H.Y.C). M.R.M. and E.A.B. acknowledge support from the
National Science Foundation Graduate Research Fellowship. A.T.S. is a Cancer
Research Institute Irvington Fellow supported by the Cancer Research Institute. B.G.G.
was supported by an IGI-AstraZeneca Postdoctoral Fellowship. M.R.C. is supported by
a grant from The Leukemia & Lymphoma Society Career Development Program. J.E.C.
was supported by the Li Ka Shing Foundation and the Heritage Medical Research
Institute. W.J.G. and A.M. are Chan Zuckerberg Biohub investigators. A.M. serves as an
advisor to Juno Therapeutics and PACT Therapeutics and the Marson lab has received
sponsored research support from Juno Therapeutics and Epinomics. A.M. and J.E.C.
are founders of Spotlight Therapeutics. Sequencing was performed by the Stanford
Functional Genomics Facility (NIH S10OD018220). We thank Agilent Technologies for
generating oligo pools for cloning of the CRISPRa gRNAs. We thank the UC Berkeley
High Throughput Screening Facility and Flow Cytometry Facility. H.Y.C. and W.J.G. are
founders of Epinomics and members of its scientific advisory board.

477
478   **AUTHOR CONTRIBUTIONS**

479
M.R.M, A.T.S., W.J.G., and H.Y.C. conceived the project. M.R.M., A.T.S., J.T., and R.L.
performed all genomics assays with help from T.N., M.R.C., N.S., and R.A.F. A.T.S.
performed all sorting for experiments. B.G.G., S.W.C., M.R.M., M.L.N., K.R.K., and
D.R.S., performed all CRISPR validation experiments. E.A.B., C.D., M.R.M., and J.X.

analyzed HiChIP data. J.G., A.T.S., and Y.W. analyzed ATAC-seq data. A.J.R. and
P.G.G. analyzed GWAS SNPs in HiChIP data. A.K., P.A.K., A.M., J.E.C., T.Q., W.J.G.,
and H.Y.C guided experiments and data analysis. M.R.M, A.T.S, E.A.B, C.D., W.J.G,
and H.Y.C. wrote the manuscript with input from all authors.
488
489

MAIN TEXT REFERENCES

1. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485,** 376–380 (2012).

2. van Arensbergen, J., van Steensel, B. & Bussemaker, H. J. In search of the determinants of enhancer-promoter interaction specificity. *Trends Cell Biol.* **24,** 695–702 (2014).

3. Lieberman-Aiden, E. *et al.* Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* **326,** 289–293 (2009).

4. Rao, S. S. P. *et al.* A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* **159,** 1665–1680 (2014).

5. Fulco, C. P. *et al.* Systematic mapping of functional enhancer-promoter connections with CRISPR interference. *Science* aag2445 (2016). doi:10.1126/science.aag2445

6. Altshuler, D., Daly, M. J. & Lander, E. S. Genetic mapping in human disease. *Science* **322,** 881–888 (2008).

7. Kumar, V. *et al.* Human Disease-Associated Genetic Variation Impacts Large Intergenic Non-Coding RNA Expression. *PLOS Genet.* **9,** e1003201 (2013).

8. Edwards, S. L., Beesley, J., French, J. D. & Dunning, A. M. Beyond GWASs: Illuminating the Dark Road from Association to Function. *Am. J. Hum. Genet.* **93,** 779–797 (2013).

9. Welter, D. *et al.* The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res.* **42,** D1001–D1006 (2014).

10. Smemo, S. *et al.* Obesity-associated variants within FTO form long-range functional connections with IRX3. *Nature* **507,** 371–375 (2014).

11. Claussnitzer, M. *et al.* FTO Obesity Variant Circuitry and Adipocyte Browning in Humans. *N. Engl. J. Med.* **373,** 895–907 (2015).

12.   Grubert, F. *et al.* Genetic Control of Chromatin States in Humans Involves Local and

Distal Chromosomal Interactions. *Cell* **162,** 1051–1065 (2015).

13.   Mifsud, B. *et al.* Mapping long-range promoter contacts in human cells with high-

resolution capture Hi-C. *Nat. Genet.* **47,** 598–606 (2015).

14.   Javierre, B. M. *et al.* Lineage-Specific Genome Architecture Links Enhancers and Non-

coding Disease Variants to Target Gene Promoters. *Cell* **167,** 1369–1384.e19 (2016).

15.   Sanjana, N. E. *et al.* High-resolution interrogation of functional elements in the

noncoding genome. *Science* **353,** 1545–1549 (2016).

16.   Mumbach, M. R. *et al.* HiChIP: efficient and sensitive analysis of protein-directed

genome architecture. *Nat. Methods* **advance online publication,** (2016).

17.   Heintzman, N. D. *et al.* Distinct and predictive chromatin signatures of

transcriptional promoters and enhancers in the human genome. *Nat. Genet.* **39,** 311–318

(2007).

18.   Creyghton, M. P. *et al.* Histone H3K27ac separates active from poised enhancers and

predicts developmental state. *Proc. Natl. Acad. Sci. U. S. A.* **107,** 21931–21936 (2010).

19.   Rada-Iglesias, A. *et al.* A unique chromatin signature uncovers early developmental

enhancers in humans. *Nature* **470,** 279–283 (2011).

20.   Ferraro, A. *et al.* Interindividual variation in human T regulatory cells. *Proc. Natl.*

Acad. Sci. U. S. A. **111,** E1111-1120 (2014).

21.   Arvey, A. *et al.* Genetic and epigenetic variation in the lineage specification of

regulatory T cells. *eLife* **4,** e07571 (2015).

22.   Bettelli, E. *et al.* Reciprocal developmental pathways for the generation of

pathogenic effector TH17 and regulatory T cells. *Nature* **441,** 235–238 (2006).

23.　　Acosta-Rodriguez, E. V. *et al.* Surface phenotype and antigenic specificity of human

interleukin 17-producing T helper memory cells. *Nat. Immunol.* **8,** 639–646 (2007).

24.　　Morita, R. *et al.* Human blood CXCR5(+)CD4(+) T cells are counterparts of T

follicular cells and contain specific subsets that differentially support antibody secretion.

Immunity **34,** 108–121 (2011).

25.　　Farh, K. K.-H. *et al.* Genetic and epigenetic fine mapping of causal autoimmune

disease variants. *Nature* **518,** 337–343 (2015).

26.　　Simeonov, D. R. *et al.* Discovery of an autoimmunity-associated IL2RA enhancer by

unbiased targeting of transcriptional activation. *bioRxiv* 091843 (2016).

doi:10.1101/091843

27.　　Engreitz, J. M. *et al.* Local regulation of gene expression by lncRNA promoters,

transcription and splicing. *Nature* **539,** 452–455 (2016).

28.　　Dao, L. T. M. *et al.* Genome-wide characterization of mammalian promoters with

distal enhancer functions. *Nat. Genet.* **49,** 1073–1081 (2017).

29.　　Ernst, J. & Kellis, M. ChromHMM: automating chromatin-state discovery and

characterization. *Nat. Methods* **9,** 215–216 (2012).

30.　　Sanborn, A. L. *et al.* Chromatin extrusion explains key features of loop and domain

formation in wild-type and engineered genomes. *Proc. Natl. Acad. Sci. U. S. A.* **112,**

E6456-6465 (2015).

31.　　Li, G. *et al.* Extensive promoter-centered chromatin interactions provide a

topological basis for transcription regulation. *Cell* **148,** 84–98 (2012).

32.   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J. Transposition

of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-

binding proteins and nucleosome position. *Nat. Methods* **10,** 1213–1218 (2013).

33.   Samstein, R. M. *et al.* Foxp3 exploits a pre-existent enhancer landscape for

regulatory T cell lineage specification. *Cell* **151,** 153–166 (2012).

34.   Ye, C. J. *et al.* Intersection of population variation and autoimmunity genetics in

human T cell activation. *Science* **345,** 1254665 (2014).

35.   Petukhova, L. *et al.* Genome-wide association study in alopecia areata implicates

both innate and adaptive immunity. *Nature* **466,** 113–117 (2010).

36.   Raj, T. *et al.* Polarization of the effects of autoimmune and neurodegenerative risk

alleles in leukocytes. *Science* **344,** 519–523 (2014).

37.   Schiering, C. *et al.* The alarmin IL-33 promotes regulatory T-cell function in the

intestine. *Nature* **513,** 564–568 (2014).

38.   Huang, H. *et al.* Fine-mapping inflammatory bowel disease loci to single-variant

resolution. *Nature* (2017). doi:10.1038/nature22969

39.   Onengut-Gumuscu, S. *et al.* Fine mapping of type 1 diabetes susceptibility loci and

evidence for colocalization of causal variants with lymphoid gene enhancers. *Nat. Genet.*

47, 381–386 (2015).

40.   O'Shea, J. J., Lahesmaa, R., Vahedi, G., Laurence, A. & Kanno, Y. Genomic views of

STAT function in CD4+ T helper cell differentiation. *Nat. Rev. Immunol.* **11,** 239–250

(2011).

41.   Miller, C. L. *et al.* Integrative functional genomics identifies regulatory mechanisms

at coronary artery disease loci. *Nat. Commun.* **7,** 12092 (2016).

42.   Nurnberg, S. T. *et al.* Coronary Artery Disease Associated Transcription Factor

TCF21 Regulates Smooth Muscle Precursor Cells That Contribute to the Fibrous Cap.

PLoS Genet. **11,** e1005155 (2015).

43.   McPherson, R. *et al.* A common allele on chromosome 9 associated with coronary

heart disease. *Science* **316,** 1488–1491 (2007).

44.   Helgadottir, A. *et al.* A common variant on chromosome 9p21 affects the risk of

myocardial infarction. *Science* **316,** 1491–1493 (2007).

45.   Clarke, R. *et al.* Genetic variants associated with Lp(a) lipoprotein level and coronary

disease. *N. Engl. J. Med.* **361,** 2518–2528 (2009).

46.   CARDIoGRAMplusC4D Consortium *et al.* Large-scale association analysis identifies

new risk loci for coronary artery disease. *Nat. Genet.* **45,** 25–33 (2013).

47.   Leung, D. *et al.* Integrative analysis of haplotype-resolved epigenomes across human

tissues. *Nature* **518,** 350–354 (2015).

48.   Dixon, J. R. *et al.* Chromatin architecture reorganization during stem cell

differentiation. *Nature* **518,** 331–336 (2015).

49.   Franzén, O. *et al.* Cardiometabolic risk loci share downstream cis- and trans-gene

regulation across tissues and diseases. *Science* **353,** 827–830 (2016).

609   **FIGURE LEGENDS**
610
Fig. 1 HiChIP identifies high-resolution chromosome conformation in primary
human T cells. (A) Primary T cell H3K27ac HiChIP experimental outline. **(B)** (Left)
FACS strategy for Naïve, T_H17, and T_{reg} cells from total peripheral blood CD4$^+$ T cells.
Number represents percent of total CD4$^+$ T cells within that gate. (Right) Knight-Ruiz
(KR) matrix-balanced interaction maps for Naïve, T_H17, and T_{reg} cells at 500 kb, 25 kb,
and 5 kb resolution, and raw interaction maps at 1 kb resolution, centered on the *KLF2*,
RBPJ, and *LRRC32* loci. **(C)** HiChIP 1D and 3D signal enrichment at the *RORC* locus in
T_H17 over Naïve T cells.
619
Fig. 2 Validation of regulatory elements identified by H3K27ac HiChIP with
CRISPR interference and activation. (A) Interaction profile of the *MYC* promoter in
K562 H3K27ac HiChIP at 10 kb resolution. K562 H3K27ac ChIP-Seq is from ENCODE.
CRISPRi-validated regulatory regions in K562 cells are indicated[5]. **(B)** Interaction profile
of the *GATA1* promoter in K562 H3K27ac HiChIP at 1 kb resolution. CRISPRi-validated
regulatory regions in K562 cells are indicated[5]. **(C)** Correlation of *MYC* K562 H3K27ac
HiChIP signal with max CRISPRi score within the HiChIP 10 kb window. (**D**) Interaction
profiles of the *MYC* promoter in GM12878 and My-La H3K27ac HiChIP at 10 kb
resolution. T cell H3K27ac ChIP-seq and ATAC-seq are from Naïve T cells. **(E)** (Top)
CRISPRi validation in GM12878 cells of GM12878 and My-La-biased *MYC* enhancers.
(Bottom) *MYC* RNA levels by qRT-PCR and cell growth rates in CRISPRi GM12878
cells targeted to cell type-biased enhancers, the *MYC* promoter, and a non-targeting
negative control (n = 3). * corresponds to p-value < 0.05, ** to p-value < 0.01, *** to p-
value < 0.001, **** to p-value < 0.0001, and n.s. to not significant. The box extends from
the 25th to 75th percentiles with a line representing the median, and the whiskers go the
minimum and maximum values. **(F)** Interaction profile of the *CD69* promoter in Jurkat
H3K27ac HiChIP at 5 kb and 1 kb resolutions. The 1 kb profile is focused on the window
of the CRISPRa tiling screen. CRISPRa-validated regulatory regions in Jurkat cells are
indicated[26]. **(G)** (Top) CRISPRa validation in Jurkat cells of *CD69* distal enhancers.
(Bottom) *CD69* RNA and protein levels in CRISPRa Jurkat cells targeted to distal
enhancers, the *CD69* promoter, the *KLRF2* promoter as a locus negative control, and a
non-targeting negative control (n = 2).
642
Fig. 3 Dynamic 3D enhancer landscapes in T cell differentiation. (A)
Conformational features observed by H3K27ac HiChIP. **(B)** HiChIP EIS in 913
differential interactions identified in T cell subtypes. Interactions are clustered by cell-
type specificity. **(C)** Cell-type specific motif identification from ATAC-seq peaks in biased
EIS anchors. **(D)** EIS bias quartiles for Naïve to T_H17 and Naïve to T_{reg} differentiation,
with corresponding differential RNA gene expression rankings. **(E)** Proportion of ATAC-
seq peaks within HiChIP differential interaction anchors that are cell-type specific (log$_2$
fold change > 1) or shared across all three subtypes. **(F)** Interaction profile of the
BACH2 promoter at 5 kb resolution, demonstrating shared accessibility signal at Naïve-

biased EIS. **(G)** Global correlation of EIS and ATAC-seq fold-change in different T cell
subset pairwise comparisons.

Fig. 4 HiChIP identifies cell type-specificity and target genes of autoimmune
diseases variants. (A) Generation of a loop set between all autoimmune SNPs and
gene promoters within a 1 Mb region. **(B)** H3K27ac ChIP and HiChIP signal bias in T
cell subtypes for SNP-TSS pairs. For each bin, PICS SNPs are tagged by H3K27ac only
in the concordant cell type for the bias tested. SNPs are grossly divided into
associations with autoimmune diseases or control, non-immune traits. **(C)** EIS Bias of
SNP-TSS loops (with nearest gene annotated) in T_H17 and T_{reg} subsets versus Naïve,
and T_H17 versus T_{reg}. **(D)** Number of HiChIP gene targets versus nearest gene
predictions for all looping nongenic autoimmune SNPs as well as SNPs for specific
diseases. **(E)** Global validation of HiChIP SNP gene targets. Synthetic SNP-TSS pairs
were generated from each $CD4^+$ eQTL SNP to its associated gene and compared to
both a distance-matched shuffled SNP-TSS pair and a liver eQTL SNP-TSS pair. **(F)**
HiChIP target gene RNA levels by qRT-PCR in CRISPRi My-La cells targeted to SNP-
containing enhancers of interest, as well as positive control sgRNAs to the HiChIP
target promoters and a non-targeting negative control (n = 3). * corresponds to p-value
< 0.05, ** to p-value < 0.01, *** to p-value < 0.001, **** to p-value < 0.0001, and n.s. to
not significant. The box extends from the 25^{th} to 75^{th} percentiles with a line representing
the median, and the whiskers go the minimum and maximum values.

Fig. 5 Fine-mapping of GWAS variants using H3K27ac HiChIP. (A) Global validation
of HiChIP signal at putatively causal SNPs versus corresponding SNPs in LD ($r^2 \geq 0.8$)
for T_H17 cells. SNP-TSS pairs were generated from published fine-mapped datasets,
compared to a distance-matched SNP-TSS pair set in the same LD block. **(B)**
Interaction profile of the *PTGER4* promoter, and a 1 kb resolution visualization of the
SNP-containing enhancer of interest. LD SNPs ($r^2 \geq 0.8$) correspond to GRASP SNPs
(genome-wide significance p-value < 10^{-8}). The highlighted SNP was identified in both
the high confidence PICS and GRASP datasets. **(C)** Interaction profiles of the *STAT1*
and *STAT4* promoters, with 1 kb resolution visualizations of the SNP-containing
enhancers of interest. Highlighted are 1D signal contributions at the *STAT1* and *STAT4*
promoters. Highlighted SNPs are PICS closest to focal EIS to *STAT4*.

Fig. 6 HiChIP identifies allelic bias to target genes for cardiovascular disease risk
variants. (A) Interaction profile of the *TCF21* gene promoter for H3K27ac HiChIP of
HCASMC and Naïve T cells. **(B)** EIS bias between HCASMC and Naïve T cells in a
union set of CARDIoGRAMplusC4D CAD and PICS autoimmune SNP-TSS loops. **(C)**
Q-Q plot of allelic EIS imbalance in high confidence loops. Allelic mapping biased loops
were identified through simulation and removed prior to EIS analysis. **(D)** EIS bias
between CAD risk variants and their alternative alleles to eQTL associated target genes.
(E) Allele-specific HiChIP interaction profiles at the 9p21.3 and *SMAD3* loci at 10 kb
resolution in order to examine the functional consequence of a risk variant compared to
its alternative allele.

METHODS
698
699 **Human Subjects**
700 This study was approved by the Stanford University Administrative Panels on Human
701 Subjects in Medical Research, and written informed consent was obtained from all
702 participants.
703
Cell Culture and Primary T cell Isolation
Mouse ES cells (v6.5, Novus Biologicals: NBP1-41162) were cultured in Knockout
DMEM (Gibco) + 15% FBS and leukemia inhibitory factor (LIF, Millipore) to 80%
confluence. GM12878 (Coriell), Jurkat, and My-La (CD4$^+$) cells (ATCC) were grown in
RPMI 1640 (Gibco) with 15% FBS to a concentration of 500,000 to 1 million cells per
mL. Normal donor human peripheral blood cells were obtained fresh from AllCells. CD4$^+$
T cells were enriched from peripheral blood using the RosetteSep Human CD4$^+$ T Cell
Enrichment Cocktail (StemCell Technology). For CD4$^+$ T helper cell subtypes, Naïve T
cells were sorted as CD4$^+$CD25$^-$CD45RA$^+$, T$_H$17 cells were sorted as CD4$^+$CD25$^-$
CD45RA$^-$CCR6$^+$CXCR5$^-$, and T$_{reg}$ cells were sorted as CD4$^+$CD25$^+$CD127^{lo}. Antibodies
used for FACS included: PerCP/Cy5.5 anti-CD45RA (Biolegend 304122), Brilliant Violet
510 anti-CD127 (Biolegend 351331), APC/Cy7 anti-CD4 (Biolegend 344616), PE anti-
CCR6 (Biolegend 353410), FITC anti-CD25 (Biolegend 302603), Brilliant Violet 421
anti-CXCR3 (Biolegend 353715), and BB515 anti-CXCR5 (BD Biosciences 564625).
For HiChIP experiments, 500,000 - 1 million cells were sorted into RPMI + 10% FCS.
For ATAC-seq experiments, 55,000 cells were sorted into RPMI + 10% FCS. Post-sort
purities of > 95% were confirmed by flow cytometry for each sample.
721
722 Primary human coronary artery smooth muscle cell (HCASMC) line derived from a
723 normal human donor heart was purchased from Cell Applications, Inc. (350-05A) and
724 cultured in smooth muscle growth medium (Lonza, CC-3182) supplemented with hEGF,
725 insulin, hFGF-b, and 5% FBS. Cells were grown according to Lonza's instructions.
726
Cell Fixation
Detached cell lines or sorted CD4$^+$ T cells were pelleted and resuspended in fresh 1%
formaldehyde (Thermo Fisher) at a volume of 1 mL formaldehyde for 1 million cells.
Cells were incubated at room temperature for 10 min with rotation. Glycine was added
at a final concentration of 125mM to quench the formaldehyde, and cells were incubated
at room temperature for 5 min with rotation. Finally, cells were pelleted and washed with
PBS, pelleted again, and stored at –80 °C or immediately taken into the HiChIP
protocol.
735
HiChIP
The HiChIP protocol was performed as previously described, using either H3K27ac
antibody (Abcam ab4729) or CTCF (Abcam ab70303)[16] with the following modifications.
For primary T cells, we performed HiChIP on as many cells as we could obtain from a
blood donation - approximately 500,000 - 1 million cells per T cell subtype per replicate.

We performed two minutes of sonication, no Protein A bead preclearing, used 4 µg of
H3K27ac antibody (Abcam ab4729), and captured the chromatin-antibody complex with
34 µL of Protein A beads (Thermo Fisher). Qubit quantification post ChIP ranged from 5
– 25 ng depending on the cell type and amount of starting material. The amount of Tn5
used and PCR cycles performed were based on the post ChIP Qubit amounts, as
previously described[16].

25m cell line libraries were generated as previously described[16]. For low cell number
mouse embryonic stem cell samples, we performed two minutes of sonication and no
Protein A bead preclearing. Either 4 µg or 2µg of H3K27ac antibody (Abcam ab4729)
was used for ChIP in 500k or 100k/50k cells, respectively, and the chromatin-antibody
complex was captured with 34 (500k cells) or 20 µL (100k/50k cells) of Protein A beads.
Post-ChIP Qubit quantification for the 25m cell samples was approximately 1.5 µg. For
lower cell numbers, quantification was 30, 10, and 5 ng for 500k, 100k, and 50k cells,
respectively. The amount of Tn5 used and PCR cycles performed were based on the
post ChIP Qubit amounts, as previously described.

HiChIP samples were size selected by PAGE purification (300-700 bp) for effective
paired-end tag mapping, and therefore were removed of all primer contamination which
would contribute to recently reported "index switching" on the Illumina HiSeq 4000
sequencer[50]. All libraries were sequenced on the Illumina HiSeq 4000 to an average
depth of 500-600M total reads.

HiChIP Data Processing
HiChIP paired-end reads were aligned to hg19 or mm9 genomes using the HiC-Pro
pipeline[51]. Default settings were used to remove duplicate reads, assign reads to MboI
restriction fragments, filter for valid interactions, and generate binned interaction
matrices. HiC-Pro filtered reads were then processed into a .hic file using the
hicpro2juicebox function. The Juicer pipeline HiCCUPS tool was used to identify high
confidence loops[4] using the same parameters as for the GM12878 *in situ* Hi-C map:
hiccups -m 500 -r 5000,10000 -f 0.1,0.1 -p 4,2 -i 7,5 -d 20000,20000 .hic_input
HiCCUPS_output. For T cell Juicer loops, performing the default Juicer calls resulted in
a high rate of false positives upon visual inspection of the interaction matrix. We
therefore called loops with the same HiCCUPS parameters in two biological replicates
for each T cell subtype and then filtered loops for those that were reproducibly called in
both replicates. In addition, we removed all loops greater than 1 Mb.

1D signal enrichment and peak calling were generated from the HiC-Pro filtered
contacts file. Intrachromosomal contacts were filtered and both anchors were extended
by 75 base pairs. The combined bed file containing both anchors was then used to
generate bigwigs for visualization in the WashU Epigenome Browser or call peaks using
MACS2.

Allele-specific HiChIP data processing was achieved using HiC-Pro's allele-specific
analysis features[51]. First, HCASMC phasing data[41] was used to mask the hg19 genome
and make indexes. HiC-Pro settings were similar as to described above, with the
exception that reads were aligned to the masked genome, and then assigned to a
specific allele based on phasing data.
789
790   **Interaction Matrices and Virtual 4C Visualization**
791   HiChIP interaction maps were generated with Juicebox using Knight-Ruiz (KR) matrix
792   balancing and visualized using Juicebox software at 500 kb, 25 kb, 10 kb, and 5 kb
793   resolutions as indicated in each analysis[4]. For 1 kb profiles, raw matrix counts were
794   visualized in Java TreeView.
795
Virtual 4C plots were generated from dumped matrices generated with Juicebox. The
Juicebox tools dump command was used to extract the chromosome of interest from the
.hic file. The interaction profile of a specific 5 kb or 10 kb bin containing the anchor was
then plotted in R. Replicate reproducibility was visualized with the mean profile shown
as a line and the shading surrounding the mean representing the standard deviation
between replicates. For the HCASMC data we observed low read coverage for allele-
specific v4Cs at loci of interest. This is due to the density of SNPs for this genotype, and
a low number of reads containing a phased SNP. We therefore could not observe
interaction profiles when visualizing separate replicates with the standard deviation. We
therefore utilized pseudoreplicates for the HCASMC v4C visualizations[52].
806
807   High confidence Juicer loop calls were loaded into the WashU Epigenome Browser
808   along with corresponding ATAC-seq profiles and publically available H3K27ac ChIP-seq
809   data from the Roadmap Epigenome Project. Browser shots from WashU track sessions
810   were then included in virtual 4C and interaction map anecdotes.
811
Reproducibility Scatter Plots and Correlations
Biological and technical reproducibility comparisons between HiChIP experiments were
generated by counting reads supporting a set of Juicer loop calls. For biological
replicate comparisons, the loop set called from the merged replicates was used. For the
comparison in mES cells between low cell number and 25 million cells, the union loop
set between the two maps was used. The Pearson correlation between replicates or
experiments was calculated from depth-normalized reads using the cor() function in R.
Scatter plots were plotted using log-transformed raw reads supporting each loop.
820
821   **Promoter/Enhancer Annotations of HiChIP Loops**
822   Promoters were defined as 1 kb regions centered at the TSS, and enhancers were
823   identified as chromHMM enhancers not overlapping with promoters in any cell type. We
824   annotated loop anchors as 'other' if the anchor did not contain a promoter or enhancer
825   as defined above.
826
Differential Analysis of HiCCUPS Loop Calls

Juicer loop calls from the three T cell subtypes were initially combined into a union set
of T cell loops. Loop signal was then obtained for the biological replicates of each T cell
subtype. Vanilla coverage square root (VCsqrt) normalized signal for the interaction
matrix of each biological replicate using the Juicebox tools dump command. Normalized
signal was then assigned to the union loop set in each replicate.

VCsqrt signal per sample was quantile-quantile normalized under the assumption that
overall signal was identically distributed across all samples. Following normalization,
samples for Naïve, T_H17 and T_{reg} cells shared Pearson correlations of 0.938, 0.942 and
0.934, respectively. PCA was performed using the prcomp function in R, which
demonstrated that the first PC, which exhibited near-identical loadings across the six
samples, explained 93% of the variance across the six samples. This was taken to
represent the shared signal across cell types. PCs 2-4 explained 2.2%, 2.0% and 1.4%,
respectively.

To study cell type-specific looping, the residual signal per loop was taken after
projecting the loop onto the unit vector along the diagonal (equal signal per cell type).
Cell type-specific and differential looping analysis were performed with the top and
bottom 5% of the distributions of either residual signal or differences between cell type
residual signals. Hierarchical clustering was performed using the union of all differential
loops in these extremes and using 1 minus the Pearson correlation as the distance
metric. QQ plots were generated by permuting residuals from the same cell type or
individual and summing them together and using this distribution to calculate p values
for the observed sums.

In parallel, differential loops were called using edgeR for both the mES and T cell
datasets. Again, biological replicate loop signal was obtained across a union set of
Juicer loops. We then used edgeR to identify loops with significant changes in signal
among pair-wise comparisons (FDR < 0.1, $\log_2FC > 1$). Importantly, inspection of
differential loops identified from the two methods revealed high concordance.

Gene density was calculated from Ensembl gene annotations. GC content was
calculated per 10 kb bin using the bedtools nuc function and aggregated as needed.
Notably, Spearman correlation between gene density of an entire chromosome and the
number of differential loop anchors (rho = 0.914) was much higher than the correlations
between the variance in cell type signal per loop anchor and number of genes per 10 kb
window (rho = 0.322) and between differential loop anchors and gene density per 100
kb section (rho = 0.083). Correlations between GC content and number of differential
loops were similar at both the chromosome (rho = 0.729) and 100 kb bin (rho = 0.148)
levels, but while local GC content is likely to confound relative abundance, it is unclear
how chromosome-wide GC content could have the same effect.

For mES analysis of H3K27ac- and cohesin-mediated HiChIP, we performed edgeR to
obtain the biased loops for each factor. To determine functional bias of the top loops,

overlap was determined between edgeR differential loop anchors and relevant ChIP-seq
peaks. Smc1a ChIP-seq peaks were obtained from a published dataset[53]. CTCF, RNA
Polymerase II, and H3K27ac ChIP-seq peaks were obtained from the mouse ENCODE
repository[54].
876
RNA Expression Analysis
Previously generated RNA-seq data[55] from Naïve, T_H17 and T_{reg} cells was downloaded
as fastqs from ArrayExpress. Illumina adaptors were trimmed using CutAdapt and
Ensembl's cDNA transcripts were quantified using kallisto. Sleuth was used to identify
transcripts that were differentially expressed across cell types with FDR controlled at
5%. The mean TPM was calculated per cell type, and TSS differential looping quantiles
at genes with with nonzero expression were correlated with differential expression
quantiles of the same genes. Only 10 kb segments of the genome that contained a
single annotated gene were considered to avoid errors in attribution of looping signal
per 10 kb bin. For genes with multiple annotated TSSs the 10 kb bin corresponding to
the median TSS was used. Significance was assessed by the cor.test function in R.
888
889   **Comparing HiChIP to Hi-C, ATAC-seq, ChIP-seq, and Capture-C datasets**
890   T cell subset ChIP-seq data was obtained from the WashU Roadmap repository. A
891   union set of peaks were called using MACS2 and peaks were quantified using Bedtools
892   intersect. Normalization for ChIP was performed using quantile normalization using
893   "preprocessCore" package in R.
894
895   Differential EIS was determined using TMM normalization in the "edgeR" package in R.
896   The significantly differential EIS (log_2 fold-change > 1 and FDR < 10%) were determined
897   for each pairwise comparison. For each differential EIS, the maximum ATAC/ChIP
898   signal peak was assessed in each 10 kb anchor (to bias against low signal peaks) and
899   then the maximum log_2 fold-change was compared to the differential EIS.
900
Differential 1D HiChIP was determined using TMM normalization in the "edgeR"
package in R. The significantly differential 1D HiChIP in EIS 10 kb anchors (log_2 fold-
change > 1 and FDR < 10%) were determined for each pairwise comparison. For each
differential 1D HiChIP anchor, the maximum ATAC/ChIP signal peak was assessed (to
bias against low signal peaks) and then the maximum log_2 fold-change was compared
to the differential 1D HiChIP.
907
908   To determine if reference cell line Hi-C data with primary T cell H3K27ac ChIP-seq
909   could recapitulate EIS in primary T cells, GM Hi-C loop anchors were binned with
910   increasing T cell subset ChIP signal as well as GM H3K27ac ChIP-seq signal (Encode).
911   Loop overlap was then determined for the different H3K27ac ChIP signal bins with
912   HiChIP loops.
913
To compare directly with CD4$^+$ Capture-C data, CHiCAGO loops were called in Naïve,
T_H17, and T_{reg} datasets. CHiCAGO calls were then combined into a union set, and loop

signal was obtained for the biological replicates of each T cell subtype using the
Juicebox tools dump command. We then used edgeR to identify loops with significant
changes in signal among pair-wise comparisons (FDR < 0.05). Total and differential
loops were then overlapped with CD4+ Capture-C CHiCAGO data.
920
Calculation of Disease-specific GWAS SNP Enrichment in Loop Anchors
We categorized GWAS SNPs into sets relevant to diseases of the immune system and,
separately, diseases with no known immune component[25]. For each disease in the
immune or non-immune set, we determined the proportion of all GWAS SNPs
associated with that disease which overlap the positions of loop anchors based on a
union set of loops identified in Naïve, T_H17, and T_{reg} cells. The ratio of the proportion of
immune and non-immune overlaps relative to a shuffled control was reported as the
enrichment of immune GWAS SNPs.
929
Distance-matched eQTL SNP-TSS Comparisons
We obtained three groups of eQTL SNP-TSS pairs within 1 Mb distance for HiChIP EIS
comparisons. The treatment group contains CD4 eQTL-TSS targets. We have two
distance-matched groups as control. The first control group contains the CD4 eQTL
SNP-random TSS pairs such that the distance between eQTL SNP and random TSS
differs by at most 5 kb with the treatment group. The second control group contains liver
eQTL SNP-TSS targets that are also distance matched with the treatment group. The
random eQTL SNP-TSS pairs were generated by individual chromosome, such that
number of control pairs and treatment pairs are the same for every chromosome. In
total, there were 158,482 distance-matched eQTL-TSS pairs. We compared the 5 kb
resolution EIS among the three eQTL SNP-TSS groups for all three T cell subtypes.
Results show that in all the cases, the EIS between CD4 eQTL-TSS targets were
significantly higher than the two control groups (p-value < 10^{-16}).
943
Distance-matched Fine-mapped SNP-TSS Comparisons
We obtained a list of putatively causal SNPs from the PICS SNP list[25] (PICS probability
> 0.5), as well as fine-mapped SNPs associated with inflammatory bowel disease
(IBD)[38] or Type 1 Diabetes (T1D)[39]. Next, we obtained all SNPs in LD with each
putatively causal SNP using European linkage disequilibrium blocks determined by all
SNPs with an $r^2 \geq 0.8$ with the SNP being considered. For the fine-mapped (T1D/IBD)
sets, using SNPs in LD with highly significant GWAS SNPs may mean that there are
several SNPs of equal or greater significance in the control set, but we still expect an
enrichment relative to the LD block.
953
We collected all the synthetic pairs between the putatively causal immune-disease
related SNPs (IBD, T1D and PICS) and nearby genes within 300 kb distance. To
perform the distance-matched EIS comparisons, for each fine-mapped SNP category,
we selected the SNP-TSS control pairs which satisfy two constraints: (1) the selected
control SNP is positioned at least 5 kb away from the fine-mapped SNP in the same
linkage disequilibrium block; (2) the distance of SNP-TSS control pair differs with fine-

mapped SNP and the target gene at most 5 kb.
961
SNP-TSS Loop Analyses
We obtained 7,747 PICS SNPs that are associated with autoimmune disease or non-
immune traits[25]. 4,331 (55.9%) were associated with autoimmune disease, and 3,416
(44.1%) were associated with non-immune traits. In addition we obtained a set of SNPs
associated with six overlapping autoimmune diseases using the GRASP catalog
(genome-wide significance p-value $< 10^{-8}$).
968
969   We constructed a synthetic loop set for immune and non-immune SNPs and any TSS
970   within 1 Mb of each SNP. We then assigned VCsqrt signal in each biological replicate of
971   the three T cell subtypes to the synthetic loop set, as described above.
972
VCsqrt signal per sample was quantile-quantile normalized as above. In this analysis,
we did not restrict to HiCCUPS-identified loops but instead examined all possible
interactions between a SNP and TSS within 1 Mb. Many of these interactions do not
exist, and therefore had little or no matrix-balanced signal supporting them. While we
removed all SNP TSS pairs below an average of 1 normalized read per sample from
subsequent analyses, in general these false interactions contributed little to the overall
differential signal for a trait.
980
981   H3K27ac data was downloaded from the WashU Roadmap repository. PICS SNPs were
982   taken from Farh, et al[25]. Rather than require strict membership within H3K27ac peaks,
983   PICS SNPs were labeled as active if they lied within 8 kb of a peak, raising the number
984   of nominally functional SNPs from ~700 to ~3200 per cell type, out of 7735 total
985   candidate SNPs.
986
Differential looping across cell types was assessed by one-sided t-test per trait and
activity partition if there were at least eight PICS SNP-TSS pairs in the partition. T_H17
bias was defined as T_H17 total loop signal minus Naïve total loop signal; T_{reg} bias was
defined as T_{reg} total loop signal minus Naïve total loop signal; and Naïve bias was
defined as Naïve total loop signal minus one-half times T_H17 and T_{reg} total loop signal.
For the main text figure, Naïve bias was assessed only using SNPs that were active in
Naïve cells, T_H17 bias from SNPs active in T_H17 and not Naïve, and T_{reg} bias from
SNPs active in T_{reg} and not Naïve. p-values were corrected for multiple hypothesis
testing by the Holm method using the p.adjust function in R. Bias assessed from SNPs
with the opposing cell type specificities (e.g. Naïve bias using SNPs active in T_H17 and
T_{reg} but not Naïve) yielded no significant hits after correction.
998
999   **SNP and TSS Connectivity Analysis**
1000   Among all immune and non-immune SNPs, 2,562 (33.1%) were located within a 25 kb
1001   region centered around TSS, 2,618 (33.8%) were located in a gene body, and the
1002   remaining 2,567 (33.1%) were located in intergenic regions.
1003

To capture the confident SNP-TSS connections for SNP-TSS pair identification, we
overlapped the anchors of significant loops that were identified by Fit-Hi-C[56] with the
SNP/TSS locations. We identified 14,738 SNP-TSS pairs that were supported by Fit-Hi-
C loops, and there were 3,046 unique SNPS connected with at least one gene. Among
those 3,046 SNPs, there were 2,181 (71.6%) and 865 (28.4%) SNPs annotated with
immune and non-immune SNPs, respectively. As expected, the immune disease SNPs
are more likely to connect with genes (Fisher's exact test, p-value = 4.8 x 10^{-85}).
1011
Phasing of HCASMC Samples
We used BEAGLE 4.1 to impute and phase recalibrated variants using 1000 Genome
phase 3 version 5a as a reference panel. The Beagle phasing algorithm was set on the
following criteria. At each iteration that the algorithm performs, phased input data are
used to build a local model of a haplotype-cluster. After the local haplotype-cluster
model is created, for each individual phased haplotypes are sampled using the induced
diploid HMM conditional on the individual's genotypes. The sampled haplotypes are
then used as the input to feed in the next iteration of the algorithm. In the final iteration,
instead of sampling haplotypes, BEAGLE uses the Viterbi algorithm to
select the conditional on the diploid HMM and the individual's genotype data and to
obtain the haplotypes for each individual that possess the greatest probability, and
these most-likely haplotypes are the final output of the BEAGLE phasing algorithm.
1024
Allelic Mapping Bias Simulation
We constructed a personal genome by editing the reference genome (hg19) according
to SNP information. SNPs labeled as "1|0" or "1|1" in the vcf file were replaced with the
alternative allele for genome 1. SNPs labeled as "0|1" or "1|1" in the vcf file were
replaced with the alternative allele for genome 2. Loop anchors were extended 100 bp
in both directions and sequences were extracted using samtools for each
genome. ~20X reads were simulated for each region, using wgsim with parameters " -e
0.01 -d 100 -s 20 -1 75 -2 75 -S -1 -h -R 0.1". Simulated reads were mapped to the
"N" masked genome. Mapping parameters were the same used by HiC-Pro ("--very-
sensitive -L 30 --score-min L, -0.6, -0.2 --end-to-end --reorder --phred33-quals). Allelic
specific reads were separated according to the SNP information and then counted for
each loop anchor using bedtools.
1037
Correction for False Positives
Despite not being originally implemented in enrichment interaction datasets, we have
previously demonstrated that KR and VC matrix balancing corrects for false positives
caused by high 1D fragment visibility[16]. Loop calls are either from HiCCUPS or Fit Hi-C,
which are KR and VC matrix balanced, respectively. For differential analysis of
HiCCUPS (Figure 3) and SNP-TSS Fit Hi-C (Figure 4c) loop calls, we are restricted
within a set of loops that were obtained from matrix balancing. Therefore, while
differential loops can be driven by both changes in looping strength and/or 1D ChIP
signal, the final interactions being observed are loops. Additionally, we performed
differential analysis of HiCCUPS loop calls using two separate methods – one using

VCsqrt normalized reads and another with non-normalized reads (edgeR) and found
high agreement. The SNP-TSS synthetic loop analysis in Figure 4b was not restricted to
loop calls, however was performed with VCsqrt matrix balanced reads to avoid false
positives. Those SNP-TSS synthetic loops were then subset by overlap with Fit Hi-C for
further differential analyses in Figure 4c. Finally, virtual 4C analysis was performed on
non-normalized reads to highlight EIS contributions of both 1D and 3D signal changes,
however HiCCUPS loop calls are included in the relevant anecdotes.
1055
ATAC-seq
Cells were isolated and subjected to ATAC-seq as previously described[16]. Briefly,
55,000 cells were pelleted, resuspended in 50 µL lysis buffer (10mM Tris-HCl, pH 7.4,
3mM MgCl2, 10mM NaCl, 0.1% NP-40 (Igepal CA-630), and immediately centrifuged at
500 rcf for 10 min at 4^0C. The nuclei pellets were resuspended in 50 µL transposition
buffer (25 µl 2X TD buffer, 22.5 µL dH$_2$0, 2.5 µL Illumina Tn5 transposase), and
incubated at 37^0C for 30 min. Transposed DNA was purified with MinElute PCR
Purification Kit (Qiagen), and eluted in 10 µL EB buffer.
1064
ATAC-seq Data Processing
Adapter sequence trimming using SeqPurge and mapping to hg19 using Bowtie2 were
performed. These reads were then filtered for mitochondrial reads, low quality, and PCR
duplicates. The filtered reads for each sample were merged and peak calling was
performed by MACS2. Each individual sample reads in peaks were quantified using
Bedtools intersect with the MACS2 narrow peaks. Peak counts were then combined into
a matrix NxM where N represents called peaks and M represents the samples and each
value Di,j represents the peak intensity for the respective peak i in sample j. This matrix
was then normalized using the "CQN" package in R to minimize bias in GC content and
length.
1075
CRISPRi Validation of HiChIP Targets
For virus production, 5×10^6 of HEK293T cells were plated per 10 cm plate. The
following day, plasmid encoding lentivirus was co-transfected with pMD2.G and psPAX2
into the cells using Lipofectamine 3000 (Thermo Fisher, L3000) according to the
manufacturer's instructions. Supernatant containing viral particles was collected 48
hours after transfection and filtered. For lentivirus encoding individual sgRNAs, virus
was concentrated 10-fold using Lenti-X concentrator (Clontech, 631232) and stored at -
80°C.
1084
In order to generate a My-La cell line expressing CRISPRi, 2×10^6 of My-La cells were
plated per T75 flask. A dCas9-BFP-KRAB-2A-Blast construct was generated by
inserting a 2A-Blast cassette into dCas9-BFP-KRAB (Addgene 46911). 24 hours after
plating, lentivirus harboring the dCas9-KRAB construct was added with polybrene (4 µg
/ mL). Media was changed 24 hours after infection, and then again 48 hours after
infection with Blasticidin (Thermo Fisher, A1113903) at 4 µg / mL. Blasticidin resistant
cells were selected for eight days with changing media every other day.

1092
Three different U6 were used for transcription of three different sgRNAs targeting the
candidate enhancers, as previously described[57]. For *MYC* locus CRISPRi experiments,
each enhancer was targeted by one guide, and therefore all *MYC* GM or My-La
enhancers together were targeted in one experiment. For the PICS SNP CRISPRi
experiments, three guides were targeted to a single SNP-containing enhancer. One of
three sgRNAs was cloned into a lentiviral vector with a human (pMJ117, Addgene
85997), mouse (pMJ179, Addgene 85996) or bovine (pMJ114, Addgene 85995) U6
promoter. These U6-sgRNA constructs were then combined into lentivirus with a
Puromycin-2A-mCherry vector, which was modified from Addgene 46914. My-La-
CRISPRi cells were infected with lentivirus harboring 3 sgRNAs and selected by
Puromycin (Thermo Fisher, A11138) at a final concentration of 1 µg / mL. Previously
reported sgRNAs targeting VPS54 or SEC24C were used for validating CRISPRi
functionality in My-La cell line[58].
1106
For readout of CRISPRi validation, we performed qRT-PCR and cell growth assays on
three biological and two technical replicates. For qRT-PCR, RNA was Trizol extracted
(Thermo Fisher, 15596026) and purified using the Zymo RNA Clean and Concentrator
kit (Zymo Research, R1016). qRT-PCR was performed with Brilliant qRT-PCR
mastermix (Agilent, 600825). Ct values were measured by using Lightcycler 480
(Roche) and relative expression level was calculated by ddCt method compared to a
GAPDH control. Primer sequences are listed in Supplementary Table 6. For cell growth,
we used the CellTiter-Glo kit (Promega, G7572) according to the manufacturers
instructions. Statistics for both RNA and cell growth changes were calculated using a
Student's t test against the non-targeting control.
1117
1118   **CRISPRa Validation of HiChIP Targets**
1119   Jurkat cells were transduced with a lentiviral dCas9-VP64-2A-GFP expression vector
1120   (Addgene 61422). Single GFP+ cells were sorted by FACS into the wells of a 96-well
1121   plate, and a clone with bright uniform GFP expression were selected for use in future
1122   experiments.
1123
sgRNAs were cloned in arrayed format for CD69 HiChIP peaks falling outside the range
of the tiling CRISPRa screen[26]. sgRNAs were chosen based on high predicted on-target
activity[59] and low predicted off-target activity[60]. sgRNAs were cloned into the lentiviral
expression vector "pCRISPRia-v2" (Addgene 84832) as described in Horlbeck et al[61].
Lentivirus was produced by transfecting HEK293T with standard packaging vectors
using *Trans*IT-LTI Transfection Reagent (Mirus, MIR 2306). Media was changed 24
hours post-transfection. Viral supernatant was harvested at 48 and 72 hr following
transfection and immediately used for infection of Jurkat-dCas9-VP64 cells.
1132
Jurkat-dCas9-VP64 cells were infected with lentiviral sgRNAs by resuspending cells in a
1:1 mix of fresh media and lentiviral supernatant at a final concentration of 0.25×10^6
cells/mL with 5 µg / mL polybrene. Cells were spinfected for 1 hour at 1000 rcf, 32 °C.

The next day, half of the media was removed and replaced with fresh lentiviral
supernatant, and the spinfection was repeated. The next day, the cells were
resuspended in fresh media with 1.5 µg / mL puromycin and cultured for 2 days to
remove uninfected cells. For readout of CRISPRa validation, we performed qRT-PCR
and FACS on two biological and two technical replicates. RNA extraction and qRT-PCR
was performed as described above. Expression of CD69 on infected cells (GFP+BFP+)
was analyzed by flow cytometry with an Attune NxT flow cytometer (Life Technologies).
Statistics for both RNA and protein level changes were calculated with a one-way
ANOVA followed by a Dunnet's multiple comparisons test against the non-targeting
control.

SUPPLEMENTARY LEGENDS

Supplementary Figure 1. H3K27ac HiChIP enriches E-P-associated chromatin contacts. (A) Schematic of chromatin contacts captured in H3K27ac HiChIP. **(B)** Loop call overlap for cohesin HiChIP and H3K27ac HiChIP in mES cells. **(C)** Contact distance distribution for loops that are biased in cohesin versus H3K27ac HiChIP. **(D)** Proportion of cohesin and H3K27ac biased HiChIP loops that have cohesin, CTCF, H3K27ac, and RNA polymerase II binding in at least one loop anchor. **(E)** Virtual 4C interaction profile of an H3K27ac-biased loop focused at the *MALAT1* promoter. **(F)** Virtual 4C interaction profile of a cohesin-biased loop with associated low transcriptional activity.

Supplementary Figure 2. H3K27ac HiChIP achieves high chromatin loop signal-over-background at low cell inputs. (A) KR balanced interaction matrices focused around the *Etv5* locus in mES cells with decreasing cellular starting material. **(B)** Read support reproducibility of loops in H3K27ac HiChIP libraries from 25 million cells compared to HiChIP in lower cell input libraries. **(C)** Aggregate peak analysis of loops in mES H3K27ac HiChIP libraries.

Supplementary Figure 3. H3K27ac HiChIP generates reproducible chromatin loop signals at low cell inputs. (A) Comparison of KR balanced interaction maps in H3K27ac HiChIP biological replicates. **(B)** Read support reproducibility of loops between H3K27ac HiChIP biological replicates. **(C)** HiCCUPS loop call overlap between H3K27ac HiChIP libraries from 25 million and 50 thousand mES cells. **(D)** Preseq library complexity estimation of H3K27ac HiChIP libraries from 25 million and 50 thousand mES cells.

Supplementary Figure 4. H3K27ac HiChIP biological replicates from primary sorted T cells are highly reproducible. (A) FACS strategy for Naïve, T_H17, and T_{reg} cells starting from human peripheral blood. Post-sort validation was used to ensure purity of T cell subtypes. Number represents percent of cells within that gate. **(B)** KR balanced interaction map of T cell subtype biological replicates. **(C)** Read support reproducibility of loops between H3K27ac HiChIP biological replicates in Naïve, T_H17, and T_{reg} cells. **(D)** Aggregate peak analysis of loops in Naïve, T_H17, and T_{reg} H3K27ac HiChIP libraries.

Supplementary Figure 5. Validation of HiChIP-identified *CD69* distal enhancers with CRISPR activation. CRISPRa validation in Jurkat cells of *CD69* distal enhancers. *CD69* protein levels are shown for individual sgRNAs tiling H3K27ac HiChIP-identified distal *CD69* enhancers relative to the *KLRF2* promoter as a locus negative control and a non-targeting negative control.

Supplementary Figure 6. Global enhancer connectome characterization in T cell differentiation. (A) ChromHMM classification of union T cell loop anchors. **(B)** Contact distance distribution of union T cell loops. **(C)** Agreement in residual signal observed

between sample signals per loop after removing imputed shared signal, calibrated by a
null distribution of random pairings of loops. QQ plot shows modest enrichment above
random pairings. PCA on residual signal clusters samples first by Naive vs Memory cell
types (PC1), and then by donor identity (PC2, 3). **(D)** Overlap of differential interactions
between Naïve, T_H17, and T_{reg} subtypes. Biased interactions were obtained by
performing pairwise comparisons between T cell types and analyzing the top 5%
enriched and top 5% depleted EIS in each T cell subtype. **(E)** ChromHMM annotation of
total loops, differential loops, and shared loops in all three T cell subtypes. O
corresponds to other loop anchors not classified as enhancer or promoter. **(F)** Number
of connections for different classes of loop elements. **(G)** Quantification of promoters
skipped before an enhancer reaches its loop target.
1235
1236   **Supplementary Figure 7. Positioning of differential HiChIP contacts in gene**
1237   **dense chromosomes. (A)** Distribution of T cell subtype differential HiChIP contacts
1238   across different chromosomes compared to the distribution of all loops. **(B)** Correlation
1239   of differential loop density with gene density and GC content.
1240
1241   **Supplementary Figure 8. Characterization of conformational landscapes**
1242   **surrounding key T cell regulatory factors. (A – C)** Virtual 4C interaction profiles
1243   anchored at the promoters of canonical Naïve, T_H17, and T_{reg} regulatory factors.
1244
1245   **Supplementary Figure 9. Chromosome conformation dynamics of canonical T**
1246   **cell regulatory factors. (A – C)** Delta interaction maps focused around known Naïve,
1247 T_H17, and T_{reg} regulatory factors.
1248
Supplementary Figure 10. Contribution of H3K27ac ChIP and chromosome
conformation to HiChIP EIS. (A) (Left) Proportion of H3K27ac ChIP peaks within EIS
differential loop anchors that are cell-type specific ($\log_2$ fold change > 1) or shared
across all three subtypes. (Right) Global correlation of EIS and H3K27ac ChIP fold-
change in different T cell subset pairwise comparisons. **(B)** Same as (A), but using
HiChIP 1D differential signal at EIS biased loop anchors. **(C)** Overlap of H3K27ac
HiChIP and bins of Hi-C loops with increasing T cell subset and GM H3K27ac ChIP-seq
signal. **(D)** Overlap of CD4[+] Capture Hi-C[14] with total and differential T cell subset
HiChIP loops. **(E)** T_{reg}-specific loops at the *LRRC32* promoter not observed in other
H3K27ac HiChIP T cell subsets nor in CD4[+] Capture Hi-C data[14].
1259
Supplementary Figure 11. Enrichment of autoimmune SNPs in T cell HiChIP loop
anchors. (A) Enrichment of specific PICS autoimmune disease and non-immune SNPs
in anchors of loops called by Juicer and Fit-Hi-C compared to a background shuffled
loop set. **(B)** Enrichment of all PICS autoimmune disease and non-immune SNPs in T
cell subset biased loop anchors and all anchors compared to a background shuffled
loop set.
1266

Supplementary Figure 12. T cell subtype HiChIP specificity of autoimmune SNPs.
(A) H3K27ac HiChIP signal bias in T cell subsets for PICS SNP-TSS pairs grouped by
each SNP's presence in cell type-specific or shared H3K27ac ChIP peaks up to 8 kb
away. **(B)** H3K27ac HiChIP signal bias in T cell subsets for PICS SNP-TSS pairs
grouped by each SNP's presence in cell type-specific or shared H3K27ac ChIP peaks
up to 2.5 kb away. **(C)** H3K27ac HiChIP signal bias in GM, K562, and My-La cell lines
for PICS SNP-TSS pairs grouped by each SNP's presence in T cell subset-specific or
shared H3K27ac ChIP peaks up to 8 kb away. **(D)** H3K27ac HiChIP signal bias in T cell
subsets for GRASP SNP-TSS pairs grouped by each SNP's presence in cell type-
specific or shared H3K27ac ChIP peaks up to 2.5 kb away. **(E)** Average number of
HiChIP gene targets for non-genic autoimmune disease and non-immune SNPs. **(F)**
Quantification of promoters skipped before a SNP reaches its gene target. **(G)**
Quantification of SNP HiChIP gene targets in autoimmune disease.
1280
1281 **Supplementary Figure 13. Validation of HiChIP signal at SNP-eQTL contacts. (A)**
1282 Validation of HiChIP signal at SNP-TSS pairs using interaction profiles of eQTL SNPs to
1283 ensure they contact their associated target gene promoter. **(B)** Interaction profiles of
1284 CRISPRi-validated loci in My-La.
1285
Supplementary Figure 14. H3K27ac HiChIP fine-mapping of GWAS variants in
haplotype blocks. (A) Global validation of HiChIP signal at putatively causal SNPs
versus corresponding SNPs in LD ($r^2 \geq 0.8$) for Naïve and T_{reg} cell subtypes. SNP-TSS
pairs were generated from published fine-mapped datasets, compared to a distance-
matched SNP-TSS pair set in the same LD block. **(B)** Interaction profile of the *SATB1*
promoter, and a 1 kb resolution visualization of the SNP-containing enhancer of interest.
Changes in bias between 5 kb and 1 kb resolution reflects EIS focused 1 kb around the
SATB1 TSS and specific SNPs within the enhancer. LD SNPs ($r^2 \geq 0.8$) correspond to
GRASP SNPs (genome-wide significance p-value $< 10^{-8}$). The highlighted SNP is a
PICS closest to focal EIS to *SATB1.*
1296
1297 **Supplementary Figure 15. Chromatin interaction landscape of the 9p21.3**
1298 **cardiovascular disease risk locus.** HCASMC v4C interaction profiles focused around
1299 the promoters of *CDKN2A*, *CDKN2B*, and *ANRIL* within the 9p21.3 locus.
1300 **Supplementary Table 1. HiChIP data processing metrics.** HiC-Pro mapping, Hi-C
1301 filtering, duplicate removal, and interaction length statistics for all HiChIP libraries.
1302
1303 **Supplementary Table 2. HiCCUPS high confidence loop calls.** High confidence loop
1304 calls for all HiChIP libraries.
1305
1306 **Supplementary Table 3. HiCCUPS differential EIS in T cell subtypes by edgeR.**
1307 High confidence loop calls in T cell subsets with edgeR significance and fold-change for
1308 pair-wise comparisons.
1309

1310 **Supplementary Table 4. HiCCUPS differential EIS in T cell subtypes.** Top 5% of
1311 EIS ranked by cell-type bias for each pair-wise comparison between different T cell
1312 subtypes.
1313
1314 **Supplementary Table 5. HiChIP gene targets of autoimmune and CAD SNPs.** SNP
1315 type (within gene, at TSS, or intergenic), associated disease, and corresponding loop
1316 annotated with the HiChIP gene target. Included are edgeR significance and fold-
1317 change for cell-type comparisons of interest.
1318
1319 **Supplementary Table 6. sgRNA and primer oligonucleotide sequences.** sgRNA
1320 and primer sequences used throughout the study.
1321
1322 **DATA AVAILABILITY STATEMENT**
1323 Raw and processed data available at NCBI Gene Expression Omnibus, accession
1324 number GSE101498.
1325
1326 T cell ATAC and HiChIP datasets can be visualized in the WashU Epigenome Browser
1327 with the following link:
1328 http://epigenomegateway.wustl.edu/browser/?genome=hg19&session=YAIzYBfrl9&stat
1329 usId=1698051079
1330
1331 **COMPETING FINANCIAL INTEREST**
1332 The authors declare no competing financial interests.

METHODS-ONLY REFERENCES

50.   Index Switching Causes "Spreading-Of-Signal" Among Multiplexed Samples In Illumina HiSeq 4000 DNA Sequencing | bioRxiv. Available at: http://biorxiv.org/content/early/2017/04/09/125724. (Accessed: 19th May 2017)

51.   Servant, N. *et al.* HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biol.* **16,** 259 (2015).

52.   Gjoneska, E. *et al.* Conserved epigenomic signals in mice and humans reveal immune basis of Alzheimer's disease. *Nature* **518,** 365–369 (2015).

53.   Kagey, M. H. *et al.* Mediator and cohesin connect gene expression and chromatin architecture. *Nature* **467,** 430–435 (2010).

54.   Yue, F. *et al.* A comparative encyclopedia of DNA elements in the mouse genome. *Nature* **515,** 355–364 (2014).

55.   Bonnal, R. J. P. *et al.* De novo transcriptome profiling of highly purified human lymphocytes primary cells. *Sci. Data* **2,** 150051 (2015).

56.   Ay, F., Bailey, T. L. & Noble, W. S. Statistical confidence estimation for Hi-C data reveals regulatory chromatin contacts. *Genome Res.* gr.160374.113 (2014). doi:10.1101/gr.160374.113

57.   Adamson, B. *et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. *Cell* **167,** 1867–1882.e21 (2016).

58.   Gilbert, L. A. *et al.* Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell* **159,** 647–661 (2014).

59.   Doench, J. G. *et al.* Optimized sgRNA design to maximize activity and minimize off-

target effects of CRISPR-Cas9. *Nat. Biotechnol.* **34,** 184–191 (2016).

60.   Hsu, P. D. *et al.* DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat.*

Biotechnol. **31,** 827–832 (2013).

61.   Horlbeck, M. A. *et al.* Nucleosomes impede Cas9 access to DNA in vivo and in vitro.

eLife **5,** (2016).

1383