1  **Enhancer connectome in primary human cells reveals target genes of disease-**
2  **associated DNA elements**
3

Maxwell R. Mumbach[1,2,12], Ansuman T. Satpathy[1,3,12], Evan A. Boyle[1,2,12], Chao Dai[1,12],
Benjamin G. Gowen[4,5], Seung Woo Cho[1], Michelle L. Nguyen[6], Adam J. Rubin[7], Jeffrey
M. Granja[1,2], Katelynn R. Kazane[4,5], Yuning Wei[1], Trieu Nguyen[8], Peyton G.
Greenside[2], M. Ryan Corces[1], Josh Tycko[2], Dimitre R. Simeonov[6,9], Nabeela Suliman[2],
Rui Li[1], Jin Xu[1], Ryan A. Flynn[1], Anshul Kundaje[2], Paul A. Khavari[7], Alexander
Marson[5,6,8], Jacob E. Corn[4,5], Thomas Quertermous[9], William J. Greenleaf[1,2,8,11,12],
Howard Y. Chang[1,7,12]
11
12 **AFFILIATIONS**
13
[1] Center for Personal Dynamic Regulomes, Stanford University School of Medicine,
Stanford, CA 94305, USA
[2] Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305,
USA
[3] Department of Pathology, Stanford University School of Medicine, Stanford, CA 94305,
USA
[4] Department of Molecular and Cell Biology, University of California, Berkeley, CA
94720, USA
[5] Innovative Genomics Institute, University of California, Berkeley, CA, 94720, USA
[6] Department of Microbiology and Immunology, University of California, San Francisco,
CA 94143, USA
[7] Program in Epithelial Biology, Stanford University School of Medicine, Stanford, CA,
94305, USA.
[8] Chan Zuckerberg Biohub, San Francisco, CA 94158, USA
[9] Department of Medicine, Division of Cardiovascular Medicine, Stanford University
School of Medicine, Stanford, California 94305, USA.
[10] Biomedical Sciences Graduate Program, University of California, San Francisco, CA
94143, USA
[11] Department of Applied Physics, Stanford University, Stanford, CA, 94305, USA
[12] These authors contributed equally
[13] These authors jointly directed this work
35
36 *To whom correspondence should be addressed. E-mail: howchang@stanford.edu and
37 wjg@stanford.edu

45  **ABSTRACT**
46
The challenge of linking intergenic mutations to target genes has limited molecular
understanding of diverse human diseases. Here, we show H3K27ac HiChIP generates
high-resolution contact maps of active enhancers and target genes in rare primary
human T cell subtypes and coronary artery smooth muscle cells. Differentiation of naïve
T cells to either T helper 17 cells or regulatory T cells create subtype-specific enhancer-
promoter interactions, specifically at regions of shared DNA accessibility. These data
provide a principled means of assigning molecular functions to autoimmune and
cardiovascular disease risk variants, linking hundreds of noncoding variants to putative
gene targets. Target genes identified with HiChIP are further supported by CRISPR
interference and activation at linked enhancers, by the presence of expression
quantitative trait loci, and by allele-specific enhancer loops in patient-derived primary
cells. The majority of disease-associated enhancers contact genes beyond the nearest
gene in the linear genome, leading to a four-fold increase of potential target genes for
autoimmune and cardiovascular diseases.

Gene expression programs are intimately linked to the hierarchical organization
of the genome. In mammalian cells, each chromosome is organized into hundreds of
megabase-sized topologically associated domains (TADs), which are conserved from
early stem cells to differentiated cell types[1]. Within this invariant TAD scaffold, cell type-
specific enhancer-promoter (E-P) interactions establish regulatory gene expression
programs[2]. Standard methods require tens of millions of cells to obtain high-resolution
interaction maps and confidently assign E-P contacts[3–5]. Thus, the principles that
govern E-P conformation in disease-relevant patient samples are incompletely
understood. This gap in understanding is particularly problematic for interpreting the
molecular functions of inherited risk factors for common human diseases, which reside
in intergenic enhancers or other non-coding DNA features in up to 90% of cases[6–9].
Such disease-relevant enhancers may not influence the expression of the nearest gene
(often reported as the default target in the literature), and instead act in a cell-type
specific manner on distant target genes residing up to hundreds of kilobases (kb)
away[2,10–14]. Recently, systematic perturbations of regulatory elements in select gene loci
have shown that effects of individual regulatory elements on gene activity can be
predicted from the combination of (i) enhancer activity [marked by histone H3 lysine 27
acetylation (H3K27ac) level] and (ii) enhancer-target looping[5,15]. Here we leverage this
insight to capture the combination of these two types of information genome-wide in a
single assay, mapping the enhancer connectome in disease-relevant primary human
cells.
110
111  **RESULTS**
112
113  **H3K27ac HiChIP identifies functional enhancer interactions in low cell number**
114  **samples**
115
We recently developed HiChIP, a method for sensitive and efficient analysis of
protein-centric chromosome conformation[16]. Cohesin HiChIP in GM12878 cells
identified similar numbers of loops as *in situ* Hi-C (~10,000) with high correlation (R =
0.83), demonstrating that HiChIP captures loops with high sensitivity and specificity.
Here, we evaluated the enhancer and promoter-associated mark H3K27ac[17–19] as a
candidate factor to selectively interrogate E-P interactions genome-wide. We performed
H3K27ac HiChIP in mouse embryonic stem (mES) cells to compare to cohesin HiChIP
in the same cell type (**Supplementary Fig. 1a, Supplementary Table 1**)[16]. 3,552 of
4,191 H3K27ac HiChIP loops in mES cells were also identified by cohesin HiChIP. The
H3K27ac-biased loops ($\log_2$ fold-change > 1 and FDR < 10%) spanned shorter
distances than cohesin-biased loops, and were enriched for H3K27ac ChIP-seq peaks
(78.9%; **Supplementary Fig. 1b-f, Supplementary Table 2**). Moreover, systematic
titration of input material showed H3K27ac HiChIP retained high signal fidelity and
reproducibility from 25 million to 50,000 cells as input material (loop signal correlation =
0.918 and loop overlap of 43%, comparable to in situ Hi-C replicate loop overlap of
53%; **Supplementary Figs. 2 and 3**). Therefore, H3K27ac HiChIP identifies high-

confidence chromatin loops focused around enhancer interactions from limited cell
numbers.
In order to capture (i) conformational change during T cell differentiation and (ii)
cell type-specific chromatin contacts of autoimmune risk variants in protective and
pathogenic T cell types, we performed H3K27ac HiChIP on primary human Naïve T
cells ($CD4^+CD45RA^+CD25^-CD127^{hi}$), regulatory T cells ($T_{reg}$; $CD4^+CD25^+CD127^{low}$) and
T helper 17 cells ($T_H17$; $CD4^+CD45RA^-CD25^-CD127^{hi}CCR6^+CXCR5^-$) directly isolated
from donors (**Fig. 1a,b and Supplementary Fig. 4a**)[20,21]. $T_H17$ cells were sorted to
include autoimmune disease-relevant pathogenic $T_H17$ cells and to exclude follicular
helper T cells with a distinct surface phenotype and immune function (**Supplementary**
**Fig. 4a**)[22–24]. Peripheral blood $CD4^+$ T cells were isolated from three healthy subjects,
isolated by FACS, and subjected to H3K27ac HiChIP. HiChIP libraries from each subset
were high quality; greater than 40% of the total reads sequenced represented unique
paired-end tags (PETs) (**Supplementary Fig. 4b-d and Supplementary Table 1**).
Furthermore, libraries exhibited high 1D signal enrichment at enhancers and promoters,
and globally recapitulated publically available H3K27ac ChIP-seq datasets (74.7%
overlap of ChIP-seq and 1D HiChIP peaks; **Fig. 1c**)[25]. Inspection of the interaction
matrix at progressively higher resolution revealed chromatin compartments, TADs, and
focal loops, as previously reported in high-resolution Hi-C and HiChIP analyses from cell
lines (**Fig. 1b**)[4,16]. Importantly, H3K27ac HiChIP maps were capable of identifying focal
interactions at 1 kb resolution, which is comparable to *in situ* Hi-C maps generated from
100-fold more cells and sequenced to 13-fold greater depth[4] (**Fig. 1b**).
Previous saturation perturbation screens demonstrated that functional enhancers
can be identified by integrating H3K27ac ChIP-seq signal with chromosome
conformation contact strength (Hi-C)[5]. Since H3K27ac HiChIP combines these two
components into one assay, we reasoned that HiChIP signal, which we term Enhancer
Interaction Signal (EIS), should identify functional regulatory elements. To validate this
prediction, we first generated H3K27ac HiChIP maps in a chronic myelogenous
leukemia cell line (K562) as a direct comparison to published high-resolution CRISPR
interference (CRISPRi) screens[5]. We then examined the 3D enhancer landscape of the
*MYC* and *GATA1* loci using virtual 4C (v4C) analysis, where a specific genomic position
is set as an anchor viewpoint, and all interactions occurring with that anchor are
visualized in 2D[16]. v4C analysis of the *MYC* promoter demonstrated that EIS in K562
cells captured all functional enhancers identified in the CRISPRi screen (**Fig. 2a**).
Analysis of the *GATA1* locus demonstrated a similar agreement between both methods
(**Fig. 2b**). Quantitatively, EIS in K562 cells was significantly correlated with CRISPRi
score in the same cell type, whereas EIS in GM12878 (GM; B cell lymphoblast) cells
was not correlated with K562 CRISPRi (Spearman's rho = 0.332 and 0.145; p-value =
$9.25 \times 10^{-5}$ and 0.1246; **Fig. 2c**).
We found the enhancer landscapes of the *MYC* promoter to be highly cell-type
specific. v4C analysis of the *MYC* promoter in GM and My-La ($CD4^+$ T cell leukemia)
cells showed dramatically different regulatory interactions with the promoter compared
to K562 cells (**Fig. 2d**). To validate EIS specificity, we performed CRISPRi experiments
in GM cells using sgRNAs targeting enhancers identified in either GM or My-La HiChIP

maps as well as a positive control sgRNA targeting the *MYC* promoter and a negative
control sgRNA targeting lambda phage sequence (**Fig. 2e**). As expected, we found that
CRISPRi of GM, but not My-La, enhancers impacted *MYC* expression and cell growth in
GM cells (**Fig. 2e**).
Finally, we focused on the *CD69* locus, where a high-resolution CRISPRa screen
identified three enhancers located approximately 4 kb, 13 kb, and 37 kb upstream of the
transcription start site[26]. These sites were also precisely identified by Naïve T cell
H3K27ac HiChIP. Moreover, HiChIP identified four additional distal enhancers that were
outside the region spanned by the sgRNA tiling array (**Fig. 2f and Supplementary Fig.
5**). To functionally validate these novel enhancers, we performed CRISPRa experiments
in Jurkat cells with sgRNAs targeting these enhancers, the *CD69* promoter, the *KLRF2*
promoter as a locus negative control, and a non-human genome-targeting negative
control. We observed a significant increase in *CD69* RNA and protein levels in the four
HiChIP enhancers compared to negative controls (**Fig. 2g and Supplementary Fig. 5**).
Interestingly, two of the four identified novel enhancers were within promoter regions of
distant genes. These findings are in line with previous reports that identified widespread
distal gene regulatory functions of promoters genome-wide[27,28]. Altogether, these results
suggest that H3K27ac HiChIP EIS identifies functional regulatory elements, and that
enhancers that regulate a gene of interest can differ significantly between cell-types.
195
196  **Landscape of enhancer interactions in primary human T cell subsets**
197
We examined global features of the enhancer connectome associated with
cellular differentiation from Naïve T cells to either $T_H17$ cells or $T_{reg}$ cells. We identified a
total of 10,706 high confidence loops in the union set of the three cell types
(**Supplementary Table 2**). Analysis of loop read support between biological replicates
demonstrated high reproducibility (**Supplementary Fig. 4c**), and ~91% of loop anchors
were associated with either a promoter or enhancer[29], as expected, with a median
distance of 130 kb (**Supplementary Fig. 6a,b**). Importantly, high-resolution E-P
connectivity maps revealed several features that could not be discerned from 1D
epigenomic data (i.e. H3K27ac ChIP-seq or ATAC-seq; **Fig. 3a**). These features
included: (i) 'enhancer skipping': enhancers that lack strong EIS with their closest
neighboring gene, or many neighboring genes, yet have strong EIS with a more distal
target promoter, (ii) higher order structures such as 'enhancer cliques' (related to loop
cliques[30]): multiple regulatory elements that have strong EIS with a single target
promoter, (iii) promoter to promoter interactions[13,31], and (iv) 'enhancer switching':
enhancers that exhibit differential EIS with a target promoter in a cell type-specific
manner (**Fig. 3a**).
We found that EIS contacts were very cell type-specific, and gene regulation in
distinct T cell populations was better understood using EIS signal, compared to 1D
enhancer activity. After quantile-quantile normalization of matrix-balanced contact reads
at high-confidence loops (correcting for false positives caused by 1D fragment visibility;
**Materials and Methods**), we focused on the top and bottom 5% of EIS ranked by cell-
type bias for each pair-wise comparison (**Supplementary Figs. 6c-g and 7,**

**Supplementary Tables 3-4**). We identified 6 clusters of EIS defined by interaction
strength patterns across T cell subtypes: Clusters 1-3 were enriched in Naïve cells,
Cluster 4 was enriched in $T_H17$ and $T_{reg}$ cells, and Clusters 5 and 6 were enriched in
$T_H17$ and $T_{reg}$ cells, respectively (**Fig. 3b**). Cell type-specific enhancer loop anchors
revealed genes encoding canonical T cell subtype TFs and effector molecules (**Fig. 3b,**
**Supplementary Figs. 8 and 9**). Deeper v4C analysis of shared and cell type-specific
loci pinpointed regulatory elements interacting with each gene promoter of interest as
well as local conformational landscape changes (**Supplementary Figs. 8 and 9**). TF
motifs located within cell type-specific loop anchors were enriched for TFs known to
drive T cell subtype differentiation and nominated novel TFs involved in regulation (**Fig.**
**3c**). Furthermore, cell type EIS bias was associated with differential expression of genes
located within corresponding EIS anchors for the same cell type (Spearman's rho =
0.242 and 0.207; p-value = 4 x $10^{-15}$ and 2 x $10^{-11}$; **Fig. 3d**).
Cell type-specific EIS may be driven by cell type-specific enhancer activation
(based on H3K27ac ChIP-seq) or stable enhancer activation with cell type-specific
looping (Hi-C) in a gene specific manner. We first examined H3K27ac ChIP-seq at
differential EIS anchors and found that many biased H3K27ac HiChIP interactions also
exhibited biased ChIP-seq signal, as expected. 58.5% of Naïve-biased loops contain at
least one Naïve-biased ChIP-seq peak ($log_2$ fold change > 1) located on the anchors.
Similarly, 66.7% of $T_H17$-biased and 67.8% of $T_{reg}$-biased interaction anchors were cell
type-specific in 1D (**Supplementary Fig. 10a**). Therefore, while on average ~64% of
the differential EIS corresponded to change in 1D data, ~36% were likely also driven by
change in 3D chromatin loop strength. To further assess the contribution of cell type-
specific 3D signal to EIS, we examined HiChIP 1D signal at differential EIS anchors. We
found that HiChIP 1D signal correlated better with ChIP-seq signal than EIS, with a
higher likelihood of differential ChIP-seq signal overlapping differential HiChIP 1D signal
compared to 3D, suggesting EIS bias is in part driven by 3D changes (**Supplementary**
**Fig. 10b**).
As a complementary analysis, we asked whether the integration of reference cell
line Hi-C data with primary T cell H3K27ac ChIP-seq could recapitulate HiChIP EIS in
primary T cells. We first binned GM Hi-C loops with increasing primary T cell ChIP-seq
signal at loop anchors and then determined the overlap of loops in each bin with loops
derived from H3K27ac HiChIP. As expected, increased ChIP-seq signal at the Hi-C
anchors led to increased overlap with the HiChIP loops. However, the overlap was lower
in all T cell subtypes compared to the same analysis performed using GM HiChIP data.
These observations demonstrate that cell-type specific 3D interactions can impact EIS
independent of differences in 1D ChIP-seq signal (**Supplementary Fig. 10c**). Similarly,
previously generated enhancer-promoter maps obtained from bulk unsorted T cells did
not identify T cell subtype-specific interactions obtained using H3K27ac HiChIP. To
assess the unique information obtained through cell type-specific interaction maps, we
compared promoter Capture Hi-C maps in bulk $CD4^+$ T cells to H3K27ac HiChIP maps
in Naïve, $T_H17$, and $T_{reg}$ cells[14]. We examined the overlap of promoter Capture Hi-C
interactions with HiChIP interactions at increasing thresholds of cell type-specificity.
Strikingly, the most cell type-specific loops in $T_H17$ and $T_{reg}$ (16-fold enriched)

demonstrated a low discovery rate in promoter Capture Hi-C T cells (11.83% in 415
loops and 13.83% in 373 loops, respectively; **Supplementary Fig. 10d**). Many of these
subset-specific interactions included genomic loci encoding functionally important
effector genes, such as *LRRC32*. The *LRRC32* locus contains $T_{reg}$-specific loops that
are neither visualized in HiChIP maps from Naïve or $T_H17$ cells nor in bulk $CD4^+$
promoter Capture Hi-C maps (**Supplementary Fig. 10e**). Since primary human $T_H17$
and $T_{reg}$ cells are present in human blood with low frequency, it would also be
challenging to generate subset-specific promoter Capture Hi-C maps with published
promoter Capture Hi-C protocols. In summary, EIS is derived from a combination of 1D
ChIP-seq and 3D interaction signal and cannot be accurately predicted from 3D maps in
reference cell lines or unsorted primary cell datasets.
Cell type-specific EIS can occur at sites of shared chromatin accessibility. Paired
chromatin accessibility profiles by Assay of Transposase-Accessible Chromatin by
sequencing (ATAC-seq)[32] from each T cell subset revealed most cell type-specific loop
anchors had equivalent chromatin accessibility across all three cell types (**Fig. 3e-g**). To
illustrate this finding, we examined the *BACH2* promoter, which exhibits shared
chromatin accessibility at enhancers, but increased EIS in Naïve cells (**Fig. 3e**).
Globally, only 14.2% of Naïve-biased loops contain at least one Naïve-biased ATAC-
seq peak ($log_2$ fold change > 1) located on the anchors. In addition, 27.8% of $T_H17$-
biased and 16.5% of $T_{reg}$-biased interaction anchors were cell type-specific for 1D
accessibility. Furthermore, the majority of cell type-specific TF motifs were observed in
shared ATAC-seq peaks within differential interactions, highlighting that these regions
are functioning in T cell differentiation (**Fig. 3f-g**). Altogether, these results suggest that
in highly related – yet functionally distinct – cell types, a portion of transcriptional control
is achieved through differential chromosome looping, rather than differential chromatin
accessibility. This finding is consistent with previous studies which demonstrated that T
cell subset-specific TFs, such as Foxp3, act predominantly at pre-accessible chromatin
sites to establish subset-specific gene expression[33].
292
293  **T cell subset-specific enhancer interactions link autoimmunity-associated DNA**
294  **elements to target genes**
295
The high specificity of EIS enabled us to identify putative target genes of
autoimmune disease risk loci in functionally relevant T cell subsets. To achieve this, we
used a previously described list of putatively causal variants associated with 21
autoimmune diseases, known as PICS SNPs, which were fine-mapped based on dense
genotyping data[25]. First, we determined that PICS autoimmune SNPs were significantly
enriched in T cell loop anchors, with specific autoimmune diseases showing greater
than 5-fold enrichment compared to a shuffled control loop set (**Supplementary Fig.**
**11**). Next, we constructed a set of all possible connections between autoimmune risk
SNPs and TSS within 1 Mb and measured the EIS for each SNP-TSS pair (**Fig. 4a**). We
aggregated these signals to determine the overall interaction activity in each T cell
subtype in each disease (**Fig. 4b**). Strikingly, we observed high interaction strength
enrichments and cell type specificity in autoimmune disease SNPs, but low enrichment

and cell specificity in non-immune traits (**Fig. 4b**). To further visualize HiChIP bias in
shared or differential enhancers, we analyzed SNP-TSS interactions grouped by their
presence near H3K27ac ChIP-seq peaks (**Supplementary Fig. 12a,b**). Interestingly,
we observed a large number of active SNP-TSS pairs that were present in regulatory
regions that were shared between T effector cell types ($T_{reg}$ and $T_H17$), while relatively
less EIS signal was observed in SNPs located in cell-type specific enhancers,
supporting the concept that many autoimmune disease variants impact common T cell
effector/activation pathways[25,34]. Notably, SNPs present in enhancers shared across all
three cell types could still be distinguished by HiChIP bias (**Supplementary Fig. 12a,b**).
For example, although we could not detect cell type bias at risk loci for Alopecia Areata
using H3K27ac ChIP-seq (**Supplementary Fig. 12a,b** and ref. 3), H3K27ac HiChIP
identified increased SNP-TSS activity in $T_{reg}$ cells among shared T cell enhancers,
consistent with several studies identifying the crucial role of this cell type in disease
pathogenesis[35]. Importantly, autoimmune signal enrichments were not readily apparent
from 1D H3K27ac ChIP-seq peaks, aggregated ChIP-seq signal within the TAD
containing the SNP, nor cell line H3K27ac HiChIP datasets (**Fig. 4b and**
**Supplementary Fig. 12c**). Therefore, examining 3D disease variant interactions may
capture cell type biases more robustly than 1D epigenomic data. Finally, to validate our
findings with an orthogonal dataset, we performed SNP-TSS EIS analysis on an
overlapping set of autoimmune disease-associated SNPs obtained from the NHLBI
GRASP catalog and observed similar enrichments of specific T cell subsets
(**Supplementary Fig. 12d**).
We leveraged HiChIP to identify potential gene targets of intergenic SNPs, which
have classically been paired to the nearest neighboring gene. We overlapped the SNP-
TSS pairs with loops to call a discrete set of target pairs. We then performed differential
analysis on the SNP-TSS loops to ascertain bias for specific T cell subsets (**Fig. 4c**).
Overall, 3.7% and 9.4% of SNP-TSS pairs were cell type-biased in a comparison of
Naïve and Memory T cells, and 2.2% and 6% were cell type-biased in a comparison of
$T_{reg}$ cells and $T_H17$ cells, respectively. Examples of biased SNP-TSS pairs included
*FOXO1* in Naïve T cells (rs9603754), *BATF* (rs2300604) in Memory T cells, *CTLA4*
(rs10186048) in $T_{reg}$ cells, and *IL2* (rs7664452) in $T_H17$ cells (**Fig. 4c**). Next, we sought
to characterize the connectivity landscape of the SNP-TSS loops. We identified an
average of 1.75 gene targets per autoimmune SNP (ranging from 0 to over 10 target
genes), while non-immune traits did not demonstrate an increase in targets (0.33 genes
per SNP; **Supplementary Fig. 12e**). We examined the SNP-nearest gene associations
and compared these to the HiChIP gene target list. For 684 autoimmune intergenic
SNPs, we identified a total of 2,597 HiChIP target genes, representing a four-fold
increase in target genes for known disease SNPs (**Fig. 4d**). Only 367 (~14%) of all
targets were the nearest gene to the SNP, while approximately ~86% of SNPs skipped
at least one gene to reach a predicted target TSS (**Supplementary Fig. 12e**).
Furthermore, approximately ~45% of SNP to HiChIP target interactions had increased
signal compared to the same SNP to nearest gene, despite distance biases.
350

**Disease-associated enhancer-target gene interactions validated by eQTL and CRISPRi**

HiChIP enhancer-target gene interactions can be validated using previously identified point mutations that alter expression at distantly located genes in T cells—i.e. expression quantitative trait loci (eQTL)[36]. For example, the celiac disease-associated SNP rs2058660 impacts the expression of the inflammatory cytokine receptor genes *IL18RAP*, *IL18R1*, *IL1RL1*, and *IL1RL2*, which are known regulators of intestinal T cell differentiation and response[37]. HiChIP EIS revealed contacts between rs2058660 and each of these predicted gene promoters (**Supplementary Fig. 13a**). Similarly, the Crohn's disease risk variant rs6890268 and the multiple sclerosis (MS) risk variant rs12946510 impact the expression of *PTGER4* and *IKZF3*, respectively, and H3K27ac HiChIP also demonstrated clear contacts between these SNPs and their predicted promoter (**Supplementary Fig. 13a**). Globally, HiChIP contact signal was increased in eQTLs in T cells compared to a distance-matched background loop set (p-value < 2.2 x $10^{-16}$; **Fig. 4e)** or to eQTLs identified in an unrelated cell type (liver; p-value < 2.2 x $10^{-16}$). The overlap of HiChIP and eQTL loci provides support for chromosome interactions as a physical basis for distal eQTLs[10–12] and further validates the HiChIP approach to assign enhancer-target gene relationships.

We next sought to directly validate HiChIP SNP-gene targets using CRISPRi in My-La cells. First, we focused on three loci of interest in primary T cells and then confirmed that the SNP-TSS loops were also present in My-La cells (**Fig. 4f and Supplementary Fig. 13b**). We then targeted sgRNAs to these SNP-containing enhancers, as well as positive control sgRNAs to the HiChIP target gene promoters and a negative control sgRNA targeting lambda phage sequence. As expected, we observed a significant reduction of RNA levels in the HiChIP target genes upon CRISPRi of its SNP-containing enhancer (**Fig. 4f**).

**Fine-mapping disease-associated DNA variants with high-resolution H3K27ac HiChIP**

Since SNP-TSS HiChIP signal is capable of identifying target genes of candidate SNPs, we asked whether TSS-SNP HiChIP signals could also be used to nominate functional causal variants within haplotype blocks in a reciprocal manner. We first performed a proof-of-principle analysis using fine-mapped SNPs associated with inflammatory bowel disease (IBD)[38] or Type 1 Diabetes (T1D)[39] as well as high confidence PICS SNPs and examined EIS from putatively causal SNPs to all gene promoters within 300 kb. EIS from putatively causal SNPs to gene promoters was significantly higher than EIS from a distance-matched set of SNPs within the same LD block to gene promoters (p-value = 2.4 x $10^{-15}$, 8.7 x $10^{-8}$, 3.9 x $10^{-3}$ for IBD fine-mapped SNPs, T1D fine-mapped SNPs, and high confidence PICS, respectively; **Fig. 5a** and **Supplementary Fig. 14a**). Next, we assessed the fine-mapping ability of HiChIP EIS at individual loci of interest. First, we focused on IBD- and MS-associated SNPs neighboring the *PTGER4* and *SATB1* loci and performed v4C analysis anchored at the

gene promoters. We calculated EIS signal at 1 kb resolution and identified specific
regions within the linkage disequilibrium (LD) blocks that contained the highest EIS to
the target promoters, positioning the likely causal SNPs within these regions (**Fig. 5b**
and **Supplementary Fig. 14b**). For example, at the *PTGER4* locus (**Fig. 5b**), the ~160
kb genomic interval spanned by LD SNPs in association with Crohn's disease is refined
to two bins of 3kb and 4kb, which both contain PICS SNPs.
We asked whether complex disease-associated loci containing more than one
gene could be fine-mapped using HiChIP. We focused on two disease-associated
enhancers in between the *STAT1* and *STAT4* gene promoters (**Fig. 5c**). These two
genes encode transcription factors with distinct and canonical roles in immune
regulation. Signal transducer and activator of transcription 1 (STAT1) is critical for type I
IFN and IFNγ signaling, whereas STAT4 is activated downstream of IL-12 signaling to
induce $T_H1$ differentiation and IFNγ expression[40]. We first investigated bias of these
enhancers to *STAT1* and *STAT4* and found that, despite comparable linear distance
and 1D signal at the promoters, the enhancers were significantly biased to interact with
*STAT4*. Next, we fine-mapped the disease associated SNPs within this locus using 1 kb
resolution EIS from the *STAT4* promoter, and narrowed down candidate functional
variants within the two enhancers (**Fig. 5c**). In summary, HiChIP EIS can nominate
functional causal variants within haplotype blocks, and two-way analysis of target gene
identification from an enhancer of interest and high-resolution interaction maps of that
enhancer with its target gene can be used to fine-map disease-associated loci
containing several candidate genes.
417
418   **Allelic target gene bias of cardiovascular disease risk variants in patient smooth**
419   **muscle cells**
420
Finally, we asked whether this approach could be applied broadly to other
categories of human disease, and whether we could directly test SNP-TSS associations
using allele-specific HiChIP. We generated high-resolution E-P maps from primary
human coronary artery smooth muscle cells (HCASMC), which can be used to inform
variants linked to cardiovascular diseases[41]. First, to validate cell type specificity, we
examined the *TCF21* gene promoter, a transcription factor required for the
differentiation of epicardial progenitor cells to HCASMC[42] and observed enrichment in
HCASMC EIS relative to Naïve T cells (**Fig. 6a**). We next examined the 9p21.3 locus,
which harbors risk associations with several cardiovascular disorders, including
coronary artery disease (CAD), myocardial infarction, and stroke[43–45]. We found that the
promoters of all three genes in the locus interact with one another and with CAD variant-
containing enhancers located approximately 100 kb upstream of the *CDKN2B* promoter
(**Supplementary Fig. 15**). We then generated SNP-TSS target lists using CAD SNPs
identified in the CARDIoGRAMplusC4D study[46]. We again performed differential
analysis on the SNP-TSS loops to ascertain bias for HCASMC versus Naïve T cells
(**Fig. 6b**). Overall, 75.1% of biased HCASMC SNP-TSS pairs were CAD SNPs, while
only 5.5% of Naïve T cell biased SNP-TSS pairs were CAD SNP-TSS loops,
demonstrating cell-type specificity of SNP-TSS interactions. Next, we examined the

connectivity of the HCASMC SNP-TSS contacts and identified 1,062 gene targets, of
which only 120 (~11%) mapped to the nearest gene. Furthermore, approximately 89%
skipped at least one gene to reach a predicted target TSS, and 64% of SNPs were
mapped to more than a single gene target.
We took advantage of genome phasing information in HCASMC to measure E-P
interactions at allele-specific CAD SNPs, allowing us to examine the functional
consequence of a risk variant compared to its alternative allele in the same nucleus.
First, 4.2% of high confidence loops in HCASMC with no observed mapping bias in the
anchors exhibited significant allelic bias (FDR < 0.05, **Fig. 6c**), consistent with
frequency of allelic imbalance of RNA expression and prior evidence of allele-specific
regulation of specific E-P interactions[49,50]. We leveraged this global E-P allelic bias to
examine the effect of a risk variant compared to its control alternative allele for a set of
CAD-associated SNP-target gene pairs (**Fig. 6d**)[49]. We found that many risk alleles
disrupt enhancer-target gene interactions, but a subset of pathogenic SNPs increased
enhancer-target gene interaction. At CAD risk variant rs1537373 in the 9p21.3 locus,
the risk allele (T) showed increased EIS to the *CDKN2A* promoter as well as an
additional enhancer within the lncRNA *ANRIL* relative to the reference allele (G) (**Fig.**
**6e**). We further observed increased EIS of the CAD risk variant rs4562997 to an
additional *SMAD3* enhancer 10 kb downstream of the TSS (**Fig. 6e**). The ability to
resolve enhancer connectomes of the risk and reference alleles in the same nucleus
demonstrates that the mutated base in the risk allele suffices to alter enhancer looping
in *cis* in disease-relevant primary cells.
461
462  **DISCUSSION**
463
Here, we developed an approach to define the high-resolution landscape of E-P
regulation in primary human cells. We find that E-P contacts are highly dynamic in
related cell types and often involve genomic elements with shared accessibility.
Accordingly, many complex features of the 3D enhancer connectome cannot simply be
predicted from 1D, which demonstrates that mapping conformation in primary cells can
identify novel regulatory connections underlying gene function in human disease. We
take advantage of this principle to chart the connectivity of autoimmune and
cardiovascular GWAS SNPs and link SNPs to hundreds of potential target genes.
Although non-genic SNPs have previously been paired with their closest neighboring
gene, we find that the majority of these variants can engage in long-distance
interactions, including skipping several promoters to predicted target genes, connecting
to multiple genes, or acting in concert with enhancer cliques to contact a single gene.
Further use of this approach will help to clarify hidden mechanisms of human disease
that are driven by genetic perturbations in non protein-coding DNA elements, which can
now be linked to their cognate gene targets in primary cells.
479
480  **AUTHOR CONTRIBUTIONS**
481

M.R.M, A.T.S., W.J.G., and H.Y.C. conceived the project. M.R.M., A.T.S., J.T., and R.L. performed all genomics assays with help from T.N., M.R.C., N.S., and R.A.F. A.T.S. performed all sorting for experiments. B.G.G., S.W.C., M.R.M., M.L.N., K.R.K., and D.R.S., performed all CRISPR validation experiments. E.A.B., C.D., M.R.M., and J.X. analyzed HiChIP data. J.G., A.T.S., and Y.W. analyzed ATAC-seq data. A.J.R. and P.G.G. analyzed GWAS SNPs in HiChIP data. A.K., P.A.K., A.M., J.E.C., T.Q., W.J.G., and H.Y.C guided experiments and data analysis. M.R.M, A.T.S, E.A.B, C.D., W.J.G, and H.Y.C. wrote the manuscript with input from all authors.

## ACKNOWLEDGEMENTS

We thank members of the Chang and Greenleaf laboratories for helpful discussions and Justin Tumey for artwork. This work was supported by National Institutes of Health (NIH) P50HG007735 (H.Y.C. and W.J.G.) and U19AI057266 (W.J.G), Human Frontier Science Program (W.J.G.), Rita Allen Foundation (W.J.G.), and the Scleroderma Research Foundation (H.Y.C). M.R.M. and E.A.B. acknowledge support from the National Science Foundation Graduate Research Fellowship. A.T.S. is a Cancer Research Institute Irvington Fellow supported by the Cancer Research Institute. B.G.G. was supported by an IGI-AstraZeneca Postdoctoral Fellowship. M.R.C. is supported by a grant from The Leukemia & Lymphoma Society Career Development Program. J.E.C. was supported by the Li Ka Shing Foundation and the Heritage Medical Research Institute. W.J.G. and A.M. are Chan Zuckerberg Biohub investigators. A.M. serves as an advisor to Juno Therapeutics and PACT Therapeutics and the Marson lab has received sponsored research support from Juno Therapeutics and Epinomics. A.M. and J.E.C. are founders of Spotlight Therapeutics. Sequencing was performed by the Stanford Functional Genomics Facility (NIH S10OD018220). We thank Agilent Technologies for generating oligo pools for cloning of the CRISPRa gRNAs. We thank the UC Berkeley High Throughput Screening Facility and Flow Cytometry Facility. H.Y.C. and W.J.G. are founders of Epinomics and members of its scientific advisory board.

**FIGURE LEGENDS**

**Fig. 1 HiChIP identifies high-resolution chromosome conformation in primary**
**human T cells. (A)** Primary T cell H3K27ac HiChIP experimental outline. **(B)** (Left)
FACS strategy for Naïve, $T_H17$, and $T_{reg}$ cells from total peripheral blood CD4$^+$ T cells.
Number represents percent of cells within that gate. (Right) Knight-Ruiz (KR) matrix-
balanced interaction maps for Naïve, $T_H17$, and $T_{reg}$ cells at 500 kb, 25 kb, and 5 kb
resolution, and raw interaction maps at 1 kb resolution, centered on the *KLF2*, *RBPJ*,
and *LRRC32* loci. **(C)** HiChIP 1D and 3D signal enrichment at the *RORC* locus in $T_H17$
over Naïve T cells.

**Fig. 2 Validation of H3K27ac HiChIP regulatory element identification with**
**CRISPR interference and activation. (A)** Interaction profile of the *MYC* promoter in
K562 H3K27ac HiChIP at 10 kb resolution. K562 H3K27ac ChIP-Seq is from ENCODE.
CRISPRi-validated regulatory regions in K562 cells are indicated[5]. **(B)** Interaction profile
of the *GATA1* promoter in K562 H3K27ac HiChIP at 1 kb resolution. CRISPRi-validated
regulatory regions in K562 cells are indicated[5]. **(C)** Correlation of *MYC* K562 H3K27ac
HiChIP signal with max CRISPRi score within the HiChIP 10 kb window. (**D**) Interaction
profiles of the *MYC* promoter in GM12878 and My-La H3K27ac HiChIP at 10 kb
resolution. T cell H3K27ac ChIP-seq and ATAC-seq are from Naïve T cells. **(E)** (Top)
CRISPRi validation in GM12878 cells of GM12878 and My-La-biased *MYC* enhancers.
(Bottom) *MYC* RNA levels by qRT-PCR and cell growth rates in CRISPRi GM12878
cells targeted to cell type-biased enhancers, the *MYC* promoter, and a non-targeting
negative control (n = 3). * corresponds to p-value < 0.05, ** to p-value < 0.01, *** to p-
value < 0.001, **** to p-value < 0.0001, and n.s. to not significant. **(F)** Interaction profile
of the *CD69* promoter in Jurkat H3K27ac HiChIP at 5 kb and 1 kb resolutions. The 1 kb
profile is focused on the window of the CRISPRa tiling screen. CRISPRa-validated
regulatory regions in Jurkat cells are indicated[26]. **(G)** (Top) CRISPRa validation in Jurkat
cells of *CD69* distal enhancers. (Bottom) *CD69* RNA and protein levels in CRISPRa
Jurkat cells targeted to distal enhancers, the *CD69* promoter, the *KLRF2* promoter as a
locus negative control, and a non-targeting negative control (n = 2).

**Fig. 3 Dynamic 3D enhancer landscapes in T cell differentiation. (A)**
Conformational features observed by H3K27ac HiChIP. **(B)** HiChIP EIS in 913
differential interactions identified in T cell subtypes. Interactions are clustered by cell-
type specificity. **(C)** Cell-type specific motif identification from ATAC-seq peaks in biased
EIS anchors. **(D)** EIS bias quartiles for Naïve to $T_H17$ and Naïve to $T_{reg}$ differentiation,
with corresponding differential RNA gene expression rankings. **(E)** Proportion of ATAC-
seq peaks within HiChIP differential interaction anchors that are cell-type specific (log$_2$
fold change > 1) or shared across all three subtypes. **(F)** Interaction profile of the
*BACH2* promoter at 5 kb resolution, demonstrating shared accessibility signal at Naïve-
biased EIS. **(G)** Global correlation of EIS and ATAC-seq fold-change in different T cell
subset pairwise comparisons.

**Fig. 4 HiChIP identifies cell type-specificity in autoimmune diseases and pairs**
**GWAS SNPs with target genes. (A)** Generation of a loop set between all autoimmune
SNPs and gene promoters within a 1 Mb region. **(B)** H3K27ac ChIP and HiChIP signal
bias in T cell subtypes for SNP-TSS pairs. For each bin, PICS SNPs are tagged by
H3K27ac only in the concordant cell type for the bias tested. SNPs are grossly divided
into associations with autoimmune diseases or control, non-immune traits. **(C)** EIS Bias
of SNP-TSS loops (with nearest gene annotated) in $T_H17$ and $T_{reg}$ subsets versus
Naïve, and $T_H17$ versus $T_{reg}$. **(D)** Number of HiChIP gene targets versus nearest gene
predictions for all looping nongenic autoimmune SNPs as well as SNPs for specific
diseases. **(E)** Global validation of HiChIP SNP gene targets. Synthetic SNP-TSS pairs
were generated from each $CD4^+$ eQTL SNP to its associated gene and compared to
both a distance-matched shuffled SNP-TSS pair and a liver eQTL SNP-TSS pair. **(F)**
HiChIP target gene RNA levels by qRT-PCR in CRISPRi My-La cells targeted to SNP-
containing enhancers of interest, as well as positive control sgRNAs to the HiChIP
target promoters and a non-targeting negative control (n = 3). * corresponds to p-value
< 0.05, ** to p-value < 0.01, *** to p-value < 0.001, **** to p-value < 0.0001, and n.s. to
not significant.
572
**Fig. 5 Fine-mapping of GWAS variants in LD using H3K27ac HiChIP. (A)** Global
validation of HiChIP signal at putatively causal SNPs versus corresponding SNPs in LD
($r^2 \geq 0.8$) for $T_H17$ cells. SNP-TSS pairs were generated from published fine-mapped
datasets, compared to a distance-matched SNP-TSS pair set in the same LD block. **(B)**
Interaction profile of the *PTGER4* promoter, and a 1 kb resolution visualization of the
SNP-containing enhancer of interest. LD SNPs ($r^2 \geq 0.8$) correspond to GRASP SNPs
(genome-wide significance p-value $< 10^{-8}$). The highlighted SNP was identified in both
the high confidence PICS and GRASP datasets. **(C)** Interaction profiles of the *STAT1*
and *STAT4* promoters, with 1 kb resolution visualizations of the SNP-containing
enhancers of interest. Highlighted are 1D signal contributions at the *STAT1* and *STAT4*
promoters. Highlighted SNPs are PICS closest to focal EIS to *STAT4*.
584
**Fig. 6 HiChIP identifies allelic bias to target genes for cardiovascular disease risk**
**variants. (A)** Interaction profile of the *TCF21* gene promoter for H3K27ac HiChIP of
HCASMC and Naïve T cells. **(B)** EIS bias between HCASMC and Naïve T cells in a
union set of CARDIoGRAMplusC4D CAD and PICS autoimmune SNP-TSS loops. **(C)**
Q-Q plot of allelic EIS imbalance in high confidence loops. Allelic mapping biased loops
were identified through simulation and removed prior to EIS analysis. **(D)** EIS bias
between CAD risk variants and their alternative alleles to eQTL associated target genes.
**(E)** Allele-specific HiChIP interaction profiles at the 9p21.3 and *SMAD3* loci at 10 kb
resolution in order to examine the functional consequence of a risk variant compared to
its alternative allele.
595
596

## MATERIALS AND METHODS

### Human Subjects

This study was approved by the Stanford University Administrative Panels on Human Subjects in Medical Research, and written informed consent was obtained from all participants.

### Availability of Sequencing Data

Raw and processed data available at NCBI Gene Expression Omnibus, accession number GSE101498.

T cell ATAC and HiChIP datasets can be visualized in the WashU Epigenome Browser with the following link:
http://epigenomegateway.wustl.edu/browser/?genome=hg19&session=YAIzYBfrl9&stat usId=1698051079

### Cell Culture and Primary T cell Isolation

Mouse ES cells (v6.5, Novus Biologicals: NBP1-41162) were cultured in Knockout DMEM (Gibco) + 15% FBS and leukemia inhibitory factor (LIF, Millipore) to 80% confluence. GM12878 (Coriell), Jurkat, and My-La (CD4$^+$) cells (ATCC) were grown in RPMI 1640 (Gibco) with 15% FBS to a concentration of 500,000 to 1 million cells per mL. Normal donor human peripheral blood cells were obtained fresh from AllCells. CD4$^+$ T cells were enriched from peripheral blood using the RosetteSep Human CD4$^+$ T Cell Enrichment Cocktail (StemCell Technology). For CD4$^+$ T helper cell subtypes, Naïve T cells were sorted as CD4$^+$CD25$^-$CD45RA$^+$, T$_H$17 cells were sorted as CD4$^+$CD25$^-$CD45RA$^-$CCR6$^+$CXCR5$^-$, and T$_{reg}$ cells were sorted as CD4$^+$CD25$^+$CD127$^{lo}$. For HiChIP experiments, 500,000 - 1 million cells were sorted into RPMI + 10% FCS. For ATAC-seq experiments, 55,000 cells were sorted into RPMI + 10% FCS. Post-sort purities of > 95% were confirmed by flow cytometry for each sample.

Primary human coronary artery smooth muscle cell (HCASMC) line derived from a normal human donor heart was purchased from Cell Applications, Inc. (350-05A) and cultured in smooth muscle growth medium (Lonza, CC-3182) supplemented with hEGF, insulin, hFGF-b, and 5% FBS. Cells were grown according to Lonza's instructions.

### Cell Fixation

Detached cell lines or sorted CD4$^+$ T cells were pelleted and resuspended in fresh 1% formaldehyde (Thermo Fisher) at a volume of 1 mL formaldehyde for 1 million cells. Cells were incubated at room temperature for 10 min with rotation. Glycine was added at a final concentration of 125mM to quench the formaldehyde, and cells were incubated at room temperature for 5 min with rotation. Finally, cells were pelleted and washed with PBS, pelleted again, and stored at –80 °C or immediately taken into the HiChIP protocol.

**HiChIP**
The HiChIP protocol was performed as previously described[16] with the following
modifications. For primary T cells, we performed HiChIP on as many cells as we could
obtain from a blood donation - approximately 500,000 - 1 million cells per T cell subtype
per replicate. We performed two minutes of sonication, no Protein A bead preclearing,
used 4 µg of H3K27ac antibody (Abcam ab4729), and captured the chromatin-antibody
complex with 34 µL of Protein A beads (Thermo Fisher). Qubit quantification post ChIP
ranged from 5 – 25 ng depending on the cell type and amount of starting material. The
amount of Tn5 used and PCR cycles performed were based on the post ChIP Qubit
amounts, as previously described[16].
651
25m cell line libraries were generated as previously described[16]. For low cell number
mouse embryonic stem cell samples, we performed two minutes of sonication and no
Protein A bead preclearing. Either 4 µg or 2µg of H3K27ac antibody (Abcam ab4729)
was used for ChIP in 500k or 100k/50k cells, respectively, and the chromatin-antibody
complex was captured with 34 (500k cells) or 20 µL (100k/50k cells) of Protein A beads.
Post-ChIP Qubit quantification for the 25m cell samples was approximately 1.5 µg. For
lower cell numbers, quantification was 30, 10, and 5 ng for 500k, 100k, and 50k cells,
respectively. The amount of Tn5 used and PCR cycles performed were based on the
post ChIP Qubit amounts, as previously described.
661
662 HiChIP samples were size selected by PAGE purification (300-700 bp) for effective
663 paired-end tag mapping, and therefore were removed of all primer contamination which
664 would contribute to recently reported "index switching" on the Illumina HiSeq 4000
665 sequencer[50]. All libraries were sequenced on the Illumina HiSeq 4000 to an average
666 depth of 500-600M total reads.
667
**HiChIP Data Processing**
HiChIP paired-end reads were aligned to hg19 or mm9 genomes using the HiC-Pro
pipeline[51]. Default settings were used to remove duplicate reads, assign reads to MboI
restriction fragments, filter for valid interactions, and generate binned interaction
matrices. HiC-Pro filtered reads were then processed into a .hic file using the
hicpro2juicebox function. The Juicer pipeline HiCCUPS tool was used to identify high
confidence loops[4] using the same parameters as for the GM12878 *in situ* Hi-C map:
hiccups -m 500 -r 5000,10000 -f 0.1,0.1 -p 4,2 -i 7,5 -d 20000,20000 .hic_input
HiCCUPS_output. For T cell Juicer loops, performing the default Juicer calls resulted in
a high rate of false positives upon visual inspection of the interaction matrix. We
therefore called loops with the same HiCCUPS parameters in two biological replicates
for each T cell subtype and then filtered loops for those that were reproducibly called in
both replicates. In addition, we removed all loops greater than 1 Mb.
681
682 1D signal enrichment and peak calling were generated from the HiC-Pro filtered
683 contacts file. Intrachromosomal contacts were filtered and both anchors were extended
684 by 75 base pairs. The combined bed file containing both anchors was then used to

685  generate bigwigs for visualization in the WashU Epigenome Browser or call peaks using
686  MACS2.
687
688  Allele-specific HiChIP data processing was achieved using HiC-Pro's allele-specific
689  analysis features[51]. First, HCASMC phasing data[41] was used to mask the hg19 genome
690  and make indexes. HiC-Pro settings were similar as to described above, with the
691  exception that reads were aligned to the masked genome, and then assigned to a
692  specific allele based on phasing data.
693
694  **Interaction Matrices and Virtual 4C Visualization**
695  HiChIP interaction maps were generated with Juicebox using Knight-Ruiz (KR) matrix
696  balancing and visualized using Juicebox software at 500 kb, 25 kb, 10 kb, and 5 kb
697  resolutions as indicated in each analysis[4]. For 1 kb profiles, raw matrix counts were
698  visualized in Java TreeView.
699
Virtual 4C plots were generated from dumped matrices generated with Juicebox. The
Juicebox tools dump command was used to extract the chromosome of interest from the
.hic file. The interaction profile of a specific 5 kb or 10 kb bin containing the anchor was
then plotted in R. Replicate reproducibility was visualized with the mean profile shown
as a line and the shading surrounding the mean representing the standard deviation
between replicates. For the HCASMC data we observed low read coverage for allele-
specific v4Cs at loci of interest. This is due to the density of SNPs for this genotype, and
a low number of reads containing a phased SNP. We therefore could not observe
interaction profiles when visualizing separate replicates with the standard deviation. We
therefore utilized pseudoreplicates for the HCASMC v4C visualizations[52].
710
711  High confidence Juicer loop calls were loaded into the WashU Epigenome Browser
712  along with corresponding ATAC-seq profiles and publically available H3K27ac ChIP-seq
713  data from the Roadmap Epigenome Project. Browser shots from WashU track sessions
714  were then included in virtual 4C and interaction map anecdotes.
715
**Reproducibility Scatter Plots and Correlations**
Biological and technical reproducibility comparisons between HiChIP experiments were
generated by counting reads supporting a set of Juicer loop calls. For biological
replicate comparisons, the loop set called from the merged replicates was used. For the
comparison in mES cells between low cell number and 25 million cells, the union loop
set between the two maps was used. The Pearson correlation between replicates or
experiments was calculated from depth-normalized reads using the cor() function in R.
Scatter plots were plotted using log-transformed raw reads supporting each loop
724
**Differential Analysis of HiCCUPS Loop Calls**
Juicer loop calls from the three T cell subtypes were initially combined into a union set
of T cell loops. Loop signal was then obtained for the biological replicates of each T cell
subtype. Vanilla coverage square root (VCsqrt) normalized signal for the interaction

matrix of each biological replicate using the Juicebox tools dump command. Normalized
signal was then assigned to the union loop set in each replicate.
731
VCsqrt signal per sample was quantile-quantile normalized under the assumption that
overall signal was identically distributed across all samples. Following normalization,
samples for Naïve, $T_H17$ and $T_{reg}$ cells shared Pearson correlations of 0.938, 0.942 and
0.934, respectively. PCA was performed using the prcomp function in R, which
demonstrated that the first PC, which exhibited near-identical loadings across the six
samples, explained 93% of the variance across the six samples. This was taken to
represent the shared signal across cell types. PCs 2-4 explained 2.2%, 2.0% and 1.4%,
respectively.
740
To study cell type-specific looping, the residual signal per loop was taken after
projecting the loop onto the unit vector along the diagonal (equal signal per cell type).
Cell type-specific and differential looping analysis were performed with the top and
bottom 5% of the distributions of either residual signal or differences between cell type
residual signals. Hierarchical clustering was performed using the union of all differential
loops in these extremes and using 1 minus the Pearson correlation as the distance
metric. QQ plots were generated by permuting residuals from the same cell type or
individual and summing them together and using this distribution to calculate p values
for the observed sums.
750
751    In parallel, differential loops were called using edgeR for both the mES and T cell
752    datasets. Again, biological replicate loop signal was obtained across a union set of
753    Juicer loops. We then used edgeR to identify loops with significant changes in signal
754    among pair-wise comparisons (FDR < 0.1, $\log_2FC > 1$). Importantly, inspection of
755    differential loops identified from the two methods revealed high concordance.
756
Gene density was calculated from Ensembl gene annotations. GC content was
calculated per 10 kb bin using the bedtools nuc function and aggregated as needed.
Notably, Spearman correlation between gene density of an entire chromosome and the
number of differential loop anchors (rho = 0.914) was much higher than the correlations
between the variance in cell type signal per loop anchor and number of genes per 10 kb
window (rho = 0.322) and between differential loop anchors and gene density per 100
kb section (rho = 0.083). Correlations between GC content and number of differential
loops were similar at both the chromosome (rho = 0.729) and 100 kb bin (rho = 0.148)
levels, but while local GC content is likely to confound relative abundance, it is unclear
how chromosome-wide GC content could have the same effect.
767
For mES analysis of H3K27ac- and cohesin-mediated HiChIP, we performed edgeR to
obtain the biased loops for each factor. To determine functional bias of the top loops,
overlap was determined between edgeR differential loop anchors and relevant ChIP-seq
peaks. Smc1a ChIP-seq peaks were obtained from a published dataset[53]. CTCF, RNA

Polymerase II, and H3K27ac ChIP-seq peaks were obtained from the mouse ENCODE
repository[54].

**Promoter/Enhancer Annotations of HiChIP Loops**
775

776  Promoters were defined as 1 kb regions centered at the TSS, and enhancers were
777  identified as chromHMM enhancers not overlapping with promoters in any cell type. We
778  annotated loop anchors as 'other' if the anchor did not contain a promoter or enhancer
779  as defined above.

**RNA Expression Analysis**
781

Previously generated RNA-seq data[55] from Naïve, $T_H17$ and $T_{reg}$ cells was downloaded
as fastqs from ArrayExpress. Illumina adaptors were trimmed using CutAdapt and
Ensembl's cDNA transcripts were quantified using kallisto. Sleuth was used to identify
transcripts that were differentially expressed across cell types with FDR controlled at
5%. The mean TPM was calculated per cell type, and TSS differential looping quantiles
at genes with with nonzero expression were correlated with differential expression
quantiles of the same genes. Only 10 kb segments of the genome that contained a
single annotated gene were considered to avoid errors in attribution of looping signal
per 10 kb bin. For genes with multiple annotated TSSs the 10 kb bin corresponding to
the median TSS was used. Significance was assessed by the cor.test function in R.

**Comparing HiChIP to Hi-C, ATAC-seq, ChIP-seq, and Capture-C datasets**
793

T cell subset ChIP-seq data was obtained from the WashU Roadmap repository. A
union set of peaks were called using MACS2 and peaks were quantified using Bedtools
intersect. Normalization for ChIP was performed using quantile normalization using
"preprocessCore" package in R.

Differential EIS was determined using TMM normalization in the "edgeR" package in R.
The significantly differential EIS ($\log_2$ fold-change > 1 and FDR < 10%) were determined
for each pairwise comparison. For each differential EIS, the maximum ATAC/ChIP
signal peak was assessed in each 10 kb anchor (to bias against low signal peaks) and
then the maximum $\log_2$ fold-change was compared to the differential EIS.

Differential 1D HiChIP was determined using TMM normalization in the "edgeR"
package in R. The significantly differential 1D HiChIP in EIS 10 kb anchors ($\log_2$ fold-
change > 1 and FDR < 10%) were determined for each pairwise comparison. For each
differential 1D HiChIP anchor, the maximum ATAC/ChIP signal peak was assessed (to
bias against low signal peaks) and then the maximum $\log_2$ fold-change was compared
to the differential 1D HiChIP.

To determine if reference cell line Hi-C data with primary T cell H3K27ac ChIP-seq
could recapitulate EIS in primary T cells, GM Hi-C loop anchors were binned with
increasing T cell subset ChIP signal as well as GM H3K27ac ChIP-seq signal (Encode).
Loop overlap was then determined for the different H3K27ac ChIP signal bins with
HiChIP loops.

817
To compare directly with CD4$^+$ Capture-C data, CHiCAGO loops were called in Naïve,
$T_H$17, and $T_{reg}$ datasets. CHiCAGO calls were then combined into a union set, and loop
signal was obtained for the biological replicates of each T cell subtype using the
Juicebox tools dump command. We then used edgeR to identify loops with significant
changes in signal among pair-wise comparisons (FDR < 0.05). Total and differential
loops were then overlapped with CD4$^+$ Capture-C CHiCAGO data.
824

**Distance-matched eQTL SNP-TSS Comparisons**
We obtained three groups of eQTL SNP-TSS pairs within 1 Mb distance for HiChIP EIS
comparisons. The treatment group contains CD4 eQTL-TSS targets. We have two
distance-matched groups as control. The first control group contains the CD4 eQTL
SNP-random TSS pairs such that the distance between eQTL SNP and random TSS
differs by at most 5 kb with the treatment group. The second control group contains liver
eQTL SNP-TSS targets that are also distance matched with the treatment group. The
random eQTL SNP-TSS pairs were generated by individual chromosome, such that
number of control pairs and treatment pairs are the same for every chromosome. In
total, there were 158,482 distance-matched eQTL-TSS pairs. We compared the 5 kb
resolution EIS among the three eQTL SNP-TSS groups for all three T cell subtypes.
Results show that in all the cases, the EIS between CD4 eQTL-TSS targets were
significantly higher than the two control groups (p-value < $10^{-16}$).
838

**Distance-matched Fine-mapped SNP-TSS Comparisons**
We obtained a list of putatively causal SNPs from the PICS SNP list[25] (PICS probability
> 0.5), as well as fine-mapped SNPs associated with inflammatory bowel disease
(IBD)[38] or Type 1 Diabetes (T1D)[39]. Next, we obtained all SNPs in LD with each
putatively causal SNP using European linkage disequilibrium blocks determined by all
SNPs with an $r^2 \geq 0.8$ with the SNP being considered. For the fine-mapped (T1D/IBD)
sets, using SNPs in LD with highly significant GWAS SNPs may mean that there are
several SNPs of equal or greater significance in the control set, but we still expect an
enrichment relative to the LD block.
848

We collected all the synthetic pairs between the putatively causal immune-disease
related SNPs (IBD, T1D and PICS) and nearby genes within 300 kb distance. To
perform the distance-matched EIS comparisons, for each fine-mapped SNP category,
we selected the SNP-TSS control pairs which satisfy two constraints: (1) the selected
control SNP is positioned at least 5 kb away from the fine-mapped SNP in the same
linkage disequilibrium block; (2) the distance of SNP-TSS control pair differs with fine-
mapped SNP and the target gene at most 5 kb.
856

**Calculation of Disease-specific GWAS SNP Enrichment in Loop Anchors**
We categorized GWAS SNPs into sets relevant to diseases of the immune system and,
separately, diseases with no known immune component[25]. For each disease in the
immune or non-immune set, we determined the proportion of all GWAS SNPs

associated with that disease which overlap the positions of loop anchors based on a
union set of loops identified in Naïve, $T_H17$, and $T_{reg}$ cells. The ratio of the proportion of
immune and non-immune overlaps relative to a shuffled control was reported as the
enrichment of immune GWAS SNPs.
865
**SNP-TSS Loop Analyses**
We obtained 7,747 PICS SNPs that are associated with autoimmune disease or non-
immune traits[25]. 4,331 (55.9%) were associated with autoimmune disease, and 3,416
(44.1%) were associated with non-immune traits. In addition we obtained a set of SNPs
associated with six overlapping autoimmune diseases using the GRASP catalog
(genome-wide significance p-value $< 10^{-8}$).
872
873  We constructed a synthetic loop set for immune and non-immune SNPs and any TSS
874  within 1 Mb of each SNP. We then assigned VCsqrt signal in each biological replicate of
875  the three T cell subtypes to the synthetic loop set, as described above.
876
VCsqrt signal per sample was quantile-quantile normalized as above. In this analysis,
we did not restrict to HiCCUPS-identified loops but instead examined all possible
interactions between a SNP and TSS within 1 Mb. Many of these interactions do not
exist, and therefore had little or no matrix-balanced signal supporting them. While we
removed all SNP TSS pairs below an average of 1 normalized read per sample from
subsequent analyses, in general these false interactions contributed little to the overall
differential signal for a trait.
884
885  H3K27ac data was downloaded from the WashU Roadmap repository. PICS SNPs were
886  taken from Farh, et al[25]. Rather than require strict membership within H3K27ac peaks,
887  PICS SNPs were labeled as active if they lied within 8 kb of a peak, raising the number
888  of nominally functional SNPs from ~700 to ~3200 per cell type, out of 7735 total
889  candidate SNPs.
890
Differential looping across cell types was assessed by one-sided t-test per trait and
activity partition if there were at least eight PICS SNP-TSS pairs in the partition. $T_H17$
bias was defined as $T_H17$ total loop signal minus Naïve total loop signal; $T_{reg}$ bias was
defined as $T_{reg}$ total loop signal minus Naïve total loop signal; and Naïve bias was
defined as Naïve total loop signal minus one-half times $T_H17$ and $T_{reg}$ total loop signal.
For the main text figure, Naïve bias was assessed only using SNPs that were active in
Naïve cells, $T_H17$ bias from SNPs active in $T_H17$ and not Naïve, and $T_{reg}$ bias from
SNPs active in $T_{reg}$ and not Naïve. p-values were corrected for multiple hypothesis
testing by the Holm method using the p.adjust function in R. Bias assessed from SNPs
with the opposing cell type specificities (e.g. Naïve bias using SNPs active in $T_H17$ and
$T_{reg}$ but not Naïve) yielded no significant hits after correction.
902
**Correction for False Positives**

Despite not being originally implemented in enrichment interaction datasets, we have
previously demonstrated that KR and VC matrix balancing corrects for false positives
caused by high 1D fragment visibility[16]. Loop calls are either from HiCCUPS or Fit Hi-C,
which are KR and VC matrix balanced, respectively. For differential analysis of
HiCCUPS (Figure 3) and SNP-TSS Fit Hi-C (Figure 4c) loop calls, we are restricted
within a set of loops that were obtained from matrix balancing. Therefore, while
differential loops can be driven by both changes in looping strength and/or 1D ChIP
signal, the final interactions being observed are loops. Additionally, we performed
differential analysis of HiCCUPS loop calls using two separate methods – one using
VCsqrt normalized reads and another with non-normalized reads (edgeR) and found
high agreement. The SNP-TSS synthetic loop analysis in Figure 4b was not restricted to
loop calls, however was performed with VCsqrt matrix balanced reads to avoid false
positives. Those SNP-TSS synthetic loops were then subset by overlap with Fit Hi-C for
further differential analyses in Figure 4c. Finally, virtual 4C analysis was performed on
non-normalized reads to highlight EIS contributions of both 1D and 3D signal changes,
however HiCCUPS loop calls are included in the relevant anecdotes.
920
921  **SNP and TSS Connectivity Analysis**
922  Among all immune and non-immune SNPs, 2,562 (33.1%) were located within a 25 kb
923  region centered around TSS, 2,618 (33.8%) were located in a gene body, and the
924  remaining 2,567 (33.1%) were located in intergenic regions.
925
To capture the confident SNP-TSS connections for SNP-TSS pair identification, we
overlapped the anchors of significant loops that were identified by Fit-Hi-C[56] with the
SNP/TSS locations. We identified 14,738 SNP-TSS pairs that were supported by Fit-Hi-
C loops, and there were 3,046 unique SNPS connected with at least one gene. Among
those 3,046 SNPs, there were 2,181 (71.6%) and 865 (28.4%) SNPs annotated with
immune and non-immune SNPs, respectively. As expected, the immune disease SNPs
are more likely to connect with genes (Fisher's exact test, p-value = 4.8 x $10^{-85}$).
933
**Phasing of HCASMC Samples**
We used BEAGLE 4.1 to impute and phase recalibrated variants using 1000 Genome
phase 3 version 5a as a reference panel. The Beagle phasing algorithm was set on the
following criteria. At each iteration that the algorithm performs, phased input data are
used to build a local model of a haplotype-cluster. After the local haplotype-cluster
model is created, for each individual phased haplotypes are sampled using the induced
diploid HMM conditional on the individual's genotypes. The sampled haplotypes are
then used as the input to feed in the next iteration of the algorithm. In the final iteration,
instead of sampling haplotypes, BEAGLE uses the Viterbi algorithm to
select the conditional on the diploid HMM and the individual's genotype data and to
obtain the haplotypes for each individual that possess the greatest probability, and
these most-likely haplotypes are the final output of the BEAGLE phasing algorithm.
946
**Allelic Mapping Bias Simulation**

We constructed a personal genome by editing the reference genome (hg19) according
to SNP information. SNPs labeled as "1l0" or "1l1" in the vcf file were replaced with the
alternative allele for genome 1. SNPs labeled as "0l1" or "1l1" in the vcf file were
replaced with the alternative allele for genome 2.  Loop anchors were extended 100 bp
in both directions and sequences were extracted using samtools for each
genome. ~20X reads were simulated for each region, using wgsim with parameters " -e
0.01  -d 100 -s 20 -1 75  -2 75 -S -1  -h -R 0.1".  Simulated reads were mapped to the
"N" masked genome. Mapping parameters were the same used by HiC-Pro ("--very-
sensitive -L 30 --score-min L, -0.6, -0.2 --end-to-end --reorder --phred33-quals).  Allelic
specific reads were separated according to the SNP information and then counted for
each loop anchor using bedtools.
959

**ATAC-seq**
Cells were isolated and subjected to ATAC-seq as previously described[16]. Briefly,
55,000 cells were pelleted, resuspended in 50 µL lysis buffer (10mM Tris-HCl, pH 7.4,
3mM MgCl2, 10mM NaCl, 0.1% NP-40 (Igepal CA-630), and immediately centrifuged at
500 rcf for 10 min at $4^0$C. The nuclei pellets were resuspended in 50 µL transposition
buffer (25 µl 2X TD buffer, 22.5 µL $dH_20$, 2.5 µL Illumina Tn5 transposase), and
incubated at $37^0$C for 30 min. Transposed DNA was purified with MinElute PCR
Purification Kit (Qiagen), and eluted in 10 µL EB buffer.
968

**ATAC-seq Data Processing**
Adapter sequence trimming using SeqPurge and mapping to hg19 using Bowtie2 were
performed. These reads were then filtered for mitochondrial reads, low quality, and PCR
duplicates. The filtered reads for each sample were merged and peak calling was
performed by MACS2. Each individual sample reads in peaks were quantified using
Bedtools intersect with the MACS2 narrow peaks. Peak counts were then combined into
a matrix NxM where N represents called peaks and M represents the samples and each
value Di,j represents the peak intensity for the respective peak i in sample j. This matrix
was then normalized using the "CQN" package in R to minimize bias in GC content and
length.
979

**CRISPRi Validation of HiChIP Targets**
For virus production, $5 \times 10^6$ of HEK293T cells were plated per 10 cm plate. The
following day, plasmid encoding lentivirus was co-transfected with pMD2.G and psPAX2
into the cells using Lipofectamine 3000 (Thermo Fisher, L3000) according to the
manufacturer's instructions. Supernatant containing viral particles was collected 48
hours after transfection and filtered. For lentivirus encoding individual sgRNAs, virus
was concentrated 10-fold using Lenti-X concentrator (Clontech, 631232) and stored at -
80°C.
988

In order to generate a My-La cell line expressing CRISPRi, $2 \times 10^6$ of My-La cells were
plated per T75 flask. A dCas9-BFP-KRAB-2A-Blast construct was generated by
inserting a 2A-Blast cassette into dCas9-BFP-KRAB (Addgene 46911). 24 hours after

plating, lentivirus harboring the dCas9-KRAB construct was added with polybrene (4 µg
/ mL). Media was changed 24 hours after infection, and then again 48 hours after
infection with Blasticidin (Thermo Fisher, A1113903) at 4 µg / mL.  Blasticidin resistant
cells were selected for eight days with changing media every other day.
996
Three different U6 were used for transcription of three different sgRNAs targeting the
candidate enhancers, as previously described[57]. For *MYC* locus CRISPRi experiments,
each enhancer was targeted by one guide, and therefore all *MYC* GM or My-La
enhancers together were targeted in one experiment. For the PICS SNP CRISPRi
experiments, three guides were targeted to a single SNP-containing enhancer. One of
three sgRNAs was cloned into a lentiviral vector with a human (pMJ117, Addgene
85997), mouse (pMJ179, Addgene 85996) or bovine (pMJ114, Addgene 85995) U6
promoter. These U6-sgRNA constructs were then combined into lentivirus with a
Puromycin-2A-mCherry vector, which was modified from Addgene 46914. My-La-
CRISPRi cells were infected with lentivirus harboring 3 sgRNAs and selected by
Puromycin (Thermo Fisher, A11138) at a final concentration of 1 µg / mL. Previously
reported sgRNAs targeting VPS54 or SEC24C were used for validating CRISPRi
functionality in My-La cell line[58].
1010
For readout of CRISPRi validation, we performed qRT-PCR and cell growth assays on
three biological and two technical replicates. For qRT-PCR, RNA was Trizol extracted
(Thermo Fisher, 15596026) and purified using the Zymo RNA Clean and Concentrator
kit (Zymo Research, R1016). qRT-PCR was performed with Brilliant qRT-PCR
mastermix (Agilent, 600825). Ct values were measured by using Lightcycler 480
(Roche) and relative expression level was calculated by ddCt method compared to a
*GAPDH* control. Primer sequences are listed in Supplementary Table 6. For cell growth,
we used the CellTiter-Glo kit (Promega, G7572) according to the manufacturers
instructions. Statistics for both RNA and cell growth changes were calculated using a
Student's t test against the non-targeting control.
1021
1022  **CRISPRa Validation of HiChIP Targets**
1023  Jurkat cells were transduced with a lentiviral dCas9-VP64-2A-GFP expression vector
1024  (Addgene 61422). Single GFP+ cells were sorted by FACS into the wells of a 96-well
1025  plate, and a clone with bright uniform GFP expression were selected for use in future
1026  experiments.
1027
sgRNAs were cloned in arrayed format for CD69 HiChIP peaks falling outside the range
of the tiling CRISPRa screen[26]. sgRNAs were chosen based on high predicted on-target
activity[59] and low predicted off-target activity[60]. sgRNAs were cloned into the lentiviral
expression vector "pCRISPRia-v2" (Addgene 84832) as described in Horlbeck et al[61].
Lentivirus was produced by transfecting HEK293T with standard packaging vectors
using *Trans*IT-LTI Transfection Reagent (Mirus, MIR 2306). Media was changed 24
hours post-transfection. Viral supernatant was harvested at 48 and 72 hr following
transfection and immediately used for infection of Jurkat-dCas9-VP64 cells.

1036
Jurkat-dCas9-VP64 cells were infected with lentiviral sgRNAs by resuspending cells in a
1:1 mix of fresh media and lentiviral supernatant at a final concentration of $0.25 \times 10^6$
cells/mL with 5 µg / mL polybrene. Cells were spinfected for 1 hour at 1000 rcf, 32 °C.
The next day, half of the media was removed and replaced with fresh lentiviral
supernatant, and the spinfection was repeated. The next day, the cells were
resuspended in fresh media with 1.5 µg / mL puromycin and cultured for 2 days to
remove uninfected cells. For readout of CRISPRa validation, we performed qRT-PCR
and FACS on two biological and two technical replicates. RNA extraction and qRT-PCR
was performed as described above. Expression of CD69 on infected cells (GFP+BFP+)
was analyzed by flow cytometry with an Attune NxT flow cytometer (Life Technologies).
Statistics for both RNA and protein level changes were calculated with a one-way
ANOVA followed by a Dunnet's multiple comparisons test against the non-targeting
control.
1050
1051    **SUPPLEMENTARY FIGURE LEGENDS**
1052
**Supplementary Figure 1. H3K27ac HiChIP enriches E-P-associated chromatin**
**contacts. (A)** Schematic of chromatin contacts captured in H3K27ac HiChIP. **(B)** Loop
call overlap for cohesin HiChIP and H3K27ac HiChIP in mES cells. **(C)** Contact distance
distribution for loops that are biased in cohesin versus H3K27ac HiChIP. **(D)** Proportion
of cohesin and H3K27ac biased HiChIP loops that have cohesin, CTCF, H3K27ac, and
RNA polymerase II binding in at least one loop anchor. **(E)** Virtual 4C interaction profile
of an H3K27ac-biased loop focused at the *MALAT1* promoter. **(F)** Virtual 4C interaction
profile of a cohesin-biased loop with associated low transcriptional activity.
1061
**Supplementary Figure 2. H3K27ac HiChIP achieves high chromatin loop signal-**
**over-background at low cell inputs. (A)** KR balanced interaction matrices focused
around the *Etv5* locus in mES cells with decreasing cellular starting material. **(B)** Read
support reproducibility of loops in H3K27ac HiChIP libraries from 25 million cells
compared to HiChIP in lower cell input libraries. **(C)** Aggregate peak analysis of loops in
mES H3K27ac HiChIP libraries.
1068
**Supplementary Figure 3. H3K27ac HiChIP generates reproducible chromatin loop**
**signals at low cell inputs. (A)** Comparison of KR balanced interaction maps in
H3K27ac HiChIP biological replicates. **(B)** Read support reproducibility of loops
between H3K27ac HiChIP biological replicates. **(C)** HiCCUPS loop call overlap between
H3K27ac HiChIP libraries from 25 million and 50 thousand mES cells. **(D)** Preseq library
complexity estimation of H3K27ac HiChIP libraries from 25 million and 50 thousand
mES cells.
1076
**Supplementary Figure 4. H3K27ac HiChIP biological replicates from primary**
**sorted T cells are highly reproducible. (A)** FACS strategy for Naïve, $T_H17$, and $T_{reg}$
cells starting from human peripheral blood. Post-sort validation was used to ensure

purity of T cell subtypes. Number represents percent of cells within that gate. **(B)** KR
balanced interaction map of T cell subtype biological replicates. **(C)** Read support
reproducibility of loops between H3K27ac HiChIP biological replicates in Naïve, $T_H17$,
and $T_{reg}$ cells. **(D)** Aggregate peak analysis of loops in Naïve, $T_H17$, and $T_{reg}$ H3K27ac
HiChIP libraries.

1085

1086 **Supplementary Figure 5. Validation of HiChIP-identified *CD69* distal enhancers**
1087 **with CRISPR activation.** CRISPRa validation in Jurkat cells of *CD69* distal enhancers.
1088 *CD69* protein levels are shown for individual sgRNAs tiling H3K27ac HiChIP-identified
1089 distal *CD69* enhancers relative to the *KLRF2* promoter as a locus negative control and a
1090 non-targeting negative control.

1091

**Supplementary Figure 6. Global enhancer connectome characterization in T cell**
**differentiation. (A)** ChromHMM classification of union T cell loop anchors. **(B)** Contact
distance distribution of union T cell loops. **(C)** Agreement in residual signal observed
between sample signals per loop after removing imputed shared signal, calibrated by a
null distribution of random pairings of loops. QQ plot shows modest enrichment above
random pairings. PCA on residual signal clusters samples first by Naive vs Memory cell
types (PC1), and then by donor identity (PC2, 3). **(D)** Overlap of differential interactions
between Naïve, $T_H17$, and $T_{reg}$ subtypes. Biased interactions were obtained by
performing pairwise comparisons between T cell types and analyzing the top 5%
enriched and top 5% depleted EIS in each T cell subtype. **(E)** ChromHMM annotation of
total loops, differential loops, and shared loops in all three T cell subtypes. O
corresponds to other loop anchors not classified as enhancer or promoter. **(F)** Number
of connections for different classes of loop elements. **(G)** Quantification of promoters
skipped before an enhancer reaches its loop target.

1106

1107 **Supplementary Figure 7. Positioning of differential HiChIP contacts in gene**
1108 **dense chromosomes. (A)** Distribution of T cell subtype differential HiChIP contacts
1109 across different chromosomes compared to the distribution of all loops. **(B)** Correlation
1110 of differential loop density with gene density and GC content.

1111

1112 **Supplementary Figure 8. Characterization of conformational landscapes**
1113 **surrounding key T cell regulatory factors. (A – C)** Virtual 4C interaction profiles
1114 anchored at the promoters of canonical Naïve, $T_H17$, and $T_{reg}$ regulatory factors.

1115

1116 **Supplementary Figure 9. Chromosome conformation dynamics of canonical T**
1117 **cell regulatory factors. (A – C)** Delta interaction maps focused around known Naïve,
1118 $T_H17$, and $T_{reg}$ regulatory factors.

1119

**Supplementary Figure 10. Contribution of H3K27ac ChIP and chromosome**
**conformation to HiChIP EIS. (A)** (Left) Proportion of H3K27ac ChIP peaks within EIS
differential loop anchors that are cell-type specific ($\log_2$ fold change > 1) or shared
across all three subtypes. (Right) Global correlation of EIS and H3K27ac ChIP fold-

change in different T cell subset pairwise comparisons. **(B)** Same as (A), but using
HiChIP 1D differential signal at EIS biased loop anchors. **(C)** Overlap of H3K27ac
HiChIP and bins of Hi-C loops with increasing T cell subset and GM H3K27ac ChIP-seq
signal. **(D)** Overlap of CD4[+] Capture Hi-C[14] with total and differential T cell subset
HiChIP loops. **(E)** T$_{reg}$-specific loops at the *LRRC32* promoter not observed in other
H3K27ac HiChIP T cell subsets nor in CD4[+] Capture Hi-C data[14].
1130
**Supplementary Figure 11. Enrichment of autoimmune SNPs in T cell HiChIP loop**
**anchors. (A)** Enrichment of specific PICS autoimmune disease and non-immune SNPs
in anchors of loops called by Juicer and Fit-Hi-C compared to a background shuffled
loop set. **(B)** Enrichment of all PICS autoimmune disease and non-immune SNPs in T
cell subset biased loop anchors and all anchors compared to a background shuffled
loop set.
1137
**Supplementary Figure 12. T cell subtype HiChIP specificity of autoimmune SNPs.**
**(A)** H3K27ac HiChIP signal bias in T cell subsets for PICS SNP-TSS pairs grouped by
each SNP's presence in cell type-specific or shared H3K27ac ChIP peaks up to 8 kb
away. **(B)** H3K27ac HiChIP signal bias in T cell subsets for PICS SNP-TSS pairs
grouped by each SNP's presence in cell type-specific or shared H3K27ac ChIP peaks
up to 2.5 kb away. **(C)** H3K27ac HiChIP signal bias in GM, K562, and My-La cell lines
for PICS SNP-TSS pairs grouped by each SNP's presence in T cell subset-specific or
shared H3K27ac ChIP peaks up to 8 kb away. **(D)** H3K27ac HiChIP signal bias in T cell
subsets for GRASP SNP-TSS pairs grouped by each SNP's presence in cell type-
specific or shared H3K27ac ChIP peaks up to 2.5 kb away. **(E)** Average number of
HiChIP gene targets for non-genic autoimmune disease and non-immune SNPs. **(F)**
Quantification of promoters skipped before a SNP reaches its gene target. **(G)**
Quantification of SNP HiChIP gene targets in autoimmune disease.
1151
1152  **Supplementary Figure 13. Validation of HiChIP signal at SNP-eQTL contacts. (A)**
1153  Validation of HiChIP signal at SNP-TSS pairs using interaction profiles of eQTL SNPs to
1154  ensure they contact their associated target gene promoter. **(B)** Interaction profiles of
1155  CRISPRi-validated loci in My-La.
1156
**Supplementary Figure 14. H3K27ac HiChIP fine-mapping of GWAS variants in**
**haplotype blocks. (A)** Global validation of HiChIP signal at putatively causal SNPs
versus corresponding SNPs in LD ($r^2 \geq 0.8$) for Naïve and T$_{reg}$ cell subtypes. SNP-TSS
pairs were generated from published fine-mapped datasets, compared to a distance-
matched SNP-TSS pair set in the same LD block. **(B)** Interaction profile of the *SATB1*
promoter, and a 1 kb resolution visualization of the SNP-containing enhancer of interest.
Changes in bias between 5 kb and 1 kb resolution reflects EIS focused 1 kb around the
SATB1 TSS and specific SNPs within the enhancer. LD SNPs ($r^2 \geq 0.8$) correspond to
GRASP SNPs (genome-wide significance p-value $< 10^{-8}$). The highlighted SNP is a
PICS closest to focal EIS to *SATB1.*
1167

1168 **Supplementary Figure 15. Chromatin interaction landscape of the 9p21.3**
1169 **cardiovascular disease risk locus.** HCASMC v4C interaction profiles focused around
1170 the promoters of *CDKN2A*, *CDKN2B*, and *ANRIL* within the 9p21.3 locus.
1171
1172 **Supplementary Table Legends**
1173
1174 **Supplementary Table 1. HiChIP data processing metrics.** HiC-Pro mapping, Hi-C
1175 filtering, duplicate removal, and interaction length statistics for all HiChIP libraries.
1176
1177 **Supplementary Table 2. HiCCUPS high confidence loop calls.** High confidence loop
1178 calls for all HiChIP libraries.
1179
1180 **Supplementary Table 3. HiCCUPS differential EIS in T cell subtypes by edgeR.**
1181 High confidence loop calls in T cell subsets with edgeR significance and fold-change for
1182 pair-wise comparisons.
1183
1184 **Supplementary Table 4. HiCCUPS differential EIS in T cell subtypes.** Top 5% of
1185 EIS ranked by cell-type bias for each pair-wise comparison between different T cell
1186 subtypes.
1187
1188 **Supplementary Table 5. HiChIP gene targets of autoimmune and CAD SNPs.** SNP
1189 type (within gene, at TSS, or intergenic), associated disease, and corresponding loop
1190 annotated with the HiChIP gene target. Included are edgeR significance and fold-
1191 change for cell-type comparisons of interest.
1192
1193 **Supplementary Table 6. sgRNA and primer oligonucleotide sequences.** sgRNA
1194 and primer sequences used throughout the study.
1195
1196 **REFERENCES**
1197

1. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of

chromatin interactions. *Nature* **485,** 376–380 (2012).

2. van Arensbergen, J., van Steensel, B. & Bussemaker, H. J. In search of the determinants of

enhancer-promoter interaction specificity. *Trends Cell Biol.* **24,** 695–702 (2014).

3. Lieberman-Aiden, E. *et al.* Comprehensive mapping of long-range interactions reveals

folding principles of the human genome. *Science* **326,** 289–293 (2009).

4. Rao, S. S. P. *et al.* A 3D map of the human genome at kilobase resolution reveals

principles of chromatin looping. *Cell* **159,** 1665–1680 (2014).

5. Fulco, C. P. *et al.* Systematic mapping of functional enhancer-promoter connections with

CRISPR interference. *Science* aag2445 (2016). doi:10.1126/science.aag2445

6. Altshuler, D., Daly, M. J. & Lander, E. S. Genetic mapping in human disease. *Science* **322,**

881–888 (2008).

7. Kumar, V. *et al.* Human Disease-Associated Genetic Variation Impacts Large Intergenic

Non-Coding RNA Expression. *PLOS Genet.* **9,** e1003201 (2013).

8. Edwards, S. L., Beesley, J., French, J. D. & Dunning, A. M. Beyond GWASs: Illuminating the

Dark Road from Association to Function. *Am. J. Hum. Genet.* **93,** 779–797 (2013).

9. Welter, D. *et al.* The NHGRI GWAS Catalog, a curated resource of SNP-trait associations.

*Nucleic Acids Res.* **42,** D1001–D1006 (2014).

10.    Smemo, S. *et al.* Obesity-associated variants within FTO form long-range functional

connections with IRX3. *Nature* **507,** 371–375 (2014).

11.    Claussnitzer, M. *et al.* FTO Obesity Variant Circuitry and Adipocyte Browning in

Humans. *N. Engl. J. Med.* **373,** 895–907 (2015).

12.    Grubert, F. *et al.* Genetic Control of Chromatin States in Humans Involves Local and

Distal Chromosomal Interactions. *Cell* **162,** 1051–1065 (2015).

13.    Mifsud, B. *et al.* Mapping long-range promoter contacts in human cells with high-

resolution capture Hi-C. *Nat. Genet.* **47,** 598–606 (2015).

14.    Javierre, B. M. *et al.* Lineage-Specific Genome Architecture Links Enhancers and Non-

coding Disease Variants to Target Gene Promoters. *Cell* **167,** 1369–1384.e19 (2016).

15.    Sanjana, N. E. *et al.* High-resolution interrogation of functional elements in the

noncoding genome. *Science* **353,** 1545–1549 (2016).

16.    Mumbach, M. R. *et al.* HiChIP: efficient and sensitive analysis of protein-directed

genome architecture. *Nat. Methods* **advance online publication,** (2016).

17.    Heintzman, N. D. *et al.* Distinct and predictive chromatin signatures of

transcriptional promoters and enhancers in the human genome. *Nat. Genet.* **39,** 311–318

(2007).

18.    Creyghton, M. P. *et al.* Histone H3K27ac separates active from poised enhancers and

predicts developmental state. *Proc. Natl. Acad. Sci. U. S. A.* **107,** 21931–21936 (2010).

19.    Rada-Iglesias, A. *et al.* A unique chromatin signature uncovers early developmental

enhancers in humans. *Nature* **470,** 279–283 (2011).

20.    Ferraro, A. *et al.* Interindividual variation in human T regulatory cells. *Proc. Natl.*

*Acad. Sci. U. S. A.* **111,** E1111-1120 (2014).

21.    Arvey, A. *et al.* Genetic and epigenetic variation in the lineage specification of

regulatory T cells. *eLife* **4,** e07571 (2015).

22.    Bettelli, E. *et al.* Reciprocal developmental pathways for the generation of

pathogenic effector TH17 and regulatory T cells. *Nature* **441,** 235–238 (2006).

23.    Acosta-Rodriguez, E. V. *et al.* Surface phenotype and antigenic specificity of human

interleukin 17-producing T helper memory cells. *Nat. Immunol.* **8,** 639–646 (2007).

24.    Morita, R. *et al.* Human blood CXCR5(+)CD4(+) T cells are counterparts of T

follicular cells and contain specific subsets that differentially support antibody secretion.

*Immunity* **34,** 108–121 (2011).

25.    Farh, K. K.-H. *et al.* Genetic and epigenetic fine mapping of causal autoimmune

disease variants. *Nature* **518,** 337–343 (2015).

26.      Simeonov, D. R. *et al.* Discovery of an autoimmunity-associated IL2RA enhancer by

unbiased targeting of transcriptional activation. *bioRxiv* 091843 (2016).

doi:10.1101/091843

27.      Engreitz, J. M. *et al.* Local regulation of gene expression by lncRNA promoters,

transcription and splicing. *Nature* **539,** 452–455 (2016).

28.      Dao, L. T. M. *et al.* Genome-wide characterization of mammalian promoters with

distal enhancer functions. *Nat. Genet.* **49,** 1073–1081 (2017).

29.      Ernst, J. & Kellis, M. ChromHMM: automating chromatin-state discovery and

characterization. *Nat. Methods* **9,** 215–216 (2012).

30.      Sanborn, A. L. *et al.* Chromatin extrusion explains key features of loop and domain

formation in wild-type and engineered genomes. *Proc. Natl. Acad. Sci. U. S. A.* **112,**

E6456-6465 (2015).

31.      Li, G. *et al.* Extensive promoter-centered chromatin interactions provide a

topological basis for transcription regulation. *Cell* **148,** 84–98 (2012).

32.      Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J. Transposition

of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-

binding proteins and nucleosome position. *Nat. Methods* **10,** 1213–1218 (2013).

33.      Samstein, R. M. *et al.* Foxp3 exploits a pre-existent enhancer landscape for

regulatory T cell lineage specification. *Cell* **151,** 153–166 (2012).

34.      Ye, C. J. *et al.* Intersection of population variation and autoimmunity genetics in

human T cell activation. *Science* **345,** 1254665 (2014).

35.      Petukhova, L. *et al.* Genome-wide association study in alopecia areata implicates

both innate and adaptive immunity. *Nature* **466,** 113–117 (2010).

36.    Raj, T. *et al.* Polarization of the effects of autoimmune and neurodegenerative risk

alleles in leukocytes. *Science* **344,** 519–523 (2014).

37.    Schiering, C. *et al.* The alarmin IL-33 promotes regulatory T-cell function in the

intestine. *Nature* **513,** 564–568 (2014).

38.    Huang, H. *et al.* Fine-mapping inflammatory bowel disease loci to single-variant

resolution. *Nature* (2017). doi:10.1038/nature22969

39.    Onengut-Gumuscu, S. *et al.* Fine mapping of type 1 diabetes susceptibility loci and

evidence for colocalization of causal variants with lymphoid gene enhancers. *Nat. Genet.*

**47,** 381–386 (2015).

40.    O'Shea, J. J., Lahesmaa, R., Vahedi, G., Laurence, A. & Kanno, Y. Genomic views of

STAT function in CD4+ T helper cell differentiation. *Nat. Rev. Immunol.* **11,** 239–250

(2011).

41.    Miller, C. L. *et al.* Integrative functional genomics identifies regulatory mechanisms

at coronary artery disease loci. *Nat. Commun.* **7,** 12092 (2016).

42.    Nurnberg, S. T. *et al.* Coronary Artery Disease Associated Transcription Factor

TCF21 Regulates Smooth Muscle Precursor Cells That Contribute to the Fibrous Cap.

*PLoS Genet.* **11,** e1005155 (2015).

43.    McPherson, R. *et al.* A common allele on chromosome 9 associated with coronary

heart disease. *Science* **316,** 1488–1491 (2007).

44.    Helgadottir, A. *et al.* A common variant on chromosome 9p21 affects the risk of

myocardial infarction. *Science* **316,** 1491–1493 (2007).

45.    Clarke, R. *et al.* Genetic variants associated with Lp(a) lipoprotein level and coronary

disease. *N. Engl. J. Med.* **361,** 2518–2528 (2009).

46.    CARDIoGRAMplusC4D Consortium *et al.* Large-scale association analysis identifies

new risk loci for coronary artery disease. *Nat. Genet.* **45,** 25–33 (2013).

47.    Leung, D. *et al.* Integrative analysis of haplotype-resolved epigenomes across human

tissues. *Nature* **518,** 350–354 (2015).

48.    Dixon, J. R. *et al.* Chromatin architecture reorganization during stem cell

differentiation. *Nature* **518,** 331–336 (2015).

49.    Franzén, O. *et al.* Cardiometabolic risk loci share downstream cis- and trans-gene

regulation across tissues and diseases. *Science* **353,** 827–830 (2016).

50.    Index Switching Causes "Spreading-Of-Signal" Among Multiplexed Samples In

Illumina HiSeq 4000 DNA Sequencing | bioRxiv. Available at:

http://biorxiv.org/content/early/2017/04/09/125724. (Accessed: 19th May 2017)

51.    Servant, N. *et al.* HiC-Pro: an optimized and flexible pipeline for Hi-C data

processing. *Genome Biol.* **16,** 259 (2015).

52.    Gjoneska, E. *et al.* Conserved epigenomic signals in mice and humans reveal immune

basis of Alzheimer's disease. *Nature* **518,** 365–369 (2015).

53.    Kagey, M. H. *et al.* Mediator and cohesin connect gene expression and chromatin

architecture. *Nature* **467,** 430–435 (2010).

54.    Yue, F. *et al.* A comparative encyclopedia of DNA elements in the mouse genome.

*Nature* **515,** 355–364 (2014).

55.    Bonnal, R. J. P. *et al.* De novo transcriptome profiling of highly purified human

lymphocytes primary cells. *Sci. Data* **2,** 150051 (2015).

56.    Ay, F., Bailey, T. L. & Noble, W. S. Statistical confidence estimation for Hi-C data

reveals regulatory chromatin contacts. *Genome Res.* gr.160374.113 (2014).

doi:10.1101/gr.160374.113

57.    Adamson, B. *et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables

Systematic Dissection of the Unfolded Protein Response. *Cell* **167,** 1867–1882.e21

(2016).

58.    Gilbert, L. A. *et al.* Genome-Scale CRISPR-Mediated Control of Gene Repression and

Activation. *Cell* **159,** 647–661 (2014).

59.    Doench, J. G. *et al.* Optimized sgRNA design to maximize activity and minimize off-

target effects of CRISPR-Cas9. *Nat. Biotechnol.* **34,** 184–191 (2016).

60.    Hsu, P. D. *et al.* DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat.*

*Biotechnol.* **31,** 827–832 (2013).

61.    Horlbeck, M. A. *et al.* Nucleosomes impede Cas9 access to DNA in vivo and in vitro.

*eLife* **5,** (2016).

1331
1332