Microtubule dynamics regulates mitochondrial fission

Kritika Mehta[1], Manjyot Kaur Chug[1], Siddharth Jhunjhunwala[1] and Vaishnavi Ananthanarayanan[1,*]

[1] *Centre for BioSystems Science and Engineering, Indian Institute of Science, Bangalore 560012*

* To whom correspondence may be addressed. Email: vaishnavi@iisc.ac.in

Abstract

Mitochondria are organised as tubular networks in the cell and undergo fission and fusion. Equilibrium between fission and fusion events is necessary to maintain normal mitochondrial function. While several of the molecular players involved in mediating the dynamics of mitochondria have been identified, the precise cellular cues that initiate fission or fusion remain largely unknown. In fission yeast, as in mammalian cells, mitochondrial positioning and partitioning are microtubule-mediated. In interphase, fission yeast mitochondria associate with microtubule bundles that are aligned along the long axis of the cell. Here, we perturbed microtubule dynamics via kinesin-like proteins to produce cells with long and short microtubules and observed that cells with long microtubules exhibited long, but fewer mitochondria, whereas cells with short microtubules exhibited short, but several mitochondria. We quantified that mitochondrial fission frequency correlated inversely with microtubule length. In agreement with these results, upon onset of mitosis wherein interphase microtubules assemble to form the spindle within the nucleus, we measured increased fission of mitochondria in the absence of interphase microtubules. We confirmed that partitioning of mitochondria at mitosis follows a stochastic, independent segregation mechanism and that the increase in mitochondrial fission prior to cell division therefore likely serves to increase the likelihood of equal partitioning of mitochondria between the two future daughter cells. Further, association of mitochondria with microtubules inhibited activity of the dynamin GTPase-related fission protein Dnm1, such that cells overexpressing Dnm1 but containing long microtubules were protected from increased fission. Thus, we demonstrate a general mechanism by which mitochondrial dynamics is dictated by the dynamics of the microtubule cytoskeleton.

Introduction

Mitochondria are double-membraned organelles whose functions range from ATP production to calcium signalling. Inside cells, mitochondrial form is dynamic and transitions from tubular networks to fragmented entities depending on the activity of the mitochondrial fission and fusion machinery. The major mitochondrial fission protein is dynamin-related Drp1 GTPase[1] (Dnm1 in yeast[2,3]). Multimeric Drp1 rings assemble around the mitochondrial membrane and utilize the energy from GTP hydrolysis to catalyse the constriction and fission of mitochondria[4,5]. Fusion of mitochondria requires two sets of proteins namely Opa1[6] for the inner membrane (Mgm1 in yeast[7]) and Mfn1/2[8] for the outer membrane (Fzo1 in yeast[9]).

The requirement for dynamic mitochondria has been attributed to two primary reasons, namely quality control and energy production[10]. Larger/longer mitochondria resulting from fusion are hypothesized to be capable of producing more energy whereas shorter/smaller mitochondria formed following a fission event are likely to undergo mitophagy[11]. In the case of the latter, fission could serve as an efficient mechanism to segregate and eliminate damaged mitochondria. Dysfunction of fission and fusion processes has been implicated in neurodegeneration[12,13], cancer[14] and cardiomyopathies[15], amongst a host of metabolic disorders.

In mammalian cells, mitochondria are transported along microtubule tracks by the activity of motor proteins kinesin-1 and dynein[16]. Kinesin-1 and dynein bind to the outer membrane of mitochondria via the Miro/Milton complex[17–19] and move mitochondria in the anterograde and retrograde direction respectively. In neuronal cells, increase in calcium levels results in the attachment of kinesin-1 motor to mitochondria via syntaphilin, which inhibits the ATPase activity of kinesin and hence leads to stationary mitochondria on neuronal microtubules[20]. About 70% of mitochondria in neuronal cells have been visualized in this stationary state[21].

In contrast to mammalian cells, mitochondria in fission yeast do not undergo motor-driven movement along microtubules. However, the protein Mmb1 has been identified to associate mitochondria with dynamic microtubules[22]. Upon microtubule depolymerisation using methyl benzimidazol-2-yl-carbamate (MBC) or deletion of Mmb1, mitochondria have been observed to undergo fragmentation[3,22,23]. Conversely, in cells overexpressing Mmb1, mitochondria seemingly undergo reduced fission and exhibit long mitochondria[22].

Cells employ several strategies to reduce partitioning error of organelles during

mitosis, such as ordered segregation mediated by spindle poles, or increasing copy

numbers of organelles prior to cell division[24]. In the latter, homogeneous distribution of the

multiple organelle copies serves to increase the probability of equal partitioning

stochastically. Mitochondrial inheritance has been observed to be microtubule-dependent

in mammalian cells[25]. In fission yeast, mitochondrial partitioning during cell division has

been proposed to be mediated by attachment of mitochondria with spindle poles[26,27], similar

to the segregation of endosomes, lysosomes and Golgi bodies in mammalian cells[28,29] .

However, only a portion of the observed mitochondria associated with the spindle poles[26].

Additionally, increased mitochondrial fragmentation upon the onset of mitosis has also been

observed[3], perhaps indicating a stochastic mechanism for mitochondrial partitioning.

While the molecular players that effect fission and fusion have been identified in

several systems, the cellular signals that regulate these events are largely elusive. Here,

we demonstrate that mitochondria piggyback on dynamic microtubules to selectively

undergo fission while microtubules depolymerise. Reorganisation of interphase

microtubules into the nucleus when cells prepare for division also provided the cue for

increased mitochondrial fission. We quantified the number of mitochondria in mother cells

immediately after formation of mitotic spindle within the nucleus and the number of

mitochondria in the resulting daughter cells, and confirmed that the partitioning was indeed

a good fit to a binomial distribution, indicating that the mitochondria were stochastically

segregated into the two daughter cells. We observed that the fission protein Dnm1 was

excluded from regions of microtubules localisation and that the presence of long and

stabilised microtubules was inhibitory to uncontrolled fission even when Dnm1 was

overexpressed.

Results

1. Perturbation of microtubule dynamics leads to changes in mitochondrial numbers

We observed that mitochondria underwent increased fission in the absence of microtubules, but did not observe their subsequent aggregation as reported previously (Extended Data Fig. 1a-c). Instead, the fragmented mitochondria were mobile and frequently in close contact with each other (Extended Data Fig. 1b, c, Supplementary Video S1). Since the depolymerisation of microtubules had a direct effect on mitochondrial fission, we set out to study the consequence of modification of microtubule dynamics on mitochondrial dynamics. To this end, we visualised the mitochondria and microtubules of fission yeast strains carrying deletions of antagonistic kinesin-like proteins, Klp5/Klp6 and Klp4 in high-resolution deconvolved images (Fig. 1a, Supplementary Videos S2, S3 and S4).

Figure 1. Mitochondrial number is inversely proportional to microtubule length. a, Maximum intensity projections of deconvolved Z-stack images of microtubules (left), mitochondria (centre) and their composite (right) in wild-type ('WT', top, strain KI001, see

Table S1), Klp5Δ/Klp6Δ ('MT$_{long}$', strain G3B, see Table S1) and Klp4Δ ('MT$_{short}$', strain
G5B, see Table S1) cells. **b,** Box plot of length of anti-parallel microtubule bundles in WT,
MT$_{long}$ and MT$_{short}$ cells (n=40, 37 and 63 bundles respectively). **c,** Box plot of number of
mitochondria per cell in WT, MT$_{long}$ and MT$_{short}$ cells (n=10, 12 and 13 cells respectively). **d,**
Box plot of the total volume of mitochondria per cell in WT, MT$_{long}$ and MT$_{short}$ cells
normalised to mean total wild-type mitochondrial volume (n=10, 12 and 13 cells
respectively). Light grey crosses represent outliers, asterisk represents significance
($p<0.05$) and 'n.s.' indicates no significant difference.
111

The heteromeric Klp5/Klp6 motor is required for maintenance of interphase
microtubule length by promoting catastrophe at microtubule plus ends[30,31]. Cells lacking
Klp5 and Klp6 exhibited long microtubule bundles ('MT$_{long}$', Fig. 1b) as reported previously
due to a decreased catastrophe rate[30]. In contrast, Klp4 is required for polarised growth in
fission yeast and has been suggested to promote microtubule growth[32,33]. As a result, in
the absence of Klp4, microtubule bundles were only about half the length of wild-type
bundles ('MT$_{short}$', Fig. 1b).

As in wild-type cells, mitochondria in Klp5Δ/Klp6Δ were in close contact with the
microtubule, whereas we observed reduced association between the short microtubules
and mitochondria in Klp4Δ cells (Fig. 1a, Supplementary Videos S3 and S4). While wild-
type cells had 4.9 ± 0.4 (mean ± s.e.m.) mitochondria per cell, we observed that
Klp5Δ/Klp6Δ contained only 2.3 ± 0.4 (mean ± s.e.m.). In contrast, Klp4Δ cells had 10 ± 0.9
mitochondria per cell (mean ± s.e.m., Fig. 1c). This indicated that the number of
mitochondria per cell was inversely related to the length of microtubule bundle. So too, cells
carrying a deletion of the EB1 homologue Mal3 exhibited small microtubule bundles[34] and
also contained several mitochondria as reported previously[22] (Extended Data Fig. 1d).
However, the decrease in the number of mitochondria in Klp5Δ/Klp6Δ cells and increase in
Klp4Δ cells were not at the expense of mitochondrial volume, since the net mitochondrial
volume in both cases was comparable to wild-type mitochondrial volume (Fig. 1d, Extended
Data Fig. 1e).
132

2. Cells with short microtubules undergo increased fission
To understand the difference in mitochondrial numbers in wild-type, Klp5Δ/Klp6Δ and
Klp4Δ cells, we acquired and analysed time-lapse videos at the single mitochondrion level
in all three cases (Fig. 2a, Supplementary Videos S5, S6 and S7). Similar to our
observations from high-resolution images, we measured 3.6 ± 0.2, 2.7 ± 0.2, and 6.9 ± 0.8

mitochondria (mean ± s.e.m.) in wild-type, Klp5Δ/Klp6Δ and Klp4Δ cells respectively (Fig.
1c and 2b). Analysis of evolution of these mitochondrial numbers revealed no significant
changes over time (Fig. 2b).

Figure 2. Microtubule catastrophe rate determines fission frequency of
mitochondria. a, Montage of maximum intensity projected confocal Z-stack images of wild-
type ('WT', strain Kl001, see Table S1), Klp5Δ/Klp6Δ ('MT$_{long}$', strain G3B, see Table S1)
and Klp4Δ ('MT$_{short}$', strain G5B, see Table S1) cells represented in the intensity map
indicated to the right of the images. White solid arrowheads point to representative fusion
events and open arrowheads to fission events. Time is indicated above the images in

mm:ss. **b,** Evolution of mitochondrial number over time indicated as mean (solid grey line)
and standard error of the mean (shaded region) for WT, MT$_{long}$ and MT$_{short}$ cells (n=21, 15
and 8 cells respectively **c,** Box plot of the fission frequency of mitochondria per second in
WT, MT$_{long}$ and MT$_{short}$ cells (n=21, 15 and 8 cells respectively). **d,** Box plot of the fusion
frequency of mitochondria per second in WT, MT$_{long}$ and MT$_{short}$ cells (n=21, 15 and 8 cells
respectively). Light grey crosses represent outliers, asterisk represents significance
($p<0.05$) and 'n.s.' indicates no significant difference. **e,** Plot of total microtubule length
normalised to total microtubule length in wild-type cells (grey) and ratio of mean fission
frequency to mean fusion frequency (blue) in WT, MT$_{long}$ and MT$_{short}$ cells.

Additionally, mitochondria in wild-type cells underwent ~1 fission and ~1 fusion event

every minute on an average, whereas Klp5Δ/Klp6Δ cells exhibited a fission frequency that

was half that of wild-type, and Klp4Δ mitochondria had a fission frequency that was almost

double that of wild-type (Fig. 2c). The fusion frequency of mitochondria in Klp4Δ cells was

slightly higher than in wild-type and Klp5Δ/Klp6Δ cells (Fig. 2d), likely due to the increased

number of mitochondria in Klp4Δ cells that could participate in fusion. However, the

resulting ratio of the mean fission frequency to the mean fusion frequency was ~1, ~0.5 and

~1.3 in wild-type, Klp5Δ/Klp6Δ and Klp4Δ cells respectively. Interestingly, these ratios

accurately matched the total microtubule lengths of wild-type, Klp5Δ/Klp6Δ and Klp4Δ cells

(19μm, 25μm and 15μm on an average) normalised to total WT microtubule length (1, 0.76

and 1.26 respectively, Fig. 2e). We therefore concluded that the difference in mitochondrial

numbers between wild-type cells and Klp5Δ/Klp6Δ and Klp4Δ arose primarily due to the

changes in fission frequencies of mitochondria. Note that the rate of mitochondrial fission

(per second per mitochondrion) was comparable to the rate of mitochondrial fusion (per

second per mitochondrion) in all three cases, which explains the constant number of

mitochondria we measured over time (Fig. 2b).

To specifically test the role of microtubules in determining the length and dynamics

of mitochondria, we depolymerised microtubules in wild-type and Klp5Δ/Klp6Δ cells and

visualised the mitochondria in time-lapse movies (Extended Data Fig. 2a). In both cases,

upon microtubule depolymerisation, we observed a switch in mitochondrial morphology to

increased numbers and increased fission that was reminiscent of Klp4Δ cells (Extended

Data Fig. 2b-h).

Increase in oxidative stress via reactive oxygen species (ROS) levels has also been

described to induce mitochondrial fission[35]. However, we measured no difference in ROS

levels between wild-type, Klp5Δ/Klp6Δ, and Klp4Δ cells (Extended Data Fig. 2i).

Figure 3. Mitochondrial sizes are predicted by partitioning the same amount of mitochondrial material among different numbers of mitochondria. **a,** Box plot of the size of mitochondria in wild-type ('WT', strain KI001, see Table S1), Klp5Δ/Klp6Δ ('MT$_{long}$', strain G3B, see Table S1) and Klp4Δ ('MT$_{short}$', strain G5B, see Table S1) cells, calculated as the length of the major axis of an ellipse fitted to each mitochondrion (n=1613, 739 and 1326 mitochondria respectively). Light grey crosses represent outliers, asterisk represents significance (p<0.05) and 'n.s.' indicates no significant difference. **b-d,** Histogram of mitochondrial sizes normalised to total size of mitochondria per cell obtained from experiments in WT, MT$_{long}$ and MT$_{short}$ cells. **e-g,** Histogram of mitochondrial sizes

194 normalised to total size of mitochondria per cell obtained from simulations using numbers of
195 mitochondria measured in WT, MT_{long} and MT_{short} cells.
196

3. Mitochondrial sizes reflect microtubule lengths

Additionally, we observed that the size of an individual mitochondrion correlated with the
length of the microtubule bundle in the cell, with Klp5Δ/Klp6Δ cells containing long
mitochondria and Klp4Δ cells having short mitochondria. Wild-type cells predictably
exhibited mitochondrial size ranges that lay between that of Klp5Δ/Klp6Δ and Klp4Δ cells
(Fig. 3a, Extended Data Fig. 3a-c). As detailed in Fig. 1d, the net mitochondrial volume was
identical in wild-type, Klp5Δ/Klp6Δ and Klp4Δ cells. We tested the possibility that the
mitochondrial size distributions we observed in wild-type, Klp5Δ/Klp6Δ and Klp4Δ cells
(Fig. 3b-d) could simply be a consequence of partitioning the same amount of mitochondria
into ~4, ~2 and ~7 mitochondria respectively. Fig. 3e-g depicts the results of simulating the
possible size distributions that mitochondria could exhibit relative to the total mitochondrial
size (see Methods).

We observed similar mitochondrial size distributions in both our experiments and
simulations reconfirming our earlier observation that the total mitochondrial content remains
unchanged between wild-type, Klp5Δ/Klp6Δ and Klp4Δ cells.

4. Mitochondria undergo stochastic partitioning during mitosis

Fission yeast undergoes closed mitosis, wherein the nuclear envelope does not undergo
breakdown during cell division[36]. Upon onset of mitosis in fission yeast, the interphase
microtubules that were previously in the cytoplasm are reorganised to form the spindle
inside the closed nucleus. This natural situation mimics the depolymerisation of
microtubules via the chemical inhibitor MBC. Therefore, we set out to study the changes in
the mitochondrial network upon cell entry into mitosis. We first obtained high-resolution
deconvolved images of the microtubule and mitochondria in fission yeast cells undergoing
cell division (Fig. 4a, Extended Data Fig. 4a, Supplementary Video S8). We observed that
dividing wild-type cells had ~4x the number of mitochondria in interphase cells (Fig. 4b).
Moreover, similar to what was seen in cells lacking microtubules or Klp4 (Fig. 1a),
mitochondria in dividing cells were shorter and more rounded (Fig. 4a, Extended Data Fig.
4a). There was no relationship between length of the mitotic spindle and the number of
mitochondria (Fig. 4b, Extended Data Fig. 4a), indicating that the increased fission likely

occurred fairly early upon entry into mitosis. Analysis of time-lapse videos of wild-type cells
before and 10min after entry into mitosis revealed a doubling of mitochondrial numbers in
this time (Fig. 4c, d, Supplementary Video S9). Note that this is likely an underestimate
since we were unable to reliably resolve some of the mitochondria in these lower-resolution
images. The fragmented mitochondria appeared more mobile and were able to traverse
distances of ~1μm in the cell presumably by Brownian motion (Fig. 4c). In this same period
of time, non-dividing interphase cells did not show any change in mitochondrial numbers
(Extended Data Fig. 4b, c).

Figure 4. Mitotic cells contain several short mitochondria. a, Maximum intensity
projections of deconvolved Z-stack images of the microtubules (top), mitochondria (centre)
and their composite (bottom) of a wild-type cell (strain KI001, see Table S1) undergoing
division. **b,** Scatter plot of the length of the mitotic spindle vs. the number of mitochondria
per cell in dividing cells (*n*=13 cells). **c,** Montage of maximum intensity projected confocal Z-
stack images of the microtubules (top) and mitochondria (bottom) in a wild-type cell (strain
KI001, see Table S1) undergoing cell division represented in the intensity map indicated to
the right of the images. White open arrowheads point to representative fission events.
Magenta arrowheads point to a representative mobile, fragmented mitochondrion. Time is
indicated above the images in mm:ss. **d,** Bar plot of the mean number of mitochondria per

cell before ('00:00') and 10mins after ('10:00') the onset of mitosis. Solid grey lines
represent data from individual cells (*n*=16 cells).

Increase in mitochondrial numbers prior to cell division could aid in increasing the

likelihood of equal partitioning of mitochondria between daughter cells, given a stochastic

mechanism. To test if mitochondria in our system underwent stochastic, independent

segregation,[24] with each mitochondrion in the mother cell having a 50% chance of

segregating to either of the future daughter cells during mitosis, we tested the fit of our data

to a binomial distribution[37] using a Chi-square test as previously described[38]. Our data

(Table S2) did not differ significantly from a binomial distribution with a Chi-square statistic

of 7.1846 with 3 degrees of freedom and p=0.0662, indicating that mitochondria in fission

yeast were indeed segregating stochastically during cell division. The increase in

mitochondrial numbers upon onset of mitosis also served to reduce the partitioning error of

mitochondria between the daughter cells as predicted by stochastic segregation (Extended

Data Fig. 4d).

5. Dnm1 rings do not localize to the microtubules

Taken together, our results suggest that association of mitochondria with microtubules

inhibits mitochondrial fission. The mitochondrial fission protein in yeast is a dynamin-related

GTPase, Dnm1[3]. Dnm1 brings about the fission of mitochondria by self-assembling into a

ring around the diameter of the mitochondria and employing its GTPase activity to effect the

scission[4]. In the absence of Dnm1, mitochondria are organised as extended, fused

'nets'[3,39]. We hypothesised that the binding of mitochondria to microtubules physically

hinders the formation of a complete ring of Dnm1 around the mitochondrion. To test this

hypothesis, we introduced GFP-tagged Dnm1 under the control of the thiamine-repressible

nmt1 promoter[40] into wild-type cells expressing fluorescent microtubules (see Table S1). C-

or N-terminal tagging of Dnm1 renders it inactive[3] and therefore the scission activity of

Dnm1 could not be directly observed (Extended Data Fig. 5a). However, we visualised the

localisation of Dnm1 'spots', representing Dnm1 assemblies. We counted a slight decrease

in Dnm1 spots in Klp5Δ/Klp6Δ, and ~2x increase in Klp4Δ cells as compared to wild-type

cells (Extended Data Fig. 5b), corroborating our previous observations indicating decreased

fission in the former and increased fission in the latter (Fig. 2c). Although almost 100% of

the Dnm1 spots colocalised with mitochondria (Extended Data Fig. 5a), 85% of the spots

did not localise on the microtubule (Fig. 5a, Supplementary Video S10), indicating that the
presence of microtubule had an inhibitory effect on the assembly of the Dnm1 ring.

Figure 5. Association with microtubules prevents uncontrolled fission of mitochondria. a, Maximum intensity projections of deconvolved Z-stack images of Dnm1 (left), microtubules (centre) and their composite (right) in wild-type cells expressing fluorescent Dnm1 and tubulin (strain MTY271 transformed with pREP41-Dnm1-Cterm-GFP, see Table S1). A majority of Dnm1 spots did not localise on the microtubule (n=14 cells). **b,** Maximum intensity projections of deconvolved Z-stack images of mitochondria in cells transformed with functional untagged Dnm1 (plasmid pREP41-Dnm1, see Table S1) in wild-type ('WT', strain FY7143, see Table S1) and Klp5Δ/Klp6Δ cells ('MT_{long}', strain FY20823, see Table S1) represented in the intensity map indicated to the bottom of the images. High overpexpression of Dnm1 is indicated with '+++' ('WT(-thia)', and 'MT_{long}(-thia)') and low overexpression with '++' ('WT(+thia)', and 'MT_{long}(+thia)'). **c,** Box plot of mitochondrial numbers in WT and MT_{long} cells with high overexpression ('+++') or low expression of Dnm1 ('++') (n= 149, 37, 63, 41 cells respectively). The mitochondrial numbers in WT and MT_{long} cells expressing normal amount of Dnm1 ('+') are depicted by the dashed lines. Light grey crosses represent outliers, asterisk represents significance (p<0.05). **d,** Model of mitochondrial dynamics mediated by microtubule dynamics. Microtubules polymerise and depolymerise at their plus ends ('+'). Absence of microtubule bundles in the cytoplasm

298 during cell division enables the fragmentation of mitochondria. **e,** When mitochondria are
299 bound to microtubules, Dnm1 ring assembly might be inhibited. Upon microtubule
300 depolymerisation, this inhibition is alleviated and Dnm1 can effectively mediate scission of
301 mitochondria.
302
To test the influence of microtubule dynamics and in essence, microtubule lengths
on the fission-fusion status of mitochondria, we transformed wild-type and Klp5Δ/Klp6Δ
cells with a plasmid expressing untagged Dnm1 under the control of the nmt1 promoter
(see Table S1) and visualized the mitochondria (Fig. 5b). In these cells, Dnm1 shows high
overexpression in the absence of thiamine and low overexpression in the presence of
0.05µM thiamine in the culture medium. We counted the number of mitochondria present in
these cells and estimated that wild-type cells highly overexpressing Dnm1 exhibited 11.6 ±
0.2 mitochondria (mean ± s.e.m), which is twice that of wild-type cells expressing normal
levels of Dnm1 (Fig. 5c). The increase in mitochondrial numbers in these cells is due to the
increase in Dnm1 numbers, that likely accelerates the kinetics of the Dnm1 ring formation
and hence, mitochondrial fission. In wild-type cells that were grown in the presence of
thiamine, the mitochondrial number was 7 ± 0.3 (mean ± s.e.m.), presumably because of
low overexpression of Dnm1. Interestingly, in Klp5Δ/Klp6Δ cells with high and low
overexpression of Dnm1, we counted only 7.5 ± 0.3 mitochondria and 5.2 ± 0.2
mitochondria (mean ± s.e.m.) respectively. In Klp5Δ/Klp6Δ, the presence of longer
microtubules than wild-type cells possibly prevented increased fission of mitochondria even
when cells overexpressed Dnm1. In fact, in Klp5Δ/Klp6Δ cells where there was low
overexpression of Dnm1, the mitochondrial number was comparable to that in wild-type
cells expressing normal levels of Dnm1 (Fig. 5c).
Extended Data Fig. 5c contains a summary of the mitochondrial parameters
measured in this study.
324
Discussion
We discovered that microtubule dynamics is a strong determinant for mitochondrial
dynamics and thereby, morphology. Mitochondrial fission frequency was proportional to
microtubule catastrophe rate. While previous studies discounted the role of motors in the
determination of mitochondrial positioning in fission yeast[23,26,41,42], we have identified
kinesin-like proteins that regulate mitochondrial morphology through their control over
microtubule length. The distribution of mitochondrial sizes that we observed in wild-type

cells and cells with long and short microtubules could be explained by the allocation of the

same amount of mitochondria between different numbers of mitochondria. We observed the

maximum number of mitochondria, and thereby the smallest mitochondrial sizes, in cells

with short microtubules.

We propose that equilibrium between fission and fusion of mitochondria in

interphase cells can be maintained solely by the inherent dynamic instability of

microtubules. Perturbation of microtubule dynamics shifts the balance between fission and

fusion frequencies to a different set point, leading to modification of mitochondrial numbers

and morphology. In mammalian cells, mitochondria employ microtubule-associated

machinery to traverse the cell. In HeLa cells, Drp1 recruitment to the mitochondria was

observed to be dependent on dynein[43]. Mobile mitochondria on microtubules have also

been seen to participate in kiss-and-run fusions that are transient[44]. Additionally, the role of

microtubule-stabilizing and destabilizing drugs has been tested on the biogenesis of

mitochondria[45]. In taxol-treated cells, mitochondria appeared more elongated whereas in

nocodazole-treated cells, mitochondria seemed fragmented[45]. However, a direct role for

microtubule localization on mitochondrial fission has not been explored.

We observed that fission yeast cells fragment mitochondria by emptying the

cytoplasm of its microtubules upon onset of mitosis, thereby ensuring stochastic

partitioning[37] of the several small mitochondria between future daughter cells (Fig. 5d). The

increase in mitochondrial numbers prior to cell division likely serves to create a well-mixed,

homogeneous cytoplasm that is primed for stochastic segregation of mitochondria.

Mitochondrial fragmentation also reduces the partitioning error between the daughter

cells[38], since mitochondrial numbers of 5 or less, which is typical of wild-type interphase

cells, exhibit errors between ~45 and 100% (Extended Data Fig. 4d). While fission yeast

cells employ this strategy, mammalian cells might additionally employ motor proteins to

actively segregate mitochondria during cell division.

We observed that Dnm1 spots were excluded from microtubules. We propose the

failure of Dnm1 ring assembly in the presence of microtubules as the nexus between

microtubule dynamics and mitochondrial dynamics (Fig. 5e). Previously, cryo-electon

tomographic analysis in fission yeast indicated a preferred separation distance of ~20nm for

mitochondria associated with microtubules[46]. So too, cryo-elctron microscopy in budding

yeast revealed that Dnm1 assembled into rings with an outer diameter of 129nm and lumen

diameter of 89nm, resulting in ~40nm-high structures around mitochondria[47]. Therefore,
mitochondria associated with microtubules might have insufficient space to accommodate
the Dnm1 ring around their diameter, thereby inhibiting fission due to simple physical
constraints.

Alternatively, increased fission upon microtubule depolymerization might be brought
about by phosphorylation of the mitochondrial fission protein Dnm1. Mammalian Drp1
undergoes phosphorylation at S616 mediated by Cdk1/cyclin B during cell division and
increases mitochondrial fission[48]. Fission yeast Dnm1 however lacks this phosphorylation
site. Drp1 also contains another phosphorylation site at S637, which is conserved in Dnm1.
However, PKA-mediated phosphorylation at S637 has been shown to inhibit fission by
Drp1[49], whereas Ca+/calmodulin-dependent protein kinase α-dependent phosphorylation of
Drp1 S637 in rat hippocampal neurons was shown to increase fission[50]. To date, yeast
Dnm1 has not been demonstrated to undergo phosphorylation at this site. Future studies
will help delineate whether the increased fission upon microtubule depolymerization is due
to a simple physical impediment to the assembly of Dnm1 ring or a more complex
biochemical signal tethered to microtubules that changes the phosphorylation status of
Dnm1 to activate it.

We also observed that the presence of long microtubules was inhibitory for fission of
mitochondria even in the overexpression of the fission protein Dnm1. Wild-type cells
contained the same total amount of mitochondria as cells with long microtubules, but
exhibited increased mitochondrial fission in the presence of increased Dnm1 levels. In
disease states such as cancer[51] and neurodegeneration[52], mitochondrial fragmentation has
been linked with overexpression of Drp1 in mammalian cells. In this work, we have
identified that cells that present increased microtubule stabilization are able to overcome
the effect of Dnm1 overexpression.

Destabilisation of microtubules is a hallmark of several neurodegenerative disorders
including Alzheimer's, Parkinson's and Huntington's disease[53]. So too, colorectal cancer
has been characterised by the absence of the tumour suppressor gene APC, which also
modulates microtubule dynamics[54]. Remarkably, all these disease states have also been
correlated with dysfunction of mitochondrial dynamics[11,55]. In future, it would be interesting
to investigate the effect of restoration of microtubule dynamics on the mitochondrial fission
and fusion in these scenarios, and consequently on the disease states.

Methods

Strains and media. The fission yeast strains used in this study are listed in Table S1. All the strains were grown in YE (yeast extract) media or Edinburg Minimal media (EMM)[56] with appropriate supplements at a temperature of 30˚C. Cells that were transformed with pREP41-Dnm1-Cterm-GFP or pREP41-Dnm1 (see below) were cultured in EMM with appropriate supplements and $0.05\mu M$ thiamine for partial induction of the nmt1 promoter.

Plasmid transformation. Transformation of strains MTY271, FY7143, FY20823 and Mcl438 with plasmid pREP41-Dnm1-Cterm-GFP and pREP41-Dnm1 (see Table S1) was carried out using the improved protocol for rapid transformation of fission yeast[57]. In brief, cells were grown overnight to log phase in low glucose EMM, pelleted and washed with distilled water. The cells were then washed in a solution of lithium acetate/EDTA (100mM LiAc, 1mM EDTA, pH 4.9) and re-suspended in the same solution. $1\mu g$ of plasmid DNA was added to the suspension, followed by addition of lithium acetate/PEG (40% w/v PEG, 100mM LiAc, 1mM EDTA, pH 4.9) and then incubated at 30°C for 30 min. This was followed by a heat shock of 15min at 42°C. Thereafter, cells were pelleted down, re-suspended in TE solution (10mM Tris-HCl, 1mM EDTA, pH 7.5) and plated onto selective EMM plates.

Preparation of yeast for imaging. For imaging mitochondria, fission yeast cells were grown overnight in a shaking incubator at 30°C, washed once with distilled water, and stained with 200nM Mitotracker Orange CMTMRos (ThermoFisher Scientific, Cat. #M7510) dissolved in EMM for 20min. Following this, cells were washed thrice with EMM and then allowed to adhere on lectin-coated (Sigma-Aldrich, St. Louis, MO, Cat. #L2380) 35mm confocal dishes (SPL, Cat. #100350) for 20min. Unattached cells were then removed by washing with EMM. In experiments where mitochondria were not imaged, staining with Mitotracker was omitted.

Microtubule depolymerisation. For depolymerisation of microtubules, cells were treated with methyl benzimidazol-2-yl-carbamate (MBC, Carbendazim 97%, Sigma Aldrich). A stock solution with a concentration of 25mg/ml was prepared in DMSO and later diluted to a working concentration of $25\mu g/ml$ in EMM.

Microscopy. Confocal microscopy was carried out using the InCell Analyzer-6000 (GE Healthcare, Buckinghamshire, UK) with 60x/0.7 N.A. objective fitted with an sCMOS 5.5MP camera having an x-y pixel separation of 108nm. For GFP and Mitotracker Orange imaging, 488 and 561nm laser lines and bandpass emission filters 525/20 and 605/52nm

respectively were employed. Time-lapses for visualisation of mitochondrial dynamics were

captured by obtaining 5 Z-stacks with a 0.5μm-step size every 12s. Deconvolution was

performed in images obtained using a Deltavision RT microscope (Applied Precision) with a

100×, oil-immersion 1.4 N.A. objective (Olympus, Japan). Excitation of fluorophores was

achieved using InsightSSI (Applied Precision) and corresponding filter selection for

excitation and emission of GFP and Mitotracker Orange. Z-stacks with 0.3μm-step sizes

encompassing the entire cell were captured using a CoolSnapHQ camera (Photometrics),

with 2X2 binning. The system was controlled using softWoRx 3.5.1 software (Applied

Precision) and the deconvolved images obtained using the built-in setting for each channel.

3D visualisation of deconvolved images. 3D views of the microtubules and mitochondria

in Supplementary Movies S2, S3, S4, S8, S9 and S10 were obtained from deconvolved

images captured in the Deltavision microscope using Fiji's '3D project' function, with the

brightest point projection method and 360˚ total rotation with 10˚ rotation angle increment.

Estimation of volume of mitochondria. Mitochondrial volume was estimated in Fiji by

integrating the areas of mitochondria in thresholded 3D stacks of cells in fluorescent

deconvolved images obtained using the Deltavision RT microscope. The total volume was

then normalised to the mean total mitochondrial volume of wild-type cells. Individual

mitochondrial volumes were estimated in the same fashion.

Analysis of mitochondrial dynamics. Individual mitochondria were identified in each

frame of the time-lapse obtained in confocal mode of the GE InCell Analyzer after projecting

the maximum intensity of the 3D stack encompassing the cell, followed by mean filtering

and visualisation in Fiji's 'mpl-magma' lookup table. Following identification of mitochondria,

the 'Measure' function of Fiji was employed to obtain the circularity, aspect ratio and

parameters of fitted ellipse. The length of the major axis of the ellipse fitted to a

mitochondrion was defined as the size of that mitochondrion. The size, circularity and

aspect ratio were estimated for mitochondria at each frame and each time point. Fission

and fusion frequencies of mitochondria were estimated by counting the number of

mitochondria identified during each frame of the time-lapse. The difference in number of

mitochondria from one frame to the next was counted, with an increase being counted as

fission event and decrease as fusion event. The total number of fission events and fusion

events per cell were estimated and divided by the total duration of the time-lapse to obtain

the fission and fusion frequencies.

Simulation of mitochondrial size distribution. Mitochondrial size distributions in wild-type, Klp5Δ/Klp6Δ and Klp4Δ cells were simulated by taking experimentally measured mitochondrial numbers in the three cases and assuming that the total mitochondrial amount was the same. Briefly, to make the system amenable to simulation using an integer-partitioning problem, the total mitochondrial amount was taken to be a constant of 10. The minimum size of mitochondria that could be present was assumed to be 1. The total mitochondria was then partitioned between the available mitochondrial numbers in wild-type, Klp5Δ/Klp6Δ and Klp4Δ cells of ~4, ~2 and ~7 in units of 1. The distributions were then normalised to total mitochondrial amount to get mitochondrial sizes as fractions of the total mitochondrial content. The minimum mitochondrial size thus obtained in this case, 0.1, is similar to experimentally obtained normalised minimum mitochondrial sizes.

Test for fit of mitochondrial partitioning during mitosis to binomial distribution. To test the fit of mitochondrial segregation during mitosis to a binomial distribution, the data were z-transformed as previously described[38]. Briefly, given n mitochondria in the mother cell just prior to cell division, x and $n - x$ mitochondria in the resulting daughter cells, z was given by $2x - n/\sqrt{n}$ to approximate the binomial distribution to a normal distribution of 0,1. The z values obtained for each x and n were binned into k bins of equal sizes and subjected to Chi-square test with $k - 1$ degrees of freedom. The z values are expected to be equally distributed among the k bins, with expected number of $1/k$ per bin.

Data analysis and plotting. Data analysis was performed in Matlab (Mathworks, Natick, MA). Box plots with the central line indicating the median and notches that represent the 95% confidence interval of the median were obtained by performing one-way ANOVA ('anova1' in Matlab). Following this, significance (p<0.05) of the means was tested using the Tukey-Kramer post-hoc test. All the plots were generated using Matlab.

References

1.   Smirnova, E., Griparic, L., Shurland, D. L. & van der Bliek, A. M. Dynamin-related protein Drp1 is required for mitochondrial division in mammalian cells. *Mol. Biol. Cell* **12,** 2245–56 (2001).

2.   Bleazard, W. *et al.* The dynamin-related GTPase Dnm1 regulates mitochondrial fission in yeast. *Nat. Cell Biol.* **1,** 298–304 (1999).

3.   Jourdain, I., Gachet, Y. & Hyams, J. S. The dynamin related protein Dnm1 fragments mitochondria in a microtubule-dependent manner during the fission yeast cell cycle. *Cell Motil. Cytoskeleton* **66,** 509–523 (2009).

4.   Ingerman, E. *et al.* Dnm1 forms spirals that are structurally tailored to fit mitochondria.

J. Cell Biol. **170,** 1021–7 (2005).

5.   Basu, K. *et al.* Molecular mechanism of DRP1 assembly studied in vitro by cryo-
electron microscopy. *PLoS One* **12,** e0179397 (2017).

6.   Delettre, C. *et al.* Nuclear gene OPA1, encoding a mitochondrial dynamin-related
protein, is mutated in dominant optic atrophy. *Nat. Genet.* **26,** 207–210 (2000).

7.   Sesaki, H., Southard, S. M., Yaffe, M. P. & Jensen, R. E. Mgm1p, a dynamin-related
GTPase, is essential for fusion of the mitochondrial outer membrane. *Mol. Biol. Cell*
14, 2342–56 (2003).

8.   Eura, Y., Ishihara, N., Yokota, S. & Mihara, K. Two Mitofusin Proteins, Mammalian
Homologues of FZO, with Distinct Functions Are Both Required for Mitochondrial
Fusion. *J. Biochem.* **134,** 333–344 (2003).

9.   Hermann, G. J. *et al.* Mitochondrial fusion in yeast requires the transmembrane
GTPase Fzo1p. *J. Cell Biol.* **143,** 359–73 (1998).

10.   Mishra, P. & Chan, D. C. Metabolic regulation of mitochondrial dynamics. *J. Cell Biol.*
212, 379–87 (2016).

11.   Chen, H. & Chan, D. C. Mitochondrial dynamics--fusion, fission, movement, and
mitophagy--in neurodegenerative diseases. *Hum. Mol. Genet.* **18,** R169-76 (2009).

12.   Hirai, K. *et al.* Mitochondrial abnormalities in Alzheimer's disease. *J. Neurosci.* **21,**
3017–23 (2001).

13.   Deng, H., Dodson, M. W., Huang, H. & Guo, M. The Parkinson's disease genes pink1
and parkin promote mitochondrial fission and/or inhibit fusion in Drosophila. *Proc.*
Natl. Acad. Sci. **105,** 14503–14508 (2008).

14.   Graves, J. A. *et al.* Mitochondrial Structure, Function and Dynamics Are Temporally
Controlled by c-Myc. *PLoS One* **7,** e37699 (2012).

15.   Ashrafian, H. *et al.* A Mutation in the Mitochondrial Fission Gene Dnm1l Leads to
Cardiomyopathy. *PLoS Genet.* **6,** e1001000 (2010).

16.   Pilling, A. D., Horiuchi, D., Lively, C. M. & Saxton, W. M. Kinesin-1 and Dynein are the
primary motors for fast transport of mitochondria in Drosophila motor axons. *Mol. Biol.*
Cell **17,** 2057–68 (2006).

17.   Stowers, R. S., Megeath, L. J., Górska-Andrzejak, J., Meinertzhagen, I. A. &
Schwarz, T. L. Axonal transport of mitochondria to synapses depends on milton, a
novel Drosophila protein. *Neuron* **36,** 1063–77 (2002).

18.   Glater, E. E., Megeath, L. J., Stowers, R. S. & Schwarz, T. L. Axonal transport of
mitochondria requires milton to recruit kinesin heavy chain and is light chain
independent. *J. Cell Biol.* **173,** 545–57 (2006).

19.   van Spronsen, M. *et al.* TRAK/Milton Motor-Adaptor Proteins Steer Mitochondrial
Trafficking to Axons and Dendrites. *Neuron* **77,** 485–502 (2013).

20.   Chen, Y. & Sheng, Z.-H. Kinesin-1-syntaphilin coupling mediates activity-dependent
regulation of axonal mitochondrial transport. *J. Cell Biol.* **202,** 351–64 (2013).

21.   Kang, J.-S. *et al.* Docking of Axonal Mitochondria by Syntaphilin Controls Their
Mobility and Affects Short-Term Facilitation. *Cell* **132,** 137–148 (2008).

22.   Fu, C., Jain, D., Costa, J., Velve-Casquillas, G. & Tran, P. T. Mmb1p binds
mitochondria to dynamic microtubules. *Curr. Biol.* **21,** 1431–1439 (2011).

23.   Li, T., Zheng, F., Cheung, M., Wang, F. & Fu, C. Fission yeast mitochondria are
distributed by dynamic microtubules in a motor-independent manner. *Sci. Rep.* **5,**
11023 (2015).

24.   Huh, D. & Paulsson, J. Random partitioning of molecules at cell division. *Proc. Natl.*
Acad. Sci. U. S. A. **108,** 15004–9 (2011).

25. Lawrence, E. & Mandato, C. Mitochondrial inheritance is mediated by microtubules in
mammalian cell division. *Commun. Integr. Biol.* **6,** e27557 (2013).
26. Yaffe, M. P., Stuurman, N. & Vale, R. D. Mitochondrial positioning in fission yeast is
driven by association with dynamic microtubules and mitotic spindle poles. *PNAS*
100, 11424–11428 (2003).
27. Krüger, N. & Tolic, I. M. Association of mitochondria with spindle poles facilitates
spindle alignment. *Curr. Biol.* **18,** 646–647 (2008).
28. Bergeland, T., Widerberg, J., Bakke, O. & Nordeng, T. W. Mitotic partitioning of
endosomes and lysosomes. *Curr. Biol.* **11,** 644–651 (2001).
29. Shima, D. T., Cabrera-Poch, N., Pepperkok, R. & Warren, G. An ordered inheritance
strategy for the Golgi apparatus: visualization of mitotic disassembly reveals a role for
the mitotic spindle. *J. Cell Biol.* **141,** 955–66 (1998).
30. Tischer, C., Brunner, D. & Dogterom, M. Force- and kinesin-8-dependent effects in
the spatial regulation of fission yeast microtubule dynamics. *Mol. Syst. Biol.* **5,** 250
(2009).
31. West, R. R., Malmstrom, T., Troxell, C. L. & McIntosh, J. R. Two related kinesins,
klp5+ and klp6+, foster microtubule disassembly and are required for meiosis in
fission yeast. *Mol. Biol. Cell* **12,** 3919–32 (2001).
32. Browning, H. *et al.* Tea2p is a kinesin-like protein required to generate polarized
growth in fission yeast. *J. Cell Biol.* **151,** 15–28 (2000).
33. Busch, K. E., Hayles, J., Nurse, P. & Brunner, D. Tea2p Kinesin Is Involved in Spatial
Microtubule Organization by Transporting Tip1p on Microtubules. *Dev. Cell* **6,** 831–
843 (2004).
34. Beinhauer, J. D., Hagan, I. M., Hegemann, J. H. & Fleig, U. Mal3, the fission yeast
homologue of the human APC-interacting protein EB-1 is required for microtubule
integrity and the maintenance of cell form. *J. Cell Biol.* **139,** 717–28 (1997).
35. Pletjushkina, O. Y. *et al.* Effect of oxidative stress on dynamics of mitochondrial
reticulum. *Biochim. Biophys. Acta - Bioenerg.* **1757,** 518–524 (2006).
36. Ding, R., West, R. R., Morphew, D. M., Oakley, B. R. & McIntosh, J. R. The spindle
pole body of Schizosaccharomyces pombe enters and leaves the nuclear envelope
as the cell cycle proceeds. *Mol. Biol. Cell* **8,** 1461–79 (1997).
37. Birky, C. W. The partitioning of cytoplasmic organelles at cell division. *Int. Rev. Cytol.*
Suppl. **15,** 49–89 (1983).
38. Hennis, A. S. & Birky, C. W. Stochastic partitioning of chloroplasts at cell division in
the alga Olisthodiscus, and compensating control of chloroplast replication. *J. Cell*
Sci. **70,** 1–15 (1984).
39. Guillou, E., Bousquet, C., Daloyau, M., Emorine, L. J. & Belenguer, P. Msp1p is an
intermembrane space dynamin-related protein that mediates mitochondrial fusion in a
Dnm1p-dependent manner in S. pombe. *FEBS Lett.* **579,** 1109–16 (2005).
40. Maundrell, K. nmt1 of fission yeast. A highly transcribed gene completely repressed
by thiamine. *J. Biol. Chem.* **265,** 10857–64 (1990).
41. Brazer, S. W., Williams, H. P. & Chappell, T. G. A fission yeast kinesin affects Golgi
membrane recycling. 149–166 (2000).
42. Chiron, S., Bobkova, A., Zhou, H. & Yaffe, M. P. CLASP regulates mitochondrial
distribution in Schizosaccharomyces pombe. *J. Cell Biol.* **182,** 41–49 (2008).
43. Varadi, A. *et al.* Cytoplasmic dynein regulates the subcellular distribution of
mitochondria by controlling the recruitment of the fission factor dynamin-related
protein-1. *J. Cell Sci.* **117,** 4389–400 (2004).

44. Liu, X., Weaver, D., Shirihai, O. & Hajnóczky, G. Mitochondrial 'kiss-and-run':
interplay between mitochondrial motility and fusion-fission dynamics. *EMBO J.* **28,**
3074–89 (2009).

45. Karbowski, M. *et al.* Opposite effects of microtubule-stabilizing and microtubule-
destabilizing drugs on biogenesis of mitochondria in mammalian cells. *J. Cell Sci.*
114, 281–291 (2001).

46. Höög, J. L. *et al.* Organization of interphase microtubules in fission yeast analyzed by
electron tomography. *Dev. Cell* **12,** 349–61 (2007).

47. Mears, J. A. *et al.* Conformational changes in Dnm1 support a contractile mechanism
for mitochondrial fission. *Nat. Struct. Mol. Biol.* **18,** 20–26 (2011).

48. Taguchi, N., Ishihara, N., Jofuku, A., Oka, T. & Mihara, K. Mitotic phosphorylation of
dynamin-related GTPase Drp1 participates in mitochondrial fission. *J. Biol. Chem.*
282, 11521–9 (2007).

49. Chang, C.-R. & Blackstone, C. Cyclic AMP-dependent protein kinase phosphorylation
of Drp1 regulates its GTPase activity and mitochondrial morphology. *J. Biol. Chem.*
282, 21583–7 (2007).

50. Han, X.-J. *et al.* CaM kinase Iα–induced phosphorylation of Drp1 regulates
mitochondrial morphology. *J. Cell Biol.* **182,** 573–585 (2008).

51. Ferreira-da-Silva, A. *et al.* Mitochondrial Dynamics Protein Drp1 Is Overexpressed in
Oncocytic Thyroid Tumors and Regulates Cancer Cell Migration. *PLoS One* **10,**
e0122308 (2015).

52. Itoh, K., Nakamura, K., Iijima, M. & Sesaki, H. Mitochondrial dynamics in
neurodegeneration. *Trends Cell Biol.* **23,** 64–71 (2013).

53. Dubey, J., Ratnakaran, N. & Koushika, S. P. Neurodegeneration and microtubule
dynamics: death by a thousand cuts. *Front. Cell. Neurosci.* **9,** 343 (2015).

54. Kroboth, K. *et al.* Lack of Adenomatous Polyposis Coli Protein Correlates with a
Decrease in Cell Migration and Overall Changes in Microtubule Stability. *Mol. Biol.*
Cell **18,** 910–918 (2007).

55. Senft, D. & Ronai, Z. A. Regulators of mitochondrial dynamics in cancer. *Curr. Opin.*
Cell Biol. **39,** 43–52 (2016).

56. Forsburg, S. L. & Rhind, N. Basic methods for fission yeast. *Yeast* **23,** 173–183
(2006).

57. Kanter-Smoler, G., Dahlkvist, A. & Sunnerhagen. Improved Method for rapid
Transformation of Intact Schizosaccharomyces pombe cells. *Biotechniques* **16,** 798–
800 (1994).

Acknowledgements

We thank J. M. Thankachan, S. S. Nuthalapati and M. Ayushman for pilot experiments;

High Content Imaging Facility at BSSE, IISc, and P. I. Rajyaguru for the use of the InCell

6000, and Deltavision RT microscopes respectively; P. Delivani, I. Jourdain, R.C. Salas, M.

Takaine, I. Tolic, P. Tran, and NBRP Japan for yeast strains and constructs; S. Jain for

CellROX reagent. The research was supported by the Department of Science and

Technology (India)-INSPIRE Faculty Award, the Department of Biotechnology (India)

Innovative Young Biotechnologist Award, and the Science and Engineering Research

Board (SERB, India) Early Career Research Award awarded to V.A., and SERB Early

Career Research Award awarded to S.J.

Author Contributions

K.M. carried out the Dnm1 experiments, M.K.C. and S. J. carried out and analysed the flow

cytometry experiments, V.A. performed all the other experiments and data analysis,

designed the project, and wrote the paper.

Competing Financial Interests

The authors declare no competing financial interests.

646
647

648
Extended Data Figure 1. Depolymerisation of microtubules increases mitochondrial
fission. a, Maximum intensity projections of deconvolved Z-stack images of Cox4-GFP
(left), Mitotracker Orange staining (centre) and their composite (right) in interphase cells
(strain PT.1650, see Table S1) showing colocalisation of both signal intensities. **b,**
Maximum intensity projections of deconvolved Z-stack images of microtubules (left),
mitochondria (centre) and their composite (right) in fission yeast cells at the onset of
mitosis. Time in mm:ss is indicated to the left of the images. **c,** Montage of maximum
intensity projected confocal Z-stack images of microtubule depolymerisation (top) and
mitochondria (bottom) of wild-type cells (strain KI001, see Table S1) treated with MBC (see
Methods) represented in the intensity map indicated at the bottom of the images. The open
white arrowhead indicates a representative fission event. **d,** Box plot of the volume of
individual mitochondria in wild-type ('WT'), Klp5Δ/Klp6Δ ('MT_{long}') and Klp4Δ ('MT_{short}') cells
normalised to mean mitochondrial volume in wild-type cells (n=49, 28 and 100 mitochondria
respectively). Light grey crosses represent outliers, asterisk represents significance
($p<0.05$) and 'n.s.' indicates no significant difference. **e,** Maximum intensity projections of
deconvolved Z-stack images of microtubule (left), mitochondria (centre) and their composite
(right) in interphase Mal3Δ cells (strain KN001, see Table S1) showing several small
mitochondrial fragments similar to those seen in Klp4Δ cells.

Extended Data Figure 2. Microtubule depolymerisation induces increased fission in Klp5Δ/Klp6Δ cells. a, Montage of maximum intensity projected confocal Z-stack images of MBC-treated wild-type (WT^{mbc}, strain KI001, see Table S1) and Klp5Δ/Klp6Δ ('MT$_{long}$^{mbc}', strain G3B, see Table S1) cells represented in the intensity map indicated to the right of the images. White open arrowheads point to representative fission events. 00:00 indicates time (mm:ss) 2min after addition of MBC. **b,** Evolution of mitochondrial number with time indicated as mean (solid grey line) and standard error of the mean (shaded region) for wild-type ('WT'), WT^{mbc}, and MT$_{long}$^{mbc} cells (*n*=21, 14 and 7 cells respectively). **c,** Box plot of the size of mitochondria in WT, WT^{mbc}, and MT$_{long}$^{mbc} cells, calculated as the length of the major

axis of an ellipse fitted to each mitochondrion (n=1613, 1765 and 886 mitochondria
respectively). **d,** Box plot of the size of individual mitochondria in WT, WT^{mbc}, and MT$_{long}^{mbc}$
cells normalised to their cell lengths (n=1613, 1765 and 886 mitochondria respectively). **e,**
Box plot of the circularity of mitochondria in WT, WT^{mbc}, and MT$_{long}^{mbc}$ cells (n=1613, 1765
and 886 mitochondria respectively). **f,** Box plot of the aspect ratio of mitochondria in WT,
WT^{mbc}, and MT$_{long}^{mbc}$ cells (n=1613, 1765 and 886 mitochondria respectively). **g,** Box plot of
the fission frequencyof mitochondria per second in WT, WT^{mbc}, and MT$_{long}^{mbc}$ cells (n=21,
14 and 7 cells respectively). **h,** Box plot of the fusion frequencyof mitochondria per second
in WT, WT^{mbc}, and MT$_{long}^{mbc}$ cells (n=21, 14 and 7 cells respectively). **i,** Bar plot of cellular
ROS measured using flow cytometry (see Supplementary Information) in live WT, MT$_{long}$
and MT$_{short}$ cells normalised to mean of the ROS intensity in WT cells (n=3 independent
experiments, ≥10,000 events each). Light grey crosses represent outliers, asterisk
represents significance (p<0.05) and 'n.s.' indicates no significant difference. Note that the
WT data represented in this figure is the identical to the wild-type data plotted in Fig. 2 and
Extended Data Fig. 2 and has been re-used for comparison.

692
Extended Data Figure 3. Microtubule dynamics determines the morphology of
mitochondria. a, Box plot of the size of individual mitochondria in wild-type ('WT'),
Klp5Δ/Klp6Δ ('MT$_{long}$') and Klp4Δ ('MT$_{short}$') cells normalised to their cell lengths (n=1613,
739 and 1326 mitochondria respectively). **b,** Box plot of the circularity of mitochondria in
WT, MT$_{long}$ and MT$_{short}$ cells (n=1613, 739 and 1326 mitochondria respectively). **c,** Box plot
of the aspect ratio of mitochondria in WT, MT$_{long}$ and MT$_{short}$ cells (n=1613, 739 and 1326
mitochondria respectively). Error bars represent standard deviation. Light grey crosses
represent outliers, asterisk represents significance (p<0.05) and 'n.s.' indicates no
significant difference.

702
Extended Data Figure 4. Mitotic spindle length does not influence number of
mitochondria. a, Maximum intensity projections of deconvolved Z-stack images of the
microtubules (left), mitochondria (centre) and their composite (right) of WT cells (strain
KI001, see Table S1) with short (top) and long (bottom) mitotic spindles. The length of the
spindle is indicated to the left of the images. **b,** Montage of maximum intensity projected
confocal Z-stack images of the microtubules (top) and mitochondria (bottom) in a non-
mitotic wild-type cell (strain KI001, see Table S1) represented in the intensity map indicated
to the right of the images.. **c,** Bar plot of the mean number of mitochondria per cell before
('00:00') and after ('10:00') the same time window for which mitotic cells were monitored in
Fig. 3c, d. Solid grey lines represent data from individual cells (*n*=7 cells). **d,** Plot of number
of mitochondria in mother cell vs. partitioning error (see Supplementary Information)
between daughter cells in simulation (grey) and experiments (blue) expressed as mean
(solid grey line) and standard error of the mean (shaded area).

Scale bar: 2μm

c

Parameter	Value (mean ± s.e.m)		
	WT	MT$_{long}$	MT$_{short}$
Number of mitochondria	4.9 ± 0.4	2.3 ± 0.4	10 ± 0.9
Normalised total volume of mitochondria	1.00 ± 0.10	1.09 ± 0.13	0.92 ± 0.07
Normalised volume of individual mitochondria	1.00 ± 0.16	2.30 ± 0.50	0.45 ± 0.04
Size of mitochondria (μm)	2.72 ± 0.03	6.53 ± 0.11	1.77 ± 0.02
Size of mitochondria normalised to cell length	0.35 ± 0.01	0.59 ± 0.01	0.23 ± 0.01
Circularity of mitochondria	0.60 ± 0.01	0.44 ± 0.01	0.71 ± 0.01
Aspect ratio of mitochondria	2.84 ± 0.04	3.55 ± 0.07	2.16 ± 0.03
Mitochondrial fission frequency (s^{-1})	0.018 ± 0.002	0.006 ± 0.001	0.032 ± 0.005
Mitochondrial fusion frequency (s^{-1})	0.017 ± 0.002	0.011 ± 0.001	0.025 ± 0.005

716
Extended Data Figure 5. Dnm1 spots colocalise with the mitochondria and their
number is inversely proportional to the length of microtubules. a, Maximum intensity
projections of deconvolved Z-stack images of Dnm1 (left), microtubules (centre) and their
composite (right) in wild-type ('WT', strain FY7143 transformed with pREP41-Dnm1-Cterm-
GFP, see Table S1), Klp5Δ/Klp6Δ ('MT$_{long}$', strain FY20823 transformed with pREP41-
Dnm1-Cterm-GFP, see Table S1) and Klp4Δ ('MT$_{short}$', strain McI438 transformed with
pREP41-Dnm1-Cterm-GFP, see Table S1) cells. **b,** Box plot of the number of Dnm1 spots
in WT, MT$_{long}$ and MT$_{short}$ cells (*n*=13, 14, and 18 cells respectively). Note that the
mitochondrial morphology in WT, MT$_{long}$ and MT$_{short}$ cells appears abnormal due to inactivity
of the GFP-tagged Dnm1. Light grey crosses represent outliers, asterisk represents
significance (p<0.05) and 'n.s.' indicates no significant difference **c,** Measured parameters
relating to mitochondrial numbers and morphology in WT, MT$_{long}$ and MT$_{short}$ interphase cells
represented as mean ± s.e.m.
730

 Supplementary Information

Methods

ROS detection and viability assay using flow cytometry. A loopful of L972, FY20823 or

Mcl438 cells were inoculated in liquid YE and incubated overnight at 30°C in a shaking

incubator. Cells were harvested using a table-top centrifuge (2000g, 30s) and washed three

times with 1X PBS. The pellets were re-suspended in 500μl of 1X PBS, containing CellROX

(5μM) dye (ThermoFisher, USA). Following incubation at 30°C for 30min, cells were

washed three times with 1X PBS. Finally, cell pellets were re-suspended in an appropriate

volume of 1X PBS containing propidium iodide (6μg/ml), incubated for 10min at room

temperature, and run on a flow cytometer (BD FACS-Celesta). A minimum of 10,000 events

of cells was collected, and data analyzed using FlowJo (FlowJo LLC, USA). ROS was

reported as median fluorescence intensity for each dataset.

Estimation of partitioning error during cell division. The partitioning error of

mitochondria at cell division between the daughter cells Q_x was calculated from the

equation[24]:

$$Q_x^2 = \frac{(L - R)^2}{x^2}$$

where L and R were mitochondrial numbers in the resulting daughter cells and x, the

number of mitochondria in the mother cell. The data in Extended Data Fig. 4d were

simulated by independently assigning x mitochondria with 50% probability to either

daughter cell. Simulations were re-run 50 times to obtain the mean and the standard error

of the mean depicted in Extended Data Fig. 4d.

Counting of Dnm1 spots. Dnm1 was visualised in cells grown in the presence of 0.05μM

thiamine for partial induction of the nmt1 promoter. Even with only partial induction, a

majority of the cells showed extremely high expression of Dnm1, with most of the signal

concentrated in a single spot in the cell. The rest of the cells showed lower expression, with

Dnm1 signal distributed in a few spots across the cell. The latter were chosen for analysis

of number of Dnm1 spots. For counting the number of Dnm1 spots in Extended Data Fig.

5b, the maximum intensity projection of deconvolved Z-stack images were used. For

identification of spots, we set the lower threshold as 60-80% of the maximum intensity in

these images. The contrast in the images in Fig. 5a and Extended Fig. 5a reflect these

threshold settings.

Table S1

Strains and constructs used in this study

Name	Phenotype	Source
FY20823	*h- leu1 ura4 his7 Δklp5::ura4+ Δklp6::ura4+*	NBRP, Japan
FY7143	*h- ura4-D18 leu1-32 ade6-M216 his7-366*	Iva Tolić, Croatia
G3B	*h- Δklp5- Δklp6 – nmt1-GFP-atb2 leu ade*	Rafael Carazo Salas, UK
G5B	*h- klp4::kanr nmt1-GFP-atb2 leu ura*	Rafael Carazo Salas, UK
KI001	*h+ sid4-GFP::kanr kanr -nmtP3-GFP-atb2+ nmt1-pCOX4RFP::leu1+ ura4-D18 ade6-M210*	Iva Tolić, Croatia
KN001	*h + Δmal3::his3+ nmt1-pCOX4-dsRed ::leu1+ nmt1-atb1-GFP:LEU2 ade6-M210 ura4-D18 leu1-32 his3D1*	Iva Tolić, Croatia
L972	*h- WT*	Iva Tolić, Croatia
Mcl438	*h+ tea2d:his3 ade6 leu1-32 ura4-D18 his3-D1*	Iva Tolić, Croatia
MTY271	*h- mCherry-atb2:hphMX6 leu1-32 ura-d18*	Masakatsu Takaine, Japan
PT1650	*h+ cox4-GFP:leu1 ade6-M210 leu1-32 ura4-D18*	Phong Tran, USA
pREP41-Dnm1-Cterm-GFP (plasmid)	*Dnm1-GFP*	Isabelle Jourdain, UK
pREP41-Dnm1 (plasmid)	*Dnm1 (untagged)*	Isabelle Jourdain, UK

766 **Table S2**
767
768 **Partitioning of mitochondria between daughter cells.**

n	x	(n-x)	N	Obs.	Exp.	Obs.-Exp.
5	2	3	2	2	1.25	0.75
6	2	4	2	2	0.94	1.06
	3	3		0	0.63	-0.63
7	4	3	3	3	1.64	1.36
8	2	6	7	1	1.53	-0.53
	3	5		4	3.06	0.94
	4	4		2	1.91	0.09
9	3	6	11	4	3.61	0.39
	4	5		7	5.41	1.59
10	3	7	6	1	1.41	-0.41
	4	6		2	2.46	-0.46
	5	5		3	1.48	1.52
11	4	7	4	1	1.29	-0.29
	5	6		3	1.80	1.20
12	5	7	8	6	3.09	2.91
	6	6		2	1.80	0.20
13	5	8	5	3	1.57	1.43
	6	7		2	2.09	-0.09
14	5	9	7	1	1.71	-0.71
	6	8		4	2.57	1.43
	7	7		2	1.47	0.53
15	6	9	5	2	1.53	0.47
	7	8		3	1.96	1.04
16	6	10	2	1	0.49	0.51
	7	9		1	0.70	0.30
	8	8		0	0.40	-0.40
17	6	11	2	1	0.38	0.62
	7	10		0	0.60	-0.60
	8	9		1	0.74	0.26
18	8	10	1	1	0.33	0.67
	9	9		0	0.37	-0.37

769 n is the number of mitochondria in the mother, x and $n-x$ in the daughter cells at cell
770 division. N is the number of cells counted with n mitochondria in the mother. *Obs.* is the
771 observed number of cells showing partition of n into x and $n-x$, and *Exp.* is the expected
772 number of cells with the same x and $n-x$ partition given N cells predicted by the binomial
773 distribution.

Supplementary video captions

Video S1. Live-cell confocal microscopy of MBC-treated interphase fission yeast cells with the mitochondria stained with Mitotracker. Warmer colours indicate higher intensities. Images were acquired at 2min intervals. Time 00:00 (mm:ss) indicates time of addition of MBC. Imaging was resumed 15min after MBC addition ('15:00'). Scale bar represents 2μm. This movie corresponds to Extended Data Fig. 1c.

Video S2. 3D projection of microtubules (green) and mitochondria (magenta) in a wild-type cell (strain KI001, see Table S1). This movie corresponds to Fig. 1a.

Video S3. 3D projection of microtubules (green) and mitochondria (magenta, stained with Mitotracker) in a Klp5Δ/Klp6Δ cell (strain G3B, see Table S1). This movie corresponds to Fig. 1a.

Video S4. 3D projection of microtubules (green) and mitochondria (magenta, stained with Mitotracker) in a Klp4Δ cell (strain G5B, see Table S1). This movie corresponds to Fig. 1a.

Video S5. Mitochondrial dynamics in a wild-type cell (strain KI001, see Table S1) stained with Mitotracker. A 5-slice Z-stack with step size of 0.5μm was obtained every 12s. The video represents the maximum-intensity projected images of the stacks obtained. Warmer colours indicate higher intensity. The closed white arrowhead points to a fusion event and the open white arrowhead points to a fission event. Scale bar represents 2μm. This movie corresponds to Fig. 2a.

Video S6. Mitochondrial dynamics in a Klp5Δ/Klp6Δ cell (strain G3B, see Table S1) stained with Mitotracker. A 5-slice Z-stack with step size of 0.5μm was obtained every 12s. The video represents the maximum-intensity projected images of the stacks obtained. Warmer colours indicate higher intensity. The closed white arrowhead points to a fusion event. Scale bar represents 2μm. This movie corresponds to Fig. 2a.

Video S7. Mitochondrial dynamics in a Klp4Δ cell (strain G5B, see Table S1) stained with Mitotracker. A 5-slice Z-stack with step size of 0.5μm was obtained every 12s. The video represents the maximum-intensity projected images of the stacks obtained. Warmer colours indicate higher intensity. The open white arrowhead points to a fission event. Scale bar represents 2μm. This movie corresponds to Fig. 2a.

Video S8. 3D projection of microtubules (green) and mitochondria (magenta) in a wild-type mitotic cell (strain KI001, see Table S1). This movie corresponds to Fig. 3a.

Video S9. Microtubule dynamics (left) and mitochondrial dynamics (right) in a wild-type cell (KI001, see Table S1) at the onset of mitosis. A 5-slice Z-stack with step size of 0.5μm was obtained every 2min. The video represents the maximum-intensity projected images of the stacks obtained. Warmer colours indicate higher intensity. Scale bar represents 2μm. The open white arrowheads point to fission events. This movie corresponds to Fig. 3b.

820   **Video S10.** 3D projection of Dnm1 (green) and microtubules (magenta) in wild-type cells
821   (strain MTY271 transformed with pREP41-Dnm1-Cterm-GFP, see Table S1). This movie
822   corresponds to Fig. 5a.