The role of stimulus similarity on the summation effect in human causal learning

Omar D. Pérez[1, 2], Rene San Martín[3, 4], Fabián A. Soto[5]

[1] Division of Humanities and Social Sciences, California Institute of Technology

[2] Nuffield College CESS Santiago, Facultad de Administración y Economía, Universidad de Santiago

[3] Facultad de Economía y Empresa, Centro de Neuroeconomía, Universidad Diego Portales

[4] Center for Cognitive Neuroscience, Duke University, Durham, NC, USA

[5] Department of Psychology, Florida International University

Address correspondence to:
Omar D. Pérez
Division of Humanities and Social Sciences
California Institute of Technology
Pasadena, California, CA 91125
odperez@caltech.edu

Abstract

Data from animal experiments suggest that the level of responding to a compound of cues that have been separately trained with a common outcome is sometimes higher than responding to each of the cues separately. Several associative models anticipate that this effect, known as summation, is modulated by the degree of similarity of the cues forming the compound. However, little is known as to whether similarity plays a role in modulating summation in human causal learning, an area in which contemporary associative models are routinely applied. In this series of studies, participants in two groups were presented with two cues with different degrees of similarity that were paired with a common outcome. During testing, they were asked to estimate the level of outcome of the compound of the two cues. In variance with the predictions of associative theory, we found that participants showed strong and similar levels of summation in spite of several manipulations of stimulus similarity. Rational theories of causal learning propose that summation is the product of an assumption of causal independence of the target cues. Experimental manipulations used to modify this assumption produced a corresponding change in summation, regardless of stimulus similarity. Our results suggest that summation in causal learning is primarily driven by factors which are most readily explained by rational models.

Keywords: generalization, causal learning, summation

SUMMATION IN CAUSAL LEARNING

In natural environments, widely different stimuli are grouped together in classes (e.g., plants, cats, birds, etc.) and related with similar outcomes (e.g., nutrition, danger of injury or death, a potential mate, etc.). As a result, encoding these relationships allows an organism to take advantage of the regularities in the environment. Take, for example, a foraging animal that has got mildly sick after eating a particular type of food. The more the animal is able to estimate the level of sickness produced by foods with similar smell or coloration, the higher will be its chances of surviving and reproducing. This process, known as *generalization*, involves the identification of a newly experienced stimulus as being related to another previously experienced, allowing the animal to extract valuable information from past events and apply this information to novel situations (Brandon, Vogel, & Wagner, 2000; Rescorla, 1976; Soto, Gershman, & Niv, 2014; Soto & Wasserman, 2010). By generalizing between current and past events, organisms learn to respond similarly to different stimuli.

A particular form of stimulus generalization, known as *compound generalization*, is also critical for organisms to thrive in natural environments. The most typical example of this type of generalization is the *summation effect*, where the animal needs to estimate the level of sickness produced in the eventual case of eating two types of food that have independently produced a mild level of sickness in the past. From a psychological standpoint, the question at issue is whether the animal will predict a similar level of sickness for the compound of the two foods than to each of the two foods alone, or a stronger level of sickness for the compound of the two foods than to each of the two foods alone. If the prediction is that the level of sickness will be stronger, then it is said that summation has occurred.

In the lab, the summation effect is studied through a design in which two different stimuli, A and B, are separately paired with a common outcome and subsequently tested in compound (AB) (Collins & Shanks, 2006b; Soto, Vogel, Castillo, & Wagner, 2009). A

summation effect is obtained, in this context, if the response to the compound AB is greater than the response to the components A or B presented separately.

Different associative models of Pavlovian conditioning make different predictions for summation studies. These models have followed two theoretical traditions, each with a different view on how the stimuli A and B are represented separately and in compound. On the one hand, elemental models (Harris & Livesey, 2010; McLaren & Mackintosh, 2002; Rescorla & Wagner, 1972; Wagner, 2003) claim that a stimulus is represented by a finite (potentially large) number of elements which separately acquire their own associative strength in predicting the outcome. Under this view, summation comes about as a consequence of the higher associative strength of the compound AB than to each of A or B alone, implying that responding to AB should be a linear additive function of the associative strengths of the individual stimuli A and B.

A fundamentally different view, known as *configural theory*, was proposed by Pearce (1987, 1994) who claimed that subjects would represent the compound AB as a different stimulus (say AB=C) during testing. According to configural theory, responding to the compound AB will depend upon the degree to which the original learning to A and B generalizes to this new stimulus representation - that is, to the extent to which the compound AB is considered to be similar to the individual stimuli A and B. The particular generalization rule proposed by Pearce (1987, 1994) anticipates that only half of the associative strength from each component is generalized to AB, yielding a level of responding to AB that is similar to the average of responding to A or B alone.

More recently, both types of models have been extended to encompass the idea that the stimulus processing is a flexible phenomenon which can be modulated as a result of specific experimental manipulations (Melchers, Shanks, & Lachnit, 2008; Thorwart, Livesey,

& Harris, 2012; Wagner, 2003). Among these, one of the critical variables is the similarity between A and B. These theories claim, in particular, that the more similar these components are, the more configural processing will be fostered. Elemental processing, in contrast, will be fostered the more dissimilar these stimuli are. A weak summation effect should thus be observed when the components A and B are similar; strong summation, on the other hand, should come about when these components are dissimilar (Harris & Livesey, 2010; Harris, 2006; Kinder & Lachnit, 2003; McLaren & Mackintosh, 2002; Soto et al., 2014; Soto, Quintana, Pérez-Acosta, Ponce, & Vogel, 2015; Thorwart et al., 2012; Wagner, 2003).

Evidence in favour of this hypothesis comes from a study by Kehoe and colleagues using rabbits (Kehoe, Horne, Horne, & Macrae, 1994). In their study, one group of subjects was trained with very dissimilar stimuli A and B coming from different modalities (e.g., A=tone, B=light) while a second group was trained with more similar stimuli A and B, coming from the same modality (e.g., A=tone, B=noise). Consistent with associative theory, these authors found that responding was lower in the group trained with tone and noise compared to the group that was trained with tone and light.

Although different modalities are rather extreme manipulations of similarity, the critical idea shared by all contemporary associative models is that more configural processing, and therefore less summation, should come about when A and B are similar, whereas elemental processing, and therefore more summation, should be observed when A and B are dissimilar.

Summation in human causal learning

In a causal learning task, a summation design involves presenting participants with A and B paired with a similar outcome (e.g., that tomato and broccoli separately produce an allergic reaction) and subsequently asking for their estimation of the likelihood or magnitude

of the outcome in the presence of AB (e.g., what is the likelihood of allergic reaction after eating both broccoli and tomato?). As in the Pavlovian conditioning case, a summation effect is obtained when participants rate the causal or predictive power of AB to be higher than each of the two cues A or B separately.

There has been considerably recent interest on the summation effect using causal learning tasks (see, for example, Collins & Shanks, 2006b; Glautier, Redhead, Thorwart, & Lachnit, 2010; Soto et al., 2009; Thorwart et al., 2012; Vadillo, Ortega-Castro, Barberia, & Baker, 2014a; Van Osselaer, Janiszewski, & Cunha Jr, 2004; Waldmann, 2007). The interest has been largely fuelled by rational theories of causal learning, which offer an explanation of summation that is fundamentally different from that of associative theory. Summation, under a rational view, comes about as a consequence of subjects' assumptions about how the potential causes A and B interact in producing the common outcome. These theories claim, in particular, that in a causal learning scenario, people will regard A and B as being independent events (Cheng, 1997; Holyoak & Cheng, 2011a; Novick & Cheng, 2004). Such an assumption implies that the likelihood of observing an outcome when A and B are present is higher than the likelihood of observing this outcome when either A or B is present in isolation. Since stimulus similarity should not affect this causal independence assumption, summation should not be mediated by this factor under a rational approach.

The goal of this study was to test if cue similarity exerts an effect on the summation effect in human causal learning, as it does in Pavlovian conditioning (Kehoe et al., 1994). In a series of experiments, we presented participants with a summation design in which two cues known to produce a certain level of allergy in a hypothetical patient were presented in compound: participants had to predict, during testing, the level of allergy produced by this compound. We manipulated properties of the cues to make them more similar or dissimilar. To do so, we adapted the widely-used "allergy prediction" task (Dickinson, Shanks, &

SUMMATION IN CAUSAL LEARNING

Evenden, 1984), in which participants are trained to predict allergies in a hypothetical

patient. In our task, participants were asked to predict allergies to drugs, which were

represented by visually-presented shapes. These shapes could vary in configuration, rotation,

spatial position, and color, factors which allowed us to easily manipulate stimulus similarity.

In all the experiments, similarity was manipulated by having cues A and B differ either by

some degree within a single dimension (i.e., intra-dimensional difference; group *intra*) or by

differing across two or more dimensions (i.e., extra-dimensional difference; group *extra*).

According to associative learning theories, these similarity manipulations should

encourage either elemental or configural processing, and therefore lead to different levels of

responding to the compound in groups *intra* and *extra*. Evidence in favour of associative

theory would appear if the summation effect is stronger for group *extra* than for group *intra*.

Experiment 1

The stimuli used for cues A and B in Experiment 1 are shown in Figure 1. Cues in

group *extra* varied in both shape and color, whereas cues in group *intra* varied only in shape.

Table 1 shows the design of this experiment. Training consisted on two different cues

which independently predicted allergic reaction (A+/B+) and two fillers which independently

predicted no allergic reaction (C-/D-). There were two further compound fillers which

predicted allergic reaction (EF+) and no allergic reaction (GH-). On training, participants

assessed whether the allergy would be present or not by using a binary response (allergy/no

allergy). Feedback was provided about the magnitude of the allergic reaction produced by the

cue. During the test stage, participants had to estimate the magnitude of allergic reaction for

different cues and combinations of cues, by entering it on a rating scale.

Associative theories predict that the relatively similar stimuli shown to group *intra*

should be processed more configurally and therefore promote lower summation than the

relatively dissimilar stimuli shown to group *extra*, which should be processed more

elementally and therefore promote higher summation.

Method

Participants

30 undergraduate students from the University of Santiago, Chile. Participants were

randomly assigned to one of two groups (*n*=15 each). All participants were tested in the same

room at the same time and did not have any previous experience with similar research. They

were compensated with course credit for their participation. All procedures were approved by

University of California Santa Barbara IRB Protocol 13-0425: *The Cognitive Neuroscience of*

Human Category Learning.

Materials

Participants were tested on Intel i7 computers running Psychopy 1.77 (Peirce, 2007)

under Windows 7. The experiment was programmed with Psychopy 1.75. Computers were

connected to 17-inch Samsung color monitors.

Procedure

At the beginning of the experiment, participants were informed that all the instructions

would be presented on the screen. They were randomly assigned to different computers and

started the experiment at the same time.

The only difference between groups *intra* and *extra* was the stimuli assigned to the roles

of A and B in the design. For group *intra*, A and B varied along a single stimulus dimension

(shape), whereas for group *extra*, A and B varied along two stimulus dimensions (color and

shape, see Figure 1). During each trial of the training phase, a stimulus or pair of stimuli were

presented at the top-center of the screen, followed by the phrase "press 'a' to indicate an allergic

reaction, and 'n' to indicate no allergic reaction" at the bottom of the screen. The stimuli were

presented in one of two positions: immediately to the left of the horizontal middle line of the

screen, or immediately to the right of that line. The position (right vs. left) of each stimulus was randomly chosen during each trial.

After the participant entered a response, feedback was provided at the bottom-center of the screen for 2 seconds. On positive trials (marked + in design table) feedback consisted of the phrase "Allergic reaction" in red, followed in the next line by the phrase "10 points of allergy over a total of 20." On negative trials, feedback consisted of the phrase "No allergic reaction" in black, followed in the next line by the phrase "0 points of allergy over a total of 20."

Participants completed 25 blocks of training, each consisting of a single presentation of each of the 6 trial types shown in Table 1. Trial types were presented in random order within each block; the assignment of stimuli to the A-H conditions was semi-random. There were two pairs of stimuli that could be assigned to the roles of A and B. The stimuli in each pair were chosen based on similarity ratings obtained in a pilot study, so that the dissimilarity between A and B in group *extra* was higher than the dissimilarity between A and B in group *intra.* There were 6 other stimuli that differed in both shape and color and that were randomly assigned to the roles of C-H for each subject. During the testing phase, a new set of instructions was presented to the participants and then each of the trial types shown in the right column of Table 1 was presented once. The test stimulus was presented at the top-center of the screen, followed by the instructions "estimate the level of allergic reaction that this drug will cause in Mr. X." At the bottom of the screen, there were 21 circles with numbers inside ranging from 0 to 20. The left side of the scale was labeled "no allergic reaction" and the right side of the scale was labeled "maximal allergic reaction."

Results and Discussion

For all the experiments reported in this paper, the following criterion was used to exclude participants from the main analysis: All participants that failed to score on average

SUMMATION IN CAUSAL LEARNING

10 (+/- 3) points of allergy to cues that predicted allergy and 0 (+3) points of allergy to cues that did not predict allergy, were left out from the analysis. The rationale behind this criterion was to include data only from participants who successfully learned to discriminate between cues that predicted allergy and those that did not predict allergy.

The statistical analyses for all the experiments reported in this paper were performed using R version 3.3.0 and RStudio version 1.0.136 (RStudio Team, 2015), as follows. After fitting a linear-mixed model using the library nlme version 3.1-131 (Pinheiro, Bates, DebRoy, Sarkar, & R Core Team, 2016), a 2 x 3 mixed analysis of variance (ANOVA) was run. The ANOVA distinguished one between-subjects factor (group, with two levels: intra and extra) and one within-subjects factor (cue, with three levels: A, B and AB). A rejection criterion of $\alpha = .05$ was used. When the F test yielded a detectable difference for any factor or interaction, a Tukey's HSD (honest significant difference) post-hoc test was performed on those factors, using Bonferroni's method to correct for multiple comparisons. Eta- squared (η^2), along with a 90% confidence interval (CI) on this estimate, was calculated as a measure of effect size for each factor and interaction. A summation effect would be revealed in the post-hoc comparisons if the compound AB is scored higher than each of A and B. Different summation effects between the groups would be revealed by a significant Cue x Group interaction.

Thirteen participants did not meet the criterion in this study. The final number of participants per group was $n_{intra} = 9, n_{extra} = 8$. The results are shown in Figure 1. Both groups reported higher scores for the compound than for the individual cues ($F_{2, 30}=907.8$, p=.00, $\eta^2=.98$ 90% CI [.97, .99]), but no effects of group were found ($F_{1, 15}=1.95$, p=.18, $\eta^2=.11$ 90% CI [.00, .36]). Importantly, there was no detectable difference in summation between the groups, as revealed by a nonsignificant interaction between group and cue ($F_{2,}$

SUMMATION IN CAUSAL LEARNING

$_{30}$=1.74, p=.19, η^2 =.10 90% CI [.00, .25]). Post hoc analysis revealed significant differences for cues A and AB, and for cues B and AB (p<.01 in both cases).

These experimental results show that summation can be obtained in causal learning when the stimuli forming the compound differ either in colour and shape (group extra) or only in shape (group intra). This expands on previous reports of summation in human causal learning (Collins & Shanks, 2006b; Glautier et al., 2010; Soto et al., 2009; Vadillo et al., 2014) and attests to the generality of this effect. However, there was no effect of stimulus similarity on the magnitude of summation obtained.

As Figure 1 shows, both groups scored the compound as being close to the linear sum of the individual predictions (around 20 points of allergy). It is possible that this is a consequence of the employment of an allergy scale which prompts participants to score the compound as the exact sum of the two individual predictions (the maximum possible rating was exactly double the magnitude of allergy predicted by cues A and B during training). In Experiment 2, the allergy scale was modified so that its maximum level was higher than 20 points of allergy, to avoid potentially prompting participants to score the compound as the maximum level of allergy on the scale. In addition, a different set of stimuli was used, varying in the dimensions of color and rotation, to increase the difference between group in similarity of A and B.

Experiment 2

The same design as in Experiment 1 was used in this experiment, except that the target stimuli now varied in color and rotation. The shape was now the same in both groups. In group *intra*, stimuli A and B were made near identical, by having them differ slightly in rotation, whereas in group *extra*, A and B were made sharply different, by having them differ not only in rotation but also in color (see Figure 3).

SUMMATION IN CAUSAL LEARNING

In Experiment 1 there was a very strong summation effect in both groups. From an associative perspective, this implies that both groups were deploying strongly elemental strategies in forming their predictions. Perhaps features of the stimuli different from similarity promoted strong elemental processing in Experiment 1. For example, cues were presented in clear spatial separation, and previous data suggest that spatial separation is a critical factor in modulating configural and elemental processing; spatially close stimuli should foster configural processing (Glautier, 2002; Melchers et al., 2008). To encourage configural processing in both groups in Experiment 2, both A and B were presented at the center of the screen.

To control for the possible effect of the rating scale on the ratings, the feedback during training showed a maximum possible outcome of 35 points of allergy. This value was explicitly chosen to be higher than the sum of the outcomes of each of the individual cues A and B, avoiding possible prompting on participants' predictions. The rating scale during testing was also changed accordingly.

Method

Participants

38 undergraduate students from the University of California Santa Barbara. Participants were randomly assigned to one of two groups (*n*=19 each). They were tested in the same way as in Experiment 1. Participants were compensated with course credit for their participation.

Materials

Participants were tested on Macintosh computers running Psychopy 1.77 (Peirce, 2007) under OSX. The experiment was programmed in Psychopy 1.75.

Procedure

The procedure was the same as in Experiment 1. The main differences were the

SUMMATION IN CAUSAL LEARNING

feedback participants received after each trial during training and the dimensions on which the stimuli varied. During training, for positive trials (marked + in Table 1) feedback consisted of the phrase "Allergic reaction" in red, followed in the next line by the phrase "10 points of allergy over a total of 35." On negative trials, feedback consisted of the phrase "No allergic reaction" in black, followed in the next line by the phrase "0 points of allergy over a total of 35."

During both phases participants were presented at the bottom of the screen with 36 circles with numbers inside ranging from 0 to 35. The left side of the scale was labeled "no allergic reaction" and the right side of the scale was labeled "maximal allergic reaction." Thus, unlike Experiment 1 in which participants had to predict the allergy using a binary entry (allergy/no allergy), in Experiment 2 they had to predict the exact level of allergy by entering a particular value on the allergy scale.

Results and Discussion

A total of 3 participants failed the pre-set criteria and were discarded from the study. The final number of participants per group was $n_{intra} = 17, n_{extra} = 18$. Figure 4 shows the results obtained in Experiment 2. Although summation was lower than in Experiment 1, both groups reported higher scores for the compound than the individual cues ($F_{2, 66}=71.39$, p<.01, $\eta^2=.68$, 90% CI [.57, .75]), but no detectable difference between groups ($F_{1, 33}=0.35$, p=.56, $\eta^2=.01$, 90% CI [.00, .12]) was found. The summation was similar between the groups ($F_{2, 66}= 0.71$, p=.49, $\eta^2=.02$, 90% CI [.00, .09]). Post hoc analysis revealed significant differences for cues A and AB (p<.01) and cues B and AB (p<.01).

Experiment 2 shows a similar pattern to that of Experiment 1. Participants in both groups predicted a higher outcome to the compound AB than to each of its constituent elements, but no detectable difference was found between the two groups in terms of the

effect. Contrary to the predictions of associative theory, the mean for group intra was higher than the mean for group *extra* in this experiment. This result was obtained regardless of our attempt to increase the difference in similarity between groups *extra* and *intra* by changing the dimensions in which the components A and B varied, and regardless of the change in the rating scale employed to assess the predictions during both phases. The present study is therefore in line with the previous experiment in that summation appears to be a strong effect in humans which does not depend on the rating scale used or the similarity of the cues forming the compound.

Experiment 3

Taken together, the previous experiments suggest that, if there is an effect of stimulus similarity on summation in human causal learning, then this effect is too small to be detected by our manipulations of brightness, color, shape and rotation. Furthermore, the summation effect appears to be strong even when configural strategies are encouraged by presenting stimuli with a high degree of spatial overlap. Still, the previous experiments did not include a manipulation of spatial distance/overlap as a potential factor determining the level of summation observed. In Experiment 1, cues A and B were spatially separated in both groups, and in Experiment 2, cues A and B were spatially close and overlapping in both groups. There was a substantial reduction in the rating to AB from Experiment 1 to Experiment 2 (from an average around 19 to an average of around 17), but it is not clear whether this reduction was due to the change in spatial separation of A and B, or to any of the other changes across experiments (maximum value of the rating scale, stimulus shapes and color, etc.). As the previous literature suggests spatial separation as a potential stimulus factor controlling the level of elemental vs. configural processing of stimuli (Glautier, 2002; Melchers et al., 2008; Soto et al., 2014), Experiment 3 was performed to further assess the potential effect of spatial separation of cues on the summation effect.

SUMMATION IN CAUSAL LEARNING

The same stimuli of Experiment 2 were used in this experiment. However, now A and B differed only in rotation in group *intra*, whereas they differed in rotation and translation (or spatial position; see Figure 5) in group *extra*. This resulted in the components A and B in the compound AB having a high degree of spatial overlap in the group *intra*, but no overlap in group *extra*. A high degree of overlap between the components should have promoted configural encoding in group *intra*. In contrast, spatially-distant stimuli in group *extra* should have promoted elemental encoding. Under these conditions, we expected the summation in group *extra* to be stronger than in group *intra*.

Method

Participants

31 undergraduate students from the University of California, Santa Barbara. Participants were randomly assigned to one of two groups ($n_{intra}=16$, $n_{extra}=15$). They received course credit for their participation in this study.

Materials

Participants were tested in the same way as in Experiment 2, but cues A and B in group *extra* differed in rotation and spatial location, as shown in Figure 5.

Procedure

The procedure was the same as in Experiment 2. The stimuli A and B had the same color (dark blue) and shape. For group *intra*, stimulus B had a slight rotation to the right with respect to stimulus A (see Figure 5), but both of them were presented in compound at the center of the screen. For group *extra*, B had a slight rotation to the right, and A and B had a substantial horizontal separation when presented in compound AB.

Results and Discussion

A total of 5 participants failed the pre-set criteria and were discarded from the study. The final number of participants per group was n_{intra}=14, n_{extra}=12. Figure 6 shows the results of Experiment 3.

As in previous experiments, groups reported higher scores for the compound than the individual cues ($F_{2, 48}$=54.41, p<.01, η^2=.69, 90% CI [.55, .76]), but no detectable difference between groups ($F_{1, 24}$=0.04, p=.41, η^2=.00, 90% CI [.00, .06]) was found. The summation effect did not differ between the groups, as revealed by a nonsignificant interaction between group and cue ($F_{2, 48}$= 0.56, p=.57, η^2= .02, 90% CI [.00, .10]).

The pattern of the two previous experiments was replicated in this experiment, with no effect of stimulus properties on the summation effect, regardless of using a different manipulation–spatial contiguity–thought to influence elemental/configural encoding. As spatial contiguity can be construed as an aspect of cue similarity (Soto et al., 2014), this experiment aligns with the previous two in showing evidence that the similarity of the cues A and B does not affect he summation effect in human causal learning. The next experiment intended to maximise the effect of stimulus properties by designing A and B in each group so that perceptual similarity was as low as possible in group *intra* and as high as possible in group *extra*.

Experiment 4

In this experiment, group *intra* was trained with stimuli designed to have only a minimal difference in one dimension, while group *extra* was trained with stimuli designed to have extreme differences in many dimensions. These manipulations should test whether maximising the stimulus dissimilarity in group *extra*, while minimising it in group *intra*, can help modulate the summation effect between groups. Using highly dissimilar stimuli should

SUMMATION IN CAUSAL LEARNING

promote elemental encoding and bring about a higher summation effect in group *extra* compared with group *intra.*

Method

Participants

72 undergraduate students from the University of California, Santa Barbara. Participants were randomly assigned to one of two groups (n=36 each). They were tested in the same way as in previous experiments and received course credit for their participation.

Materials

The materials were the same as in previous experiments. For group intra, stimuli A and B varied only slightly in rotation, whereas for group *extra* they had extreme differences in rotation (the object's main axes were orthogonal to one another), color (dark blue vs. red), spatial position and configuration (see Figure 7).

Procedure

The procedure was the same as in previous experiments.

Results and discussion

A total of 23 participants failed the pre-set criteria and were discarded from the analysis. The final number of participants per group was n_{intra}=35, n_{extra}=14. Figure 8 presents the results for Experiment 5. As in previous experiments, the mean causal rating to the novel compound AB was greater than each of the cues A and B separately.

Groups reported higher scores for the compound than the individual cues ($F_{2, 92}$ =126.47 p<.01, η^2=.73, 90% CI [.65, .78]), but no detectable difference between groups ($F_{1, 47}$= 0.56, p=.46, η^2=.01, 90% CI [.00, .10]) was found. The summation effect did not differ between the groups ($F_{2, 92}$ = 0.80, p=0.45, η^2=.02, 90% CI [.00, .07]). Post hoc analysis revealed significant differences for cues A and AB (p<.01) and cues B and AB (p<.05).

SUMMATION IN CAUSAL LEARNING

Results from this experiment reinforce the idea that participants are insensitive to stimulus' similarity when estimating the causal status of the compound AB. Importantly, the present experiment shows that summation does not differ between groups *intra* and *extra* even when A and B differ in several dimensions and are explicitly designed to maximise elemental processing in group *extra*, while A and B differ only slightly in rotation and are presented with a high degree of overlapping as a compound in group *intra*, a design that should have maximised configural processing.

Experiment 5

None of the previous experiments found any evidence of an effect of stimulus similarity on the summation effect. However, in all previous experiments the filler cues that predicted no reward, C and D, differed from A and B in many ways (color, shape and configuration). Perhaps stimulus similarity has an effect on the summation effect, but that effect can only be observed when participants are explicitly trained to pay attention to features of A and B that highlight either their similarity or their dissimilarity. Take the stimuli shown in Figure 9. In the row labelled "Intra," cues A and B differ from cues C and D only in brightness. To successfully categorize these cues into classes that produce and do not produce allergy, people can learn to attend only to stimulus brightness and respond according to a very simple rule: "darker produces allergy". In the row labelled "Extra", cues A and B differ from cues C and D in more than just brightness. No single-dimensional rule allows to categorize A and B into one class, and C and D into a different class. There is one darker and one brighter stimulus in each class, and there is one "more circular" and one "more rhomboid" stimulus in each class. To successfully categorize these cues into classes that produce and do not produce allergy, people need to learn the identity of each specific stimulus. This could be done by memorizing the value of each stimulus in a single dimension, but is much easier to solve the task by paying attention to differences across both dimensions.

Instead of using the same design as in previous experiments, in Experiment 5 participants were required to solve the discrimination problems shown in Figure 9. For group *intra*, the stimuli A and B differed from stimuli C and D only in brightness; participants could therefore discriminate between those causing allergy from those not causing it by only paying attention to this dimension. For group *extra*, all stimuli differed both in color and shape. In this case, participants had to pay attention to differences along both dimensions to solve the task in an optimal way. It is important to note that stimuli in both groups can be described as points in a two-dimensional space created from a shape dimension (circular to rhomboid) and a brightness dimension. The stimuli were selected as to foster attention to one or two stimulus dimensions, but using similar linearly-separable classification tasks in both groups. While a non-linearly-separable task would have *forced* participants in the *extra* group to pay attention to both dimensions, this would have added a confound to our design. The demands of non-linearly-separable tasks, like negative patterning and biconditional discrimination, are thought to foster configural processing independently of stimulus factors (see Melchers et al., 2008). Our manipulation achieves the goal of promoting attention to different stimulus properties, highlighting the similarity or difference between cues A and B, without adding the confound of task demands to our design.

Method

Participants

28 undergraduate students from the University of California, Santa Barbara. Participants were randomly assigned to one of two groups (n=14). They were tested in the same way as in previous experiments and they received course credit for their participation.

Materials

The materials were the same as in previous experiments, but the stimuli used for cues A, B, C and D were as shown in Figure 9.

SUMMATION IN CAUSAL LEARNING

Procedure

The procedure was the same as in all previous experiments. The main difference was that the stimuli used in this experiment required participants to differentiate between target cues and fillers in a discrimination task (see Figure 9). The feedback and scales presented to participants were the same as in the previous experiment.

Results and discussion

A total of 8 participants failed the pre-set criteria and were discarded from the analysis. The final number of participants per group was n_{intra}=8, n_{extra}=12. Figure 10 shows the main results. Both groups estimated a higher allergy magnitude for the compound AB than to each of its constituent elements A and B.

Groups reported higher scores for the compound than the individual cues ($F_{2, 36}$ = 126.7, p= .00, η^2=.88, 90% CI [.80, .91]), but no detectable difference between groups ($F_{1, 18}$ = 2.51, p=.13, η^2=.12, 90% CI [.00, .35]) was found. The summation effect did not differ between the groups ($F_{2, 36}$ = 2.35, p=.11, η^2=.12, 90% CI [.00, .26]). Post hoc analysis revealed significant differences for cues A and AB (p<.01) and cues B and AB (p<.05).

These results show that even when people are trained in a task highlighting either similarities or differences between cues A and B, there is no effect of similarity on the observed summation effect. Even when participants are capable of estimating the levels of the outcome correctly, they nevertheless seem to disregard similarity in forming their predictions.

The results of Experiments 1-6 are largely unfavourable to contemporary associative learning models. As noted in the introduction, most current models designed to explain compound generalization phenomena, such as summation, predict changes in the magnitude of the summation effect when similarity is manipulated, as we have done in all previous experiments. At odds with this prediction, we have found a weak effect of similarity in the

SUMMATION IN CAUSAL LEARNING

difference of the AB scores between the groups *extra* and *intra.* Moreover, the difference

obtained was in some experiments numerically opposite to the predictions of associative

theory.

One could argue that many of these experiments used rather small sample sizes, leading to

under-powered statistical tests of our main hypothesis. One way in which these data can be

queried through a more powerful test, and provide us with better information as to the size of

the effect of similarity in summation is by collapsing all the previous studies in a single

estimated statistic. The approach we take here is one suggested by Rouder (2011), who

proposes a meta-analytic t-test based on Bayesian principles. This method combines the

evidence obtained in each experiment in terms of the difference in scores for AB in groups

extra and *intra* and determines the overall evidence in favor of the null in Experiments 1-5 by

calculating a Bayes Factor (BF_{01}) that compares evidence for the null hypothesis of no

difference against the alternative hypothesis of there being a difference in summation

between the groups. In our case, the value of BF_{01} for Experiments 1-5 was 7.34, implying

that the data are 7.34 times more likely to be obtained if there was no difference in

summation between the groups than if there was a difference between them. According to

Jeffreys (1961), this value can be interpreted as giving substantial evidence for the

hypothesis of no difference in summation between groups.

Experiment 6

One of the central topics in the study of learning is determining if the processes

underlying many of the most widely-observed phenomena in animal and human learning can

be explained by rational principles rather than associative mechanisms (Lovibond & Shanks,

2002; Mitchell, De Houwer, & Lovibond, 2009; Shanks, 2010). For example, rational theory

can offer a non-associative explanation for the summation effect by assuming that

participants rely on simple prior assumptions regarding the causal independence of A and B

in producing the outcome. If A and B are assumed to be independent, the predicted

magnitude of an outcome when AB is presented would follow an additive rule similar to that

present in elemental associative theory. Independence is, in fact, one of the critical

assumptions of many influential rational models of causal induction (Cheng, 1997; Cheng &

Novick, 1992; Holyoak & Cheng, 2011b), and a notion that has also been adopted and

discussed extensively in contemporary approaches to learning that attribute all associative

phenomena as arising from propositional rules (Declercq, De Houwer, & Baeyens, 2008;

Lovibond & Shanks, 2002; Mitchell et al., 2009). For example, the influential Power PC

theory proposed by Cheng and colleagues (Cheng, 1997; Holyoak & Cheng, 2011) assumes

that the joint predictive power of two candidate causes that independently predict an outcome

is given by the noisy-OR rule. If q_A and q_B represent, respectively, the probability of

observing the outcome when A or B are present, the noisy-OR rule states that the joint

predictive power of the candidate causes A and B will be given by $q_A + q_B - q_A q_B$. Because

q_A and q_B are both less than one, the theory predicts a higher causal power –or summation–

for the combination of the cues than for each of the cues separately.

From the perspective of rational theory, there is no reason to expect stimulus

properties to have an influence on the summation effect. As long as participants use the

assumption of independence, they should show the same level of summation across changes

in stimulus similarity, as found in Experiments 1-5. The ratings during the test stage would

reflect this assumption and be close to a linear sum of each of their single predictions,

provided an equal weight to each of the candidate causes ("A and B are independent,

therefore A and B together should cause near 20 points of allergy").

If the assumption of independence is the main mechanism driving summation in

human causal learning, then manipulations other than changes in cue similarity are likely

bring about rational rules that may impact on the level of summation obtained. Previous research shows, for example, that the application of these type of rules can influence compound generalization. Shanks and Darby (1998) gave participants training with cues A and B followed by the outcome in isolation (A+, B+), but not in compound (AB-), and cues C and D followed by the outcome in compound (CD+), but not in isolation (C-, D-). The critical manipulation was to present participants with the target cues W and X followed by the outcome (W+, X+), and cues Y and Z not followed by the outcome (Y-, Z-). Associative theories would predict that the compound WX will be a stronger predictor of the outcome than YZ, since the latter two have never been followed by the outcome. Surprisingly, this is not what Shanks and Darby (1998) found: participants predicted that the compound WX would be less likely to be followed by the outcome. This is, of course, in sharp contrast with the predictions of any standard associative model, which only considers the associative strength acquired in training by Y, Z, W and X, independently of the relationship between other cues (e.g., A and B) and their compounds (AB). In the Shanks and Darby's study, however, participants appear to have formed a general rule that made them believe that the relationship between A, B, and C, D with their respective compounds AB and CD during training ("the compound predicts an outcome opposite of that predicted by the individual cues") should be the same as that of W, X, and Y, Z and their respective compounds WX and YZ during testing.

The aim of this experiment was to explore to what extent a simple rational rule grounded on causal independence was driving participants' behaviour in previous experiments. To this end, participants were presented during training with examples of cues which predicted the same level of allergy both when presented as a single stimulus and in compound (X+/XX+), and others that predicted no allergy both when presented as a single stimulus and in compound (Y-/YY-). This manipulation prompted participants to learn the

rule that that extremely similar (i.e. identical in all aspects, except spatial position) cues combine according to a non-additive rule. In addition, we included the same stimulus manipulation as in Experiment 1 for groups *intra* and *extra*, to determine whether stimulus similarity might have an effect on the summation effect when people assume a non-additive combination rule for extremely similar stimuli. That is, we expected that the non-additive combination rule, learned from extremely similar stimuli, would be applied more readily to similar stimuli in group *intra*, rather than to dissimilar stimuli in group *extra*. Table 2 shows the design of Experiment 6. The only difference with the previous experiments was the inclusion of trials involving the two additional filler cues, X and Y.

Method

Participants

46 undergraduate students from the University of Santiago. Participants were randomly assigned to one of two groups (*n*=23). Participants were tested in the same way as in Experiment 1 and they received course credit for their participation.

Materials

The materials were the same as in previous experiments.

Procedure

The same stimuli as in Experiment 1 were used. The only difference with Experiment 1 was the addition of the two new fillers, X and Y. Cue X predicted an outcome of 10 points of allergy both when presented as a single stimulus (X+) and in compound (XX+). Cue Y predicted 0 points of allergy both when single (Y-) and in compound (YY-). Participants completed 25 blocks of training, each consisting of a single presentation of each of the 8 trial types shown in Table 2. Trial types were presented in random order within each block. The total number of training trials was 200. During training and testing we presented participants with the same rating scale as in Experiments 2 and 3 (i.e. a 0-35 points allergy scale).

SUMMATION IN CAUSAL LEARNING

Results and discussion

A total of 5 participants failed the pre-set criteria and were discarded from the analysis. The final number of participants per group was $n_{intra}=19$, $n_{extra}=22$. Figure 11 shows the results from the main summation test.

Groups reported higher scores for the compound than the individual cues ($F_{2, 78} = 9.89$, p<.01, $\eta^2=.20$, 90% CI [.07, .31]), but no detectable difference between groups ($F_{1, 39} = 0.00$, p=.93, $\eta^2=.00$, 90% CI [.00, .00]) was found. The summation effect did not differ between the groups, as revealed by a nonsignificant interaction between group and cue ($F_{2, 78} = 0.021$, p=.98, $\eta^2=.00$, 90% CI [.00, .01]). Post hoc analysis revealed significant differences for cues A and AB (p<.01) and cues B and AB (p<.05).

Consistent with our prediction that fostering the adoption of a non-linear combination rule would modulate the summation effect, this effect seemed much weaker for both groups in this experiment than in all previous experiments. This observation was supported by a comparison of the estimated size of the summation effect for the present experiment and a meta-analytic estimate of the size of the summation effect for all previous experiments. Figure 12 presents a forest plot for Cohen's D (and 95% CI) for the summation effect observed in Experiments 1-5, calculated as the difference between the ratings for the compound AB and the average rating for A and B, collapsing across groups. The overall effect size for the prior five experiments was 1.99, 95% CI [1.57, 2.42], which is four times larger than the effect of 0.5, 95% CI [0.25, 1.14] obtained for the present experiment. The fact that the CI from this experiment did not overlap with the meta-analytic CI from all previous experiments suggests that our manipulation was successful in decreasing the magnitude of summation obtained across both groups.

SUMMATION IN CAUSAL LEARNING

In sum, a manipulation of stimulus similarity was again ineffective in influencing the summation effect, even when participants were explicitly trained to assume a non-additive combination rule for extremely similar stimuli. Rather than showing a stronger summation effect for more dissimilar stimuli, participants simply applied the same non-additive rule regardless of the properties of cues A and B. These results suggest that a manipulation aimed at modifying participants' assumption of causal independence successfully influences the magnitude of the summation effect, but does not interact with stimulus similarity.

General discussion

In the series of experiments reported here, a summation effect was observed across a wide range of experimental manipulations. These results are in accord with previous studies looking into different aspects of summation in causal learning (Collins & Shanks, 2006b; Glautier et al., 2010; Soto et al., 2009) and indicate that summation is a robust effect in human causal learning. Manipulating the perceptual similarity and other properties of the components A and B did not, however, produce any difference in the magnitude of summation between the groups. Experiment 1 demonstrated equivalent summation in groups *intra* and *extra*, despite the fact that in the former cues A and B differed only in brightness, while in the latter cues A and B differed both in brightness and shape. Experiment 2 extended this conclusion, suggesting that the weak effect obtained in Experiment 1 cannot be attributed to features of the rating scale or the spatial distance between components A and B during testing –a factor thought to be critical for obtaining configural or elemental stimulus processing. Furthermore, the fact that changes in rotation and colour–rather than shape and brightness–did not change the result, suggests that the lack of an effect of similarity on summation cannot be attributed to the dimensions chosen to manipulate the stimuli. In Experiment 3, a variable thought to foster different generalization strategies, spatial separation (Melchers et al., 2008), was further explored. In group *intra*, A and B were

SUMMATION IN CAUSAL LEARNING

presented at the center of the screen and differed only slightly in rotation but had a high

degree of overlapping, a manipulation that should have encouraged configural processing.

The components A and B in group *extra* also differed slightly in rotation, but their spatial

separation during the test was large, a manipulation that should encourage elemental

processing. The results replicated those from the previous two experiments, in that

differences in summation failed to emerge even when spatial separation was minimized in

group intra, and maximized in group *extra.*

In Experiment 4, the stimuli were designed so that A and B in group *extra* were

extremely different, with extreme variations in colour and shape. As in Experiment 3, their

distance was also maximal on the screen. In contrast, group *intra* was presented with the

same stimuli as in Experiment 3. Once again, we did not detect differences in summation

between the groups.

Experiment 5 explored the possibility that stimulus similarity may have an effect on

the summation effect, but that effect can only be observed when participants are explicitly

trained to pay attention to features of A and B that highlight either their similarity or their

dissimilarity. Although people were trained in a task highlighting either similarities or

differences between cues A and B, there was no effect of similarity on the observed

summation effect.

The results from these five experiments are largely unfavourable to all contemporary

models of Pavlovian conditioning which assume that perceptual similarity should impact the

stimulus processing strategy deployed by subjects. To recapitulate, these theories anticipate

elemental processing and therefore summation with dissimilar stimuli, but configural

processing and lower summation with more similar stimuli (Harris & Livesey, 2010; Harris,

2006; Kinder & Lachnit, 2003; McLaren & Mackintosh, 2002; Soto et al., 2014, 2015;

SUMMATION IN CAUSAL LEARNING

Thorwart et al., 2012; Wagner, 2003). Modulation of other classic Pavlovian conditioning phenomena, such as blocking, are also anticipated by this idea (Soto, 2016).

The mechanisms by which these phenomena are modulated in different models vary, but the idea of similarity as a critical factor has been widely accepted in the field. The emphasis does not only pertain to associative theories. In a recent paper, Soto et al. (2014) proposed a rational model which also anticipates an effect of similarity on summation. The model combines the notion of consequential regions proposed by Shepard (1987) with contemporary Bayesian approaches to generalization (Courville, 2005; Courville, Daw, & Touretzky, 2006), and makes a similar prediction to that of associative models: similar stimuli should produce a weaker summation effect than dissimilar stimuli.

To be fair, most of these models were developed to deal specifically with Pavlovian conditioning phenomena, an area in which the influence of stimulus similarity and other stimulus properties on stimulus processing (elemental vs. configural) seems to be well documented (see Melchers et al., 2008; Wagner, 2003, 2008). However, it seems surprising that contemporary models of Pavlovian conditioning have given such a great importance to stimulus similarity in modulating summation, given the scarce evidence supporting this claim. In fact, most of the evidence is indirect. To the best of our knowledge, the most direct evidence comes from the study of Kehoe and colleagues (1994) in rabbits, and, as noted in the introduction, similarity was not directly manipulated in this study; the stimuli employed in this study were from either a different or the same modality, a design that should have maximized the perceptual difference in the group trained with different modalities while minimizing the difference in the group trained with stimuli from the same modality. In the present experiments, however, systematic manipulations of similarity failed to modulate the summation effect.

The fact that we could only found a weak impact of similarity on summation prompted us to hypothesise that other, non-associative factors needed to be explored to explain the data from Experiments 1 to 5. We thought the most reasonable interpretation for these experiments could be found in rational inductive theories (Cheng, 1997; Cheng & Novick, 1992; Mitchell et al., 2009) and propositional approaches to associative learning (Lovibond et al., 2003; Mitchell et al., 2009; Weidemann, Satkunarajah, & Lovibond, 2016). These theories regard summation as a consequence of a rational account in which participants' causal attribution to the cues forming a compound is based on a statistical independence assumption. A simple rule that takes each of A and B as a single entity that independently predicts the outcome (the noisy-OR rule) can readily explain the fact that summation was consistently obtained in spite of several manipulation of stimulus properties and procedural variables.

Experiment 6 explored these ideas. Here, stimuli were designed so that identity (the maximum level of similarity possible) did not necessarily yield a higher outcome in compound than each of the elements in isolation. After this manipulation, participants' score for the compound AB was much lower than in all previous experiments, suggesting that the addition of X+/XX+ and Y-/ YY- trials made participants to form a rule which implied that the relation between A and AB, and B and AB, was similar to that of X and XX or that of Y and YY. The rule, however, was equally applied by both groups, irrespective of the level of similarity of the components A and B.

The present data is most readily integrated within the current debate contrasting associative and rational approaches to causal learning in humans (Lovibond & Shanks, 2002; Mitchell et al., 2009; Shanks, 2010). Proponents of the rational view either rule out associative mechanisms, or consider associative predictions as a "null hypothesis against which to assess cognition" (Dickinson, 2012, pp. 2734). In the present study, the possibility

of an associative process playing a significant role on summation was abandoned in favour of

a rational interpretation which was evident when manipulations of causal independence were

performed. Indeed, previous studies found that such manipulations produce a similar effect

on other associative phenomena, like the blocking effect (Beckers et al., 2005; Mitchell et al.,

2009).

While it is possible that some modification of associative theory could provide an

explanation for the data reported in this paper, this series of studies provides good evidence in

favour of the view that summation in human causal learning is more readily explained by

rational approaches to causal learning. Contrary to the widely-held assumption of

contemporary models of Pavlovian conditioning, similarity does not appear to be a factor in

modulating this effect.

References

Borenstein, M., Hedges, L. V, Higgins, J., & Rothstein, H. R. (2009). Introduction to meta-analysis. Wiley Online Library.

Brandon, S. E., Vogel, E. H., & Wagner, A. R. (2000). A componential view of configural cues in generalization and discrimination in Pavlovian conditioning. *Behavioural Brain Research*, *110*(1–2), 67–72.

Cheng, P. W. (1997). From covariation to causation: A causal power theory. *Psychological Review*, *104*(2), 367–405. https://doi.org/10.1037//0033-295X.104.2.367

Cheng, P. W., & Novick, L. R. (1992). Covariation in natural causal induction. *Psychological Review*, *99*(2), 365–382.

Collins, D. J., & Shanks, D. R. (2006b). Summation in causal learning: elemental processing or configural generalization? *Quarterly Journal of Experimental Psychology (2006)*, *59*(9), 1524–1534. https://doi.org/10.1080/17470210600639389

Courville, A. C., Daw, N. D., & Touretzky, D. S. (2006). Bayesian theories of conditioning in a changing world. *Trends in Cognitive Sciences*, *10*(7), 294–300. https://doi.org/10.1016/j.tics.2006.05.004

Courville, A., Daw, N., & Touretzky, D. (2004). Similarity and discrimination in classical conditioning: A latent variable account. *Nips*, 614.

Declercq, M., De Houwer, J., & Baeyens, F. (2008). Evidence for an expectancy-based theory of avoidance behaviour. *Quarterly Journal of Experimental Psychology (2006)*, *61*(12), 1803–1812. https://doi.org/10.1080/17470210701851214

Dickinson, A. (2012). Associative learning and animal cognition. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *367*(1603), 2733–2742. https://doi.org/10.1098/rstb.2012.0220

Dickinson, A., Shanks, D. R., & Evenden, J. (1984). Judgement of act-outcome contingency: The role of selective attribution. *The Quarterly Journal of Experimental Psychology*, *36A*(1), 37–41. https://doi.org/10.1080/14640748408401502

Glautier, S. (2002). Spatial separation of target and competitor cues enhances blocking of human causality judgements. *The Quarterly Journal of Experimental Psychology*, *55*(2), 121–135. https://doi.org/10.1080/02724990143000207

Glautier, S., Redhead, E., Thorwart, A., & Lachnit, H. (2010). Reduced summation with common features in causal judgments. *Experimental Psychology*, *57*(4), 252–259. https://doi.org/10.1027/1618-3169/a000030

Harris, J. a. (2006). Elemental representations of stimuli in associative learning. *Psychological Review*, *113*(3), 584–605. https://doi.org/10.1037/0033-295X.113.3.584

Harris, J. A., & Livesey, E. J. (2010). An attention-modulated associative network. *Learning & Behavior*, *38*(1), 1–26.

Holyoak, K. J., & Cheng, P. W. (2011). Causal learning and inference as a rational process: the new synthesis. *Annual Review of Psychology*, *62*, 135–163. https://doi.org/10.1146/annurev.psych.121208.131634

Jeffreys, H. (1961). *The theory of probability*. OUP Oxford.

Kehoe, E. J., Horne, A. J., Horne, P. S., & Macrae, M. (1994). Summation and configuration between and within sensory modalities in classical conditioning of the rabbit. *Animal Learning & Behavior*, *22*(1), 19–26. https://doi.org/10.3758/BF03199952

Kinder, A., & Lachnit, H. (2003). Similarity and discrimination in human Pavlovian conditioning. *Psychophysiology*, *40*(2), 226–234.

Lovibond, P. E. F., Been, S.-L., Mitchell, C. J., Bouton, M. E., & Frohardt, R. (2003). Forward and backward blocking of causal judgment is enhanced by additivity of effect magnitude. *Memory & Cognition*, *31*(1), 133–142.

Lovibond, P. F., & Shanks, D. R. (2002). The role of awareness in Pavlovian conditioning: Empirical evidence and theoretical implications. *Journal of Experimental Psychology: Animal Behavior Processes*, *28*(1), 3–26. https://doi.org/10.1037/0097-7403.28.1.3

McLaren, I. P. L., & Mackintosh, N. J. (2002). Associative learning and elemental representation: II. Generalization and discrimination. *Animal Learning & Behavior*, *30*(3), 177–200. https://doi.org/10.3758/BF03192828

Melchers, K. G., Shanks, D. R., & Lachnit, H. (2008). Stimulus coding in human associative learning: flexible representations of parts and wholes. *Behavioural Processes*, *77*(3), 413. https://doi.org/10.1016/j.beproc.2007.09.013

Mitchell, C. J., De Houwer, J., & Lovibond, P. F. (2009). The propositional nature of human associative learning. *The Behavioral and Brain Sciences*, *32*(2), 183–246. https://doi.org/10.1017/S0140525X09000855

Morey, R. D., & Rouder, J. N. (2015). BayesFactor: Computation of Bayes Factors for Common Designs.

Novick, L. R., & Cheng, P. W. (2004). Assessing interactive causal influence. *Psychological Review*, *111*(2), 455–485. https://doi.org/10.1037/0033-295X.111.2.455

Pearce, J. M. (1987). A model for stimulus generalization in Pavlovian conditioning. *Psychological Review*, *94*(1), 61–73.

Pearce, J. M. (1994). Similarity and discrimination: a selective review and a connectionist model. *Psychological Review*, *101*(4), 587–607.

Peirce, J. W. (2007). PsychoPy—psychophysics software in Python. *Journal of Neuroscience Methods*, *162*(1), 8–13.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team. (2016). nlme: Linear and Nonlinear Mixed Effects Models.

Rescorla, R. (1976). Stimulus generalization: some predictions from a model of Pavlovian conditioning. *Journal of Experimental Psychology. Animal Behavior Processes*, *2*(1), 88–96.

Rescorla, R., & Wagner, A. R. (1972). A theory of Pavlovian conditioning: Variations in the effectiveness of reinforcement and nonreinforcement. *Classical Conditioning II: Current Research and Theory*, *2*, 64–99.

Morey, R. D., & Rouder, J. N. (2011). Bayes factor approaches for testing interval null hypotheses. *Psychological methods*, *16*(4), 406.

RStudio Team. (2015). RStudio: Integrated Development Environment for R. Boston, MA.

Schwarzer, G. (2007). meta: An R package for meta-analysis. *R News*, *7*(3), 40–45.

Shanks, D. R. (2010). Learning: from association to cognition. *Annual Review of Psychology*, *61*, 273–301. https://doi.org/10.1146/annurev.psych.093008.100519

Shanks, D. R., & Darby, R. J. (1998). Feature-and rule-based generalization in human associative learning. *Journal of Experimental Psychology: Animal Behavior Processes*, *24*(4), 405.

Shepard, R. N. (1987). Toward a universal law of generalization for psychological science. *Science*, *237*(4820), 1317-1323.

Soto, F. A. (in press). Contemporary associative learning theory predicts failures to obtain blocking . Comment on Maes et al . (2016). *Journal of Experimental Psychology: General.*

Soto, F. A., Gershman, S. J., & Niv, Y. (2014). Explaining compound generalization in associative and causal learning through rational principles of dimensional generalization. *Psychological review*, *121*(3), 526.

Soto, F. A., Quintana, G. R., Pérez-Acosta, A. M., Ponce, F. P., & Vogel, E. H. (2015). Why are some dimensions integral? Testing two hypotheses through causal learning experiments. *Cognition*, *143*, 163–177. https://doi.org/10.1016/j.cognition.2015.07.001

Soto, F. A., Vogel, E. H., Castillo, R. D., & Wagner, A. R. (2009). Generality of the summation effect in human causal learning. *Quarterly Journal of Experimental Psychology (2006)*, *62*(5), 877–889. https://doi.org/10.1080/17470210802373688

Soto, F. A., & Wasserman, E. A. (2010). Error-driven learning in visual categorization and object recognition: a common-elements model. *Psychological Review*, *117*(2), 349–381. https://doi.org/10.1037/a0018695

Thorwart, A., Livesey, E. J., & Harris, J. A. (2012). Normalization between stimulus elements in a model of Pavlovian conditioning: Showjumping on an elemental horse. *Learning & Behavior*, *40*(3), 334–346.

Vadillo, M. a, Ortega-Castro, N., Barberia, I., & Baker, a G. (2014). Two Heads Are Better Than One, but How Much? *Experimental Psychology*, 1–12. https://doi.org/10.1027/1618-3169/a000255

Van Osselaer, S. M. J., Janiszewski, C., & Cunha Jr, M. (2004). Stimulus generalization in two associative learning processes. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *30*(3), 626.

Wagner, A. R. (2003). Context-sensitive elemental theory. *The Quarterly Journal of Experimental Psychology. B, Comparative and Physiological Psychology*, *56*(1), 7–29. https://doi.org/10.1080/02724990244000133

Waldmann, M. R. (2007). Combining versus analyzing multiple causes: How domain assumptions and task context affect integration rules. *Cognitive Science*, *31*(2), 233–256.

Weidemann, G., Satkunarajah, M., & Lovibond, P. F. (2016). I Think, Therefore Eyeblink: The Importance of Contingency Awareness in Conditioning. *Psychological Science*. https://doi.org/10.1177/0956797615625973

Appendix

The two meta-analyses reported here were carried out using R version 3.3.0 and

RStudio version 1.0.136 (RStudio Team, 2015), extended with the *BayesFactor* 0.9.12

(Morey & Rouder, 2015) and *meta* 4.8.3 (Schwarzer, 2007) packages.

Meta-analysis 1: Bayesian meta-analytic t-test.

For the first analysis, we obtained the difference between the mean scores for the AB

compound between groups *extra* and *intra* and calculated an independent t-test for

Experiments 1-5. These values were then used, together with the sample sizes for groups

intra and *extra* on each study, to obtain an overall Bayesian Factor (BF_{01}) for these studies.

The value for the

Meta-analysis 2: Overall summation effect.

Our selected effect size was Cohen's *D*, as it allowed us to standardize the difference

between the mean scores obtained in each study. Although it could be argued that calculating

a raw difference between means would have sufficed to analyse the overall effect of these

experiments, the fact that there were changes in the rating scale presented to participants led

us to favor the selection of *D* as our effect size of interest.

Given the similarity of Experiments 1-5 in terms of the methods and question

involved, it is reasonable to assume that they all estimate the same underlying effect. In other

words, the estimated population effect size can be regarded as fixed, and all the variability

observed by randomness between experiments. In meta-analysis terminology, this is known

as a *fixed effect model* (Borenstein, Hedges, Higgins, & Rothstein, 2009). This was the

method chosen for our analysis.

For the analysis, the overall summation effect, SEF, was calculated as $SEF = AB - mean(A, B)$, where AB represents the mean score for AB, and $mean(A, B)$ the mean of A and B together, collapsed across both groups. The value of D for each of the studies was then calculated as $D = \frac{SEF}{s_{avg}}$, where s_{avg} represents the average of the standard deviations obtained for AB and $mean(A.B)$, respectively, given by $s_{avg} = \sqrt{\frac{s_{AB} + s_{mean(A,B)}}{2}}$. The values obtained where then entered into the *meta* package to obtain the results.

SUMMATION IN CAUSAL LEARNING

Table 1. Design of Experiment 1. The only difference between the group

intra and *extra* was the dimensions in which the stimuli A and B differed

during training.

Training	Test
Stimuli	A
A+	B
B+	EF
EF+	AB
Fillers	C
C-	D
D-	GH
GH-	CD

Note: Letters represent different chemical shapes which could be followed

(+) or not followed (-) by allergic reaction in a hypothetical patient.

SUMMATION IN CAUSAL LEARNING

Table 2. Design of Experiment 6.

Training	Test
Stimuli	A
A+	B
B+	EF
EF+	AB
Fillers	C
C-	D
D-	GH
GH-	CD
X+	X
XX+	XX
Y-	Y
YY -	YY

SUMMATION IN CAUSAL LEARNING

Figure 1. Target stimuli used in Experiment 1. Minimal variations in shape were used in both groups. Maximal variations in colour were used in addition for group *extra.* The positions of stimuli A and B in the compound AB were counterbalanced during both training and testing.

SUMMATION IN CAUSAL LEARNING

Figure 2. Mean causal ratings for cues A, B and AB for groups *extra* and *intra* in Experiment

1. Error bars represent 95% confidence intervals.

SUMMATION IN CAUSAL LEARNING

Figure 3. Target stimuli used in Experiment 2. Minimal variations in rotation were used in both groups. Variations in colour were used in addition for group *extra.*

SUMMATION IN CAUSAL LEARNING

Figure 4. Mean Causal ratings for cues A, B and AB for groups *extra* and *intra* in

Experiment 2. Error bars represent 95% confidence intervals.

SUMMATION IN CAUSAL LEARNING

Figure 5. Target stimuli used in Experiment 3. The position of stimuli A and B in the compound AB was counterbalanced in training. For group intradimensional, A and B were presented at the center of the screen; for group extradimensional they were presented to the left and to the right of the center of the screen. A and B respected these positions to form the compound AB during testing.

Group	A	B	AB
Extra			
Intra			

SUMMATION IN CAUSAL LEARNING

Figure 6. Mean Causal ratings for cues A, B and AB for groups extradimensional and

intradimensional in Experiment 3. Error bars represent 95% confidence intervals.

SUMMATION IN CAUSAL LEARNING

Figure 7. Target stimuli used in Experiment 4. The position of stimuli A and B in the compound AB was counterbalanced in training. For group intradimensional, A and B were presented at the center of the screen; for group extradimensional they were presented to the left and to the right of the center. A and B respected these positions to form the compound AB during testing.

SUMMATION IN CAUSAL LEARNING

Figure 8. Mean Causal ratings for cues A, B and AB for groups extradimensional and

intradimensional in Experiment 4. Error bars represent 95% confidence intervals.

SUMMATION IN CAUSAL LEARNING

Figure 9. Stimuli used in Experiment 5. In group intra, A and B differed from C and D only

in color. In group extra, all stimuli differed both in color and shape.

SUMMATION IN CAUSAL LEARNING

Figure 10. Mean Causal ratings for cues A, B and AB for groups extradimensional and

intradimensional in Experiment 5. Error bars represent 95% confidence intervals.

SUMMATION IN CAUSAL LEARNING

Figure 11. Mean Causal ratings for cues A, B and AB for groups *extra* and *intra* in

Experiment 6. Error bars represent 95% confidence intervals.

SUMMATION IN CAUSAL LEARNING

Figure 12. Meta-analysis for the summation effect (score for AB minus score for the average

of A and B in both groups) from Experiments 1 to 5. Each dot and error bar represent,

respectively, Cohen's D and 95% CI. The diamond at the bottom represents the overall effect

size across all experiments. The size of each dot represents the importance of each study in

the overall effect (bigger dots representing higher weighting factors). See Appendix for

details.