Title: A unifying motif for spatial and directional surround suppression

Abbreviated title: Surround suppression in MT

Author names and affiliation: Liu D. Liu[1], Kenneth D. Miller[2], and Christopher C. Pack[1]

[1]Department of Neurology and Neurosurgery, Montreal Neurological Institute, McGill

University, Montreal, Quebec H3A 2B4, Canada

[2]Department of Neuroscience, Center for Theoretical Neuroscience, Swartz Program in

Theoretical Neuroscience, Kavli Institute for Brain Science, College of Physicians and Surgeons,

Columbia University, New York, NY 10032, USA

Corresponding author: Christopher C. Pack;

christopher.pack@mcgill.ca;

Montreal Neurological Institute, Montreal, Quebec H3A 2B4, Canada

Number of pages: 42

Number of figures: 6

Number of words for Abstract, Introduction, and Discussion: 245, 650, and 1487

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: This work was supported by grants from the Canadian Institutes of Health

Research to C.C.P. (PJT-148488) and L.D.L. (CGSD-121719), and NIH R01-EY11001 and the

Gatsby Charitable Foundation (K.D.M.). We would like to thank Julie Coursol and the staff of

the Animal Care Facility (Montreal Neurological Institute) for excellent technical support.

Abstract

In the visual system, the response to a stimulus in a neuron's receptive field can be modulated by

stimulus context, and the strength of these contextual influences vary with stimulus intensity.

Recent work has shown how a theoretical model, the stabilized supralinear network (SSN), can

account for such modulatory influences, using a small set of computational mechanisms. While

the predictions of the SSN have been confirmed in primary visual cortex (V1), its computational

principles apply with equal validity to any cortical structure. We have therefore tested the

generality of the SSN by examining modulatory influences in the middle temporal area (MT) of

the primate visual cortex, using electrophysiological recordings and pharmacological

manipulations. We developed a novel stimulus that can be adjusted parametrically to be larger or

smaller in the space of all possible motion directions. We found, as predicted by the SSN, that

MT neurons integrate across motion directions for low-contrast stimuli, but that they exhibit

suppression by the same stimuli when they are high in contrast. These results are analogous to

those found in visual cortex when stimulus size is varied in the space domain. We further tested

the mechanisms of inhibition using pharmacologically manipulations of inhibitory efficacy. As

predicted by the SSN, local manipulation of inhibitory strength altered firing rates, but did not

change the strength of surround suppression. These results are consistent with the idea that the

SSN can account for modulatory influences along different stimulus dimensions and in different

cortical areas.

Significance Statement

Visual neurons are selective for specific stimulus features in a region of visual space known as the receptive field, but can be modulated by stimuli outside of the receptive field. The SSN model has been proposed to account for these and other modulatory influences, and tested in V1. As this model is not specific to any particular stimulus feature or brain region, we wondered whether similar modulatory influences might be observed for other stimulus dimensions and other regions. We tested for specific patterns of modulatory influences in the domain of motion direction, using electrophysiological recordings from MT. Our data confirm the predictions of the SSN in MT, suggesting that the SSN computations might be a generic feature of sensory cortex.

Introduction

Spiking activity is the main mechanism of communication in the nervous system.

Electrophysiological recordings provide a means of studying the relationships between spiking

activity and sensory stimuli, but they generally do not provide direct insight into the circuits that

give rise to these relationships. Such circuits involve an interplay of excitation and inhibition.

Recent work suggests that spiking activity in the visual cortex is modulated by a type of

circuit known as the Stabilized Supralinear Network (SSN) (Rubin et al., 2015). The SSN

involves recurrent excitation that amplifies weak inputs and inhibition that stabilizes network

activity. In V1, one type of modulatory influence is surround suppression, which is a decrease in

a neuron's firing rate when the size of a stimulus exceeds that of the receptive field (Allman et

al., 1985; Jones et al., 2001; Cavanaugh et al., 2002). Surround suppression is stronger for high-

contrast stimuli than for low-contrast stimuli (Sceniak et al., 1999; Pack et al., 2005; Tsui and

Pack, 2011).

For low-contrast stimuli, the SSN exhibits weak activation in the receptive field center,

while recurrent inputs primarily provide excitation. When the center is strongly activated,

recurrent interactions become stronger, and dominated by inhibition. The local network's

balance is then tilted towards inhibition, which suppresses both excitatory and inhibitory neurons

(Tsodyks et al., 1997; Ozeki et al., 2009). Thus, contrast-dependent surround suppression

emerges from the dynamics of recurrent activity, without the need for explicit assumptions about

different thresholds for excitation and inhibition (Rubin et al., 2015).

Although the model has been primarily tested with data from V1, the underlying

principles are generic (Ozeki et al., 2009; Rubin et al., 2015; Miller, 2016). That is, one should

be able to predict the response properties of neurons in other cortical regions based on the

structure of the SSN and the topography of the area from which the recordings are obtained. In

particular, if the connection strength between neurons decreases with their distance in a feature

space (e.g., preferred orientation in V1, (Cossell et al., 2015); or preferred direction in MT), then

the SSN model predicts that there should be contrast-dependent surround suppression in that

feature space, just as in retinotopic space (Rubin et al., 2015). Area MT provides a

straightforward means of testing the genericity of the SSN: MT contains a local columnar

structure based on selectivity for visual motion (Albright, 1984), so that nearby neurons encode

similar motion directions (Born and Bradley, 2005). The SSN therefore makes the prediction, by

analogy with V1 surround suppression, that MT firing rates should be decreased when stimuli

activate MT neurons with a wider range of motion preferences. Furthermore, in the SSN model,

such intracortical interactions vary with the overall level of activation, so that direction-domain

suppression should be stronger at higher contrasts. Finally, a counterintuitive aspect of the SSN

is that the strength of suppressive interactions are changed little by local blockade of GABAergic

inputs (Ozeki et al., 2004; Ozeki et al., 2009), because the suppression is caused by a withdrawal

of excitatory input that is not disrupted by local manipulations of inhibition.

We tested these ideas by designing a stimulus that could be manipulated parametrically to

be larger or smaller in the space of directions, while maintaining a fixed size in visual space. We

found that responses in MT were indeed suppressed by stimuli with a wider range of motion

directions, but only when the stimulus was high in contrast. At low contrast, neurons integrated

over a larger spread of motion directions, as has been observed for spatial integration (Levitt and

Lund, 1997; Kapadia et al., 1999; Sceniak et al., 1999). This provided support for a key

prediction of the SSN. In addition, we confirmed a counterintuitive property of the SSN, namely

that blockage of GABAergic inhibition does not reduce neuronal surround suppression. These

results are consistent with the idea that the SSN is a generic mechanism of cortical computation

(Miller, 2016).

Materials and Methods

Electrophysiological Recordings and Visual Stimuli

Two adult female rhesus monkeys (*Macaca mulatta*, both 7 kg) were used for

electrophysiological recordings in this study. Before training, under general anesthesia, an MRI-

compatible titanium head post was attached to each monkey's skull. The head posts served to

stabilize their heads during subsequent training and experimental sessions. For both monkeys,

eye movements were monitored with an EyeLink1000 infrared eye tracking system (SR

Research) with a sampling rate of 1,000 Hz. All procedures conformed to regulations established

by the Canadian Council on Animal Care and were approved by the Institutional Animal Care

Committee of the Montreal Neurological Institute.

Area MT was identified based on an anatomical MRI scan, as well as depth, prevalence

of direction-selective neurons, receptive field size to eccentricity relationship, and white matter

to grey matter transition from a dorsal-posterior approach. We recorded single units using linear

microelectrode arrays (V-Probe, Plexon) with 16 contacts.

Neural signals were thresholded online, and spikes were assigned to single units by a

template-matching algorithm (Plexon MAP System). Offline, spikes were manually sorted using

a combination of automated template matching, visual inspection of waveform, clustering in the

space defined by the principle components, and absolute refractory period (1 ms) violations

(Plexon Offline Sorter).

Visual motion stimuli were displayed at 60 Hz at a resolution of 1,280 by 800 pixels; the

viewing area subtended $60° \times 40°$ at a viewing distance of 50 cm. Stimuli consisted of random

dot stimuli displayed on a gray background (luminance of 98.8 cd/m^2). Half the dots were black,

and half the dots were white, resulting in a constant mean luminance across stimulus conditions.

At 100% contrast, the black dots had luminance of 0.4 cd/m^2, and the white dots had luminance

of 198 cd/m^2. The intermediate contrasts were defined as a percentage of the luminance

difference from the gray background luminance, contrast = |(luminance - 98.8 cd/m^2) / 98.8

cd/m^2|. Animals were trained to fixate on a small dot at the center of the screen. Stimuli were

shown after 300 ms of fixation. Each stimulus was presented for 500 ms, and the animals were

required to maintain fixation throughout the stimulus and for another 300 ms after the end of the

stimulus to receive a liquid reward. In all trials, gaze was required to remain within 2º of the

fixation point in order for the reward to be dispensed. Data from trials with broken fixation were

discarded.

The direction tuning and contrast response of the single units were quantified using 100%

coherent dot patches placed inside the receptive fields. Offline the receptive field locations were

further quantified by fitting a spatial Gaussian to the neuronal response measured over a 5 x 5

grid of stimulus positions. The grid consisted of moving dot patches centered on the initially

hand-mapped receptive field locations. We confirmed that all neurons included in our analysis

had receptive field centers within the stimulus patch used.

Size Tuning Stimuli in Direction Space

We designed a stimulus that would allow us to study surround suppression in the motion domain

in a manner that was analogous to studies in the spatial domain. In this conception, the input to

the receptive field "center" is the strength of motion in a range about the neuron's preferred

direction. The "surround" is then motion in other directions, and the bandwidth of the center plus

surround is the size of the stimulus in direction space. That is, a stimulus that contains motion in

a range of directions spanning 180° is larger than a stimulus that spans a range of 60°. For these

experiments we did not manipulate the spatial size of the stimulus, but rather fixed it according

to the size of the hand-mapped spatial receptive field.

Our stimuli made use of random dots, each of which could be assigned to either a noise

or a signal pool. The noise dots moved in random directions. The signal dots moved in a range of

directions that straddled the preferred direction of each neuron. All dots moved at the same fixed

speed of 8 or 16°/s, depending on the speed preference of the neuron. In all cases, dot patches

were centered on the receptive fields determined by hand mapping. All conditions were

interleaved randomly, and each stimulus was repeated 20 times.

We wished to change the size of the stimulus in direction space without changing other

stimulus variables to which the neurons were sensitive. However, changing the size in direction

space entails changing other low-level stimulus parameters (e.g., total number of dots or total

amount of motion energy), which could confound our interpretation of the data. We therefore

used two different methods to vary the stimulus bandwidth in direction space, each of which

entailed changing a different low-level aspect of the stimulus.

In the first method, we kept the total number of stimulus dots fixed, and increased the

motion bandwidth by drawing dots from a noise pool. Thus the total number of dots was

identical for all stimuli, across variations in direction bandwidth. We constructed stimuli that

contained signal dots moving in 1, 2, 3, and 4 directions, and each increase in the number of

motion directions involved recruiting 25% of the noise dots to move coherently in the new

direction (Fig. 2A). For the 1-direction stimulus, 25% of the dots moved in the neuron's

preferred direction, and 75% of the noise dots moved in random directions. For the 2-direction

stimulus, we took 25% of the total number of dots and assigned them motion directions in equal

numbers to 30° clockwise and counterclockwise from the preferred direction; to keep the total

number of dots constant we decreased the number of noise dots accordingly. This gives a

stimulus direction bandwidth of 60º (Fig. 2B). Similarly, for stimuli with 3 and 4 signal

directions, the noise dots pool was 25% and 0%, and the additional 25% of dots moved in

directions +/-60° and +/-90° from the preferred direction. This gives a stimulus direction

bandwidth of 120º and 180º, respectively (Fig. 2B). This paradigm thus allowed us to test the

influence of size in direction space for stimuli comprised of a fixed number of dots and a fixed

amount of overall motion energy.

However, in this approach, increases in motion bandwidth are yoked to decreases in

noise, which might be expected to affect the strength of inhibitory inputs on their own (Hunter

and Born, 2011). Thus, we also tested neurons using a second method, in which there was no

noise pool, and we increased the size in direction space by simply adding more dots that moved

in different directions. In this case the center stimulus strength (i.e. the strength of motion in the

preferred direction) was constant across conditions, but the total number of dots (and hence the

total motion energy) increased with stimulus size. The lowest dot density used was 2

dots/degree^2, which is beyond the density at which MT responses typically saturate, at least for

100% coherence stimuli (Snowden et al., 1992).

We again tested four different direction conditions (Fig. 3A). As before, stimuli with one

direction contained signal dots that all moved in the neuron's preferred direction, and dot density

was 2 dots/degree^2. For stimuli containing more signal directions, additional dots were added,

and these moved at 30° clockwise and counterclockwise from the preferred direction, raising the

dot density to 4 dots/degree^2. Similarly, for stimuli with 3 and 4 signal directions, the dot density

was 6 dots/degree^2 and 8 dots/degree^2, with additional directions being added at 30° increments.

For all size tuning experiments in direction space, we tested each of the 4 sizes at high

and low contrasts. High contrast was defined as 100% contrast, and the low contrast was chosen

online to be around the c_{50} of the contrast response function obtained with the 100% coherent dot

patch. Offline, we eliminated neurons for which the response at any tested contrast was below 2

standard deviations of the spontaneous baseline firing rate.

Grating, plaid, and pattern selectivity

We tested a subset of MT neurons (n = 65) with a standard measure of motion integration, the

plaid stimulus (Movshon et al., 1985). Direction selectivity for each neuron was first measured

with a 100% contrast drifting sinusoidal grating of spatial frequency of 0.5 cycles/°. Stimulus

size and temporal frequency were matched to the neuron's preferences. Plaid stimuli were

constructed by superimposing two gratings (Fig. 4A).

We used the standard approach to quantify the component and pattern selectivity of each

neuron (Smith et al., 2005). The partial correlations for the pattern and component predictions

were calculated as,

203
$$PC_p = \frac{r_p - r_c r_{pc}}{\sqrt{(1 - r_c^2)(1 - r_{pc}^2)}}$$

204
$$PC_c = \frac{r_c - r_p r_{pc}}{\sqrt{(1 - r_p^2)(1 - r_{pc}^2)}}$$

205   Where r_p and r_c are the correlations between the plaid response and the pattern and component

206   predictions, respectively, and r_{pc} is the correlation between the pattern and component

207   predictions. The partial correlations are z-scored as,

208
$$Z_p = 0.5ln\left(\frac{(1 + PC_p)/(1 - PC_p)}{\sqrt{1/(n - 3)}}\right)$$

209
$$Z_c = 0.5 ln \left(\frac{(1 + PC_c)/(1 - PC_c)}{\sqrt{1/(n-3)}} \right)$$

Where $n = 12$ is the number of directions. The pattern index was calculated as $Z_p - Z_c$.

Pharmacological Injections

The pharmacological injection system has been previously described (Liu and Pack, 2017).

Briefly, our linear electrode arrays contained a glass capillary with an inner diameter of 40 µm.

One end of the capillary was positioned at the opening between contacts 5 and 6 of the array

(contact 1 was most dorsal-posterior). The other end of the capillary was connected via plastic

tubing to a Hamilton syringe for the injection of pharmacological agents with a minipump.

To effectively manipulate neuronal responses without compromising isolation, we

typically used injections of 0.1-0.2 µL at 0.05 µL/min. For GABA, we used a concentration of

25 mM, which reduced neural activity without silencing it completely (Bolz and Gilbert, 1986;

Nealey and Maunsell, 1994). For gabazine, the concentration was 0.05 mM, and we used

injections of approximately 0.5 µL at 0.05 µL/min. In a few cases, this induced unstable and

synchronized responses in the nearby neurons (Chagnac-Amitai and Connors, 1989). The

electrophysiological recordings in those sessions were not further analyzed here.

Data Analysis

MT direction tuning curves $r(x_d)$ were characterized by fitting a Gaussian function to the mean

responses using the least-squares minimization algorithm (lsqcurvefit in MATLAB). The

Gaussian function is

230
$$r(x_d) = ae^{-0.5 d(\theta, x_d)^2 / b^2} + m$$

231   where a scales the height of the tuning curve; b determines the tuning curve width; x_d is the

232   motion direction; θ is the preferred direction of motion; and m is the baseline firing rate of the

233   cell. $d(\theta, x_d)$ is the shortest distance around the 360 degree circle between θ and x_d.

234   The contrast response functions $r(x_c)$ were fitted with a Naka-Rushton function,

235
$$r(x_c) = R_{max} \frac{x_c^n}{x_c^n + c_{50}^n} + m$$

236   where R_{max} scales the height of the contrast response function; n determines the slope; c_{50} is the

237   contrast at which the response function achieves half of its maximum response; and m is the

238   baseline firing rate of the cell. x_c is the contrast.

239   The neuronal size tuning curves $r(x_s)$ in retinotopic space were fitted by a Difference of

240   Error functions (DoE) (Sceniak et al., 1999; DeAngelis and Uka, 2003),

241
$$r(x_s) = A_e erf\left(\frac{x_s}{s_e}\right) - A_i erf\left(\frac{x_s}{s_e + s_i}\right) + m$$

where A_e and A_i scale the height of the excitatory center and inhibitory surround, respectively. s_e

and s_i are the excitatory and inhibitory sizes, and m is the baseline firing rate of the cell. x_s is the

stimulus size.

The size suppression index (SI_S) for each neuronal size tuning curve was calculated as

$SI_S = (R_m - R_L)/R_m$, where R_m is the maximum across responses to different stimulus sizes and

R_L is the response observed at the largest size. Since using the raw responses is sensitive to noise

at both the maximum response and the response at the largest size, we used the values from the

DoE fits for SI calculations.

Since we only measured the response at 4 sizes in the directional space, we were unable

to fit a DoE function to the directional size tuning curves. Instead, to capture potential

suppressive influences in the direction domain, we calculated a direction integration index from

the raw data $II_D = (R_L - R_S) / (R_L + R_S)$, where R_L is the response observed at the largest size and

R_S is the response observed at the smallest size.

SSN Model Simulations

We first simulated a 1D ring model, which captures putative interactions among neurons

representing different motion directions (Fig. 1A). Details of this model can be found elsewhere

(Rubin et al., 2015). Our model differs in that the ring is 360 degrees in extent (vs. 180 degrees

in Rubin et al., 2015), representing all possible motion directions. There is an excitatory (E) and

inhibitory (I) neuron at every integer position $x_i = 0^o$, 1^o, …, 359^o, where x_i represents the

preferred direction of the corresponding E and I cells. We can write the model equation in matrix

notation as,

$$\tau \frac{d}{dt} r(x_i) = -r(x_i) + k([W^*r(x_i) + ch(x_i)]_+)^n$$

where $r(x_i)$ is the vector of firing rates of the excitatory and inhibitory neurons with preferred

motion direction x_i, $W(y)$ is the weight matrix of E → E, E → I, I → E, and I → I connections

between neurons separated by angular distance y (measured as shortest distance around the 360

degree circle). The connections $W_{ab}(y) = J_{ab}G\sigma_{dir}(y)$, where $J_{EE} = 0.44$, $J_{EI} = 0.23$, $J_{IE} = 0.042$, J_{II}

$= 0.018$, $G\sigma_{dir}(y)$ is a Gaussian function with standard deviation of 64^o. $W^*r(x_i)$ is the

convolution $\sum_j W(x_i\text{-}x_j)r(x_j)$ where the sum is over all preferred directions x_j , $h(x_i)$ is the

vector of external input to the E and I neurons preferring x_i, and c is the strength (monotonically

related to contrast) of the input. The elements of the vector of input to the neuron,

$W^*r(x_i) + ch(x_i)$, are thresholded at zero before being raised to the power n:

$[z]_+ = 0\ if\ z < 0, = z\ if\ z \geq 0$ (the operations of thresholding and raising to a power are

applied separately to each element of the vector). k and n are identical for E and I neurons, with k

= 0.04 and n = 2. τ is a diagonal matrix of the time constant for E cells, τ_E = 20 ms, and for I

cells, τ_I = 10 ms.

We simulated network responses to random dot field stimuli of variable coherence. We

assumed that a coherent dot stimulus of a given direction gives input to MT neurons proportional

to a Gaussian function, of standard deviation 60º, of the difference (shortest distance around a

360º circle) between the neuron's preferred direction and the stimulus direction. The non-

coherent (noise) dots gave equal input, proportional to 1/360, to neurons of all preferred

directions. The strength of the stimulus is given by a parameter c, identified as the "contrast" in

Figure 1. As in our electrophysiological experiments, we used stimuli corresponding to 4

different sizes in direction space (Fig. 2A). Thus for the smallest size, 25% of the input, $\boldsymbol{h}$, was

modelled as a Gaussian distribution around the preferred direction (peak of the Gaussian = $c/4$),

while the remaining 75% was spread equally around the ring (uniform distribution of size (3/4) ×

c/360). At 2 directions, an additional 25% was taken from the non-coherent input and added to

Gaussian spreads about +/-30° from the preferred direction (these two Gaussians have peak =

c/8; noise amplitude becomes (1/2) × c/360). 3 and 4 directions followed in a similar manner

while the total input strength was kept constant across sizes. We also simulated the same set of

stimuli except without a noise background (so that the total input strength grew with increasing

number of directions), and the results were qualitatively similar as presented in Results.

Experimental design and statistical analysis

We used two female rhesus monkeys (*Macaca mulatta*) for electrophysiological recordings in

this study; this is standard for electrophysiological studies involving monkeys. We used the

Wilcoxon rank-sum test to evaluate the difference between the Integration Index at low and high

contrast, and the difference between Direction Tuning Width and Suppression Index before and

after injection of Gabazine. We calculated Pearson correlation coefficients to evaluate the

relationship between the Pattern Index and Direction Tuning Width with the Integration Index.

We used built-in MATLAB functions and custom scripts to perform the analyses. The complete

results of the statistical analyses for each experiment can be found in the corresponding Results

section.

Results

In this section, we first present simulation results for the SSN. We then test a crucial model

prediction with neurophysiological recordings from MT neurons in awake and behaving

macaques. The theoretical and empirical results show that surround suppression in the motion

domain behaves similarly to surround suppression in the space domain, with integration at low

contrasts switching to suppression at high contrasts (Figs. 2 and 3). We also find that pattern-

selective cells (as assayed from plaid responses) show greater motion integration than

component-selective cells (Fig. 4). Finally, as predicted by the SSN model, local

pharmacological manipulation of inhibition does not alter spatial surround suppression, although

our methods had the expected effects on directional tuning width (we did not examine effects on

surround suppression in the direction domain) (Figs. 5 and 6).

Stabilized supralinear network predicts contrast-dependent surround suppression in the

direction domain in MT

Previous instantiations of the SSN have considered a model in which connections are defined

either across a retinotopic sheet of the kind found in V1 or across a ring of preferred orientations

(Ahmadian et al., 2013; Rubin et al., 2015; Miller, 2016). Like orientation, motion direction is a

circular variable, but it takes values over 360º rather than 180º as for orientation. Thus to

examine the properties of the SSN in this circular space, we first simulated a ring model ((Rubin

et al., 2015); Fig. 1A) of motion direction space. This represents neurons of varying preferred

directions sharing a common location in retinotopic space.

In general, the SSN predicts that contrast-dependent surround suppression should occur

in any stimulus feature dimension, provided certain minimal connectivity conditions are met, e.g.

average connection strength between neurons decreases with the dimensional distance between

them. We accordingly assumed that the strengths of connections between neurons on the ring

decreased with increasing difference in their preferred directions. By analogy with the study of

size-tuning in the spatial domain, we tested the SSN with stimuli of different motion-domain

sizes. We increased the size of the stimulus in direction space by including stimuli at

increasingly wider ranges of directions about the preferred direction (the "center" of the

receptive field). As described in Methods, we considered size or bandwidth 0° (preferred-

direction stimulus only), 60° (adding stimuli at $+/- 30^\circ$ about the preferred), 120° (adding

additional stimuli at $+/- 60^\circ$), and 180° (additional stimuli at $+/- 90^\circ$). For each motion size, we

examined different levels of stimulus contrast, represented as scaling the strengths of all inputs.

The simulation results (Fig. 1B) show that the model predicts strong direction-domain

surround suppression at high contrast, but not at low contrast. Specifically, at low contrasts (red),

increasing the range of motion directions leads to increased responses with a hint of suppression

for the largest stimulus size, while at high contrasts larger motion-domain stimulus sizes lead to

strong suppression (blue). Intermediate contrasts give an intermediate result (black). These

results change very little with changes in the total number of dots in the stimulus (Fig. 1C), a

factor that we consider in our experiments below (Fig. 3). Thus the model consistently predicts

direction-domain suppression that is analogous to space-domain surround suppression. In the

SSN, the dependence of surround suppression on contrast arises generically from the dynamics

of the SSN in summing inputs, rather than by the assumption of a higher contrast threshold for

inhibition, as in previous models (Somers et al., 1998; Huang et al., 2008; Schwabe et al., 2010;

Carandini and Heeger, 2012).

Figure 1. Stabilized supralinear network can account for surround suppression in both spatial

and direction domains. **A,** Schematic of the 1D SSN ring model as a direction space analogue of

the visual space model. In the visual space model (top), stimuli of different sizes in visual space

(gray circles) are simulated as input, *h(x)*, of varying width, to a linear 1D grid of excitatory (E,

red) and inhibitory (I, blue) units. The grid positions represent visual space positions. In the

direction space (bottom), there are 360 E and I units, with coordinates on the ring as preferred

directions. A dot stimulus, *h(x)*, moving at a single direction is a Gaussian-shaped input with

standard deviation of 60º. Stimuli including multiple directions simply add such input for each

direction. We considered two methods of adding directions: including a "noise pool" stimulus of

equal input to all directions, and subtracting from the noise pool as we added directions to keep

total input strength unchanged (Fig. 2A); or simply adding additional input as we added

directions, without a noise pool (Fig. 3A). **B,** Directional surround suppression at high contrast,

but not at low contrast, arises from the dynamics of the model. This simulation result is for the

first method of taking dots from a noise pool to add further directions about the preferred (Fig.

2A). The response at each contrast is normalized to the peak response. **C,** The simulation result

for the second method of adding dots to further directions about the preferred without a noise

pool (Fig. 3A). The response at each contrast is normalized to the peak response.

Surround suppression in direction domain of MT

We tested the model predictions by recording from individual MT neurons, using the same

stimuli as in the simulations. We first show results for the first type of stimulus described above,

in which there was a noise pool of dots moving in random directions. For each neuron we fixed

the physical size of each stimulus according to an estimate of the classical receptive field size.

We then varied stimulus size in the motion domain, as well as dot contrast. Thus for the smallest

stimulus, all the coherent dots moved in the preferred direction of the neuron (Fig. 2A, left), with

the remaining dots in the noise pool moving in random directions. To increase the size of stimuli

in the motion space, we recruited dots from the noise pool and added them to directions around

the preferred direction (Fig. 2A). This manipulation kept the total motion energy and dot density

of the stimulus constant across sizes.

Figure 2B shows the firing rate of an example MT neuron for stimuli of different

contrasts and motion sizes. For the low contrast stimulus (red), firing rate increased with motion

size, while for higher contrasts (blue, black) firing rate decreased with motion size. Thus the

pattern of firing rates for this neuron was consistent with the SSN prediction that MT neurons

would shift from motion-domain integration to suppression as the stimulus contrast was

increased (Fig. 1B). Indeed, just as in the space domain, for large stimuli it is possible to increase

firing rates by lowering contrast (Fig. 2B; Pack et al., 2005).

To examine these effects across the MT population, we calculated the directional

integration index (II_D, the difference between responses to the largest and smallest sizes divided

by the sum of these responses; see Methods) for data of the kind shown in Figure 2B for 125

neurons. The II_D captures the integration of signals across motion directions, with larger II_D

values indicating more integration. Across the population (Fig. 2C) the II_D was frequently below

zero, indicating a suppression of the response when dots activated the directional surround.

Overall the II_D was significantly decreased at high contrast compared to low contrast, consistent

with reduced integration at high contrasts ($p < 0.001$, rank sum test). Note that this is not due to a

failure of the low contrast stimuli to elicit a response from the neurons, as all neurons except one

showed responses to the lowest contrast tested that were significantly above baseline. The one

neuron that failed to meet this criterion was eliminated from further analysis. Overall, these

results are similar to previous results in the space domain in MT (Pack et al., 2005; Tsui and

Pack, 2011). However, the mechanisms of spatial and directional integration for a given cell

appeared to be independent, as there was no correlation between the degree of spatial surround

suppression and directional surround suppression measured at high contrast in the same neurons

(Pearson's $r = -0.06$, $p = 0.46$, N = 124).

Figure 2. Surround integration and suppression in the direction domain. **A,** Illustration of the

stimulus that engages directional surround suppression in MT while the dot density is fixed. **B,**

Surround suppression occurs in direction space at high contrast, but not at low contrast for an

example neuron. **C,** Contrast response function for the same example neuron using 100%

coherent dots in the preferred direction. The line indicates the Naka-Rushton function fit. **D,**

Population data for direction surround integration. Scatter plot of the integration index, II_D, at

low contrast against the II_D at high contrast (rank sum test, $p < 0.001$). The marginal distributions

are histograms of the II_D (Median at low contrast = -0.003; Median at high contrast = 0.051).

Dashed lines in the histograms show location of $II_D = 0$.

We also tested 46 neurons using a second stimulus in which there was no noise pool, and

we increased the total number of stimulus dots with size in the direction domain (Fig. 3A). This

stimulus was designed to control for a potential confound in the previous experiment, which kept

the total number of dots constant across stimulus size. In the latter configuration, increases in

direction-domain size were yoked to decreases in the number of noise dots, and because noise

includes motion in all directions, this can be viewed as reduction in the strength of the directional

surround, analogous to the far surround in retinal space (Angelucci and Bullier, 2003; Angelucci

and Bressloff, 2006). The new stimulus was directly analogous to that typically used in size

tuning experiments, in which the stimulus is simply expanded to probe the influence of the

surround.

We tested this subpopulation of MT neurons with both stimuli, and the results are shown

in Figures 3B and 3C. For the control stimulus, the II_D is still significantly higher at low contrast

than at high contrast (Fig. 3B; $p = 0.04$, rank sum test). Thus integration across direction space

was greater at low contrast, regardless of how size was manipulated. For these neurons, we also

replicated the previous result using the stimulus with a constant total number of dots (Fig. 3C; p

< 0.001, rank sum test). The contrast modulation of II_D was not significantly different for the two

stimulus types (rank sum test, $p = 0.45$).

Figure 3. Additional controls for direction surround integration and suppression. **A,** Illustration

of the stimulus that engages directional surround suppression in MT while the dot density

increases with directional size. **B,** Population data for direction surround integration. Scatter plot

of the directional integration index (II_D) at low contrast against the SI at high contrast (rank sum

test, $p = 0.04$). The marginal distributions are histograms of the II_D (Median at low contrast = -

0.012; Median at high contrast = 0.018). Dashed lines in the histograms show location of $II_D = 0$.

C, The contrast modulation of II_D for the same 46 neurons as in B, when the number of dots is

held fixed by drawing from a noise pool (as in Fig. 2). The conventions are the same as in panel

B (Median at low contrast = 0.003; Median at high contrast = 0.065).

444   Of the complete MT population, 65 were also tested with a standard probe of direction-

domain integration, the plaid stimulus (Movshon et al., 1985). Our plaid stimuli consisted of two

superimposed sine-wave gratings, moving in directions separated by 120° (Fig. 4A); stimulus

size was again matched to the classical receptive field, and contrast was 100%. From the

resulting data we computed a pattern index (see Methods; Smith et al., 2005), which captures the

extent to which MT neurons integrate the two motion directions; higher values indicate greater

integration (Fig. 4B and C). We found that the pattern index was significantly correlated with the

directional II_D, as measured in our direction-size-tuning experiments at both low (Fig. 4D;

Pearson's $r = 0.33$, $p = 0.01$) and high contrasts ($r = 0.27$, $p = 0.03$). That is, cells with higher

pattern indices showed less surround suppression in direction space – greater motion integration

-- both at low and high stimulus contrasts. This suggests that area MT might use similar

mechanisms to integrate motion signals for dot stimuli and grating stimuli. We also found that

there was no correlation between the directional motion integration index and the width of the

direction tuning curve, as measured using responses to standard stimuli of drifting dots moving

coherently in a single direction (Fig. 4E; Pearson's $r = -0.08$, $p = 0.38$ for low contrast, $r = 0.05$,

$p = 0.57$ for high contrast).

Figure 4. Direction integration with plaid stimuli. **A,** Illustration of the grating (left) and plaid stimuli (right). **B,** Direction tuning curve for an example neuron in response to drifting gratings. **C,** Direction tuning curve for the same neuron in response to moving plaids. The dashed line indicates the component prediction, which is the expected result if the neuron fails to integrate the motion of the plaid. **D,** Population data for motion integration. Scatter plot of the pattern index against the directional integration index (II$_D$) at low contrast ($r = 0.33$, $p = 0.01$). **E,** Scatter plot of the direction tuning width against the directional integration index (II$_D$) at high contrast ($r = -0.08$, $p = 0.38$).

GABAergic influence on neuronal direction tuning and surround suppression in the spatial domain

Another prediction of the SSN is that local changes in the strength of inhibition should have little or no effect on surround suppression, because surround suppression is a result of withdrawal of network excitation (as well as inhibition), and a local blockade of inhibition will not change these network dynamics (Ozeki et al., 2009). This is different from conventional models, which posit that suppression is induced by an increase in the inhibition that a cell receives, so that a reduction in the inhibition to a given neuron will reduce its surround suppression (Tsui and Pack, 2011). Previous work has confirmed the SSN predictions in anesthetized cat V1, using iontophoretic injection of GABA antagonists: inhibitory blockade did not reduce surround suppression (Ozeki et al., 2004). In this section, we examine the effects of pharmacological manipulation of GABA in MT of awake monkeys.

We first confirmed that gabazine, a $GABA_A$ receptor antagonist, robustly modulated neuronal firing in MT (Thiele et al., 2012). We measured direction tuning using random-dot stimuli of fixed spatial size, with all dots moving coherently in a single direction (Fig. 5A). We found that injection of gabazine non-specifically increased firing rates across all directions (Fig. 5C), leading to increases in direction tuning width, as found previously (Thiele et al., 2004; Thiele et al., 2012). In contrast, injections of GABA decreased firing rates across all directions (Fig. 5E), leading to narrower tuning (Leventhal et al., 2003). Figure 6A summarizes the influence of gabazine on direction tuning widths for a population of 38 MT cells: Tuning width significantly increased following the injection (rank sum test, $p = 0.04$). We did not have enough data from the GABA experiments to perform statistical analyses, but in all 5 experiments,

direction tuning width decreased following injection. Overall these results show that local

manipulations of GABA concentration had the expected effects on direction tuning in MT.

To test the influence of GABA concentrations on surround suppression, we performed

standard (space-domain) measurements of size tuning, using random-dot stimuli (100%

coherence) of different physical extents, with all dots moving in the neuron's preferred direction

(Fig. 5B). Previous work has shown that these stimuli elicit surround suppression in the upper

and lower layers in MT, but not in layer 4, suggesting that the suppression is generated through

intrinsic connections within MT (Born and Tootell, 1992; Raiguel et al., 1995). This property

makes such stimuli useful for testing the predicted role of inhibitory inputs in the SSN.

Figure 5D shows size tuning curves from the same MT neuron as in Figure 5C. The pre-

injection data (black line) show that the neuron exhibited substantial surround suppression, as the

response was reduced significantly with increasing stimulus size. As for the direction tuning

curve, injection of gabazine increased firing rates in a non-specific manner. However, in this

neuron there was no apparent reduction in surround suppression (Fig. 5D), and this result was

generally true for the MT population (n = 38): The size suppression index (SI_S), defined as the

difference between the peak response and the response to the largest stimulus divided by the

peak response, was similar before and after injection of gabazine (Fig. 6B; rank sum test, $p =$

0.98). These results are similar to those found in V1 of anesthetized cats (Ozeki et al., 2004),

despite the much larger volume of gabazine used here. In a smaller sample (n = 5), we found that

injection of GABA did not increase surround suppression, despite a strong overall reduction in

firing rate (Fig. 5F).

Figure 5. Effect of GABA on motion direction and size tuning. **A and B,** 100% coherent random

dot patches were used to probe the direction and size tuning of MT neurons. **C and E,** Direction

tuning curve for an example neuron before (black) and after injection of gabazine (C, red) or

GABA (E, blue). The points are the mean responses for each direction. The lines indicate

Gaussian function fits. Direction tuning width (DW) was defined as full width at half maximum

of the fit. **D and F,** Size tuning curves for an example neuron, plotting the firing rate (mean ±

s.e.m.) as a function of patch size before (black) and after injection of gabazine (D, red) or

GABA (F, blue). The lines indicate difference of error functions fits. The horizontal lines show

the spontaneous firing rate.

524 **Figure 6.** Population data on the effects of gabazine on direction and size tuning. **A,** Scatter plot

525 of the direction tuning width before the injection of gabazine against the tuning width after

526 injection (rank sum test, $p = 0.04$). Red and black lines represent the medians of the respective

527 marginal distributions. **B,** Scatter plot of the neuronal size suppression index (SI_S) before the

528 injection of gabazine against the neuronal SI_S after injection (rank sum test, $p = 0.98$).

Discussion

Through electrophysiological recordings in awake monkeys, we have found contrast-dependent

surround suppression in MT in a space defined by motion directions. In addition, we found that

local manipulation of the efficacy of GABAergic inhibition had little influence on standard

measures of surround suppression. Both results are consistent with predictions of the stabilized

supralinear network (SSN), previously tested in V1 (Rubin et al., 2015).

SSN as a unifying motif for normalization in multiple cortical areas

The contrast dependence of surround suppression in the space domain has been observed in both

V1 and MT (Polat et al., 1998; Kapadia et al., 1999; Sceniak et al., 1999; Pack et al., 2005;

Schwabe et al., 2010; Tsui and Pack, 2011). These results have previously been modeled under

the assumption that inhibitory neurons have higher contrast thresholds than excitatory neurons

(Somers et al., 1998; Huang et al., 2008; Schwabe et al., 2010; Carandini and Heeger, 2012).

However, there is little experimental support for this assumption, and some data that contradict it

(Contreras and Palmer, 2003; Song and Li, 2008).

In the SSN, the excitatory and inhibitory units can have the same properties (Rubin et al.,

2015). Each unit has a power-law input/output function, but is stabilized by network inhibition

(Ozeki et al., 2009; Ahmadian et al., 2013; Rubin et al., 2015). With low contrast inputs, the

recurrent interactions within the network are weak, so neurons act relatively independently,

summing their feedforward inputs and responding according to their transfer functions. With

higher-contrast inputs, strong recurrent connections within the network provide contrast- and

size-dependent suppression, with size in the spatial and feature (direction) domains playing

similar roles.

The SSN also predicts that the local blockade of GABA$_A$ receptors should not reduce

surround suppression (Ozeki et al., 2009). In the SSN, surround suppression is not a result of an

increase in inhibitory GABAergic input, but a withdrawal of both excitation and inhibition. In

contrast, in models in which surround suppression results from an increase in the inhibition

received by suppressed neurons (e.g., Tsui and Pack, 2011), local blockade of inhibition should

reduce or prevent surround suppression.

Modulatory influences in visual cortex are often modeled within the normalization

framework, which is hypothesized to be a generic computation with equal validity across brain

regions and stimulus modalities (Carandini et al., 1997; Reynolds and Heeger, 2009; Carandini

and Heeger, 2012; Krause and Pack, 2014). The normalization model as typically conceived, is a

phenomenological rather than circuit model, in which some form of unnormalized neuronal

response is suppressed by the sum of unnormalized responses in other neurons that constitute the

"normalization pool". The precise form of normalization, for example whether the normalizing

pool constitutes all neurons or is restricted in some way based on neuronal tuning, must be

matched to fit the particular experiments modeled.

The SSN can be regarded as a circuit instantiation of the normalization model, in that

many SSN results closely match the results of an appropriately constructed normalization model

(Rubin et al., 2015). In the circuit implementation, the form of normalization is determined by

the connectivity. For example, in the SSN, orientation-specific long-range horizontal

connectivity leads to the orientation-selectivity of surround suppression (Rubin et al., 2015); in a

normalization model, this would be explained by assuming that the normalization pool consists

of neurons of similar preferred orientations to the normalized cell. The normalization model does

not explain the mechanism of suppression, and alternative mechanisms yield different

predictions. For example, if the normalization pool exerted suppression by adding inhibition to

the normalized cells, then one would expect increased inhibition and increased conductance in

normalized (e.g., surround-suppressed) cells, and local GABAergic blockade would reduce or

eliminate the normalization. In the SSN mechanism, normalization typically results from a

decrease in both excitation and inhibition and thus a decreased conductance (Rubin et al., 2015).

Relationship to motion integration in MT

In MT, the integration of different motion directions has frequently been probed with the plaid

stimuli (Movshon et al., 1985; Smith et al., 2005), comprised of superimposed gratings moving

in different directions. Previous work has distinguished between pattern cells, which respond to

the plaid motion direction, and component cells, which respond to the individual grating motion

directions (Movshon et al., 1985).

In the terminology used here, a plaid stimulus moving in a neuron's preferred direction

entails component motion confined to the directional surround. Thus for a high-contrast plaid,

the component gratings should suppress the neuron's response, and this could contribute to the

observed responses of component neurons. Furthermore, component-selective neurons have

small direction centers (i.e. narrow tuning width), so that they do not integrate input from two

gratings moving in very different directions (Rust et al., 2006; Tsui et al., 2010; Khawaja et al.,

2013).

Pattern cells have broader direction tuning than component cells (Rust et al., 2006;

Khawaja et al., 2013). Direction tuning, measured from the responses to individual motion

directions, corresponds to the "minimal response field" in visual space, the region in which small

stimuli can activate the cell; this measure does not change with contrast (Song and Li, 2008). Our

measure of motion integration is not correlated with direction tuning width (Fig. 4E), and is best

related to the "summation field size" in visual space, the size of a stimulus that best drives a cell

before further size increases cause surround suppression. The summation field size, like our

measure of motion integration, shrinks with contrast (Sceniak et al., 1999). We found a weak

correlation between our motion integration index and the pattern index, which quantifies

integration of plaid stimuli (Fig. 4D). These results suggest that the motion-domain summation

field and pattern selectivity are linked, but that summation on its own is insufficient to account

for pattern selectivity.

Pattern cells also show stronger suppression than component cells by stimuli moving

opposite to their preferred directions (Rust et al., 2006). This suggests a direction-domain

analogue of the "far surround" suppression that is found in the space domain; such suppression is

also regulated by contrast both in the direction domain in MT (Pack et al., 2005) and in spatial

surrounds in V1 (Schwabe et al., 2010). Our stimuli did not contain null-direction motion, and so

they would not have probed this component of the MT receptive fields. Nevertheless, an

inference from the existing data is that pattern cells in MT have both larger directional

summation fields and larger (or stronger) directional surrounds.

It can be argued that random-dot stimuli are larger than gratings in the direction domain,

as they activate a broader range of columns in V1 (Simoncelli and Heeger, 1998). Thus stimuli

composed of multiple dots fields moving in different directions might elicit stronger suppression

than grating stimuli containing a similar number of directions. Evidence in support of this idea

comes from studies that use transparent motion stimuli, comprised of overlapping dot fields

moving in two different directions. These stimuli evoke responses in MT that seem to reflect a

suppression of responses to stimuli that straddle the preferred direction (Xiao and Huang, 2015),

particularly for pattern cells (McDonald et al., 2014). One prediction of the current work is that

such suppression should be weaker for low-contrast stimuli.

Functional correlates of integration and suppression

A number of psychophysical studies have drawn a close link between contrast-dependent

responses in MT and visual motion perception. For simple motion discrimination tasks,

performance mirrors spatial processing in MT: for high-contrast stimuli, performance is worse

for large than for small stimuli (Tadin et al., 2003; Liu et al., 2016). Similarly, motion perception

can decrease at high contrasts when the stimulus speed is low, mirroring the contrast-dependent

suppression found in MT (Pack et al., 2005; Seitz et al., 2008). In the direction domain, MT

neurons exhibit higher null-direction suppression when the stimulus is high in contrast (Pack et

al., 2005). This suggests further that suppressive influences are stronger for high-contrast stimuli,

and there is some evidence that motion perception can worsen as the size of the stimulus

increases in the direction domain (Treue et al., 2000; Dakin et al., 2005). Conversely, motion

discrimination with noisy dots can sometimes improve at low contrast (Tadin et al., 2003). Our

results predict the ability to integrate motion signals in the direction domain should

systematically improve at low contrast, as has been found with manipulations of stimulus speed

(Seitz et al., 2008) and spatial size (Tadin et al., 2003).

Conclusion

A growing body of evidence points to a set of generic computations that are similar across brain

regions (Creutzfeldt, 1977; Barlow, 1985; Miller, 2016) and across sensory modalities

(Mountcastle, 1978; Pack and Bensmaia, 2015). Although this idea is attractive from a

theoretical standpoint, it remains somewhat speculative. In this work, we have provided an

experimental test of the genericity of one computational model by comparing results in MT with

those obtained previously in V1. The qualitative pattern of results is similar, supporting the

possibility that this model provides a more general framework for modulatory responses and

integration in cortex.

References

Ahmadian Y, Rubin DB, Miller KD (2013) Analysis of the stabilized supralinear network.

Neural Comput 25:1994-2037.

Albright TD (1984) Direction and orientation selectivity of neurons in visual area MT of the

macaque. J Neurophysiol 52:1106-1130.

Allman J, Miezin F, McGuinness E (1985) Stimulus specific responses from beyond the classical

receptive field: neurophysiological mechanisms for local-global comparisons in visual neurons.

Annu Rev Neurosci 8:407-430.

Angelucci A, Bullier J (2003) Reaching beyond the classical receptive field of V1 neurons:

horizontal or feedback axons? J Physiol Paris 97:141-154.

Angelucci A, Bressloff PC (2006) Contribution of feedforward, lateral and feedback connections

to the classical receptive field center and extra-classical receptive field surround of primate V1

neurons. Prog Brain Res 154:93-120.

Barlow HB (1985) The twelfth Bartlett memorial lecture: the role of single neurons in the

psychology of perception. Q J Exp Psychol A 37:121-145.

Bolz J, Gilbert CD (1986) Generation of end-inhibition in the visual cortex via interlaminar

connections. Nature 320:362-365.

Born RT, Tootell RB (1992) Segregation of global and local motion processing in primate

middle temporal visual area. Nature 357:497-499.

Born RT, Bradley DC (2005) Structure and function of visual area MT. Annu Rev Neurosci

28:157-189.

Carandini M, Heeger DJ (2012) Normalization as a canonical neural computation. Nat Rev

Neurosci 13:51-62.

Carandini M, Heeger DJ, Movshon JA (1997) Linearity and normalization in simple cells of the

macaque primary visual cortex. Journal of Neuroscience 17:8621.

Cavanaugh JR, Bair W, Movshon JA (2002) Nature and interaction of signals from the receptive

field center and surround in macaque V1 neurons. J Neurophysiol 88:2530-2546.

Chagnac-Amitai Y, Connors BW (1989) Horizontal spread of synchronized activity in neocortex

and its control by GABA-mediated inhibition. J Neurophysiol 61:747-758.

Contreras D, Palmer L (2003) Response to contrast of electrophysiologically defined cell classes

in primary visual cortex. J Neurosci 23:6936-6945.

Cossell L, Iacaruso MF, Muir DR, Houlton R, Sader EN, Ko H, Hofer SB, Mrsic-Flogel TD

(2015) Functional organization of excitatory synaptic strength in primary visual cortex. Nature

518:399-403.

Creutzfeldt OD (1977) Generality of the functional structure of the neocortex.

Naturwissenschaften 64:507-517.

Dakin SC, Mareschal I, Bex PJ (2005) Local and global limitations on direction integration

assessed using equivalent noise analysis. Vision Res 45:3027-3049.

DeAngelis GC, Uka T (2003) Coding of horizontal disparity and velocity by MT neurons in the

alert macaque. J Neurophysiol 89:1094-1111.

Huang X, Albright TD, Stoner GR (2008) Stimulus dependency and mechanisms of surround

modulation in cortical area MT. J Neurosci 28:13889-13906.

Hunter JN, Born RT (2011) Stimulus-dependent modulation of suppressive influences in MT. J

Neurosci 31:678-686.

Jones HE, Grieve KL, Wang W, Sillito AM (2001) Surround suppression in primate V1. J

Neurophysiol 86:2011-2028.

Kapadia MK, Westheimer G, Gilbert CD (1999) Dynamics of spatial summation in primary

visual cortex of alert monkeys. Proc Natl Acad Sci U S A 96:12073-12078.

Khawaja FA, Liu LD, Pack CC (2013) Responses of MST neurons to plaid stimuli. J

Neurophysiol 110:63-74.

Krause MR, Pack CC (2014) Contextual modulation and stimulus selectivity in extrastriate

cortex. Vision Res 104:36-46.

Leventhal AG, Wang Y, Pu M, Zhou Y, Ma Y (2003) GABA and its agonists improved visual

cortical function in senescent monkeys. Science 300:812-815.

Levitt JB, Lund JS (1997) Contrast dependence of contextual effects in primate visual cortex.

Nature 387:73-76.

Liu LD, Pack CC (2017) The Contribution of Area MT to Visual Motion Perception Depends on

Training. Neuron 95:436-446 e433.

Liu LD, Haefner RM, Pack CC (2016) A neural basis for the spatial suppression of visual motion

perception. Elife 5.

McDonald JS, Clifford CW, Solomon SS, Chen SC, Solomon SG (2014) Integration and

segregation of multiple motion signals by neurons in area MT of primate. J Neurophysiol

111:369-378.

Miller KD (2016) Canonical computations of cerebral cortex. Curr Opin Neurobiol 37:75-84.

Mountcastle VB (1978) Brain mechanisms for directed attention. J R Soc Med 71:14-28.

Movshon JA, Adelson EH, Gizzi MS, Newsome WT (1985) The analysis of visual moving

patterns. In: Pattern Recognition Mechanisms (Chagas C, Gattass R, Gross C, eds), pp 117-151.

New York: Springer.

Nealey TA, Maunsell JH (1994) Magnocellular and parvocellular contributions to the responses

of neurons in macaque striate cortex. J Neurosci 14:2069-2079.

Ozeki H, Finn IM, Schaffer ES, Miller KD, Ferster D (2009) Inhibitory stabilization of the

cortical network underlies visual surround suppression. Neuron 62:578-592.

Ozeki H, Sadakane O, Akasaki T, Naito T, Shimegi S, Sato H (2004) Relationship between

excitation and inhibition underlying size tuning and contextual response modulation in the cat

primary visual cortex. J Neurosci 24:1428-1438.

Pack CC, Bensmaia SJ (2015) Seeing and Feeling Motion: Canonical Computations in Vision

and Touch. PLoS Biol 13:e1002271.

Pack CC, Hunter JN, Born RT (2005) Contrast dependence of suppressive influences in cortical

area MT of alert macaque. J Neurophysiol 93:1809-1815.

Polat U, Mizobe K, Pettet MW, Kasamatsu T, Norcia AM (1998) Collinear stimuli regulate

visual responses depending on cell's contrast threshold. Nature 391:580-584.

Raiguel S, Van Hulle MM, Xiao DK, Marcar VL, Orban GA (1995) Shape and spatial

distribution of receptive fields and antagonistic motion surrounds in the middle temporal area

(V5) of the macaque. Eur J Neurosci 7:2064-2082.

Reynolds JH, Heeger DJ (2009) The normalization model of attention. Neuron 61:168-185.

Rubin DB, Van Hooser SD, Miller KD (2015) The stabilized supralinear network: a unifying

circuit motif underlying multi-input integration in sensory cortex. Neuron 85:402-417.

Rust NC, Mante V, Simoncelli EP, Movshon JA (2006) How MT cells analyze the motion of

visual patterns. Nat Neurosci 9:1421-1431.

Sceniak MP, Ringach DL, Hawken MJ, Shapley R (1999) Contrast's effect on spatial summation

by macaque V1 neurons. Nat Neurosci 2:733-739.

Schwabe L, Ichida JM, Shushruth S, Mangapathy P, Angelucci A (2010) Contrast-dependence of

surround suppression in Macaque V1: experimental testing of a recurrent network model.

Neuroimage 52:777-792.

Seitz AR, Pilly PK, Pack CC (2008) Interactions between contrast and spatial displacement in

visual motion processing. Curr Biol 18:R904-906.

Simoncelli EP, Heeger DJ (1998) A model of neuronal responses in visual area MT. Vision Res

38:743-761.

Smith MA, Majaj NJ, Movshon JA (2005) Dynamics of motion signaling by neurons in macaque

area MT. Nat Neurosci 8:220-228.

Snowden RJ, Treue S, Andersen RA (1992) The response of neurons in areas V1 and MT of the

alert rhesus monkey to moving random dot patterns. Exp Brain Res 88:389-400.

Somers DC, Todorov EV, Siapas AG, Toth LJ, Kim DS, Sur M (1998) A local circuit approach

to understanding integration of long-range inputs in primary visual cortex. Cereb Cortex 8:204-

217.

Song XM, Li CY (2008) Contrast-dependent and contrast-independent spatial summation of

primary visual cortical neurons of the cat. Cereb Cortex 18:331-336.

Tadin D, Lappin JS, Gilroy LA, Blake R (2003) Perceptual consequences of centre-surround

antagonism in visual motion processing. Nature 424:312-315.

Thiele A, Distler C, Korbmacher H, Hoffmann KP (2004) Contribution of inhibitory

mechanisms to direction selectivity and response normalization in macaque middle temporal

area. Proc Natl Acad Sci U S A 101:9810-9815.

Thiele A, Herrero JL, Distler C, Hoffmann KP (2012) Contribution of cholinergic and

GABAergic mechanisms to direction tuning, discriminability, response reliability, and neuronal

rate correlations in macaque middle temporal area. J Neurosci 32:16602-16615.

Treue S, Hol K, Rauber HJ (2000) Seeing multiple directions of motion-physiology and

psychophysics. Nat Neurosci 3:270-276.

Tsodyks MV, Skaggs WE, Sejnowski TJ, McNaughton BL (1997) Paradoxical effects of

external modulation of inhibitory interneurons. J Neurosci 17:4382-4388.

Tsui JM, Pack CC (2011) Contrast sensitivity of MT receptive field centers and surrounds. J

Neurophysiol 106:1888-1900.

Tsui JM, Hunter JN, Born RT, Pack CC (2010) The role of V1 surround suppression in MT

motion integration. J Neurophysiol 103:3123-3138.

Xiao J, Huang X (2015) Distributed and Dynamic Neural Encoding of Multiple Motion

Directions of Transparently Moving Stimuli in Cortical Area MT. J Neurosci 35:16180-16198.

775