Parallel evolution of gene expression between trophic specialists despite divergent genotypes and morphologies

Joseph A. McGirr[1], Christopher H. Martin[1,]

Article Type: Letter

[1]Department of Biology, University of North Carolina at Chapel Hill, 1250 South Rd., Chapel Hill, NC 27514-3280

Running Title: Parallel evolution in divergent species

Keywords: RNA-seq, adaptive radiation, hard sweep, genomics, development, parallel evolution

Correspondence: Joseph A. McGirr. Department of Biology, University of North Carolina at Chapel Hill, Genome Science 2244, 250 Bell Tower Drive., NC, 27514, USA

Email: jmcgirr@email.unc.edu

Word count abstract: 204 words; Word count main text: 6,823; Figures: 3, Tables: 1.

Abstract

Parallel evolution of gene expression commonly underlies convergent niche specialization, but

parallel changes in expression could also underlie divergent specialization. We investigated

divergence in gene expression and whole-genome genetic variation across three sympatric

Cyprinodon pupfishes endemic to San Salvador Island, Bahamas. This recent radiation consists

of a generalist and two derived specialists adapted to novel niches – a 'scale-eater' and a 'snail-

eater.' We sampled total mRNA from all three species at two early developmental stages and

compared gene expression with whole-genome genetic differentiation among all three species in

42 resequenced genomes. 80% of genes that were differentially expressed between snail-eaters

and generalists were up or downregulated in the same direction between scale-eaters and

generalists; however, there were no fixed variants shared between species underlying these

parallel changes in expression. Genes showing parallel evolution of expression were enriched for

effects on metabolic processes, whereas genes showing divergent expression were enriched for

effects on cranial skeleton development and pigment biosynthesis, reflecting the most divergent

phenotypes observed between specialist species. Our findings reveal that even divergent niche

specialists may exhibit convergent adaptation to higher trophic levels through shared genetic

pathways. This counterintuitive result suggests that parallel evolution in gene expression can

accompany divergent ecological speciation during adaptive radiation.

Impact Summary

Adaptations that result in unique forms of ecological specialization are central to research in

evolutionary biology, yet little is known about their molecular foundations. We combined

transcriptome sequencing with whole-genome divergence scans to study the molecular evolution

of two specialist *Cyprinodon* pupfish species – a 'scale-eater' and a 'snail-eater' – that rapidly

diverged from a sympatric generalist ancestor within the last 10,000 years. While parallel

evolution of gene expression driving convergent niche specialization seems common, we

present, to our knowledge, the first example of significant parallel changes in expression

coinciding with divergent niche specialization. 80% of genes that were differentially expressed

between snail-eaters and generalists showed the same direction of expression in scale-eaters

relative to generalists. Furthermore, parallel evolution in expression seem to be controlled by

unique genetic variants in each specialist species. Genes showing parallel changes in expression

were enriched for metabolic processes that may facilitate adaptation to a higher trophic level,

while genes showing divergent expression likely shape the striking morphological differences

between specialists. These findings contribute to a more nuanced understanding of convergent

adaptations that arise during speciation, and highlight how species can evolve similar expression

profiles adapted to divergent niches.

Introduction

Abundant research on the genetic basis of adaptive traits has revealed an overarching pattern in nature – when species are faced with similar selective pressures, they often respond with the same adaptive solutions (Conte et al. 2012). For example, parallel changes in gene expression underlying convergent adaptive traits are a well-documented evolutionary phenomenon, with examples from experimental evolution studies imposing uniform selection pressures on replicate populations (Cooper et al. 2003; Riehle et al. 2003), studies in natural systems between closely related taxa (Reid et al. 2016; Derome and Bernatchez 2006; Chan et al. 2010; Nagai et al. 2011; Reed et al. 2011; Manousaki et al. 2013; Zhao et al. 2015), and distantly related taxa (Shapiro et al. 2006; Miller et al. 2007a; Shen et al. 2012). This work has shown that parallelism at the level of gene expression is common in many cases of phenotypic convergence, particularly when divergence time between species is short (Losos 2011; Conte et al. 2012).

However, few studies have investigated the extent of parallel changes in gene expression contributing to species divergence, largely because most expression studies focus on only two species and are either concerned with divergent expression giving rise to divergent phenotypes (Poelstra et al. 2014; Uebbing et al. 2016; Davidson and Balakrishnan 2016) or parallel expression of specific loci (Shapiro et al. 2006; Miller et al. 2007a; Quin et al. 2010) (but see Enard et al. 2002; Ahi et al. 2014). Furthermore, while many genetic and demographic factors are thought to influence the probability of parallel evolution (Rosenblum et al. 2014.; Conte et al. 2012), there are no theoretical expectations for the amount of parallel genetic variation contributing to parallel changes in gene expression during ecological speciation (Schluter et al. 2004; Pavey et al. 2010).

Here we ask whether both parallel and divergent changes in expression underlie novel phenotypes by measuring transcriptomic and genomic divergence between three sympatric species of *Cyprinodon* pupfishes endemic to hypersaline lakes on San Salvador Island, Bahamas. This recent radiation consists of a dietary generalist species (*C. variegatus*) and two novel specialists: a 'snail-eater' (*C. brontotheroides*) and a 'scale-eater' (*C. desquamator*). Scale-eaters have large jaws and elongated bodies, whereas snail-eaters have short, thick jaws and a protruding nasal region that may function in crushing hard-shelled mollusks. These specialists are more morphologically diverged from one another than either is from their sympatric

generalist sister species, and occupy higher trophic levels than the generalist species (Martin and

Wainwright 2013a; Martin 2016a, Martin et al. 2017, Hernandez et al. 2017). Scale-eating and

snail-eating rapidly evolved within saline lakes on San Salvador Island, Bahamas. These lakes

filled within the past 10,000 years after the last glacial maximum (Mylroie and Hagey 1995;

Turner et al. 2008), suggesting that speciation occurred rapidly. Scale-eaters and snail-eaters

have only been found on San Salvador, and likely diverged from a generalist common ancestor

based on phylogenetic analyses of outgroup species (Martin and Wainwright 2011; Martin and

Wainwright 2016b). Pupfish populations on many neighboring Bahamian islands and throughout

the Caribbean are dietary generalists (Martin and Wainwright 2011; Martin 2016a) and these

specialist niches appear unique within atherinomorph and cyprinodontiform fishes (Martin and

Wainwright 2011). Indeed, the scale-eating pupfish is separated by 168 million years from other

scale-eating fishes (Martin and Wainwright 2013b).

We performed total mRNA sequencing to examine gene expression in lab-reared

individuals of all three San Salvador pupfish species from different lake populations at two

developmental stages. We also searched 42 whole genomes for SNPs unique to each specialist

and determined whether fixed variants near differentially expressed genes showed signs of hard

selective sweeps (Pavlidis et al. 2013). We found significant parallelism at the level of gene

expression in specialists, but did not find evidence for shared fixed variants underlying parallel

changes in expression. We tested whether this counterintuitive result of parallel changes in

expression between divergent trophic specialists may be due to 1) decreased pleiotropic

constraint for genes showing parallelism or that 2) specialists experience parallel selective

environments and adapted to higher trophic levels using similar genetic pathways. Finally, we

identified genes differentially expressed between generalists and scale-eaters that contain fixed

genetic variants within regions that were previously associated with jaw size variation and

showed signs of experiencing a recent hard selective sweep (McGirr and Martin 2017). These

regions with fixed variants represent promising *cis*-regulatory elements underlying divergent jaw

size – the most rapidly diversifying trait in the San Salvador pupfish radiation (Martin and

Wainwright 2013b).

Methods

Study system and sample collection

Individuals were caught from hypersaline lakes on San Salvador Island, Bahamas using a hand

net or seine net in 2011, 2013, and 2015. Whole genome resequencing was performed for wild-

caught individuals from a total of nine isolated lakes on San Salvador (Great Lake, Stout's Lake,

Oyster Lake, Little Lake, Crescent Pond, Moon Rock, Mermaid's Pond, Osprey Lake, and

Pigeon Creek). 14 scale-eaters were sampled from six populations; 11 snail-eaters were sampled

from four populations; and 13 generalists were sampled from eight populations on San Salvador.

Outgroup samples included one *C. laciniatus* from Lake Cunningham, New Providence Island,

Bahamas, one *C. bondi* from Etang Saumautre lake in the Dominican Republic, one *C. diabolis*

from Devil's Hole in California, and captive-bred individuals of *C. simus* and *C. maya* from

Laguna Chicancanab, Quintana Roo, Mexico. Sampling is further described in (McGirr and

Martin 2017; Richards and Martin 2017). Fish were euthanized in an overdose of buffered MS-

222 (Finquel, Inc.) following approved protocols from the University of California, Davis

Institutional Animal Care and Use Committee (#17455) and University of California, Berkeley

Animal Care and Use Committee (AUP-2015-01-7053) and stored in 95-100% ethanol.

RNA sequencing and alignment

Juvenile pupfish were derived from either F_0 wild caught or F_1 lab raised individuals that were

held in a common laboratory environment and fed identical diets (Table S1; 25-27^o C, 10-15 ppt

salinity, pH 8.3). We collected larvae at two developmental stages: 8-10 and 17-20 days post-

fertilization (dpf). The variation in sampling time is due to uncertainty in precise spawning times

since eggs were fertilized naturally within breeding tanks and collected on the same day or

subsequent day following egg laying. However, we sampled hatched larvae in a haphazard

manner over multiple spawning intervals and it is unlikely that sampling time varied consistently

by species. Larvae were euthanized in an overdose of buffered MS-222, and stored in RNA later

(Ambion, Inc.) at 4^o C for one day, followed by long-term storage at -20^o C for up to one year.

We extracted whole-larvae RNA using RNeasy kits (Qiagen) from 15 larvae (8-10 dpf) (Three F_2

generalists and F_2 snail-eaters from Crescent Pond, three F_1 generalists and F_2 snail-eaters from

Little Lake, and three F_1 scale-eaters from Little Lake; Table S1). We also dissected 14 larvae

(17-20 dpf) to isolate tissues from the anterior craniofacial region containing the dentary, angular

articular, maxilla, premaxilla, palatine, and associated craniofacial connective tissues using fine-

tipped tweezers washed with RNase AWAY (Three F_2 generalists and F_2 snail-eaters from

Crescent Pond, three F_1 generalists and F_2 snail-eaters from Little Lake, and two F_1 scale-eaters

from Little Lake; Table S1).

Libraries were prepared using the KAPA stranded mRNA-seq kit (KAPA Biosystems

2016) at the High Throughput Genomic Sequencing Facility at UNC Chapel Hill. Stranded

sequencing on one lane of Illumina 150PE Hiseq4000 resulted in 677 million raw reads. We

filtered raw reads using Trim Galore (v. 4.4, Babraham Bioinformatics) to remove Illumina

adaptors and low-quality reads (mean Phred score < 20). We mapped these reads to the

Cyprinodon reference genome using the RNA-seq aligner STAR (v. 2.5 (Dobin et al. 2013)). We

used the featureCounts function of the Rsubread package (Liao et al. 2014) requiring paired-end

and reverse stranded options to generate read counts across previously annotated features. We

assessed mapping and count quality using MultiQC (Ewels et al. 2016).

Differential expression analyses

We quantified differences in gene expression between all three species at two developmental

stages. Our raw counts determined by featureCounts were normalized with DESeq2 (v. 3.5 (Love

et al. 2014)) which uses counts to calculate a geometric mean for each gene across samples,

divides individual gene counts by this mean, and uses the median of these ratios as a size factor

for each sample. Next, we used DESeq2 to perform pairwise tests pooling species across lakes to

identify differentially expressed genes between generalists *vs.* snail-eaters and generalists *vs.*

scale-eaters at 8-10 dpf and 17-20 dpf (Table. S1). Genes with fewer than two read counts were

discarded from all analyses (n = 1,570), along with genes showing low normalized counts at a

threshold determined by DESeq2 (Love et al. 2014). Wald tests determined significant

differences in expression between species by comparing normalized posterior log fold change

estimates and correcting for multiple testing using the Benjamini–Hochberg procedure with a

false discovery rate of 0.05 (Benjamini and Hochberg 1995).

We performed two analyses to test whether specialist species exhibited nonrandom

patterns of parallel changes in expression relative to their generalist sister species. We used a

Fisher's exact test to determine whether there was a significant overlap between genes that

showed differential expression in both comparisons (i.e. genes that were differentially expressed

between generalists *vs.* snail-eaters and generalists *vs.* scale-eaters). A gene that was

differentially expressed in both comparisons could either show the same direction of expression

in specialists or opposite directions of expression. We performed 10,000 permutations sampling

from a binomial distribution to estimate the expected number of genes showing shared and

opposite directions of expression. Under this null model of gene expression evolution, a strong

positive deviation from 50% of genes showing a shared direction of expression in specialists

would indicate significant parallel changes in expression.

Our scale-eater sample sizes were lower than generalist and snail-eater samples for each

pairwise comparison (see above). We used a down sampling procedure to test whether sample

size affected patterns of parallel changes in expression. We analyzed differential expression for

generalists *vs.* snail eaters and generalists *vs.* scale-eaters in 1,000 permutations where

generalists and snail-eaters were randomly sampled from our full dataset to match scale-eater

sample sizes (n = 3 for 8-10 dpf comparisons; n = 2 for 17-20 dpf). Next, we identified the

number of genes differentially expressed between generalists *vs.* snail eaters and generalists *vs.*

scale-eaters in each permutation and calculated the proportion of those genes that showed the

same direction of expression in specialists relative to generalists. A strong positive deviation

from 50% of genes showing a shared direction of expression across permutations would indicate

that parallel evolution of expression in specialists is robust to variation in sample size.

Gene ontology enrichment analyses

We performed gene ontology (GO) enrichment analyses for differentially expressed genes using

GO Consortium resources available at geneontology.org (Ashburner et al. 2000; GO Consortium

2017). We used BlastP (v. 2.6 (Camacho et al. 2009)) to identify zebrafish protein orthologs with

high similarity (*E*-value < 1) to NCBI protein accessions for genes that we identified as

differentially expressed between *Cyprinodon* species. Orthology was established using one-way

best hits, where a protein sequence in *Cyprinodon* was the best match to a sequence in zebrafish,

and reciprocal best blast hits, where a sequence in *Cyprinodon* was the best match to a sequence

in zebrafish and vice versa. While reciprocal best hits robustly predict orthology with high

precision, it is highly conservative and fails to detect many true orthologs in duplication rich

clades such as teleosts (Altenhoff and Dessimoz 2009; Salichos and Rokas 2011; Dalquen and

Dessimoz 2013). Thus, we performed GO enrichment analyses using orthologs defined as one-

way best hits, and compare these results to enrichment analyses using more conservative

orthologs defined as reciprocal best hits.

Genes were either differentially expressed between generalists and snail-eaters,

generalists and scale-eaters, or in both comparisons. Thus, we performed two GO enrichment

analyses for: 1) genes that were differentially expressed in both companions, and 2) genes

differentially expressed in one comparison. We grouped enriched GO categories into similar

representative terms using the REVIGO clustering algorithm (Tomislav 2011). REVIGO groups

semantically similar terms to reduce the size and redundancy of lists from GO enrichment

analyses, where grouping is guided by *P*-values corrected for multiple comparisons (Tomislav

2011). When similar terms show similar enrichment, they are assigned to a single representative

term. We measured differences in the proportion of representative terms describing metabolic

and developmental processes between genes showing parallel and divergent changes in

expression between specialists.

Measuring pleiotropy for differentially expressed genes

The probability of parallel evolution of gene expression may be higher for genes that are less

constrained by pleiotropy (Cooper and Lenski 2000; Manceau et al. 2010; Rosenblum et al.

2014). High gene pleiotropy is correlated with participation in more protein-protein interactions

(PPIs), which in turn effects multiple biological processes (He and Zhang 2006; Safari-

alighiarloo et al. 2014). Genes that act across multiple developmental stages are also more

pleiotropic (Stern and Orgogozo 2008). We used one-way best hits zebrafish orthologs to

estimate pleiotropy for differentially expressed genes based on their number of associated GO

biological processes, PPIs, and developmental stages when they are known to be expressed

(Papakostas et al. 2014). We again used GO Consortium resources (Ashburner et al. 2000; GO

Consortium 2017) to determine the number of biological processes associated with each gene.

We examined biological process annotations only for genes from ZFIN (zfin.org) with

experimental evidence (GO evidence code EXP). The String protein database (v. 10; (Szklarczyk

et al. 2015)) calculates a combined score measuring confidence in protein interactions by

considering known interactions (experimentally determined and from manually curated

databases) and predicted interactions. We used the String database to quantify PPIs for protein

products of differentially expressed genes, focusing only on interactions with experimental

evidence (i.e. non-zero experimental evidence scores). Next, we determined the number of

developmental stages where a gene is known to be expressed using the Bgee expression call

database for zebrafish (v. 14.0 (Bastian et al. 2008)). We considered eight developmental stages

from larval day five to juvenile day 89 from the Zebrafish Stage Ontology (ZFS) that were

deemed 'gold quality,' meaning there was no contradicting call of absence of expression for the

same gene, in the same developmental stage (Bastian et al. 2008).

We tested whether genes showing parallel changes in expression between specialists

showed lower levels of pleiotropy than genes showing divergent changes in expression by fitting

a generalized linear model on count data for pleiotropy estimates (negative binomial family;

glm.nb function in the R library "MASS"). We did not measure pleiotropy for genes expressed at

17-20 dpf due to the low number of zebrafish orthologs matched for genes with parallel

expression in craniofacial tissues (11 out of 23).

Genomic variant discovery and population genetic analyses

SNP variants were called using previously outlined methods (McGirr and Martin 2017; Richards

and Martin 2017). Briefly, 42 individual DNA samples extracted from muscle tissue were

fragmented, barcoded with Illumina indices, and quality checked using a Fragment Analyzer

(Advanced Analytical Technologies, Inc.). Sequencing on four lanes of Illumina 150PE

Hiseq4000 resulted in 2.8 billion raw reads that were mapped from 42 individuals to the

Cyprinodon reference genome (NCBI, *C. variegatus* Annotation Release 100, total sequence

length = 1,035,184,475; number of scaffold = 9,259, scaffold N50, = 835,301; contig N50 =

20,803; (Lencer et al. 2017)). We followed Genome Analysis Toolkit (v 3.5) best practices and

hard filter criteria to call and refine our SNP variant dataset (QD < 2.0; FS < 60; MQRankSum <

-12.5; ReadPosRankSum < -8 (DePristo et al. 2011)). We filtered our final SNP dataset to

include individuals with a genotyping rate above 90% (no individuals were excluded by this

filter) and SNPs with minor allele frequencies higher than 5%, resulting in 16 million variants

with a mean sequencing coverage of 7× per individual (range: 5.2–9.3×).

We identified SNPs that were fixed in each specialist species. We calculated genome

wide F_{st} using VCFtools' 'weir-fst-pop' function for two different population comparisons

involving samples collected from San Salvador: generalists (n = 13) *vs.* snail-eaters (n = 11) and

generalists (n = 13) *vs.* scale-eaters (n = 9). Differences in sample sizes made our analyses biased

to detect more fixed variation between generalists vs. scale-eaters (n = 13 vs. 9) than between

generalists vs. snail-eaters (n = 13 vs. 11). We also performed 1,000 permutations calculating

genome wide F_{st} between randomly subsampled groups in order to identify non-randomly

differentiated genomic regions between species. We calculated the 99th percentile estimates of

F_{st} across all SNPs between randomly sampled generalists and snail-eaters (n = 13 vs. n = 11)

and between randomly sampled generalists and scale-eaters (n = 13 vs. n = 9). We took the 99th

percentile of these distributions to set a threshold defining significantly divergent outliers (Fig.

S7).

Our SNP dataset included 14 scale-eaters, however, we split our scale-eater population

into two groups (large-jawed scale-eaters, n = 9 and small-jawed scale-eaters, n = 5) based on

previous evidence that these two populations are genetically distinct (McGirr and Martin 2017;

Richards and Martin 2017). This allowed us to identify SNPs unique to large-jawed scale-eaters

(i.e. *C. desquamator* (Martin and Wainwright 2013a)), which were the only type of scale-eater

we sampled for RNA-seq. We identified which of these SNPs resided in gene regions (either

exonic, intronic, or within 10kb of the first or last exon) for genes showing differential

expression. We determined whether these regions showed signatures of hard selective sweeps

using SweeD ((Pavlidis et al. 2013); methods previously described in (McGirr and Martin

2017)). Briefly, SweeD sections scaffolds into 1,000 windows of equal size and calculates a

composite likelihood ratio (CLR) using a null model where the site frequency spectrum of each

window does not differ from that of the entire scaffold. We previously estimated ancestral

effective population sizes of San Salvador pupfishes using MSMC (Schiffels and Durbin 2014;

McGirr and Martin 2017) and used these estimates to correct the expected neutral site frequency

spectrum for the inferred recent population bottleneck in Caribbean pupfishes using SweeD.

Windows with fixed SNPs that showed CLRs above the 95[th] percentile across their respective

scaffolds (>10,000bp) under the assumptions of a recent population bottleneck were interpreted

as regions that recently experienced a hard sweep.

Results

Differential expression between generalists and each specialist

Total mRNA sequencing across all 29 samples resulted in 677 million raw reads, which was

reduced to 674 million reads after quality control and filtering. 81.2% of these reads successfully

aligned to the reference genome and 75.5% of aligned reads mapped to annotated features with

an average read depth of 309× per sample. The number of reads mapping to annotated features

was comparable across generalists, snail-eaters, and scale-eaters (ANOVA; 8-10 dpf $P = 0.47$;

17-20 dpf $P = 0.33$; Fig. S1).

Snail-eaters and scale-eaters occupy novel niches among over 2,000 species of

atherinomorph fishes (Martin and Wainwright 2011), and these trophic specialist species likely

evolved from a generalist common ancestor within the past 10,000 years (Mylroie, J.E, Hagey

1995; Turner et al. 2008). We analyzed transcriptomic changes underlying rapid trophic

divergence by comparing specialist species gene expression against their sympatric generalist

sister species. We used DESeq2 to identify genes that were differentially expressed between

generalists *vs.* snail-eaters and generalists *vs.* scale-eaters at 8-10 days post-fertilization (whole

body tissue) and 17-20 dpf (craniofacial tissue).We measured expression across 22,183 genes

with greater than two read counts out of 24,383 total genes annotated for the *Cyprinodon*

variegatus assembly (NCBI, *C. variegatus* Annotation Release 100, (Lencer et al. 2017)).

At 8-10 dpf, we found 1,014 genes differentially expressed between generalists *vs.* snail-

eaters and 5,982 genes differentially expressed between generalists *vs.* scale-eaters (Fig. 1A and

C; Fig. 2A) (Benjamini and Hochberg adjusted $P \leq 0.05$). 818 genes were differentially

expressed in both comparisons, which is a significantly larger amount of overlap than expected

by chance (Fisher's exact test, $P < 1.0 \times 10^{-16}$). Remarkably, 815 of these 818 genes showed the

same direction of expression in specialists relative to generalists (Fig. 2B). Specifically, 441

differentially expressed genes showed lower expression in both specialist species compared to

generalists, while 374 showed higher expression in specialists. Only three genes showed opposite

directions of expression (Fig. 2B). Two genes showed higher expression in snail-eaters and

lower expression in scale-eaters while one gene showed higher expression in scale-eaters (Table

S2). This is significantly more parallel change in expression between specialists than would be

expected under a null model of gene expression evolution, where a gene has an equal chance of

showing a shared or opposite direction of expression in specialists relative to generalists (10,000

permutations, $P < 1.0 \times 10^{-4}$; Fig. S2). Parallel evolution of expression in specialists was

consistent at both the gene and isoform level (Fig. S2, S3).

Craniofacial morphology is the most rapidly diversifying phenotypic axis measured so far

within the San Salvador radiation (Martin and Wainwright 2013b; Martin 2016a). In order to

detect genes expressed during jaw development, we compared expression within craniofacial

tissue at the 17-20 dpf stage. We found a similar pattern of parallel changes in gene expression at

this developmental stage (Fig. S4). 120 genes were differentially expressed between generalists

vs. snail-eaters and 1,903 genes differentially expressed between generalists *vs.* scale-eaters (Fig.

1B and D). Again, we saw a significant amount of overlap between comparisons with 23 genes

differentially expressed in both comparisons (Fisher's exact test, $P < 1.0 \times 10^{-5}$). 22 of these 23

genes showed the same direction of expression in specialists relative to generalists (Fig. S4).

Specifically, 10 genes showed lower expression in both specialist species compared to

generalists, while 12 showed higher expression in specialists (Fig. S4). Only one gene (*mybpc2*)

showed opposite directions of expression, with higher expression in snail-eaters and lower

expression in scale-eaters (Table S2).

Our sample sizes for scale-eater species were lower for comparisons at 8-10 dpf (n = 3)

and 17-20 dpf (n = 2) relative to snail-eaters and generalists (n = 6 for both stages). We measured

differential expression for generalists *vs.* snail eaters and generalists *vs.* scale-eaters in 1,000

permutations where we randomly down-sampled generalists and snail-eaters from our full

dataset to match scale-eater sample sizes. Figure 2C shows the proportion of genes differentially

expressed at 8-10 dpf in both comparisons that showed the same direction of expression in

specialists relative to generalists across 1,000 permutations. The total number of differentially

expressed genes in each permutation was variable (Fig. S5A and C, median number of genes

common to both comparisons = 61). Despite this variability, we found that the parallel evolution

of expression in specialists was robust to smaller sample size, with greater than 90% of genes

showing parallel evolution of expression in 90% of permutations (Fig. 2C). However, at 17-20

dpf parallel changes in expression were not as consistent across permutations (Fig. S4C and

S5F).

Genes showing parallel changes in expression are enriched for metabolic processes

We performed GO enrichment analyses with one-way blast hit zebrafish orthologs for genes

showing parallel changes in expression between specialists (n = 620) and genes showing

divergent expression patterns in snail-eaters (n = 102) and scale-eaters (n = 3,349). We restricted

these analyses to genes expressed at 8-10 dpf because the number of genes showing parallel

expression in specialists at 17-20 dpf (n = 23) was low and did not show enrichment for any

biological process.

386   We grouped enriched GO categories into similar representative terms using the REVIGO

387   clustering algorithm (Tomislav 2011). Genes showing parallel changes in expression between

388   specialists were enriched for metabolic processes (20% of representative terms; Fig. 3A; Table

389   S3). In contrast, genes with divergent expression patterns in specialists were enriched for cranial

390   skeletal development and pigment biosynthesis (7% and 3% of terms, respectively) while only

391   11% of enriched categories described metabolic processes (Table S4).

392   We also performed GO enrichment analyses using orthologs that were established using a

more conservative reciprocal best hit approach, where a sequence in *Cyprinodon* was the best

match to a sequence in zebrafish and vice versa. As expected, we identified fewer reciprocal

best hits than one-way hits (615 genes showing parallel changes in expression between

specialists, 95 genes showing divergent expression unique to snail-eaters, and 2,150 genes

showing divergent expression unique to scale-eaters). Encouragingly, we still found that genes

showing parallel changes in expression were enriched for metabolic processes (26% of

representative terms), whereas genes showing divergent expression showed less enrichment for

metabolic processes (20% of representative terms). However, we did not see any enrichment for

cranial development or pigment biosynthesis for genes showing divergent expression using

reciprocal best hit orthologs.

We tested whether genes showing parallel changes in expression were less constrained by
pleiotropy than genes showing divergent expression between specialists. We estimated
pleiotropy for orthologs of differentially expressed genes based on their number of protein-
protein interactions (PPIs), associated GO biological processes, and developmental stages when
they are known to be expressed. However, we did not find any difference in pleiotropy for genes
showing parallel changes in expression compared to genes showing divergent expression using
any of these three metrics (GLM; biological processes: $P = 0.67$; PPIs: $P = 0.09$; developmental
stages: $P = 0.89$) (Fig. S6).

Genetic variation underlying parallel changes in expression

We identified 79 SNPs fixed between generalists *vs.* snail-eaters and 1,543 SNPs fixed between
generalists *vs.* scale-eaters (also see our previous study on genome-wide association mapping
jaw length in these species). None of these fixed variants were shared between specialists. Next,
we determined which of these fixed SNPs fell within gene regions (either exonic, intronic, or
within 10kb of the first or last exon; Table 1). 26 SNPs fixed in snail-eaters overlapped with 17
gene regions, whereas 1,276 SNPs fixed in scale-eaters overlapped with 245 gene regions.

Next, we identified fixed variants near genes that showed differential expression. We
found 319 SNPs fixed in scale-eaters within 71 gene regions that showed differential expression
between generalists and scale-eaters at 8-10 dpf and 118 SNPs within 26 gene regions
differentially expressed between generalists and scale-eaters at 17-20 dpf. We suspect that some
of these fixed variants are within *cis*-regulatory elements responsible for species-specific
expression patterns that ultimately give rise to phenotypic differences in scale-eaters.
Conversely, we only identified a single SNP fixed in snail-eaters within a gene (*tmprss2*) that
was differentially expressed between generalists and snail-eaters at 8-10 dpf. We did not find any
fixed variants near genes differentially expressed between generalists and snail-eaters at 17-20
dpf, possibly suggesting that fixed variants regulate expression divergence at an earlier
developmental stage.

Since we did not find any variants that were fixed between snail-eaters and generalists
that were also fixed between scale-eaters and generalists, we searched for shared variation at a

lower threshold of genetic divergence. We calculated the 99th percentile outlier F_{st} estimates

between randomly subsampled groups of each species across 1,000 permutations to create two

null distributions of genome-wide divergence. We took the 99th percentile of these distributions

as an estimate of significantly high divergence (Fst > 0.36 for generalists *vs.* snail-eaters; Fst >

0.42 for generalists *vs.* scale-eaters; Fig. S7). We found 4,410 SNPs above this lower threshold

of divergence near 134 genes showing parallel changes in expression between specialists at 8-10

dpf. The most differentiated SNPs near genes showing parallel changes in expression show F_{st} <

0.8 between generalists *vs.* snail-eaters and generalists *vs.* scale-eaters. Overall, these results

suggest it is unlikely that the parallel evolution of gene expression in specialists is controlled by

shared variation that is fixed or nearly fixed in specialist populations.

The genetic basis of extreme craniofacial divergence

We previously described 30 candidate gene regions containing variants fixed between trophic

specialist species associated with variation in jaw length. These candidates also showed

signatures of a recent hard selective sweep (McGirr and Martin 2017). Encouragingly, we found

ten of these genes differentially expressed between generalists and scale-eaters (eight at 8-10 dpf

and two at 17-20 dpf) and one between generalists and snail-eaters (8-10 dpf; Table S5).

We searched for signatures of hard selective sweeps across the 84 gene regions

containing fixed variation in specialists (Table 1). Interestingly, 80% of these gene regions

showed signs of a hard sweep (estimated by SweeD; CLR > 95th percentile across their

respective scaffolds; Table S6). All of these gene regions contained SNPs that were either fixed

between generalists *vs.* snail-eaters or generalists *vs.* scale-eaters and showed differential

expression at 8-10 dpf, 17-20 dpf, or both. Finally, we compared this list of genes experiencing

selection to those annotated for cranial skeletal system development (GO:1904888) and muscle

organ development (GO:0007517). While this search was limited to zebrafish orthologs

identified as one-way best hits, we were able to identify three genes containing fixed variation in

scale-eaters that likely influence craniofacial divergence through cis-acting regulatory

mechanisms (*loxl3b* (annotated for cranial effects)*; fbxo32* and *klhl40a* (annotated for muscle

effects)).

Discussion

We combined RNA sequencing with genome-wide divergence scans to study the molecular
evolution of two trophic specialist species that rapidly diverged from a generalist common
ancestor within the last 10,000 years. We examined how gene expression and SNP variation
influence snail-eater and scale-eater niche adaptations using comparisons between each specialist
and their generalist sister species. We found a significant amount of parallelism at the level of
gene expression yet no parallelism at the level of fixed genetic variation within specialists.
Specifically, 80% of genes that were differentially expressed between snail-eaters and generalists
were up or downregulated in the same direction when comparing expression between scale-
eaters and generalists (Fig. 2A). We explored two possible explanations for this pattern: 1)
reduced pleiotropic constraints made these genes likely targets for parallelism or 2) convergent
processes drove parallel gene expression evolution in this highly divergent pair of specialist
species due to shared adaptations to a higher trophic level.

1. Pleiotropic constraints do not explain parallel changes in gene expression

Genes that effect one or a few traits are less constrained than genes with many phenotypic
effects, perhaps making them simpler shared targets for expression divergence during adaptive
evolution between independently evolving lineages. Indeed, theory predicts that the probability
of parallel evolution of gene expression should be higher for genes with minimal pleiotropic
effects (Manceau et al. 2010, Rosenblum et al. 2014). We predicted that genes showing parallel
changes in expression between specialists would show lower degrees of pleiotropy than
divergently expressed genes. We estimated three measures of gene pleiotropy (number of
associated GO biological processes, protein-protein interactions (PPIs), and developmental
stages when they are known to be expressed) and found no significant difference in any measure
for genes showing parallel versus divergent changes in expression patterns (Fig. S6). This
finding is consistent with some empirical evidence and theoretical models of gene expression
evolution that found pleiotropy constrains the variability of gene expression within species, but
does not hinder divergence between species (Tulchinsky et al. 2014; Uebbing et al. 2016).

2. Parallel changes in gene expression underlie convergent metabolic adaptations to a higher
trophic level in each specialist

While the specialists are more morphologically diverged from one another than either is from the
generalist species, particularly in their craniofacial phenotype and male reproductive coloration
(Martin and Wainright 2013a; Martin et al. 2017) (Fig. 3B and C), dietary isotope analyses
show that they both occupy a higher trophic level than generalists (Martin 2016b). Fish scales
and mollusks contribute to more nitrogen-rich diets in specialists compared to generalist species
that primarily consume algae and detritus (Martin 2016b). Perhaps the same metabolic processes
required for this type of diet are adaptive at higher trophic levels for both scale-eaters and snail-
eaters, which might explain patterns of parallel changes in expression. Thus, we predicted that
genes showing parallel changes in expression would affect metabolic processes that may be
similar between specialists, whereas genes showing divergent expression between specialists
would affect morphological development.

GO enrichment analyses using one-way best-hit zebrafish orthologs support both
hypotheses. We found that 20% of GO terms enriched for genes showing parallel changes in
expression described metabolic processes, and zero described cranial skeletal development or
pigment biosynthesis (Fig. 3A; Table S3). In contrast, 10% of terms showing enrichment in the
divergently expressed gene set described developmental processes (cranial skeletal development
and pigment biosynthesis) and only 11% described metabolic processes (Fig 3A, Table S4). GO
enrichment analyses using more conservatively defined reciprocal best hit orthologs confirmed
that genes showing parallel changes in expression were highly enriched for metabolic processes
(26% of representative terms). These results suggest that the parallel evolution of expression in
specialists confers adaptation to a higher trophic level. Snail-eating and scale-eating may present
similar metabolic requirements relative to the lower trophic level of algivorous generalists. This
is consistent with the high macroalgae content of generalist diets relative to both specialist
species (Martin and Wainright 2013b) and the shorter intestinal lengths observed in both
specialists relative to the generalist (CHM and JAM personal observation).

Enrichment analyses using one-way best hit orthologs indicate that genes showing
divergent expression in specialists are responsible for shaping divergent cranial and pigmentation

phenotypes between species (Fig. 3), but we did not find enrichment for these processes using

reciprocal best hit orthologs. This may be because up to 60% of orthologous relationships are

missed by the reciprocal best-hit criterion in lineages with genome duplications, including

teleosts (Dalquen and Dessimoz 2013). Finally, both approaches we used to establish orthology

indicated that genes showing divergent expression in specialists were moderately enriched for

metabolic processes (Fig. 3A; Table S4). While parallel changes in expression may broadly

influence adaptation to a higher trophic level, these divergently expressed metabolic genes likely

play a role in dietary specialization unique to each species.

Parallel changes in gene expression despite unshared genetic variation

We find significant parallel evolution of gene expression across genes that are annotated for

effects on metabolism, yet shared expression patterns do not seem to be driven by the same fixed

variants. This is surprising in this young radiation given that the probability of shared genetic

variation underlying phenotypic convergence increases with decreasing divergence time

(Schluter et al. 2004; Conte et al. 2012; Martin and Orgogozo 2013). Although 80% of

differentially expressed gene regions containing fixed SNPs show signs of experiencing a

selective sweep, and almost none of these variants were in exons, it is still possible that fixed

alleles do not regulate parallel changes in expression for metabolic genes. Indeed, we found

4,410 SNPs that showed significant differentiation between generalists *vs.* snail-eaters and

generalists *vs.* scale-eaters near 134 genes showing parallel changes in expression. These shared

variants all showed $F_{st} < 0.8$, suggesting that parallel expression is not controlled by shared

variation that is fixed or nearly fixed in specialist populations. However, our results do not rule

out a role for fixed variation influencing the parallel evolution of expression through long-range

chromosome interactions or during earlier critical developmental stages, such as neural crest cell

migration at approximately 48 hpf.

Many studies that show parallel adaptation at the gene level describe convergence within

pigmentation and skeletal development pathways (Miller et al. 2007b; Reed et al. 2011; Conte et

al. 2012; Kronforst et al. 2012). Perhaps the architecture of metabolic adaptation is more flexible,

having more mutational targets or employing more late-acting developmental regulatory

networks that are less constrained than early-acting networks (Kalinka et al. 2010; Garfield et al.

2013; Martin and Orgogozo 2013; Reddiex et al. 2013; Ferna et al. 2014; Comeault et al. 2017).

Our findings highlight the importance of understanding convergence across different biological

levels of organization.

Candidate genes influencing trophic adaptations

We found many genes affecting metabolism that were differentially expressed in the same

direction in specialists relative to their generalist sister species. While the metabolism ontology

includes a broad class of proteins with a variety of biological functions, we find many with

distinct effects on dietary metabolism. For example, the gene *asl* (argininosuccinate lyase) is

important for nitrogen excretion. Variants of *asl* are associated with argininosuccinic aciduria

and citrullinemia, conditions involving an accumulation of ammonia in the blood (Saheki et al.

1987; Hu et al. 2015). This gene, along with some of 274 other genes we found annotated for

nitrogen metabolism, may show parallel changes in expression between specialists as an

adaptation to nitrogen-rich diets (Martin 2016b).

We also identified candidate genes influencing cranial divergence that were differentially

expressed between scale-eaters and generalists, contain SNPs fixed in scale-eaters, and showed

signs of a hard selective sweep. *loxl3b* is highly expressed in scale-eaters at 8-10 dpf and

annotated for cranial effects (Table S6). The protein encoded by this gene (lysyl oxidase 3b)

controls the formation of crosslinks in collagens, and is vital to cartilage maturation during

zebrafish craniofacial development (Van Boxtel et al. 2011). Mutations in *loxl3b* are associated

with Stickler Syndrome, which is characterized by cranial anomalies and cleft palate (Alzahrani

et al. 2015). *fbxo32* and *klhl40a* are both expressed at lower levels in scale-eaters at 8-10 dpf

relative to generalists and may influence skeletal muscle divergence between species (Table S6).

High expression of *fbxo32* is associated with muscle atrophy, while mutations in *klhl40a* cause

nemaline myopathy (muscle weakness) (Ravenscroft et al. 2013; Mei et al. 2015). Variants fixed

in scale-eaters near these genes, along with fixed variation near differentially expressed genes

previously associated with large jaw size (McGirr and Martin 2017; Table S5) represent strong

cis-acting regulatory candidates potentially influencing scale-eater cranial traits.

Caveats to gene expression analyses and the robustness of parallel evolution

We compared the transcriptomes of derived trophic specialists to a contemporary generalist sister

species to identify gene expression divergence important for the evolution of trophic traits.

However, the generalist transcriptome represents an approximation of the putative ancestral

state, and has also evolved over the past 10,000 years (Holtmeier 2001; Turner et al. 2008;

Martin and Wainwright 2011; Martin 2016a). We chose to sample RNA at 8-10 dpf and 17-20

dpf to identify transcriptional variation that influences larval development, however, some

activation of parallel gene networks is likely specified at pre-hatching developmental stages

(Garfield et al. 2013; Ferna et al. 2014). It is also possible that we did not have the power to

identify subtle differences in expression for genes that showed high divergence between

specialists and generalists. Detecting differential expression of transcripts is notoriously difficult

when read counts are low and variance within treatment groups is high (Conesa et al. 2016; Lin

et al. 2016). We were able to detect differential expression for genes with a mean normalized

count as low as 1.6 (median = 150) and $\log_2$ fold change as low as 0.2 (median = 1.11).

Furthermore, our scale-eater sample sizes (8-10 dpf n = 3; 17-20 dpf n = 2) were lower than that

of generalists and snail-eaters (n = 6 at both stages; Table S1). Nonetheless, down sampling

analyses suggest that patterns of parallel expression are robust to smaller sample sizes for 8-10

dpf tissue (Fig. 2C), but less so for 17-20 dpf tissue (Fig. S4C).

Finally, our novel results are consistent with a recently published independent analysis of

gene expression in San Salvador pupfishes that identified many of the same genes we found

divergently expressed between specialists (Lencer et al. 2017). We examined this dataset using

the same significance thresholds for differentially expressed genes as described in Lencer *et al.*

for mRNA extracted from all three species at 8 dpf and 15 dpf ($P < 0.1$ and $|\text{Log}_2$ fold change$| >$

0.2). We found that 40% of genes divergently expressed between specialists in this dataset were

divergently expressed in our own dataset. Importantly, Lencer *et al.* only examined cranial

tissues at both of these developmental stages and they did not choose to examine parallel

evolution of expression. We also searched for evidence of parallel change in expression for

mRNA extracted from all three species at 8 dpf in the Lencer et al. dataset. 28.8% of genes that

were differentially expressed between snail-eaters and generalists were up or downregulated in

the same direction between scale-eaters and generalists. This is a lower proportion of parallel

change in expression than we identified (Fig. 2), but this is most likely because Lencer *et al.* only

sampled RNA from cranial tissues at 8 dpf, unlike our sampling of whole larvae. Thus, the

majority of parallel changes in expression between specialists likely occurs in non-cranial

tissues, consistent with our shared metabolic hypothesis.

Conclusion

Here we find significant parallel evolution of gene expression between two highly divergent

specialist species relative to their generalist sister species. While there are many cases of parallel

changes in expression underlying parallel specialization, to our knowledge, this represents the

first case of parallel expression underlying divergent specialization. Numerous studies have

shown that shared genetic variation underlying phenotypic convergence is more likely when

divergence times between species are short (Schluter et al. 2004; Conte et al. 2012; Martin and

Orgogozo 2013). Scale-eating and snail-eating species have evolved rapidly within the last

10,000 years, yet we do not find the same variants fixed in both species underlying parallel

changes in expression. We show that parallel evolution of expression likely reveals convergent

adaptation to a higher trophic level in each specialist, despite their highly divergent resource use

and morphology.

Acknowledgements

This study was funded by the University of North Carolina at Chapel Hill and the Miller Institute

for Basic Research in the Sciences to CHM. We thank Sara Suzuki, Jelmer Poelstra, and Emilie

Richards for valuable discussions and computational assistance; the High Throughput Genomic

Sequencing Facility at UNC Chapel Hill for performing RNA library prep and Illumina

sequencing; the Gerace Research Centre for accommodation; and the Bahamian government

BEST Commission for permission to conduct this research. All datasets used for this study will

be deposited in the NCBI Short Read Archive associated with BioProject PRJNA391309.

Author Contributions

JAM wrote the manuscript. JAM extracted the RNA samples, and conducted all bioinformatic

and population genetic analyses. Both authors contributed to the conception and development of

the ideas and revision of the manuscript.

Data Accessibility

All genomic and transcriptomic raw sequence reads are available on the NCBI BioProject

database. Title: Craniofacial divergence in Caribbean Pupfishes. Accession: PRJNA391309.

Competing interests

We declare we have no competing interests.

References

Ahi, E. P., K. H. Kapralova, A. Pálsson, V. H. Maier, J. Gudbrandsson, S. S. Snorrasonet al.
2014. Transcriptional dynamics of a conserved gene expression network associated with
craniofacial divergence in Arctic charr. Evodevo. 5(1):40.

Altenhoff, A. M., C. Dessimoz. 2009. Phylogenetic and functional assessment of orthologs
inference projects and methods. PLoS Comput Biol. 5(1):e1000262

Alzahrani, F., S.A. Al Hazzaa, H. Tayeb, and F.S. Alkuraya. 2015. LOXL3, encoding lysyl
oxidase-like 3, is mutated in a family with autosomal recessive Stickler syndrome. Hum Genet.
Apr;134(4):451-3.

Ashburner, M., C. A. Ball, J. A. Blake, D. Bolstein et al. 2000. Gene ontology: tool for the
unification of biology. Nat. Genet. 25(1):25-9.

Bastian, F., G. Parmentier, J. Roux, S. Moretti et al. 2008. Bgee: Integrating and comparing
heterogeneous transcriptome data among species. DILS. 124–131.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. J R Stat Soc Series B. 57(1):289-300.

Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, and T. L. Madden.
2009. BLAST +□: architecture and applications. BMC Bioinformatics. 10:421.

Chan, Y. F., M. E. Marks, F. C. Jones, G. V. Jr, M. D. Shapiro, S. D. Brady et al. 2010. Adaptive
evolution of pelvic reduction in sticklebacks by recurrent deletion of a Pitx1 enhancer. Science.
327(5963):302-5.

Comeault, A. A., A. Serrato-capuchina, D. A. Turissini, P. J. Mclaughlin, J. R. David, and D. R.
Matute. 2017. A nonrandom subset of olfactory genes is associated with host preference in the
fruit fly Drosophila orena. Evolution Letters. 1(2):73-85

Conesa, A., P. Madrigal, S. Tarazona, D. Gomez-cabrero, A. Cervera, A. Mcpherson et al. 2016.
A survey of best practices for RNA-seq data analysis. Genome Biol. 17:13.

Conte, G. L., M. E. Arnegard, C. L. Peichel, and D. Schluter. 2012. The probability of genetic
parallelism and convergence in natural populations. Proc. R. Soc. B Biol. Sci. 279:5039–5047.

Cooper, T. F., D. E. Rozen, and R. E. Lenski. 2003. Parallel changes in gene expression after
20,000 generations of evolution in Escherichia coli. Proc. Natl. Acad. Sci. U.S.A. 100(3):1072-7.

Dalquen, D. A., C. Dessimoz. 2013. Bidirectional best hits miss many orthologs in duplication-
rich clades such as plants and animals. Genome Biol Evol. 5(10):1800-6.

Davidson, J. H., and C. N. Balakrishnan. 2016. Gene Regulatory Evolution During Speciation in
a Songbird. G3. 6:1357–1364.

DePristo, M. A., E. Banks, R. Poplin, K. V Garimella, J. R. Maguire, C. Hartl et al. 2011. A
framework for variation discovery and genotyping using next-generation DNA sequencing data.
Nat. Genet. 43:491–8.

Derome, N., and L. Bernatchez. 2006. The Transcriptomics of Ecological Convergence between
2 Limnetic Coregonine Fishes (Salmonidae). Mol Biol Evol. 23(12):2370-8

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson,
and T. R. Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15–21.

Enard, W., P. Khaitovich, J. Klose, F. Heissig, P. Giavalisco, K. Nieselt-struwe et al. 2002. Intra-
and Interspecific Variation in Primate Gene Expression Patterns. Science 296:340–344.

Ewels, P., S. Lundin, and K. Max. 2016. Data and text mining MultiQC: summarize analysis
results for multiple tools and samples in a single report. Bioinformatics 32:3047–3048.

Ferna, A., J. J. Tena, C. Gonza, H. Parra-acero, J. W. Cross, P. W. J. Rigby et al. 2014.
Comparative epigenomics in distantly related teleost species identifies conserved cis -regulatory
nodes active during the vertebrate phylotypic period. Genome Res. 24(7):1075-85.

Garfield, D. A., D. E. Runcie, C. C. Babbitt, R. Haygood, W. J. Nielsen, and G. A. Wray. 2013.
The Impact of Gene Expression Variation on the Robustness and Evolvability of a
Developmental Gene Regulatory Network. PLoS Biol. 11(10):e1001696.

Gene Ontology Consortium. 2017. Expansion of the Gene Ontology knowledgebase and
resources. Nucleic Acids Res. 45:331-338

He, X., and J. Zhang. 2006. Toward a Molecular Understanding of Pleiotropy. Genetics. 1891:1885–1891.

Hernandez, L. P., M. Adriaens, C. H. Martin, P. C. Wainwright, B. Masschaele, M. Dierick. 2017. Building trophic specializations that result in substantial niche partitioning within a young adaptive radiation. J. Anat. doi: 10.1111/joa.12742.

Holtmeier, C.L. 2001. Heterochrony, maternal effects, and phenotypic variation among sympatric pupfishes. Evolution 55:330–338.

Hu L., A.V. Pandey, C. Balmer, S. Eggimann, V. Rüfenacht , J. M. Nuoffer et al. 2015. Unstable argininosuccinate lyase in variant forms of the urea cycle disorder argininosuccinic aciduria. J Inherit Metab Dis. 38(5):815-27

Kalinka, A. T., K. M. Varga, D. T. Gerrard, S. Preibisch, D. L. Corcoran, J. Jarrells et al. 2010. Gene expression divergence recapitulates the developmental hourglass model. Nature 468:811–814.

Kronforst, M. R., G. S. Barsh, A. Kopp, J. Mallet, A. Monteiro, S. P. Mullen et al. 2012. Unraveling the thread of nature's tapestry: the genetics of diversity and convergence in animal pigmentation. Pigment Cell Melanoma Res. 25(4):411-33

Lencer, E. S., W. C. Warren, R. Harrison, and A. R. Mccune. 2017. The Cyprinodon variegatus genome reveals gene expression changes underlying differences in skull morphology among closely related species. BMC Genomics 18(1):424.

Liao, Y., G. K. Smyth, and W. Shi. 2014. Sequence analysis featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 30:923–930.

Lin, Y., K. Golovnina, Z. Chen, H. N. Lee, Y. L. S. Negron, H. Sultana et al. 2016. Comparison of normalization and differential expression analyses using RNA-Seq data from 726 individual Drosophila melanogaster. BMC Genomics. 17:28.

Losos, J. B. 2011. Convergence, adaptation, and constraint. Evolution. 65(7)1827–1840.

Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Gen. Biol. 15:550.

Manousaki, T., P. M. Hull, H. Kusche, G. Machado-schiaffino, P. Franchini, and C. Harrod. 2013. Parsing parallel evolution : ecological divergence and differential gene expression in the adaptive radiations of thick-lipped Midas cichlid fishes from Nicaragua. Mol Ecol. 22(3):650-69.

Manceau M, V. S. Domingues, C. R. Linnen, E. B. Rosenblum, H. E. Hoekstra. 2010. Convergence in pigmentation at multiple levels: mutations, genes and function. Philos. Trans. R. Soc. B-Biol. Sci. 365(1552):2439–50

Martin, C. H., P. A. Erickson, and C. T. Miller. 2017. The genetic architecture of novel trophic specialists : larger effect sizes are associated with exceptional oral jaw diversification in a pupfish adaptive radiation. Mol Ecol. 26(2):624–638.

Martin, A., and V. Orgogozo. 2013. The Loci of repeated evolution: a catalog of genetic hotspots
of phenotypic variation. Evolution. 67(5):1235-50.

Martin, C. H. 2016a. The cryptic origins of evolutionary novelty : 1000-fold faster trophic
diversification rates without increased ecological opportunity or hybrid swarm. Evolution.
70(11):2504-2519.

Martin, C. H. 2016b. Context-dependence in complex adaptive landscapes : frequency and trait-
dependent selection surfaces within an adaptive radiation of Caribbean pupfishes. Evolution
70(6):1265-1282.

Martin, C. H., and P. C. Wainwright. 2013a. A remarkable species flock of Cyprinodon
pupfishes endemic to San Salvador Island, Bahamas. Bull. Peabody Museum Nat. Hist. 54:231–
241.

Martin, C. H., and P. C. Wainwright. 2013b. On the measurement of ecological novelty: scale-
eating pupfish are separated by 168 my from other scale-eating fishes. PLoS One 8:e71164.

Martin, C. H., and P. C. Wainwright. 2011. Trophic novelty is linked to exceptional rates of
morphological diversification in two adaptive radiations of Cyprinodon pupfish. Evolution
65:2197–212.

McGirr, J. A., and C. H. Martin. 2017. Novel Candidate Genes Underlying Extreme Trophic
Specialization in Caribbean Pupfishes. Mol. Biol. Evol. 34:873–888.

Mei, Z., D. Zhang, B. Hu, J. Wang, X. Shen, W. Xiao. 2015. FBXO32 Targets c-Myc for
Proteasomal Degradation and Inhibits c-Myc Activity. J Biol Chem. 290(26):16202-14.

Miller, C. T., S. Beleza, A. A. Pollen, D. Schluter, R. A. Kittles, M. D. Shriver, and D. M.
Kingsley. 2007a. cis -Regulatory Changes in Kit Ligand Expression and Parallel Evolution of
Pigmentation in Sticklebacks and Humans. Cell 131(6):1179-89.

Miller, C. T., M. E. Swartz, P. A. Khuu, M. B. Walker, J. K. Eberhart, and C. B. Kimmel. 2007b.
mef2ca is required in cranial neural crest to effect Endothelin1 signaling in zebrafish. Dev. Biol.
308:144–157.

Mylroie, J.E, Hagey, F. M. 1995. Terrestrial and Shallow Marine Geology of the Bahamas and
Bermuda. Geological Society of America, Boulder, CO.

Nagai, H., Y. Terai, T. Sugawara, H. Imai, H. Nishihara et al. 2011. Reverse Evolution in RH1
for Adaptation of Cichlids to Water Depth in Lake Tanganyika Research article Determination of
Sequences Upstream. Mol. Biol. Evol. 28:1769–1776.

Papakostas, S., L. A. Vøllestad, M. Bruneaux M et al. 2014. Gene pleiotropy constrains gene
expression changes in fish adapted to different thermal conditions. Nature. Comm. 5:4071.

Pavey, S. A., P. Nosil, and S. M. Rogers. 2010. The role of gene expression in ecological
speciation. Ann. N. Y. Acad. Sci. 1206:110–129.

Pavlidis, P., D. Živković, A. Stamatakis, and N. Alachiotis. 2013. SweeD: Likelihood-based detection of selective sweeps in thousands of genomes. Mol. Biol. Evol. 30:2224–2234.

Poelstra, J. W., N. Vijay, C. M. Bossu, H. Lantz, B. Ryll, I. Müller et al. 2014. The genomic landscape underlying phenotypic integrity in the face of gene flow in crows. Science 344:1410–4.

O'Quin, K. E., C. M. Hofmann, H. A. Hofmann, and K. L. Carleton. 2010. Parallel Evolution of Opsin Gene Expression in African Cichlid Fishes Research article. Mol Biol Evol. 27(12):2839-54.

Ravenscroft, G., S. Miyatake, V.L. Lehtokari, E.J Todd, P. Vornanen, K.S Yau et al. Mutations in KLHL40 are a frequent cause of severe autosomal-recessive nemaline myopathy. Am J Hum Genet. 93(1):6-18.

Reddiex, A. J., T. P. Gosden, R. Bonduriansky, and S. F. Chenoweth. 2013. Sex-Specific Fitness Consequences of Nutrient Intake and the Evolvability of Diet Preferences. Am Nat. 182(1):91-102.

Reed, R. D., R. Papa, A. Martin, H. M. Hines, M. R. Kronforst, R. Chen et al. 2011. optix drives the repeated convergent evolution of butterfly wing pattern mimicry. Science. 2011 333(6046):1137-41

Reid, N. M., D. A. Proestou, B. W. Clark, W. C. Warren, J. K. Colbourne, J. R. Shaw et al. 2016. The genomic landscape of rapid repeated evolutionary adaptation to toxic pollution in wild fish. Science 354(6317):1305-1308.

Richards, E. J., and C. H. Martin. 2017. Adaptive introgression from distant Caribbean islands contributed to the diversification of a microendemic adaptive radiation of trophic specialist pupfishes. PLoS Genet. 13(8):e1006919.

Riehle, M. M., A. F. Bennett, R. E. Lenski, A. D. Long, M. Michelle, A. F. Bennett, et. al. 2003. Evolutionary changes in heat-inducible gene expression in lines of Escherichia coli adapted to high temperature. Physiol. Genomics. 14(1):47-58.47–58.

Rosenblum, E. B., C. E. Parent, and E. E. Brandt. 2014. The Molecular Basis of Phenotypic Convergence. Ann. Rev. Eco. Evo. Sys. 45:203-26.

Safari-alighiarloo, N., M. Taghizadeh, M. Rezaei-tavirani, B. Goliaei, and A. A. Peyvandi. 2014. Protein-protein interaction networks (PPI) and complex diseases. Gastroenterol. Hepatol. Bed. Bench. 7:17–31.

Saheki, T., K. Kobayashi , H. Ichiki, S. Matuo, M. Tatsuno, Y. Imamura et al. 1987. Molecular basis of enzyme abnormalities in urea cycle disorders. With special reference to citrullinemia and argininosuccinic aciduria. Enzyme. 38(1-4):227-32.

Salichos, L., A. Rokas. 2011. Evaluating ortholog prediction algorithms in a yeast model clade. PLoS One. 6(4):e18755

Schiffels, S., and R. Durbin. 2014. Inferring human population size and separation history from
multiple genome sequences. Nat. Publ. Gr. 46:919–925.

Schluter, D., E. A. Clifford, M. Nemethy, and J. S. Mckinnon. 2004. Parallel Evolution and
Inheritance of Quantitative Traits. Am Nat. 163(6):809-22.

Shapiro, M. D., M. A. Bell, and D. M. Kingsley. 2006. Parallel genetic origins of pelvic
reduction in vertebrates. PNAS. 103:13753–13758.

Shen, Y., L. Liang, G. Li, R. W. Murphy, and Y. Zhang. 2012. Parallel Evolution of Auditory
Genes for Echolocation in Bats and Toothed Whales. PLoS Genet. 8(6):e1002788.

Stern, D. L., V. Orgogozo. 2008. The loci of evolution: How predictable is genetic evolution?
Evolution 62(9):2155–77.

Szklarczyk, D., A. Franceschini, S. Wyder, K. Forslund et al. 2015. STRING v10: protein –
protein interaction networks, integrated over the tree of life. Nucleic Acids Res. 43:447–452.

Tomislav, S. 2011. REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms.
PLoS One. 6(7):e21800.

Turner, B.J., D. D. Duvernell, T.M. Bunt, M.G.Barton. 2008 Reproductive isolation among
endemic pupfishes (Cyprinodon) on San Salvador Island, Bahamas: microsatellite evidence.
Proc. Roy. Soc. Lond. B. 95:566–582.

Uebbing, S., A. Künstner, H. Mäkinen, N. Backström, P. Bolivar, R. Burri et al. 2016.
Divergence in gene expression within and between two closely related flycatcher species. Mol
Ecol. 25(9):2015-28.

Van Boxtel, A.L., J.M. Gansner, H.W Hakvoort, H. Snell, J. Legler, J.D. Gitlin. 2011. Lysyl
oxidase-like 3b is critical for cartilage maturation during zebrafish craniofacial development.
Matrix Biol. 30(3):178-87

Zhao, L., J. Wit, N. Svetec, and D. J. Begun. 2015. Parallel Gene Expression Differences
between Low and High Latitude Populations of Drosophila melanogaster and D . simulans. PLoS
Genet. 11(5):e1005184.

Table 1. Genomic distribution of fixed variants. The first five columns show the total number
of fixed SNPs in each species comparison and how many fall within exons, introns, 10kb of the

first or last exon of a gene, and outside of 10kb from the first or last exon of a gene. Final two
columns show the number of genes with fixed SNPs within the gene and/or within 10kb of the
first or last exon. The last column shows the number of differentially expressed (DE) genes near
fixed SNPs that includes DE genes from 8-10 dpf and 17-20 dpf comparisons.
846
847

comparison	fixed SNPs	exonic	intronic	within 10kb of coding region	>10kb away from coding region	number of genes near fixed SNPs	number of DE genes near fixed SNPs
generalist vs. snail-eater	79	0	16	10	53	17	1
generalist vs. scale-eater	1,543	140	512	624	267	245	83

Fig. 1. Differential gene expression between generalists and trophic specialists. Red points represent genes that are differentially expressed in 8-10 dpf whole-larvae tissue (A, C) and 17-20 dpf craniofacial tissue (B, C) between generalists *vs.* scale-eaters (A, B) and generalist *vs.* snail-eaters (C, D). Bottom panels show the top two principal components accounting for a combined 52% (8-10 dpf; E) and 48% (17-20 dpf; F) of the total variation between samples across 413 million reads mapped to annotated features. Triangles represent samples from Little Lake and circles represent samples from Crescent Pond on San Salvador Island.

Fig. 2. Parallel evolution of gene expression between specialists despite divergent trophic adaptation. A) Circles illustrate genes differentially expressed in 8-10 dpf whole-larvae tissue for generalists *vs.* scale-eaters (left) and generalists *vs.* snail-eaters (right). Genes showing differential expression in both comparisons are shown in blue, and those showing divergent expression patterns unique to each specialist are green. Significantly more genes show differential expression in both comparisons than expected by chance (Fisher's exact test, $P < 1.0 \times 10^{-16}$). B) Significantly more genes show the same direction of expression in specialists relative to generalists than expected by chance (10,000 permutations; $P < 1.0 \times 10^{-4}$; Fig. S2). C) Distribution of the proportion of genes differentially expressed in the same direction between specialists relative to generalists after 1,000 down sampling permutations show that parallel expression is robust to variation in sample size (median number of genes common to both comparisons = 61).

Fig. 3. Parallel gene expression underlies metabolic adaptations while divergent expression underlies trophic morphology A) Genes showing parallel changes in expression between specialists (blue) and genes showing divergent expression (green) are contrastingly enriched for terms describing metabolic processes (parallel: 20% of enriched terms; divergent: 11% of terms). Genes showing divergent expression are enriched for cranial skeleton development (7% of terms) and pigment biosynthesis (3% of terms). B) μCT scans show drastic craniofacial divergence between snail-eaters (top) and scale-eaters (bottom) (modified from Hernandez et al. 2017). Bottom panels show male breeding coloration characteristic of light snail-eaters (C) and dark scale-eaters (D).