MicroRNAs from the parasitic plant *Cuscuta campestris* target host messenger

RNAs

Saima Shahid[1,2], Gunjune Kim[3], Nathan R. Johnson[1,2], Eric Wafula[2], Feng Wang[1,2,4],

Ceyda Coruh[1,2,5], Vivian Bernal-Galeano[3], Tamia Phifer[6], Claude W. dePamphilis[1,2],

James H. Westwood[3], and Michael J. Axtell[1,2]

[1] Intercollege Ph.D. Program in Plant Biology, Huck Institutes of the Life Sciences, The

Pennsylvania State University, University Park, PA 16802 USA

[2] Department of Biology, The Pennsylvania State University, University Park, PA 16802

USA

[3] Department of Plant Pathology, Physiology and Weed Science, Virginia Polytechnic

Institute and State University, Blacksburg, VA 24061, USA.

[4] Current Address: Department of Biology, Indiana University, Bloomington IN 47405,

USA

[5] Current Address: Salk Institute for Biological Studies, La Jolla, CA 92037, USA

[6] Knox College, Galesburg, IL 61401, USA

First paragraph:

Dodders (*Cuscuta* spp.) are obligate parasitic plants that obtain water and

nutrients from the stems of host plants via specialized feeding structures called

haustoria. Dodder haustoria facilitate bi-directional movement of viruses, proteins, and

mRNAs between host and parasite[1], but the functional effects of these movements are

not clear. Here we show that *C. campestris* haustoria accumulate high levels of many

novel microRNAs (miRNAs) while parasitizing *Arabidopsis thaliana* hosts. Many of

these miRNAs are 22 nts long, a usually rare size of plant miRNA associated with

amplification of target silencing through secondary small interfering RNA (siRNA)

production[2]. Several *A. thaliana* mRNAs are targeted by *C. campestris* 22 nt miRNAs

during parasitism, resulting in mRNA cleavage, secondary siRNA production, and

decreased mRNA accumulation levels. Hosts with mutations in two of the targets

supported significantly higher growth of *C. campestris.* Homologs of target mRNAs from

diverse plants also have predicted target sites to induced *C. campestris* miRNAs, and

the same miRNAs are expressed and active against host targets when *C. campestris*

parasitizes a different host, *Nicotiana benthamiana*. These data show that *C. campestris*

miRNAs act as *trans*-species regulators of host gene expression, and suggest that they

may act as virulence factors during parasitism.

Host-induced gene silencing (HIGS) involves plant transgenes that produce

siRNAs which can silence targeted pathogen/parasite mRNAs in *trans*[3,4]. Plant-based

HIGS is effective against fungi[5], nematodes[6], insects[7], and the parasitic plant *Cuscuta*

pentagona[8]. The apparent ease of engineering HIGS suggests that plants might

exchange naturally occurring small RNAs during pathogen/parasite interactions.

Consistent with this hypothesis, small RNAs from the plant pathogenic fungus *Botrytis*

cinerea target host mRNAs during infection[9], and HIGS targeting *B. cinerea* Dicer-Like

mRNAs reduces pathogen virulence[10]. Conversely, the conserved miRNAs miR159 and

miR166 can be exported from cotton into the fungal pathogen *Verticillium dahliae* where

they target fungal mRNAs encoding virulence factors[11]. However, to date, no examples

of naturally occurring *trans*-species miRNAs have been described for plant-to-plant

interactions.

Cuscuta haustoria facilitate bi-directional movement of viruses, proteins, and

mRNAs[1], but the functional effects of these movements are unclear. *Cuscuta* is

susceptible to HIGS, so we hypothesized that naturally occurring small RNAs might be

exchanged across the *C. campestris* haustorium and affect gene expression in the

recipient species. To test this hypothesis, we profiled small RNA expression from *C.*

campestris grown on *A. thaliana* hosts using high-throughput small RNA sequencing

(sRNA-seq). Two biological replicates each from three tissues were analyzed: Parasite

stem (PS), comprising a section of *C. campestris* stem above the site of haustorium

formation; Interface (I), comprising *C. campestris* stem with haustoria with associated *A.*

thaliana stem tissue; and Host stem (HS), comprising sections of *A. thaliana* stems

above the interface region, as previously described[12]. Small RNA-producing loci from

both organisms were identified, classified, and subject to differential expression

analyses (Supplementary Data 1).

As expected due to dilution of parasite RNA with host RNA, *C. campestris* small

RNA loci were generally down-regulated in I relative to PS (Figure 1A). However, 76 *C.*

campestris small RNA loci were significantly (FDR <= 0.05) higher in I relative to PS. 43

of these (57%) were *MIRNA* loci as determined by canonical accumulation of a discrete

miRNA/miRNA* pair from predicted stem-loop precursors (Figure 1B, Supplementary

Data 2-4). RNA blots confirmed I-specific expression (Figure 1C). One of the 43

*MIRNA*s is a member of the conserved *MIR164* family; the other 42 up-regulated

*MIRNA*s have no obvious sequence similarity to previously annotated *MIRNA* loci, and

none of their mature miRNAs or miRNA*s were perfectly alignable to the *A. thaliana*

genome (Supplementary Data 5). Several of the key *MIRNA* loci were detected by PCR

of *C. campestris* genomic DNA prepared from four-day old seedlings that had never

interacted with a host plant (Extended Data Figure 1). The majority of the induced *C.*

campestris MIRNA loci (26/43) produced a 22 nt mature miRNA. 22 nt plant miRNAs

are usually less frequent than 21nt miRNAs, and they are strongly associated with

secondary siRNA accumulation from their targets[13,14]. Secondary siRNAs are thought to

amplify the strength of miRNA-directed gene silencing[2].

81

Figure 1. *C. campestris* miRNAs induced at the haustorial interface. **a)** MA plot of *C.*

campestris small RNA loci comparing interface (I) to parasite stem (PS) samples.

Significantly up-regulated loci (alternative hypothesis: true difference > 2-fold, FDR <=

0.05) are highlighted. **b)** Predicted secondary structures of example *MIRNA* hairpin

precursors of *C. campestris* 22 nt miRNAs. Color-coding shows depth of small RNA-seq

coverage at each nucleotide. **c)** RNA blots of 22 nt I-induced miRNAs. PS: parasite

stem, I: interface, HS: host stem, CS: control stem.

We hypothesized that the induced 22 nt miRNAs would cause secondary siRNA

formation from targeted host mRNAs. Therefore we searched for *A. thaliana* mRNAs

that had one or more plausible miRNA complementary sites and accumulation of

secondary siRNAs specifically in the I small RNA-seq samples. Six *A. thaliana* mRNAs

that met both criteria were found: *TIR1*, *AFB2*, and *AFB3*, which encode related and

partially redundant auxin receptors[15], *BIK1*, which encodes a plasma membrane-

localized kinase required for both pathogen-induced and developmental signaling[16,17],

SEOR1, which encodes a major phloem protein that accumulates in filamentous

networks in sieve tube elements and reduces photosynthate loss from the phloem upon

injury[18,19], and *SCZ/HSFB4*, which encodes a predicted transcriptional repressor that is

required for the formation of ground tissue stem cells in roots[20-22]. The induced siRNAs

from these mRNAs resembled other examples of secondary siRNAs in their size

distributions, double-stranded accumulation, and phasing (Figure 2A-B; Extended Data

Figure 2). *TIR1*, *AFB2*, and *AFB3*, are also known to be targeted by the 22 nt miR393

and to produce secondary siRNAs downstream of the miR393 complementary site[23]. In

parasitized stems the location and phase register of the *TIR1*, *AFB2*, and *AFB3*

secondary siRNAs shift upstream, proximal to the complementary sites to the *C.*

campestris miRNAs (Extended Data Figure 2), implying that the *C. campestris* miRNAs,

not miR393, are triggering the I-specific secondary siRNAs. The predominant 21 nt

phase register at several loci was shifted +1 to +2 relative to the predictions. This is

consistent with the 'phase drift' seen at other phased siRNA loci[24,25] and likely due to the

presence of low levels of 22nt siRNAs, causing the register to be shifted forward.

Analysis of uncapped mRNA fragments using 5'-RNA ligase-mediated rapid

amplification of cDNA ends found strong evidence for miRNA-directed cleavage at all of

the complementary sites to *C. campestris* miRNAs, specifically from interface samples

but not from control stem samples (Figure 2; Extended Data Figure 2). We did not find

any induced miRNAs or siRNAs from the *A. thaliana* host capable of targeting these six

mRNAs. We also did not find any endogenous *C. campestris* secondary siRNA loci

corresponding to any of the induced miRNAs. Some *C. campestris* orthologs of

TIR/AFB, SCZ/HSFB4, and *BIK1* had possible, but very poorly complementary, miRNA

target sites (Extended Data Figure 3). These observations suggest that the induced *C.*

campestris miRNAs have evolved to avoid targeting 'self' transcripts. We conclude that

22 nt miRNAs from *C. campestris* act in a *trans*-species manner to target *A. thaliana*

mRNAs.

Figure 2. *C. campestris* miRNAs cause slicing and phased siRNA production from host

mRNAs. **a)** Small RNA-seq coverage (reads per million) for the *A. thaliana SEOR1*

transcript for host stem (HS), interface (I), and parasite stem (PS) samples. miRNA

complementary site with 5'-RLM-RACE data and expected 21nt phasing register is

shown. **b)** Length and polarity distribution of *SEOR1*-mapped siRNAs from the I

samples. **c)**. Radar chart showing fraction of I-derived siRNAs in each of the 21

possible phasing registers. **d)** 5'-RLM-RACE products from nested amplifications for the

indicated cDNAs. I: interface, CS: control stem. *ARF17* is a positive control.

Accumulation of five of the six secondary siRNA-producing targets was

significantly reduced in stems parasitized by *C. campestris* compared to un-parasitized

stems (Figure 3A), consistent with miRNA-mediated repression. The true magnitude of

repression for these targets could be even greater, since many miRNAs also direct

translational repression. Accumulation of many known *A. thaliana* secondary siRNAs is

often partially dependent on the endonuclease Dicer-Like 4 (DCL4) and wholly

dependent on RNA-Dependent RNA polymerase 6 (RDR6/SGS2/SDE1)[2]. Accumulation

of an abundant secondary siRNA from *TIR1* was eliminated entirely in the *sgs2-1*

mutant, and reduced in the *dcl4-2t* mutant (Figure 3B). Thus host *DCL4* and

RDR6/SGS2/SDE1 are required for secondary siRNA production. This implies that the

C. campestris derived miRNAs are active inside of host cells and hijack the host's own

silencing machinery to produce secondary siRNAs.

In repeated trials with varying methodologies we did not observe consistent

significant differences in parasite fresh weight using *dcl4-2t* and *sgs2-1* mutants as

hosts (Extended Data Figure 4). Thus, loss of induced secondary siRNAs is not

sufficient to detectably affect parasite biomass accumulation in this assay. Similarly,

there were no significant differences in parasite fresh weights when *scz2* or *tir1-1/afb2-3*

plants were used as hosts (Figure 3C). Significantly less *C. campestris* biomass was

observed using *bik1* mutants as hosts. However, interpretation of this result is

complicated by the weak, frequently lodging stems of the *bik1* mutant[16]. Significantly

more *C. campestris* biomass was observed when grown on *seor1* or *afb3-4* mutant

hosts (Figure 3C). Therefore, both *SEOR1* and *AFB3* function to restrict *C. campestris*

growth on *A. thaliana*. This observation is consistent with the hypothesis that both

SEOR1 and *AFB2* are *trans*-species miRNA targets of biological relevance.

Figure 3. Effects of *C. campestris* miRNAs and their targets. **a)** *A. thaliana* mRNA

accumulation levels in I (interface) vs. CS (control stems) during *C. campestris*

parasitism, assessed by qRT-PCR. Data from 7 or 8 biological replicates are plotted

(dots), and boxplots indicate the median (horizontal lines), 1st-3rd quartile range

(boxes), and up to 1.5 x the interquartile range (whiskers). Numbers indicate p-values

comparing CS and I samples (Wilcoxon rank-sum test, unpaired, one-tailed). *TIP41-I* :

control mRNA. **b)** RNA blots from *C. campestris* infestations of the indicated *A. thaliana*

genotypes. I: interface, HS: host stem, PS: parasite stem, CS: control stem. **c)**

Accumulation of *C. campestris* biomass on *A. thaliana* hosts of the indicated genotypes

18 days post-attachment. P-values (Wilcoxon rank-sum tests, unpaired, two-tailed) from

comparison of mutant to wild-type (Col-0) are shown. Boxplot conventions as in panel a.

n=11, 8, 11, 10, 14, and 7 for Col-0, *seor1, bik1, scz2, tir1-1/afb2-3*, and *afb3*,

respectively.

C. campestris has a broad host range among eudicots[26]. Therefore, we searched

for miRNA complementary sites for the interface-induced *C. campestris* miRNAs in

eudicot orthologs of the targeted *A. thaliana* mRNAs. Probable orthologs of *BIK1,*

SEOR1, TIR/AFB, and *SCZ/HSFB4* were predicted targets of interface-induced miRNAs

in many eudicot species, while only one species had predicted targets for the negative

control orthologs of *GAPDH* (Figure 4A, Extended Data Table 1). We conclude that the

induced *C. campestris* miRNAs collectively would be able to target *TIR/AFB, SEOR1,*

SCZ/HSFB4, and *BIK1* orthologs in many eudicot species.

We performed additional small RNA-seq from *C. campestris* on *A. thaliana* hosts,

and from *C. campestris* on *Nicotiana benthamiana* hosts. Both sets of experiments were

designed identically to the original small RNA-seq study (two biological replicates each

of HS, I, and PS samples). The I-induced set of *C. campestris MIRNA* loci was highly

reproducible across both of the *A. thaliana* experiments as well as the *N. benthamiana*

experiment (Extended Data Figure 5). Induction of several *C. campestris* miRNAs

during *N. benthamiana* parasitism was confirmed by RNA blots (Figure 4B). Several *N.*

benthamiana mRNAs had both plausible target sites for *C. campestris* miRNAs and

accumulation of phased, secondary siRNAs in the I samples, including orthologs of

TIR/AFB and *BIK1* (Extended Data Figure 6). Analysis of uncapped RNA ends provided

strong evidence for miRNA-directed cleavage of one of the *N. benthamiana TIR1/AFB*

orthologs (Figures 4C-D; Extended Data Figure 7). This directly demonstrates that the

same *C. campestris* miRNAs target orthologous host mRNAs in multiple species. None

of the interface-induced miRNAs we tested were detectable from *C. campestris* pre-

haustoria sampled from seedling tips that had coiled around dead bamboo stakes

instead of a live host (Figure 4B; Extended Data Figure 8). This suggests that

expression of these miRNAs requires prior contact with a living host.

Figure 4. Conservation of host mRNA targeting by *C. campestris*. **a)** Predicted targets

of the induced *C. campestris* miRNAs/miRNA*s among indicated orthologs. **b)** RNA

blots from interface (I) and control stem (CS) samples of *C. campestris* infested *N.*

benthamiana (*N.b.*) and *A. thaliana* (*A.t.*) hosts, as well as *C. campestris* pre-haustoria

(PH; Extended Data Figure 8). **c)** 5'-RLM-RACE products for the indicated *N.*

benthamiana cDNAs. I: interface, CS: control stem. *N.b. ARF* is a positive control.

Image cropped to remove irrelevant lanes; uncropped image: Extended Data Figure 7.

d) Complementary site and 5'-RLM-RACE results from a *N. benthamiana TIR/AFB*

ortholog.

These data demonstrate that *C. campestris* induces a large number of miRNAs

at the haustorium, and that some of them target host mRNAs and reduce their

accumulation. Many of the induced miRNAs are 22 nts long, and associated with

secondary siRNA production from their host targets using the host's own secondary

siRNA machinery. Several of the targets are linked to plant pathogenesis: Manipulation

of *TIR1/AFB2/AFB3* accumulation levels affects bacterial pathogenesis and defense

signaling[27], and BIK1 is a central regulator of pathogen-induced signaling[28]. Perhaps

the most intriguing target is *SEOR1*, which encodes a very abundant protein present in

large agglomerations in phloem sieve-tube elements[18]. *seor1* mutants have an

increased loss of sugars from detached leaves[19], and our data show that *seor1* mutants

also support increased *C. campestris* biomass accumulation. A key function of the

haustorium is to take nutrients from the host phloem; targeting *SEOR1* could be a

strategy to increase sugar uptake from host phloem. Overall, these data suggest that *C.*

campestris trans-species miRNAs might function as virulence factors to remodel host

gene expression to its advantage during parasitism.

Methods [separate online only document].

References

1.   Kim, G. & Westwood, J. H. Macromolecule exchange in Cuscuta-host plant

interactions. *Curr Opin Plant Biol* **26,** 20–25 (2015).

2.   Fei, Q., Xia, R. & Meyers, B. C. Phased, Secondary, Small Interfering RNAs in

Posttranscriptional Regulatory Networks. *Plant Cell* **25,** 2400–2415 (2013).

3.   Baulcombe, D. C. VIGS, HIGS and FIGS: small RNA silencing in the interactions

of viruses or filamentous organisms with their plant hosts. *Curr Opin Plant Biol* **26,**

141–146 (2015).

4.   Weiberg, A., Bellinger, M. & Jin, H. Conversations between kingdoms: small

RNAs. *Curr. Opin. Biotechnol.* **32,** 207–215 (2015).

5.   Nowara, D. *et al.* HIGS: host-induced gene silencing in the obligate biotrophic

fungal pathogen Blumeria graminis. *Plant Cell* **22,** 3130–3141 (2010).

6.   Huang, G., Allen, R., Davis, E. L., Baum, T. J. & Hussey, R. S. Engineering broad

root-knot resistance in transgenic plants by RNAi silencing of a conserved and

essential root-knot nematode parasitism gene. *Proc Natl Acad Sci USA* **103,**

14302–14306 (2006).

7.   Baum, J. A. *et al.* Control of coleopteran insect pests through RNA interference.

Nat Biotechnol **25,** 1322–1326 (2007).

8.   Alakonya, A. *et al.* Interspecific RNA interference of SHOOT MERISTEMLESS-

like disrupts Cuscuta pentagona plant parasitism. *Plant Cell* **24,** 3153–3166

(2012).

9.   Weiberg, A. *et al.* Fungal small RNAs suppress plant immunity by hijacking host

RNA interference pathways. *Science* **342,** 118–123 (2013).

10. Wang, M. *et al.* Bidirectional cross-kingdom RNAi and fungal uptake of external

RNAs confer plant protection. *Nature Plants* **2,** 16151 (2016).

11.   Zhang, T. *et al.* Cotton plants export microRNAs to inhibit virulence gene

expression in a fungal pathogen. *Nature Plants* **2,** 16153 (2016).

12.   Kim, G., LeBlanc, M. L., Wafula, E. K., dePamphilis, C. W. & Westwood, J. H.

Plant science. Genomic-scale exchange of mRNA between a parasitic plant and

its hosts. *Science* **345,** 808–811 (2014).

13.   Chen, H.-M. *et al.* 22-Nucleotide RNAs trigger secondary siRNA biogenesis in

plants. *Proc Natl Acad Sci USA* **107,** 15269–15274 (2010).

14.   Cuperus, J. T. *et al.* Unique functionality of 22-nt miRNAs in triggering RDR6-

dependent siRNA biogenesis from target transcripts in Arabidopsis. *Nat Struct*

Mol Biol **17,** 997–1003 (2010).

15.   Dharmasiri, N. *et al.* Plant development is regulated by a family of auxin receptor

F box proteins. *Dev Cell* **9,** 109–119 (2005).

16.   Veronese, P. *et al.* The membrane-anchored BOTRYTIS-INDUCED KINASE1

plays distinct roles in Arabidopsis resistance to necrotrophic and biotrophic

pathogens. *Plant Cell* **18,** 257–273 (2006).

17.   Lin, W. *et al.* Inverse modulation of plant immune and brassinosteroid signaling

pathways by the receptor-like cytoplasmic kinase BIK1. *Proc Natl Acad Sci USA*

110, 12114–12119 (2013).

18.   Froelich, D. R. *et al.* Phloem ultrastructure and pressure flow: Sieve-Element-

Occlusion-Related agglomerations do not affect translocation. *Plant Cell* **23,**

4428–4445 (2011).

19.   Jekat, S. B. *et al.* P-proteins in Arabidopsis are heteromeric structures involved in

283   rapid sieve tube sealing. *Front Plant Sci* **4,** 225 (2013).

284 20. Mylona, P., Linstead, P., Martienssen, R. & Dolan, L. SCHIZORIZA controls an

285   asymmetric cell division and restricts epidermal identity in the Arabidopsis root.

286   *Development* **129,** 4327–4334 (2002).

287 21. Pernas, M., Ryan, E. & Dolan, L. SCHIZORIZA controls tissue system complexity

288   in plants. *Curr Biol* **20,** 818–823 (2010).

289 22. Hove, ten, C. A. *et al.* SCHIZORIZA encodes a nuclear factor regulating

290   asymmetry of stem cell divisions in the Arabidopsis root. *Curr Biol* **20,** 452–457

291   (2010).

292 23. Si-Ammour, A. *et al.* miR393 and secondary siRNAs regulate expression of the

293   TIR1/AFB2 auxin receptor clade and auxin-related development of Arabidopsis

294   leaves. *Plant Physiol* **157,** 683–691 (2011).

295 24. Allen, E., Xie, Z., Gustafson, A. M. & Carrington, J. C. microRNA-Directed

296   Phasing during Trans-Acting siRNA Biogenesis in Plants. *Cell* **121,** 207–221

297   (2005).

298 25. Axtell, M. J., Jan, C., Rajagopalan, R. & Bartel, D. P. A two-hit trigger for siRNA

299   biogenesis in plants. *Cell* **127,** 565–577 (2006).

300 26. Dawson, J. H., Musselman, L. J., Wolswinkel, P. & Dörr, I. Biology and control of

301   Cuscuta. *Reviews of Weed Science* **6,** 265–317 (1994).

302 27. Robert-Seilaniantz, A. *et al.* The microRNA miR393 re-directs secondary

303   metabolite biosynthesis away from camalexin and towards glucosinolates. *Plant J*

304   **67,** 218–231 (2011).

305 28. Lu, D. *et al.* A receptor-like cytoplasmic kinase, BIK1, associates with a flagellin

receptor complex to initiate plant innate immunity. *Proc Natl Acad Sci USA* **107,**

496–501 (2010).

Acknowledgements

We thank the Penn State & Huck Institutes Genomics Core Facility for small RNA-seq

services. We thank Hervé Vaucheret, Michael Knoblauch, Gabriele Monshausen,

Tesfaye Mengiste, and Renze Heidstra for gifts of *A. thaliana* mutant seed. We thank

Beth Johnson for advice on growing conditions for *C. campestris*. Purchase of the

Illumina HiSeq2500 used for small RNA-seq was funded by a major research

instrumentation award from the US National Science Foundation [1229046 to MJA and

CWD]. This research was supported in part by awards from the US National Science

Foundation [1238057 to JHW and CWD; 1339207 to MJA] and the National Institute of

Food and Agriculture [135997 to JHW].

Author Contributions: SS and MJA performed most bioinformatics analysis. SS, MJA,

and NRJ prepared figures and tables. GK and JHW cultivated and harvested plant

specimens used for initial small RNA-seq experiments. NRJ, SS, TP, and MJA

cultivated and harvested plant specimens for other experiments. EW, GK, CWD, and

JHW performed genome and transcriptome sequencing and preliminary assemblies.

FW, SS, and NRJ performed RNA blots. SS and MJA performed 5'-RLM-RACE and

qRT-PCR. CC and TP constructed small RNA-seq libraries. NRJ and VB-G performed

growth assays. MJA and JHW conceived of the project. MJA wrote and revised the

manuscript with input from all other authors.

Author Information

Correspondence and requests for materials should be addressed to Michael J. Axtell at

mja18@psu.edu

Extended Data

Extended Data Figure 1. PCR of *C. campestris MIRNA* loci. The template for PCR was

genomic DNA isolated from *C. campestris* seedlings four days after germination; the

seedlings had never attached to nor been near a host plant, ruling out host DNA

contamination. trnL-F: Positive control plastid locus.

Extended Data Figure 2. *C. campestris* miRNAs cause slicing and phased siRNA

production from host mRNAs. Small RNA-seq coverage across the indicated *A. thaliana*

transcripts are shown in blue for host stem (HS), interface (I), and parasite stem (PS)

samples. For display, the two biological replicates of each type were merged. y-axis is

in units of reads per million. Red marks and vertical lines show position of

complementary sites to *C. campestris* miRNAs, with the alignments shown above.

Fractions indicate numbers of 5'-RLM-RACE clones with 5'-ends at the indicated

positions; the locations in red are the predicted sites for miRNA-directed slicing

remnants. Bar charts show the length and polarity distribution of transcript-mapped

siRNAs. Radar charts show the fractions of siRNAs in each of the 21 possible phasing

registers; the registers highlighted in magenta are the ones predicted by the miRNA

target sites.

Cp_v0.1_Contig11111_02898.1 .. A *TIR/AFB* ortholog
```
5' CUGGUCUGCAAGUCCUGGUACG 3' Transcript: Cp_v0.1_Contig11111_02898.1:82-103 Slice Site:94
   || ||||o||o|| ||||||o
3' GAGCAGAUGUUUAGUACCAUGU 5' Query: Cluster_115407_miRNA-star

5' CUGGUCUGCAAGUCCUGGUACG 3' Transcript: Cp_v0.1_Contig11111_02898.1:82-103 Slice Site:94
   |||| ||||||o|| ||||||o
3' AACCAUACGUUUAGUACCAUGU 5' Query: Cluster_67631_miRNA
```

Cp_v0.1_Contig122651_07831.1 .. A *BIK1* ortholog
```
5' AUGGGUACGCCGCUCCCGAGUA 3' Transcript: Cp_v0.1_Contig122651_07831.1:668-689 Slice Site:680
   |||| ||||||||| || |||||
3' UACCGAUGCGGCGCGGACUCAU 5' Query: Cluster_54651_miRNA
```

Cp_v0.1_Contig297862_09445.1 .. A *TIR/AFB* ortholog
```
5' CUGGUCUGCAAGUCCUGGUACG 3' Transcript: Cp_v0.1_Contig297862_09445.1:82-103 Slice Site:94
   || ||||o||o|| ||||||o
3' GAGCAGAUGUUUAGUACCAUGU 5' Query: Cluster_115407_miRNA-star

5' CUGGUCUGCAAGUCCUGGUACG 3' Transcript: Cp_v0.1_Contig297862_09445.1:82-103 Slice Site:94
   |||| ||||||o|| ||||||o
3' AACCAUACGUUUAGUACCAUGU 5' Query: Cluster_67631_miRNA
```

Cp_v0.1_Contig3449_00601.1 ..A *TIR/AFB* ortholog
```
5' CUCGUCUGCAAGUCGUGGUACC 3' Transcript: Cp_v0.1_Contig3449_00601.1:259-280 Slice Site:271
   |||||||o||o||o||||||
3' GAGCAGAUGUUUAGUACCAUGU 5' Query: Cluster_115407_miRNA-star
```

Cp_v0.1_Contig370259_15766.1 .. An *SCZ/HSFB4* ortholog
```
5' CAGACAGCUCAACACAUACGGG 3' Transcript: Cp_v0.1_Contig370259_15766.1:225-246 Slice Site:237
   |||||||||||| ||||| ||o
3' CUCUGUCGAGUUAUGUAUACCU 5' Query: Cluster_57001_miRNA
```

Cp_v0.1_Contig501179_37185.1 .. An *SCZ/HSFB4* ortholog
```
5' CAGACAGCUCAACACAUACGGG 3' Transcript: Cp_v0.1_Contig501179_37185.1:213-234 Slice Site:225
   |||||||||||| ||||| ||o
3' CUCUGUCGAGUUAUGUAUACCU 5' Query: Cluster_57001_miRNA
```

Cp_v0.1_Contig70142_36225.1 .. An *SCZ/HSFB4* ortholog
```
5' CAGGCAGCUCAACACUUAUGGA 3' Transcript: Cp_v0.1_Contig70142_36225.1:300-321 Slice Site:312
   ||o||||||||| || ||||||
3' CUCUGUCGAGUUAUGUAUACCU 5' Query: Cluster_57001_miRNA
```

Cp_v0.1_Contig81842_24454.1 .. A *BIK1* ortholog
```
5' UUGUAUAUGAAUUUAUGCAGA 3' Transcript: Cp_v0.1_Contig81842_24454.1:608-628 Slice Site:619
   |||| |||||||| o||||||||
3' AACAAAUACUUAUGUACGUCU 5' Query: Cluster_105389_miRNA-star

5' CUUGUAUAUGAAUUUAUGCAGA 3' Transcript: Cp_v0.1_Contig81842_24454.1:607-628 Slice Site:619
   || || ||o|||||||||||o|
3' GAUCAGAUGCUUAAAUACGUUU 5' Query: Cluster_105391_miRNA

5' AAGGCUAUGCUGCCCCUGAGUA 3' Transcript: Cp_v0.1_Contig81842_24454.1:893-914 Slice Site:905
   |||||o||o|| |||||||||
3' UACCGAUGCGGCGCGGACUCAU 5' Query: Cluster_54651_miRNA
```

360   **Extended Data Figure 3.** Possible miRNA target sites within endogenous *C.*

361   *campestris* mRNAs. Note that none of these mRNAs had evidence of any secondary

362   siRNA accumulation, and the complementarity of these sites was generally poor.

Extended Data Figure 4. Growth of *C. campestris* on *A. thaliana sgs2-1* and *dcl4-2t*

mutants with varying methodologies, as indicated. P-values (Wilcoxon rank-sum tests,

unpaired, two-tailed) from comparison of mutant to wild-type (Col-0) are shown. Dots

show all data points. Boxplots represent medians (horizontal lines), the central half of

the data (boxes), and other data out to 1.5 times the interquartile range (whiskers). **a)**

n=16 and 9 for Col-0 and *sgs2-1*, respectively. **b)** n=10 and 8 for Col-0 and *dcl4-2t*,

respectively. **c)** n=11 and 10 for Col-0 and *dcl4-2t*, respectively. **d)** n=8 and 9 for Col-0

and *dcl4-2t*, respectively. **e)** n=14, 14, and 12 for Col-0, *sgs2-1*, and *dcl4-2t*,

respectively.

Extended Data Figure 5. Highly reproducible induction of *C. campestris MIRNA*s on

different hosts. **a)** MA plot from original experiment on *A. thaliana* hosts of *C. campestris*

small RNA loci comparing interface (I) to parasite stem (PS) samples. Significantly up-

regulated loci (alternative hypothesis: true difference > 2-fold, FDR <= 0.05) are

highlighted. Reproduced from Figure 1A. **b-c)** As in a, except for a new set of *A.*

thaliana hosts (b) or from an experiment using *Nicotiana benthamiana* as hosts (c). **d)**

Area-proportional Euler diagram showing overlaps of up-regulated *C. campestris* 21-22

nt *MIRNA* loci between the three small RNA-seq experiments. The locations of the six

MIRNA loci of special interest are highlighted in green.

Extended Data Figure 6. *C. campestris* miRNAs cause slicing and phased siRNA production from *Nicotiana benthamiana* mRNAs. Small RNA-seq coverage across the indicated *N. benthamiana* transcripts are shown in blue for host stem (HS), interface (I), and parasite stem (PS) samples. For display, the two biological replicates of each type were merged. y-axis is in units of reads per million. Red marks and vertical lines show position of complementary sites to *C. campestris* miRNAs, with the alignments shown above. Fraction indicates numbers of 5'-RLM-RACE clones with 5'-ends at the indicated positions; the locations in red are the predicted sites for miRNA-directed slicing remnants. Barcharts show the length and polarity distribution of transcript-mapped siRNAs. Radar charts show the fractions of siRNAs in each of the 21 possible phasing registers; the registers highlighted in magenta are the ones predicted by the miRNA target sites.

402
Extended Data Figure 7. Uncropped image of *N. benthamiana* 5'-RLM-RACE products.

Lanes with 'X' are irrelevant to this study. This is the uncropped version of the image in

Figure 4C.

Extended Data Figure 8. *C. campestris* pre-haustoria. **a)** *C. campestris* seedling

wound around a bamboo stake. **b)** The same seedling, removed from the stake to show

the prominent pre-haustorial bumps. Seedling was scarified, germinated on moist paper

towels for three days at ~28C, and then placed next to bamboo stake for four days with

far-red LED lighting. Approximately 30 such seedlings were used for the 'PH' RNA in

Figure 4B. Scales bars: 1mm.

Phytozome 11 species code	Species	BIK1	SEOR1	SCZ / HSFB4	TIR1	GAPDH
Org_Acoerulea	Auilegia coerulea	0	1	0	1	0
Org_Alyrata	Arabidopsis lyrata	1	1	0	1	0
Org_Athaliana	Arabidopsis thaliana	1	1	1	1	0
Org_BrapaFPsc	Brassica rapa	1	0	1	1	0
Org_Bstricta	Boechera stricta	1	0	0	1	0
Org_Cclementina	Citrus clementina	0	1	0	0	0
Org_Cgrandiflora	Capsella grandiflora	1	0	0	1	0
Org_Cpapaya	Carica papaya	1	0	1	1	0
Org_Crubella	Capsella rubella	1	0	0	1	0
Org_Csativus	Cucumus sativus	0	0	0	0	0
Org_Csinensis	Citrus sinensis	1	1	0	1	0
Org_Egrandis	Eucalyptus grandis	0	1	1	1	0
Org_Esalsugineum	Eutrema salsugineum	1	0	0	1	0
Org_Fvesca	Fragaria vesca	0	NA	NA	1	1
Org_Gmax	Glycine max	1	1	1	1	0
Org_Graimondii	Gossypium raimondii	1	0	1	1	0
Org_Kmarnieriana	Kalanchoe marnieriana	0	1	1	1	0
Org_Lusitatissimum	Linum usitatissimum	0	1	0	0	0
Org_Mdomestica	Malus domestica	0	NA	0	1	0
Org_Mesculenta	Manihot esculenta	1	1	0	1	0
Org_Mguttatus	Mimulus gattus	1	NA	0	1	0
Org_Mtruncatula	Medicago truncatula	1	1	0	1	0
Org_Ppersica	Prunus persica	0	NA	0	1	0
Org_Ptrichocarpa	Populus trichocarpa	1	1	1	1	0
Org_Pvulgaris	Phaseolus vulgaris	0	1	0	1	0
Org_Rcommunis	Ricinus communis	1	1	0	1	0
Org_Slycopersicum	Solanum lycopersicum	0	1	1	1	0
Org_Spurpurea	Salix purpurea	0	1	1	0	0
Org_Stuberosum	Solanum tuberosum	1	NA	1	1	0
Org_Tcacao	Theobroma cacao	0	1	1	1	0
Org_Vvinifera	Vitis vinifera	0	1	1	1	0

Extended Data Table 1. Predicted miRNA targets un multiple plant species. Targets

were predicted using targetdinder.pl, keeping all hits with a score of 4.5 or less.

Probable orthologs of the indicated Arabidopsis thaliana genes were found using

BLASTP against the 31 eudicot species present in Phytozome 11, simply keeping up to

the top 100 BLAST hits. miRNA queries were all mature miRNAs and miRNA*'s from C.

campestris interface-induced MIRNA loci. An entry of 'NA' means no probable orthologs

were recovered from a given species. A 1 means there was one or more predicted

target in that species, a 0 means there were 0 predicted targets. GAPDH orthogroup:

negative control.

METHODS

Germplasm

Cuscuta was initially obtained from a California tomato field, and seed stocks derived from self-pollination through several generations in the Westwood laboratory. The isolate was initially identified as *Cuscuta pentagona* (Engelm.) *C. pentagona* is very closely related to *C. campestris* (Yunck.), and the two are distinguished by microscopic differences in floral morphology; because of this they have often been confused[1]. We subsequently determined that our isolate is indeed *C. campestris*. *Arabidopsis thaliana* *sgs2-1* mutants[2] were a gift from Hervé Vaucheret (INRA Versailles, France). *A. thaliana dcl4-2t* mutants (GABI_160G05[3]) were obtained from the Arabidopsis Biological Resource Center (The Ohio State University, USA). *A. thaliana seor* mutants (GABI-KAT 609F04[4]) were a gift from Michael Knoblauch (Washington State University, USA). The *A. thaliana tir1-1/afb2-* and *afb3*-4 mutants[5] were a gift from Gabriele Monshausen (The Pennsylvania State University, USA). The *bik1* mutant[6] was a gift from Tesfaye Mengiste (Purdue University, USA). The *scz2* mutant[7] was a gift from Renze Heidstra (Wageningen University, The Netherlands). All *A. thaliana* mutants were in the Col-0 background.

Growth conditions and RNA extractions

For initial experiments (small RNA-seq and RNA blots in Figure 1) *A. thaliana* (Col-0) plants were grown in a growth room at 18-20°C with 12-h light per day, illuminated (200 µmol m^{-2}s^{-1}) with metal halide (400W, GE multi-vapor lamp) and spot-gro (65W, Sylvania) lamps. *C. campestris* seeds were scarified in concentrated sulfuric

acid for 45 min, followed by 5-6 rinses with distilled water and dried. *C. campestris*

seeds were placed in potting medium at the base of four-week-old *A. thaliana* seedlings

and allowed to germinate and attach to hosts. The *C. campestris* plants were allowed to

grow and spread on host plants for an additional three weeks to generate a supply of

uniform shoots for use in the experiment. Sections of *C. campestris* shoot tip (~10 cm

long) were placed on the floral stem of a fresh set of *A. thaliana* plants. Parasite shoots

coiled around the host stems and formed haustorial connections. Tissues from plants

that had established *C. campestris* with at least two coils around healthy host stems and

clear parasite growth were used in these studies. Control plants were grown under the

same conditions as parasitized plants, but were not exposed to *C. campestris*.

For the preparation of tissue-specific small RNA libraries, tissues were harvested

after *C. campestris* cuttings had formed active haustorial connections to the host. This

was evidenced by growth of the *C. campestris* shoot to a length of at least 10 cm

beyond the region of host attachment (7-10 d after infection). Three tissues were

harvested from the *A. thaliana-C. campestris* associations: 1) 2.5 cm of *A. thaliana* stem

above the region of attachment, 2) *A. thaliana* and *C. campestris* stems in the region of

attachment (referred to as the interface), 3) 2.5 cm of the parasite stem near the point of

attachment. To remove any possible cross-contamination between *A. thaliana* and *C.*

campestris, harvested regions of the parasite and host stem were taken 1 cm away

from the interface region and each harvested tissue was surface cleaned by immersion

for 5 min in 70% ethanol, the ethanol was decanted and replaced, the process was

repeated three times and the stems were blotted dry with a Kimwipe after the final rinse.

All three sections of tissue were harvested at the same time and material from 20

attachments were pooled for small RNA extraction. Small RNA was extracted from ~

100 mg of each tissue using the mirPremier microRNA Isolation Kit (Sigma-Aldrich, St.

Louis, MO, USA) according to the manufacturer's protocol. Small RNA was analyzed

using an Agilent small RNA Kit on a 2100 Bioanalyzer platform.

Samples used for RNA ligase-mediated 5' rapid amplification of cDNA ends (5'-

RLM-RACE; Figure 2D), quantitative reverse-transcriptase polymerase chain reaction

(qRT-PCR; Figure 3A) analyses of *A. thaliana* targets were prepared as described

above with the following modifications: Col-0 *A. thaliana* hosts were cultivated in a

growth room with 16 hr. days, 8 hr. nights, at ~23C, under cool-white fluorescent lamps,

attachment of *C. campestris* cuttings was promoted by illumination with far-red LED

lighting for 3-5 days, and total RNA was extracted using Tri-reagent (Sigma) per the

manufacturer's suggestions, followed by a second sodium-acetate / ethanol precipitation

and wash step. Samples used for RNA blots of secondary siRNA accumulation from *A.*

thaliana mutants and replicate small RNA-seq libraries were obtained similarly, except

that the samples derived from the primary attachments of *C. campestris* seedlings on

the hosts instead of from cuttings. In these experiments, scarified *C. campestris*

seedlings were first germinated on moistened paper towels for three days at ~28C, then

placed adjacent to the host plants with their radicles submerged in a water-filled

0.125ml tube.

C. campestris pre-haustoria (Extended Data Figure 8) were obtained by

scarifying, germinating and placing seedlings as described above, next to bamboo

stakes in soil, under illumination from cool-white fluorescent lights and far-red emitting

LEDs. Seedlings coiled and produced pre-haustoria four days after being placed, and

were harvested and used for total RNA extraction (used for RNA blot in Figure 4B)

using Tri-reagent as described above. *Nicotiana benthamiana* was grown in a growth

room with 16 hr. days, 8 hr. nights, at ~23C, under cool-white fluorescent lamps. Three-

to four-week old plants served as hosts for scarified and germinated *C. campestris*

seedlings. Attachments were promoted by three-six days with supplementation by far-

red emitting LEDs. Under these conditions, *C. campestris* attached to the petioles of the

N. benthamiana hosts, not the stems. Interfaces and control petioles from un-

parasitized hosts were collected 7-8 days after successful attachments, and total RNA

(used for RNA blots in Figure 4B and small RNA-seq libraries) recovered using Tri-

reagent as described above.

small RNA-seq

The initial small RNA-seq libraries were constructed using the Illumina Tru-Seq

small RNA kit per the manufacturer's protocol and sequenced on an Illumina HiSeq2500

instrument. Subsequent small RNA-seq libraries (replicate two using *A. thaliana* hosts,

and the *N. benthamiana* experiments) instead used New England Biolabs NEBnext

small RNA library kit, following the manufacturer's instructions. Raw sRNA-seq reads

were trimmed to remove 3'-adapters, and filtered for quality and trimmed length >= 16

nts using cutadapt[8] version 1.9.1 with settings "-a TGGAATTCTCGGGTGCCAAGG --

discard-untrimmed -m 16 --max-n=0". For experiments where *A. thaliana* was the host,

trimmed reads that aligned with zero or one mismatch (using bowtie[9] version 1.1.2,

settings "-v 1") to the *A. thaliana* plastid genome, the *C. gronovii* plastid genome (*C.*

gronovii was the closest relative to *C. campestris* that had a publically available

completed plastid genome assembly available), *A. thaliana* rRNAs, tRNAs, snRNAs, or

snoRNAs were removed. Similarly, for experiments where *N. benthamiana* was the

host, the reads were cleaned against the *C. gronovii* plastid genome, the *N. tabacum*

plastid genome and rRNAs, and a set of tRNAs predicted from the *N. benthamiana*

genome using tRNAscanSE.

For the original *A. thaliana* host data, the 'clean' reads were aligned and

analyzed with reference to the combined TAIR10 *A. thaliana* reference genome and a

preliminary version 0.1 draft genome assembly of *C. campestris* using ShortStack[10]

(version 3.8.3) using default settings. The resulting annotated small RNA loci

(Supplementary Data 1) were analyzed for differential expression (I vs. PS) using

DESeq2[11], with a $\log_2$ fold threshold of 1, alternative hypothesis of "greaterAbs", and

alpha of 0.05. p-values were adjusted for multiple testing using the Benjamini-Hochberg

procedure, and loci with an adjusted p-value of <= 0.05 (equivalent to an FDR of <=

0.05) were called up-regulated in I relative to PS. Among the up-regulated loci, those

annotated by ShortStack as *MIRNA*s deriving from the *C. campestris* genome which

produced either a 21nt or 22nt mature miRNA (Supplementary Data 2) were retained

and further analyzed. The predicted secondary structures and observed small RNA-seq

read coverage was visualized (Supplementary Data 3-4) using strucVis (version 0.3;

https://github.com/MikeAxtell/strucVis).

For analysis of mRNA-derived secondary siRNAs, the 'clean' small RNA-seq

reads from the original *A. thaliana* experiment were aligned to the combined TAIR10

representative cDNAs from *A. thaliana* and our preliminary version 0.1 transcriptome

assembly for *C. campestris*, using ShortStack[10] version 3.8.3, with settings --

mismatches 0, --nohp, and defining the full length of each mRNA as a 'locus' using

option --locifile. The resulting counts of small RNA alignments for each mRNA were

used for differential expression analysis, comparing I vs. HS, using DESeq2 [11] as

described above. *A. thaliana* mRNAs with significantly up-regulated (FDR <= 0.05)

small RNAs comparing I vs. HS were retained for further analysis. The cDNA

sequences of these loci were retrieved, and used for miRNA target predictions using

GSTAr (version 1.0; https://github.com/MikeAxtell/GSTAr); the full set of mature

miRNAs and miRNA*'s (Supplementary Data 2) from the I-induced *C. campestris*

MIRNA loci were used as queries.

Analysis of the second set of *A. thaliana* - *C. campestris* small RNA-seq data

aligned the cleaned reads to the combined *A. thaliana* and *C. campestris* reference

genomes as described above, except that the list of loci derived in the analysis of the

original data (Supplementary Data 1) was used as a "--locifile" in the ShortStack

analysis. Differential expression analysis was then performed using DESeq2 as

described above. Analysis of the *N. benthamiana* - *C. campestris* small RNA-seq data

began with a ShortStack analysis of the cleaned reads against the combined *N.*

benthamiana (version 0.4.4) genome and the preliminary assembly of the *C. campestris*

genome, using default settings. The *de novo N. benthamiana* loci obtained from this run

were retained. The resulting alignments were used to quantify small RNA abundance

from the *C. campestris* small RNA loci defined with the original data. The resulting read

counts were then used for differential expression analysis with DESeq2 as described

above. Analysis of secondary siRNAs derived from *N. benthamiana* mRNAs was

performed similarly as the *A. thaliana* mRNA analysis described above, except that the

combined transcriptomes were from *C. campestris* and *N. benthamiana* (version 0.4.4

annotations).

RNA blots

Small RNA gel blots were performed as previously described[12] with

modifications. For the blots in Figure 1B, small RNAs (1.8 micrograms) from each

sample were separated on 15% TBE-Urea Precast gels (Bio-Rad), transblotted onto the

Hybond NX membrane and cross-linked using 1-ethyl-3-(3-dimethylamonipropyl)

carbodiimide[13]. Hybridization was carried out in 5×SSC, 2×Denhardt's Solution, 20 mM

sodium phosphate (pH 7.2), 7% SDS with 100 µg/ml salmon testes DNA (Sigma-

Aldrich). Probe labeling, hybridization and washing were performed as described[12].

Radioactive signals were detected using Typhoon FLA 7000 (GE Healthcare).

Membranes were stripped in between hybridizations by washing with 1% SDS for 15

min at 80°C and exposed for at least 24 h to verify complete removal of probe before re-

hybridization. Sequences of probes are listed below. Blots in Figures 3B and 4B were

performed similarly, except that 12 micrograms of total RNA were used instead. Probe

sequences are listed in Supplementary Data 6.

5' RNA ligase-mediated rapid amplification of cDNA ends (5'-RLM-RACE)

Five micrograms of total RNA were ligated to one microgram of a 44 nucleotide

RNA adapter (Supplementary Data 6) using a 20ul T4 RNA ligase 1 reaction (NEB) per

the manufacturer's instructions for a one-hour incubation at 37C. The reaction was then

diluted with 68ul of water and 2ul of 0.5M EDTA pH 8.0, and incubated at 65C for 15

minutes to inactivate the ligase. Sodium acetate pH5.2 was added to a final

concentration of 0.3M, and the RNA precipitated with ethanol. The precipitated and

washed RNA was resuspended in 10ul of water. 3.33ul of this sample was used as the

template in a reverse transcription reaction using random primers and the Protoscript II

reverse transcriptase (NEB) per the manufacturer's instructions. The resulting cDNA

was used as template in first round PCR using a 5' primer matching the RNA adapter

and a 3' gene-specific primer (Supplementary Data 6). 1ul of the first round PCR

product was used as the template for nested PCR with nested primers (Supplementary

Data 6). Gene-specific primers for *A. thaliana* cDNAs were based on the representative

TAIR10 transcript models, while those for *N. benthamiana* cDNAs were based on the

version 0.4.4 transcripts (Sol Genomics Network[14]). In Figure 4c, *N. benthamiana*

TIR/AFB is transcript ID *NbS00011315g0112.1*; *N. benthamiana ARF* is transcript ID

NbS00059497g0003.1. Bands were gel-purified from agarose gels and cloned into

pCR4-TOPO (Life Tech). Inserts from individual clones were recovered by colony PCR

and subject to Sanger sequencing.

Quantitative reverse-transcriptase PCR (qRT-PCR)

Total RNA used for qRT-PCR was first treated with DNaseI (RNase-free; NEB)

per the manufacturer's instructions, ethanol precipitated, and resuspended. 2

micrograms of treated total RNA was used for cDNA synthesis using the High Capacity

cDNA Synthesis Kit (Applied Biosystems) per the manufacturer's instructions. PCR

reactions used PerfeCTa SYBR Green FastMix (Quanta bio) on an Applied Biosystems

StepONE-Plus quantitative PCR system per the manufacturer's instructions. Primers

(Supplementary Data 6) were designed to span the miRNA target sites, to ensure that

only uncleaved mRNAs were measured. Three reference mRNAs were used: *ACTIN*,

PP2A (PP2A sub-unit PDF2; *At1g13320*), and *TIP41-l* (*TIP41-like; At4g34270*) [15]. Raw

Ct values were used to calculate relative normalized expression values to each

reference mRNA separately, and the final analysis took the median relative expression

values between the *ACTIN-* and *TIP41-l* normalized data.

C. campestris growth assays

C. campestris seedlings were scarified, pre-germinated, and placed next to hosts

in 0.125ml water-filled tubes under cool-white fluorescent lighting supplemented with

far-red emitting LEDs (16hr day, 8hr night) at ~ 23C as described above. After a single

attachment formed (4 days), far-red light supplementation was removed to prevent

secondary attachments. After 18 more days of growth, entire *C. campestris* vines were

removed and weighed (Figure 3C). Multiple additional growth trials were performed

specifically on the *dcl4-2t* and *sgs2-1* mutant hosts under varying conditions (Extended

Data Figure 4).

miRNA target predictions

To find probable orthologs for *Arabidopsis thaliana* genes of interest, the *A.*

thaliana protein sequences were used as queries for a BLASTP analysis of the 31

eudicot proteomes available on Phytozome 11

(https://phytozome.jgi.doe.gov/pz/portal.html#). Transcript sequences for the top 100

hits were retrieved. In some cases no hits from a particular species were found; these

are 'N/A' on Figure 4a. The miRNA query set was all mature miRNAs and miRNA*s

from the I-induced, *C. campestris*-derived 21nt or 22nt *MIRNA*s (Supplementary Data

2). Targets were predicted from the probable 31-species with a maximum score of 4.5

using targetfinder.pl (https://github.com/MikeAxtell/TargetFinder/) version 0.1.

N. benthamiana orthologs of *A. thaliana* proteins were found based on BLAST-P

searches against the version 0.4.4 *N. benthamiana* protein models at Sol Genomics

Network[14], and miRNA target sites predicted using targetfinder.pl as above.

Code availability

ShortStack[10] (small RNA-seq analysis), strucVis (visualization of predicted RNA

secondary structures with overlaid small RNA-seq depths), and

Shuffler.pl/targetfinder.pl (prediction of miRNA targets controlling for false discovery

rate) are all freely available at https://github.com/MikeAxtell. Cutadapt version 1.9.1[8] is

freely available at http://cutadapt.readthedocs.io/en/stable/index.html. The R package

DESeq2[11] is freely available at http://www.

bioconductor.org/packages/release/bioc/html/DESeq2.html.

Data Availability

Small RNA-seq data from this work are available at NCBI GEO under accession

GSE84955 and NCBI SRA under project PRJNA408115. The draft, preliminary *C.*

campestris genome and transcriptome assemblies used in this study are available at

the Parasitic Plant Genome Project website at http://ppgp.huck.psu.edu.

References (Methods)

1.   Costea, M., García, M. A., Baute, K. & Stefanović, S. Entangled evolutionary

history of Cuscuta pentagona clade: A story involving hybridization and Darwin in

the Galapagos. *Taxon* **64,** 1225–1242 (2015).

2.   Elmayan, T. *et al.* Arabidopsis mutants impaired in cosuppression. *Plant Cell* **10,**

1747–1758 (1998).

3.   Xie, Z., Allen, E., Wilken, A. & Carrington, J. C. DICER-LIKE 4 functions in trans-

acting small interfering RNA biogenesis and vegetative phase change in

Arabidopsis thaliana. *Proc Natl Acad Sci USA* **102,** 12984–12989 (2005).

4.   Froelich, D. R. *et al.* Phloem ultrastructure and pressure flow: Sieve-Element-

Occlusion-Related agglomerations do not affect translocation. *Plant Cell* **23,**

4428–4445 (2011).

5.   Parry, G. *et al.* Complex regulation of the TIR1/AFB family of auxin receptors.

Proc Natl Acad Sci USA **106,** 22540–22545 (2009).

6.   Veronese, P. *et al.* The membrane-anchored BOTRYTIS-INDUCED KINASE1

plays distinct roles in Arabidopsis resistance to necrotrophic and biotrophic

pathogens. *Plant Cell* **18,** 257–273 (2006).

7.   Hove, ten, C. A. *et al.* SCHIZORIZA encodes a nuclear factor regulating

asymmetry of stem cell divisions in the Arabidopsis root. *Curr Biol* **20,** 452–457

(2010).

8.   Martin, M. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet. journal* (2011).

9.   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-

efficient alignment of short DNA sequences to the human genome. *Genome Biol*

10, R25 (2009).

10.   Johnson, N. R., Yeoh, J. M., Coruh, C. & Axtell, M. J. Improved Placement of

Multi-Mapping Small RNAs. *G3 (Bethesda)* (2016). doi:10.1534/g3.116.030452

11.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* **15,** 550 (2014).

12.   Cho, S. H., Coruh, C. & Axtell, M. J. miR156 and miR390 regulate tasiRNA

accumulation and developmental timing in Physcomitrella patens. *Plant Cell* **24,**

4837–4849 (2012).

13.   Pall, G. S. & Hamilton, A. J. Improved northern blot method for enhanced

detection of small RNA. *Nat Protoc* **3,** 1077–1084 (2008).

14.   Bombarely, A. *et al.* A draft genome sequence of Nicotiana benthamiana to

enhance molecular plant-microbe biology research. *Mol Plant Microbe Interact* **25,**

1523–1530 (2012).

15.   Czechowski, T., Stitt, M., Altmann, T., Udvardi, M. K. & Scheible, W.-R. Genome-

wide identification and testing of superior reference genes for transcript

normalization in Arabidopsis. *Plant Physiol* **139,** 5–17 (2005).

SI Guide

Supplementary Data 1: *A. thaliana* and *C. campestris* small RNA loci. Output from ShortStack 3.8.3 (https://github.com/MikeAxtell/ShortStack) showing all small RNA loci identified in this study. (.xlsx format)

Supplementary Data 2: Mature miRNAs and miRNA*s from *C. campestris* Interface-induced *MIRNA* loci. (.xlsx format)

Supplementary Data 3: Details of *C. campestris MIRNA* loci: Text-based sequences, predicted secondary structures, and aligned small RNA reads (all six libraries). Lower-case letters indicate small RNA bases that are mismatched to the genomic sequence. Plain-text (ASCII) format.

Supplementary Data 4: Images showing *C. campestris* hairpins overlaid with color-codes representing total read-depth (all six 'original' *C. campestris* x *A. thaliana* small RNA libraries). (pdf, 43 pages).

Supplementary Data 5: Alignments of mature miRNAs and miRNA*s (from Supplementary Data 2 -- the induced *C. campestris* miRNAs) against the *A. thaliana* genome. Note that most have three or more mismatches. (SAM format).

Supplementary Data 6: Oligonucleotide sequences. Excel (.xlsx) format.