The molecular mechanism of N-acetylglucosamine side-chain attachment to the

Lancefield group A Carbohydrate in *Streptococcus pyogenes*.

Jeffrey S. Rush[a], Rebecca J. Edgar[a], Pan Deng[b], Jing Chen[a], Haining Zhu[a],

Nina M. van Sorge[c], Andrew J. Morris[b], Konstantin V. Korotkov[a], and Natalia

Korotkova[a#]

Department of Molecular and Cellular Biochemistry, University of Kentucky,

Lexington, Kentucky, USA[a]; Division of Cardiovascular Medicine and the Gill Heart

Institute, University of Kentucky Lexington, Kentucky, USA[b]; Department of Medical

Microbiology, University Medical Center Utrecht, Utrecht, The Netherlands[c].

[#]Address correspondence to Natalia Korotkova, nkorotkova@uky.edu

Running Head: Mechanism of cell wall modification with GlcNAc in GAS

The word count for the abstract: 196

The word count for the text: 12387

Abstract

Streptococcus pyogenes or Group A *Streptococcus* (GAS) causes numerous diseases in humans, ranging from minor skin and throat infections, to life-threatening systemic conditions such as bacteremia, necrotizing fasciitis, and toxic-shock syndrome. GAS synthesizes a key antigenic surface polymer — the Lancefield group A Carbohydrate (GAC). GAC is attached to peptidoglycan and consists of a polyrhamnose polymer, with N-acetylglucosamine (GlcNAc) side chains, that is an essential virulence determinant. The molecular details of the mechanism of polyrhamnose modification with GlcNAc are currently unknown. In this report we demonstrate that GAC biosynthesis requires two distinct undecaprenol-linked GlcNAc-lipid intermediates: GlcNAc-pyrophosphoryl-undecaprenol (GlcNAc-P-P-Und) produced by GacO and GlcNAc-phosphate-undecaprenol (GlcNAc-P-Und) produced by GacI. The GAC polyrhamnose backbone is assembled on GlcNAc-P-P-Und. Furthermore, our data suggests that GlcNAc-P-Und is used by the membrane glycosyltransferase GacL to transfer GlcNAc from GlcNAc-P-Und to the polyrhamnose polysaccharide. In addition, GacJ, a small membrane protein, forms a complex with GacI and significantly stimulates its catalytic activity. We show that GlcNAc modification of polyrhamnose protects GAS from amidase-induced lysis. Thus, our study significantly expands our understanding of the biosynthesis of GAS cell wall polysaccharide and points to the functional importance of polysaccharide modifications in protection of peptidoglycan from lytic enzymes.

IMPORTANCE

In many species of Lactobacillales peptidoglycan is decorated by polyrhamnose polysaccharides that are critical for cell envelope integrity and cell shape and represent key antigenic determinants. Despite the biological importance of these polysaccharides, their biosynthetic pathways have received limited study. The important human pathogen, *Streptococcus pyogenes*, synthesizes a cell wall-associated polyrhamnose polysaccharide with N-acetylglucosamine (GlcNAc) side chains. The GlcNAc side chain is an important virulence determinant used by bacteria to evade the host innate immune defense. Here we identify the molecular mechanism of polyrhamnose modification with GlcNAc in *S. pyogenes*. We show that GacI synthesizes GlcNAc-phosphate-undecaprenol (GlcNAc-P-Und) aided by GacJ. GlcNAc-P-Und is apparently used by

GacL to transfer GlcNAc to polyrhamnose. We demonstrate that GacI homologs

perform a similar function in *Streptococcus agalactiae* and *Enterococcus faecalis*. Thus,

the elucidation of polysaccharide biosynthesis in *S. pyogenes* enhances our

understanding of how other Gram-positive bacteria produce essential components of

their cell wall.

INTRODUCTION

The cytoplasmic membrane of Gram-positive bacteria is surrounded by a thick cell

wall consisting of multiple peptidoglycan layers decorated with proteins and a variety of

carbohydrate-based polymers. Rhamnose (Rha) is the main component of the

carbohydrate structures in many species of the Lactobacillales order (1). These

polymers play essential roles in maintaining and protecting bacterial cell envelopes and

in pathogenesis of infections caused by *Streptococcus pyogenes* and *Enterococcus*

faecalis (1). It is of paramount importance to understand the molecular mechanisms of

Rha-containing cell wall polysaccharide biosynthesis for developing novel therapeutics

against these bacterial pathogens.

S. pyogenes or Group A *Streptococcus* (GAS) is associated with numerous

diseases in humans ranging from minor skin and throat infections such as impetigo and

pharyngitis to life-threatening invasive infections such as streptococcal toxic syndrome

and necrotizing fasciitis (2). The main component of GAS cell wall is the Lancefield

group A Carbohydrate (GAC) that comprises about 40-60% of the total cell wall mass

(3). Serological grouping of beta-hemolytic streptococci (A, B, C, E, F, and G groups),

introduced by Rebecca Lancefield in 1933, is based on the detection of carbohydrate

antigens present on the cell wall (4). GAC is presumably covalently linked to N-

acetylmuramic acid (MurNAc) of peptidoglycan, although the molecular structure of the

linkage unit has not been established (5). All serotypes of GAS produce GAC consisting

of a polyrhamnose backbone with N-acetylglucosamine (GlcNAc) side chains (6, 7). The

average molecular mass of GAC has been reported to be 8.9±1.0 kDa, corresponding

to an average of 18 repeating units of [→3)α-Rha(1→2)[β-GlcNAc(1→3)]α-Rha(1→] (8).

Since Rha is absent in mammalian cells, GAC is an attractive candidate for a universal

GAS vaccine (8, 9). Moreover, the GlcNAc side chains are an important virulence

determinant in GAS (5). GAS mutants lacking GlcNAc are susceptible to innate immune

clearance by neutrophils and antimicrobial agents, and are significantly attenuated in

animal models of GAS infection (5).

Similar to the biosynthesis of other cell-envelope polymers — peptidoglycan,

lipopolysaccharide, wall teichoic acid (WTA), and capsular polysaccharide — the

synthesis of Rha-containing carbohydrate structures is likely initiated on the inside of

the cytoplasmic membrane and proceeds through several steps including the

attachment of the first sugar residue to a lipid carrier, undecaprenyl phosphate (Und-P),

followed by elongation of the polysaccharide through stepwise addition of activated

sugar residues to the lipid carrier (1). After translocation of the polysaccharide across

the membrane and further modifications, it is presumably attached to peptidoglycan via

a phosphate ester linkage (1). The first membrane step of O-antigen biosynthesis in

enterobacteria and WTA biosynthesis in *Bacillus subtilis* and *Staphylococcus aureus*

(10) is catalyzed by the UDP-GlcNAc:Und-P GlcNAc-1-phosphate transferase encoded

by WecA homologs (11, 12). WecA transfers GlcNAc-phosphate from UDP-GlcNAc to

Und-P, forming GlcNAc-pyrophosphoryl-undecaprenol (GlcNAc-P-P-Und). Although

GAS strains do not produce WTA, all GAS genomes contain a *wecA* homolog, *gacO*

(5). Significantly, GAC biosynthesis is sensitive to tunicamycin, a known inhibitor of

WecA (5) and the *Streptococcus mutans* GacO homolog, RgpG, has been shown to

complement WecA activity in *E. coli* (13). These observations support an essential role

for GacO in the biosynthesis of GlcNAc-P-P-Und and suggest GlcNAc-P-P-Und may

function as a lipid-anchor in the initiation of GAC biosynthesis.

Other genes required for GAC biosynthesis and transport are found in a separate

location on the GAS chromosome and comprise a 12-gene locus (*gacA-gacL*) (Fig.1A).

The first three genes, *gacA-gacC* together with *gacG,* are conserved in many species of

the Lactobacillales order (1). *GacA* encodes a dTDP-4-dehydrorhamnose reductase,

the enzyme responsible for dTDP-rhamnose biosynthesis (14). *GacB*, *gacC*, and *gacG*

encode putative cytoplasmic rhamnosyltransferases. In *S. mutans* homologous genes

rgpA, *rgpB* and *rgpF* are involved in rhamnan backbone biosynthesis (15). In GAS,

gacA-gacC are essential for viability and cannot be deleted (5). However, deletion

mutants have been obtained in other genes of the GAC gene cluster (5). It has been

shown that *gacI*, *gacJ* and *gacK* are non-essential for viability, but are required for

GlcNAc side-chain addition to the polyrhamnose (5). They encode a putative

cytoplasmic glycosyltransferase, a small membrane protein and a protein with homology

to the Wzx family of membrane proteins involved in the export of O-antigen and teichoic

acids, respectively. In contrast, inactivation of *gacD*, *gacE*, *gacF*, *gacG*, *gacH* or *gacL*

has no effect on GAS viability and the GAC produced by these mutants is reported to

display a wild type (WT) antigenic profile, indicating the presence of the

immunodominant GlcNAc side chains (5). *GacD* and *gacE* encode the components of

an ABC transport system. In *S. mutans* the *gacD* and *gacE* homologs are responsible

for rhamnan polysaccharide transport (15). *GacF*, *gacH* and *gacL* encode a cytosolic

glycosyltransferase, a putative membrane-associated glycerol phosphate transferase

and a membrane-associated glycosyltransferase, respectively.

In this study we address the molecular mechanism of GlcNAc attachment to

polyrhamnose. We demonstrate that GacI catalyzes formation of GlcNAc-P-Und, and

GacJ stimulates the catalytic activity of GacI. Subsequently GacL transfers GlcNAc from

GlcNAc-P-Und to the polyrhamnose backbone of GAS polysaccharide. Moreover, we

confirm GacO function in GlcNAc-P-P-Und formation and demonstrate the role of this

GlcNAc-lipid in initiation of polyrhamnose biosynthesis.

RESULTS

GacL is required for GlcNAc attachment to polyrhamnose. GacL, a polytopic (12–

13 transmembrane segments) membrane protein, is reported to be dispensable for

GlcNAc attachment to polyrhamnose (5). To investigate the function of GacL in GAC

biogenesis, we disrupted *gacL* in the hyperinvasive *S. pyogenes* M1T1 serotype strain,

MGAS5005 (16), creating the 5005Δ*gacL* mutant. In agreement with published data (5)

5005Δ*gacL* did not display any detectable growth phenotype in comparison to the WT

strain. Surprisingly, we found that 5005Δ*gacL* cells failed to bind GlcNAc-specific anti-

GAC antibodies, suggesting a loss of the GlcNAc antigenic epitope (Fig. 1B). The

5005Δ*gacL* phenotype was restored by expressing the WT copy of *gacL* on the mutant

chromosome (Fig. 1B). To confirm that 5005Δ*gacL* cells are deficient in GlcNAc addition

to polysaccharide, we measured the binding of fluorescently labeled succinylated wheat

germ agglutinin (sWGA), a lectin that specifically binds non-reducing terminal β-GlcNAc residues, to WT and 5005Δ*gacL* cells. Deletion of *gacL* led to a significant decrease in binding of WGA to the 5005Δ*gacL* strain as compared with the WT (Fig. 1C), indicating that the mutant has substantially less GlcNAc-containing saccharides on the cell surface. Furthermore, direct compositional analysis of cell wall purified from WT and 5005Δ*gacL* cells confirmed a significant decrease of GlcNAc content in the 5005Δ*gacL* cell wall sample (Fig.1D, E and F). These data strongly support a role for GacL in GlcNAc side-chain attachment to polyrhamnose.

A *gacL* deletion mutant accumulates GlcNAc-P-Und. To investigate further the effect of GacL deletion in *S. pyogenes*, we isolated the phospholipid fractions from the membranes of WT and 5005Δ*gacL* cells. TLC analysis revealed accumulation of a previously unidentified phosphoglycolipid in 5005Δ*gacL* (Fig. 2A). The novel lipid was found to be stable to mild alkaline methanolysis, but sensitive to mild acid (0.1 N HCl, 50 ºC, 50% isopropanol) and reacted with orcinol spray (17), Dittmer-Lester phospholipid spray reagent (18) and with anisaldehyde, an isoprenol-specific reagent (19), consistent with its tentative identification as a glycophosphoprenol (data not shown). High-resolution negative ion ESI-MS analysis of the purified lipid identified a molecular ion $[M-H]^{-1}$ of M/z = 1048.73621 (Fig. 2B). In addition, prominent high-resolution fragment ions characteristic of PO_3, H_2PO_4, N-acetyl hexosamine-phosphate and Und-P were also found (Table 1). These data are consistent with a glycolipid comprised of N-acetyl hexosamine-phosphate-undecaprenol. In experiments described in detail below, we found that when membrane fractions from GAS are incubated with UDP-$[^3H]$GlcNAc, a $[^3H]$GlcNAc-lipid which co-chromatographs on TLC with this novel lipid is rapidly formed. Importantly, the product ion spectra shown in Figure 2B does not contain a fragment ion containing $[Und-PO_4-C_2H_2NHCOCH_3]^-$ (M/z=929.7), arising from a cross-ring fragmentation reaction, suggesting that the hydroxyl of the anomeric carbon and the nitrogen at the 2-position of the glycosyl ring are *trans* to the plane of the ring (20, 21). This observation strongly suggests that the anomeric hydroxyl is present in the β configuration. Furthermore, isolated $[^3H]$GlcNAc-P-Und phospholipid (described in detail below) is extremely sensitive to incubation with 50% phenol at 68 °C — a property that is consistent with β-GlcNAc-P-Und (22) (Table S4). Taken together, these data

identify the novel glycolipid accumulating in the 5005Δ*gacL* strain as β-GlcNAc-P-Und

and suggest that GacL is a GlcNAc transferase using β-GlcNAc-P-Und as GlcNAc

donor for the addition of the GlcNAc side-chains to GAC.

GAS synthesizes two GlcNAc-lipids *in vitro*. To investigate the biosynthetic

origin of the novel glycolipid accumulating in 5005Δ*gacL* cells, membrane fractions from

MGAS5005 were incubated with UDP-[³H]GlcNAc and analyzed for lipid products as

described in Material and Methods. Experiments showed that [³H]GlcNAc was efficiently

transferred from UDP-[³H]GlcNAc into two detectable GlcNAc-lipids (Fig. 3A, Table 2).

The major lipid product co-migrates on TLC with the novel lipid accumulating in the

5005Δ*gacL* strain and is identified as β-[³H]GlcNAc-P-Und, as described above. The

minor product is assumed to be [³H]GlcNAc-P-P-Und, since it co-migrates with authentic

GlcNAc-P-P-Und synthesized in *B. cereus* membranes (Fig. 3B) (23) and its formation

is potently inhibited by tunicamycin (Fig. S1).

GacI encodes a UDP-GlcNAc:Und-P GlcNAc transferase activity. The GAS

polysaccharide gene cluster contains a gene*, gacI*, reported to be required for GlcNAc

addition to cell wall polysaccharide (5) and annotated as a GT-A type

glycosyltransferase. The following studies were conducted to determine if GAS GacI

might be responsible for the synthesis of GlcNAc-P-Und. To determine if GacI function

was essential for formation of GlcNAc-P-Und, we generated a deletion of *gacI* in the

MGAS5005. In agreement with published data (5) 5005Δ*gacI* lost reactivity with anti-

GAC antibodies, indicating a loss of GlcNAc modification in polyrhamnose (Fig. 1B).

When 5005Δ*gacI* membrane fractions were incubated with [³H]UDP-GlcNAc no

incorporation into [³H]GlcNAc-P-Und was observed and only the minor lipid,

[³H]GlcNAc-P-P-Und was found (Fig. 3C, Table 2), strongly supporting an essential role

for GacI in GlcNAc-P-Und synthesis. The observation that the 5005Δ*gacL* strain

synthesizes normal levels of [³H]GlcNAc-P-Und *in vitro* (Fig. 3D) indicates that

5005Δ*gacL* does not fail to add GlcNAc side-chains to polyrhamnose due to a lack of

GlcNA-P-Und and supports the conclusion that GacL may be the GlcNAc-P-

Und:polyRhamnan GlcNAc transferase.

To confirm the role of GacI in GlcNAc-P-Und biosynthesis we expressed GAS GacI

in *E. coli*. When membrane fractions from *E. coli* carrying an empty vector were

incubated with UDP-[^{3}H]GlcNAc and analyzed by silica gel TLC, a small peak of
[^{3}H]GlcNAc-P-P-Und was found (Table 2, Fig. S4A). In contrast, the membranes of
recombinant *E. coli* expressing GacI accumulated two products corresponding to a
small amount of GlcNAc-P-P-Und and a very large amount of GlcNAc-P-Und (Table 2,
Fig. S4B). Thus, our data strongly indicate that GacI is the GlcNAc-P-Und synthase.

GacI homologs in GBS and *E. faecalis* function in the biosynthesis of GlcNAc-P-Und. The presence of GlcNAc-P-P-Und and GlcNAc-P-Und have been previously
reported in *B. cereus* and *Bacillus megaterium* (12, 23). Our *in vitro* studies show that
the MGAS5005 strain synthesizes two [^{3}H]GlcNAc-lipids that exactly co-migrate on TLC
with the previously reported [^{3}H]GlcNAc-lipids from *B. cereus* (Fig. 3A and 3B).
Moreover, an analysis of bacterial genomes using GacI in a BLAST search identified a
GacI homolog (60% identity) in *B. cereus* and *B. megaterium*, suggesting its role in the
GlcNAc-P-Und biosynthesis in these bacteria. Furthermore, *gacI* homologs were
identified in the polysaccharide biosynthesis gene clusters of the important human
pathogens *E. faecalis* (*epaI* with 48% sequence identity) (24) and *S. agalactiae* (Group
B Streptococcus or GBS) (SAN_1536 gene with 46% sequence identity) (Fig. S2). To
investigate whether the GBS homolog of GacI also catalyzes the formation of GlcNAc-P-Und, we engineered a knock out of the GacI homolog in GBS COH1. Although GBS
COH1 membranes synthesize both [^{3}H]GlcNAc-P-P-Und and [^{3}H]GlcNAc-P-Und (Fig.
4B and Fig. S3A), GBS COH1Δ*gacI* membranes no longer synthesize [^{3}H]GlcNAc-P-Und (Table 2 and Fig. S3B).

To confirm that the GacI homolog detected in *E. faecalis*, EpaI (24), also possesses
GlcNAc-P-Und synthase activity, EpaI was expressed exogenously in *E. coli* and found
to actively catalyze the formation of [^{3}H]GlcNAc-P-Und (Table 2, Fig. S4C). Altogether,
our results are consistent with the function of GacI homologs from GAS, GBS and *E.
faecalis* in the transfer of GlcNAc from UDP-GlcNAc to Und-P forming GlcNAc-P-Und.

ATP stimulates GlcNAc-P-Und biosynthesis. Preliminary enzymatic properties
for the synthesis of [^{3}H]GlcNAc-P-Und established that Mg^{2+} was the preferred divalent
cation and that the formation of GlcNAc-P-Und was substantially stimulated by
exogenously added Und-P (as a dispersion in CHAPS detergent). In early studies of
GAC biosynthesis in GAS (25), it was shown that the addition of ATP dramatically

stimulated incorporation of radioactive UDP-[^{3}H]GlcNAc into [^{3}H]GlcNAc membrane

lipids and [^{3}H]GlcNAc-polysaccharide. We confirmed that inclusion of 1 mM ATP

significantly stimulated the incorporation of [^{3}H]GlcNAc into [^{3}H]GlcNAc-P-Und (Fig. 5

and Fig. S5), as well as [^{3}H]GlcNAc-P-P-Und (Fig. S5). However, the inclusion of ATP

did not stimulate GacI activity in *in vitro* assays of CHAPS-soluble, affinity purified GacI

solely dependent on exogenously added Und-P as acceptor. These data suggest that

the effect of ATP addition is most likely due to formation of Und-P, *in situ,* by

phosphorylation of endogenous undecaprenol by undecaprenol kinase (26).

GacJ forms a complex with GacI and enhances its catalytic efficiency. *GacJ* is

located immediately downstream of *gacI* in the GAS GAC biosynthesis gene cluster and

encodes a small 113 aa membrane protein (Fig. 1A). We hypothesized that GacI might

form an obligate complex with GacJ. This hypothesis is based on the observation that

gacJ and *gacI* are frequently located adjacent to each other on bacterial chromosomes

and are sometimes fused to form a single polypeptide (accession numbers: GAM11018,

ADH85075, ALC15489 and ADU67183). To test this hypothesis, we solubilized

membranes from *E. coli* co-expressing GacJ and amino-terminal His-tagged GacI with

the zwitterionic detergent CHAPS, and isolated GacI complexes using Ni-NTA

chromatography (Fig. 6A). SDS-PAGE of the affinity-purified sample revealed two

bands corresponding to the anticipated molecular sizes of GacI and GacJ (Fig. 6B).

Proteomics analysis of the excised protein bands confirmed the identities of the

recovered proteins. This result indicates that GacI and GacJ form a stable, CHAPS-

soluble complex and co-purify during affinity chromatography.

To investigate if GacJ performs any catalytic function in association with GacI, we

tested the proteins for GlcNAc-P-Und synthase activity *in vitro*, individually and in

combination. When GacI was expressed singly in *E. coli*, enzymatically active protein

was found in the membrane fraction (Table 2, Fig. S4B), indicating that GacI does not

require GacJ for activity or membrane association. GacJ was found to be catalytically

inactive when expressed by itself in *E. coli*, consistent with the observation that

5005Δ*gacI* membranes show no residual GlcNAc-P-Und synthase activity. Kinetic

analysis of GlcNAc-P-Und synthase activity for Und-P in membrane fractions of *E. coli*

expressing GacI revealed an apparent Km of 18.7 μM (Table 3, Fig. S6A). However, co-

expression of GacJ dramatically lowered the apparent Km of GacI for Und-P to 1.1 μM,

suggesting a significant change in affinity for the lipid acceptor. Moreover, the V_{max} of

GacI activity increased from 54.2 pmol/min/mg to 18.5 nmol/min/mg in the presence of

GacJ (Table 3, Fig. S6B). A kinetic analysis of GlcNAc-P-Und synthase for Und-P in

MGAS5005 strain gave an apparent Km of 6.4 μM and a V_{max} of 333 pmol/min/mg

(Table 3, Fig. S6C), similar to the enzymatic parameters of GacI co-expressed with

GacJ in *E. coli*. These enzymatic parameters are in marked contrast to those of the

GlcNAc-P transferase that synthesizes GlcNAc-P-P-Und. When a similar kinetic

analysis was performed in 5005Δ*gacI*, so that only GlcNAc-P-P-Und synthesis could be

scored, an apparent Km for Und-P of 19.3 μM was obtained and the V_{max} was only 2.75

pmol/min/mg (Table 3, Fig. S6D). Clearly, GacI displays a much greater catalytic

efficiency for the synthesis of GlcNAc-P-Und than the enzyme synthesizing GlcNAc-P-

P-Und, which is reflected in the dramatic difference in the rates of synthesis of the two

glycolipids. In summary, our data indicate that GacJ forms a stable association with

GacI and stimulates the catalytic activity of GacI.

GlcNAc-P-P-Und is required for initiation of polyrhamnose polysaccharide

biosynthesis. The GAS genome contains a close relative, *gacO*, of *E. coli wecA*, the

tumicamycin-sensitive GlcNAc-phosphate transferase responsible for the synthesis of

GlcNAc-P-P-Und in many bacteria (11, 12). Significantly, rhamnopolysaccharide

synthesis in GAS is inhibited by tunicamycin (5) and a close homolog of GacO, *S.*

mutans RpgP, has been shown to be required for rhamnopolysaccharide synthesis and

can complement WecA deficiency in *E. coli* (13). Our analysis of UDP-[³H]GlcNAc

incorporation in GAS and GBS membrane lipids, revealing the presence of two GlcNAc-

lipids, GlcNAc-P-Und and GlcNAc-P-P-Und, prompted us to investigate the role of

GlcNAc-P-P-Und in the initiation of GAC biosynthesis. First, we investigated the role of

GacO in the synthesis of GlcNAc-P-P-Und. When membrane fractions from the WecA

deficient *E. coli* strain (CLM37) expressing GacO, CLM37:GacO, were incubated with

UDP-[³H]GlcNAc and Und-P (added as a dispersion in 1 % CHAPS), [³H]GlcNAc-P-P-

Und was formed at an enzymatic rate that is similar to that found in the WecA over-

expressor strain, PR4019 (Fig. 7A). CLM37 carrying an empty vector synthesizes no

detectable GlcNAc-P-P-Und under these conditions (Fig. 7A).

To test whether GlcNAc-P-P-Und might function as a membrane anchor for the

synthesis of the polyrhamnose chain of the rhamnopolysaccharide, membrane fractions

from MGAS5005 or the 5005Δ*gacI* mutant were pre-incubated with UDP-[^{3}H]GlcNAc

and ATP, to form [^{3}H]GlcNAc-P-P-Und *in situ*, and chased with non-radioactive dTDP-

rhamnose. The formation of the resultant [^{3}H]GlcNAc-lipids in 5005Δ*gacI* was analyzed

by TLC on silica gel G after 30 min incubation as shown in (Fig. 7B and C). In the

absence of TDP-rhamnose, 5005Δ*gacI* membranes produced only [^{3}H]GlcNAc-P-P-Und

(Fig. 7B), whereas in the presence of TDP-rhamnose two radioactive products,

[^{3}H]GlcNAc-P-P-Und and an additional product with slower mobility on TLC was

observed (Fig 7C). Figure S7 shows the time course of accumulation of the new

GlcNAc-lipid product. Significantly, parallel incubations with MGAS5005 showed that

GlcNAc-P-Und was not glycosylated further. These results strongly support the

possibility that GlcNAc-P-P-Und is an acceptor for rhamnosyl units in GAS and may

function as the lipid anchor for polyrhamnosyl polysaccharide synthesis.

The GacL mutant displays increased sensitivity to peptidoglycan amidases.

The Gram-positive cell wall protects interior structures, plasma membrane and

peptidoglycan, from host defense peptides and hydrolytic and antimicrobial enzymes. It

has been shown that the GacI mutant which is GlcNAc-deficient is more sensitive to LL-

37-induced killing (5). To test the hypothesis that the loss of GlcNAc decorations in GAC

alters cell wall permeability, we investigated the sensitivity of 5005Δ*gacL* to

peptidoglycan amidases: PlyC (27), PlyPy (28) and CbpD (29). When 5005Δ*gacL* cells

were grown in increasing concentrations of CbpD (Fig. 8A), PlyPy (Fig. 8B) or PlyC (Fig.

8C) cellular growth was dramatically inhibited compared to MGAS5005, indicating

increased sensitivity to the presence of amidases.

DISCUSSION

In almost all Gram-positive bacteria, cell wall-attached glycopolymers are critical for cell

envelope integrity and their depletion is lethal (30). Most streptococcal species including

two human pathogens GAS and GBS do not synthesize WTA and instead produce Rha-

containing glycopolymers as functional homologs of WTA (1). GAC is the major cell wall

component of GAS and plays important roles in bacterial physiology and pathogenesis.

The polyrhamnose core of GAC is modified with GlcNAc in an approximately 2:1 ratio

(3, 7) of rhamnose to GlcNAc. Collectively, the results of our study suggest a molecular

mechanism of GAC biosynthesis in which rhamnan polymer is assembled at the

cytoplasmic face of the plasma membrane, translocated to the cell surface and modified

by GlcNAc on the outer side of the membrane as illustrated in Figure 9. We report that a

lipid carrier, GlcNAc-P-P-Und, synthesized by GAS GacO, is a potential acceptor for

initiation of rhamnan backbone biosynthesis. We speculate that the next step of

rhamnan biosynthesis involves the action of the GacB, GacC, GacG and GacF

glycosyltransferases. The lipid-anchored polyrhamnose is then translocated across the

membrane by the ABC transporter encoded by *gacD* and *gacE*. This hypothesis is

supported by studies of rhamnan biosynthesis in *S. mutans* (13, 31, 32).

Biosynthesis of the rhamnan-backbone of GAC is likely essential for GAS viability

because the GacA enzyme involved in dTDP-rhamnose biosynthesis (14), and GacB

and GacC glycosyltranferases are indispensable in GAS (5). In contrast, deletion of

genes required for GlcNAc attachment to the rhamnan backbone does not affect GAS

viability (5). This observation supports our hypothesis that biosynthesis of polyrhamnose

and its translocation to the cell surface occur separately from the pathway involved in

polyrhamnose modification with GlcNAc. In this study, we found that GlcNAc

modification of rhamnan requires GlcNAc-P-Und synthesis. Previously, GlcNAc-P-Und

was isolated from various *Bacilli* membranes (12, 23). However the enzyme required for

GlcNAc-P-Und biosynthesis was not identified and the biological function of this lipid

remained unknown. Our *in vitro* analysis of UDP-GlcNAc incorporation into GlcNAc-

lipids by GAS membranes showed that GacI was required for the biosynthesis of

GlcNAc-P-Und. Moreover CHAPS-soluble, affinity-purified GacI protein catalyzed the

transfer of GlcNAc from UDP-GlcNAc to Und-P yielding GlcNac-P-Und in an *in vitro*

reaction system.

Interestingly, we found that, in contrast to *B. cereus*, GAS and GBS membranes

incubated with UDP-[³H]GlcNAc synthesized primarily [³H]GlcNAc-P-Und. Since GacI

and GacO utilize a common pool of Und-P, this phenomenon is presumably due to a

much higher apparent affinity of GacI for Und-P compared to GacO. Significantly, we

observed a measurable increase in [³H]GlcNAc incorporation into [³H]GlcNAc-P-P-Und

in the GacI deletion mutant. This observation further confirmed that the relative

synthetic rates of GlcNAc-P-Und and GlcNAc-P-P-Und biosynthesis are determined

largely by Und-P availability. Furthermore, we found that the formation of GlcNAc-P-Und

and GlcNAc-P-P-Und in GAS membranes was significantly stimulated by ATP. Since

GacI activity is not stimulated directly by ATP, it is likely that the effect of ATP is due to

increased formation of Und-P, *in situ*, via undecaprenol kinase activity. In *S. mutans,*

membrane-associated undecaprenol kinase catalyzes the ATP-dependent

phosphorylation of undecaprenol to Und-P (26). It is likely that the homolog of this

enzyme encoded by M5005_Spy_0389 is responsible for Und-P biosynthesis in GAS.

Wzx flippases translocate Und-P-P-linked oligosaccharides from the cytoplasmic

side to the periplasmic side of membranes in bacteria (33). We suggest that GacK may

function to transport GlcNAc-P-Und to the extracellular space for utilization as GlcNAc

donor in the GlcNAc modification of polyrhamnose. This hypothesis is based on the

GlcNAc-deficient phenotype of the GacK mutant (5) and the high domain homology

between GacK and proteins of the Wzx family of flippases. However, further research

will be required to confirm this hypothesis. The phenotypes of the GacL mutant:

absence of the GlcNAc side-chains in GAC and accumulation of GlcNAc-P-Und in the

membrane is consistent with GacL functioning in the transfer of GlcNAc from GlcNAc-P-

Und to polyrhamnose yielding GlcNAc-modified polysaccharide. Bioinformatics

searches with the HHpred program against available protein structures identified the

GT-C superfamily of glycosyltrasferases as the closest structural homologs of GacL.

The members of the GT-C superfamily consist of proteins characterized with multiple

transmembrane domains and a large periplasmic loop close to the N-terminus that

contains a metal-binding motif (DxD, ExD, DDx or DEx) (34, 35). The GT-C enzymes

act on the outside of the plasma membrane by transferring sugars from the lipid carrier,

Und-P, onto either glycolipid or protein. GacL contains 12 transmembrane segments

and the highly conserved DEX motif in the first extracellular loop. Thus, the predicted

structural homology of GacL is in agreement with the proposed GacL function in GlcNAc

transfer from GlcNAc-P-Und to polyrhamnose.

Thus, we suggest that the proposed mechanism of GlcNAc attachment to

polyrhamnose in GAS is similar to the mechanism of O-antigen modification in *Shigella*

flexneri in which the GtrB glycosyltransferase, together with GtrA flippase and GtrV

membrane protein, are required for addition of a glycosyl group to rhamnose of the O-

antigen backbone polysaccharide, which is a part of lipopolysaccharide (36). The

modification with glycosyl residues takes place after O-unit assembly and before

transfer of the mature O-polysaccharide to the lipid A-core region of the

lipopolysaccharide. Moreover, in *Mycobacterium tuberculosis* the modification of cell

wall arabinogalactan with galactosamine occurs through similar mechanism involving N-

acetyl galactosaminyl-phosphate-undecaprenol synthase, PpgS, and a membrane

associated GT-C enzyme, Rv3779, that transfers a galactosamine residue from

galactosaminyl-phosphate-undecaprenol to arabinogalactan (37).

The last step of GAC biosynthesis is probably similar to the last step of WTA

biosynthesis: it involves attachment of GAC to certain MurNAc residues in

peptidoglycan via a phosphate ester linkage (38). It is likely catalyzed by members of

LytR-CpsA-Psr (LCP) phosphotransferase family encoded by M5005_Spy_1099 and

M5005_Spy_1474.

GacI is predicted by the HHpred server to have structural homology to

polyisoprenyl-glycosyltransferase GtrB from *Synechocystis* which is a homolog of *S.*

flexneri GtrB (39). GtrB homologs in bacteria and eukaryotes belong to GT-A

superfamily of glycosyltransferases and they are responsible for the addition of UDP-

glucose to the bactoprenol carrier in the cytoplasm yielding the Und-P-glucose

precursor in bacteria and dolichol phosphate-glucose precursor in eukaryotes (34, 35).

Our BLAST search using the GacI sequence as query found homologs of this gene in

many bacterial species, with the broadest diversity observed in the phylum Firmicutes

(Supplemental File). To determine evolutionary relationships between the GacI

homologs, these sequences were analyzed using CLANS (CLuster ANalysis of

Sequences) (40) (Fig. 10). It is noteworthy that the *B. cereus* GacI has two homologs

clustered in two different groups. Our bioinformatics analysis shows that these *gacI*

homologs are located in different gene clusters encoding proteins involved in

polysaccharide biosynthesis and transport. Moreover, the GAS GacI subgrouped with

one *B. cereus* GacI. The GBS GacI and *E. faecalis* GacI homologs are located in the

cluster well-separated from the GAS GacI subgroup. Thus, this observation points to

possible horizontal gene transfer between GAS and *Bacillus*. Analysis of UDP-GlcNAc

incorporation into GlcNAc-lipids in a GBS *gacI* knock-out strain confirmed the function of

this gene in the biosynthesis of GlcNAc-P-Und in GBS. Additionally, we demonstrated

that the *E. faecalis* GacI homolog (EpaI) functions in the biosynthesis of GlcNAc-P-Und.

Significantly, the presence of GacI homologs in GBS and *E. faecalis* matches the

reported occurrence of GlcNAc residues in the cell wall polysaccharides of these

bacteria (41-44).

GacJ, a small membrane protein with three transmembrane α-helices, is required

for GlcNAc side-chain attachment to polyrhamnose (5). GacJ belongs to DUF2304

family of proteins according to the Pfam protein family database (45). In *Geobacter* sp.,

Desulfuromonas sp., *Desulfurivibrio alkaliphilus, Desulfuromonas soudanensis* and

Desulfurispirillum indicum, DUF2304 homologous domains are fused with *gacI*

homologs (Fig. 10 and supplemental file). In *M. tuberculosis* a GacJ homolog, Rv3632,

is co-transcribed with the gene encoding N-acetyl galactosaminyl-phosphate-

undecaprenol synthase, PpgS, and is found to stimulate PpgS activity (37). Consistent

with this finding we showed that GacI and GacJ form a complex and co-expression of

GacI with GacJ significantly enhanced GacI catalytic activity. Further work is under way

to characterize the mechanism of GacJ action on GacI activity.

The identification of the mechanism of GlcNAc attachment to polyrhamnose raises

the question of how this modification functions biologically in bacteria. It has been

previously found that the Δ*gacI* mutant is hypersusceptible to human antimicrobial

peptide LL-37 and the antimicrobial action of factors released by thrombin-activated

platelets, suggesting a role of GlcNAc modification in protecting the plasma membrane

from antimicrobial agents (5). Our study identified the importance of GlcNAc for

protection of GAS peptidoglycan from amidase-induced lysis. In *E. faecalis* the GacI

homolog, EpaI, is involved in biosynthesis of a cell wall-attached polysaccharide (24),

however the polysaccharide structure of the mutant has not been elucidated. The Δ*epaI*

mutant was defective in conjugative transfer of a plasmid and resistance of bacteria to

detergent and bile salts (24).

In conclusion, our study provides a platform for elucidation of novel pathways of

glycopolymer biosynthesis in other bacterial pathogens including important drug-

resistant bacteria such as *E. faecalis* and *Clostridium sordellii* that possess GacI and

GacJ homologs. Since the enzymes involved in the biosynthesis of cell wall attached

glycopolymers are promising targets for novel antimicrobials and the glycopolymers

represent important features for diagnostics and vaccine targets, our data may provide

opportunities for developing novel therapeutics against antibiotic-resistant bacterial

pathogens.

MATERIALS AND METHODS

Bacterial strains and growth conditions. All plasmids, strains and primers used in

this study are listed in Tables S1 and S2 in the supplemental material. The strains used

in this study were GAS M1-serotype strain MGAS5005 (16), *Bacillus cereus* (ATCC

14579), *Streptococcus agalactiae* COH1 (Group B *Streptococcus* or GBS), *E. coli*

CLM37 (46), *E. coli* PR4019 (12), *E. coli* DH5α and *E. coli* Rosetta (DE3). GAS and

GBS cultures were grown in Todd-Hewitt broth (BD) supplemented with 0.2% yeast

extract (THY), or on THY agar plates at 37 °C. *E. coli* and *B. cereus* strains were grown

in Luria-Bertani (LB) medium or on LB agar plates at 37 °C. When required, antibiotics

were included at the following concentrations: ampicillin at 100 μg ml^{-1} for *E. coli*;

streptomycin at 100 μg ml^{-1} for *E. coli*; erythromycin at 500 μg ml^{-1} for *E. coli* and 1 μg

ml^{-1} for GAS and GBS; chloramphenicol at 10 μg ml^{-1} for *E. coli* and 5 μg ml^{-1} for GAS;

spectinomycin at 200 μg ml^{-1} for *E. coli* and 100 μg ml^{-1} for GAS and GBS.

DNA techniques. Plasmid DNA was isolated from *E. coli* by commercial kits

(Qiagen) according to the manufacturer's instructions and used to transform *E. coli*,

GAS and GBS strains. Plasmids were transformed into GAS and GBS by

electroporation as described previously (47). Chromosomal DNA was purified from GAS

and GBS as described in (48). To construct single-base substitutions or deletion

mutations, we used the QuikChange® II XL Site-Directed Mutagenesis Kit (Stratagene)

according to the manufacturer's protocol. Constructs containing mutations were

identified by sequence analysis. Primers for site-directed mutagenesis are listed in

Table S3. All constructs were confirmed by sequencing analysis (Eurofins MWG

Operon).

Construction of the *gacI* deletion mutant in GAS. For construction of strain

5005Δ*gacI*, MGAS5005 chromosomal DNA was used as a template for amplification of

two DNA fragments using two primers pairs: GacIm-BamHI-f/GacIdel-r and GacIdel-

f/GacIm-XhoI-r (Table S2). Primer GacIdel-f is complementary to primer GacIdel-r. The

two gel-purified PCR products containing complementary ends were mixed and

amplified using a PCR overlap method (49) with primer pair GacIm-BamHI-f/GacIm-

XhoI-r to create the deletion of *gacI*. The PCR product was digested with BamHI and

XhoI and ligated into BamHI/SalI-digested temperature-sensitive shuttle vector

pJRS233 (50). The plasmid was designated pJRS233Δ*gacI* (Table S1). The resulting

plasmid was transformed into MGAS5005, and erythromycin resistant colonies were

selected on THY agar plates at 30 °C. Integration was performed by growth of

transformants at 37 °C with erythromycin selection. Excision of the integrated plasmid

was performed by serial passages in THY media at 30 °C and parallel screening for

erythromycin-sensitive colonies. Mutants were verified by PCR sequencing of the loci.

Construction of the *gacL* deletion mutant in GAS. To create BglII and XhoI

cloning sites in the polylinker region of pUC19, site-directed mutagenesis of the

plasmid, using the primer pair listed in Table S3 was carried out. The plasmid was

designated pUC19BX. The nonpolar *aadA* spectinomycin resistance cassette was

amplified from pLR16T (Table S2) using primers Spec-SalI-f/Spec-BamH-r (Table S2),

digested with SalI/BamHI, and ligated into SalI/BamHI-digested pUC19BX to yield

pUC19BXspec. MGAS5005 chromosomal DNA was used as a template for

amplification of two DNA fragments using two primers pairs: gacLup-BglII-f/gacLup-SalI-

r and gacLdown-BamHI-f/gacLdown-XhoI-r (Table S2). The first PCR product was

digested with BglII/SalI and ligated into BglII/SalI-digested pUC19BXspec. The resultant

plasmid, pUC19BXspecL1, was digested with BamHI/XhoI and used for ligation with the

second PCR product that was digested with BamHI/XhoI. The resultant plasmid,

pUC19BXspecL2, was digested with BglII and XhoI to obtain a DNA fragment

containing *aadA* flanked with the *gacL* upstream and downstream regions. The DNA

fragment was ligated into pHY304 vector (51) and digested with BamHI/XhoI to yield

pHY304Δ*gacL*. The resulting plasmid was transformed into MGAS5005, and

erythromycin resistant colonies were selected on THY agar plates at 30 °C. The

mutants were isolated as describe above. 5005Δ*gacL* mutants were screened for

sensitivity to spectinomycin and verified by PCR sequencing of the loci.

Complementation of the 5005Δ*gacL* mutant with *gacL* (5005Δ*gacL gacL*⁺).

To construct the plasmid for complementation of the 5005Δ*gacL* mutant, MGAS5005

chromosomal DNA was used as a template for amplification of a wild-type copy of *gacL*

using the primer pair GacL-XhoI-f and GacL-BglII-r (Table S2). The PCR products were

digested with XhoI and BglII and cloned in pBBL740 (Table S1) previously digested with

the respective enzymes. The integrational plasmid pBBL740 does not have a replication

origin that is functional in GAS, so the plasmid can be maintained only by integrating

into the GAS chromosome through homologous recombination. The resultant plasmid

pGacL was transformed into 5005Δ*gacL* by electroporation and transformants were

selected on agar plates containing chloramphenicol. Several chloramphenicol resistant

colonies were selected and *gacL* integration into the chromosome was confirmed by

sequencing a PCR fragment.

Construction of the *gacI* (SAN_1536) deletion mutant in GBS. GBS COH1

chromosomal DNA was used as a template for amplification of two DNA fragments

using two primers pairs: gbsIup-BglII-f/gbsIup-SalI-r and gbsId-BamHI-f/gbsId-XhoI-r

(Table S2). The plasmid for *gacI* (SAN_1536) knock-out was constructed using the

same strategy described for *gacL* deletion (see above). The resultant plasmid,

pHY304GBSΔI, was transformed into GBS COH1, and erythromycin resistant colonies

were selected on THY agar plates at 30 °C. The mutants were isolated as described

above. GBS COH1 Δ*gacI* mutants were screened for sensitivity to spectinomycin and

verified by PCR sequencing of the loci.

Construction of the plasmids for *E. coli* expression of GacI, GacJ and GacO.

To create a vector for expression of GacI from GAS, the gene was amplified from

MGAS5005 chromosomal DNA using the primer pair GacI-NcoI-f and GacI-XhoI-r

(Table S2). The PCR product was digested with NcoI and XhoI, and ligated into

NcoII/XhoI-digested pRSF-NT vector (Table S1). The resultant plasmid, pGacI,

contained *gacI* fused at the N-terminus with a His-tag followed by a TEV protease

recognition site. To create a vector for expression of GacI and GacJ, the bicistronic

DNA fragment was amplified from MGAS5005 chromosomal DNA using the primer pair

GacI-NcoI-f and GacJ-XhoI-r (Table S2). The vector was constructed as described

above. The plasmid was designated pGacIJ. pET21_NESG plasmid for expression of

the GacI homolog, EpaI, from *Enterococcus faecalis* was obtained from the DNASU

repository (52). The construct was confirmed by sequencing analysis. The plasmids

were transferred into competent *E. coli* Rosetta (DE3) (Novagen) using the

manufacturer's protocol.

To create a vector for expression of GacO, the gene was amplified from MGAS5005

chromosomal DNA using the primer pair GacO-Xba-f and GacO-HindIII-r (Table S2).

The PCR product was digested with XbaI and HindIII, and ligated into XbaI/HindIII-

digested pBAD33 vector. The resultant plasmid, pGacO, was transferred into competent

E. coli CLM37 strain that has a deletion of the *wecA* gene.

Construction of the plasmids for *E. coli* expression of PlyPy, CbpD and PlyC.

To create a vector for expression of CbpD amidase (29), the gene was amplified from

MGAS5005 chromosomal DNA using the primer pair 28-NcoI-f and 28-stop-r (Table

S2). The PCR product was digested with NcoI and XhoI, and ligated into NcoI/XhoI-

digested pRSF-NT vector. The resultant plasmid, pCbpD, contained *cbpD* fused at the

N-terminus with a His-tag followed by a TEV protease recognition site. To create a

vector for expression of PlyPy amidase (28), the gene was amplified from MGAS5005

chromosomal DNA using the primer pair PlyPy-NcoI-f and PlyPy-XhoI-r. The PCR

product was digested with NcoI and XhoI, and ligated into NcoI/XhoI-digested pET-21d

vector. The resultant plasmid, pPlyPy contained the gene fused at the C terminus with a

His-tag.

To create a vector for expression of PlyC amidase (53), a DNA fragment spanning

bicistronic operon which encodes *plyCA* and *plyCB* (27) was synthesized by

ThermoFisher Scientific. The plasmid was digested with NcoI and XhoI, and ligated into

NcoI/XhoI-digested pET21d vector. The resultant plasmid, pPlyC contained *plyCA*

followed by *plyCB* fused at the C terminus with a His-tag.

Expression and purification of GacI, EpaI and GacI/GacJ complex. For

expression of GacI, EpaI and GacI/GacJ complex, *E. coli* Rosetta (DE3) cells carrying

the respective plasmid were grown to an OD_{600} of 0.4-0.6 and induced with 1 mM IPTG

at 18 °C for approximately 16 hours. The cells were lysed in 20 mM Tris-HCl pH 7.5,

300 mM NaCl with two passes through a EmulsiFlex-C5 microfluidizer cell disrupter

(Avestin, Inc., Ottawa Ontario,CA). The lysate was centrifuged at 7000 x g for 30

minutes, 4°C. The supernatant was centrifuged at 30,000 x g for 60 minutes, 4°C to isolate the membrane fraction.

To isolate GacI/GacJ complexes the membrane proteins were solubilized in 2.5% CHAPS, 20 mM Tris-HCl pH 7.5, 300 mM NaCl for 60 minutes, rotating at room temperature. Insoluble material was removed by centrifugation at 30,000 x g for 60 minutes, 4°C. Solubilized GacI/GacJ were purified by Ni-NTA chromatography with washes of 2.5% CHAPS, 20 mM Tris-HCl pH 7.5, 300 mM NaCl and 2.5% CHAPS, 20 mM Tris-HCl pH7.5, 300 mM NaCl, 10 mM imidazole, and elution with 2.5% CHAPS, 20 mM Tris-HCl pH 7.5, 300 mM NaCl, 250 mM imidazole. The GacI/GacJ complex was further purified by size exclusion chromatography on a Superdex 200 (GE Biosciences) column in 2.5% CHAPS, 20 mM HEPES pH 7.5, 100 mM NaCl, with monitoring for protein elution at 280 nm.

Expression of GacO. For expression of GacO, *E. coli* CLM37 cells carrying the pertinent plasmid were grown to an OD_{600} of 0.8 and induced with 13 mM L-arabinose at 25 °C for approximately 3 hours. The cells were lysed in 20 mM Tris-HCl pH 7.5, 300 mM NaCl with two passes through a microfluidizer cell disrupter. The lysate was centrifuged at 1000 x g for 15 minutes, 4 °C. The supernatant was centrifuged at 40,000 x g for 60 minutes, 4 °C to isolate the membrane fraction.

Expression and purification of PlyPy, CbpD and PlyC. For expression and purification of PlyPy, CbpD and PlyC, *E. coli* Rosetta (DE3) cells carrying the respective plasmids were grown to an OD_{600} of 0.4-0.6 and induced at 18 °C with 1 mM IPTG for approximately 16 hours. The cells were lysed in 20 mM Tris-HCl pH 7.5, 300 mM NaCl with two passes through a microfluidizer cell disrupter. The soluble fraction was purified by Ni-NTA chromatography with washes of 20 mM Tris-HCl pH 7.5, 300 mM NaCl and 20 mM Tris-HCl pH7.5, 300 mM NaCl, 10mM imidazole, and elution with 20 mM Tris-HCl pH 7.5, 300 mM NaCl, 250 mM imidazole. The PlyPy and CbpD eluate was further purified by size exclusion chromatography on a Superdex 200 column in 20 mM HEPES pH 7.5, 100 mM NaCl for PlyPy, or 20 mM MOPS pH 6.5, 100 mM NaCl for CbpD. PlyC was further purified by anion exchange chromatography on a MonoQ 5/50 GL (GE Biosciences) column in 10 mM sodium phosphate pH 6.0, and a 20 column volume elution gradient of 0-500 mM NaCl.

Isolation of GAS and GBS membranes. Bacteria were grown at 37 °C to an OD_{600} of 0.8. To obtain GAS cell membranes, cell pellet was re-suspended in phosphate-buffered saline (PBS) and incubated 1 h with PlyC lysin as described in (54). To obtain GBS membranes, the cell pellet was re-suspended in acetate buffer (50 mM CH_3COONa, 10 mM $CaCl$, 50 mM NaCl, pH 5) and incubated with mutanolysin (200 U/ml, Sigma–Aldrich) for 2 h at 37 °C. After hydrolytic enzyme treatment the bacterial suspension was sonicated using a Fisher Scientific™ Model 505 Sonic Dismembrator, 15 times ×15 s. After centrifugation at 8,000 g for 10 min at 4 °C, the supernatant was collected and centrifuged at 40,000 g for 60 min. The pellet was collected as the membrane fraction.

Mass-spectrometry analysis of GacI and GacJ. LC-MS/MS for proteomic analysis was performed using an LTQ-Orbitrap mass spectrometer (Thermo Fisher Scientific) coupled with an Eksigent Nanoflex cHiPLC system (Eksigent) through a nanoelectrospray ionization source. The LC-MS/MS data were subjected to database searches for protein identification using Proteome Discoverer software V. 1.3 (Thermo Fisher Scientific) with a local MASCOT search engine.

Dot-blot analysis. Bacterial cells (1 ml) from exponential phase cultures (OD_{600}=0.8) was centrifuged, washed with PBS, resuspended in 100 μl of PBS and incubated with 1 μl PlyC (1.5 mg/ml) for 1 h at 37°C. After centrifugation at 16,000 g for 2 min, 5 μl of the supernatant was spotted on a nitrocellulose membrane. The membrane was blocked 1 h with 7% skim milk in PBS with 0.1% Tween 20, and incubated overnight with an anti-GAC antibody diluted 1:5,000 (ab9191; Abcam). Bound antibodies were detected with a peroxidase-conjugated goat anti-rabbit immunoglobulin G antibody, and the Amersham ECL (enhanced chemiluminescence) Western blotting system.

Binding of succinylated wheat germ agglutinin (sWGA) to bacteria. MGAS5005 and 5005Δ*gacL* cells were collected during mid-exponential phase (OD_{600}=0.6), washed three times with BSA-saline solution (0.5% BSA, 0.15 M NaCl), resuspended in BSA-saline solution and incubated for 30 minute at 37 °C with mixing. GlcNAc specific fluorescein-sWGA was added to final concentrations of 0, 12.5, 25, 50, 100 μg/ml. After 1 h of incubation at 37 °C with mixing, the cells were centrifuged, washed twice, and

resuspended in BSA-saline. Bound fluorescein-sWGA was quantified in a fluorimeter

SpectraMax M5 (Molecular Devices) using an excitation of 544 nm and emission of 590

nm.

Isolation of cell wall from GAS. MGAS5005 and 5005Δ*gacL* cell wall was

prepared from exponential phase cultures (OD_{600}=0.8) by the SDS-boiling procedure as

described for *Streptococcus pneumoniae* (55). Purified cell wall samples were

lyophilized and used for composition analysis.

Carbohydrate composition analysis. Carbohydrate composition analysis was

performed at the Complex Carbohydrate Research Center (Athens, GA) by combined

gas chromatography/mass spectrometry (GC/MS) of the per-O-trimethylsilyl (TMS)

derivatives of the monosaccharide methyl glycosides produced from the sample by

acidic methanolysis as described previously by (56).

Assay of lytic activity of PlyPy, CbpD and PlyC. Aliquots of MGAS5005 and

5005Δ*gacL* frozen stocks [prepared as described in (57)] were inoculated into THY

medium 1:20. To a 96 well plate, 200 µL aliquots of culture were dispensed in duplicate,

per condition. The plates were grown at 37 °C without aeration for 2 hours. An assay

plate was set up containing PlyPy (8 µg mL^{-1}, 16 µg mL^{-1}, 32 µg mL^{-1}), CbpD (0.31 µg

mL^{-1}, 0.63 µg mL^{-1}, 1.26 µg mL^{-1}) or PlyC (1.6 ng mL^{-1}, 3.1 ng mL^{-1}, 6.2 ng mL^{-1}) in a

volume of 100 µL. To the assay plate, 100 µL of culture was added to each well giving

final concentrations of: PlyPy (4 µg mL^{-1}, 8 µg mL^{-1}, 16 µg mL^{-1}), CbpD (0.16 µg mL^{-1},

0.31 µg mL^{-1}, 0.63 µg mL^{-1}), or PlyC (0.8 ng mL^{-1}, 1.6 ng mL^{-1}, 3.1 ng mL^{-1}). After

mixing, the absorbance at 600 nm was measured for each well (t_0) and the plate was

incubated at 37 °C for 1 h. The absorbance at 600 nm was then measured again (t_1).

The percentage difference (t_1-t_0) in growth relative to 0 mM of enzyme/antimicrobial, for

each concentration was calculated (i.e. 0 mM PlyPy = 100%).

Extraction and characterization of bacterial glycolipids. Two liters of bacterial

cells from exponential phase cultures (OD_{600}=0.8) were recovered by sedimentation at

10,000 x g for 30 min, and washed with ice-cold PBS 2x. Cells were resuspended in 50

ml PBS, sensitized by incubation with 10 ng mL^{-1} PlyC for 1 h at 37 °C and stirred

vigorously with two volumes of CH_3OH and one volume of $CHCl_3$ for 30 min at room

temperature. The mixture was divided into 5 mL aliquots in 12 mL glass centrifuge

tubes. Insoluble material was removed by centrifugation at 200 g and the organic
extract was transferred to a separatory funnel. The insoluble residue was further
extracted with 1 mL of $CHCl_3/CH_3OH$ (2:1) per tube, two times, and the extracts were
combined with the previous organic phase. The organic extract was supplemented with
$CHCl_3$ and 0.9% NaCl to give a final composition of $CHCl_3/CH_3OH/0.9\%$ NaCl (3:2:1),
mixed vigorously, and allowed to stand in the cold until phase separation was achieved.
The organic phase was drained off into a second separatory funnel and the organic
layer was washed with 1/3 volume of $CHCl_3/CH_3OH/0.9\%$ saline (3:48:47), two times.
The aqueous layers were discarded. The organic extract was dried on a vacuum rotary
evaporator, dissolved in a small volume of $CHCl_3/CH_3OH$ (2:1) and transferred to a 12.5
x 100 mm screw cap glass tube (with Teflon lined cap). The organic extract was dried
under a stream of nitrogen gas and the glycerolipids were destroyed by deacylation in
0.1 M KOH in toluene/CH_3OH (1:3) at 0 °C, 60 min. Following deacylation, the reactions
were neutralized with acetic acid, diluted with two volumes of $CHCl_3$, 1 volume of
$CHCl_3/CH_3OH$ (2:1) and 1/5 volume of 0.9% NaCl/10 mM EDTA. The two-phase mixture
was mixed vigorously and centrifuged to separate the phases. The organic phase was
dried under nitrogen, spotted on a 20 x 20 cm sheet of Silica Gel G and developed in
$CHCl_3/CH_3OH/H_2O/NH_4OH$ (65:25:4:1). Bacterial lipids were visualized by staining with
iodine vapors and pertinent spots were scraped from the thin layer plate, eluted from the
silica gel with $CHCl_3/CH_3OH$ (2:1) and reserved for further analysis.

Mass spectrometry analysis of a phospholipid isolated from 5005Δ*gacL*. A
phospholipid accumulated by 5005Δ*gacL* was purified as described above using
preparative TLC in silica gel. The compound was analyzed by LC-MS using a Q-
exactive mass spectrometer and an Ultimate 3000 ultra high performance liquid
chromatography system (Thermo Fisher Scientific, San Jose, CA) on a Kinetex C18
reversed-phase column (2.6 mm × 100 mm, 2.1 µm, Phenomenex, USA). Two solvents
were used for gradient elution: (A) acetonitrile/water (2:3, v/v), (B)
isopropanol/acetonitrile (9:1, v/v). Both A and B contained 10 mM ammonium formate
and 0.1% formic acid. The column temperature was maintained at 40 °C, and the flow
rate was set to 0.25 ml/min. Mass spectrometric detection was performed by
electrospray ionization in negative ionization mode with source voltage maintained at

4.0 kV. The capillary temperature, sheath gas flow and auxiliary gas flow were set at

330 °C, 35 arb and 12 arb, respectively. Full-scan MS spectra (mass range m/z 400 to

1500) were acquired with resolution R = 70,000 and AGC target 5e5. MS/MS

fragmentation was performed using high-energy C-trap dissociation with resolution R =

35,000 and AGC target 1e6. The normalized collision energy was set at 30.

Assay for incorporation of GlcNAc into polyisoprenol-linked glycolipid

intermediates. Reaction mixtures for measuring the incorporation of [^{3}H]GlcNAc into

lipids contained 50 mM Tris, pH 7.4, 0.25 M sucrose, 20 mM $MgCl_2$, 0.5 mM β-

mercaptoethanol, 5 μM UDP-[6-^{3}H]GlcNAc [(100-2000 cpm/pmol) American

Radiolabelled Chemicals] and bacterial membrane suspension (50-250 μg bacterial

membrane protein) in a total volume of 10 to 100 μL. In some experiments, 1 mM ATP

was included, as indicated. Following incubation at 30 °C, the enzymatic reactions were

terminated by the addition of 40 volumes of $CHCl_3$/CH_3OH (2:1), thoroughly mixed and

incubated for 5 min at room temperature. Insoluble material was sedimented at 200 *g*

and the organic extract was transferred to a 12x100 mm glass tube. The residue was

re-extracted with 1 ml $CHCl_3$/CH_3OH (2:1), two times, and the organic extracts were

combined. The pooled organic extracts were freed of unincorporated radioactivity by

sequential partitioning with 1/5 volume of 0.9% saline/10 mM EDTA and then with 1/3

volume of $CHCl_3$/CH_3OH/0.9% saline (3:48:47) 3 times, discarding the aqueous phase

each time. The washed organic phases were dried under nitrogen and re-dissolved in

$CHCl_3$/CH_3OH (2:1). A carefully measured aliquot was removed and analyzed for

radioactivity by liquid scintillation spectrometry after drying. The remainder of the

sample was analyzed by thin layer chromatography on Silica Gel G, developed in

$CHCl_3$/CH_3OH/H_2O/NH_4OH (65:25:4:1), using a BioScan AR2000

radiochromatoscanner. Incorporation of [^{3}H]GlcNAc into individual [^{3}H]GlcNAc-lipids

was calculated using the peak integration values obtained from the BioScan.

Degradation of GlcNAc-lipids by mild acid and mild alkaline treatment.

Bacterial lipids were subjected to mild acid hydrolysis (50% isopropanol, 0.1 M HCl, 50

°C, 1 h) and mild alkaline methanolysis (0.1 M KOH in toluene/CH_3OH (1:3), 0 °C, 1 h).

Following incubation, the reactions were neutralized with either 1 M Tris or concentrated

acetic acid, diluted with $CHCl_3$, CH_3OH and 0.9% NaCl to give a final composition of

3:2:1, respectively and partitioned as described above. The organic phases were dried

under nitrogen and either quantified for radioactivity or analyzed by thin layer

chromatography and detected by iodine staining.

Phenolysis of GlcNAc-P-Und. Purified [^{3}H]GlcNAc-P-Und was dried under

nitrogen in a 12x100 mm conical screw-cap tube and heated to 68 °C in 0.2 ml 50%

aqueous phenol as described by Murazumi et al. (22). Following phenolysis, 0.1 ml

water was added and the samples were thoroughly mixed. The aqueous and phenolic

layers were separated, dried and analyzed for radioactivity by scintillation spectrometry.

Analytical methods. Membrane protein concentrations were determined using the

BCA protein assay kit (Pierce Chemical Co.) employing the method of Ruiz et al. (33).

Radioactivity was quantified by liquid scintillation spectrometry on a Packard Tri-Carb

Liquid Scintillation Spectrometer using Econosafe Complete Counting Cocktail

(Research Products International, Inc., Elk Grove IL).

Phylogenetic analysis of GacI homologs. Sequences of GacI homologs were

retrieved from the non-redundant database using BLAST (58). One thousand

sequences were downloaded and were manually curated to remove sequences with

shorter that 90% sequence length overlap. The redundancy was reduced using CD-HIT

with 0.98 cut off (59). Sequences were manually curated to correct misannotated

translation start codons. The pair-wise similarities were analyzed and visualized using

CLANS with an E-value cut off 1 e^{-80} (40, 60, 61).

Bioinformatics tools. The TOPCONS (http://topcons.net/) (62) web server was

employed to predict trans-membrane regions of GacI, GacJ and GacL. Homology

detection and structure prediction were performed by the HHpred server

(https://toolkit.tuebingen.mpg.de/#/tools/hhpred) (63).

Statistical Analysis. Unless otherwise indicated, statistical analysis was carried out

from at least three independent experiments. Quantitative data was analyzed using the

paired Student's t-test. A P value equal to or less that 0.05 is considered statistically

significant.

Supplemental Material

Additional Supplemental Material may be found in the online version of this article at the

publisher's web-site.

Table S1

Table S2

Table S3

Table S4

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

Figure S7

Supplemental File 1

ACKNOWLEDGEMENTS

J.S.R., R.J.E., H.Z., A.J.M., N.M.vS., K.V.K. and N.K. designed the experiments. J.S.R.,

R.J.E. and N.K. performed biochemical experiments. P.D., J.C., H.Z. and A.J.M.

performed MS analysis. N.K. constructed plasmids and isolated mutants. J.S.R., R.J.E.,

K.V.K. and N.K. wrote the manuscript.

Funding Information

This work was supported by NIH grants R21AI113253 from the National Institute of

Allergy and Infectious Diseases (to N.K.) and R01GM102129 (to Dr. Charles J.

Waechter and J.S.R), 1S10OD021753 to A.J.M. and by the Center of Biomedical

Research Excellence (COBRE) Pilot Grant (to K.V.K, N.K. and J.S.R.) supported by

NIH grant P30GM110787 from the National Institute of General Medical Sciences.

N.M.vS. is supported by VIDI grant 91713303 from the Netherlands Organization for

Scientific Research (NWO).

Carbohydrate composition analysis at the Complex Carbohydrate Research Center

was supported by the Chemical Sciences, Geosciences and Biosciences Division,

Office of Basic Energy Sciences, U.S. Department of Energy grant (DE-FG02-

93ER20097) to Parastoo Azadi.

The funders had no role in study design, data collection and interpretation, or the

decision to submit the work for publication

REFERENCES

1. **Mistou MY, Sutcliffe IC, van Sorge NM.** 2016. Bacterial glycobiology: rhamnose-containing cell wall polysaccharides in Gram-positive bacteria. FEMS Microbiol Rev **40:**464-479.

2. **Carapetis JR, Steer AC, Mulholland EK, Weber M.** 2005. The global burden of group A streptococcal diseases. Lancet Infect Dis **5:**685-694.

3. **McCarty M.** 1952. The lysis of group A hemolytic streptococci by extracellular enzymes of Streptomyces albus. II. Nature of the cellular substrate attacked by the lytic enzymes. J Exp Med **96:**569-580.

4. **Lancefield RC.** 1933. A Serological Differentiation of Human and Other Groups of Hemolytic Streptococci. J Exp Med **57:**571-595.

5. **van Sorge NM, Cole JN, Kuipers K, Henningham A, Aziz RK, Kasirer-Friede A, Lin L, Berends ET, Davies MR, Dougan G, Zhang F, Dahesh S, Shaw L, Gin J, Cunningham M, Merriman JA, Hutter J, Lepenies B, Rooijakkers SH, Malley R, Walker MJ, Shattil SJ, Schlievert PM, Choudhury B, Nizet V.** 2014. The classical lancefield antigen of group a Streptococcus is a virulence determinant with implications for vaccine design. Cell host & microbe **15:**729-740.

6. **Coligan JE, Kindt TJ, Krause RM.** 1978. Structure of the streptococcal groups A, A-variant and C carbohydrates. Immunochemistry **15:**755-760.

7. **McCarty M.** 1956. Variation in the group-specific carbohydrate of group A streptococci. II. Studies on the chemical basis for serological specificity of the carbohydrates. J Exp Med **104:**629-643.

8.   **Kabanova A, Margarit I, Berti F, Romano MR, Grandi G, Bensi G, Chiarot E,**

Proietti D, Swennen E, Cappelletti E, Fontani P, Casini D, Adamo R, Pinto V,

Skibinski D, Capo S, Buffi G, Gallotta M, Christ WJ, Campbell AS, Pena J,

Seeberger PH, Rappuoli R, Costantino P. 2010. Evaluation of a Group A

Streptococcus synthetic oligosaccharide as vaccine candidate. Vaccine **29:**104-

114.

9.   **Sabharwal H, Michon F, Nelson D, Dong W, Fuchs K, Manjarrez RC, Sarkar**

A, Uitz C, Viteri-Jackson A, Suarez RS, Blake M, Zabriskie JB. 2006. Group A

streptococcus (GAS) carbohydrate as an immunogen for protection against GAS

infection. J Infect Dis **193:**129-135.

10.   **Soldo B, Lazarevic V, Karamata D.** 2002. tagO is involved in the synthesis of all

anionic cell-wall polymers in Bacillus subtilis 168. Microbiology **148:**2079-2087.

11.   **Meier-Dieter U, Barr K, Starman R, Hatch L, Rick PD.** 1992. Nucleotide

sequence of the Escherichia coli rfe gene involved in the synthesis of

enterobacterial common antigen. Molecular cloning of the rfe-rff gene cluster. J

Biol Chem **267:**746-753.

12.   **Rush JS, Rick PD, Waechter CJ.** 1997. Polyisoprenyl phosphate specificity of

UDP-GlcNAc:undecaprenyl phosphate N-acetylglucosaminyl 1-P transferase

from E.coli. Glycobiology **7:**315-322.

13.   **Yamashita Y, Shibata Y, Nakano Y, Tsuda H, Kido N, Ohta M, Koga T.** 1999.

A novel gene required for rhamnose-glucose polysaccharide synthesis in

Streptococcus mutans. J Bacteriol **181:**6556-6559.

14.   **van der Beek SL, Le Breton Y, Ferenbach AT, Chapman RN, van Aalten DM,**

Navratilova I, Boons GJ, McIver KS, van Sorge NM, Dorfmueller HC. 2015.

GacA is essential for Group A Streptococcus and defines a new class of

monomeric dTDP-4-dehydrorhamnose reductases (RmlD). Mol Microbiol **98:**946-

962.

15.   **Shibata Y, Yamashita Y, Ozaki K, Nakano Y, Koga T.** 2002. Expression and

characterization of streptococcal rgp genes required for rhamnan synthesis in

Escherichia coli. Infect Immun **70:**2891-2898.

16.   **Sumby P, Porcella SF, Madrigal AG, Barbian KD, Virtaneva K, Ricklefs SM,**

Sturdevant DE, Graham MR, Vuopio-Varkila J, Hoe NP, Musser JM. 2005.

Evolutionary origin and emergence of a highly successful clone of serotype M1

group a Streptococcus involved multiple horizontal gene transfer events. J Infect

Dis **192:**771-782.

17.   **Bevenue A, Williams KT.** 1951. Further evidence indicating the specificity of the

orcinol spray reagent for ketoheptoses on paper chromatograms. Arch Biochem

Biophys **34:**225-227.

18.   **Dittmer JC, Lester RL.** 1964. A Simple, Specific Spray for the Detection of

Phospholipids on Thin-Layer Chromatograms. Journal of lipid research **5:**126-

127.

19.   **Danilov LL, Druzhinina TN, Kalinchuk NA, Maltsev SD, Shibaev VN.** 1989.

Polyprenyl phosphates: synthesis and structure-activity relationship for a

biosynthetic system of Salmonella anatum O-specific polysaccharide. Chemistry

and physics of lipids **51:**191-203.

20.   **Wolucka BA, de Hoffmann E, Rush JS, Waechter CJ.** 1996. Determination of

the anomeric configuration of glycosyl esters of nucleoside pyrophosphates and

polyisoprenyl phosphates by fast-atom bombardment tandem mass

spectrometry. Journal of the American Society for Mass Spectrometry **7:**541-549.

21.   **Wolucka BA, Rush JS, Waechter CJ, Shibaev VN, de Hoffmann E.** 1998. An

electrospray-ionization tandem mass spectrometry method for determination of

the anomeric configuration of glycosyl 1-phosphate derivatives. Analytical

biochemistry **255:**244-251.

22.   **Murazumi N, Yamamori S, Araki Y, Ito E.** 1979. Anomeric configuration of N-

acetylglucosaminyl phosphorylundecaprenols formed in Bacillus cereus

Membranes. J Biol Chem **254:**11791-11793.

23.   **Yamamori S, Murazumi N, Araki Y, Ito E.** 1978. Formation and function of N-

acetyloglucosamine-linked phosphoryl- and pyrophosphorylundecaprenols in

membranes from Bacillus cereus. J Biol Chem **253:**6516-6522.

24.   **Dale JL, Cagnazzo J, Phan CQ, Barnes AM, Dunny GM.** 2015. Multiple roles

for Enterococcus faecalis glycosyltransferases in biofilm-associated antibiotic

resistance, cell envelope integrity, and conjugative transfer. Antimicrob Agents

Chemother **59:**4094-4105.

25.   **Reusch VM, Jr., Panos C.** 1977. Synthesis of "group polysaccharide" by

membranes from Streptococcus pyogenes and its stabilized L-form. J Bacteriol

129:1407-1414.

26.   **Lis M, Kuramitsu HK.** 2003. The stress-responsive dgk gene from

Streptococcus mutans encodes a putative undecaprenol kinase activity. Infect

Immun **71:**1938-1943.

27.   **Nelson D, Schuch R, Chahales P, Zhu S, Fischetti VA.** 2006. PlyC: a

multimeric bacteriophage lysin. Proc Natl Acad Sci U S A **103:**10765-10770.

28.   **Lood R, Raz A, Molina H, Euler CW, Fischetti VA.** 2014. A highly active and

negatively charged Streptococcus pyogenes lysin with a rare D-alanyl-L-alanine

endopeptidase activity protects mice against streptococcal bacteremia.

Antimicrob Agents Chemother **58:**3073-3084.

29.   **Berg KH, Ohnstad HS, Havarstein LS.** 2012. LytF, a novel competence-

regulated murein hydrolase in the genus Streptococcus. J Bacteriol **194:**627-635.

30.   **Chu M, Mallozzi MJ, Roxas BP, Bertolo L, Monteiro MA, Agellon A,**

Viswanathan VK, Vedantam G. 2016. A Clostridium difficile Cell Wall

Glycopolymer Locus Influences Bacterial Shape, Polysaccharide Production and

Virulence. PLoS Pathog **12:**e1005946.

31.   **Tsukioka Y, Yamashita Y, Nakano Y, Oho T, Koga T.** 1997. Identification of a

fourth gene involved in dTDP-rhamnose synthesis in Streptococcus mutans. J

Bacteriol **179:**4411-4414.

32.   **Yamashita Y, Tsukioka Y, Tomihisa K, Nakano Y, Koga T.** 1998. Genes

involved in cell wall localization and side chain formation of rhamnose-glucose

polysaccharide in Streptococcus mutans. J Bacteriol **180:**5803-5807.

33.   **Ruiz N.** 2015. Lipid Flippases for Bacterial Peptidoglycan Biosynthesis. Lipid

insights **8:**21-31.

34.   **Gomez-Casati DF, Martin M, Busi MV.** 2013. Polysaccharide-synthesizing

907   glycosyltransferases and carbohydrate binding modules: the case of starch

908   synthase III. Protein and peptide letters **20:**856-863.

909   35.   **Lairson LL, Henrissat B, Davies GJ, Withers SG.** 2008. Glycosyltransferases:

910   structures, functions, and mechanisms. Annu Rev Biochem **77:**521-555.

911   36.   **Korres H, Mavris M, Morona R, Manning PA, Verma NK.** 2005. Topological

912   analysis of GtrA and GtrB proteins encoded by the serotype-converting cassette

913   of Shigella flexneri. Biochem Biophys Res Commun **328:**1252-1260.

914   37.   **Skovierova H, Larrouy-Maumus G, Pham H, Belanova M, Barilone N,**

915   **Dasgupta A, Mikusova K, Gicquel B, Gilleron M, Brennan PJ, Puzo G, Nigou**

916   **J, Jackson M.** 2010. Biosynthetic origin of the galactosamine substituent of

917   Arabinogalactan in Mycobacterium tuberculosis. J Biol Chem **285:**41348-41355.

918   38.   **Schaefer K, Matano LM, Qiao Y, Kahne D, Walker S.** 2017. In vitro

919   reconstitution demonstrates the cell wall ligase activity of LCP proteins. Nature

920   chemical biology **13:**396-401.

921   39.   **Ardiccioni C, Clarke OB, Tomasek D, Issa HA, von Alpen DC, Pond HL,**

922   **Banerjee S, Rajashankar KR, Liu Q, Guan Z, Li C, Kloss B, Bruni R,**

923   **Kloppmann E, Rost B, Manzini MC, Shapiro L, Mancia F.** 2016. Structure of

924   the polyisoprenyl-phosphate glycosyltransferase GtrB and insights into the

925   mechanism of catalysis. Nat Commun **7:**10175.

926   40.   **Frickey T, Lupas A.** 2004. CLANS: a Java application for visualizing protein

927   families based on pairwise similarity. Bioinformatics **20:**3702-3704.

41.   **Hancock LE, Gilmore MS.** 2002. The capsular polysaccharide of Enterococcus

faecalis and its relationship to other polysaccharides in the cell wall. Proc Natl

Acad Sci U S A **99:**1574-1579.

42.   **Michon F, Chalifour R, Feldman R, Wessels M, Kasper DL, Gamian A,**

Pozsgay V, Jennings HJ. 1991. The alpha-L-(1----2)-trirhamnopyranoside

epitope on the group-specific polysaccharide of group B streptococci. Infect

Immun **59:**1690-1696.

43.   **Pritchard DG, Gray BM, Dillon HC, Jr.** 1984. Characterization of the group-

specific polysaccharide of group B Streptococcus. Arch Biochem Biophys

235:385-392.

44.   **Teng F, Singh KV, Bourgogne A, Zeng J, Murray BE.** 2009. Further

characterization of the epa gene cluster and Epa polysaccharides of

Enterococcus faecalis. Infect Immun **77:**3759-3767.

45.   **Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC,**

Punta M, Qureshi M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A.

2016. The Pfam protein families database: towards a more sustainable future.

Nucleic Acids Res **44:**D279-285.

46.   **Linton D, Dorrell N, Hitchen PG, Amber S, Karlyshev AV, Morris HR, Dell A,**

Valvano MA, Aebi M, Wren BW. 2005. Functional analysis of the

Campylobacter jejuni N-linked protein glycosylation pathway. Mol Microbiol

55:1695-1703.

47.   **Hoff JS, DeWald M, Moseley SL, Collins CM, Voyich JM.** 2011. SpyA, a C3-

like ADP-ribosyltransferase, contributes to virulence in a mouse subcutaneous

model of Streptococcus pyogenes infection. Infection and immunity **79:**2404-

2411.

48.   **Caparon MG, Scott JR.** 1991. Genetic manipulation of pathogenic streptococci.

Methods Enzymol **204:**556-586.

49.   **Horton RM, Hunt HD, Ho SN, Pullen JK, Pease LR.** 1989. Engineering hybrid

genes without the use of restriction enzymes: gene splicing by overlap extension.

Gene **77:**61-68.

50.   **Ruiz N, Wang B, Pentland A, Caparon M.** 1998. Streptolysin O and adherence

synergistically modulate proinflammatory responses of keratinocytes to group A

streptococci. Mol Microbiol **27:**337-346.

51.   **Chaffin DO, Beres SB, Yim HH, Rubens CE.** 2000. The serotype of type Ia and

III group B streptococci is determined by the polymerase gene within the

polycistronic capsule operon. J Bacteriol **182:**4466-4477.

52.   **Seiler CY, Park JG, Sharma A, Hunter P, Surapaneni P, Sedillo C, Field J,**

Algar R, Price A, Steel J, Throop A, Fiacco M, LaBaer J. 2014. DNASU

plasmid and PSI:Biology-Materials repositories: resources to accelerate

biological research. Nucleic Acids Res **42:**D1253-1260.

53.   **Nelson D, Loomis L, Fischetti VA.** 2001. Prevention and elimination of upper

respiratory colonization of mice by group A streptococci by using a bacteriophage

lytic enzyme. Proc Natl Acad Sci U S A **98:**4107-4112.

54.   **Raz A, Fischetti VA.** 2008. Sortase A localizes to distinct foci on the

Streptococcus pyogenes membrane. Proc Natl Acad Sci U S A **105:**18549-

18554.

55.   **Bui NK, Eberhardt A, Vollmer D, Kern T, Bougault C, Tomasz A, Simorre JP,**

Vollmer W. 2012. Isolation and analysis of cell wall components from

Streptococcus pneumoniae. Analytical biochemistry **421:**657-666.

56.   **Santander J, Martin T, Loh A, Pohlenz C, Gatlin DM, 3rd, Curtiss R, 3rd.**

2013. Mechanisms of intrinsic resistance to antimicrobial peptides of

Edwardsiella ictaluri and its influence on fish gut inflammation and virulence.

Microbiology **159:**1471-1486.

57.   **Edgar RJ, Chen J, Kant S, Rechkina E, Rush JS, Forsberg LS, Jaehrig B,**

Azadi P, Tchesnokova V, Sokurenko EV, Zhu H, Korotkov KV, Pancholi V,

Korotkova N. 2016. SpyB, a Small Heme-Binding Protein, Affects the

Composition of the Cell Wall in Streptococcus pyogenes. Frontiers in cellular and

infection microbiology **6:**126.

58.   **Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ.** 1990. Basic local

alignment search tool. J Mol Biol **215:**403-410.

59.   **Huang Y, Niu B, Gao Y, Fu L, Li W.** 2010. CD-HIT Suite: a web server for

clustering and comparing biological sequences. Bioinformatics **26:**680-682.

60.   **Alva V, Nam SZ, Soding J, Lupas AN.** 2016. The MPI bioinformatics Toolkit as

an integrative platform for advanced protein sequence and structure analysis.

Nucleic Acids Res **44:**W410-415.

61.   **Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K,**

Madden TL. 2009. BLAST+: architecture and applications. BMC bioinformatics

10:421.

62.   **Tsirigos KD, Peters C, Shu N, Kall L, Elofsson A.** 2015. The TOPCONS web

997   server for consensus prediction of membrane protein topology and signal

998   peptides. Nucleic Acids Res **43:**W401-407.

999   63.   **Soding J, Biegert A, Lupas AN.** 2005. The HHpred interactive server for protein

1000   homology detection and structure prediction. Nucleic Acids Res **33:**W244-248.

1001

1002

1003   Tables

1004   Table 1. The major fragments of the precursor ion at *m/z* 1048

m/z	Formula	Mass error (ppm)
78.95731	PO_3	-8.19
96.96791	H_2PO_4	-6.31
300.04883	$C_8H_{15}O_9NP$	3.12
845.65894	$C_{58}H_{88}ONP$	-1.02

1005   Fragment ions originating from the precursor molecular ion of 1048 were subjected to

1006   MS/MS fragmentation as described in materials and methods. The proposed molecular

1007   formulae of the product ions are supported by the observed isotope patterns for the

1008   relevant ions.

1009

1010

1011　Table 2. GlcNAc-lipid synthesis in membrane fractions from various bacterial strains.[a]

1012

Enzyme Source	GlcNAc-P-Und, pmol/mg	GlcNAc-P-P-Und, pmol/mg
MGAS5005	259.4±8.2	7.8±1.8
5005Δ*gacI*	Not Detected	15.9±2.5
GBS COH1	62±0.4	5.6±0.8
GBSΔ*gacI*	Not Detected	12±2
E. coli Rosetta (DE3)	Not Detected	19.5±0.3
E. coli:GacI	102.8±3.3	5.4±0.05
E. coli:EpaI	189±3.9	7±1.5

1013

[a]Reaction mixtures contained 50 mM Tris-HCl, pH 7.4, 5 mM 2-mercaptoethanol, 20

mM $MgCl_2$, 1 mM ATP, 5 µM UDP-[^{3}H]GlcNAc (486 cpm/pmol) and bacterial membrane

suspension (50-100 µg membrane protein) in a total volume of 0.01 ml. Following a 10

minute pre-incubation at 30°C, GlcNAc-lipid synthesis was initiated by the addition of

UDP-[^{3}H]GlcNAc. After 5 min, reactions were processed for GlcNAc-lipid synthesis as

described in Materials and Methods. The results are representative of at least three

separate experiments.

1021

1022 Table 3. Kinetic parameters of GlcNAc-lipid synthases in membrane fractions from

1023 various bacterial strains.[a]

1024

Enzyme source	GlcNAc-P-Und		GlcNAc-P-P-Und	
	Km, μM	Vmax, pmol/min/mg	Km, μM	Vmax, pmol/min/mg
MGAS5005	6.4	333	ND	ND
5005ΔgacI	ND	ND	19.3	2.75
E. coli:GacI	18.7	54.2	ND	ND
E. coli:GacI/J	1.1	18,500	ND	ND

[a]Membrane fractions from the indicated bacterial strains were assayed for the formation

of [^{3}H]GlcNAc-lipids in the presence of increasing amounts of Und-P, added as a

dispersion in 1% CHAPS. Apparent kinetic parameters for Und-P were calculated from

Michaelis-Menton plots (Fig. S5) using the linear regression algorithm in Sigma Plot ver.

12 (Systat Software, Inc. San Jose, CA). R squared values were 0.9953 for GAS GacI

expressed in *E. coli*; 0.983 for GAS GacI co-expressed with GacJ in *E. coli*, 0.9976 for

GacI in MGAS5005 and 0.931 for GacO in MGAS5005. The results are representative

of at least two separate experiments. ND = Not Detected.

1033

 Figure legends:

Figure 1. Map of *S. pyogenes* genes involved in Group A Carbohydrate (GAC)

biosynthesis and analysis of 5005Δ*gacI* and 5005 Δ*gacL* deletion mutants. (A) GAC

biosynthesis gene cluster. Numbers below represent MGAS5005 gene designations (B)

Representative immunoblot analysis of cell-wall fractions isolated from MGAS5005,

5005Δ*gacI*, 5005Δ*gacL* and 5005Δ*gacL gacL*$^+$. Data are representative of biological

triplicates. (C) Binding of N-acetyl glucosamine specific fluorescein-succinylated-WGA

to whole MGAS5005 and 5005Δ*gacL* was measured. Data are the average of three

replicates ± standard deviation. (D) Rhamnose and GlcNAc mole percentage of total

carbohydrate was determined by GC-MS for cell wall material isolated from MGAS5005

and 5005Δ*gacL* following methanolysis as described in Materials and Methods. Data

are the average of four replicates ± standard deviation. The asterisk indicate statistically

different values (* $p < 0.05$, ** $p < 0.01$) as determined by the Student's t-test. (E and F)

GC-MS chromatograms for glycosyl composition analysis of cell wall isolated

MGAS5005 and 5005Δ*gacL*. The deduced schematic structure of the repeating unit of

GAC is shown for each strain. The chromatograms are representative of four separate

analyses performed on two different cell wall preparations.

Figure 2. Purification and identification of GlcNAc-phosphate-undecaprenol in

5005Δ*gacL.* (A) Thin layer chromatography (TLC) analysis of phospholipids isolated

from MGAS5005 and 5005Δ*gacL*. Phospholipids extracted from bacterial strains were

separated by TLC on Silica Gel G in $CHCl_3/CH_3OH/H_2O/NH_4OH$ (65:25:4:1). Position of

the novel, alkaline-resistant and acid-labile, phospholipid accumulating in 5005Δ*gacL* is

indicated by the arrow. The results are representative of three separate experiments.

(B) ESI-MS/MS analysis of the novel phospholipid isolated from 5005Δ*gacL*. The

spectrum is assigned to GlcNAc-phosphate-undecaprenol.

Figure 3. Thin layer chromatography of [^{3}H]GlcNAc-lipids from *in vitro* incubations of

GAS mutants and *Bacillus cereus* membranes with UDP-[^{3}H]GlcNAc. Membrane

fractions from MGAS5005 (panel A), *B. cereus* (Panel B), 5005Δ*gacl* (Panel C), or

5005Δ*gacL* (Panel D) were incubated with UDP-[^{3}H]GlcNAc and analyzed for

[^{3}H]GlcNAc lipid synthesis by thin layer chromatography. Reaction mixtures contained

50 mM Tris-Cl, pH 7.4, 5 mM 2-mercaptoethanol, 20 mM MgCl$_2$, 1 mM ATP, 5 μM UDP-

[^{3}H]GlcNAc (486 cpm/pmol) and bacterial membrane suspension (100-200 μg

membrane protein) in a total volume of 0.02 ml. Following a 10 minute pre-incubation at

30°C, GlcNAc-lipid synthesis was initiated by the addition of UDP-[^{3}H]GlcNAc. After 10

min, reactions were processed for GlcNAc-lipid synthesis as described in Materials and

Methods. The organic layers were dried, dissolved in a small volume of CHCl$_3$/CH$_3$OH

(2:1) and a portion was removed and assayed for radioactivity by liquid scintillation

spectrometry. The remainder was spotted on 10 x 20 cm plate of silica gel G and

developed in CHCl$_3$/CH$_3$OH/NH$_4$OH/H$_2$O (65:25:1:4). [^{3}H]GlcNAc-lipids were detected

by scanning with an AR2000 Bioscan Radiochromatoscanner. The results are

representative of three separate experiments.

Figure 4. Thin layer chromatography of [^{3}H]GlcNAc-lipids from *in vitro* incubations of

GAS and GBS membranes with UDP-[^{3}H]GlcNAc. Membrane fractions from WT

MGAS5005 (panel A) or GBS COH1 (Panel B) were incubated with UDP-[^{3}H]GlcNAc

and analyzed for [^{3}H]GlcNAc-lipid synthesis by thin layer chromatography. Reaction

mixtures were exactly as described in the legend to Figure 3. After 5 min incubation with

UDP-[^{3}H]GlcNAc, reactions were processed for GlcNAc-lipid synthesis as described in

Materials and Methods. The organic layers were dried, dissolved in a small volume of

CHCl$_3$/CH$_3$OH (2:1) and a portion was removed and assayed for radioactivity by liquid

scintillation spectrometry. The remainder was spotted on 10 x 20 cm plate of silica gel G

and developed in CHCl$_3$/CH$_3$OH/NH$_4$OH/H$_2$O (65:25:1:4). [^{3}H]GlcNAc-lipids were

detected by scanning with an AR2000 Bioscan Radiochromatoscanner. The results are

representative of three separate experiments.

Figure 5. Effect of ATP on GlcNAc-P-Und synthesis *in vitro* in MGAS5005 membrane

fractions. GlcNAc-P-Und synthesis in MGAS5005 membranes was assayed after the

indicated time at 30 °C in the presence (○) or absence (●) of 1 mM ATP. Reaction

mixtures were identical to those described in Figure 3 except for the presence of ATP.

Following incubation, incorporation into [^{3}H]GlcNAc-P-Und was determined as

described in Materials and Methods. The results are representative of three separate

Figure 6. GacI and GacJ exist as a detergent-stable complex in the membrane. GacJ

and His-tagged GacI were co-expressed in *E. coli* Rosetta DE3 cells and extracted from

the membrane fraction in 2.5% CHAPS. The proteins were purified using Ni-NTA

agarose in the presence of 2.5% CHAPS. (A) Fractions collected during Ni-NTA

purification were analyzed by immunoblot using anti-His antibodies. (B) The eluted

proteins were analyzed by SDS-PAGE. The results are representative of three separate

experiments.

Figure 7. Analysis of GacO function in MGAS5005. (A) GacO catalyzes the synthesis of

GlcNAc-P-P-Und in *E. coli* membranes**.** Reaction mixtures contained 50 mM Tris-HCl,

pH 7.4, 5 mM 2-mercaptoethanol, 20 mM MgCl$_2$, 0.5 % CHAPS, 20 μM Und-P

(dispersed by ultrasonication in 1 % CHAPS), 5 μM UDP-GlcNAc (452 cpm/pmol) and

E. coli membrane fraction from either PR4019, CLM37 or CLM37:GacO strains.

Following incubation for 10 min at 30 ºC, incorporation of [^{3}H]GlcNAc into [^{3}H]GlcNAc-P-

P-Und was determined as described in Materials and Methods. Data are the average of

three replicates ± standard deviation.

(B and C) GlcNAc-P-P-Und can function as an acceptor substrate for rhamnosylation to

form Rhamnosyl-GlcNAc-P-P-Und in 5005Δ*gacI* membranes**.** Incubation conditions

were as described in the legend to Figure 3. (B) After 35 minutes incubation with UDP-

[^{3}H]GlcNAc, the reactions were analyzed for formation of [^{3}H]GlcNAc lipids as described

in Figure 3. (C) After 5 minutes incubation with UDP-[^{3}H]GlcNAc, the reactions were

incubated with 20 μM TDP-Rhamnose for an additional 30 min and analyzed for

formation of [^{3}H]GlcNAc lipids as described in Figure 3. The results are representative

of three separate experiments.

Figure 8. The absence of GacL increases sensitivity to cell wall amidases. Mid-

exponential phase MGAS5005 and 5005Δ*gacL* were grown in the indicated

concentrations of (A) CbpD, (B) PlyPy, and (C) PlyC. The change in growth is

represented as a percentage of growth where no amidase was present. Data are the

average of three replicates ± standard deviation. The asterisk indicate statistically

different values (** $p < 0.01$) as determined by the Student's t-test.

Figure 9. Schematic diagram of GAC biosynthesis. GAC is anchored to peptidoglycan

presumably via phosphodiester bond. GAC biosynthesis is initiated on the inner leaflet

of the plasma membrane where GacO produces GlcNAc-P-P-Und which serves as a

membrane-anchored acceptor for polyrhamnose synthesis catalyzed by the GacB,

GacC, GacF and GacG rhamnosyltransferases. Following polymerization polyrhamnose

is transferred to the outer leaflet of the membrane presumably by the GacD/ GacE ABC

transporter. Also in the inner leaflet of the membrane, GacI aided by GacJ produces

GlcNAc-P-Und which then diffuses across the plasma membrane to the outer leaflet

aided by GacK. Subsequently, GacL transfers GlcNAc to polyrhamnose using GlcNAc-

P-Und as glycosyl donor. Lastly, protein members of LytR-CpsA-Psr

phosphotransferase family presumably attach GAC to peptidoglycan. Several details of

this biosynthetic scheme are still speculative and further research will be required to

definitively confirm this hypothetical pathway, but the overall organization is consistent

with other isoprenol-mediated capsular polysaccharide pathways.

Figure 10. Sequence relationship of GacI family of proteins. Homology between GacI

homologs is graphically displayed using CLANS analysis (40). Dots correspond to

individual protein sequences selected as described in Material and Methods and

provided as Supplementary File. Selected homologs of *S. pyogenes* are highlighted by

colored dots.

GlcNAc-P-P-Und   GlcNAc-P-Und

- 🔴 *Streptococcus pyogenes*
- 🟡 *Streptococcus agalactiae*
- 🔵 *Enterococcus faecalis*
- 🔵 *Bacillus cereus*
- 🟣 *Clostridium sordellii*
- 🟢 DUF2304