Somatic maintenance alters selection acting on mutation rate

Authors:

*Andrii Rozhok[1], *James DeGregori[1,2,3,4]*

Affiliations:

[1]Department of Biochemistry and Molecular Genetics,

[2]Integrated Department of Immunology,

[3]Department of Pediatrics,

[4]Department of Medicine, Section of Hematology, University of Colorado School of
Medicine, Aurora, CO 80045

Emails:

Andrii Rozhok: Andrii.Rozhok@ucdenver.edu

James DeGregori: James.DeGregori@ucdenver.edu

Abstract

The evolution of multi-cellular animals has produced a conspicuous trend toward
increased body size. This trend has introduced at least two novel problems: an elevated risk of
somatic disorders, such as cancer, and declining evolvability due to reduced population size,
lower reproduction rate and extended generation time. Low population size is widely
recognized to explain the high mutation rates in animals by limiting the presumed universally

negative selection acting on mutation rates. Here, we present evidence from stochastic

modeling that the direction and strength of selection acting on mutation rates is highly

dependent on the evolution of somatic maintenance, and thus longevity, which modulates the

cost of somatic mutations We argue that this mechanism may have been critical in facilitating

animal evolution.

Keywords: somatic maintenance, longevity, body size, mutation rate, selection

Introduction

Increasing body size has been one of the major trends in animal evolution across many taxa, as formulated in Cope's rule (*1, 2*). The evolution of larger bodies introduces some fundamentally new evolutionary challenges. The carrying capacity of ecosystems limits biomass per group/species, so larger body size leads to reduced population size. Furthermore, large animals generally demonstrate lower reproduction rates and longer generation times. In aggregate, such changes weaken selection that can act on a population and thus negatively affect evolvability. This general reduction in evolvability should, however, be at least partially alleviated by diversity facilitated by sexual reproduction.

The mutation rate (MR) is another critical evolvability parameter. It is believed that selection generally acts to lower MR (*3-5*), and the significantly higher MRs observed in animals compared to unicellular organisms have been argued to result from the reduced power of selection imposed by small population sizes (*6-8*). Germline (gMR) and somatic (sMR) mutation rates are linked, as they employ the same basic DNA replication and repair machinery (*9-11*). While elevated gMR improves evolvability, the ensuing higher sMR should elevate the risk of somatic disorders, such as cancer (*12*). For cancer, increasing body size is expected to increase the frequency of oncogenic mutations by increasing the number of target cells (*13*). Somatic mutations also contribute to aging and a variety of aging-related diseases (*14*). The increased cost of sMR should thus exert negative selective pressure on gMR in larger animals.

Recent evidence demonstrates that the sMR in some animal tissues can be significantly higher than the rate inferred from observed mutations, because somatic purifying selection is very effective at eliminating damaged somatic cells (*15*). Many mechanisms, such as various tumor suppressor gene functions (including DNA damage induced apoptosis) (*16*), autophagy (*17*), purifying somatic selection (*15, 18*), and immune surveillance (*19*), should buffer the costs of somatic mutation and in aggregate promote lifespan extension by maintaining tissue

integrity. We will collectively call these mechanisms – the *somatic maintenance program* (SMP).

We present theoretical evidence from Monte Carlo modeling indicating that somatic maintenance not only improves individuals' survival for large animals by reducing sMR costs, but should have played a crucial role in animal evolution by substantially modifying selection acting on gMR. We show that positive selection for increased body size promotes positive selection for extended longevity by improving SMP. Our results also indicate that positive selection acting on traits that do not impact somatic risks also promotes selection for an improved SMP. In both cases, positive selection acting to increase gMR was observed because of the reduced sMR cost, which dramatically improved evolvability of the simulated population.

Results

Theoretical introduction to the modeling. We built a stochastic model of evolution in animal populations, incorporating reproduction and survival (**Fig. 1**), whereby each individual's trait is inherited with variance proportional to gMR (for code, see **Supplements: Section 1a**). Traits are assumed to be polygenic and exhibit phenotypic variation in the population. In particular, MR is assumed to be a highly polygenic trait, given the many genes responsible for DNA repair, DNA replication, damage avoidance (e.g. anti-oxidant defenses), and mutagen detoxification, which in aggregate can determine MR. The evolution of body size, somatic maintenance and germline mutation rate was then tracked under various regimens of selection (see also **Methods: Model algorithm**).

The model should reasonably approximate a sexually reproducing population. The model operates with single-parent reproduction model so that each individual descends from one parent. In this regard, technically it is tempting to view it as a model of an asexual population. However, at a higher level of abstraction the fundamental difference between sexual and

asexual populations (aside from the issue of purging deleterious mutations) is the amount of
variation produced per the same size population per generation. Variance of inheritance in our
model is too high to be assumed as being generated by mutations accumulating along a clonal
lineage and equals 10% of a trait's value per generation within 95 percentile. As the modeled
traits are assumed to be *multigenic* and have a continuous phenotypic range in the population,
we did not need to simulate the processes of allelic segregation by recombination in order to
reconstruct a sexual population. Moreover, to model allelic segregation would require
assumptions regarding the number of genes and alleles underlying a trait, the dominance of
these alleles, and their relative contributions to the phenotype. As such, the model only
operates with the net ultimate change of a trait over generations. The assumed multigenic
nature of the simulated traits also means that both segregation of alleles by recombination and
aggregation of alleles by co-selection are impeded. The efficiency of allele segregation for
multigenic traits is inversely proportional to the number of genes encoding a trait. We therefore
assume that the net co-evolution of a pair of multigenic traits will ultimately depend on
selection acting on these traits that can overcome allele segregation effects.

The model incorporates three major factors of mortality, including aging. Human life tables
indicate that aging proceeds exponentially, whereby mortality and diseases accelerate at
advanced ages (e.g. https://www.ssa.gov, https://seer.cancer.gov). The combined action of
SMP mechanisms provides for an extended early period of high body fitness with little to no
decline. We generalized this complex program in a curve that describes modeled animal
mortality of physiological causes schematically shown in **Fig. 2A** and based on the following
equation:

$$D_A = M \times e^{A^{Som}} \qquad (1)$$

where D_A is the probability of dying of physiological causes <u>at</u> age A, M is mutation rate,
and *Som* is a composite parameter that determines SMP efficiency. The cumulative
distribution function of D_A, or the probability of dying of physiological causes <u>by</u> age A,
resembles human mortality (**Fig. 2B**). The equation should thus provide a robust model for
aging-related mortality, reflecting the extended period of high fitness and the late-life

accelerating mortality. **Fig. 2A** also demonstrates the relative effects of MR, which is a linear

contributor, and the *Som* parameter, which stands for the total damage buffering capacity of

the SMP (for details and theory see **Methods: The somatic maintenance program paradigm**). It

is important to keep in mind that the *M* parameter (mutation rate) in **Eq. 1** is responsible for the

somatic costs of MR (higher MR in **Fig. 2A** accelerates aging-related mortality). Improved

SMP, just as body size or other trait, may come at a cost on a short evolutionary time scale,

which is later diminished by further adaptation. We did not include this cost in the modeling,

since if a trait responds to directional selection this means that the benefit outweighs the cost.

Since the amount of change of a trait as a result of positive selection in our model is arbitrary

(not exactly copying any particular natural species), we can conclude that this amount of

change could already incorporate the net benefit minus cost. In other words, if the benefit of an

evolutionary change exceeds its cost, then modeling benefit and cost on an arbitrary scale is

mathematically equivalent to modeling only benefit.

The evolution of SMP promotes selection for increased body size through better tolerance

of MR. In our simulations, positive selection for increased body size (**Fig. 2C, green**) led to a

concurrent selection for elevated gMR (**Fig. 2D, green**) and improved SMP (**Fig. 2E, green**).

Artificially blocking SMP evolution by fixing SMP at the initial value (**Fig. 2E, blue**) significantly

slowed the evolution of body size (**Fig. 2C, blue; p << 0.001**) and triggered selection for lower

gMR (**Fig. 2D, blue**). We implemented the ecosystem carrying capacity by setting a maximum

biomass for the population; therefore, increasing body size led to a corresponding decline in

population numbers, amplifying the power of drift (**Fig. 2F,G**). When SMP was allowed to

evolve, however, the population entered a "drift zone" when its size decreased to ~4,000

individuals, which shortly thereafter was overcome by selection for even larger body size,

visible also by a continuing decline in population numbers (**Fig. 2F**). When we artificially

blocked SMP, however, the drift zone was more profound. It occurred earlier at the population

size of ~6,000-7,000 individuals, and the population was not able to escape from it (for ~1,000

generations) and restore its initial rates of evolution (**Fig. 2G**), indicating an important role of

SMP evolution in maintaining evolvability. We further generated a population with two

simulated genotypes – Genotype A that could evolve SMP (10% of the population) and

Genotype B with SMP fixed at the initial value (90%). We set a maximum population size and

removed the maximum biomass limit to rule out body mass effects on population size and

selection, and tracked Genotype A and Genotype B frequencies under positive selection for

increased body size (for code see **Supplements: Section 1b**). Despite the initial abundance,

Genotype B (with fixed SMP) lost the competition in less than 200 generations, reflecting a

direct competitive advantage of the capacity to evolve enhanced SMP (**Fig. 2H**). Hereafter, we

will call the setting with positive selection for increased body size and freely evolving SMP and

gMR the *standard condition* (usually shown in green, unless otherwise indicated) used in

comparisons with other selection regimens.

Abrogating selection for increased body size reduces selection for gMR and SMP. In the

absence of positive selection for increased body mass (**Fig. 3A, blue**), both gMR (**Fig. 3B, blue**)

and SMP (**Fig. 3C, blue**) demonstrate early positive selection, which appeared to have been

caused by rapid evolution of reproductive parameters (see **Supplement: Section 2**). Overall,

gMR demonstrates a significant general decrease (non-overlapping confidence intervals (CIs)

at the beginning relative to the end of the simulation), and SMP undergoes a significantly

smaller improvement compared to the standard condition (green; $p \ll 0.001$). Blocking the

evolution of body mass (**Fig. 3D, blue**) and SMP (**Fig. 3F, blue**) expectedly led to strong

selection for lower gMR (**Fig. 3E, blue**) compared to the standard condition ($p \ll 0.001$), which

we interpret as being driven by the sMR costs in the absence of benefits of high gMR. In other

words, mutation rate is selected against because of its somatic costs and the absence of

benefits of higher gMR in static conditions. In natural populations that are under stabilizing

selection, gMR will have costs due to greater phenotypic variance from a well-adapted state

that are independent of sMR, but we do not model stabilizing selection in this study.

Decoulpling sMR cost from gMR enhances the evolution of larger bodies. To investigate

the role of the putative gMR benefit versus sMR cost balance in evolution, we further

decoupled gMR and sMR by allowing gMR to evolve but making sMR cost fixed and

independent of gMR (see **Methods: Model variations**). Decoupling sMR cost from gMR

significantly accelerated the evolution of body size (**Fig. 3G, blue**) relative to the standard

condition (green; p = 0.0052), revealing that sMR costs can limit the evolution of larger body

size. During the early fast evolution of body mass, gMR (**Fig. 3H, blue**) and SMP (**Fig. 3I, blue**)

demonstrate a corresponding positive response. Later, further body mass evolution becomes

impeded (likely because of the severe depletion in population numbers), coinciding with

selection for lower gMR. SMP plateaus during this second phase at a significantly lower level

compared to the standard condition (p << 0.001), indicating that the somatic costs of mutation

rate stimulate the evolution of more robust SMP.

Selection acting on a somatic cost-unrelated trait still promotes selection for increased

gMR and SMP. As we have seen under blocked selection for increased body size (**Fig. 3B,C,**

blue), SMP demonstrates an early phase of positive selection (**Fig. 3C, blue**) that is apparently

reflected in a corresponding positive selection for higher gMR (**Fig. 3B, blue**). This observation

suggests that both SMP and gMR may also respond to selection acting on some other traits,

e.g. reproductive parameters (**Supplements: Section 2**). This raises the question whether SMP

and gMR evolution would be sensitive to strong selection for a trait that does not affect somatic

risks (greater body size increases the target size for somatic mutations). We simulated a

condition that was similar to the standard condition, except positive selection was applied to a

trait that did not affect sMR related somatic costs (see **Methods: Model variations**); e.g. if SMP

improvement is solely a response to the increased sMR cost imposed by larger body, selection

acting on an sMR cost unrelated trait should not drive improvements in SMP. As shown in **Fig.**

3J (blue), unimpeded by increased sMR costs and declining population size, the evolution of

an sMR cost unrelated trait is significantly faster compared to the evolution of increased body

size (p << 0.001). Interestingly, gMR (**Fig. 3K, blue**) also demonstrated an early phase of

positive selection during early rapid evolution of the selected trait and remains above the initial

gMR throughout the entire simulation. As anticipated, SMP is positively selected; however, in

the absence of an increasing sMR cost associated with larger bodies, SMP's improvement is

significantly smaller (**Fig. 3L, blue,** p << 0.001). Notably, even with much less enhanced SMP,

gMR is still under positive selection in response to positive selection for the sMR cost

unrelated trait (**Fig. 3K, blue**), consistent with the sMR/gMR cost/benefit ratio being an
important factor regulating selection acting on gMR. Regardless, the results demonstrate that
both gMR and SMP are responsive to selection for somatic risk unrelated traits, which
indicates that high mutation rate is beneficial in positively selective conditions.

SMP enables maintenance of gMR when directional selection is weak. As we have seen in
Fig. 3D-F, in the absence of strong positive selection for increased body size and SMP
efficiency, selection acts to lower gMR. **Fig. 4** shows, however, that this selection is
significantly modified by the efficiency of SMP. Stronger SMPs (lower *Som* value) relax
selection for lower gMR when directional selection is weak (non-overlapping CIs between the
standard (red) and either of the improved SMPs). As will be explained further below, this
observation may have significant implication on long-term species survival in relatively static
environments.

Modeling competition between a wildtype and mutator phenotypes. Under strong positive
selection, whether for increased body mass (**Fig. 2A-C, blue**) or a sMR cost unrelated trait (**Fig.
3H,I, blue**, and **Fig. 3K,L, blue**), we observed consistent signs of positive selection for higher
gMR. However, because gMR and sMR are linked, higher gMR is a trait that should negatively
impact individual fitness and therefore be under negative selection. To investigate this
question, we mixed two simulated genotypes, one "wild-type" (50%) and one "mutator" (50%)
in a population of stable size and under positive selection for a sMR cost unrelated trait. We
then observed the genotypes' frequencies in the population using varying strength of mutators.
Fig. 5A demonstrates that while the mutator's fitness initially is lower compared to wild-type,
eventually the mutator outcompetes its wild-type counterpart. Interestingly, with increased
mutation rate, the magnitude of the mutator's initial decline increases, but so does the speed at
which it subsequently overtakes the population. This result provides a clue for how higher
mutation rate, being a trait with negative impact on fitness, can be selected for. Because net
organismal fitness is a composite trait impacted by the fitness value of many individual traits,
the initial fitness of the "mutator" is lower because, all other traits equal, higher MR incurs
increased sMR cost. However, in response to selection, mutator is capable of more rapidly

developing other (adaptive) traits (**Fig. 5B**) and thus its overall fitness soon becomes higher compared to wild-type.

Discussion

Our study demonstrates that positive selection for increased body size triggers a concurrent selection for improved somatic maintenance to mitigate the increased somatic risks of larger bodies. Improved somatic maintenance, in turn, promotes selection for higher germline mutation rates by reducing the cost of somatic mutations and thus altering the sMR/gMR cost/benefit ratio. Conditions of strong positive selection for somatic cost independent traits, as our model shows, can also alter this balance by elevating the benefits of higher gMR. Under stable conditions, alternatively, the sMR/gMR cost/benefit balance is altered by the existing cost of somatic mutations and by the increased cost and absent/reduced benefits of gMR itself, which ultimately favors lower mutations rates. Under stasis, gMR exerts a cost independent of somatic risks by increasing deviation of progeny phenotypes from population mean/median and thus reducing their fitness. Our study thus demonstrates that the evolution of mutation rate is not under a universal population size-dependent selection acting to lower it, but is highly tunable and governed by selection acting on other traits. Importantly, our modeling indicates that under certain conditions elevated mutation rate, unlike perhaps any other trait, can be positively selected despite its negative effects on individual fitness (as explained in **Fig. 5**).

Mutation rate in eukaryotes is a highly polygenic trait encoded by multiple genes involved in DNA replication, repair, damage avoidance, and cell division machineries (*9, 11*). Animals mostly reproduce sexually, which should generate an extensive population allelic diversity for these genes. This diversity should provide for a relatively continuous distribution of mutation rate in populations, rather than being a uniform trait marked with sporadic monogenic mutants, as may occur in asexual populations (*20-22*). Such intra-population variation (*23, 24*), as well as the ability of mutation rate to rapidly evolve (*25*), has been shown for humans . However,

sexual reproduction would be supposed to effectively segregate alleles contributing to mutation
rate from alleles for other (e.g. adaptive) traits. It has been argued based on other evidence
that the efficiency of such segregation in sexual populations is limited (*26*). In particular, as
argued in **Theoretical introduction to the modeling**, the multi-genic nature of the gMR trait
should substantially slow segregation of gMR from other traits.

It also appears from our results that animal evolution, with the macroscopic trend toward
larger bodies, should have driven a concurrent evolution of extended longevity, the latter being
determined by the efficiency of species-specific somatic maintenance programs. Even though
extended longevity tentatively appears to be a benefit on its own, e.g. due to extended
reproduction period, our model demonstrates that somatic maintenance (and thus longevity) is
under a much weaker positive selection in the absence of other positively selected traits. This
observation can explain why extended longevity demonstrates significant deviations across
animal taxa from the general rule larger body → longer lifespan. Our results indicate that the
evolution of longevity (as a function of somatic maintenance efficiency) should be greatly
impacted by the rate of evolution of other traits, and not necessarily body size.

Interestingly, our study predicts an important evolutionary role for the mechanisms of
somatic maintenance in addition to their evolution as a means of improving individual survival
of large animals (*13, 18*). Our results demonstrate that selection for enhanced somatic
maintenance goes well beyond the evolution of body size and is promoted by strong directional
selection acting on any trait. This result indicates that SMPs may have had an important role in
the evolution of large animals. Selection for higher gMR following improved SMP may be an
important mechanism "rescuing" the reduced evolvability imposed by reduced population size,
extended generation times and lower reproduction rates. Therefore, SMPs and longevity may
have an important contribution to species' long-term survival. For example, a prolonged
evolutionary stasis (*27-30*) should trigger selection for lower mutation rates. By relaxing
selection for lower mutation rate and thus maintaining evolvability (as shown in **Fig. 4**),
enhanced SMPs can ensure better survival of animal groups facing rapid evolutionary
transitions or drastically changed environments after such relatively static periods. All other

traits equal, species with extended longevity may survive such transitions with higher
probabilities.

Lynch and colleagues have provided extensive arguments supporting the idea that the
higher MRs in animals compared to unicellular organisms are likely to be caused by reduced
population sizes that limit the ability of selection to lower mutation rate (*6-8*). In conjunction
with population size, in large animals the strength of selection will be further attenuated by
lower reproduction rates and extended generation times. Based on our results, Lynch's theory
can be extended by recognizing that somatic maintenance programs (and longevity) should
have substantial influence on the general relationship between population size and mutation
rates, and on the strength and directionality of selection acting on mutation rates. For example,
in our simulation, populations of the same initial size but with different SMP efficiencies
demonstrate profound differences in the effects of population size driven weakening of
selection (**Fig. 2F,G**, as well as discrepant selection for mutation rates (**Fig. 2D**).

Selection for higher mutation rates has been shown experimentally in bacteria (*20-22, 31*),
whereby engineered or spontaneous mutants with higher mutation rates have been shown to
have advantages over wild-type under positively selective conditions. The "mutator hitchhiker
hypothesis" explains such selection by the higher probability that adaptive mutations will
appear in a mutator cell (*22*). Once such a mutation occurs, the mutator genotype spreads to
fixation by being genetically linked to the adaptive phenotype. Modeling studies demonstrate
that evolution of evolvability, including varying selection on mutation rates, should be possible
in sexually reproducing organisms (*26, 32, 33*). Yet robust experimental corroboration of such
a possibility appears to be lacking.

In conclusion, our results raise the question of whether the evolution of large body size in
animals would be possible without such a complex pattern of selection acting on mutation rate,
and whether such a complex relationship is necessary to explain the evolution of large
animals. The evolution of large bodies has entailed the cost of losing the ability to evolve via all
major parameters that define this ability, such as population size, reproduction rate and

generation time, <u>except</u> mutation rate (which increased). Therefore, one scenario could have

been that this cost has been so prohibitive for many species that positive selection for mutation

rate was necessary to allow evolution of large animals. Alternatively, mutation rate could have

been high enough to maintain evolvability at the selection/drift barrier point where selection

was no longer able to reduce it further (*6*). Understanding which of these scenarios prevails in

the evolution of large animals requires more research.

Methods

Software. The model was created and all simulations were run in the Matlab environment
(MathWorks Inc, MA) version R2014a.

Model algorithm. The model is a stochastic Monte Carlo type model (the exact algorithm can
be found in **Supplements: Section 1a**) that runs a total of 1,005,000 updates ("time" in arbitrary
units, AU) unless otherwise stated, which represents ~1000 generations of the simulated
animal population (see **Fig. 1** for the flow chart). The simulation starts with building an initial
population of 10,000 individuals. Each individual has a number of simulated traits: 1) ID, which
is 1 (monogenotypic population) or 1 and 2 (in experiments with competition between two
genotypes in a mixed population to indicate genotypes); 2) current age, which increments by 1
at each simulation update; 3) inherited body mass, which is inherited with variation by an
individual and will be reached by adulthood (at age ~1000) and equals 5000 AU in the initial
population; 4) current body mass, which changes during individual growth, following a growth
curve, and plateaus at the inherited body mass in adults; 5) inherited birth mass, which in
individuals of the initial population is 300 AU; 6) inherited mutation rate of 10^{-9} AU (explained
below); 7) inherited reproduction rate, which is the period with variation between successive
reproductions in adult individuals and equals ~600 in the initial population; 8) inherited litter
size (initially 1), which is the number of progeny produced per individual per reproduction; 9)
inherited parameter of somatic maintenance, which determines the strength of the somatic
maintenance program as further explained below; 10) age of first reproduction, which dictates
that an individual begins reproducing when its current body mass reaches 0.9693 of its
inherited adult body mass (the number is derived so that in the initial population maturity is
reached at age ~1000 based on the growth curve).

Each inherited trait varies in progeny relative to parental. This variation was produced by
multiplying the inherited mutation rate by the parameter of inherited variance (*inhvar* =
250,000,000) and the product was used as the standard deviation (STD) of the normally
distributed variation in inheritance. This transformation was not necessary, as the *inhvar*
parameter is constant throughout simulation and it simply determines the magnitude of the
mutation rate's effects in the germline, which is imaginary and in the initial population simply

produces 0.000000001 x 25,000,000 = 0.025 that serves as the STD parameter for the normal

distribution from which inheritance variation is drawn. However, we kept this two-parametric

model for inheritance because mutation rate is also separately used in the equation of the

somatic maintenance program (as will be explained later).

Each newborn individual grows, reaches maturity, then reproduces over the rest of its lifetime

and eventually dies. The model is asynchronous, so that at every time-point of the simulation

the population contains individuals of various ages whose lifecycles develop independently.

And finally, three factors of mortality were modelled in the simulations. First, at every timepoint

of the simulation, an individual could die of somatic causes with a certain probability. This

probability is small at the beginning of life (but still can be caused by some imaginary inherited

genetic defects) and increases exponentially with age based on the paradigm of the aging

curve, which is primarily determined by an individual's inherited somatic maintenance program

(SMP). In humans, the aging curve also depends on lifestyle, however we assume in this

model that in a wild animal population lifestyle distribution is sufficiently uniform to be

neglected. More detailed description of the somatic maintenance paradigm that we applied will

be explained further below. Secondly, the simulated animals had a chance of dying of external

hazards, such as predators. We applied the Lotka-Volterra model of predator-prey interactions

(*34, 35*) to implement the dynamics of predator pressure (effectively the chance of dying of an

external hazard cause per timeunit). Here we should mention that smaller individuals and

juveniles had higher chances of dying of external hazards, which effectively created positive

selection for increased body size and also reflected the typical high mortality rates among

juveniles observed in natural populations. And lastly, individuals could die of intra-specific

competition. We implemented such competition by setting the upper limit of population's total

biomass, which in nature is imposed by the ecosystem's carrying capacity. Therefore, in the

simulated population biomass produced over the biomass limit caused additional mortality, so

that stochastically, population total biomass never exceeded the limit. Larger individuals also

had lower probability of dying of intra-specific competition, based on the assumption that

competition for resources and mates (the failure to reproduce is effectively an evolutionary

death) will typically favor larger individuals and this should have been one of the forces that

has been driving the macroscopic animal evolutionary trend towards increasing body size. The

advantage of size in this mortality model also created additional positive selective pressure for

body size. The total age-dependent mortality of all causes in our model did approximate a

typical wild animal mortality curve (**Supplements: Section 3**).

The somatic maintenance program paradigm. In order to replicate natural mortality caused by

physiological aging, such as cancer, decreased immune defense and lower ability to avoid

predators or to succeed in intra-specific competition, we made use of the aging curve, or

somatic maintenance, concept. Modern humans (in developed nations) and captive animal

mortality curves (**Fig. 2B** for human) differ from wild animal mortality curves in very high early

life survival with most mortality significantly delayed into advanced ages (*36, 37*). This

difference is caused by many reasons, such as much lower mortality caused by external

hazards and better nutrition and general healthcare. It therefore can be assumed that the

human and captive animal mortality curves are close representations of the physiological aging

curve. As longevity depends on multiple mechanisms of maintaining the soma, we can also

call this curve *the somatic maintenance curve*. In order to reconstruct this curve, we assumed

that somatic maintenance depends on the interaction of two opposing forces: 1) the

accumulation of genetic and structural damage in the soma that promotes aging and 2) the

somatic maintenance program consisting of a number of mechanisms that prevent or buffer

the effects of genetic and structural damage. The exact mathematical relationship between

these two forces and age is not known, however an example of cancer development can be

used as a proxy to explain the equation we derived for it.

Oncogenic mutations (including oncogenic epigenetic changes) are the ultimate necessary

condition for cancer to develop. The frequency of oncogenic mutations linearly depends on

mutation rate on a per cell division basis. Therefore, we assume that linear changes in

mutation rate will have linear effects on the odds of the occurrence of oncogenic mutations. An

oncogenic mutation provides the initiated cells with a linear change in their fitness relative to

normal cells. However, over time an advantageous clone with a constant linear fitness

advantage will proliferate exponentially. Therefore, we can already assume that mutation rate

should have a linear effect on the cancer curve, while time/age adds an exponential

component revealed in an exponential growth of a tumor. We can reasonably assume further

that a strong SMP will efficiently suppress such a clone, slowing or even preventing its growth

(*38*). A weaker SMP will allow the clone to proliferate faster. Therefore, SMP strength can

modulate the effects of mutations and time on cancer risk. The exact relationship between

SMP strength and physiological risk factors is not known. However, we know that their

interaction leads to a net exponent in physiological decline and disease risk.

We therefore reconstructed the human aging curve by maintaining the general principal

relationship between these factors as shown in **Eq. 1**. As seen from the equation, mutation rate

is a linear contributor to aging. Age itself contributes exponentially, and the somatic

maintenance composite parameter *Som* is, in turn, in power relationship to age. The

cumulative distribution function of D_A (**Eq. 1**) produces *D(A)* – the probability of dying of

somatic/physiological causes <u>by</u> age *A* and yields a shape close to the human mortality curve

(**Fig. 2A,B**). We cannot claim that these three factors are in the exact relationship predicted by

Eq. 1, as it is unknown. As seen in **Fig. 2A**, changes in the *Som* parameter have substantially

greater effects on the resulting mortality curve than mutation rate, with mutation rate still

having a sizeable effect as well. Yet claims are still made (e.g. (*39*)) that mutation rate is a

larger factor in aging than we assume in this model. Validation of our assumption in general

comes from the body of solid evidence that up to 50% of mutations in humans accumulate

during body growth by the age 18-20 (*40-42*). If mutation accumulation had a significant effect

on aging on its own, we should age rapidly until age 18-20 (half-way) and then the rate of

aging should decelerate. However, in reality the opposite happens, indicating that the

combined strength of the SMP has an overpowering effect in modulating the effects of genetic

damage on aging. As a result, we reason that **Eq. 1** might reasonably approximate the natural

relationships of these three factors. Therefore, based on an individual's aging curve we

calculated the D_A parameter at each simulation time-point (using the individual's mutation rate,

age and *Som* parameter) and applied it in a binomial trial as the probability of that individual's

dying of somatic/physiological causes in an age-dependent manner. As further explained in

Supplements: Section 4, the exact relationship between the *Som* parameters and each of the

other two (mutation rate and age) has no effect on the model, as the model represents SMP

and its variation by using area under the mortality curve. Therefore the sole purpose of **Eq. 1** in

the model is to generate an age-dependent curve of physiological mortality whose cumulative

function (probability of dying <u>by</u> a certain age) resembles in shape the human mortality/aging

curve (see **Supplements: Section 4** for detailed explanation and illustration).

Model variations. A number of model variations used in simulation experiments are

employed. *Fixed trait values* involved simply fixing the initial trait value without inherited

variation throughout the entire simulation. *Dislinking of somatic and germline mutation rate* was

done by making the value M in **Eq. 1** independent of an individual's mutation rate, which

resulted in somatic costs independent of transgenerational variation of mutation rate

(effectively from germline mutation rate). *Selection for a trait that did not affect somatic risks*

was achieved by transforming the "body mass" trait's effects by removing the trait from

calculations of the risk of death by somatic causes (unlike body size, it did not influence the

risk), then removing the population biomass limit and setting maximum population size (unlike

body mass, other traits do not directly affect population numbers) and fixing the growth rate

curve so that it reached the initial body mass of 5,000 AU (the current body mass parameter in

the model; the inherited body mass variation did not exist and the inherited body mass

parameter was replaced with the somatic risk unrelated trait). These manipulations made the

selected trait a proxy for a trait unrelated to somatic risks (e.g. hair color). *Competitive assays*

included individuals with different ID parameters, such as 1 and 2 to indicate different

"genotypes"; traits of the "genotypes" then were tracked and stored separately.

Data processing. Processing of primary data included removal of outliers (see

Supplements: Section 5). Occasionally the simulations generated "NaN" (not a number) values

in individual parameters, which were rare but quickly propagated if left in the population. We

immediately deleted individuals from the population if "NaN" values appeared in any of their

parameters. Based on the rarity of such events, we can assume that they had the effect of rare

early lethal mutations and affected the population at random. Thus, we assume these did not

affect the principal results.

Statistics and data presentation. Most simulation experiments were made with 25 repeats.

Due to heavy skews in sample distributions (inferred by D'Agostino-Pearson test for normality

of a distribution), all figure panels represent medians (thick lines) and 95 percentiles on each

tail (color-shaded areas). Statistical differences between experimental conditions were

calculated as follows. We first calculated the sum of all values in each run throughout the

entire evolution of a trait (typically 1,005,000 time points). In this way, given the small

increment over a long time the sum essentially approximated the area under the curve of a

trait's evolution. These sums (usually 25 repeats in one experiment/sample) were then

compared by applying the Matlab implementation of the Wilcoxon rank sum test, which is

considered equivalent to the Mann-Whitney U-test. P-values <= 0.05 were considered as

indicating significant difference.

Supplementary Materials

Section 1. Model code.

a. General model for positive selection for body size

```matlab
for iteration = 1 : 25
    disp(iteration);

    newrun = true;

    if(newrun)
        clearvars -except newrun iteration
        fname = 1;
        timeun = 1;
        iter = 1;

        % OUTPUT STORAGE MATRICES
        sommortality = []; % counts of mortality for somatic reasons
        extmortality = []; % counts of mortality caused by external hazard
        capmortality = []; % counts of mortality imposed by ecosystem's
                           %  carrying capacity (intra-specific competition)
        biomassdyn = [];    % population biomas dynamics over time
        popsizedyn = [];    % population size dynamics over time
        births = [];        % counts of new births over time

        bodymassevol = []; % population's average bodymass over time
        birthmassevol = []; % population's average birthmass over time
        littersizeevol = []; % population's average litter size over time
        mutrateevol = []; % population's average mutation rate over time
        rrateevol = []; % population's average reproduction rate over time
        lifespanevol = []; % population's average somatic maintenance
                           %  coefficient over time

    else
        timeun = timeunit + 1;
        if(fname == size(filenames, 2))
            fname = 1;
        else
            fname = fname + 1;
        end
    end
    filenames = ['a' 'b' 'c' 'd' 'e' 'f' 'g' 'h' 'i' 'j' 'k' 'l' 'm' 'n'...
        'o' 'p' 'q' 'r' 's' 't' 'u' 'v' 'w' 'x' 'y' 'z'];

    % GENERAL MODEL PARAMETERS:
    totaltime = 1005000; % total # of simulation updates ("time")
```

```matlab
516        popsize = 10000; % initial population size
517        mutrate = 0.000000001;
518        inhvar = 25000000; % a multiplier of mutation rate determining
519                               %  variance in trait inheritance (var=inhvar*mutrate)
520                               %  so that inhertance variance is proportional to
521                               %  mutation rate
522        bodymass = 5000; % initial adult bosymass
523        birthmass = 300; % initial body mass at birth
524
525        repbodymass = 0.9693; % multiplier determining at what body mass
526                               %  as a fraction of the individual's inherited
527                               %  adult body mass the individual begins to
528                               %  reproduce
529        rrate = 600; % initial time (in # simulation updates)
530                       %  between successive reproductions
531        littersize = 1; % initial # progeny per reproduction per individual
532        littervar = (0.1*littersize)/littersize; % variance of littersize
533        rratevar = (0.1*rrate)/rrate; % variance of reproduction rate
534        growthrate = 57; % coefficient of body growth rate
535
536        somdeath = 0.34; % an exponential coefficient of the somatic maintenance
537    equation
538        somEnergy = 2231.81365913237; % initial energy invested in somatic
539         %  maintenance when somdeath=0.34
540                                       %  see SUPPLEMENTS
541
542        aging = mutrate*exp([1:1000000].^somdeath); % the aging function -
543                       % probability of dying for somatic reasons over time
544
545        risk = cumsum(aging); % cumulative sum function of the aging curve
546        [c riskage] = min(abs(risk-1)); % riskage - age at which risk = 1;
547                                         %  explained in METHODS
548
549
550
551        %INITIAL BODY GROWTH FUNCTION
552        growthcurve = [birthmass]; % curve for body size distribution in the initial
553    population
554                                     % initial population is generated with ages
555                                     % ranging from 1 to riskage, they are
556                                     % aasigned their current body mass according
557                                     % to growthfunction
558        for i = 2 : riskage
559            growthcurve(i) = growthcurve(i-1) + 0.3*growthrate*(1 - (growthcurve(i-
560    1)/bodymass));
561        end
562
563        [c reprodage] = min(abs(growthcurve-(bodymass*repbodymass))); % age
564                       % of beginning to reproduce when body weight reaches
565                       %   bodymass*repbodymass (slightly smaller than adult)
566
567
568
569        repenergy = birthmass*littersize/bodymass; % a koefficient of investment
570                       % into reproduction
```

```matlab
571                             % used for balancing how much energy an
572                             % individual can invest into
573                             % different reproductive
574                             % parameteres
575
576
577        exthaz = 0.0001; % koefficient affecting the chance of dying of external
578                             % hazards
579        a = 1;
580        b = 1; % exthaz, a and b are used in the Lotka-Volterra equation that
581                 % regulates external hazard pressure
582
583        % INITIAL POPULATION
584        initpop(1, 1:popsize) = 1:popsize; %1.
585        initpop(2, 1:popsize) = randi([1, riskage], 1, popsize); %2.
586        initpop(3, 1:popsize) = ones(1, popsize).*bodymass; %3.
587        initpop(4, 1:popsize) = growthcurve(1, initpop(2, :)); %4.
588        initpop(5, 1:popsize) = birthmass; %5.
589        initpop(6, 1:popsize) = mutrate; %6.
590        initpop(7, 1:popsize) = rrate; %7.
591        initpop(8, 1:popsize) = littersize; %8.
592        initpop(9, 1:popsize) = somdeath; %9.
593        initpop(10, 1:popsize) = reprodage; %10.
594        initpop(11, 1:popsize) = somEnergy; %11.
595
596        %1. individual ID (used in mixed genotype experiments to identify genotype)
597        %2. current age
598        %3. inherited body mass
599        %4. current body mass
600        %5. inherited birth mass
601        %6. inherited mutation rate
602        %7. inherited reproduction rate
603        %8. inherited litter size
604        %9. parameter of somatic death probability function (somdeath) in the aging
605             % function
606        %10. age when beginning to reproduce
607        %11. energy invested in somatic maintenance (explained in METHODS)
608
609        if(newrun) % initial population is created at the beginning of simulation
610             population = initpop;
611        end
612
613        %THE CORE SIMULATION RUN
614        for timeunit = timeun : totaltime
615
616             disp(timeunit);
617
618             % STORAGE MATRICES KEEP TRACK OF POPULATION PARAMETERS THROUGHOUT
619    SIMULATION
620             biomassdyn(timeunit) = sum(population(4, :))/sum(initpop(4, :));
621             popsizedyn(timeunit) = size(population, 2)/size(initpop,2);
622             bodymassevol(timeunit) = mean(population(3, :));
623             birthmassevol(timeunit) = mean(population(5, :));
624             littersizeevol(timeunit) = mean(population(8, :));
625             mutrateevol(timeunit) = mean(population(6, :));
626             rrateevol(timeunit) = mean(population(7, :));
```

```matlab
627            lifespanevol(timeunit) = 1/mean(population(9, :));
628
629
630
631
632            %======================= REPRODUCTION ==========================
633
634            % potreprodpop (potentially reproducing population) collects mature
635     % subpopulation
636            potreprodpop = population(:, population(2, :)-population(10, :)>0);
637
638            % variance is introduced in time between reproductions
639            reprodvars = round(normrnd(rrate, rratevar));
640
641            % reprodpop (reproducing population) collects individuals that are
642            % past their period between reproduction and are due reproducing
643     % (+ some additional variance)
644            reprodpop = potreprodpop(:, rem(potreprodpop(2, :)-potreprodpop(10, :),
645     reprodvars) == 0);
646
647
648            % copies of their parent individual are created as their progeny -
649            % newgen
650            newgen = zeros(size(reprodpop, 1), 1);
651            for i = 1 : size(reprodpop, 2)
652                if(~isempty(reprodpop))
653                    progeny = repmat(reprodpop(1:size(reprodpop, 1), i), 1,
654     round(normrnd(littersize, littervar)));
655                    newgen = [newgen, progeny];
656                end
657            end
658            newgen = newgen(:, 2:end);
659
660            % number of new offspring is collected into a storage matrix
661            births(timeunit) = size(newgen, 2);
662
663            % inherited variance (proportional to parent's mutation rate)
664            % modifies parental parameters producing varying offspring
665            newgen(2, :) = 1;
666            newgen(3, :) = real(newgen(3, :) + (normrnd(0, newgen(6, :)*inhvar).*
667     newgen(3, :)));
668            newgen(4, :) = real(newgen(5, :) + (normrnd(0, newgen(6, :)*inhvar).*
669     newgen(5, :)));
670            newgen(5, :) = real(newgen(5, :) + (normrnd(0, newgen(6, :)*inhvar).*
671     newgen(5, :)));
672            newgen(5, newgen(5, :) > 0.5.*newgen(3, :)) = 0.5.*newgen(3, newgen(5, :) >
673     0.5.*newgen(3, :));
674            newgen(6, :) = real(newgen(6, :) + (normrnd(0, newgen(6, :)*inhvar).*
675     newgen(6, :)));
676            newgen(8, :) = real(newgen(8, :) + (normrnd(0, newgen(6, :)*inhvar).*
677     newgen(8, :)));
678            newgen(7, :) = real(newgen(5, :).*newgen(8, :)./newgen(3,
679     :)./repenergy.*rrate);
680            newgen(10, :) = 0;
```

```matlab
681            newgen(11, :) = real(newgen(11, :) + (normrnd(0, newgen(6, :)*inhvar).*
682     newgen(11, :)));
683
684            % the somatic maintenance (somdeath) parameter of the aging
685            % function is calculated based on the somatic maintenance energy
686            % investment with inherited variance (see METHODS)
687            newgen(9, :) = real((0.00000072523237903965.*(log(newgen(11, :)).^6))...
688                -(0.0000458064654458169.*(log(newgen(11, :)).^5))...
689                +(0.00123267215690707.*(log(newgen(11, :)).^4))...
690                -(0.0183381238349637.*(log(newgen(11, :)).^3))...
691                +(0.162769338153511.*(log(newgen(11, :)).^2))...
692                -(0.863957066277595.*(log(newgen(11, :)).^1))...
693                + 2.46992883606531000000);
694
695            % new offpsring is added to the population
696            population = [population, newgen];
697
698
699
700            %======================= MORTALITY =========================
701
702            %MORTALITY CAUSED BY SOMATIC/PHYSIOLOGICAL FACTORS
703
704            % individual probabilities of dying of somatic causes during this update
705            probsdeath = [];
706
707            % version 1 (standard) = death rates are affected by body mass
708             % (increased somatic risk)
709             % and the performance of the somtic maintenance program in
710             % mitigating somatic risk
711            probsdeath = population(6, :).*(population(4, :)/bodymass)...
712                .*exp(population(2, :).^population(9, :));
713
714
715            % version 2 = somatic cost unrelated
716        % (used when the "body mass" parameter is converted into
717             % a trait that is selected for but does not affect somatic risks)
718     %       probsdeath = population(6, :)...
719     %           .*exp(population(2, :).^population(9, :));
720
721            probsdeath(probsdeath > 1) = 0;
722            probsdeath(probsdeath < 0) = 0;
723
724            % individuals actually dying of somatic causes during this update
725            % based on binomial trials using probsdeath
726            death = [];
727            death = binornd(1, probsdeath(1, :));
728
729
730            % data on mortality of somatic causes is stored in a storage matrix
731            sommortality = [sommortality, population(2, death(1, :) == 1)];
732            %-----------------------
733
734            % dead individuals are eliminated from the population
```

```matlab
735            population(:, death(1, :) == 1) = 0;
736            population = population(:, population(1, :) > 0);
737
738
739
740            % MORTALITY CAUSED BY EXTERNAL HAZARDS (predation, disease, etc)
741            % (the Lotka-Voterra model of predator-prey dynamics was used as a basis)
742
743            % population size-dependent external hazard pressure (exthazard)
744            exthazard = exthaz...
745                +((a*popsizedyn(timeunit)*exthaz) - (b*exthaz));
746
747            % probabilities of dying of external hazards (development of bodymass
748             % or other selected trait reduces chances of dying
749             % of external hazards)
750            extprobs = [];
751            extprobs = exthazard.*(bodymass./population(4, :));
752            extprobs(1, extprobs > 1) = 1;
753            extprobs(1, extprobs < 0) = 0;
754
755            % individuals actually dying of causes related to external hazards
756             % based on binomial trials using extprobs
757            extdeath = [];
758            extdeath = binornd(1, extprobs(1, :));
759
760            % data on mortality caused by external hazards is stored in a storage
761    matrix
762            extmortality = [extmortality, population(2, extdeath(1, :) == 1)];
763            %------------------------
764
765            % dead individuals are eliminated from the population
766            population(:, extdeath(1, :) == 1) = 0;
767            population = population(:, population(1, :) > 0);
768
769
770
771            % MORTALITY IMPOSED BY ECOSYSTEM'S CARRYING CAPACITY
772             % (essentially reflects mortality caused by intra-specific competition)
773
774            % Version 1 = used when maximum biomass is kept stable
775             % (in experiments when body mass evolves)
776             % (development of body mass reduces the chances of dying
777             % in intra-specific competition)
778            overkill = sum(population(4, :))/ sum(initpop(4, :));
779            invs = 1./population(4, :);
780            capprobs = invs/sum(invs);
781            capprobs = capprobs-(mean(capprobs));
782            capprobs = capprobs+(1-(1/overkill));
783
784            % Version 2 = used when population size is kept stable
785             % (in experiments when "body mass" is trasformed
786             % into another selected trait)
787             % (development of body mass or other selected trait
788             % reduces the chances of dying in intra-specific competition)
789    %        overkill = size(population, 2) / size(initpop, 2);
```

```matlab
790  %           invs = 1./population(4, :);
791  %           capprobs = invs/sum(invs);
792  %           capprobs = capprobs-(mean(capprobs));
793  %           capprobs = capprobs+(1-(1/overkill));
794
795            capprobs(capprobs < 0) = 0;
796            capprobs(capprobs > 1) = 1;
797
798            % individuals actually dying in intra-specific competitioN
799             % based on binomial trials using extprobs
800            capdeath = [];
801            capdeath = binornd(1, capprobs(1, :));
802
803
804
805            % data on mortality caused by intra-specific competition
806             % is stored in a storage matrix
807            capmortality = [capmortality, population(2, capdeath(1, :) == 1)];
808
809            % dead individuals are eliminated from the population
810            population(:, capdeath(1, :) == 1) = 0;
811            population = population(:, population(1, :) > 0);
812
813
814
815
816            % ============UPDATING AGE AND BODY MASS DUE TO GROWTH==============
817            population(2, :) = population(2, :) + 1;
818            population(4, :) = population(4, :) + 0.3*growthrate*(1 - (population(4,
819  :)./population(3, :)));
820
821
822
823            % =============ASSIGNING MATURITY AGES FOR THE NEW OFFSPRING========
824            newborns = find(population(10, :) == 0);
825            grownnewborns = find(population(4, :)./population(3, :) >= repbodymass);
826            mature = intersect(newborns, grownnewborns);
827            population(10, mature) = population(2, mature);
828
829
830
831            % ================SAVING VARIABLES INTO FILES======================
832            if(fname > size(filenames, 2))
833                fname = 1;
834                iter = iter+1;
835            end
836            if(rem(timeunit, 15000) == 0)
837                its(1:iter) = 'z';
838                save(['D:\' its filenames(fname) '.mat']);
839            end
840            if(rem(timeunit, 30000) == 0)
841                fname = fname + 1;
842            end
843
```

```matlab
844
845            % =========REMOVAL OF OCCASIONAL NaNs==============================
846            population(:, isnan(sum(population(:, :)))) = 0;
847            population = population(:, population(1, :) > 0);
848
849        end
850
851
852        % =========ENTIRE SIMULATION RUN IS SAVED IN A FILE====================
853        these(1:iteration) = '0';
854        save(['D:\' these 'zzh.mat']);
855    end
856
857    % ======TOTAL SIMULATION TIME MEASURES====================================
858    time = toc;
859    hours = floor(time / 3600);
860    time = time - hours * 3600;
861    mins = floor(time / 60);
862    secs = time - mins * 60;
863    secs = round(secs);
864
865    fprintf('Execution time (HH:MM:SS) - %d:%d:%d        \n\n', hours, mins, secs);
866
```

b. *Competitive model for competition between two genotypes*.

```matlab
868
869    for iteration = 1 : 25
870        disp(iteration);
871
872        newrun = true;
873
874        if(newrun)
875            clearvars -except newrun iteration
876            fname = 1;
877            timeun = 1;
878            iter = 1;
879
880            % OUTPUT STORAGE
881            sommortality = []; % counts of mortality for somatic reasons
882            extmortality = []; % counts of mortality caused by external hazard
883            capmortality = []; % counts of mortality imposed by ecosystem's
884                               %  carrying capacity (intra-specific competition)
885            biomassdyn = [];   % population biomas dynamics over time
886            popsizedyn = [];   % population size dynamics over time
887            births = [];       % counts of new births over time
888
889            fracspec1 = []; % fraction of genotype 1
890
891            % individual parameters for genotype 1
892            bodymassevol1 = []; % population's average bodymass over time
893            birthmassevol1 = []; % population's average birthmass over time
894            littersizeevol1 = []; % population's average litter size over time
895            mutrateevol1 = []; % population's average mutation rate over time
896            rrateevol1 = []; % population's average reproduction rate over time
897            lifespanevol1 = []; % population's average somatic maintenance
```

```matlab
898                                    %  coefficient over time
899
900          % individual parameters for genotype 2
901              bodymassevol2 = []; % population's average bodymass over time
902              birthmassevol2 = []; % population's average birthmass over time
903              littersizeevol2 = []; % population's average litter size over time
904              mutrateevol2 = []; % population's average mutation rate over time
905              rrateevol2 = []; % population's average reproduction rate over time
906              lifespanevol2 = []; % population's average somatic maintenance
907                                    %  coefficient over time
908       else
909           timeun = timeunit + 1;
910           if(fname == size(filenames, 2))
911               fname = 1;
912           else
913               fname = fname + 1;
914           end
915       end
916       filenames = ['a' 'b' 'c' 'd' 'e' 'f' 'g' 'h' 'i' 'j' 'k' 'l' 'm' 'n'...
917           'o' 'p' 'q' 'r' 's' 't' 'u' 'v' 'w' 'x' 'y' 'z'];
918
919     % GENERAL MODEL PARAMETERS:
920      totaltime = 1005000; % total # of simulation updates ("time")
921      popsize = 10000; % initial population size
922      mutrate = 0.000000001;
923      inhvar = 25000000; % a multiplier of mutation rate determining
924                          %  variance in trait inheritance (var=inhvar*mutrate)
925                          %  so that inhertance variance is proportional to
926                          %  mutation rate
927      bodymass = 5000; % initial adult bosymass
928      birthmass = 300; % initial body mass at birth
929
930      repbodymass = 0.9693; % multiplier determining at what body mass
931                            %  as a fraction of the individual's inherited
932                            %  adult body mass the individual begins to
933                            %  reproduce
934      rrate = 600; % initial time (in # simulation updates)
935               %  between successive reproductions
936      littersize = 1; % initial # progeny per reproduction per individual
937      littervar = (0.1*littersize)/littersize; % variance of littersize
938      rratevar = (0.1*rrate)/rrate; % variance of reproduction rate
939      growthrate = 57; % coefficient of body growth rate
940
941      somdeath = 0.34; % an exponential coefficient of the somatic maintenance
942  equation
943      somEnergy = 2231.81365913237; % initial energy invested in somatic
944       %  maintenance when somdeath=0.34
945                                    %  see SUPPLEMENTS
946
947      aging = mutrate*exp([1:1000000].^somdeath); % the aging function -
948                    % probability of dying for somatic reasons over time
949
950      risk = cumsum(aging); % cumulative sum function of the aging curve
951      [c riskage] = min(abs(risk-1)); % riskage - age at which risk = 1;
952                                       %  explained in METHODS
953
```

```matlab
954        %INITIAL BODY GROWTH FUNCTION
955        growthcurve = [birthmass]; % curve for body size distribution in the initial
956    population
957                                     % initial population is generated with ages
958                                     % ranging from 1 to riskage, they are
959                                     % aasigned their current body mass according
960                                     % to growthfunction
961        for i = 2 : riskage
962            growthcurve(i) = growthcurve(i-1) + 0.3*growthrate*(1 - (growthcurve(i-
963    1)/bodymass));
964        end
965
966        [c reprodage] = min(abs(growthcurve-(bodymass*repbodymass))); % age
967                          % of beginning to reproduce when body weight reaches
968                          %   bodymass*repbodymass (slightly smaller than adult)
969
970
971
972        repenergy = birthmass*littersize/bodymass; % a koefficient of investment
973                               % into reproduction
974                               % used for balancing how much energy an
975                               % individual can invest into
976                               % different reproductive
977                               % parameteres
978
979
980        exthaz = 0.0001; % koefficient affecting the chance of dying of external
981                           % hazards
982        a = 1;
983        b = 1; % exthaz, a and b are used in the Lotka-Volterra equation that
984               % regulates external hazard pressure
985
986        %initial population
987        initpop(1, 1:ceil(popsize/2)) = 1; %1 genotype 1
988        initpop(1, ceil(popsize/2)+1:popsize) = 2; %1 genotype 2
989        initpop(2, 1:popsize) = randi([1, maxage], 1, popsize); %2
990        initpop(3, 1:popsize) = ones(1, popsize).*bodymass; %3
991        initpop(4, 1:popsize) = growthcurve(1, initpop(2, :)); %4
992        initpop(5, 1:popsize) = birthmass; %5
993        initpop(6, 1:ceil(popsize/2)) = mutrate/10; %6 genotype 1
994        initpop(6, ceil(popsize/2)+1:popsize) = mutrate; %6 genotype 2
995        initpop(7, 1:popsize) = rrate; %7
996        initpop(8, 1:popsize) = littersize; %8
997        initpop(9, 1:popsize) = somdeath; %9
998        initpop(10, 1:popsize) = reprodage; %10
999        initpop(11, 1:popsize) = somEnergy; %11
1000
1001       %1. individual ID (used in mixed genotype experiments to identify genotype)
1002       %2. current age
1003       %3. inherited body mass
1004       %4. current body mass
1005       %5. inherited birth mass
1006       %6. inherited mutation rate
1007       %7. inherited reproduction rate
1008       %8. inherited litter size
1009       %9. parameter of somatic death probability function (somdeath) in the aging
```

```matlab
1010             % function
1011         %10. age when beginning to reproduce
1012         %11. energy invested in somatic maintenance (explained in METHODS)
1013
1014
1015
1016         if(newrun) % initial population is created at the beginning of simulation
1017             population = initpop;
1018         end
1019
1020
1021         %THE CORE SIMULATION RUN
1022         for timeunit = timeun : totaltime
1023             disp(timeunit);
1024
1025             % STORAGE MATRICES KEEP TRACK OF POPULATION PARAMETERS THROUGHOUT
1026     SIMULATION
1027             biomassdyn(timeunit) = sum(population(4, :))/sum(initpop(4, :));
1028             popsizedyn(timeunit) = size(population, 2)/size(initpop,2);
1029
1030             bodymassevol1(timeunit) = mean(population(3, population(1,:)==1));
1031             birthmassevol1(timeunit) = mean(population(5, population(1,:)==1));
1032             littersizeevol1(timeunit) = mean(population(8, population(1,:)==1));
1033             mutrateevol1(timeunit) = mean(population(6, population(1,:)==1));
1034             rrateevol1(timeunit) = mean(population(7, population(1,:)==1));
1035             lifespanevol1(timeunit) = 1/mean(population(9, population(1,:)==1));
1036
1037             bodymassevol2(timeunit) = mean(population(3, population(1,:)==2));
1038             birthmassevol2(timeunit) = mean(population(5, population(1,:)==2));
1039             littersizeevol2(timeunit) = mean(population(8, population(1,:)==2));
1040             mutrateevol2(timeunit) = mean(population(6, population(1,:)==2));
1041             rrateevol2(timeunit) = mean(population(7, population(1,:)==2));
1042             lifespanevol2(timeunit) = 1/mean(population(9, population(1,:)==2));
1043
1044             fracspec1(timeunit) = numel(population(1, population(1,:) ==
1045     1))/size(population, 2)*100;
1046
1047             %======================== REPRODUCTION =========================
1048
1049             % potreprodpop (potentially reproducing population) collects mature
1050             % subpopulation
1051             potreprodpop = population(:, population(2, :)-population(10, :)>0);
1052
1053             % variance is introduced in time between reproductions
1054             reprodvars = round(normrnd(rrate, rratevar));
1055
1056             % reprodpop (reproducing population) collects individuals that are
1057             % past their period between reproduction and are due reproducing
1058             % (+ some additional variance)
1059             reprodpop = potreprodpop(:, rem(potreprodpop(2, :)-potreprodpop(10, :),
1060     reprodvars) == 0);
1061
1062
1063             % copies of their parent individual are created as their progeny -
```

```matlab
        % newgen
        newgen = zeros(size(reprodpop, 1), 1);
        for i = 1 : size(reprodpop, 2)
            if(~isempty(reprodpop))
                progeny = repmat(reprodpop(1:size(reprodpop, 1), i), 1,
round(normrnd(littersize, littervar)));
                newgen = [newgen, progeny];
            end
        end
        newgen = newgen(:, 2:end);

        % number of new offspring is collected into a storage matrix
        births(timeunit) = size(newgen, 2);

        % inherited variance (proportional to parent's mutation rate)
        % modifies parental parameters producing varying offspring
        newgen(2, :) = 1;
        newgen(3, :) = real(newgen(3, :) + (normrnd(0, newgen(6, :)*inhvar).*
newgen(3, :)));
        newgen(4, :) = real(newgen(5, :) + (normrnd(0, newgen(6, :)*inhvar).*
newgen(5, :)));
        newgen(5, :) = real(newgen(5, :) + (normrnd(0, newgen(6, :)*inhvar).*
newgen(5, :)));
        newgen(5, newgen(5, :) > 0.5.*newgen(3, :)) = 0.5.*newgen(3, newgen(5, :) >
0.5.*newgen(3, :));
        % mutation rates are fixed and differ between two genotypes
        % newgen(6, newgen(1, :) == 1) = real(newgen(6, newgen(1, :) == 1) +
(normrnd(0, newgen(6, newgen(1, :) == 1)*inhvar).* newgen(6, newgen(1, :) == 1)));
        newgen(8, :) = real(newgen(8, :) + (normrnd(0, newgen(6, :)*inhvar).*
newgen(8, :)));
        newgen(7, :) = real(newgen(5, :).*newgen(8, :)./newgen(3,
:)./repenergy.*rrate);
        newgen(10, :) = 0;
        newgen(11, :) = real(newgen(11, :) + (normrnd(0, newgen(6, :)*inhvar).*
newgen(11, :)));

    % the somatic maintenance (somdeath) parameter of the aging
     % function is calculated based on the somatic maintenance energy
     % investment with inherited variance (see METHODS)
     newgen(9, :) = real((0.00000072523237903965.*(log(newgen(11, :)).^6))...
         -(0.0000458064654458169.*(log(newgen(11, :)).^5))...
         +(0.00123267215690707.*(log(newgen(11, :)).^4))...
         -(0.0183381238349637.*(log(newgen(11, :)).^3))...
         +(0.162769338153511.*(log(newgen(11, :)).^2))...
         -(0.863957066277595.*(log(newgen(11, :)).^1))...
         + 2.46992883606531000000);

     % new offpsring is added to the population
     population = [population, newgen];

        %========================= MORTALITY =========================

        %MORTALITY CAUSED BY SOMATIC/PHYSIOLOGICAL FACTORS
```

```matlab
1119
1120            % individual probabilities of dying of somatic causes during this update
1121            probsdeath = [];
1122
1123            % version 1 (standard) = death rates are affected by body mass
1124             % (increased somatic risk)
1125             % and the performance of the somtic maintenance program in
1126             % mitigating somatic risk
1127            probsdeath = population(6, :).*(population(4, :)/bodymass)...
1128                .*exp(population(2, :).^population(9, :));
1129
1130
1131            % version 2 = somatic cost unrelated
1132        % (used when the "body mass" parameter is converted into
1133             % a trait that is selected for but does not affect somatic risks)
1134    %       probsdeath = population(6, :)...
1135    %           .*exp(population(2, :).^population(9, :));
1136
1137            probsdeath(probsdeath > 1) = 0;
1138            probsdeath(probsdeath < 0) = 0;
1139
1140            % individuals actually dying of somatic causes during this update
1141            % based on binomial trials using probsdeath
1142            death = [];
1143            death = binornd(1, probsdeath(1, :));
1144
1145
1146            % data on mortality of somatic causes is stored in a storage matrix
1147            sommortality = [sommortality, population(2, death(1, :) == 1)];
1148            %------------------------
1149
1150            % dead individuals are eliminated from the population
1151            population(:, death(1, :) == 1) = 0;
1152            population = population(:, population(1, :) > 0);
1153
1154
1155
1156            % MORTALITY CAUSED BY EXTERNAL HAZARDS (predation, disease, etc)
1157            % (the Lotka-Voterra model of predator-prey dynamics was used as a basis)
1158
1159            % population size-dependent external hazard pressure (exthazard)
1160            exthazard = exthaz...
1161                +((a*popsizedyn(timeunit)*exthaz) - (b*exthaz));
1162
1163            % probabilities of dying of external hazards (development of bodymass
1164             % or other selected trait reduces chances of dying
1165             % of external hazards)
1166            extprobs = [];
1167            extprobs = exthazard.*(bodymass./population(4, :));
1168            extprobs(1, extprobs > 1) = 1;
1169            extprobs(1, extprobs < 0) = 0;
1170
1171            % individuals actually dying of causes related to external hazards
1172             % based on binomial trials using extprobs
```

```matlab
1173             extdeath = [];
1174             extdeath = binornd(1, extprobs(1, :));
1175
1176             % data on mortality caused by external hazards is stored in a storage
1177   matrix
1178             extmortality = [extmortality, population(2, extdeath(1, :) == 1)];
1179             %------------------------
1180
1181             % dead individuals are eliminated from the population
1182             population(:, extdeath(1, :) == 1) = 0;
1183             population = population(:, population(1, :) > 0);
1184
1185
1186
1187             % MORTALITY IMPOSED BY ECOSYSTEM'S CARRYING CAPACITY
1188              % (essentially reflects mortality caused by intra-specific competition)
1189
1190             % Version 1 = used when maximum biomass is kept stable
1191              % (in experiments when body mass evolves)
1192              % (development of body mass reduces the chances of dying
1193              % in intra-specific competition)
1194             overkill = sum(population(4, :))/ sum(initpop(4, :));
1195             invs = 1./population(4, :);
1196             capprobs = invs/sum(invs);
1197             capprobs = capprobs-(mean(capprobs));
1198             capprobs = capprobs+(1-(1/overkill));
1199
1200             % Version 2 = used when population size is kept stable
1201              % (in experiments when "body mass" is trasformed
1202              % into another selected trait)
1203              % (development of body mass or other selected trait
1204              % reduces the chances of dying in intra-specific competition)
1205   %          overkill = size(population, 2) / size(initpop, 2);
1206   %          invs = 1./population(4, :);
1207   %          capprobs = invs/sum(invs);
1208   %          capprobs = capprobs-(mean(capprobs));
1209   %          capprobs = capprobs+(1-(1/overkill));
1210
1211             capprobs(capprobs < 0) = 0;
1212             capprobs(capprobs > 1) = 1;
1213
1214             % individuals actually dying in intra-specific competitioN
1215              % based on binomial trials using extprobs
1216             capdeath = [];
1217             capdeath = binornd(1, capprobs(1, :));
1218
1219
1220
1221             % data on mortality caused by intra-specific competition
1222              % is stored in a storage matrix
1223             capmortality = [capmortality, population(2, capdeath(1, :) == 1)];
1224
1225             % dead individuals are eliminated from the population
1226             population(:, capdeath(1, :) == 1) = 0;
1227             population = population(:, population(1, :) > 0);
```

```matlab
1228
1229
1230
1231
1232            % ============UPDATING AGE AND BODY MASS DUE TO GROWTH==============
1233            population(2, :) = population(2, :) + 1;
1234            population(4, :) = population(4, :) + 0.3*growthrate*(1 - (population(4,
1235    :)./population(3, :)));
1236
1237
1238
1239            % =============ASSIGNING MATURITY AGES FOR THE NEW OFFSPRING========
1240            newborns = find(population(10, :) == 0);
1241            grownnewborns = find(population(4, :)./population(3, :) >= repbodymass);
1242            mature = intersect(newborns, grownnewborns);
1243            population(10, mature) = population(2, mature);
1244
1245
1246
1247            % ================SAVING VARIABLES INTO FILES=====================
1248            if(fname > size(filenames, 2))
1249                fname = 1;
1250                iter = iter+1;
1251            end
1252            if(rem(timeunit, 15000) == 0)
1253                its(1:iter) = 'z';
1254                save(['D:\' its filenames(fname) '.mat']);
1255            end
1256            if(rem(timeunit, 30000) == 0)
1257                fname = fname + 1;
1258            end
1259
1260
1261            % =========REMOVAL OF OCCASIONAL NaNs==============================
1262            population(:, isnan(sum(population(:, :)))) = 0;
1263            population = population(:, population(1, :) > 0);
1264
1265        end
1266
1267
1268        % =========ENTIRE SIMULATION RUN IS SAVED IN A FILE====================
1269        these(1:iteration) = '0';
1270        save(['D:\' these 'zzh.mat']);
1271    end
1272
1273    % ======TOTAL SIMULATION TIME MEASURES====================================
1274    time = toc;
1275    hours = floor(time / 3600);
1276    time = time - hours * 3600;
1277    mins = floor(time / 60);
1278    secs = time - mins * 60;
1279    secs = round(secs);
1280
1281    fprintf('Execution time (HH:MM:SS) - %d:%d:%d         \n\n', hours, mins, secs);
```

1282

1283

1284

1285 **Section 2. Evolution of reproductive traits under fixed adult body mass.** As shown in Fig. S1,

1286 the early simulation period is linked with rapid evolution of reproduction rate and body mass at

1287 birth, which is likely to have caused positive selection for gMR shown in Fig. 2B. Litter size,

1288 however, in our simulations did not show any consistent evolution under this condition.

1289

1290

1291

1292 Fig. S1. Evolution of reproductive parameters in simulations with fixed adult body mass.

1293

1294   **Section 3. All-cause age-dependent mortality in the model.** The model recapitulates a typical

1295   age-dependent mortality chart for wild animals (Fig. S2). Early life is accompanied with the

1296   very high mortality rates which drop until maturity. Fig. S3 demonstrates natural log data.

1297

1298

1299   Fig. S2. Total mortality by age in absolute numbers.

1300

1301

1302

1303 Fig. S3. Total mortality by age in log-absolute numbers.

1304

1305

Section 4. The aging curve calculations. In order to model inherited variation of SMP strength, we needed a method of linearly varying SMP (e.g. +1%, -5% etc). Since the *Som* parameter in Eq. 1 is in a complex non-linear relationship with the resulting aging curve, this parameter is not suitable for such manipulation. We therefore reasoned that the best representation of the efficiency of SMP is using the area under the physiological mortality curve as a measure of the general efficiency of SMP over lifetime. Eq. 1 generates the probability D_A of dying of physiological causes <u>at</u> age A. Its cumulative probability function generates probability *D(A)* of dying <u>by</u> age A. *D(A)* thus is directly related with longevity (like the human mortality curve). However, the *D(A)* function decelerates as the cumulative probability of dying approaches 1 (it can be seen in the human mortality curve in Fig. 1B and is a general property of cumulative probability functions). In order to avoid these effects, we did not use the area under the *D(A)* function as a measure of SMP strength, but instead we applied the area under the cumulative sum function of the D_A probability, as shown in the figure below, starting from the simulated age 1 and until the age at which this function reaches 1. In Fig. S4, the green curve represents extended longevity compared to the blue curve, since the sum of its probabilities of dying accumulates more slowly (slower aging). As a result, the area under the green curve is larger, corresponding to a stronger SMP program. In order to model inherited variation in SMP, we used this area as a representation of the SMP strength. The area was stochastically varied from generation to generation as explained in Methods, and its new value in progeny was used to calculate the *Som* parameter for Eq. 1 (determines the probability of dying <u>at</u> age A). The calculation was based on the observation that the area shown in Fig. S4 demonstrates a strong non-linear log-log relationship with the *Som* parameter (polynomial regression of the 6th order; $R^2 > 0.99999$) as shown in Fig. S5.

1329

1330   Fig. S4. Area under cumulative sum function of Eq. 1 as a measure of the relative

1331   efficiency of SMP.

1332

1333

1334   Fig. S5. Relationship between log-area in Fig. S4 (log-efficiency of SMP) and the *Som*

1335   parameter in Eq. 1 (probability of dying <u>at</u> age A).

1336

1337

1338

1339

1340

1341

1342

1343   **Section 5. Removal of outliers.** Occasionally the model demonstrated unnatural "spikes" in the

1344   evolution of some traits under some conditions. We had to apply the following code to remove

1345   them:

```
1346    input = aaccbmbllsev_lit(22,:); % a certain problematic model run
1347
1348    threshold = 0.15; % arbitrary value
1349
1350    for row = 1 : size(input, 1)
1351        for col = 2 : size(input, 2)
1352            if input(row, col-1)/input(row, col) > 1+threshold ||...
1353                    input(row, col-1)/input(row, col) < 1-threshold
1354                input(row, col) = input(row, col-1);
1355            end
1356        end
1357    end
1358
```

1359   The illustration below demonstrates an example (from a standard condition run) of the result:

1360

1361

1362   The parameter "threshhold" required manual alteration until the spike was cleaned by the code

1363   above. Such spikes were visibly outstanding from the normal trend, so that the trend in the

1364   evolving trait continued after the spike with values similar to those immediately preceding the

1365   spike, indicating that the spikes were some artifacts that neither related to nor influenced the

1366   modeled trait evolution. We were not able to determine the source of such spikes

1367

1368

1369

1370

1371

1372

References

1373   1.   N. A. Heim, M. L. Knope, E. K. Schaal, S. C. Wang, J. L. Payne, Animal evolution. Cope's rule in the evolution of marine animals. *Science* **347**, 867-870 (2015).

2.   J. Baker, A. Meade, M. Pagel, C. Venditti, Adaptive evolution toward larger size in mammals. *Proc Natl Acad Sci U S A* **112**, 5093-5098 (2015).

3.   M. Kimura, On the evolutionary adjustment of spontaneous mutation rates. *Genetical Research* **9**, 23-34 (1967).

4.   C. F. Baer, M. M. Miyamoto, D. R. Denver, Mutation rate variation in multicellular eukaryotes: causes and consequences. *Nat Rev Genet* **8**, 619-631 (2007).

5.   K. J. Dawson, The dynamics of infinitesimally rare alleles, applied to the evolution of mutation rates and the expression of deleterious mutations. *Theor Popul Biol* **55**, 1-22 (1999).

6.   M. Lynch, Evolution of the mutation rate. *Trends Genet* **26**, 345-352 (2010).

7.   M. Lynch, The lower bound to the evolution of mutation rates. *Genome Biol Evol* **3**, 1107-1118 (2011).

8.   M. Lynch *et al.*, Genetic drift, selection and the evolution of the mutation rate. *Nat Rev Genet* **17**, 704-714 (2016).

9.   J. Pothof *et al.*, Identification of genes that protect the C. elegans genome against mutations by genome-wide RNAi. *Genes Dev* **17**, 443-448 (2003).

10.   E. Marcon, P. B. Moens, The evolution of meiosis: recruitment and modification of somatic DNA-repair proteins. *Bioessays* **27**, 795-808 (2005).

11.   D. Galetzka *et al.*, Expression of somatic DNA repair genes in human testes. *J Cell Biochem* **100**, 1232-1239 (2007).

12.   D. Hanahan, R. A. Weinberg, The hallmarks of cancer. *Cell* **100**, 57-70 (2000).

13.   A. F. Caulin, C. C. Maley, Peto's Paradox: evolution's prescription for cancer prevention. *Trends Ecol Evol* **26**, 175-182 (2011).

14.   C. Lopez-Otin, M. A. Blasco, L. Partridge, M. Serrano, G. Kroemer, The hallmarks of aging. *Cell* **153**, 1194-1217 (2013).

15.   S. J. Pfau, R. E. Silberman, K. A. Knouse, A. Amon, Aneuploidy impairs hematopoietic stem cell fitness and is selected against in regenerating tissues in vivo. *Genes Dev* **30**, 1395-1408 (2016).

16.   C. J. Sherr, Principles of tumor suppression. *Cell* **116**, 235-246 (2004).

17.   D. Glick, S. Barth, K. F. Macleod, Autophagy: cellular and molecular mechanisms. *J Pathol* **221**, 3-12 (2010).

18.   A. I. Rozhok, J. DeGregori, The evolution of lifespan and age-dependent cancer risk. *Trends Cancer* **2**, 552-560 (2016).

19.   J. B. Swann, M. J. Smyth, Immune surveillance of tumors. *J Clin Invest* **117**, 1137-1146 (2007).

20.   E. C. Cox, T. C. Gibson, Selection for high mutation rates in chemostats. *Genetics* **77**, 169-184 (1974).

21.   T. C. Gibson, M. L. Scheppe, E. C. Cox, Fitness of an Escherichia coli mutator gene. *Science* **169**, 686-688 (1970).

22.   P. D. Sniegowski, P. J. Gerrish, R. E. Lenski, Evolution of high mutation rates in experimental populations of E. coli. *Nature* **387**, 703-705 (1997).

23.   K. Harris, Evidence for recent, population-specific evolution of the human mutation rate. *Proc Natl Acad Sci U S A* **112**, 3439-3444 (2015).

24.   D. F. Conrad *et al.*, Variation in genome-wide mutation rates within and between human families. *Nat Genet* **43**, 712-714 (2011).

25.   K. Harris, J. K. Pritchard, Rapid evolution of the human mutation spectrum. *Elife* **6**, (2017).

26.   J. Draghi, G. P. Wagner, Evolution of evolvability in a developmental model. *Evolution* **62**, 301-315 (2008).

27.   M. J. Benton, P. N. Pearson, Speciation in the fossil record. *Trends Ecol Evol* **16**, 405-411 (2001).

28. N. Eldredge, S. J. Gould, in *Models in Paleobiology,* T. J. M. Schopf, Ed. (Freeman Cooper, San Francisco,
1972), pp. 82-115.
29. S. J. Gould, N. Eldredge, Punctuated equilibrium comes of age. *Nature* **366**, 223-227 (1993).
30. C. Venditti, A. Meade, M. Pagel, Multiple routes to mammalian diversity. *Nature* **479**, 393-396 (2011).
31. E. Loh, J. J. Salk, L. A. Loeb, Optimization of DNA polymerase mutation rates during bacterial evolution.
Proc Natl Acad Sci U S A **107**, 1154-1159 (2010).
32. A. G. Jones, R. Burger, S. J. Arnold, Epistasis and natural selection shape the mutational architecture of
complex traits. *Nat Commun* **5**, 3709 (2014).
33. A. G. Jones, S. J. Arnold, R. Burger, The mutation matrix and the evolution of evolvability. *Evolution* **61**,
727-745 (2007).
34. A. J. Lotka, *Elements of Physical Biology.* (Williams and Wilkins, Baltimore, 1925).
35. V. Volterra, Variazioni e fluttuazioni del numero d'individui in specie animali conviventi. *Mem Acad*
Lincei Roma **2**, 31-113 (1926).
36. M. E. Hochberg, R. J. Noble, A framework for how environment contributes to cancer risk. *Ecol Lett* **20**,
117-134 (2017).
37. T. Madsen *et al.*, Cancer Prevalence and Etiology in Wild and Captive Animals. *Ecology and Evolution of*
Cancer, 11 (2017).
38. A. I. Rozhok, J. DeGregori, Toward an evolutionary model of cancer: Considering the mechanisms that
govern the fate of somatic mutations. *Proc Natl Acad Sci U S A* **112**, 8914-8921 (2015).
39. S. R. Kennedy, L. A. Loeb, A. J. Herr, Somatic mutations in aging, cancer and neurodegeneration. *Mech*
Ageing Dev **133**, 118-126 (2012).
40. B. A. Finette *et al.*, Determination of hprt mutant frequencies in T-lymphocytes from a healthy pediatric
population: statistical comparison between newborn, children and adult mutant frequencies, cloning
efficiency and age. *Mutat Res* **308**, 223-231 (1994).
41. H. Giese *et al.*, Age-related mutation accumulation at a lacZ reporter locus in normal and tumor tissues
of Trp53-deficient mice. *Mutat Res* **514**, 153-163 (2002).
42. S. Horvath, DNA methylation age of human tissues and cell types. *Genome Biol* **14**, R115 (2013).

1447

1448 **Acknowledgments**

1449 We would like to thank Mark Johnston, L. Alex Liggett and David Pollock of the University

1450 of Colorado, and Robert Gatenby and Andriy Marusyk of the Moffitt Cancer Center for critical

1451 review of the manuscript. These studies were supported by National Cancer Institute grant

1452 R01CA180175 to J.D.

1453

1454

1455

 Figures and Legends

1457

Fig. 1. A scheme of the model simulations. (A) Stages of an individual simulated lifespan. (B) At each timepoint during the simulation the modeled population undergoes 5 main updates: 1. Individuals that have not reached maturity increase their body mass following their growth curve, starting from the initial birth mass and up until they reach their inherited body mass (parent body mass with variation proportional to parent mutation rate); mature individuals remain at the same body mass. 2. Each individual past maturation reproduces with a certain inherited frequency of reproduction, producing on average an inherited number of progeny per litter, each progeny's birth body mass is inherited from parent with variation proportional to parent's mutation rate. Each

individual is tried in a binomial trial with a small probability (at each timepoint) of dying of three main causes: 3.

Death following limitations imposed by ecosystem carrying capacity which allows for a certain maximum

population size and promotes intra-specific and inter-specific competition for resources if population numbers

exceed this capacity. 4. Death caused by predation is modelled based on the Lotka-Volterra model of predator-

prey interaction (*34, 35*). 5. Death caused by physiological aging, such as due to cancer, frailty or other age-

related causes; the probability is negilgible early in life but increases exponentially with age; the speed of increase

of the probability of death caused by aging depends on an individual's aging profile which is determined by the

aging curve as explained in **Fig. 2A**, "Theoretical introduction to the modeling" subsection of **Results** and "The

somatic maintenance program paradigm" subsection of **Methods.**

1474

1475

1476

Fig. 2. The effect of SMP evolution on the evolution of body mass and mutation rate. (A) physiological/aging related mortality curves generated based on the cumulative distribution function of D_A (Eq. 1). Colors represent the effect of the *Som* (SMP) parameter (Eq. 1). Dotted lines were generated by elevating mutation rate 2-fold. (B) modern human mortality in the U.S.A (https://www.ssa.gov). (C-E) evolution of life history traits under positive selection for body size. (F,G) population size dynamics when SMP can evolve (corresponds to green in *C-E*) or SMP evolution is blocked (blue in *C-E*); colors indicate individual populations. (H) relative frequency of Species B (SMP evolution blocked, blue in *C-E*) in a mixed population with Species A (SMP can evolve, green in *C-E*). For

 (*C*), (*D*), (*E*) and (*H*) (and similar graphs in other figures), 25 simulations are combined, with the dark line

1485 reflecting the mean and shaded area denoting the 95% confidence intervals.

1486

Fig. 3. Evolution of body mass, gMR and SMP under various regimens of selection. Separate experiments are stacked as indicated in their subtitles. The layout: left – body size, middle – gMR, right – SMP (the *Som*

1489   parameter in Eq. 1) is maintained as in Fig. 2C-E. Green – the standard condition (as green in Fig. 2C-E); blue –

1490   alternative conditions with fixed values of a trait (blue horizontal line in *A,D,F*), when gMR and sMR are dislinked

1491   so that the somatic cost is fixed while gMR can evolve (blue in G-I), and under selection for a somatic risk

1492   unrelated trait (blue in *J-L*).

1493

Fig. 4. The evolution of gMR in in the absence of positive selection for body mass and SMP. The SMP's

Som parameter was fixed at 0.34 (red), 0.24 (green; enhanced 10X) and 0.2 (blue; enhanced 40X); a linear

decrease in the *Som* value results in a substantially improved SMP, so that the green SMP is ~10X more efficient

compared to red, and the blue is a ~4X more efficient SMP than the green. The standard (red) SMP leads to a

significantly stronger selection for lower gMR (non-overlapping 95% CIs); however, the absence of difference

between the 10X (green) and 40X (blue) improved SMPs indicates that overly improved SMPs might not provide

any further difference for how selection acts on gMR.

1501

1502

Fig. 5. Positive selection for mutators. (**A**) frequency of a mutator phenotype in a mixed competitive population with "wild-type" species. Red (1.4X), orange (2X) and green (10X) are mutators of different fold increase in MR relative to the competitor as indicated by the respective numbers. (**B**) positive selection for a somatic cost neutral trait demonstrates faster evolution (and so adaptation) of mutators. Colors and MR fold increase as in (*A*).

1509