Title: EnsembleNet: ensemble gene function predictions for *Arabidopsis thaliana*

Authors: Bjoern Oest Hansen[1,2*], Etienne H. Meyer[1*], Camilla Ferrari[1], Neha Vaid[1], Sara

Movahedi[3,4], Klaas Vandepoele[3], Zoran Nikoloski[1,5], Marek Mutwil[1,6#]

[1]Max Planck Institute of Molecular Plant Physiology, Am Muehlenberg 1, 14476 Potsdam,

Germany

[2]Institut für Medizinische Informatik, Universitätsmedizin Göttingen, Robert-Koch-Str. 40

37075 Göttingen, Germany

[3]VIB Center for Plant Systems Biology, Department of Plant Biotechnology and

Bioinformatics Ghent University, Technologiepark 927, B-9052 Gent, Belgium

[4]Rijk Zwaan Breeding B.V., Burgemeester Crezéelaan 40, PO Box 40, 2678 ZG De Lier, The

Netherlands

[5]Bioinformatics Group, Institute of Biochemistry and Biology, University of Potsdam, Karl-

Liebknecht-Str. 24-25, 14476 Potsdam-Golm, Germany

[6]School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive,

Singapore 637551, Singapore

#Corresponding author:

Marek Mutwil

Max Planck Institute of Molecular Plant Physiology,

Am Muehlenberg 1,

14476 Potsdam,

Germany

Email: Mutwil@mpimp-golm.mpg.de

*Equal contribution

None of the authors have any competing interests.

Abstract:

Despite increasing availability of sequenced genomes, accurate characterization of gene

functions is needed to close the genotype-phenotype gap. Recent advances in gene function

prediction rely on ensemble approaches that integrate the results from multiple inference

methods to produce superior predictions. Yet, these developments remain largely unexplored

in plants. We present Neighbor Counting Ensemble, a gene function prediction method which

integrates eleven gene co-function networks for *Arabidopsis thaliana,* and produces more

accurate gene function predictions for a larger fraction of genes with unknown function. We

used these predictions to identify genes involved in mitochondrial complex I formation, and

for five of them we confirmed the predictions experimentally. The ensemble predictions are

provided as a user-friendly online database, EnsembleNet, available at

http://www.gene2function.de/ensemblenet.html.

Keywords: gene function prediction, ensemble prediction, co-function network, complex I,

Arabidopsis thaliana

Background:

Extracting useful knowledge from genomes is dependent on our ability to correctly assign

biological functions to gene products. However, since proteins may have more than one

function that are carried out under different condition (e.g. developmental stage,

environment), experimental characterization of protein function is resource-intensive.

Therefore, the challenge posed by the sheer number of uncharacterized gene products has

rendered gene function prediction as one of the most pressing bioinformatics research efforts

([1,2], reviewed in [3]). Apart from providing an overview of the functional repertoire of an

organism, gene function predictions in plants are invaluable for experimentalist, as they

provide the basis for a rapid reverse genetic approach to identify relevant genes. For example,

several studies used *in silico* predictions to identify genes involved in plant viability [4], seed

germination [5], shade avoidance [6], cyclic electron flow [7], cell division [8], drought

sensitivity and lateral root development [9].

Current protein function prediction methods largely rely on the Gene Ontology [10] to

communicate and describe functional annotation, and to describe relationships between those

annotations. The Gene Ontology (GO) uses three functional domains to specify gene

function: Cellular Component (CC, subcellular component/protein complex where the gene

product acts), Molecular Function (MF, biochemical activity of the gene product), and

Biological Process (BP, purpose of the gene product). For example, *At1g72370*, a 40S

ribosomal protein SA [11], is an "mRNA binding" protein (GO:0003729, MF), which is part

of "cytosolic small ribosomal subunit" (GO:0022627, CC), and is involved in "mature

ribosome assembly" (GO:0042256, BP, www.arabidopsis.org). Importantly, GO also

provides evidence codes useful in specifying if an annotation is based on an experiment (e.g.

IEP: inferred from expression pattern) or a prediction (e.g. ISA: inferred from sequence

alignment), which is needed to construct a "gold standard" dataset of experimentally verified

genes, used in turn for propagating function from characterized to uncharacterized genes

([1,2], reviewed in [3]).

Gene function prediction, regardless of the data used, requires three components: (i) a

dataset that specifies features of the analyzed genes, (ii) a method to identify functionally

related genes in the given dataset and (iii) a method to transfer functional annotation from the

gold standard to uncharacterized genes ([1,2], reviewed in [3]). For example, sequence

homology-based function predictions (e.g. InterProScan, [12]) use (i) a database of

functionally annotated protein sequences (the gold standard), to (ii) find functionally related

proteins that are identified via BLAST, with the aim to (iii) transfer functional information

from a similar gold standard protein to an uncharacterized protein. The gene function

prediction methods can be divided into three "generations": (i) single, (ii) integrative and (iii)

ensemble (reviewed in [13]). However, regardless of the prediction method, all approaches

are based on the guilt-by-association (GBA) principle, whereby genes with similar features

are likely to have same function ([14], reviewed in [3]).

First generation methods use a single type of data to identify functionally related

genes (reviewed in [3]). Genomic GBA approaches use sequence similarity (e.g.

InterProScan, [12]), genomic co-occurrence (i.e. gene families that are co-gained or co-lost

across species have same function, [15]), genomic neighbor (i.e. proximity on chromosome

implies same function, [16]), gene fusion (i.e. two separate gene products that are found as

one protein in a species, are interacting, [17]) and domain co-occurrence (i.e. protein domains

that frequently co-occur on proteins are functionally related, [18]) to identify functionally

related genes. Transcriptomic GBA approaches utilize co-response to perturbation and co-

expression to predict gene function. For instance, transcripts that are up-regulated upon cold

stress may be involved in cold adaptation [19] and functionally related genes tend to show

similar expression profiles across treatments [20]. Similarly, protein-protein interaction GBA

states that interaction of two proteins generally implicates that they are functionally related

(e.g. subunits of a ribosome, [21]).

Second generation methods integrate different data types, thereby addressing

important shortcomings of the first generation methods ([22], reviewed in [3]). These

shortcomings include limited coverage (e.g. the Arabidopsis ATH1 microarray captures only

60% of protein coding transcripts [23], or <10% of the predicted Arabidopsis interactome has

been elucidated, [24]), technical artifacts (e.g. bimolecular fluorescence complementation

(BiFC) interaction studies produce false positive results due to protein over-expression, [25])

and technical limitation (e.g. yeast-two-hybrid (Y2H) only works on proteins that can be

imported into the nucleus). Integration of different datasets can result in increased coverage

(e.g. combining BiFC and Y2H can increase the size of inferred interactome) and confidence

(e.g. protein interactions reported by both BiFC and Y2H are less likely to be false positives).

Indeed, through Bayesian integration of protein-protein interactions, literature text mining,

co-expression, genomic neighbor, gene function and domain co-occurrence, Lee and

colleagues constructed a co-function network with prediction accuracy higher than any of the

single methods [9]. Furthermore, since different subset of biological information is captured

by each method, integrating the diverse biological data can result in a more complete

functional description [26].

Third generation ensemble (also called community) methods produce a prediction by

integrating the outputs of multiple first and second generation methods. The concept is

similar to the "Ask the audience" lifeline (91% success rate) in the Who Wants to Be a

Millionaire television quiz, which performs better than "phone a friend" (65% success rate)

lifeline ("The Wisdom of Crowds", James Surowiecki, 2004). Indeed, integration of 29 gene

regulatory network inference methods in yeast and *E. coli* generated an ensemble prediction

that outperformed all of the individual methods [27]. The ensemble from as few as three input

networks emphasized the advantages and diminished the limitations of each of the 29

methods.

In plants, ensemble approaches are largely unexplored. SUBA3 provides ensemble

sub-cellular predictions based on 22 prediction programs [28]. Furthermore, by integrating

four different methods, a recent study predicted an ensemble gene regulatory network in

Arabidopsis [29]. To our knowledge, ensemble gene function predictions are not available for

Arabidopsis thaliana or other plants, which we set to remedy in this study.

RESULTS

Benchmarking of eleven gene co-function networks

In this study, we obtained and standardized eleven gene co-function networks (see Methods). AraNet v2, GeneMANIA and STRING are second generation methods integrating multiple data sources (Table 1). The remaining eight networks are obtained from first-generation methods based on co-expression (ATTED-II, CC-AFM, PlaNet-microarrays, PlaNet-RNAseq, PODC) or protein-protein interactions (BIOGRID, CorNet, TAIR-PPI). To determine and evaluate the predictions, we have obtained the gold standard Gene Ontology annotations, based on experimental evidence for 14,032 *Arabidopsis thaliana* genes (www.arabidopsis.org, [30]). Analysis of the gold standard revealed that 31% of the 28,392 Arabidopsis genes in BP domain have been experimentally characterized (Figure S1A), while less than 12% of genes have all three GO domains experimentally characterized (Figure S1B).

To predict gene function, we used the Neighbor Counting (NC) approach as the simplest method for network-based function prediction [31]. NC predicts function of a gene by identifying the most frequent function(s) found in the network neighborhood of the gene (Figure 1A). Furthermore, since all networks (apart from PPI networks) contain edge weights that can be scaled to range from 0 (weakest association between genes) to 1 (strongest association), a network can perform differently at a specific edge cut-off. The performance of a network at a given edge cut-off is specified by the performance of the subnetwork obtained upon removal of all edges with weight smaller than the specified cut-off. Therefore, too low an edge cut-off could connect functionally unrelated genes, resulting in an incorrect prediction (Figure 1A, first network), or partially correct prediction (Figure 1A, second network). Conversely, too high a cut-off could disconnect functionally related genes and thus hinder any prediction (Figure 1A, fourth network).

To provide a fair evaluation and comparison of the networks, we first set to find an

edge threshold where a given network produces the highest number of correct predictions. To

this end, we used the F-measure which represents the harmonic mean between precision (i.e.

average fraction of correctly predicted terms) and recall (i.e. average fraction of correctly

predicted terms over all relevant terms, [1]), as a measure of performance for prediction

methods in Critical Assessment of Functional Annotation [1]. We applied a leave-one-out

cross validation, where the function of an experimentally characterized gene is compared to

the Neighbor Counting-predicted function of the gene, treating its function as unknown. The

F-measure can range from 0 (worst value) to 1 (best value) and provides a number that can be

used to compare the performance of the networks (Table S1A).

We illustrate this type of analysis with the STRING network, for which we evaluated

the performance for the BP domain at a given edge cut-off. Not surprisingly, as a

consequence of genes becoming disconnected in the network, the number of made

predictions dropped as the edge cut-off increased (Figure 1B, red line, right y-axis). The F-

measure peaked at 0.88 edge cut-off, identifying the F_{max} value where STRING produces

highest prediction accuracy for BP domain (Figure 1B, blue line, left y-axis). Interestingly,

F_{max} for the BP, CC and MF domain were found at 0.88, 0.6 and 0.65, respectively,

suggesting that one edge cut-off may not be suitable to produce most accurate predictions for

the three domains (Table S1B).

To compare the 11 co-function networks, we plotted the F_{max} values and the number

of made predictions (Figure 1C). As expected from networks integrating multiple data

sources, AraNet v2, GeneMANIA and STRING showed higher F_{max} values than co-

expression and PPI-based methods (Figure 1C, gray bars, left y-axis), but the two co-

expression networks with highest F_{max} (ATTED-II and PlaNet) showed higher number of

made predictions (Figure 1C, red bars, right y-axis).

Generation and benchmarking of ensemble gene function predictions

To construct community predictions from the 11 co-function networks, we explored two different approaches to integrate these networks: the first approach, termed Neighbor Counting Ensemble (NCE) expands the NC rule to multiple networks. More precisely, NCE (i) uses Neighbor Counting to predict function of a gene in each of the input networks and (ii) identifies the most frequent term found by NC in the previous step (ensemble majority vote, Figure 2A). To investigate which combination of the co-function networks produces F_{max} for NCE, we tested all combinations of 2 to 11 networks at their F_{max} threshold, resulting in 2,047 tested combinations for each GO domain (Table S2).

The second approach, based on STATIS [32], termed STATISnet, determines the similarity between each pair of the scaled network adjacency matrices, by calculating Rv coefficient of the adjacency matrices. The principal eigenvector is then used to build a weighted sum of the scaled adjacency matrices, which yields the adjacency matrix of STATISnet (Figure 2B). The F_{max} for STATISnet was then calculated as for the 11 input networks (Table S1).

The analysis revealed that NCE produced highest F_{max} and number of made predictions (prediction coverage) than the best single method (GeneMANIA), and STATISnet for all three GO domains (Figure 2C, Table S2). While F_{max} of STATISnet was lower than F_{max} of GeneMANIA, the prediction coverage of the former was modestly higher for BP and CC domains (Figure 2C). Interestingly, NCE based on all 11 networks produced a lower value for the F-measure than NCE based on GeneMANIA, AraNet v2 and ATTED-II, indicating that the combination of selected networks may have a negative effect on the prediction performance (Table S2). Furthermore, while F_{max} NCE for BP and CC domains used GeneMANIA, AraNet v2 and ATTED-II, NCE for MF utilized GeneMANIA and

AraNet v2, suggesting that different types of information may increase prediction accuracy of

different GO domains.

To further gain insight into how network selection can influence NCE predictions, we

plotted the F-measures of the 2,047 tested combinations (y-axis, Figure 2D) versus the

number of made predictions for each combination (x-axis). The plot revealed that while

integrating all 11 co-function networks increases the number of made predictions (Figure 2D,

bright yellow color), the combinations that produce high F-measures included fewer than 6

networks (Figure 2D, blue color), which was observed for the three GO domains (Figure S2,

Table S2).

In addition to the leave-one-out cross validation to benchmark the networks, we also

performed a timestamp analysis [1,2], where predictions based on gold standard from

November 2015 were compared to functional information available on June 2017.

Comparison of these two gold standards revealed that within 20 months, 488 previously

unannotated genes gained functional information, while 2,588 genes gained additional GO

terms. We then calculated the F-measure of our predictions based on the new functional

information. Similarly to the previous comparison, the timestamp analysis revealed that NCE

had higher F-measure than the best single performing method (GeneMANIA) for all three

GO domains (Figure 2E, Table S3).

Influence of the gold standard completeness on the prediction accuracy and coverage

Since the gold standard is used to propagate functional information to uncharacterized genes

in gene function prediction studies, the size of the gold standard should strongly influence the

quality and number of predictions.

We first investigated how the amount of available functional information influences

the accuracy of the predictions. To this end, we incrementally removed 10% randomly

selected genes from the gold standard of 14,032 genes, and calculated the F-measure for the

remaining genes with leave-one-out analysis. Not surprisingly, when only 10% (1,403) of

gold standard genes retained functional information, the mean value for the F-measure over

100 repetitions decreased for NCE and AraNet v2, GeneMANIA and STRING (Figure 3A).

However, NCE and GeneMANIA, and to a lower degree AraNet v2 and STRING,

approached their highest F-measures at gold standard containing ~50% of genes (Figure 3A).

Since NC uses network neighbors to predict gene function, the number of made predictions

showed strong dependency on the size of the gold standard (Figure 3B). This was also

observed for the CC and MF domains (Figure S3). Taken together, the completeness of the

gold standard has a strong effect on the number and the accuracy of made predictions.

Negative correlation between gene age and gene function predictability

Our previous analyses revealed that younger genes tend to be less functionally characterized

than older genes, as the gold standard is composed mostly of older gene families [33]. While

the cause for this phenomenon can only be speculated (e.g. younger genes have weaker, easy

to miss mutant phenotypes), the amount of functional information almost perfectly reflects

the order of appearance of the gene families. We ordered the gene families from oldest (i.e.

Green Plants, representing gene families that appeared in the ancestors of green algae) to

youngest (i.e. *Arabidopsis thaliana*, representing gene families specific to *Arabidopsis*

thaliana) based on specific time (termed phylostrata) at which they appeared. We found that

the percentage of genes with experimentally verified annotation decreases as the phylostrata

become younger (Figure 3C).

To examine if prediction coverage also depends on the gene age, we calculated the

percentage of genes with predicted function for each phylostrata. Interestingly, the percentage

of genes with predicted function decreases for younger phylostrata (Figure 3C), suggesting

that function of younger genes is more difficult to predict. Since Neighbor Counting uses

network neighbors with experimentally verified annotation to predict gene function (Figure

1A), we investigated the connectivity of the genes in the different phylostrata. The analysis

revealed that younger genes tend to be less connected than older genes, as exemplified for

GeneMANIA (Figure 3D) and the remaining networks (Figure S4), explaining why fewer

predictions can be made for younger genes (Figure 3C).

Implementation of an online tool to access the ensemble predictions

To provide easy access to NCE predictions with the aim to facilitate hypothesis generation,

we have implemented an online tool, EnsembleNet, available at http://aranet.mpimp-

golm.mpg.de/ensemblenet.html. The tool provides three approaches to retrieve functional

information of genes: (i) by using a query gene to retrieve other genes with similar function

(gene-centric search), (ii) by using a GO term to retrieve genes assigned to the term (gene

ontology-centric search), and (iii) by using a group of query genes to retrieve genes with

similar function (gene set-centric search). However, to enable these analyses, we had to

systematically (i) predict which functions a given gene has and (ii) estimate which genes are

functionally related.

To predict gene function, we set to investigate which genes are associated to a given

Gene Ontology query term (see Methods). Briefly, (i) NC was used to count the number of

networks predicting that given gene is associated to the query term, followed by (ii) a

permutation analysis to derive an empirical p-value reflecting how well the prediction could

estimate correct GO term-gene associations (exemplified in Figure S5, see Methods). The

predicted gene-GO associations are available as Additional File 1. To identify functionally

related genes, we used Jaccard index (also called intersection over union) to calculate the

similarity of GO terms of any two genes. The Gene Ontology Jaccard Index (GOJI) value

ranges from 0 (two genes have no GO terms in common) to 1 (two genes have identical GO

terms). We then estimated GOJI>0.178 as a threshold capturing biologically relevant values,

as <1% of randomly selected gene pairs are found above this threshold (Figure S6, see

Methods). The resulting GOJI network is available as Additional File 2.

The gene-centric search of EnsembleNet starts by entering the ID of a query gene

(Figure 4A). The analysis returns a list of genes sorted according to the GOJI value, which

indicates how functionally similar the genes are to the query gene. This information is also

shown as an interactive GOJI network, which visualizes GOJI relationships between the

query gene (large node) and the functionally related genes. Furthermore, a table with the

predicted GO terms of the query gene indicates how much evidence (i.e. experimental

annotation and co-function network predictions) support this term prediction. The reliability

of the predicted GO term-gene association is also indicated by the empirical p-value

(exemplified in Figure S5, see Methods). Taken together, the gene-centric analysis can (i)

reveal genes that are functionally related to the query and (ii) predict the function of the

query.

The GO-centric analysis reveals which genes are assigned to the query GO term.

EnsembleNet first shows the p-value which indicates how well the predictions could correctly

predict the genes experimentally associated to the query term, thus indicating the reliability of

the predictions for the term. Furthermore, for each gene associated to the term, the table

shows how many evidences (experimental and prediction) support this association (Figure

4B). Taken together, the GO-centric search can elucidate novel genes that are involved in a

given biological process.

Finally, the gene set-centric analysis accepts several query genes to identify other

genes that are functionally related to the queries. The analysis returns a GOJI intersection

network, which shows genes that are connected to at least two queries (Figure 4C). By color

coding genes according to the number of queries they are connected to, the network can

highlights relevant genes that are highly connected to the queries. This information is also

presented as a gene table, which is sorted by the number of queries to which a given gene is

connected. To summarize, the gene set-centric search can reveal genes that are functionally

related to multiple input genes.

Identification of five complex I assembly mutants

The NADH dehydrogenase complex (complex I) is essential for cellular respiration because it

serves as the main site for electron insertion into the mitochondrial electron transport chain,

and can provide up to 40% of the protons for mitochondrial ATP formation [34–36]. The

number of complex I subunits varies between different eukaryotes due to appearance of some

lineage-specific accessory subunits (e.g. *Chlamydomonas reinhardtii*: 42 subunits; *Yarrowia*

lipolytica: 42; *Bos taurus*: 45; *Arabidopsis thaliana*: 49) [37–40]. Despite decades of research

about complex I, new subunit compositions of the complex are still being discovered [40–

42].

We demonstrate how we used the gene-, gene ontology- and gene set-centric

approaches of EnsembleNet to identify genes involved in mitochondrial respiratory chain

function, and how we arrived at five candidates: *At3g07480*, *At2g44620*, *At1g05205*,

At2g20820 and *At5g64350*. However, any of the three analyses was sufficient to uncover

these candidates, and can serve as a stand-alone approach to uncover new, relevant genes.

We first performed a gene-centric search using a NADH dehydrogenase (ubiquinone)

gene *At2g02510* (www.gene2function.de/responder.py?name=ens!at2g02510), which is

annotated as a subunit of mitochondrial complex I [43]. The analysis revealed 174

functionally related genes with GOJI values above the significance threshold (Table S4).

While many of these genes are known components of complex I (e.g. *At5g47890*,

At3g18410, *At3g03070*, and others [43], the table also contains several uncharacterized

genes, including the five candidates (highlighted in Table S4A). Furthermore, the GOJI

network of the top 50 genes contained the five candidates and revealed that most of the genes

are fully interconnected, further indicating that these genes are functionally related (Figure

5A, light blue nodes). Finally, the tables below the GOJI network can show predicted BP, CC

and MF terms of the query gene. Indeed, respiratory chain complex I (GO:0045271) is one of

the top terms found in the CC table of *At2g02510*, supported by all three co-function

networks (AraNet v2, ATTED-II and GeneMANIA), and a p-value<0.001 (Table S4B),

which is in line with the complex I function of *At2g02510.*

Next, we used "mitochondrial respiratory chain complex I" as a term-centric search

(www.gene2function.de/responder.py?name=ens!GO:0005747). The analysis returned 1,761

gene candidates, sorted according to the number of evidences (i.e. experimental annotation,

and predictions from AraNet v2, ATTED-II and GeneMANIA) supporting the association of

a gene to the GO term, which can be used to assign confidence to the prediction. Of the 1761

predictions, 32, 113, 344 and 1272 of the genes were supported by 4, 3, 2 and 1 evidences

(Table S5). The five gene candidates were found to be associated to the query term by 2, 3,

3, 2 and 3 evidences for *At3g07480, At2g44620, At1g05205, At2g20820* and *At5g64350*,

respectively (Table S5, candidates highlighted in green).

Finally, we conducted a gene set-centric search, by using four genes with

experimentally verified association to "mitochondrial respiratory chain complex I":

At5g63510, *At5g52840*, *At1g67350* and *At2g42210* [43]. The analysis revealed genes ranked

according to the number of queries they are connected to. For example, the GOJI intersection

network of top 100 connected genes showed candidates connected to four (green nodes) or

three (blue nodes) queries (Figure 5B, queries shown as large nodes). A table below the

network revealed that 98, 108 and 121 genes are connected to 4, 3 and 2 of the queries,

respectively (Table S6). The five candidates were connected to the four queries in the GOJI

intersection network (Figure 5B, nodes with gene IDs) and the table (Table S6, highlighted

genes).

361   To confirm that the five candidates are involved in complex I formation, we analyzed

knock-out mutants of these genes. Phenotypic characterization revealed that two of the

mutants showed decreased plant size (*at1g05205* and *at5g64350*, Figure 5C). To see if

complex I amount is perturbed in the mutants, we performed a blue native polyacrylamide gel

electrophoresis (BN-PAGE) analysis on the mitochondrial proteome (Figure 5D), which

allowed us to estimate the abundance of complex I, by calculating the complex I:complex III

and complex I:complex V ratios (Figure 5E, [44]). The analysis revealed that the five mutants

have significantly different CI/CV and CI/CIII ratios from wild type (Col-0, p-value<0.01,

Table S7), where four of the mutants have lower ratio of complex I, while *at2g20820* has

higher amount (Figure 5E). Interestingly, while the dwarf phenotypes of *at1g05205* and

at5g64350 are associated with a decrease of complex I ratio, the strong decrease of the ratio

in *at2g44620* did not produce any observable growth phenotype (Figure 5C). This was also

observed for the *ca2* mutant, where 90% decrease of complex I produced no growth

phenotype [45].

Discussion and conclusion

Gene function prediction is one of the most active fields of bioinformatics, and recent

publications show a constant improvement of the predictive power of these approaches [1,2].

In this paper, we studied ensemble gene function prediction, a powerful, but unexplored

approach in the plant field.

381   We used Neighbor counting [31] and the F-measure [1,2] to benchmark the 11

networks. As expected, the second generation integrative approaches perform better than the

first generation networks (Figure 1). However, while the two top networks (GeneMANIA and

AraNet v2) use different data sources and methods to integrate them (regression vs. Bayesian,

respectively), the performance of the two networks was similar (Figure 1C), suggesting that

the performance of the second generation methods might have reached it's peak.

We next established an ensemble prediction by using two different approaches

(Figure 2). The neighbor counting ensemble (NCE) integrates the predictions to arrive at an

ensemble prediction, while the STATISnet integrates the co-function networks to arrive at an

ensemble network. While integration of the networks via STATISnet produced an inferior

prediction than GeneMANIA, NCE showed the highest accuracy and coverage (Figure 2C).

In contrasts to conclusions from previous study [27], integration of all 11 methods did not

result in highest accuracy, but rather highest prediction coverage at the cost of accuracy

(Figure 2C-D). To arrive at the highest accuracy, a combination of GeneMania, AraNet v2

and ATTED-II was needed to address BP and CC, while combination of GeneMANIA and

AraNet v2 was better suited to address MF domain. This suggests that careful combination of

the inference methods might be needed to achieve highest accuracy for different domains of

gene function.

We next investigated how the amount of available information influences the

accuracy and coverage of the predictions. Upon removing genes from the gold standard, we

observed that the prediction accuracy dropped substantially for gold standard size of <50%,

followed by a more modest increase for gold standard size of >50% (Figure 3A, Figure S3).

The number of made predictions shows a strong dependency on the size of the gold standard,

as in contrast to accuracy, the prediction coverage shows a substantial increase for gold

standard size >50%. These results are expected, as function of the uncharacterized genes in

this study is transferred from their characterized direct network neighbors, which strongly

depends on the fraction of gold standard genes in the network. To partially circumvent this

dependency, density-based methods which identify highly connected network modules could

be used to annotate genes that are not directly connected [31].

Are functions of certain genes more difficult to predict? Our previous study revealed

that older gene families are more functionally characterized than younger gene families

(Figure 3C, [33]). Surprisingly, we observed that functions of older gene families can be

more readily predicted than for younger genes (Figure 3C), which could be attributed to

higher disconnectedness of younger genes (Figure 3D, Figure S4). Interestingly, we observed

this in BIOGRID protein-protein interaction networks and to a weaker degree for co-

expression networks based on RNA-sequencing data (Figure S4). Higher connectivity of

older genes in protein-protein interaction networks could be explained by selection bias (e.g.

older genes have stronger mutant phenotypes, and are more likely to be used as bait in yeast-

two-hybrid experiments), but we observed the same effect for co-expression networks based

on unbiased genome-wide RNA-seq analysis, which indicates that the disconnectedness is an

inherent property of young genes. Since it is not possible to predict gene function of

disconnected genes, function prediction of the young genes still remains a challenge.

To provide an easy access to the predictions, we present the EnsembleNet database

which allows gene-, gene ontology- and gene set-centric searches (Figure 4). These versatile

features enabled us to identify 5 uncharacterized genes important for complex I assembly

(Figure 5). The high granularity of Gene Ontology terms which allows interrogating the

polygenic composition of specific processes (e.g. GO:0035619: "root hair tip"), together with

cutting-edge performance, represents a major step towards the goal of computationally

bridging the gap between genotype and phenotypes in plants.

Methods

Retrieval and normalization of gene co-function networks

The most recent versions of networks available for *Arabidopsis thaliana* on 21.11.2015 were

downloaded for AraNet2 (http://www.inetbio.org/aranet/downloadnetwork.php), STRING

(https://string-db.org/cgi/download.pl), GeneMANIA (http://genemania.org/data/), ATTED-II

(http://atted.jp/download.shtml), PlaNet (http://aranet.mpimp-golm.mpg.de/download.html),

Conserved Co-expression Arabidopsis Functional Modules (CC-AFM), PODC

(http://plantomics.mind.meiji.ac.jp/podc/bulkDownload.html), TAIR-PPI

(ftp://ftp.arabidopsis.org/home/tair/Proteins/Protein_interaction_data/), BIOGRID

(https://thebiogrid.org/download.php) and CORNET

(https://bioinformatics.psb.ugent.be/cornet/). CC-AFM refers to Arabidopsis co-expression

edges (top 300 per gene based on Pearson Correlation Coefficient) filtered with conserved

co-expression information from six other plants, including *Glycine max*, *Medicago*

truncatula, *Oryza sativa*, *Populus trichocarpa*, *Vitis vinifera* and *Zea mays* from [26].

Furthermore, normalized expression data derived from RNA-seq experiments from [46] was

used to construct Highest Reciprocal Rank (HRR) co-expression networks, as done in PlaNet

(HRR cut-off=30, [4]).

Edges of each of the networks were scaled to range between 0 and 1, where 1 indicates the

strongest association between a gene pair, while 0 indicates the weakest association.

Retrieval of gold standard - experimentally verified functional annotation

Experimentally verified Gene Ontology annotations were downloaded from TAIR

(www.arabidopsis.org), as ATH_GO_GOSLIM.txt file on 21.11.2015 (used to construct and

benchmark EnsembleNet) and 10.08.2017 (used to perform time-stamp analysis). A gene was

considered experimentally annotated if it was associated with GO terms having EXP, IDA,

IPI, IMP, IGI, IEP or TAS evidence codes. To avoid evaluating general terms, (e.g.

"cellular_component", "binding", "biological regulation"), terms that were assigned to more

than 5% of the 14,031 experimentally annotated genes were removed. This removed 105 out

of 7,352 terms. For each gene, all terms assigned to a gene were recursively propagated

towards the root of the terms (excluding the terms present in more than 5% of the

experimentally annotated genes), regardless of the type of relationship between terms.

Calculating F-measure

The main evaluation metric in this study was F-measure, as defined by Critical Assessment of

protein Function Annotation (CAFA) experiment [1,2].

For a given target gene i and network edge threshold $t \in [0,1]$, the precision and recall were

calculated as:

$pr_i(t) = \frac{|P_t \cap T|}{|P_t|}$ and $rc_i(t) = \frac{|P_t \cap T|}{|T|}$

where $|P_t|$ is the number of predicted terms at edge threshold t, $|T|$ is the number of

experimentally determined terms, and $|P_t \cap T|$ is the number of terms appearing in both P_t and

T.

Performance of a network at a given edge threshold is obtained by averaging precision and

recall across genes. Precision at edge threshold t is calculated as:

$pr(t) = \frac{1}{m_{(t)}} * \sum_{i=1}^{m(t)} pr_i(t),$

where $m(t)$ is the number of genes for which at least one prediction was made, above edge

threshold t. Precision is calculated only for the genes for which a prediction could be made. A

prediction cannot be made for genes that are not connected to genes with any experimentally

determined annotations at a given t.

Conversely, recall is calculated over all n genes in a target set, as:

$rc(t) = \frac{1}{n} * \sum_{i=1}^{n} rc_i(t),$

regardless of whether a prediction can be made at a given threshold t.

The F-measure (a harmonic mean between precision and recall) provides a number useful to

compare performances of the networks at a given threshold. An edge threshold where a given

network performs highest in terms of the F-measure, can be identified by obtaining the

maximum F-measure (F_{max}) over all thresholds, given by:

$$F_{max} = max_t \left\{ \frac{2 * pr(t) * rc(t)}{pr(t) + rc(t)} \right\}$$

Associating genes to Gene Ontology terms

To predict which genes are associated with a given GO term (query term), we first removed

GO terms assigned to more genes that the query term from the gold standard. The reason for

removing these terms is that more frequent terms (e.g. "binding", "enzymatic activity") are

by chance more likely to be more abundant in network neighborhoods than less frequent

terms. Next, for each gene in the used networks (i.e. GeneMANIA and AraNet v2 for MF

domain), NCE was used to predict genes with the query term.

Calculating the empirical p-value to evaluate gene - Gene Ontology term associations

To estimate how accurately genes assigned to a given Gene Ontology term can be predicted

by NCE, we first retrieved genes experimentally assigned to the term of interest. Then, for

each of these genes, we counted the number of networks for which the neighbor counting

made a correct prediction (Figure S2). This number, $pred_{observed}$, is indicative of the extent to

which a term can be accurately predicted in a network setting. See Figure S5A and S5B for

toy examples where a term can be predicted well ($\text{pred}_{\text{observed}} = 9$, out of 9 predictions) and

poorly ($\text{pred}_{\text{observed}} = 2$, out of 9 predictions), respectively.

Next, for each gene that is experimentally assigned to the term, we sampled a random

prediction from the pool of all predicted gene-term associations, and counted the number of

networks which assigned a term to a gene ($\text{pred}_{\text{sampled}}$). The empirical p-value is defined as:

$$p - value_{term} = \frac{\sum_{n=1}^{N} I(pred_{observed} \leq pred_{sampled)}}{N}$$

Where $\text{pred}_{\text{observed}}$ is the observed number of correct predictions, $\text{pred}_{\text{sampled}}$ is the number of

correct calls from sampled predictions, N is the number of permutations, and I is an indicator

function. To account for multiple hypothesis testing, we applied False Discovery Rate (FDR)

correction to the p-values [47]. The predicted gene-GO term associations and p-values are

available as Supplemental Data 1.

Calculating STATIS compromise network

To combine the 11 symmetric (i.e., undirected) weighted networks, we applied an approach

similar to STATIS [32]. We first determined the Rv coefficient for each pair of the scaled

networks (represented as scaled adjacency matrices), gathered in the cosine matrix C. The

entries in the first eigenvector of C, after rescaling to unit norm, were used as weights to

determine the contribution of each scaled adjacency matrix to the STATIS network. We note

that the larger the value for the weight, the bigger the contribution of the respective data set to

the adjacency matrix of the STATIS network. The STATIS network was obtained as the

weighted sum of the scaled adjacency matrices. The resulting adjacency matrix *S* was also

scaled by dividing with its maximum value. The STATIS network is available for download

at http://aranet.mpimp-golm.mpg.de/download.html.

Identifying functionally related genes

To estimate which genes are functionally related, we first retrieved the predicted terms for

each gene, and calculated the Jaccard Index between all possible gene pairs, defined as:

$J(X,Y) = |X \cap Y| / |X \cup Y|$, where X and Y are predicted GO terms of gene x and y. To estimate

which gene pairs are functionally related, we calculated the Jaccard Index between 10.000

random gene pairs and found the J threshold (0.178) below which 99% of the random pairs

were found (Figure S6). Thus, gene pairs with J>0.178 were deemed to be functionally

related. The resulting GOJI network is available as Supplemental Data 2.

Plant material, plant growth and genotyping

Seeds were surface sterilized with 70 % (v/v) ethanol containing 0.5 % (v/v) Triton X100,

sown on synthetic media (0.5× MS, 1 % sucrose, 0.7 % agar) and incubated under long day

photoperiod: 16 h light (150 µE, 22 °C)/8 h dark (20 °C). After 2 weeks, seedlings were

transferred to soil and grown under long day photoperiod. Plant DNA was isolated as

described in [48], and genotyping PCR was performed as in [49]. The genotyping primers for

detecting the wild type and mutant alleles are given in Table S8.

Mitochondria isolation, Blue-Native gel electrophoresis and quantification of complex I.

The aerial organs of 6-week-old plants were harvested and mitochondria were isolated

according to [50]. For Blue-Native gel electrophoresis, total mitochondria proteins (50 µg)

were solubilized with dodecylmaltoside (1% [w/v] final) in ACA buffer (750 mM amino

caproic acid, 0.5 mM EDTA, and 50 mM Tris-HCl, pH 7.0) and incubated 20 min at 4°C.

After centrifugation for 10 min at 20000g, the supernatant was recovered and supplemented

with Coomassie Blue G250 (0.2 % [w/v] final). The samples were loaded onto a 4.5% to 16%

(w/v) polyacrylamide gel (ratio acrylamide/bisacrylamide: 29/1, gel buffer: 0.25 M

aminocaproic acid, 25 mM Bis-Tris-HCl pH 7.0). The migration of the samples was

performed at 4°C using cathode buffer (50 mM Tricine, 15 mM Bis-Tris-HCl pH 7.0, 0.02%

[w/v] Coomassie Blue G250) and anode buffer (50 mM Bis-Tris-HCl pH 7.0) in two steps,

first 100 V for 45 min then 15 mA (limiting voltage: 500 V) for 10 h. After migration the gel

was stained using colloidal blue (3% [v/v] orthophosphoric acid, 6% [w/v] ammonium

sulphate, 0.1% [w/v] Coomassie Blue G250) and destained with ultrapure water. The gel was

scanned using a flatbed scanner. The gel analyzer tool from ImageJ was used to quantify the

bands corresponding to complexes I, III and V.

Declarations

Availability of data and materials

The datasets analyzed in this study are available from the repositories given in the Methods.

Competing interests

The authors do not have any competing interests

Funding

This work was supported by the Max-Planck-Gesellschaft (B.H, E.M., C.F., N.V., Z.N.,

M.M.).

Author contributions

M.M. conceived the project. M.M. and B.H. performed the bioinformatical analyses with

input from Z.N.. K.V and S.M. provided CC-AFM network. E.M. performed the mutant

analyses. M.M., B.H., E.M., C.F., N.V., K.V. and Z.N. wrote the article with help from all

authors.

Acknowledgements

We would like to express our gratitude to the creators of ATTED-II, GeneMANIA, STRING,

AraNet v2, BIOGRID, CORNET, PODC and TAIR for providing these excellent tools and

networks. We would like to thank the employees of the Max Planck Institute of Molecular

Plant Physiology, and especially Dr. Arun Sampathkumar, for their feedback regarding

EnsembleNet tool. Marek Mutwil would like to thank Sue Rhee for the great discussions

regarding gene prediction function. This work was supported by the Max-Planck-Gesellschaft

(B.H, E.M., C.F., N.V., Z.N., M.M.).

References

1. Radivojac P, Clark WT, Oron TR, Schnoes AM, Wittkop T, Sokolov A, et al. A large-scale

evaluation of computational protein function prediction. Nat. Methods. 2013;10:221–7.

2. Jiang Y, Oron TR, Clark WT, Bankapur AR, D'Andrea D, Lepore R, et al. An expanded

evaluation of protein function prediction methods shows an improvement in accuracy.

Genome Biol. [Internet]. 2016;17:184. Available from:

http://genomebiology.biomedcentral.com/articles/10.1186/s13059-016-1037-6

3. Rhee SY, Mutwil M. Towards revealing the functions of all genes in plants. Trends Plant

Sci. 2014;19:212–21.

4. Mutwil M, Usadel B, Schütte M, Loraine A, Ebenhöh O, Persson S. Assembly of an

interactive correlation network for the Arabidopsis genome using a novel heuristic clustering

algorithm. Plant Physiol. 2010;152:29–43.

5. Bassel GW, Lan H, Glaab E, Gibbs DJ, Gerjets T, Krasnogor N, et al. Genome-wide

network model capturing seed germination reveals coordinated regulation of plant cellular

phase transitions. Proc. Natl. Acad. Sci. [Internet]. 2011;108:9709–14. Available from:

http://www.pnas.org/content/108/23/9709.abstract%5Cnhttp://www.pnas.org/content/108/23/

9709.full.pdf

6. Jiménez-Gómez JM, Wallace AD, Maloof JN. Network analysis identifies ELF3 as a QTL

for the shade avoidance response in arabidopsis. PLoS Genet. 2010;6.

7. Takabayashi A, Ishikawa N, Obayashi T, Ishida S, Obokata J, Endo T, et al. Three novel

subunits of Arabidopsis chloroplastic NAD(P)H dehydrogenase identified by bioinformatic

and reverse genetic approaches. Plant J. 2009;57:207–19.

8. Takahashi N, Lammens T, Boudolf V, Maes S, Yoshizumi T, De Jaeger G, et al. The DNA

replication checkpoint aids survival of plants deficient in the novel replisome factor ETG1.

EMBO J. [Internet]. 2008;27:1840–51. Available from:

http://emboj.embopress.org/content/27/13/1840.abstract

9. Lee I, Ambaru B, Thakkar P, Marcotte EM, Rhee SY. Rational association of genes with

traits using a genome-scale gene network for Arabidopsis thaliana. Nat. Biotechnol.

[Internet]. 2010;28:149–56. Available from: http://dx.doi.org/10.1038/nbt.1603

10. Ashburner M, Ball CA, Blake JA, Butler H, Cherry JM, Eppig JT, et al. Creating the

Gene Ontology resource: Design and implementation. Genome Res. 2001;11:1425–33.

11. Garcia-Hernandez M, Daviesl E, Staswicksii PE. Arabidopsis p40 Homologue. HE J.

Biol. Chem. 1994;269:20744–9.

12. Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, et al.

InterProScan: Protein domains identifier. Nucleic Acids Res. 2005;33.

13. Proost S, Mutwil M. Tools of the trade: Studying molecular networks in plants. Curr.

Opin. Plant Biol. 2016. p. 130–40.

14. Pavlidis P, Gillis J. Progress and challenges in the computational prediction of gene

function using networks. F1000Research [Internet]. 2012; Available from:

http://f1000research.com/articles/1-14/v1

15. Pellegrini M, Marcotte EM, Thompson MJ, Eisenberg D, Yeates TO. Assigning protein

functions by comparative genome analysis: Protein phylogenetic profiles. Proc. Natl. Acad.

Sci. [Internet]. 1999;96:4285–8. Available from:

http://www.pnas.org/cgi/doi/10.1073/pnas.96.8.4285

16. Lee JM, Sonnhammer ELL. Genomic gene clustering analysis of pathways in eukaryotes.

Genome Res. 2003;13:875–82.

17. Nakamura Y, Itoh T, Martin W. Rate and polarity of gene fusion and fission in Oryza

sativa and Arabidopsis thaliana. Mol. Biol. Evol. 2007;24:110–21.

18. Wang Z, Zhang XC, Le MH, Xu D, Stacey G, Cheng J. A protein domain Co-occurrence

network approach for predicting protein function and inferring species phylogeny. PLoS One.

2011;6.

19. Lee B -h. The Arabidopsis Cold-Responsive Transcriptome and Its Regulation by ICE1.

PLANT CELL ONLINE [Internet]. 2005;17:3155–75. Available from:

http://www.plantcell.org/cgi/doi/10.1105/tpc.105.035568

20. Wu LF, Hughes TR, Davierwala AP, Robinson MD, Stoughton R, Altschuler SJ. Large-

scale prediction of Saccharomyces cerevisiae gene function using overlapping transcriptional

clusters. Nat. Genet. [Internet]. 2002;31:255–65. Available from:

http://www.nature.com/doifinder/10.1038/ng906

21. Vazquez A, Flammini A, Maritan A, Vespignani A. Global protein function prediction

from protein-protein interaction networks. Nat. Biotechnol. [Internet]. 2003;21:697–700.

Available from: http://www.nature.com/doifinder/10.1038/nbt825

22. Heyndrickx KS, Vandepoele K. Systematic identification of functional plant modules

through the integration of complementary data sources. Plant Physiol [Internet].

2012;159:884–901. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22589469

23. Mutwil M, Klie S, Tohge T, Giorgi FM, Wilkins O, Campbell MM, et al. PlaNet:

combined sequence and expression comparisons across plant networks derived from seven

species. Plant Cell. 2011;23:895–910.

24. Arabidopsis Interactome Mapping Consortium. Evidence for network evolution in an

Arabidopsis interactome map. Science. 2011;333:601–7.

25. Kudla J, Bock R. Lighting the Way to Protein-Protein Interactions: Recommendations on

Best Practices for Bimolecular Fluorescence Complementation Analyses. Plant Cell

[Internet]. 2016;28:1002–8. Available from:

http://www.plantcell.org/lookup/doi/10.1105/tpc.16.00043

26. Heyndrickx KS, Vandepoele K. Systematic Identification of Functional Plant Modules

through the Integration of Complementary Data Sources. PLANT Physiol. 2012;159:884–

901.

27. Marbach D, Costello JC, Küffner R, Vega NM, Prill RJ, Camacho DM, et al. Wisdom of

crowds for robust gene network inference. Nat. Methods. Nature Publishing Group, a division

of Macmillan Publishers Limited. All Rights Reserved.; 2012;9:796–804.

28. Tanz SK, Castleden I, Hooper CM, Vacher M, Small I, Millar H a. SUBA3: A database

for integrating experimentation and prediction to define the SUBcellular location of proteins

in Arabidopsis. Nucleic Acids Res. 2013;41:1185–91.

29. Vermeirssen V, De Clercq I, Van Parys T, Van Breusegem F, Van de Peer Y. Arabidopsis

Ensemble Reverse-Engineered Gene Regulatory Network Discloses Interconnected

Transcription Factors in Oxidative Stress. Plant Cell [Internet]. 2014;26:4656–79. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/25549671

30. Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, et al. The

Arabidopsis Information Resource (TAIR): Improved gene annotation and new tools. Nucleic

Acids Res. 2012;40.

31. Sharan R, Ulitsky I, Shamir R. Network-based prediction of protein function. Mol. Syst.

Biol. 2007;3:88.

32. Abdi H, Williams LJ, Valentin D, Bennani-Dosse M. STATIS and DISTATIS: Optimum

multitable principal component analysis and three way metric multidimensional scaling.

Wiley Interdiscip. Rev. Comput. Stat. 2012. p. 124–67.

33. Ruprecht C, Proost S, Hernandez-Coronado M, Ortiz-Ramirez C, Lang D, Rensing SA, et

al. Phylogenomic analysis of gene co-expression networks reveals the evolution of functional

modules. Plant J. [Internet]. 2017;n/a-n/a. Available from:

http://dx.doi.org/10.1111/tpj.13502

34. Hunte C, Zickermann V, Brandt U. Functional Modules and Structural Basis of

Conformational Coupling in Mitochondrial Complex I. Science (80-.). [Internet].

2010;329:448–51. Available from:

http://www.sciencemag.org/cgi/doi/10.1126/science.1191046

35. Braun H-P, Binder S, Brennicke A, Eubel H, Fernie AR, Finkemeier I, et al. The life of

plant mitochondrial complex I. Mitochondrion [Internet]. 2014;19:295–313. Available from:

http://linkinghub.elsevier.com/retrieve/pii/S1567724914000208

36. Hirst J. Mitochondrial Complex I. Annu. Rev. Biochem. [Internet]. 2013;82:551–75.

Available from: http://www.annualreviews.org/doi/10.1146/annurev-biochem-070511-

103700

37. Cardol P, Vanrobaeys F, Devreese B, Van Beeumen J, Matagne RF, Remacle C. Higher

plant-like subunit composition of mitochondrial complex I from Chlamydomonas reinhardtii:

31 Conserved components among eukaryotes. Biochim. Biophys. Acta - Bioenerg.

2004;1658:212–24.

38. Carroll J, Fearnley IM, Skehel JM, Shannon RJ, Hirst J, Walker JE. Bovine complex I is a

complex of 45 different subunits. J. Biol. Chem. 2006;281:32724–7.

39. Angerer H, Zwicker K, Wumaier Z, Sokolova L, Heide H, Steger M, et al. A scaffold of

accessory subunits links the peripheral arm and the distal proton-pumping module of

mitochondrial complex I. Biochem. J. [Internet]. 2011;437:279–88. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3273332&tool=pmcentrez&rende

rtype=abstract

40. Peters K, Belt K, Braun H-P. 3D Gel Map of Arabidopsis Complex I. Front. Plant Sci.

[Internet]. 2013;4:153. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3671202&tool=pmcentrez&rende

rtype=abstract

41. Guerrero-Castillo S, Cabrera-Orefice A, Huynen MA, Arnold S. Identification and

evolutionary analysis of tissue-specific isoforms of mitochondrial complex I subunit

NDUFV3. Biochim. Biophys. Acta - Bioenerg. 2017;1858:208–17.

42. Bridges HR, Mohammed K, Harbour ME, Hirst J. Subunit NDUFV3 is present in two

distinct isoforms in mammalian complex I. Biochim. Biophys. Acta - Bioenerg.

2017;1858:197–207.

43. Klodmann J, Braun H-P. Proteomic approach to characterize mitochondrial complex I

from plants. Phytochemistry [Internet]. 2011;72:1071–80. Available from:

http://www.sciencedirect.com/science/article/pii/S0031942210004334

44. Peters K, Nießen M, Peterhänsel C, Späth B, Hölzle A, Binder S, et al. Complex I-

complex II ratio strongly differs in various organs of Arabidopsis thaliana. Plant Mol. Biol.

2012;79:273–84.

45. Perales M, Eubel H, Heinemeyer J, Colaneri A, Zabaleta E, Braun HP. Disruption of a

nuclear gene encoding a mitochondrial gamma carbonic anhydrase reduces complex I and

supercomplex I+III2 levels and alters mitochondrial physiology in Arabidopsis. J. Mol. Biol.

2005;350:263–77.

46. Giorgi FM, Del Fabbro C, Licausi F. Comparative study of RNA-seq- and Microarray-

derived coexpression networks in Arabidopsis thaliana. Bioinformatics. 2013;29:717–24.

47. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B. [Royal Statistical Society,

Wiley]; 1995;57:289–300.

48. Weigel D, Glazebrook J. Dellaporta miniprep for plant DNA isolation. Cold Spring Harb.

Protoc. 2009;4.

49. O'Malley RC, Barragan CC, Ecker JR. A user's guide to the arabidopsis T-DNA insertion

mutant collections. Plant Funct. Genomics Methods Protoc. Second Ed. 2015. p. 323–42.

50. Kühn K, Obata T, Feher K, Bock R, Fernie AR, Meyer EH. Complete Mitochondrial

Complex I Deficiency Induces an Up-Regulation of Respiratory Fluxes That Is Abolished by

Traces of Functional Complex I. Plant Physiol. [Internet]. 2015;168:1537–49. Available

from: http://www.plantphysiol.org/lookup/doi/10.1104/pp.15.00589

51. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al.

STRING v10: protein-protein interaction networks, integrated over the tree of life. Nucleic

Acids Res. 2015;43:D447-52.

52. Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, et al. The

GeneMANIA prediction server: biological network integration for gene prioritization and

predicting gene function. Nucleic Acids Res. [Internet]. 2010;38:W214–20. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2896186&tool=pmcentrez&rende

rtype=abstract%5Cnhttp://nar.oxfordjournals.org/lookup/doi/10.1093/nar/gkq537

53. Lee T, Yang S, Kim E, Ko Y, Hwang S, Shin J, et al. AraNet v2: an improved database of

co-functional gene networks for the study of Arabidopsis thaliana and 27 other nonmodel

plant species. Nucleic Acids Res. 2015;43:D996-1002.

54. Aoki Y, Okamura Y, Tadaka S, Kinoshita K, Obayashi T. ATTED-II in 2016: A Plant

Coexpression Database Towards Lineage-Specific Coexpression. Plant Cell Physiol.

[Internet]. 2015;2:pcv165-. Available from:

http://pcp.oxfordjournals.org/content/early/2016/01/11/pcp.pcv165.full

55. Proost S, Mutwil M. PlaNet: Comparative Co-Expression Network Analyses for Plants.

Methods Mol. Biol. [Internet]. 2017 [cited 2017 Feb 17]. p. 213–27. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/27987173

56. Ohyanagi H, Takano T, Terashima S, Kobayashi M, Kanno M, Morimoto K, et al. Plant

Omics Data Center: an integrated web repository for interspecies gene expression networks

with NLP-based curation. Plant Cell Physiol. 2015;56:e9.

57. Chatr-Aryamontri A, Oughtred R, Boucher L, Rust J, Chang C, Kolas NK, et al. The

BioGRID interaction database: 2017 update. Nucleic Acids Res. 2017;45:D369–79.

58. De Bodt S, Hollunder J, Nelissen H, Meulemeester N, Inzé D. CORNET 2.0: integrating

plant coexpression, protein-protein interactions, regulatory interactions, gene associations and

functional annotations. New Phytol. 2012;195:707–20.

Tables:

Table 1. Used co-function networks.

Network	Data source	Integration scheme	Number of nodes (genes)	Number of edges
STRING [51]	PPI, TM , CE, CO, N, F	Bayesian	24283	5318676
GeneMANIA [52]	PPI, CE, CL, D, GI	Regression	25215	11333378
AraNet v2 [53]	PPI, TM, CE, GN, GF, DC	Bayesian	22894	895000
ATTED-II [54]	CE	None	20835	719050

PlaNet [55]	CE	None	21051	105955
CC-AFM [26]	CE	None	16494	685666
PlaNet: RNA-seq	CE	None	25183	166677
PODC [56]	CE	None	8864	753072
TAIR-PPI [30]	PPI	None	7178	72267
BIOGRID [57]	PPI	None	9529	35641
CORNET [58]	PPI	None	3014	51885

Type of data and integration scheme. PPI (protein-protein interaction), TM (text mining), CE

(co-expression), CO (genomic co-occurrence), GN (genomic neighbor), GF (gene fusion), CL

(co-localization), D (shared protein domains), GI (genetic interaction), DC (domain co-

occurrence).

Figures:

Figure 1. Evaluation of the eleven gene co-function networks. A) Illustration of the

Neighbor Counting method and influence of the edge cutoff on predictions, exemplified by

four toy examples. The toy examples include the query gene (large node) connected to other

genes in the co-function network (smaller nodes). The edges connecting genes have weights

ranging from weak (solid gray edge) to strong (solid black edge). Gene functions assigned to

a gene are depicted as red, blue and green colors. Neighbor Counting method assigns function

to the query gene (large node) that occur most frequently in its neighbors. The accuracy of

the prediction can be estimated by comparing the known functions of the query with the

predicted function. B) Influence of the edge cutoff on the F-measure (blue line) and made

predictions (red line) for STRING BP predictions. The edge cutoff where F-measure is

highest (F_{max}) is indicated. C) Performance of the eleven co-function network expressed as

F_{max} value (gray barcharts, left y-axis) and the number of made predictions (red bars, right y-

axis).

Figure 2. Performance of ensemble gene function prediction methods. A) Outline of the

Neighbor Counting Ensemble (NCE) approach, exemplified by a toy example. Function for

the query gene (large white node) is estimated by neighbor counting, predicting orange, red

and red function from networks a, b and c, respectively. In the next step, similarly to the

Neighbor Counting, the ensemble prediction is obtained by selecting the most frequent

function from the previous step, assigning red function to the query gene. B) Outline of the

STATIS method. C) Performance summary of the methods for BP (red), CC (blue) and MF

(green). The left y-axis (bars) indicates the F-measure, while the right y-axis (black line)

indicates the number of made predictions. D) Analysis of the 11 gene co-function networks,

resulting in 2047 tested combinations. The x-axis indicates the number of made predictions,

while the y-axis shows the F-measure. Point color indicates the number of integrated

networks, ranging from 1 network (dark blue) to 11 (bright yellow). E) Time-stamp

comparison of the Ensemble method and GeneMANIA predictions. The y-axis indicates the

F-measure.

Figure 3. Influence of available functional information on prediction performance. A)

Relationship between the F-measure (y-axis) and % of genes with available GO terms (x-

axis) in the three best performing co-function networks (dashed lines) and NCE (solid red

line). The lines represent bootstrapped 68% confidence interval plots of 100 permutations. B)

Relationship between the number of made predictions (y-axis) and % of genes with available

GO terms (x-axis). C) Percentage (y-axis) of genes from the different phylostrata (given by

color legend in Figure 3D), for experimentally characterized genes, Ensemble predictions and

GeneMANIA predictions. D) Relationship between the percentage of genes with a neighbor

in a network (y-axis) and the phylostrata of the genes (x-axis). The example is given for

GeneMANIA, BP.

Figure 4. Features of EnsembleNet. A) The query gene-centric can reveal genes that are

functionally related to the query, and predict function of the query. Functionally related genes

are identified by i) Gene Ontology Jaccard Index (GOJI) table and ii) network. Predicted

Gene Ontology (GO) terms of the query are given in the tables below the network. B) Gene

Ontology-centric analysis reveals which genes are assigned to a given GO term. For each

gene, the table shows how many evidences (experimental annotation and co-function network

predictions) support this association. C) Gene set-centric analysis shows i) GOJI intersection

network, which reveals genes that are connected to at least two of the query genes. In this

example, large nodes correspond to query genes, small nodes represent genes connected to

the queries, while edges represent GOJI associations above the significance threshold. Note

that genes a and d are removed from the network, as they are connected to only one query

gene. ii) Gene table shows to how many queries the found genes are connected to.

Figure 5. Functional characterization of *at3g07480*, *at2g44620*, *at1g05205*, *at2g20820*

and *at5g64350*. A) GOJI network of At4g30010 (large node). Smaller nodes represent genes

connected to the At4g30010. The edges connecting the nodes represent GOJI values above

the significance threshold. Genes selected for functional characterization are indicated as dark

blue nodes. B) Gene set analysis using queries *At5g63510*, *At5g52840*, *At1g67350* and

At2g42210 (large nodes). 100 Genes connected to four (green nodes) or three (blue nodes) of

the queries are shown. The five genes selected for functional analysis are indicated. C) Top

view of 5 week old rosettes. D) Blue Native Polyacrylamide Gel Electrophoresis (BN-PAGE)

analysis of the five mutants and wild type (Col-0). Bands corresponding to complexes I, III

and V are indicated to the left. E) Calculated ratio of complex I/complex V (CI/CV) and

complex I/complex III (CI/CIII). The bars represent means, while the error bars represent

standard deviation (n=4).

Supplemental figures:

Figure S1. Status of the functional annotation of the 28392 *Arabidopsis thaliana* genes on

November 2015. A) The pie charts show the fraction of experimentally validated (green),

predicted (blue) and unknown function (gray) genes for BP, CC and MF. B) Venn diagram

showing the overlap of functionally characterized genes in the three domains of GO.

Figure S2. Combinatorial analysis of the 11 gene co-function networks for BP (top), CC

(middle) and MF (bottom). The x-axis indicates the number of made predictions, while the

y-axis shows the F-measure. Point color indicates the number of integrated networks, ranging

from 1 network (dark blue) to 11 (bright yellow).

Figure S3. Influence of available functional information on F-measure and the number

of made predictions. Relationship between the F-measure (y-axis) and % of genes with

available GO terms (x-axis) given for BP (top left), CC (middle left) and MF (bottom left).

Relationship between the number of made predictions (y-axis) and % of genes with available

GO terms (x-axis) for BP (top right), CC (middle eight) and MF (bottom right). The lines

represent bootstrapped 68% confidence interval plots of 100 permutations.

Figure S4 Relationship between the percentage of genes with a neighbor in a network

(y-axis) and the phylostrata of the genes (x-axis). The example is given for ATTED-II

(top), AraNet v2 (middle) and GeneMANIA (bottom), BP.

Figure S5. Estimating the performance of Gene Ontology term predictions. A) An

example of a GO term that can be predicted well. Three genes (1-3) experimentally assigned

to the term are predicted to be assigned to this term in 9 cases (indicated by green +). B) An

example of a GO term that cannot be predicted well. Only 2 out of 9 networks predicted that

the experimentally assigned genes A-C are assigned to this term.

Figure S6 Estimating the significant Gene Ontology Jaccard Index (GOJI) value. The

histogram shows the relationship between GOJI values (x-axis) and frequency of random

gene pairs showing the particular value. The red line indicates the GOJI threshold (0.178)

below which 99% of the random pairs are found.

Supplemental tables:

Table S1. F-measure, precision, recall, number of made predictions of the 11 co-

function networks at the different edge cutoffs. The values are given for the three domains

of GO. STATISnet is included.

Table S2. F-measure, precision, recall, number of made predictions for all combinations

of the 11 co-function networks at the F_{max} edge cutoffs. Tables S2A, S2B and S2C

correspond to BP, CC and MF, respectively.

Table S3. Timestamp F-measure, precision, recall, number of made predictions of the

11 co-function networks and NCE. The method with highest F-measure in a given domain

is in bold.

Table S4. Gene-centric analysis using *At2g02510*. A) Output of the GOJI table, where

genes are sorted according to the decreasing GOJI value. The five candidates are highlighted

in green. B) Predicted BP domain terms of *At2g02510*. The number of evidences supporting

this function is a sum of evidences (Experimentally verified function, AraNet v2, ATTED-II

and GeneMANIA) that support (Y) this association.

Table S5. GO-centric analysis using *GO:0005747* ("mitochondrial respiratory chain

complex I"). The table is sorted according to the number of evidences (Experimentally

verified function, AraNet v2, ATTED-II and GeneMANIA) supporting this association (Y).

The five candidates are highlighted in green.

Table S6. Gene set-centric analysis using queries *At5g63510, At5g52840, At1g67350 and*

***At2g42210*.** The table is sorted according to the number of the queries a given gene is

connected to with a GOJI edge. The five candidates are highlighted in green.

Table S7. Statistical analysis of complex I/V and complex I/III ratios. The ratios of the

mutant and wild type (Col-0) indicate the t-test statistic (n=4).

Table S8. Primers used to test for wild type allele (RP+LP) and mutant allele (RP+LB)

for the five knock out lines.

Additional files:

Additional File 1. NCE predicted gene-GO term associations for BP, CC and MF. The

columns in each file show the GO ID, empirical p-value reflecting how well the prediction

could estimate correct GO term-gene associations, gene ID, and indication if a given co-

function network predicted the association (1), did not predict this association (0), or could

not make a prediction for this gene (-).

Additional File 2. Gene Ontology Jaccard Index (GOJI) network. The first two columns

contain gene IDs, while the third column shows the GOJI value. Only gene pairs above the

significance threshold are shown.

A Neighbor counting

Most frequent term ③
Outcome: Incorrect

Most frequent terms ② ② ②
Outcome: Correct
1 false positive: ●

Most frequent terms ② ②
Outcome: Correct
0 false positives

Most frequent terms None
Outcome: Incorrect

B STRING Biological Process

C Biological Process

A Query gene-centric search

- Entry: gene ID (e.g. At4g32410) or keyword (e.g. cellulose)
- Answers: Which other genes have a similar function?
- What is the function of my gene?

i) **GOJI table:** Identify genes with similar function

GOJI value	Gene ID
0.308	at5g64740
0.302	at5g05170
0.293	at4g39350
...	...

ii) **GOJI network:** Visualize GOJI relationships between the functionally related genes

Functionally related gene (small node)

Query gene (large node)

GOJI edge

iii) **GO predictions:** Show which GO terms are associated to the query gene

# evidences supporting this function	P-value	GO ID	Description	Experimental evidence	Network A	Network B
3	0.01	GO:0000271	...	✔	✔	✔
2	0.03	GO:0005975	...	✔		✔
2	0.06	GO:0030244	...		✔	✔
1	0.13	GO:0044238	...		✔	

B Gene Ontology-centric search

- Entry: GO ID (e.g. GO:0000271) or keyword (e.g. cellulose)
- Answers: which genes have this function?

Gene table: Reveal genes assigned to the term of interest

Gene ID	# evidences supporting this function	Experimental evidence	Network A	Network B
A	3	✔	✔	✔
B	2		✔	✔
C	2	✔	✔	
D	1		✔	

C Gene set-centric search

- Entry: multiple gene IDs
- Answers: which genes are functionally related to the entered genes?

i) **GOJI intersection network:** Visualize genes connected to the queries.

ii) **Gene table:** Show genes connected to the queries

Gene ID	# of queries connected to this gene	Q1	Q2	Q3
E	3	✔	✔	✔
B	2	✔		✔
C	2		✔	✔