Human macrophages survive and adopt activated genotypes in living zebrafish

Colin D Paul[1][#], Alexus Devine[1][#], Kevin Bishop[2], Qing Xu[3], William J Wulftange[4], Hannah Burr[1], Kathryn M Daly[1], Chaunte Lewis[1], Daniel S Green[5], Jack R Staunton[1], Swati Choksi[3], Zheng-Gang Liu[3], Raman Sood[2], and Kandice Tanner[1][*]

[1] Laboratory of Cell Biology, Center for Cancer Research, National Cancer Institute, National Institutes of Health

[2] Zebrafish Core, National Human Genome Research Institute, National Institutes of Health

[3] Laboratory of Immune Cell Biology, Center for Cancer Research, National Cancer Institute, National Institutes of Health

[4] National Institute of Biomedical Imaging and Bioengineering, National Institutes of Health

[5] Women's Malignancy Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health

[#] co-first authors

[*] To whom correspondence should be addressed: Kandice Tanner, Ph.D., 37 Convent Dr., Bethesda, MD 20852. Email: kandice.tanner@nih.gov

Abstract

The inflammatory response, modulated both by tissue resident macrophages and recruited

monocytes from peripheral blood, plays a critical role in human diseases such as cancer and

neurodegenerative disorders. Here, we sought a model to interrogate human immune behavior *in*

vivo. We determined that primary human monocytes and macrophages survive in zebrafish for up

to two weeks. Flow cytometry revealed that human monocytes cultured at the physiological

temperature of the zebrafish survive and differentiate comparable to cohorts cultured at human

physiological temperature. Moreover, key genes that encode for proteins that play a role in tissue

remodeling were also expressed. Human cells migrated within multiple tissues at speeds

comparable to zebrafish macrophages. Analysis of gene expression of *in vivo* educated human

macrophages confirmed expression of activated macrophage phenotypes. Here, human cells

adopted phenotypes relevant to cancer progression, suggesting that we can define the real time

immune modulation of human tumor cells during the establishment of a metastatic lesion in

zebrafish.

Keywords

Zebrafish, immune cells, live imaging, flow cytometry, human xenograft, cell migration, brain

microenvironment, activated phenotype, gene expression

Macrophages represent a mature population of terminally differentiated cells of myeloid-lineage

found in all tissues(1, 2). They are often categorized by distinct functional properties, cell surface

markers, and the cytokine profile of the microenvironment. Highly plastic, macrophages adopt

diverse phenotypic and functional states to regulate tissue homeostasis, tissue patterning,

branching morphogenesis, wound repair and immunity(2). They respond to environmental cues

within tissues such as damaged cells, activated lymphocytes, or microbial products to differentiate

into distinct functional phenotypes(3). However, macrophages may adopt functions that aid and

promote disease due to environmental cues that arise as a result of abnormal physiological states

such as obesity, fibrosis, brain neurodegenerative disorders and cancer(1, 4-7). In particular, one

of the "hallmarks" of cancer and predictors of aggressive metastatic disease is the chronic presence

of activated myeloid cells, such as tumor associated macrophages (TAMs), within primary

tumors(8-10). Probing the role of the inflammatory response in the earliest stages of malignant

transformation remains technically and ethically difficult in human subjects. Nevertheless, the

broad importance of immune cell biology necessitates appropriate models to adequately study

implications in human disease.

A number of efforts have been made to "humanize" animal model systems to study human

homeostasis and disease *in vivo*, especially in the context of blood cancers(11-17). However, the

role of hematopoietic stem cells (HSCs) and their derivative lineages, such as myeloid cells, have

also been identified as regulators of tumor progression for solid cancers(4, 8). There are several

classes of macrophages that may be found around tumors(2, 10, 18). One class of macrophages

includes the classical "M1" macrophage phenotype, which is characterized by the production and

secretion of elevated levels of pro-inflammatory cytokines, mediation of resistance to pathogens

with strong microbicidal properties, high production of reactive nitrogen and oxygen

intermediates, and evocation of Th1 responses. In contrast, M2 macrophages are characterized by

their involvement in tissue remodeling, immune regulation, tumor promotion and efficient

phagocytic activity(2, 10). A second class of macrophages that mediate metastasis, metastasis-

associated macrophages (MAMs) is thought to drive therapeutic resistance and establish

macrometastases at distant sites(19, 20). However, probing the role of the inflammatory response

on the etiology of metastasis, namely when one or few cells have initiated the formation of a distant

lesion in vivo, is challenging. Of all the animal models of oncoimmunology, mice remain the

standard model, as one can interrogate human cells using immune-compromised strains and

similarly examine syngeneic cancer cell lines in the appropriate immune competent strain(11, 16,

21). However, directly visualizing the colonization of multiple murine organs is expensive as it

requires many mice to achieve sufficiently powered statistics. Simply put, a model system that

recapitulates physiologically relevant aspects of metastatic disease in which cells can be observed

at the single-cell level would profoundly benefit metastatic cancer research(22).

Zebrafish larvae are optically transparent at larval stages and conducive for non-invasive

imaging *in vivo*. This, coupled with the fact that many processes involved in embryonic

development, cell biology, and genetics are conserved across vertebrates, makes zebrafish an

attractive model to study tumor-immune interactions during early stages of organ colonization(23,

24). In addition, organ architecture, composition and regulatory signaling networks for several

organs, including the brain, are well conserved in embryonic zebrafish(25, 26). As early as the first

few days after fertilization, key components of the Central Nervous System (CNS) such as

neurons, astrocytes, and microglia, as well as specialized structures such as the blood-brain barrier

and choroid plexus, have been identified in larval zebrafish. The innate immune system, comprised

primarily of neutrophils, tissue specific macrophages and peripheral derived

monocytes/macrophages, is also conserved at this early developmental stage (15, 25, 26). Hence,

the embryonic zebrafish offers a system in which cell dynamics can be observed while sharing key

cellular and structural characteristics with the mammalian organs. This model can be used to

delineate tissue resident vs. recruited circulating monocytes in the context of inflammation and

tumor education for either primary brain tumors or in the case of early metastatic colonization.

However, a key feature missing in studies of immune response in zebrafish is a careful

demonstration that human immune cells can be introduced in this system with conserved survival

and cell response.

Here, we introduced human monocytes/macrophages into the zebrafish, both directly into

circulation and in an organ-specific manner. We determined that both monocytes and macrophages

(differentiated in culture) survive in innate immune competent zebrafish for up to two weeks post-

injection despite the lower physiological temperature of the zebrafish (28.5°C). Similar results

were obtained *in vitro*, where flow cytometry and RNAseq analysis revealed that human

monocytes cultured at the physiological temperature of the zebrafish survive, differentiate, and

show surface markers associated with mature macrophages in response to cytokines comparably

to cohorts cultured at human physiological temperature (37°C). We also determined that human

cells are transformed by zebrafish astrocytes by employing transgenic fish where fluorescently

tagged astrocytes can be visualized *in vitro* and *in vivo*. Gene expression analysis of *in vivo*

educated human macrophages revealed gene expression associated with activation. In summary,

these results characterized the function of human immune cells in the *in vivo* environment and

physiological temperature of *Danio rerio*. Thus, this model system allows for examination of the

contribution of specific immune cells within specialized organ microenvironments.

Results

Zebrafish are reared at the lower temperature of 28.5ºC compared to 37ºC, the

physiological temperature of mammals(27). Zebrafish also have an innate and adaptive immune

response beginning from two and nine days post fertilization (dpf), respectively. We thus asked if

human immune cells can survive at the physiological temperature of zebrafish in the presence of

the innate and immature adaptive immunity in an organ specific manner. Human primary

monocytes that were differentiated into macrophages in culture were injected into the hindbrain of

the zebrafish. Longitudinal imaging of the same fish one day and 7 days post injection (dpi; 3 dpf

and 9 dpf, respectively) revealed that human macrophages survive for at least one week after

injection and can be found dispersed in the brain **(Figure 1a-c).** Similar examination of human

macrophages injected into zebrafish where the host microglia are fluorescently tagged revealed

survival and distribution within the brain **(Supplementary Figure 1a).** We next examined the

survival of an immortalized human monocyte cell line under the same conditions. Similar survival

and morphologies were observed within the brain (**Supplemental Figure 1b**). Examination of

zebrafish two weeks post injection revealed that a few human macrophages can persist *in vivo* up

to this time point **(Figure 1d, Supplemental Figure 1c).** We next asked if the human cells

migrated within the parenchyma when directly injected into the brain. Serial imaging revealed that

human macrophages were widely dispersed within the zebrafish brain and were often found in

close vicinity to blood vessels (**Figure 2**). As immune cells are involved in tissue remodeling and

surveillance, we next asked if the introduced human cells show comparable motilities. We

quantified the host immune cells movement by tracking neutrophils and macrophages in addition

to introduced human monocytes within the brain at 3 dpf. We determined that neutrophils move

with an average speed of 7 microns/minute, which was significantly greater than that of either the

zebrafish brain resident macrophages (P<0.0001) and the introduced monocytes (P<0.0001). There

was no significant difference between the slower monocytes/zebrafish macrophages (P=0.8263),

where an average speed of 0.5-1 microns/minutes was observed **(Supplemental Figure 2).**

138   Immune cells are able to move from circulation to different tissues. Therefore, we asked if

human cells would adopt similar phenotypes and survive in zebrafish upon access to the circulatory

system. Human macrophages differentiated at either 28.5°C or 37°C and then injected directly to

the zebrafish circulation were found throughout the entire zebrafish as early as ~3 hrs post injection

and 1 day post injection **(Supplemental Movies 1-2, Figure 3a-c).** They were observed to be

motile and moved within intersegmental vessels and within different tissues **(Supplemental**

Movies 1-2). Imaging of zebrafish at one day and 7 days post injection (3 dpf and 9 dpf,

respectively) revealed that human macrophages survive for at least one week after injection into

the circulation and can be found dispersed in multiple tissues, including the gills, intersegmental

vessels **(Figure 3d)**, and caudal tissue (data not shown)**.** Interestingly, a subset of primary human

macrophages injected to the zebrafish circulation at 2 dpf can disperse through the zebrafish and

be found in the head and tail at 14 days post injection (**Figure 3e**).

150   Macrophages represent a class of terminally differentiated monocytes, and we reasoned

that survival may differ as a function of differentiation status when cultured at the physiological

temperature of zebrafish. Thus, we directly compared survival of human monocytes cultured at the

two different temperatures. A subset of human monocytes were injected into the hind brain of the

zebrafish, while others were cultured in vitro at either 28.5°C or 37°C. We imaged larva over the

course of 3 days post injection at 24 hour intervals **(Figure 4a).** Analysis of zebrafish imaged

serially over the course of 1 day intervals revealed that 25-50% of cells survived in multiple

transgenic lines after 3 days post injection (3 dpi) **(Figure 4b).** There was no regional preference

for survival as cells were observed throughout the fore, mid and hind brain of each animal

(**Supplemental Figure 3a,b**). Additionally, the zebrafish strain into which monocytes were

injected was not a significant factor on monocyte survival (P=0.4917, **Supplementary Figure**

3c).We determined that a similar percentage of monocytes cultured at the zebrafish physiological

temperature survived in vitro. Approximately 40% of monocytes from de-identified human donors

survived at the zebrafish physiological temperatures and were viable up to 7 days of *in vitro* culture

in only culture medium. In contrast, these same cells cultured at mammalian physiological

temperatures showed rapid cell death and viability, where ~50% of cells underwent apoptosis after

2 days of *in vitro* culture in culture medium and only 10% of cells were present 7 days after plating

(Figure 4c).

We then investigated whether these human cells cultured at zebrafish physiological

temperature are capable of differentiating into functional macrophages. Human monocytes

differentiate into macrophages in response to *in vitro* addition of human macrophage colony

stimulating factor (H-M-CSF)(28). We observed that monocytes cultured in the presence of H-M-

CSF at the lower temperature were not adherent and remained spherical in suspension (**Figure 5a**).

This spherical morphology was similar to that seen in control monocytes cultured without addition

of H-M-CSF at both temperatures. In comparison, cells that were cultured at the higher

temperature adopted the classical morphology of adherent elongated cells following treatment of

H-M-CSF for five days **(Figure 5a).** Immune cells respond to cytokines secreted by other cell

types. To determine if monocytes/macrophages respond to zebrafish stroma, we co-cultured these

immune cells with zebrafish astrocytes using transwell assays. Quantitation of cells that migrated

in transwell assays revealed that a greater fraction of cells migrated in response to zebrafish

astrocytes for each culture condition than under control conditions **(Figure 5b).** We determined

that monocytes cultured in the presence or absence of differentiation media showed an increased

survival at the zebrafish physiological temperatures, where ~40% of cells without H-M-CSF, and

~70% of cells with H-M-CSF, were viable up to 5 days of *in vitro* culture. In contrast, these same

cells cultured at mammalian physiological temperatures showed reduced viability when cultured

with or without differentiation media **(Figure 5c).** Macrophages derived from monocyte

precursors adopt specific phenotypes depending on cues from the local tissue environment. We

next analyzed cell surface markers using flow cytometry to monitor macrophage maturation. Cells

that were cultured with H-M-CSF at both temperatures showed expression for the surface marker

CD14 **(Figure 5d)**(29). This surface protein, CD14, is an indicator of macrophage maturation.

Surface marker expression for differentiated human macrophages is observed after 5, 8 and 11

days of exposure for both physiological temperatures. However, when cells are cultured at the

lower temperature, there is a temporal delay in the expression of CD14 compared to cells cultured

at the higher temperature, and a subset of these cells remain CD14-negative **(Figure 5d).**

Monocytes that are subjected to M-CSF will differentiate into macrophage populations marked by

expression of CD86, CD163 and CD206(30, 31). We thus asked if monocytes differentiated at the

lower temperature will also show functional plasticity. Cells positive for CD14 expression were

then assessed for markers, CD86, CD163 and CD206 at each temperature. Surface marker

expression for differentiated human macrophages is observed after 5, 8 and 11 days of exposure

for both physiological temperatures. However, when cells are cultured at the lower temperature,

there is a temporal delay in the expression of CD86, CD163 and CD206, compared to cells cultured

at the higher temperature **(Figure 5e).** Functionally, monocytes differentiated into macrophages at

the lower temperature were motile *in vivo* and showed similar motility to macrophages that were

differentiated at the higher temperature **(Supplemental Movies 1 and 2).**

For the first three weeks of development the zebrafish innate immune system is the primary

defense against invading pathogens. This is in contrast to mice, which have circulating T cells at

7 days post-partum(32, 33). Hence, we examined the innate immune response to the introduced

human cells. Neutrophils are conserved in the axis of innate immunity in zebrafish(23). They form

the first line of defense and produce a sustained inflammatory response following introduction of

foreign entities. In the brain, the resident macrophages are the microglia(26). Innate zebrafish

neutrophils and macrophages are present throughout 3 dpf zebrafish, including in the brain (**Figure**

6a,b). To determine if there was a differential zebrafish neutrophil or microglia response,

differentiated human macrophages cultured at each temperature were injected into the hindbrain

of 2 dpf zebrafish where either host neutrophils/ macrophages and vasculature were fluorescently

labeled (**Figure 6c**). Serial imaging of the same zebrafish one day and 7 days post injection (3 dpf

and 9 dpf, respectively) revealed that macrophages differentiated at each temperature survive for

at least one week after injection and can be found dispersed in the zebrafish brain as observed in

previous transgenic lines **(Figure 6d-f).** We determined that similar numbers survived in either

line **(**two-way ANOVA with one degree of freedom, P=0.4575 for interaction, P=0.3072 for

temperature effect, P=0.4822 for zebrafish line effect; **Figure 6d),** with survival rates comparable

to those observed in previous experiments. Injected macrophages adopt elongated and spherical

phenotypes and are seen along blood vessels and in the brain parenchyma **(Figure 6,**

Supplemental Figure 4a,b). No swarming of neutrophils was observed in the vicinity of the

human monocytes at 1 day or 7 days post injection. Micrographs show that a few neutrophils are

in close proximity to human macrophages, and neutrophil distributions are similar to those in

uninjected embryos. We observed some co-localization between the host and human macrophages

as determined by overlapping in the micrographs of the zebrafish imaged 1 day post injection.

However, very little overlap is observed one week post injection.

In addition to tissue surveillance, macrophages remodel the tissue microenvironment.

Thus, we asked if genes that encode for the matrisome were similarly expressed following in vitro

differentiation at 28.5°C or 37°C. RNAseq revealed that 295/474 matrisome proteins were

similarly expressed, with 179 others having statistically significantly different log10(normalized

count) numbers. Notably, proteoglycan 2 was expressed at higher levels for macrophages

differentiated at the physiological temperature of zebrafish. Moreover, examination of a cohort of

genes associated with macrophage polarization revealed similar trends, with 71/164 genes

similarly expressed and 93 having statistically significantly different log10(normalized count)

numbers. Il-6 was expressed at higher levels for macrophages differentiated at the physiological

temperature of zebrafish. Macrophages can polarize into different phenotypes in response to cues

from the local tissue microenvironment(29). Thus, we compared gene expression of macrophages

exposed to known cytokines in vitro at the two temperatures. We determined that similar

expression of TNF-α, CD163 and VEGF for human macrophages cultured *in vitro* in the presence

of recombinant EGF and TNF-α for 24 hours at both physiological temperatures. However, there

was a greatly reduced expression of CCL18 for cells cultured at physiological temperature of

zebrafish than at physiological temperature of humans **(Figure 7c).** We then compared gene

expression of human cells that were differentiated and then injected in the zebrafish brain for 24

hours prior to RNA purification.sfnalysis revealed similar expression of TNF-α and CD163 for

human macrophages cultured in vitro at both physiological temperatures prior to injection.

However, there was a reduced expression of VEGF, CCL17, CCL18, and CD24 for cells cultured

at physiological temperature of zebrafish compared to physiological temperature of humans,

though these differences were not significantly different. In contrast, IL 6 was expressed at higher

levels in cells cultured at the physiological temperature of zebrafish than cells cultured at

physiological temperature of humans, confirming the in vitro RNAseq result. **(Figure 7d).**

Discussion

Here, we introduce components of the human innate immune system to the zebrafish

microenvironment by introducing human monocytes/macrophages. We first determined that

human monocytes differentiate into functional macrophages at the physiological temperature of

zebrafish. We showed that these human monocytes are able to survive for at least two-weeks post

injection in vivo in immune competent zebrafish. Specifically, in transgenic zebrafish where

fluorophores coupled to promoters of either myeloid-specific peroxidase (*mpx)*, a homologue of

myeloperoxidase, or macrophage expressed gene 1 (*mpeg1*), allow specificity in distinguishing

neutrophils from macrophages despite their common lineage. We used this distinction in lineage

to analyze the survival and response of human immune cells introduced to embryonic zebrafish,

demonstrating that human monocytes and macrophages injected into the zebrafish hindbrain or

circulation survive and are functional for up to two weeks post-injection.

Introduction of human hematopoietic stem cells (HSCs) and hematopoietic cancers have

been successfully performed in fetal sheep, immunocompromised mice, and zebrafish(10, 11, 16,

17, 24). In these animal models, the HSCs grafted successfully to the appropriate in vivo niche,

bone marrow in mammals and kidney in the zebrafish, with varying lengths of survival in vivo.

Moreover, the HSCs retained the ability to differentiate into multiple blood lineages, allowing the

study of normal tissue homeostasis. This has greatly assisted in delineating what goes awry during

transformation to cancers of the blood. Tissue grafting becomes more complicated in the case of

cross species transplantation due to the mismatch in immune compatibility (11, 34). One solution

involves sub-lethal irradiation of immune competent animals allowing for a temporal window

whereby the donor components can be introduced (11, 34). However, there may be adverse and

off-target effects due to the depletion of the recipient immune response. The zebrafish innate

immune system comprised primarily of neutrophils and macrophages shows excellent

conservation with mammals (15, 25). Here, we show that human monocytes/macrophages can

survive in vivo in the larval zebrafish in the presence of innate immunity and immature adaptive

immunity (**Figures 1-4, Supplemental Figures 1-3**). Interestingly, the introduced macrophages

migrate through multiple tissues mimicking tissue surveillance, a key aspect of macrophage

function **(Supplemental Movies 1 and 2).** Furthermore, they co-exist seemingly unmolested by

host neutrophils and macrophages **(Figure 6, Supplementary Figure 4).** Thus, there is sufficient

conservation in signaling proteins that regulate immune survival. This suggests that this model

organism can provide a comparable study of metastases in the same manner as observed in mice

between host tissues of zebrafish and human tumor cells.

In adult mammals, innate immune cells, monocytes and macrophages share a common

myeloid progenitor in the bone marrow (2, 10, 11, 35, 36). These cells differ greatly in their

lifespan, where peripheral blood derived monocytes (PBDM) survive just a few days before

undergoing apoptosis. In contrast, differentiated monocytes such as macrophages may have a

lifespan of months. While these immune subsets are conserved in zebrafish innate immunity, the

major difference is that they exist at different physiological temperatures. This study showed that

human monocytes also survive on the order of days at the physiological temperature of zebrafish

and surprisingly, with a greater survival than a cohort cultured at the physiological temperature of

human in vitro and in vivo **(Figure 4).** Monocytes respond to different chemokines following a

strictly regulated schedule (2, 10, 11, 35, 36). Following dissemination from the bone marrow,

circulating monocytes are CCR2 negative. One example is that they become CCR2 positive

roughly 1- 2 days post release and successive tissue extravasation. Functional analysis revealed

that human monocytes differentiate into M1 and M2 macrophages as dictated by high CD86,

CD163 and CD206 positive expressions at both temperatures in a similar time frame. However,

monocytes cultured at the lower temperature differentiate into mature macrophages with a

temporal delay compared to monocytes culture at human temperatures. In addition, the cells

displayed differences in morphology, where cells cultured at zebrafish physiological temperature

remained in suspension and did not adhere, indicating that there are subtle differences due to the

mismatch of culture temperature. One reason could be reduced metabolism at the lower

temperature, or that the human cells follow the differentiation steps that zebrafish

monocytes/macrophages adopt. Nevertheless, the functional status is achieved **(Figures 5 and 7).**

Within a growing tumor, resident and recruited monocytes receive chemical cues from

tumor-secreted chemokines and physical cues from the microenvironment such as hypoxia(37).

Consequently, they become tumor-educated and adopt phenotypes that help drive tumor growth

through promotion of neoangiogenesis, aberrant tissue remodeling, and development of

immunosuppressive microenvironments(2, 4, 8). These tumor-associated macrophages (TAMs)

are abundantly found within or adjacent to primary tumors in several types of human cancer.

TAMs are regulated in part by colony stimulating factor (CSF)-1, mediated by the CSF-1 receptor

(CSF1R), which drives macrophage maturation, tissue recruitment, and activation (38). Activated

macrophages in turn release epidermal growth factor (EGF). At the primary tumor, this signaling

pathway leads breast cancer cells and macrophages to form complexes that promote intravasation

into the bloodstream (21, 39). There are several classes of macrophages that may be found around

tumors. M1-type macrophages are thought to antitumorigenic and are characterized by high CD80,

MHC-II, and PDL1 expression and low CD163 and CSF1R expression. In contrast, macrophages

with high expression of CD68, CD163, CD206, and CSF1R are tumorigenic M2-type

macrophages(20, 40) (18). Macrophages have been implicated in the clearance of circulating

tumor cells during early metastatic dissemination (41, 42). Importantly, another class of

macrophages, termed metastasis-associated macrophages (MAMs), may mediate metastasis,

though the origins and recruitment of such macrophages are unclear(19, 20, 43). Here, we

determined that in response to recombinant cytokines and brain-conditioning in vivo, human

macrophages adopt gene expression that forms part of the identification of M1 and M2 phenotypes

as measured by TNF-A, CD163 and VEG-F expression **(Figure 7).** M2 macrophages have been

classified into subdivisions, a, b and c(44, 45). The M2a subtype can also be defined as alternative

activated macrophages, the M2b as type 2 macrophages, and the M2c as deactivated macrophages.

The deactivated terminology refers to the in vitro ability of macrophages to adopt M2 activation

following M1 activation, thus deactivating the M1-like gene transcription. Interestingly, CCL18 is

differentially expressed as a function of culture conditions in vivo and in vitro **(Figure 7).** CCL18

production is associated with the M2c subdivision for human monocytes and macrophages(45,

46). These data suggest that human monocytes differentiated at the lower temperature may not

show the plasticity to switch between phenotypes compared to the macrophages differentiated at

the human temperature. Interleukin-6 is a pleiotropic cytokine overexpressed in response to injury,

inflammation, and infection(47). IL-6 is produced by many cells including osteoblasts, monocytes

and macrophages(48). In our system, we observe that macrophages differentiated at lower

temperatures express higher expression than macrophages differentiated at higher temperatures.

Furthermore, this effect persists when cells are introduced in vivo. Interleukin-6 has a direct

growth stimulatory effect on many tumor cells through the activation of several signaling

pathways(48). Thus, in future studies, it may be more desirable to use human macrophages

cultured at the physiological temperature of humans for co-injection of human tumor cells and

macrophages.

Brain metastasis remains a challenge in clinical settings as patient survival is still measured

in weeks or months (49). Emerging data indicate that immune-mediated signaling plays an

important role in the establishment of brain metastasis (50). However, what is less understood is

the role of tissue specific microenvironment of the immune cells at early stages of colonization

restricted to lesions well below the current methods of detection. In a spontaneous mouse model

of melanoma, not only were disseminated tumor cells detected before clinically detected primary

tumors, it was determined that metastatic growth at distal organs was regulated by a tissue specific

immune response (51). Recently, in studies of mouse models of brain malignancies and confirmed

in patient data, that in addition to tissue resident macrophages microglia, an abundance of bone

marrow derived monocytes (BDMs) can be found even in an immune privileged organ such as the

brain (52). Furthermore, they uncovered that the resultant tumor associated populations were

distinct based on the macrophage ontogeny within the brain. Here, we determined that human

macrophages survive and become activated in the zebrafish brain environment. This model allows

for examination of the contribution of immune signaling in a tissue specific manner. These findings

open up the possibility of studying early metastatic events of human cancers in zebrafish. More

importantly, it allows us to add the complexity of human tumor-human immune interactions whilst

maintaining single cell resolution.

Methods

Animal studies

Animal studies were conducted under protocols approved by the National Cancer Institute, and the

National Institutes of Health Animal Care and Use Committee. In our experiments, we employed

several transgenic lines, i.e. Tg(mpeg1:GFP), Tg(gfap:GFP) and Tg(mpx:GFP),

Tg(kdrl:GFP)la116 and Tg(kdrl:mCherry-CAAX)[y171], kindly provided by Brant Weinstein

(NICHD, Bethesda, MD), and the Casper strain, a kind gift from David Langenau (Harvard

MGH)(53-55). Dual labeled vasculature and immune/astrocyte cells were generated by crossing

Tg(kdrl:mCherry-CAAX)[y171] and GFAP:GFP, mpeg:GFP or mpx:GFP. Progeny were screened

using fluorescence microscopy to identify those that expressed fluorescent markers for vasculature

and astrocytes, macrophages and neutrophils respectively. From this pool, founders were selected

for continuation of the line. Zebrafish were maintained at 28.5°C on a 14-hour light/10-hour dark

cycle according to standard procedures. Embryos obtained from natural spawning, raised at 28.5°C

and maintained in egg water containing 0.6 g sea salt per liter of DI water were checked for normal

development. At day 5 regular feeding commenced.

Primary human monocytes- Maintenance and Macrophage differentiation protocol and

functionality assays

Elutriated monocytes from 8 de-identified healthy human donors were acquired from the NIH,

Department of Transfusion Medicine (DTM) (according to NIH protocol 99CC0168). Donors

within the age group of 21-50 of either sex were selected for all experiments. Primary monocytes

were cultured in DMEM high glucose, 10% fetal bovine serum and 1% penicillin/streptomycin.

Cells were either cultured at physiological temperature of humans (37°C) or at physiological

temperature of zebrafish (28.5°C). These cells were cultured at a concentration of 1.1×10^6 cells

/cm^2 for minimum of five days to facilitate macrophage differentiation in the presence of

recombinant human M-CSF (25 ng/ml). Media was replenished every 3 days. Samples were

prepared for a minimum of three technical replicates for each donor sample. For comparison, a

human monocyte cell line, U937, was cultured in suspension in RPMI media (ThermoFisher

Scientific, Waltham, MA) supplemented with final volume of 10 % (v/v) Fetal Bovine Serum

(FBS), Penicillin (100 U/mL) and Streptomycin (100 µg/mL). Cells were cultured at 37°C with

5% CO_2 and relative humidity maintained at 95%. Media was refreshed every 2-3 days.

Primary monocytes that had been cultured in the presence of human M-CSF for eight days were

cultured at each temperature (28.5 vs. 37°C). Cells cultured at each temperature were then cultured

for an additional 24 hrs with a vehicle control and in the presence of either recombinant EGF at a

concentration of 10 ng/ml or TNF-α at a concentration of 20 ng/ml. In addition, 2-5 nL of a sub

set of the originally differentiated cells were then injected into 2 dpf fli:GFP embryos at a

concentration of 1×10^7 cells/mL for comparison. After 24 hrs, RNA was extracted for the *in vitro*

educated macrophages. Similarly, RNA was extracted from pooled samples of 50 3 dpf embryos

for each temperature (28.5 vs. 37°C) and age matched uninjected embryos.

Quantification of in vitro monocyte survival at different physiological temperatures

For cell survival, human monocytes from 5 donors were seeded at a concentration of 1.1×10^6 cells

/cm^2 and cultured at either 28.5°C or 37°C in base media. Human monocytes were collected and

centrifuged at 1000 rpm for 5 minutes. Supernatant was removed and cells were resuspended in

1mL of sterile $1\times$ PBS. Cells were collected at 2, 3, 5 and 7 days after plating and analyzed for cell

number and viability using Nexcelom Bioscience Cell Counter. Technical triplicates were

collected for each donor sample for each temperature and day to obtain an average fraction survival

per donor. Data were then displayed using GraphPad Prism 7.0a to graph cell number and viability

as a function of days and temperature. Two-way ANOVA was used to determine statistical

significance. Bright field and epi-fluorescent images were obtained on a Zeiss Axiovert 200 with

an AxioCam MRm, using AxioVision LE 4.8.2 software. Images were acquired using 10× air

objective with numerical aperture, 0.25, where each individual image had lateral dimensions of

1388×1040 square pixels corresponding to 0.448×0.335 mm^2.

Quantification of in vivo monocyte survival

Image processing was performed via ImageJ. Spectral crosstalk was removed by subtracting

contributions of the red channel from the far-red detection channel. Images were then median

(radius of 3 pixels) and Gaussian (sigma of 2) filtered to minimize background noise. Binary masks

of monocytes were generated using Huang and Otsu thresholding methods on corrected images.

The volume occupied by monocytes was then calculated using ImageJ's built in 'analyze particles'

function and the 3D shapes plugin. Number of monocytes was estimated by dividing the total

volume of the binary monocyte mask by the expected average volume of a human monocyte (10

μm in diameter, volume of 104.7 μm^3). Loss of monocytes was calculated for each larva by

subtracting final number of monocytes from the initial number of monocytes (24 hours post

injection) then normalizing by the initial number of monocytes. Percent survival was reported as

1 minus loss of monocytes multiplied by one hundred.

As an example: **[100* (1- ((#24hpi - #48hpi) / #24hpi)]**

Data were then displayed using GraphPad Prism 7.0a to graph cell number and viability as a

function of days. All error bars are 95% confidence intervals where D'Agostino and Pearson

normality test was performed prior to either unpaired t-test or Kruskal-Wallis test to determine

statistical analysis.

Flow cytometry to determine cell surface expression to determine phenotype

Cells cultured in the absence and presence of human M-CSF were processed at minimum five days

after plating for flow cytometry. Cells that remained in suspension were collected whereas

adherent cells were detached with StemPro Accutase (ThermoFisher). For both conditions, the cell

suspension was centrifuged and the supernatant removed as previously described. The cell pellets

were re-suspended in ice-cold PBS with the following antibodies to assess surface expression for

determination of macrophage polarization. Cells were incubated on ice with CD14-FITC (BD),

CD86-PE (BD), and CD206-APC (BD), or with CD14-FITC, CD163-PE, and CD206-APC

antibodies for 1 hour and cells were washed twice before processing for flow cytometry. Unstained

cells and cells stained with single antibodies were used for gating cell size and setting instrument

parameters. Samples were processed on a BD FACSCanto II (BD Biosciences) and analyzed with

FlowJo software (Tree Star). PE- or APC-conjugated IgG stained cells were used to indicate

background fluorescence and to set quadrants before calculating the percentage of positive cells.

Fluorescence-activated cell sorting (FACS).

Embryos at 2 days post fertilization (dpf) with labeled vasculature and astrocytes cells were

dechorionated in pronase solution (Roche Life Science #10165921001) as necessary (less than

5minutes). Embryos were then protease dissociated for FACS as previously described(56). Cell

sorting was performed on a FACSAria Fusion instrument (BD Biosciences, San Jose, CA) using

85 µm nozzle at 45 psi pressure. Cells were gated on forward and side light scatter and separated

based on GFP and/or mCherry-positive fluorescence signals defined with non-fluorescent negative

controls.

In vitro **migration assays.**

Astrocytes isolated from FACS were centrifuged at 1000 rpm for 5 minutes, resuspended and

counted. A 24-well plate was coated with 10 µg/ml laminin (Cytoskeleton Inc.) at room

temperature for 1 h prior to triplicate wash with PBS. Fluorescence-activated cell sorted astrocytes

were then added to the wells at 1 x 10^5 cells per mL (1 ml/well in 24-well plate). Zebrafish

astrocytes were maintained overnight in DMEM (Thermo Fisher) supplemented with 10% fetal

bovine serum, 1% penicillin-streptomycin and 1% L-glut at 28.5°C. The following day, cells were

prepared for transwell assay. Astrocytes were washed once with serum free RPMI medium. 700µl

of serum-free media was placed on top of astrocytes. A sterile transwell membrane with 8 µm

pores (Corning #3422) was inserted and 300µl of human macrophages were placed on top of

membrane at 5 x 10^4 cells per membrane. Transwells were incubated overnight at 33°C. Three

membranes were run in parallel per condition (temperature and presence/absence of zebrafish

astrocytes). Transwells were fixed in 4% paraformaldehyde for 10 minutes at room temperature.

Membranes were washed twice in 1X PBS. They were permeabilized using 0.1% triton-x 100 for

5 minutes at room temperature, followed by a second wash step and then processed for

immunofluorescence. Staining mixture consisted of Hoechst 33258 (10mg/mL, Thermo Fisher)

was used at 1µg/mL in 1% BSA-PBS along with Phalloidin 565 (Sigma) used at 20µl/mL of

Hoechst mixture. The membranes were imaged using Zeiss 780 LSM to acquire 5 x 5 tile scan

images of each membrane, using a 10× air objective of 0.3 NA where each individual image

comprised 2046×2046 square pixels corresponding to 1416×1416 square µm for a total Z

distance of 100 µm with 5 µm step size. Three randomly selected 500 x 500µm fields of view were

used for data analysis. We simultaneously excited our sample with the 405 nm from an argon ion

laser with a power of < 3 % (total power 30 mW) and 546 nm from a solid-state laser (power

of < 10 %). A secondary dichroic mirror, SDM 560, was employed in the emission pathway to

delineate the red (band-pass filters 560–575 nm) and blue channels (480-495 nm) at a gain of 400

on the amplifier. The laser power for the 543 nm setting was set at < 3 % of the maximum power

and the gain on the detectors was set to 450. The Hoechst-stained cell nuclei were counted. The

lower Z represents the bottom of the membrane and higher Z represents the top of the membrane.

Nuclei counted as through were on the bottom or in pores. Nuclei counted on top were used in

total cell count. Data were displayed using GraphPad Prism 7.0a to graph cell migration with

given attractants. Two-way ANOVA with Tukey's multiple comparisons post-test among all

conditions was used to determine statistical significance.

Zebrafish injections

For all experiments, at 24 hours post fertilization (hpf), embryos were transferred to egg water

supplemented with phenylthiourea (PTU, Sigma P5272), suspended at 7.5% w/v in DMSO, at 1

part in 4500 to inhibit melanin formation for increased optical transparency. Embryos were then

returned to the incubator at 28.5°C and checked for normal development. Zebrafish embryos at 2

days post fertilization (2 dpf) were anesthetized using 0.4% buffered tricaine. Human

monocytes/macrophages were labeled with cell tracker (Green or Deep Red, Invitrogen) at a final

concentration of 1 µM for 30 minutes. Cells were then centrifuged and the supernatant removed

and pellet resuspended in PBS with a final cell concentration of 1×10^7 cells/mL before injection.

2-5 nL of labeled cells were injected into the hindbrain of the embryo. Zebrafish were then reared

at 28.5°C with 5% CO_2 and relative humidity maintained at 95% for two days. At 5 days post

fertilization, some zebrafish were then returned to system water and regular feeding at 28.5 °C for

long-term survival studies. For injection to the circulation, zebrafish were anesthetized in 0.4%

buffered tricaine and oriented in a lateral orientation on an agarose bed. Cells were injected directly

in the circulation via the posterior cardinal vein using a pulled micropipette. Zebrafish were

screened within 24 h of injection to check for successful introduction of cells to the circulatory

system

Intravital microscopy

For live cell imaging, embryos were anesthetized using 0.4% buffered tricaine, then embedded in

a lateral orientation in 1% low melting point agarose (NuSieve GTG agarose, Lonza), and allowed

to polymerize in with cover glass (no. 1.5 thickness) as previously described (57). Egg water

supplemented with tricaine was added to the agarose hydrogel for the entire time of data

acquisition. Zebrafish were imaged on Zeiss 710 or 780 laser scanning confocal microscopes.

Initial overview experiments were taken at 20×, with 1 µm Z steps, as tile scans over the entire

length and height of the zebrafish. Images in the head and tail of the zebrafish were acquired at

10× magnification every 10 min for 14 h. Z stacks were acquired using a tiled approach and a

10× air objective of 0.3 NA where each individual image comprised 2046×2046 square pixels

corresponding to 1416×1416 square µm for a total Z distance of 276 µm. One-photon, confocal,

2-dimensional images of 512×512 pixels (lateral dimensions) were acquired with a 1.4 NA

$40 \times$ oil-immersion objective. We simultaneously excited our sample with the 405 nm, 488 nm

lines from an argon ion laser with a power of < 3 % (total power 30 mW) and 546 nm from a solid-

state laser (power of < 10 %). A secondary dichroic mirror, SDM 560, was employed in the

emission pathway to delineate the red (band-pass filters 560–575 nm) and green (band-pass filters

505–525 nm) and blue channels (480-495 nm) at a gain of 400 on the amplifier. The laser power

for the 543 nm setting was set at < 3 % of the maximum power and the gain on the detectors was

set to 450.

RNA isolation, RNAseq Analysis and Real-time PCR

Total RNA was extracted with Trizol according to the manufacturer's guidelines (Invitrogen). Any

remaining DNA was removed with the DNA-free kit (Ambion) and was re-purified with the

RNAeasy kit (Qiagen). Illumina Stranded mRNA kit (Ref: 15056200) and the Illumina NeoPrep™

Library Prep System (Ref: 20003841) were used to prepare libraries from total RNA. Libraries

were then sequenced using the Illumina NextSeq500 with 75-bp paired-end reads. Sequencing

reads were trimmed for quality control using Trimmomatic (58) and quality was further verified

with FASTQC (59). Trimmed reads were then aligned to human genome 19 (hg19) using STAR

(60). Gene abundances were then determined using featureCounts (61) and normalized with TMM

(62). Differential expression between samples was then determined using EdgeR (63, 64) and

visualized with pheatmap (65). Genes associated with expression of proteins that comprise the

matrisome and macrophage polarization were examined(66, 67). Genes plotted in Figure 7 were

those from the matrisome and macrophage polarization profile for which p<0.05 and FDR<0.01.

This cut-off was reached for 179/474 matrisome genes and 93/164 polarization genes.

Taqman real-time gene expression assays were run on an ABI StepOne Plus system according to

manufacturer's protocol (Applied Biosystems). Gene expression was normalized to that of

GAPDH or β-actin. For analysis of gene expression in vitro, two biological replicates were used.

Per biological replicate, cycle thresholds for each gene were normalized against a corresponding

GAPDH replicate for each of two technical replicates, and technical replicates were averaged to

determine a mean ΔcT and fold change compared to control macrophages differentiated at 37°C

in the absence of EGF or TNFα. Results from the two biological replicates were then used to

calculate the average fold changes and standard deviations. For analysis of gene expression from

in vivo educated macrophages, cycle thresholds for each gene were normalized against a

corresponding GAPDH threshold. Two technical replicates were then used to calculate mean fold

changes and associated standard deviations. For each gene target, results were normalized to

expression for in vivo educated macrophages differentiated at 37°C. All Taqman real-time primers

used for gene-expression analysis were pre-designed and confirmed either from Integrated DNA

Technologies: human TNFa (Hs.PT.45.14765639), VEGFA (Hs.PT.58.21234833), GAPDH

(Hs.PT.39a.22214836), IL-6 (Hs.PT.58.39866843.g), CD24 (Hs.PT.58.39572682.g) or from Life

Technologies: human CD163 (Hs 00174705), CCL17 (Hs01128674).

Intravital cell tracking

Time-lapse microscopy images were exported to FIJI for analysis. To adjust for zebrafish growth

during imaging, images of zebrafish injected with U937 cells were first registered using the Correct

3D Drift plugin, with the vasculature of the zebrafish used as a topographical reference. Innate

immune cells in the head of 3 dpf Tg(mpx:GFP/flk:mcherry) or Tg(mpeg:GFP) zebrafish were

imaged every minute. Immune cells were then tracked in 3D using the TrackMate plugin for FIJI.

Images were segmented in the cell fluorescence channel in each frame using a Laplacian of

Gaussian (14) detector with a 15 µm estimated particle diameter. An initial threshold of 1.0 was

set, and the sub-pixel localization and median filter options in TrackMate were activated.

Segmentation was then further refined by manual adjustment of the threshold to minimize the non-

macrophage particles selected. Segmented objects were linked from frame to frame with a Linear

Assignment Problem (LAP) tracker with 30 µm maximum frame-to-frame linking distance. Tracks

were visually inspected for completeness and accuracy over the entire acquisition period and were

manually edited to ensure that point-to-point tracks were generated for the entire time that a cell

was in the field of view.

Calculation of Cell Speed

Temporal and spatial information for each cell track was exported to MATLAB. For each cell,

frame-to-frame speed was calculated by dividing the displacement of the cell by the time interval

between frames. An average speed for that cell over the course of imaging was then calculated by

averaging these frame-to-frame speeds. The ensemble of cell speeds for 1 zebrafish per condition

(N=28 neutrophils, N=17 macrophages, N=13 monocytes) was compared in GraphPad Prism 7

using one-way ANOVA with Tukey's multiple comparisons post-test following D'Agostino and

Pearson normality test.

Statistical Analysis

All statistical analysis was carried out in GraphPad Prism 7 (GraphPad Software, La Jolla, CA,

USA). No statistical methods were used to predetermine sample size. Zebrafish were randomly

selected from larger clutches prior to injection with human monocytes or macrophages. For

analysis of *in vitro* monocyte survival, two-way ANOVA was used to determine statistical

significance. Survival of monocytes *in vivo* for all zebrafish strains injected (monocyte survival

data grouped for all zebrafish strains tested) was assessed by successive unpaired t tests, with

significance levels adjusted by Bonferroni correction. For analysis of monocyte survival between

gfap:gfp/flk:mCherry and mpeg:GFP zebrafish, an unpaired t test was used to assess monocyte

survival following a D'Agostino and Pearson normality test. Similarly, survival of primary

macrophages *in vivo* was assessed from 3 larvae per condition (temperature and zebrafish strain)

by two-way ANOVA with one degree of freedom and Tukey's multiple comparisons post-test.

Macrophage transmigration through Transwell pores *in vitro* for three membranes per condition

(temperature and presence of zebrafish astrocytes) was assessed by two-way ANOVA with one

degree of freedom and Tukey's multiple comparisons post-test among all conditions. To compare

cell migration speeds *in vivo*, the ensemble collection of cell speeds were checked for normality

using a D'Agostino and Pearson normality test, and average speeds between conditions were then

assessed by Tukey's multiple comparisons test. To assess gene expression in vitro by PCR, two-

way ANOVA with Tukey's multiple comparisons test was used across temperatures and cytokine

treatments for each target. For PCR gene expression analysis following RNA isolation from

macrophages in vivo, expression differences were assessed by two-way ANOVA with Sidak's

multiple comparisons test for each gene at 28.5°C and 37°C differentiation temperature.

Statistically significant differences are indicated in figures with asterisks, where *, P<0.05, **,

P<0.01, ***, P<0.001, and ****, P<0.0001.

Data Availability

The authors declare that all data supporting the findings of this study are available within the paper

and its Supplementary Information. All raw images and data generated in this work, including the

representative images provided in the manuscript, are available from the corresponding author.

Acknowledgements

This research was supported by the Intramural Research Program of the National Institutes of

Health, the National Cancer Institute. We would like to thank Susan Garfield and Langston Lim,

CCR Confocal Microscopy Core Facility, Laboratory of Cancer Biology and Genetics, NCI for

use of the core microscopes. Cell sorting was performed by the NCI LGI Flow Cytometry Core .

We would also like to thank the CCR Genomics Core, Center for Cancer Research, National

Cancer Institute, National Institutes of Health. We would like to thank Ashley Williams for

assistance in preliminary monocyte differentiation experiments. We would like to thank Zeiss

microscopy for demonstration of a light sheet microscope used to generate Figure 6a,b using our

samples.

Author Contributions

K.T., Z.L, S.C, R.S., D.G. designed and discussed experiments, K.T., K.B., C.P., Q.A.D., C.L

performed experiments, K.T., C.P., A.D., K.D., Q.X., C.L, J.S, H.B., W.W. performed data

analysis. K.T., C.P., S.C. wrote the main manuscript text. K.T., C.P, Q.X., K.D., J.S., H.B., W.W.

prepared figures. All authors reviewed the manuscript.

622   **Materials & Correspondence**

623   Correspondence and material requests should be addressed to Kandice Tanner, Ph.D., 37 Convent

624   Dr., Bethesda, MD 20852. Email: kandice.tanner@nih.gov.

Figures

Figure 1- Human macrophages survive *in vivo* for up to two weeks post-
injection following brain injection. (a) Schematic of experimental design: primary
monocytes were differentiated into macrophages before injection into the zebrafish
brain at age 2 days post fertilization (dpf) and imaged at 1, 7 and 14 days post
injection (dpi). (b) Micrographs of representative whole larva at 3 dpf (left) and 3D
projections showing distribution and survival of human primary macrophages (blue)
injected into the hind brain of transgenic mpx:GFP (neutrophils-green)/flk:mCherry
(vessels-red) zebrafish embryos at 1 dpi (3 dpf) (right). (c) Micrographs of
representative whole larva at 9dpf (left) and 3D projections showing distribution and
survival of human primary macrophages (blue) injected into the hind brain of
transgenic mpx:GFP (neutrophils-green)/flk:mCherry (vessels-red) zebrafish
embryos at 7 dpi (9 dpf) (right). (d) Micrographs show that cells can persist for up
to 2 weeks after injection at 16 dpf. Top panel: representative zebrafish at 16 dpf.
Left panel: micrograph shows tiled image of transgenic mpx:GFP (neutrophils-
green)/flk:mCherry (vessels-red)16 dpf zebrafish, white square highlights region of
interest in the zebrafish brain. Right panel: micrograph of the inset where the white
arrows indicate human cells. Scales are indicated on each image.

Figure 2- Human macrophages survive and interact with zebrafish stromal cells

post-injection following brain injection. (a) Schematic of experimental design:

primary monocytes were differentiated into macrophages before injection into the

zebrafish brain at age 2 days post fertilization (dpf) and imaged at 1 and 2 days post

injection (dpi). (b) 3D projections of whole head at 3 dpf (left) and 4 dpf (middle)

showing distribution and survival of human primary macrophages (blue) injected

into the hind brain of flk:mCherry (vessels-red) (top) and GFAP:GFP (astrocytes-

656 green)/flk:mCherry (vessels-red) (bottom) zebrafish larvae. Boxes indicate positions

657 of insets shown in right panels. Scales are indicated.

Figure 3- Human macrophages survive *in vivo* for up to two weeks post-injection following injection directly into the circulation of the zebrafish. (a) Schematic showing experimental set up, where human macrophages were directly injected into the circulation of 2 days post-fertilization (dpf) embryos, after being differentiated from primary human monocytes. Injected embryos were imaged at 1, 7 and/or 14 days post injection (dpi). (b-c) Micrographs of whole larva at 3 dpf (left) showing distribution and survival of human primary macrophages (blue)

differentiated at either at physiological temperature of zebrafish (28.5°C) or at

physiological temperature of humans (37°C) before injection into the caudal vein of

transgenic fli:GFP (vessels-red) zebrafish embryos. Micrographs display

macrophage distribution at 1 dpi. Bottom panels: Micrographs of 3D projections at

higher magnification of insets highlighted in tiled images showing distribution and

survival of human primary macrophages (blue). (d) Top panel: micrograph shows

tiled image of the 9 dpf zebrafish, 7dpi, showing distribution and survival of human

primary macrophages differentiated at human physiological temperature (37°C)

(blue) injected into transgenic fli:GFP (vessels-red) zebrafish embryo. Bottom panel:

Micrographs of 3D projections at higher magnification of insets highlighted in tiled

image showing distribution and survival of human primary macrophages (blue). (e)

Top panel: micrograph shows representative tiled image of a 16 dpf zebrafish.

Bottom panel: Micrographs of 3D projections showing distribution and survival of

human primary macrophages (blue) at 14 dpi following injection to transgenic

mpx:gfp (neutrophils-green)/flk:mCherry (vessels-red) zebrafish embryos at 2 dpf.

Arrows indicate macrophage positions.

Figure 4- Primary human monocytes survive at physiological temperature of the zebrafish in vitro and in vivo. (a) Primary human monocytes (blue) were injected into the hind brain of transgenic GFAP:GFP (astrocytes-green)/flk:mCherry (vessels-red) zebrafish embryos at 2 dpf. Micrographs show representative images where zebrafish was imaged serially over the course of 3 days post-injection at interval of 24 hours. (b) Plot (mean±95% CI) of average in vivo survival calculated for primary human monocytes obtained from larvae where the numbers of cells that survived over the course of 3 days for each larvae were normalized to the initial numbers measured one day post injection. Statistical analysis where * indicates a p value of p<0.025 and *** indicates a p value of p<0.0005 by successive t tests with Bonferroni correction, with n=41 larvae for Day 1, n=41 larvae for Day 2, and n=4 larvae for Day 3. Scatter plots show values for individual zebrafish. Not all zebrafish

survived to 5 days post injection. (c) Cells were either cultured at physiological

temperature of humans (37°C) or at physiological temperature of zebrafish (28.5°C).

Plot (mean±SD) of average survival calculated for primary human monocytes

obtained from n=4 donors where the numbers of cells that survived over the course

of 7 days were normalized to the original seeding density. Days in culture was a

statistically significant factor in survival by two-way ANOVA with one degree of

freedom (P<0.0001), as was the temperature of culture (P=0.0018).

Figure 5- Flow cytometry determined that primary human monocytes can survive and differentiate into macrophages at physiological temperature of the zebrafish in vitro. (a) Schematic showing that human primary monocytes were cultured at physiological temperature of zebrafish (28.5°C) or at physiological temperature of humans (37°C) with or without differentiation by cytokine human macrophage colony stimulating factor (H-M-CSF) for 5, 8, and 11 days. Bottom panel shows micrographs show the morphology of cells in the presence or absence of cytokine at each temperature. (b) Graphs (mean±SD) showing fraction of migrated cells in response or absence of isolated zebrafish astrocytes in a transwell

assay. *, P=0.0410 by Tukey's multiple comparisons post-test, n=3

membranes/condition for astrocytes isolated on a single day. (c) Primary monocytes

from 5 donors were either cultured at physiological temperature of humans (37°C)

or at physiological temperature of zebrafish (28.5°C) with or without H-M-CSF. Plot

(mean±SD) of average survival calculated for primary human

monocytes/macrophages that survived over the course of 5 days were normalized to

the original seeding density. Significantly more cells survived at 28.5°C vs. 37°C in

the presence or absence of H-M-CSF by two-way ANOVA. (d) Human primary

monocytes were cultured at physiological temperature of zebrafish (28.5°C, left

panels) or at physiological temperature of humans (37°C, right panels) with cytokine

human macrophage colony stimulating factor (H-M-CSF) for five, eight and 11

days. Expression of CD14 was assessed by flow cytometry. Light grey curve

indicates control where shift in dark grey curve indicates expression. (e) Cells

positive for CD14 expression were then assessed for markers, CD86, CD163 and

CD206 by flow cytometry at each temperature and time point.

Figure 6- Primary human macrophages can survive at physiological temperature of the zebrafish in vivo and do not cause a sustained inflammatory response. (a) Representative light sheet micrographs of uninjected, control whole larva at 3 days post-fertilization (dpf). Both transgenic mpeg:GFP (macrophages-green)/flk:mCherry (vessels-red) (top) and mpx:GFP (neutrophils-

green)/flk:mCherry (vessels-red) (bottom) are shown. (b) Representative lightsheet

micrographs of head regions of uninjected, control embryos at 3 dpf. Both transgenic

mpeg:GFP (macrophages-green)/flk:mCherry (vessels-red) (top) and mpx:GFP

(neutrophils-green)/flk:mCherry (vessels-red) (bottom) are shown. (c) Schematic of

experimental design: primary monocytes were differentiated into macrophages

before injection into the zebrafish brain at age 2 dpf and imaged at 1 and 7 days post

injection (dpi). (d) Plot (mean±SD) of average *in vivo* survival calculated for

primary human macrophages differentiated at physiological temperatures of the

zebrafish or human obtained from 3 larvae each, where the numbers of cells that

survived over the course of 7 days were normalized to the initial numbers measured

one day post injection. Differences in survival were not significant by two-way

ANOVA with Tukey's multiple comparisons post-test. The temperature of

macrophage differentiation (P=0.3072), fish line injected (P=0.4822), and

interaction between factors (P=0.4575) were not significant sources of variation by

two-way ANOVA with one degree of freedom. Monocytes that had been cultured in

the presence of human macrophage colony stimulating factor (H-M-CSF) for eight

days (e) at physiological temperature of (28.5°C) and (f) at physiological

temperature of humans (37°C). Primary human macrophages (blue) were injected

into the hind brain of transgenic mpeg:GFP (macrophages-green)/flk:mCherry

(vessels-red) or mpx:GFP (neutrophils-green)/flk:mCherry (vessels-red) zebrafish

embryos at 2 dpf. Micrographs of 3D projections showing distribution and survival

of human primary macrophages 1 dpi, when larvae are 3 dpf, and 7 dpi, when larvae

are 9 dpf. The same fish is shown in the left and right panels.

a Matrisome

b Polarization markers

Differentiation temperature (°C)

Differentiation temperature (°C)

c Gene expression *in vitro*

TNF   CD163   CCL18   VEGF

Cytokine treatment   Cytokine treatment   Cytokine treatment   Cytokine treatment

Differentiation temperature
■ 28.5°C ■ 37°C

d Gene expression *in vivo*

Primary human monocytes → Primary human macrophages

Differentiation - 8 days at 28.5°C or 37°C

Brain injection at 2 dpf

Primary human macrophages → RNA extraction from pooled embryos + PCR

24 h post-injection

Differentiation temperature
■ 28.5°C ■ 37°C

Figure 7- Primary human macrophages show gene expression of activated phenotypes in vitro and in vivo when cultured at the physiological temperature of the zebrafish. Human primary monocytes were cultured at physiological temperature of zebrafish (28.5°C) or at physiological temperature of humans (37°C) with human macrophage colony stimulating factor (H-M-CSF) for eight days prior to RNA isolation for RNAseq or further treatment with the cytokines EGF or TNFα prior to PCR. (a) Heatmap showing the log10 normalized counts of matrisome-associated genes. (b) Heatmap showing the log10 normalized counts for genes associated with macrophage polarization. In (a, b), heatmaps display genes for which $p < 0.05$ and $FDR < 0.01$ between temperatures. (c) Graphs show relative gene expression in specific markers for TNF−α, CD163, CCL18, and VEGF for human macrophages cultured *in vitro* in the presence or absence of recombinant EGF or TNF−α for 24 hours at physiological temperature of zebrafish (28.5°C) or at physiological temperature of humans (37°C) by real-time PCR. Plots are normalized to untreated macrophages differentiated at 37°C. (d) Human primary monocytes were cultured at physiological temperature of zebrafish (28.5°C) or at physiological temperature of humans (37°C) with human macrophage colony stimulating factor (H-M-CSF) for eight days prior to injection to the zebrafish brain. RNA was extracted from injected embryos 24 hours after injection. Graph shows *in vivo* relative gene expression in specific markers for TNF−α, CD163, CCL18, VEGF,

781   CCL17, CD24, and IL-6 for human macrophages 24 hours after injection into the

782   brain of the zebrafish. Plots are normalized to expression in macrophages

783   differentiated at 37°C prior to injection and subsequent RNA isolation for each gene.

784   For (c, d), *, $p<0.05$, ***, $p<0.001$, ****, $p<0.0001$ by two-way ANOVA with

785   Sidak's or Tukey's multiple comparisons test.

786
References

1.   Wynn TA, Chawla A, & Pollard JW (2013) Macrophage biology in development,
homeostasis and disease. *Nature* 496(7446):445-455.
2.   Pollard JW (2009) Trophic macrophages in development and disease. *Nat Rev*
Immunol 9(4):259-270.
3.   Murray PJ, *et al.* (2014) Macrophage activation and polarization: nomenclature and
experimental guidelines. *Immunity* 41(1):14-20.
4.   Coussens LM & Werb Z (2002) Inflammation and cancer. *Nature* 420(6917):860-867.
5.   Kingwell K (2012) Neurodegenerative disease: Microglia in early disease stages. *Nat*
Rev Neurol 8(9):475.
6.   Perry VH, Nicoll JA, & Holmes C (2010) Microglia in neurodegenerative disease. *Nat*
Rev Neurol 6(4):193-201.
7.   Lumeng CN & Saltiel AR (2011) Inflammatory links between obesity and metabolic
disease. *Journal of Clinical Investigation* 121(6):2111-2117.
8.   DeNardo DG, Johansson M, & Coussens LM (2008) Immune cells as mediators of solid
tumor metastasis. *Cancer Metast Rev* 27(1):11-18.
9.   Hanahan D & Weinberg RA (2011) Hallmarks of cancer: the next generation. *Cell*
144(5):646-674.
10.   Vesely MD, Kershaw MH, Schreiber RD, & Smyth MJ (2011) Natural innate and
adaptive immunity to cancer. *Annu Rev Immunol* 29:235-271.
11.   Zitvogel L, Pitt JM, Daillere R, Smyth MJ, & Kroemer G (2016) Mouse models in
oncoimmunology. *Nat Rev Cancer* 16(12):759-773.
12.   AlmeidaPorada G, Ascensao JL, & Zanjani ED (1996) The role of sheep stroma in
human haemopoiesis in the human/sheep chimaeras. *Brit J Haematol* 93(4):795-802.
13.   Langenau DM & Zon LI (2005) The zebrafish: A new model of T-cell and thymic
development. *Nature Reviews Immunology* 5(4):307-317.
14.   Morton JJ, *et al.* (2016) XactMice: humanizing mouse bone marrow enables
microenvironment reconstitution in a patient-derived xenograft model of head and
neck cancer. *Oncogene* 35(3):290-300.
15.   Renshaw SA & Trede NS (2012) A model 450 million years in the making: zebrafish
and vertebrate immunity. *Dis Model Mech* 5(1):38-47.

16.   Theocharides AP, Rongvaux A, Fritsch K, Flavell RA, & Manz MG (2016) Humanized
819   hemato-lymphoid system mice. *Haematologica* 101(1):5-19.
820   17.   Zanjani ED, AlmeidaPorada G, & Flake AW (1996) The human/sheep xenograft
821   model: A large animal model of human hematopoiesis. *Int J Hematol* 63(3):179-192.
822   18.   Ruffell B, Affara NI, & Coussens LM (2012) Differential macrophage programming in
823   the tumor microenvironment. *Trends Immunol* 33(3):119-126.
824   19.   Joyce JA & Pollard JW (2009) Microenvironmental regulation of metastasis. *Nat Rev*
825   *Cancer* 9(4):239-252.
826   20.   Wang J, *et al.* (2015) Novel mechanism of macrophage-mediated metastasis revealed
827   in a zebrafish model of tumor development. *Cancer Res* 75(2):306-315.
828   21.   Wyckoff JB, *et al.* (2007) Direct visualization of macrophage-assisted tumor cell
829   intravasation in mammary tumors. *Cancer Res* 67(6):2649-2656.
830   22.   Kim J & Tanner K (2015) Recapitulating the Tumor Ecosystem Along the Metastatic
831   Cascade Using 3D Culture Models. *Front Oncol* 5:170.
832   23.   Renshaw SA & Trede NS (2012) A model 450 million years in the making: zebrafish
833   and vertebrate immunity. *Disease Models & Mechanisms* 5(1):38-47.
834   24.   White R, Rose K, & Zon L (2013) Zebrafish cancer: the state of the art and the path
835   forward. *Nat Rev Cancer* 13(9):624-636.
836   25.   Meeker ND & Trede NS (2008) Immunology and zebrafish: spawning new models of
837   human disease. *Dev Comp Immunol* 32(7):745-757.
838   26.   Oosterhof N, Boddeke E, & van Ham TJ (2015) Immune cell dynamics in the CNS:
839   Learning from the zebrafish. *Glia* 63(5):719-735.
840   27.   Sanders GE (2012) Zebrafish housing, husbandry, health, and care: IACUC
841   considerations. *ILAR J* 53(2):205-207.
842   28.   Gordon S & Martinez FO (2010) Alternative activation of macrophages: mechanism
843   and functions. *Immunity* 32(5):593-604.
844   29.   Zhang Y, *et al.* (2013) ROS play a critical role in the differentiation of alternatively
845   activated macrophages and the occurrence of tumor-associated macrophages. *Cell*
846   *Res* 23(7):898-914.
847   30.   Tarique AA, *et al.* (2015) Phenotypic, functional, and plasticity features of classical
848   and alternatively activated human macrophages. *Am J Respir Cell Mol Biol* 53(5):676-
849   688.
850   31.   Weiss M, Blazek K, Byrne AJ, Perocheau DP, & Udalova IA (2013) IRF5 is a specific
851   marker of inflammatory macrophages in vivo. *Mediators Inflamm* 2013:245804.
852   32.   Langenau DM & Zon LI (2005) The zebrafish: a new model of T-cell and thymic
853   development. *Nat Rev Immunol* 5(4):307-317.
854   33.   Adkins B (1999) T-cell function in newborn mice and humans. *Immunol Today*
855   20(7):330-335.
856   34.   Macchiarini F, Manz MG, Palucka AK, & Shultz LD (2005) Humanized mice: are we
857   there yet? *J Exp Med* 202(10):1307-1311.
858   35.   Gordon S & Taylor PR (2005) Monocyte and macrophage heterogeneity. *Nat Rev*
859   *Immunol* 5(12):953-964.
860   36.   Montali RJ (1988) Comparative pathology of inflammation in the higher vertebrates
861   (reptiles, birds and mammals). *J Comp Pathol* 99(1):1-26.

37.   DeNardo DG & Coussens LM (2007) Inflammation and breast cancer. Balancing
immune response: crosstalk between adaptive and innate immune cells during breast
cancer progression. *Breast cancer research : BCR* 9(4):212.
38.   Sullivan AR & Pixley FJ (2014) CSF-1R signaling in health and disease: a focus on the
mammary gland. *J Mammary Gland Biol Neoplasia* 19(2):149-159.
39.   Wyckoff J, *et al.* (2004) A paracrine loop between tumor cells and macrophages is
required for tumor cell migration in mammary tumors. *Cancer Research* 64(19):7022-
7029.
40.   Ries CH, *et al.* (2014) Targeting tumor-associated macrophages with anti-CSF-1R
antibody reveals a strategy for cancer therapy. *Cancer Cell* 25(6):846-859.
41.   Gul N, *et al.* (2014) Macrophages eliminate circulating tumor cells after monoclonal
antibody therapy. *J Clin Invest* 124(2):812-823.
42.   Pucci F, *et al.* (2016) SCS macrophages suppress melanoma by restricting tumor-
derived vesicle-B cell interactions. *Science* 352(6282):242-246.
43.   Quail DF & Joyce JA (2013) Microenvironmental regulation of tumor progression and
metastasis. *Nat Med* 19(11):1423-1437.
44.   Roszer T (2015) Understanding the Mysterious M2 Macrophage through Activation
Markers and Effector Mechanisms. *Mediat Inflamm*.
45.   Mantovani A, *et al.* (2004) The chemokine system in diverse forms of macrophage
activation and polarization. *Trends Immunol* 25(12):677-686.
46.   Schraufstatter IU, Zhao M, Khaldoyanidi SK, & Discipio RG (2012) The chemokine
CCL18 causes maturation of cultured monocytes to macrophages in the M2 spectrum.
Immunology 135(4):287-298.
47.   Tanaka T, Narazaki M, & Kishimoto T (2014) IL-6 in inflammation, immunity, and
disease. *Cold Spring Harb Perspect Biol* 6(10):a016295.
48.   Ara T & Declerck YA (2010) Interleukin-6 in bone metastasis and cancer progression.
Eur J Cancer 46(7):1223-1231.
49.   Steeg PS (2016) Targeting metastasis. *Nat Rev Cancer* 16(4):201-218.
50.   Erez N & Coussens LM (2011) Leukocytes as paracrine regulators of metastasis and
determinants of organ-specific colonization. *International journal of cancer. Journal*
international du cancer 128(11):2536-2544.
51.   Eyles J, *et al.* (2010) Tumor cells disseminate early, but immunosurveillance limits
metastatic outgrowth, in a mouse model of melanoma. *J Clin Invest* 120(6):2030-2039.
52.   Bowman RL, *et al.* (2016) Macrophage Ontogeny Underlies Differences in Tumor-
Specific Education in Brain Malignancies. *Cell Rep* 17(9):2445-2459.
53.   Renshaw SA, *et al.* (2006) A transgenic zebrafish model of neutrophilic inflammation.
Blood 108(13):3976-3978.
54.   Ellett F, Pase L, Hayman JW, Andrianopoulos A, & Lieschke GJ (2011) mpeg1 promoter
transgenes direct macrophage-lineage expression in zebrafish. *Blood* 117(4):e49-56.
55.   Bernardos RL & Raymond PA (2006) GFAP transgenic zebrafish. *Gene Expr Patterns*
6(8):1007-1013.
56.   Manoli M & Driever W (2012) Fluorescence-activated cell sorting (FACS) of
fluorescently tagged cells from zebrafish larvae for RNA isolation. *Cold Spring Harb*
Protoc 2012(8).

57. Blehm BH, Devine A, Staunton JR, & Tanner K (2016) In vivo tissue has non-linear rheological behavior distinct from 3D biomimetic hydrogels, as determined by AMOTIV microscopy. *Biomaterials* 83:66-78.

58. Bolger AM, Lohse M, & Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30(15):2114-2120.

59. Andrew S (2010) FastQC: a quality control tool for high throughput sequence data).

60. Dobin A*, et al.* (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29(1):15-21.

61. Liao Y, Smyth GK, & Shi W (2014) featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30(7):923-930.

62. Robinson MD & Oshlack A (2010) A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol* 11(3):R25.

63. McCarthy DJ, Chen Y, & Smyth GK (2012) Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res* 40(10):4288-4297.

64. Robinson MD, McCarthy DJ, & Smyth GK (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26(1):139-140.

65. Kolde R (2012) Pheatmap: pretty heatmaps. *R package version* 61.

66. Naba A*, et al.* (2016) The extracellular matrix: Tools and insights for the "omics" era. *Matrix Biol* 49:10-24.

67. Martinez FO, Gordon S, Locati M, & Mantovani A (2006) Transcriptional profiling of the human monocyte-to-macrophage differentiation and polarization: new molecules and patterns of gene expression. *J Immunol* 177(10):7303-7311.