Long title: Human demographic history has amplified the effects of background selection across the genome

Short title: Background selection and demography in humans

Raul Torres[1], Zachary A. Szpiech[2], and Ryan D. Hernandez[2,3,4,5*]

[1] Biomedical Sciences Graduate Program, University of California San Francisco, San Francisco, CA, United States of America

[2] Department of Bioengineering and Therapeutic Sciences, University of California San Francisco, San Francisco, CA, United States of America

[3] Institute for Human Genetics, University of California San Francisco, San Francisco, CA, United States of America

[4] Institute for Computational Health Sciences, University of California San Francisco, San Francisco, CA, United States of America

[5] Quantitative Biosciences Institute, University of California San Francisco, San Francisco, CA, United States of America

* Corresponding author

E-mail: ryan.hernandez@ucsf.edu (RDH)

Abstract

Natural populations often grow, shrink, and migrate over time. Demographic processes such as these can impact genome-wide levels of genetic diversity. In addition, genetic variation in functional regions of the genome can be altered by natural selection, which drives adaptive mutations to higher frequency or purges deleterious ones. Such selective processes impact not only the sites directly under selection but also nearby neutral variation through genetic linkage. While there is extensive literature examining the impact of linked selection (i.e., genetic hitchhiking in the context of positive selection and background selection in the context of deleterious variants) at demographic equilibrium, less is known about how non-equilibrium demographic processes impact patterns of linked selection. Utilizing a global sample of human whole-genome sequences from the Thousand Genomes Project and extensive simulations, we investigate how non-equilibrium demographic processes magnify and dampen the consequences of background selection (BGS) across the human genome. We observe that, compared to Africans, non-African populations have experienced larger proportional decreases in neutral genetic diversity in regions of strong BGS. We replicate these findings in admixed populations by showing that non-African ancestral components of the genome have been impacted more severely in regions of strong BGS. We attribute these differences to the strong population bottlenecks that non-Africans experienced as they migrated out of Africa and throughout the globe. Furthermore, we observe a strong correlation between F_{ST} and BGS, suggesting a stronger rate of genetic drift in regions of strong BGS. Forward simulations of human demographic history and BGS support these observations. Our results show that non-equilibrium demography significantly alters the conse-

24 quences of BGS and support the need for more work investigating the dynamic process

25 of multiple evolutionary forces operating in concert.

Author summary

Patterns of genetic diversity within a species are impacted at broad and fine

scales by population size changes ("demography") and natural selection. From both

population genetics theory and observation of genomic sequence data, it is known that

population size changes can impact genome-wide average neutral genetic diversity.

Additionally, natural selection can impact neutral genetic diversity regionally across the

genome through the process of linked selection. During this process, natural selection

acting on adaptive or deleterious variants in the genome will also impact diversity at

linked neutral sites. However, less is known about the dynamic changes to diversity that

occur in regions impacted by linked selection when a population undergoes a size

change. We characterize these dynamic changes using human sequence data, focusing

on regions of the genome experiencing linked selection that is caused by deleterious

variation (called "background selection"). We find that the population size changes ex-

perienced by humans have shaped the consequences of linked selection in the human

genome. In particular, population contractions, such as those experienced by non-

Africans, have disproportionately decreased neutral diversity in regions of the genome

experiencing strong BGS, resulting in large differences between African and non-African

populations.

Introduction

Genetic diversity in a species is determined through the complex interplay of mutation, demography, genetic drift, and natural selection. These evolutionary forces operate in concert to shape patterns of diversity at both the local scale and genome-wide scale. For example, in recombining species, levels of genetic diversity are distributed heterogeneously across the genome as peaks and valleys that are often correlated with recombination rate and generated by past or ongoing events of natural selection [1]. But at the genome-wide scale, average levels of genetic diversity are primarily impacted by population size changes, yielding signatures of diversity that are a function of a population's demographic history [2]. These patterns of diversity may also yield information for inferring past events of natural selection and population history, giving valuable insight into how populations have evolved over time [3–8]. With recent advancements in sequencing technology yielding whole-genome data from thousands of individuals from species with complex evolutionary histories [9,10], formal inquiry into the interplay of demography and natural selection and testing of whether demographic effects act uniformly across the genome as a function of natural selection is now possible.

In the past decade, population genetic studies have shed light on the pervasiveness of dynamic population histories in shaping overall levels of genetic diversity across different biological species. For example, multiple populations have experienced major population bottlenecks that have resulted in decreased levels of genome-wide diversity. Evidence for population bottlenecks exists in domesticated species such as cattle [11], dogs [12], and rice [13], and in natural populations such as *Drosophila melanogaster* [14–16], rhesus macaque [17], and humans [18,19]. Notably, population bottlenecks

leave inordinately long lasting signatures of decreased diversity, which may be de-

pressed long after a population has recovered to its ancestral size [20].

Locally (i.e., regionally) across the genome, the action of natural selection can

also lead to distinct signatures of decreased genetic diversity (although some forms of

selection, such as balancing selection, can increase genetic diversity [21]). For exam-

ple, mutations with functional effects may be removed from the population due to purify-

ing selection or fix due to positive selection, thereby resulting in the elimination of genet-

ic diversity at the site. But while sites under direct natural selection in the genome rep-

resent only a small fraction of all sites genome-wide, the action of natural selection on

these selected sites can have far-reaching effects across neutral sites in the genome

due to linkage. Under positive selection, genetic hitchhiking [22] causes variants lying

on the same haplotype as the selected allele to rise to high frequency during the selec-

tion process. Conversely, under purifying selection, background selection [23] causes

linked neutral variants to decrease in frequency or to be removed from the population.

Both of these processes of linked selection result in decreased neutral genetic diversity

around the selected site. Recombination can decouple neutral sites from selected sites

in both cases and neutral diversity tends to increase toward its neutral expectation as

genetic distance from selected sites increases [24].

Evidence for genetic hitchhiking and background selection has been obtained

from the genomes of several species, including *Drosophila melanogaster* [25–30], wild

and domesticated rice [31,32], nematode [33,34], humans [3,6,35–39], and others (see

[1] for a review). While the relative contributions of genetic hitchhiking and background

selection to shaping patterns of human genomic diversity have been actively debated

[37,40–42], the data strongly support a large role for background selection in shaping

genome-wide patterns of neutral genetic variation [38,39]. Indeed, recent arguments

have been made in favor of background selection being treated as the null model when

investigating the impact of linked selection across recombining genomes [1,29,42–44].

To build on this movement, our investigation will focus on aspects of background selec-

tion (BGS) in humans.

The impact of BGS has been quantified across the human genome [6], which

suggests that neutral genetic diversity in the human autosomes has been reduced 19-

26% because of BGS. However, it is not obvious whether populations that have experi-

enced different demographic histories, such as African and non-African humans, should

exhibit similar effects in regions of BGS. Early work resulted in the expression $\pi \approx$

$4f_0 N_e \mu$ [23], which suggests that the expected level of diversity with background selec-

tion would be proportional to the neutral expectation (with proportionality constant f_0 be-

ing a function of the rates of deleterious mutation and recombination). While demo-

graphic forces will impact N_e, this classic model implies that the relative reduction in di-

versity due to BGS may be insensitive to different demographic histories. Recent work

has demonstrated that this assumption may hold if selection is strong enough (or popu-

lations are large enough) such that mutation-selection balance is maintained [45,46].

However, humans and several other natural populations likely do not meet such criteria,

and are rarely at demographic equilibrium. Therefore, other evolutionary forces, such as

genetic drift, may perturb genetic diversity in regions of BGS that are unaccounted for in

these models.

While little is known about the impact of demography on BGS' effects, recent

studies have suggested that alleles directly under natural selection experience non-

linear dynamics in the context of non-equilibrium demography. In general, the equilibri-

um frequency of an allele is dependent on its fitness effect, with deleterious alleles hav-

ing lower equilibrium frequencies than neutral alleles. After a population size change,

deleterious alleles tend to change frequency faster than neutral alleles, allowing them to

reach their new equilibrium frequency at a faster rate [47,48]. This can result in relative

differences in deleterious allele frequencies when comparing populations with different

demographic histories. Such effects are especially apparent in populations suffering

bottlenecks [49] and have been tested and observed between different human popula-

tions [50–52]. We hypothesized that these non-equilibrium dynamics could also amplify

the effects of background selection, with classic models being poorly suited to describe

such impacts in non-equilibrium populations. In the case of population bottlenecks, re-

gions under BGS may respond more rapidly than neutral regions to such demographic

events, if the rate of removal of genetic diversity in regions under BGS is also higher.

To investigate the impact of non-equilibrium dynamics in regions of BGS, we

measure patterns of average pairwise neutral genetic diversity (π), or neutral heterozy-

gosity if the population was admixed, as a function of the strength of BGS within a glob-

al set of human populations from phase 3 of the Thousand Genomes Project (TGP) [9].

We focus particularly on the ratio of neutral diversity in regions of strong BGS to regions

of weak BGS, which we term "relative diversity." We also investigate how genetic differ-

entiation between TGP populations (as measured by F_{ST}) is shaped by BGS strength.

We find substantial differences in relative diversity between populations, which we at-

tribute to their non-equilibrium demographics. We confirm that the interplay of demogra-

phy and BGS can explain the differences of relative diversity across human populations

with simulations incorporating a parametric demographic model of human history [7]

with and without a model of BGS. Specifically, our simulations of non-African human

demography show that the out-of-Africa population bottleneck and European-Asian split

population bottleneck each coincide with a distinct drop in relative diversity. Further-

more, we demonstrate that back migration from Europeans and Asians into Africa re-

introduces sufficient deleterious variation to impact BGS strength, leading to decreased

relative diversity in Africans. These results demonstrate the strong impact that changing

demography has on perturbing levels of diversity in regions under BGS and have impli-

cations for population genetic studies seeking to characterize linked selection across

any species or population that is not at demographic equilibrium.

Results

Differential impact of BGS across human populations

We measured mean pairwise genetic diversity (π) among the 20 non-admixed

populations from the phase 3 TGP data set, consisting of 5 populations each from 4

continental groups: Africa (AFR), Europe (EUR), South Asia (SASN), and East Asia

(EASN) (Table S11 in Supporting information for population labels and grouping). After

a set of stringent filters were applied to all 20 populations to identify a high-quality set of

putatively neutral sites in the genome (see Materials and Methods), sites were divided

into quantile bins based on their BGS coefficient (termed "*B*"), which represents the in-

ferred strength of BGS from McVicker et al. [6] (see Materials and Methods). For our

initial set of analyses, we focused on the bins corresponding to the 1% of sites under

strongest BGS (or lowest B) and the 1% of sites under weakest BGS (or highest B).

Mean diversity (normalized by divergence with rhesus macaque) within these bins for

each population is shown in Figs 1A-B. As expected, normalized diversity was highest

in African populations and lowest in East Asian populations across both 1% B quantile

bins.

To obtain estimates of the reduction in diversity in regions of strong BGS for each

population, we calculated a statistic called "relative diversity" (analogous to π/π_0 in the

BGS literature; [23,53]), which we defined as the ratio of normalized diversity in the

strongest 1% BGS bin to normalized diversity in the weakest 1% BGS bin. Fig 1C

shows that relative diversity was lower in non-African populations (0.348-0.365 for non-

Africans, 0.396-0.408 for Africans), suggesting higher rates of genetic diversity reduc-

tion in strong BGS regions in non-African populations compared to African populations.

Furthermore, the greatest reduction was observed among East Asian populations (CDX,

JPT, and CHS populations).

To characterize these effects across a broader distribution of BGS strengths, we

grouped populations together according to their continental group (i.e., African, Europe-

an, South Asian, and East Asian, see Table S11 in Supporting information for a detailed

description) and estimated relative diversity at neutral sites for each of the continental

groups in bins corresponding to the bottom 1%, 5%, 10%, and 25% quantiles of B (note

these partitions were not disjoint). As expected, relative diversity increased for all conti-

nental groups as the bins became more inclusive (Fig 2B), reflecting a reduced impact

on the reduction of diversity due to BGS. We also observed that non-African continental

groups consistently had a lower relative diversity compared to African groups, demon-

strating that the patterns we observed in the most extreme BGS regions also held for

weaker BGS regions. Interestingly, we observed a consistent trend of rank order for rel-

ative diversity between the different continental groups for each quantile bin, with the

East Asian group experiencing the greatest reduction of relative diversity, followed by

the South Asian, European, and African groups. This result suggested a stronger effect

for demography on the diversity-reducing effect of BGS for those populations experienc-

ing the strongest bottlenecks. However, the observed differences in relative diversity

between non-African and African continental groups became less pronounced as the

bins became more inclusive (Fig 2B). These effects remained even after we controlled

for the effects of GC-biased gene conversion and recombination hotspots (Figs S2 and

S4) or if we did not normalize diversity by divergence (Figs S3 and S5).

Recent admixture has not altered the impact of BGS

We then investigated whether the effects of BGS have remained consistent

across human populations that have experienced recent admixture. To do so, we tested

for the effect of BGS on normalized and relative diversity in the 6 admixed TGP popula-

tions (ASW, ACB, CLM, MXL, PEL, and PUR). We first used the local ancestry tracks

inferred by the phase 3 TGP consortium to divide up admixed samples into genomic

segments that are homozygous for a specific local ancestry (i.e., African, European, or

Native American). These segments were then collated across all admixed samples (see

Materials and Methods), irrespective of their source population. We then calculated

normalized and relative heterozygosity across the collated segments as a function of

the BGS quantile bins described in the previous section. For comparison, heterozygosi-

ty was also calculated for each TGP continental group. Across all BGS quantile bins,

normalized diversity (heterozygosity/divergence) in African and European ancestry

segments closely matched the values observed in their non-admixed counterparts (Fig

3A). However, normalized diversity was significantly lower in the Native American an-

cestry segments of admixed individuals than in the East Asian continental group (Fig

3A). This was expected given the more recent divergence of Native American popula-

tions and the strong population bottleneck they experienced migrating into the Americas

[18,54,55].

Patterns of relative diversity for each of the ancestries also largely recapitulated

the patterns observed in their continental group counterparts across all BGS quantile

bins, with the largest reductions in relative diversity occurring for the Native American

and European ancestral segments (Fig 3B). These patterns were similar to the broader

analyses of the 20 non-admixed populations described above, with a consistent rank

order of decreasing relative diversity observed for African, European, and Native Ameri-

can ancestral segments. However, for the strongest 1% BGS quantile bin, relative di-

versity in Native American ancestry segments was observed to be greater than for the

European continental group or European local ancestry segments, which was incon-

sistent with the other BGS quantile bins.

BGS has shaped patterns of population differentiation

To test if greater genetic drift in regions of strong BGS has contributed to the ob-

served greater losses of relative diversity in non-African populations and to understand

how BGS has impacted local patterns of population differentiation, we measured F_{ST} as

a function of B. It has been shown that estimates of F_{ST} are affected by SNP ascertain-

ment as well as the choice of computation method [56].To limit such biases, we followed

the procedures recommended by Bhatia et al. [56] (see Materials and Methods) and on-

ly analyzed SNPs that were polymorphic in an outgroup human population (the Khoe-

San). After filtering, a total of 3,497,105 sites were left for estimating F_{ST}. Sites were di-

vided into 2% quantile bins based on the genome-wide distribution of B, and F_{ST} was

calculated within each bin for all population pairs in which both populations were from

different continental groups (150 total pairwise comparisons). We performed simple lin-

ear regression using B as an explanatory variable and F_{ST} as our dependent variable

with the linear model $F_{ST} = \beta_0 + \beta_1 B + \varepsilon$. We found that B was significantly associated

with an increase in local F_{ST} across the genome (Fig 4A, Table 1). When considering all

150 population comparisons (referred to as the "Global" estimate in Table 1), B ex-

plained 26.9% of the change in F_{ST} across the most extreme B values (i.e., when com-

paring weak BGS [$B = 1$] to the strong BGS [$B = 0$]). To test if the linear model was bi

ased by outliers or bins with high influence, we also performed robust linear regression

but still observed a significant contribution of B to F_{ST}, with a 28.3% change between the

highest and lowest B value (Table S6 in Supporting information).

257 **Table 1. Regression coefficient estimates for linear regression of F_{ST} on 2% quan-**
258 **tile bins of B**.

	AFR vs. EASN	AFR vs. EUR	AFR vs. SASN	EUR vs. SASN	EUR vs. EASN	SASN vs. EASN	Global
β_0	0.2044	0.1716	0.1596	0.0455	0.1216	0.0903	0.1322
± SEM	± 0.0039	± 0.0031	± 0.0029	± 0.0011	± 0.0029	± 0.0023	± 0.0019
(p-value)	(< 1e-04)	(< 1e-04)	(< 1e-04)	(< 1e-04)	(< 1e-04)	(< 1e-04)	(< 1e-04)
β_1	-0.0434	-0.0358	-0.0355	-0.0098	-0.0173	-0.0261	-0.0280
± SEM	± 0.0046	± 0.0037	± 0.0034	± 0.0013	± 0.0035	± 0.0027	± 0.0022
(p-value)	(< 1e-04)	(< 1e-04)	(< 1e-04)	(< 1e-04)	(< 1e-04)	(< 1e-04)	(< 1e-04)
r	-0.8363	-0.7441	-0.7794	-0.3847	-0.6220	-0.5968	-0.1292
± SEM	± 0.0295	± 0.0362	± 0.0332	± 0.0414	± 0.0785	± 0.0348	± 0.0098

259
The first two rows give the regression coefficients for the linear model $F_{ST} = \beta_0 + \beta_1 B +$
ε, where B represents the mean background selection coefficient for the bin being test-
ed and F_{ST} is the estimated F_{ST} for all population comparisons within a particular pair of
continental groups (given in the column header). The final column, "Global", gives the
regression coefficients for the linear model applied to all pairwise population compari-
sons (150 total). The correlation coefficient, r, between B and F_{ST} for each comparison
is shown in the bottom row. Standard errors of the mean (SEM) for β_0, β_1, and r were
calculated from 1,000 bootstrap iterations (see Materials and Methods). P-values are
derived from a two-sided t-test of the t-value for the corresponding regression coeffi-
cient.
270

Earlier studies using SNP array data have shown that F_{ST} and recombination rate

are correlated in humans [57]. We could only partially replicate these findings when we

conducted linear regression of F_{ST} as a function of recombination rate (ρ) (measured in

2% recombination rate quantile bins) with the linear model $F_{ST} = \beta_0 + \beta_1\rho + \varepsilon$. We ob-

served that recombination rate only significantly predicts a change in F_{ST} across the ge-

nome for comparisons between South Asian and East Asian populations (Fig 4B, Table

2). This result remained unchanged when performing robust linear regression for the

model (Table S7 in Supporting information).

279
280

Table 2. Regression coefficient estimates for linear regression of F_{ST} on 2% quantile bins of recombination rate.

	AFR vs. EASN	AFR vs. EUR	AFR vs. SASN	EUR vs. SASN	EUR vs. EASN	SASN vs. EASN	Global
β_0 ± SEM (p-value)	0.1688 ± 0.0007 (< 1e-04)	0.1422 ± 0.0006 (< 1e-04)	0.1305 ± 0.0006 (< 1e-04)	0.0373 ± 0.0002 (< 1e-04)	0.1070 ± 0.0006 (< 1e-04)	0.0688 ± 0.0004 (< 1e-04)	0.1091 ± 0.0003 (< 1e-04)
β_1 ± SEM (p-value)	-0.0009 ± 0.0026 (0.7073)	0.0005 ± 0.0022 (0.8454)	0.0005 ± 0.0021 (0.8196)	-0.0015 ± 0.0007 (0.3906)	0.0005 ± 0.0021 (0.7002)	-0.0050 ± 0.0014 (0.0363)	-0.0010 ± 0.0012 (0.8842)
r ± SEM	-0.0106 ± 0.0287	0.0055 ± 0.0257	0.0065 ± 0.0253	-0.0243 ± 0.0119	0.0109 ± 0.0379	-0.0592 ± 0.0159	-0.0017 ± 0.0021

The first two rows give the regression coefficients for the linear model $F_{ST} = \beta_0 + \beta_1\rho + \varepsilon$, where ρ represents the mean recombination rate for the bin being tested and F_{ST} is the estimated F_{ST} for all population comparisons within a particular pair of continental groups (given in the column header). The final column, "Global", gives the regression coefficients for the linear model applied to all pairwise population comparisons (150 total). When performing the regression, ρ was first scaled to between 0 and 1, such that 1 represents the maximum observed recombination rate (126.88 cM/Mb) and 0 represents the minimum observed recombination rate (0.0 cM/Mb). The correlation coefficient, r, between ρ and F_{ST} for each comparison is shown in the bottom row. Standard errors of the mean (SEM) for β_0, β_1, and r were calculated from 1,000 bootstrap iterations (see Materials and Methods). P-values are derived from a two-sided t-test of the t-value for the corresponding regression coefficient.

Since the correlation between F_{ST} and recombination rate was previously documented as being strongest in coding regions [57], where the impact of BGS and other modes of linked selection is also expected to be strongest, we sought to disentangle the roles of BGS and recombination rates as explanatory variables for F_{ST} by using multiple linear regression. This served as a test for any additional effect recombination rate may have on F_{ST} that B failed to capture. To do so, we first split the genome into 2% recombination rate quantile bins and further subdivided each of these bins into 4% B quantile bins (50×25 = 1,250 bins total). We then measured F_{ST} within each bin. We also partitioned sites in the reverse order (2% B bins followed by 4% recombination rate bins)

306 and repeated all analyses. Our choice in total number of bins resulted in a minimum of

307 320 SNPs per bin for estimating F_{ST} between any two populations, which should be suf-

308 ficient to avoid errors when estimating F_{ST} across multiple loci [58]. After performing

309 multiple linear regression of F_{ST} on B, recombination rate (ρ), and an interaction term

310 between the two ($B\rho$) with the linear model $F_{ST} = \beta_0 + \beta_1 B + \beta_2 \rho + \beta_3 B\rho + \varepsilon$, we ob-

311 served that B was a statistically significant predictor ($p < 1e\text{-}04$) for F_{ST} across all popu-

312 lation comparisons regardless of how we partitioned sites (Table S8 in Supporting in-

313 formation). This result remained unchanged when performing robust regression. In con-

314 trast, recombination rate exhibited sporadic significance as an explanatory variable for

315 F_{ST} across population comparisons and was dependent upon how sites were partitioned

316 (i.e., whether we first partitioned by B or by recombination rate) (Table S8 in Supporting

317 information). Furthermore, strong differences between the two binning schemes were

318 observed for the magnitude of the recombination rate regression coefficient for certain

319 population comparisons (e.g., African vs. East Asian and South Asian vs. East Asian),

320 while the coefficients for B were consistent across binning schemes. The direction in

321 which recombination rate explained F_{ST} was also inconsistent across different popula-

322 tion comparisons, with European vs. South Asian and European vs. East Asian compar-

323 isons showing a significant positive change in F_{ST} as a function of increasing recombi-

324 nation rate. This result was contrary to an expectation of decreasing F_{ST} as a function of

325 increasing recombination rate [57]. We also failed to observe consistent effects from the

326 interaction term for B and recombination rate on F_{ST} across population comparisons or

327 binning schemes (Table S8 in Supporting information). Performing robust regression on

328 the model did not change these results. However, in contrast to recombination rate,

when the model was performed utilizing all TGP populations (i.e., the "Global" estimate),

the interaction term was significant in explaining F_{ST} across both types of binning

schemes.

To aid in visualizing the results of our multidimensional linear model, we plotted

F_{ST} for each population comparison as a function of recombination rate (across 4%

quantile bins) while conditioning on B (Fig 5A). We also plotted points in the reciprocal

direction, with F_{ST} being plotted as a function of B while conditioning on recombination

rate (Fig 5B). These data points were derived from the same points used as input for

the multiple linear regression model described above. The results for F_{ST} between Afri-

can and South Asian populations showed that B separated different levels of F_{ST} across

most recombination rate bins (Fig 5A, Table S9 in Supporting information). Furthermore,

regardless of how B was conditioned on recombination rate, it still exhibited a strong

trend of increasing F_{ST} as the strength of BGS increased (Fig 5B, Table S10 in Support-

ing information). These patterns were imperfect though, and statistical significance was

not always attained, especially for comparisons between non-African populations (Fig

S6, Table S10 in Supporting information). However, greater separation in F_{ST} was gen-

erally achieved when conditioning recombination rate on B and the slope was always

negative when plotting F_{ST} across B, regardless of which recombination rate percentile

bin B was conditioned on. These results suggested that B is a dominant predictor for

F_{ST}, while recombination rate plays a minor role.

Demographic inference in putatively neutral regions of the genome

One consequence of BGS driving patterns of neutral variation within and be-

tween human populations is that demographic inference could be substantially biased

[59,60]. To assess the degree of bias, we inferred a 13-parameter demographic model

of African, European, and East Asian demography using only putatively neutral regions

of the genome under the weakest effects of BGS ($B \geq 0.994$) from a subset of TGP indi-

viduals with high coverage whole genome sequence data (see Materials and Methods).

Our demographic model followed that of Gutenkunst et al. [7], with an ancient human

expansion in Africa and a single out-of-Africa bottleneck followed by European- and

East Asian-specific bottlenecks, as well as exponential growth in both non-African popu-

lations and migration between all populations. To make comparisons to previous studies

that have used sequence data from coding regions or genes [7,61,62], which may be

under strong BGS, we also inferred demographic parameters using coding four-fold de-

generate synonymous sites. Our inferred parameters for human demography were strik-

ingly different between the two sets of sequence data (Fig S1, Table S1 in Supporting

information). Notably, inferred effective population size parameters were larger for con-

temporary population sizes when using four-fold degenerate synonymous sites versus

weak BGS neutral regions, with N_e inferred to be 22%, 23%, and 29% larger for AFR,

EUR, and EASN populations, respectively. This is despite the fact that the ancestral N_e

was inferred to be lower for four-fold degenerate synonymous sites (N_e = 18,449 and

17,118, for weak BGS neutral regions and four-fold degenerate sites, respectively). This

result may stem from the expected decrease in N_e going into the past in regions of

BGS, which can lead to inflated estimates of recent population growth [60] and has also

been shown in simulation studies of synonymous sites under BGS [59].

Simulations confirm that demographic effects can impact background selection

Using the demographic parameters inferred from weak BGS neutral regions, we simulated the joint effects of demography and BGS on neutral diversity for African, European, and East Asian populations (see Materials and Methods). We also ran an identical set of simulations without BGS in order to generate an expectation of the effect of demography only on neutral diversity. To measure the relative impact of BGS for each population, we then took the ratio of neutral diversity from BGS simulations (π) and neutral diversity from simulations without BGS (π_0) to calculate relative diversity (π/π_0). As expected, we found that BGS reduced relative diversity ($\pi/\pi_0 < 1$) for all three populations in our simulations. However, non-African populations experienced a proportionally larger decrease in π/π_0 compared to the African population (π/π_0 = 0.43, 0.42, 0.41 in AFR, EUR, and EASN respectively). To understand how this dynamic process occurs, we sampled all simulated populations every 100 generations through time to observe the effect of population size change on π/π_0 (Fig 6). We observed that there is a distinct drop in π/π_0 at each population bottleneck experienced by non-Africans, with East Asians suffering the largest drop overall, followed by Europeans. Interestingly, Africans also experienced a large drop in π/π_0 (but less than non-Africans) even though they did not experience any bottlenecks. This was attributable to migration between non-Africans and Africans and this pattern disappeared when we ran simulations using an identical demographic model with BGS but without migration between populations (Fig S7). This finding highlights an evolutionary role that deleterious alleles can play when they are transferred across populations through migration (see Discussion).

Our simulations were based on the functional density found in the region of the

human genome with the strongest effect of BGS (chr3: 48,600,000-50,600,000), where

20.46% of sites were either coding or conserved non-coding (see Materials and Meth-

ods). Thus, the fraction of the genome experiencing deleterious mutation in our simula-

tions of strong BGS (which we define as U) was 0.2046. The patterns we observed in

these simulations likely represent an upper bound on the strength of BGS in the human

genome. We therefore lowered the impact of BGS by reducing U (see Materials and

Methods). When U is decreased 2-4 fold, we continued to observe a stepwise decrease

in π/π_0 similar to our simulations of U = 0.2046, with the specific rank order of African,

followed by European, and then East Asian populations (Fig S8). As expected, π/π_0 in-

creased overall for all populations as the fraction of sites that were simulated as delete-

rious was decreased (π/π_0 = 0.641 vs. 0.802, 0.62 vs. 0.777, and 0.611 vs. 0.777 for

AFR, EUR, and EASN when U = 0.1023 and U = 0.05115, respectively).

Discussion

In our analyses of thousands of genomes from globally distributed human popu-

lations, we have confirmed that neutral variation in humans is governed largely by the

processes of both linked selection and demography. While this observation is not unex-

pected, we have characterized for the first time the dynamic consequences of non-

equilibrium demographic processes in regions of BGS. We find that demography's ef-

fect in these regions do not result in the same relative magnitude of decreased neutral

diversity across populations. Instead, relative diversity in regions of BGS is highly de-

pendent on a population's demographic history, with bottlenecks playing a particularly

strong role. For example, the strongest overall decreases in relative diversity occur in

the East Asian continental group, which is expected if they also suffered the strongest

population bottlenecks. To remove any possible biases that would influence our results,

we controlled for functional effects of mutations, variability in mutation along the ge-

nome, potential sequencing artifacts, GC-based gene conversion, and the potential mu-

tagenic effects of recombination hotspots. None of these factors qualitatively affected

our results.

We do recognize that one caveat of our controls is the fact that divergence itself

is not independent of BGS [63], and this may present biases when using divergence to

control for variation in mutation rate along the genome. This is because the rate of coa-

lescence in the ancestral population of two groups will be faster in regions of strong

BGS compared to regions of weak BGS due to the lower N_e of the former, leading to a

decrease in overall divergence in those regions. To limit the contribution of such biases

in ancestral N_e to divergence, we use rhesus macaque since it is more distantly related

to humans than other primate species such as orangutan or chimpanzee (human-

rhesus divergence: 29.6 MYA; [64]). However, biases in divergence due to BGS can be

still be observed between species as distantly related as human and mouse [63] (hu-

man-mouse divergence: 75 MYA; [65]). Despite this, the apparent correlation between

BGS and divergence should not qualitatively affect our results of relative diversity. Ra-

ther, it is likely that differences in normalized diversity between different BGS bins are

greater than what we observe here. Normalizing by the lower divergence that is charac-

teristic of strong BGS bins and the higher divergence that is characteristic of weak BGS

bins should make any differences between the two smaller, not greater. In fact, for our

calculations of relative diversity in which we skip the normalization step, the differences

in diversity between the strongest 1% and weakest 1% BGS bins are greater and give a

lower ratio of relative diversity (π/π_0 for AFR is 0.373 without the divergence step and

0.402 with the divergence step). A similar pattern is also observed for other continental

groups (compare Fig 2 and Fig S5). More importantly though, we should not expect the

potential biases of our divergence step to contribute to the differences in relative diversi-

ty between each of the continental groups since biases in divergence across the ge-

nome should impact all human populations equally.

We also note that the estimates of B by McVicker et al. [6] may be biased by

model assumptions concerning mutation rates and the specific sites subject to purifying

selection, with the exact values of B also unlikely to be precisely inferred. However, a

similar study by Comeron et al. [29] that investigated BGS in *Drosophila* and utilized the

same model of BGS as McVicker et al. found that biases presented by model assump-

tions or mis-inference on the exact value of B do not significantly change the overall

rank order for the inferred strength of BGS across the genome. Thus we, expect

McVicker et al.'s inference of B to provide good separation between the weakest and

strongest regions of BGS in the human genome, with model misspecification unlikely to

change our empirical results.

The increased population differentiation we observe in regions of strong BGS are

consistent with the increased action of genetic drift occurring in genomic regions with

characteristically low N_e, which has been predicted theoretically [66,67] and observed in

previous studies [57,68,69]. Here, we replicate this pattern by measuring F_{ST} as a func-

tion of B across a global set of populations, providing a rich description of global human

population differentiation across the genome. Furthermore, unlike previous studies of

F_{ST} in humans that have been plagued by biases due to SNP ascertainment or allele

frequency [56], we make use of SNP data from publically available KhoeSan genomes

to select a set of truly informative SNPs for estimating F_{ST}. Our results show that popu-

lation differentiation can be dramatically influenced by BGS, with estimates of F_{ST} at

neutral sites in the weakest 2% BGS bin and strongest 2% BGS bin in the genome dif-

fering by 23.6%-70.1% when we compare all pairs of continental groups (Fig 4A). When

accounting for differences using recombination rate alone, pairs of continental groups

showed little change at all (Fig 4B), thus demonstrating the added value of using explicit

models of linked selection when resolving properties of neutral variation across the ge-

nome.

While we focus specifically on the effects of BGS on generating local patterns of

diversity across humans, there is still a possibility that the effects of positive selection,

through recurrent-hitchhiking, soft-sweeps, and classic selective sweeps, are contrib-

uting to these patterns as well. Models taking into account both the effects of purifying

selection and positive selection will be necessary to provide a full description of the im-

pact of linked selection across the human genome and its behavior as a function of dif-

ferent population demographic histories [42]. Development of mathematical models

characterizing the joint effects of both BGS and genetic hitchhiking [24] as well as their

recent application to different species [30,44], gives promise towards this goal. Recent-

ly, a joint model of classic selective sweeps and BGS was applied to *Drosophila* and

predicted that BGS has had a ~1.6 to 2.5-fold greater effect on neutral genetic diversity

than classic selective sweeps [30]. We should expect this magnitude to be even greater

for humans, since classic selective sweeps were found to be rare in recent human evo-

lution [38] and adaptive substitutions in the human genome are much less frequent

than *Drosophila* [5,70,71]. Despite this, we still attempt to alleviate any contribution from

other modes of linked selection by masking out regions of hard-sweeps or soft-sweeps

using a novel H12-like method [72] that uses an integrated haplotype homozygosity

framework [36] to scan for population specific signatures of these events in the genome

([73]; S2 Appendix). While there may still be some residual contribution of these events

to our estimates of π/π_0, if such signatures do exist, they are likely to localize where

BGS is also expected to act. Thus, it is possible that our empirical results describe the

consequences of non-equilibrium demography in the broader context of both positive

and purifying modes of linked selection. Indeed, our simulations of BGS fail to capture

the complete effects of linked selection on reducing π/π_0 in different human populations

(compare Figs 2B and 6C) and the additional contribution of hitchhiking, which we did

not simulate, may explain this discrepancy.

Non-equilibrium demography has also been been recently investigated with

regard to its impact on patterns of deleterious variation across human populations. Initial

work comparing non-synonymous and synonymous sites in European-Americans and

African-Americans found that, while African-Americans have a greater number of

segregating non-synoymous and synonymous sites than European-Americans,

European-Americans have an overall greater amount of non-synonymous and

synonymous homozygous derived alleles [50]. Other work showed similar findings

[62,74], with Henn et al. [75] demonstrating that the number of deleterious derived

alleles per genome increased with distance from Africa. In addition, similar results in

human founder populations [76,77], *Arabadopsis* [78], and domesticated species such

as dogs [12] and sunflowers [79] further demonstrate the pervasive impact that

demography has on influencing the relative amount of deleterious variation across a

variety of populations and species. Since BGS itself is a function of deleterious

variation, it is not surprising that we also witness differences in π/π_0 across human

populations that have experienced different demographic histories. These effects are

likley ubiquitous across other species as well. However, there has been recent

contention about whether the previously described patterns of increased deleterious

variation are driven by a decrease in the efficacy of natural selection or are solely

artifacts of the response of deleterious variation to demographic change [51,52,80–82].

Following a strong population bottleneck, a transient increase in the relative ratio of

deleterious variation to neutral variation is expected, that is independent of the efficacy

of selection [47,48]. Since our own investigation focuses solely on neutral variants, we

can only invoke differences in population demography and drift on causing the

differences between populations, supporting the conjecture that demographic processes

may transiently dominate patterns of diversity at sites that may be under the direct

influence (or in the case of BGS, indirect influence) of purifying selection.

Recently, Koch et al. [49] investigated the temporal dynamics of human

demography on selected sites and observed that after a population contraction,

heterozygosity at selected sites can undershoot its expected value at equilibrium as

low-frequency variants are lost at a quicker rate before the recovery of intermediate

frequency variants can occur. In the context of BGS, which skews the site frequency

spectrum of linked neutral mutations towards rare variants [23,83], we also expect a

transient decrease in diversity as low-frequency variants are lost quickly during a

population contraction. This effect may be driving the results of the greater losses of

relative diversity in non-African populations, but further work is needed to better

understand the impact of population contractions on perturbing the site-frequency

spectrum of regions under BGS. Koch et al. demonstrate that this effect is only

temporary and that long-term diversity at selected sites approaches greater values once

equilibrium is reached. We stress that these temporal effects also impact the patterns of

neutral diversity caused by BGS and likely explain the differences of π/π_0 that we

observed in humans. We predict that as evolution continues forward through time, the

magnitude and rank order of relative diversity among populations may change.

However, this will be highly dependent on the demographic effects that different

populations experience. Although more flexible modeling frameworks that can account

for BGS under changing population size have been recently developed [45,46], they as-

sume that selection is strong enough that the impact of demography does not perturb

mutation-selection balance, and they also ignore the additional impact that demography

itself has on perturbing neutral variants.

A greater contemporary N_e in non-Africans could theoretically result in a greater

efficacy of purifying selection and, consequently, a stronger impact of BGS. However, it

is very unlikely that this is occuring or driving the observed patterns of relative diversity

that we see. The greater contemporary population size of non-Africans has transpired

only in the very recent past, with accelerated growth in Europeans occurring within the

last few hundred generations [62,84–87]. Thus, most population specific mutations in

non-Africans have arisen only during this recent population expansion and are both

young and very rare [88,89]. It is doubtful that these population-specific variants have

had enough time to exercise a large effect on the absolute strength of BGS itself.

Rather, our simulations indicate that the response of π in regions under BGS is driven

by population contractions, with reductions in π/π_0 occurring concomitantly with the out-

of-Africa bottleneck and European-East Asian split bottleneck events (Fig 6) and contin-

uing even after the European and East Asian expansion events. Interestingly though,

our simulations reveal an additional factor that can influence the impact of BGS within

populations – migration between populations. We observe that the exchange of delete-

rious variants between populations can impact linked selection and lead to decreases in

π/π_0 in populations of constant size, such as the recent history inferred for Africans (Fig

6B). In particular, a decrease in π/π_0 for Africans coincides with decreases in π/π_0 for

non-Africans. This trend disappears, though, once migration is removed (Fig S7). This

result suggests that the migration of nearly neutral or mildly deleterious alleles in Euro-

peans/East Asians into Africans results in those variants being more strongly selected in

Africans due to their historically larger N_e, resulting in stronger BGS and lower π. How-

ever, more work is needed to definitively test this. Conversely, migration of African al-

leles into Europeans/East Asians has no noticeable effect on π in our simulations with

BGS (compare solid lines in Fig 6B and Fig S7B). In simulations where U is lower than

our base level of 0.2046, we also witness differences in π/π_0 between simulations with

and without migration. Some of these involve rank order changes in π/π_0 between Afri-

cans, Europeans, and East Asians (Fig S8, $U = 0.02046$ with migration vs. no migra-

tion), which is likely a consequence of the increased stochasticity that migration be-

tween populations introduces, as simulations without migration restored the consistent

patterns of π/π_0. These results demonstrate the complexities that migration introduces to

patterns of diversity in regions of linked selection and to the dynamic changes of π/π_0

over time.

While we describe here the differential effects of non-equilibrium demography on

neutral diversity in regions under strong and weak BGS, it is worth mentioning that dif-

ferences in the reduction of neutral diversity in the genome between different popula-

tions have also been investigated at the level of entire chromosomes. In particular,

analyses of neutral diversity comparing autosomes to non-autosomes (i.e., sex chromo-

somes and the mitochondrial genome [mtDNA]) have been conducted. Interestingly,

these studies have shown that population contractions have impacted the relative re-

duction of neutral diversity between non-autosomes and autosomes in a similar fashion

to what we have observed between regions of strong BGS and weak BGS, with the

greatest losses occurring in bottlenecked populations. This was demonstrated in both

humans and *Drosophila* [90,91] and later modeled [92], showing that stronger genetic

drift due to the lower N_e of non-autosomes causes diversity to be lost more quickly in

response to population size reductions. Recent work in humans has confirmed such

predictions by showing that relative losses of neutral diversity in the non-autosomes are

greatest for non-Africans [93–95]. These studies, plus others [96], have also shown that

there is strong evidence for a more dominant effect of linked selection on the sex chro-

mosomes relative to the autosomes in humans.

Since linked selection is a pervasive force in shaping patterns of diversity across

the genomes in a range of biological species [1], it has been provided as an argument

for why neutral diversity and estimates of N_e are relatively constrained across species in

spite of the large variance in census population sizes that exist [44,97]. However, since

population bottlenecks are common among species and have an inordinate influence on

N_e [20], demography has also been argued as a major culprit for constrained diversity

[2,97–99]. Yet, as we show in humans, it is likely that patterns of neutral diversity are in

fact jointly impacted by both of these forces, magnifying one another to deplete levels of

diversity beyond what is expected by either one independently. In our work, we also

identify a potentially substantial role for migration from smaller populations that harbor

more strongly deleterious alleles on patterns of linked neutral diversity in large popula-

tions. Together, these combined effects may help provide additional clues for the puz-

zling lack of disparity in genetic diversity among different species [100].

Our results give caution that not properly accounting for the impacts of demogra-

phy on patterns driven by linked selection may lead to spurious rejections of certain

models of linked selection in favor of others. For example, when comparing humans to

other primates, a recent study by Nam et al. [101] found a greater relative decrease of

neutral genetic diversity in regions close to genes (where linked selection is greatest)

versus regions far away from genes in non-human primate species. The authors sug-

gest that these patterns are best described by the action of selective sweeps since non-

human primate populations usually exhibit a larger N_e and thus should experience a

greater influx of adaptive alleles per unit time. However, certain orangutan, gorilla, and

chimpanzee species have also experienced large population contractions in their recent

history that are comparable in time and potentially more extreme in scale to what hu-

mans have also experienced [102–105]. Thus, the consequences of strong demograph-

ic change to regions of BGS can provide an alternative explanation for why these pri-

mate species exhibit a larger relative reduction in neutral diversity near genes, despite

their larger historical N_e. While Nam et al. also test for BGS using simulations with de-

mography, their model assumed a more limited population contraction (50% population

reduction) over a shorter time scale than the demographic model we test.

Finally, our results also have implications for human medical genetics research,

since selection may be acting on functional regions contributing to disease susceptibil-

ity. Since different populations will have experienced different demographic histories,

the action of linked selection may result in disparate patterns of genetic variation (with

elevated levels of drift) near causal loci. Recent work has already demonstrated that

BGS's consequence of lowering diversity impacts power for disease association tests

[106]. Our results indicate that this impact may be even further exacerbated by demog-

raphy in bottlenecked populations, leading to potentially larger discrepancies in power

between different populations. Overall, this should encourage further scrutiny for tests

and SNP panels optimized for one population since they may not be easily translatable

to other populations. It should also further motivate investigators to simultaneously ac-

count for demography and linked selection when performing tests to uncover disease

variants within the genome [88,106,107].

Materials and methods

Data

2,504 samples from 26 populations in phase 3 of the Thousand Genomes Project

(TGP) [9] were downloaded from ftp://ftp.ncbi.nlm.nih.gov/1000genomes/. vcftools

(v0.1.12a) [108] and custom python scripts were used to gather all bi-allelic SNP sites

from the autosomes of the entire sample set.

A subset of TGP samples that were sequenced to high coverage (~45X) by

Complete Genomics (CG) were downloaded from

ftp://ftp.ncbi.nlm.nih.gov/1000genomes/. After filtering out related individuals via pedi-

gree analyses, we analyzed 53 YRI, 64 CEU, and 62 CHS samples (Table S2). The

cgatools (v1.8.0) listvariants program was first used to gather all SNPs from the 179

samples using their CG ASM "Variations Files" (CG format version 2.2). Within each

population, the number of reference and alternate allele counts for each SNP was then

calculated using the cgatools testvariants program and custom python scripts. Only al-

lele counts across high quality sites (i.e., those classified as VQHIGH variant quality by

CG) were included. Low quality sites (i.e., those with VQLOW variant quality) were

treated as missing data. Only autosomes were kept. Non-bi-allelic SNPs and sites vio-

lating Hardy-Weinberg equilibrium (HWE) (p-value < 0.05 with a Bonferroni correction

for multiple SNP testing) were also removed.

We collected 13 whole-genome sequenced KhoeSan samples (sequence-

coverage: 2.5-50X, see Table S3 in Supporting information) from 3 studies [75,109,110]

and used the processed vcf files from each of those respective studies to gather all bi-

allelic polymorphic SNPs (i.e., the union of variants across all vcf files). SNPs were only

retained if they were polymorphic within the 13 samples (i.e., sites called as alternate

only within the sample set were ignored).

Filtering and ascertainment scheme

Positions in the genome were annotated for background selection by using the

background selection coefficient, *B*, which was inferred by McVicker et al. [6] and down-

loaded from http://www.phrap.org/othersoftware.html. *B* was inferred by applying a clas-

sical model of BGS [53], which treats its effects as a simple reduction in N_e at neutral

sites as a function of their recombination distance from conserved loci, the strength of

purifying selection at those conserved loci, and the deleterious mutation rate. *B* can be

interpreted as the reduced fraction of neutral genetic diversity at a particular site along

the genome that is caused by BGS, with a value of 0 indicating a near complete remov-

al of neutral genetic diversity due to BGS and a *B* value of 1 indicating little to no effect

of BGS on neutral genetic diversity ($B = \pi/\pi_0 = N_e/N_0$). Positions for *B* were lifted over

from hg18 to hg19 using the UCSC liftOver tool. Sites that failed to uniquely map from

hg18 to hg19 or failed to uniquely map in the reciprocal direction were excluded. Sites

lacking a *B* value were also ignored. We focused our analyses on those regions of the

genome within the top 1%, 5%, 10%, and 25% of the genome-wide distribution of *B* and

within the bottom 1% of the genome-wide distribution of *B*. These quantiles correspond

to the *B* values 0.095, 0.317, 0.463, 0.691, and 0.994, respectively.

A set of 13 filters (referred to as the "13-filter set") were used to limit errors from

sequencing and misalignments with rhesus macaque and to remove regions potentially

under the direct effects of natural selection and putative selective sweeps (we ignore

the linked selection effects of background selection). These filters were applied to all

samples in phase 3 TGP (all filters are in build hg19) for all sets of analyses (see Table

S4 in Supporting information for the total number of Mb that passed the described filters

below for each particular *B* quantile):

1. Coverage/exome: For phase 3 data, regions of the genome that were part of

696   the   high   coverage   exome   were   excluded   (see

ftp://ftp.ncbi.nlm.nih.gov/1000genomes/ftp/technical/reference/exome_pull_do

wn_targets/20130108.exome.targets.bed.README). This was done to limit

biases due to differing levels of coverage across the genome and to remove

likely functional sites within the exome.

2. phyloP: Sites with phyloP [111] scores > 1.2 or < -1.2 were removed to limit

the effects of natural selection due to conservation or accelerated evolution.

Scores   were   downloaded   from

http://hgdownload.cse.ucsc.edu/goldenPath/hg19/phyloP46way/.

3. phastCons: Regions in the UCSC conservation 46-way track (table:

phastCons46wayPlacental) [112] were removed to limit the effects of natural

selection due to conservation.

4. CpG: CpG islands in the UCSC CpG islands track were removed because of

their potential role in gene regulation and/or being conserved.

5. ENCODE blacklist: Regions with high signal artifacts from next-generation

sequencing experiments discovered during the ENCODE project [113] were

removed.

6. Accessible genome mask: Regions not accessible to next-generation se-

quencing using short reads, according to the phase 3 TGP "strict" criteria,

were   removed   (downloaded   from

ftp://ftp.ncbi.nlm.nih.gov/1000genomes/ftp/release/20130502/supporting/acce

ssible_genome_masks/StrictMask/).

7. Simple repeats: Regions in the UCSC simple repeats track were removed

due to potential misalignments with outgroups and/or being under natural se-

lection.

8. Gaps/centromeres/telomeres: Regions in the UCSC gap track were removed,

including centromeres and telomeres.

9. Segmental duplications: Regions in the UCSC segmental dups track [114]

were removed to limit potential effects of natural selection and/or misalign-

ments with rhesus macaque.

10. Transposons: Active transposons (HERVK retrotransposons, the AluY sub-

family of Alu elements, SVA elements, and L1Ta/L1pre-Ta LINEs) in the hu-

man genome were removed.

11. Recent positive selection: Regions inferred to be under hard and soft selec-

tive sweeps (using iHS and iHH12 regions from selscan v1.2.0 [73]; S2 Ap-

pendix) within each phase 3 population were removed.

12. Non-coding transcripts: Non-coding transcripts from the UCSC genes track

were removed to limit potential effects of natural selection.

13. Synteny: Regions that did not share conserved synteny with rhesus macaque

(rheMac2) from UCSC syntenic net filtering were removed (downloaded from

http://hgdownload.soe.ucsc.edu/goldenPath/hg19/vsRheMac2/syntenicNet/).

Additionally, an extra set of filters was applied, but only for those estimates of diversity

that controlled for GC-biased gene conversion and recombination hotspots:

14. GC-biased gene conversion (gBGC): Regions in UCSC phastBias track [115]

from UCSC genome browser were removed to limit regions inferred to be un-

der strong GC-biased gene conversion.

15. Recombination hotspots: All sites within 1.5 kb (i.e., 3 kb windows) of sites

with recombination rates ≥ 10 cM/Mb in the 1000G OMNI genetic maps for

non-admixed populations (downloaded from ftp://ftp-

trace.ncbi.nih.gov/1000genomes/ftp/technical/working/20130507_omni_reco

mbination_rates/) and the HapMap II genetic map [116] were removed. 1.5 kb

flanking regions surrounding the center of hotspots identified by Ref. [117]

(downloaded from

http://science.sciencemag.org/content/sci/suppl/2014/11/12/346.6211.125644

2.DC1/1256442_DatafileS1.txt) were also removed, except for the cases in

which the entire hotspot site was greater than 3 kb in length (in which case

just the hotspot was removed).

To generate a set of four-fold degenerate synonymous sites, all polymorphic

sites that we retained from the high-coverage Complete Genomic samples were anno-

tated using the program ANNOVAR [118] with Gencode V19 annotations. ANNOVAR

and Gencode V19 annotations were also used to gather an autosome-wide set of four-

fold degenerate sites, resulting in 5,188,972 total sites.

Demographic inference

The inference tool dadi (v1.6.3) [7] was used to fit, via maximum likelihood, the 3-

population 13-parameter demographic model of Gutenkunst et al. [7] to the 179 YRI,

CEU, and CHS samples from the high coverage CG dataset of TGP. This sample set

consisted of 53 YRI (African), 64 CEU (European), and 62 CHS (East Asian) samples.

The demographic model incorporates an ancient human expansion in Africa and a sin-

gle out-of-Africa bottleneck followed by European- and East Asian-specific bottlenecks,

as well as exponential growth in both non-African populations and migration between

populations. During the inference procedure, each population was projected down to

106 chromosomes, corresponding to the maximum number of chromosomes available

in the CG YRI population. Sites were polarized with chimpanzee to identify putative an-

cestral/derived alleles using the chain and netted alignments of hg19 with panTro4

(http://hgdownload.soe.ucsc.edu/goldenPath/hg19/vsPanTro4/axtNet/), and the correc-

tion for ancestral misidentification [119] option in dadi was used. The 13-filter set de-

scribed previously was applied to the CG data set, and an additional filter keeping only

the weakest ~1% of autosomal sites under background selection ($B \geq 0.994$) was also

applied in order to mitigate potential biases in inference due to BGS [59,60] or other

forms of linked selection [120]. After site filtering and correction for ancestral misidentifi-

cation, a total of 110,582 segregating sites were utilized by dadi for the inference proce-

dure. For optimization, grid points of 120, 130, and 140 were used, and 15 independent

optimization runs were conducted from different initial parameter points to ensure con-

vergence upon a global optimum. An effective sequence length (L) of 7.15 Mb was cal-

culated from the input sequence data after accounting for the fraction of total sites re-

moved due to filtering. In addition to the 13-filter set, this filtering included sites violating

HWE, sites without B value information, sites that did not have at least 106 sampled

chromosomes in each population, sites with more than two alleles, sites that did not

have tri-nucleotide information for the correction for ancestral misidentification step, and

sites treated as missing data. For calculating the reference effective population size, a

mutation rate (μ) of 1.66×10^{-8} (inferred from Ref. [121]) was used. Using the optimized

θ from dadi after parameter fitting, the equation $\theta = 4N_e\mu L$ was solved for N_e to generate

the reference effective population size, from which all other population N_e's were calcu-

lated. This same procedure was also used to infer demographic parameters from four-

fold degenerate synonymous sites across the same set of samples. After site filtering

(note that B and the 13-filter set were not included in the filtering step for four-fold de-

generate synonymous sites), 41,260 segregating sites were utilized by dadi for the in-

ference procedure, and an effective sequence length of 2.37 Mb was used for calculat-

ing the reference effective population size.

Simulations

Forward simulations incorporating the results from the demographic inference

procedure described above and a model of background selection were conducted using

SFS_CODE [122]. For the model of background selection, the recombination rate, ρ,

and the fraction of the genome experiencing deleterious mutation, U, was calculated us-

ing the 2 Mb region of chr3: 48,600,000-50,600,000, which has been subject to the

strongest amount of BGS in the human genome (mean B = 0.002). A population-scaled

recombination rate (ρ) of 6.0443 x 10^{-5} was calculated for this region using the HapMap

II GRCh37 genetic map [116]. For generating U, the number of non-coding "functional"

sites in this region was first calculated by taking the union of all phastCons sites and

phyloP sites with scores > 1.2 (indicating conservation) that did not intersect with any

coding exons. This amount totaled to 270,348 base pairs. Additionally, the number of

coding sites was calculated by summing all coding exons within this region from GEN-

CODE v19, which totaled to 138,923 base pairs. From these totals, a U of 0.2046 was

generated.

The background selection model was simulated using a middle 30 kb neutral re-

gion flanked by two 1 Mb regions under purifying selection. From the calculated U de-

scribed above, 20.46% of sites in the two 1 Mb flanking regions were simulated as be-

ing deleterious. Two distributions of fitness effects were used for the deleterious sites,

with 66.06% of deleterious sites using the gamma distribution of fitness effects inferred

across conserved non-coding regions by Ref. [123] (β = 0.0415, α = 0.00515625) and

33.94% of deleterious sites using the gamma distribution of fitness effects inferred

across coding regions by Ref. [5] (β = 0.184, α = 0.00040244). The relative number of

sites receiving each distribution of fitness effects in our simulations was determined by

the relative number of non-coding "functional" sites and coding exons described above.

Gamma distribution parameters are scaled to the ancestral population size of the de-

mographic models used in Refs. [5,123]. To simulate varying levels of background se-

lection strength, different total fractions of our calculated U were used (5%, 10%, 25%,

50%, and 100% of 0.2046) but with the same relative percentage of non-coding and

coding sites just described. To simulate only the effects of demography without back-

ground selection (i.e., U = 0), only the 30 kb neutral region was simulated. 2,000 inde-

pendent simulations were conducted for each particular U (12,000 total). Simulations

output population genetic information every 100 generations and also at each genera-

tion experiencing a population size change (22,117 total generations were simulated),

for which mean pairwise nucleotide diversity (π) was calculated across the 2,000 simu-

lations.

Calculating recombination rate across the genome

We used recombination rate estimates from the HapMap II GRCh37 genetic

map. To annotate sites in phase 3 that were not in HapMap II, recombination rates were

interpolated to the midway point between the preceding and following positions in Hap-

Map II. If the difference between successive HapMap II positions was greater than

18,848 base pairs (the first standard deviation for the distribution of distances between

positions in HapMap II), then the recombination rate was only extended out 9,424 base

pairs beyond the focal position. Positions beyond this distance were then ignored during

analysis in which the recombination rate was used. Recombination rate quantiles were

calculated using the genome-wide distribution of recombination rates (i.e., the distribu-

tion of recombination rates across all sites, including those that are not polymorphic in

the data set) resulting from the procedure described above.

Population-specific calculations of diversity

Mean pairwise genetic diversity (π) was calculated as a function of the B quantile

bins described in "Filtering and ascertainment scheme" for each of the 20 non-admixed

populations in phase 3 TGP and across 4 broad populations that grouped the 20 non-

admixed populations together by continent (African, European, South Asian, and East

Asian, see Table S11 in Supporting information). Additionally, only regions of the ge-

nome passing the 13-filter set were used in the diversity calculations (see Table S4 in

Supporting information for total number of Mb used in diversity calculations for each B

quantile). For estimates of diversity controlling for gBGC or recombination hotspots, the

additional corresponding filters described in "Filtering and ascertainment scheme" were

also used. Only 100 kb regions of the genome with at least 10 kb of divergence infor-

mation with Rhesus macaque were used in the diversity calculations (see "Normaliza-

tion of diversity and divergence calculations with Rhesus macaque" below).

Ancestry specific calculations of diversity

To calculate genetic diversity as a function of local ancestry across the admixed

phase 3 population samples, we used the ancestry deconvolution results generated by

the   1000   Genomes   Project   Consortium   (see

ftp://ftp.ncbi.nlm.nih.gov/1000genomes/ftp/technical/working/20140818_ancestry_decon

volution/README_20140721_phase3_ancestry_deconvolution).   Briefly,   the   local-

ancestral inference tool, RFMix [124], was run across the ACB, ASW, CLM, MXL, PEL,

and PUR phase 3 TGP samples. For the reference panel, 50 unrelated shapeit2 [125]

trio-phased YRI and CEU samples each (from phase 3 TGP) and 43 shapeit2 popula-

tion-phased Native American samples (from Ref. [126]) were used. We utilized local an-

cestry tracks that were inferred by RFMix using "trio-phased" mode.

Admixed samples were then parsed for all genomic segments homozygous for

each particular ancestry (i.e., African, European, or Native American). These homozy-

gous segments were also filtered according to the 13-filter set described in "Filtering

and ascertainment scheme." Heterozygosity was calculated across admixed samples

for each set of homozygous ancestries and *B* quantile bins described previously. Sam-

ples were included in the analyses only if the total length of their genome that passed all

filters for the particular ancestry and *B* quantile bin was greater than 1 Mb. Additionally,

only 100 kb segments that had at least 10 kb of divergence information with Rhesus

macaque were used (see below). Per-site heterozygosity estimates for each ancestry

and *B* quantile set were averaged across all admixed samples, regardless of their TGP

population of origin. See Table S5 in Supporting information for total number of Mb used

in the analyses. Additionally, heterozygosity was also calculated across the 4 continen-

tal groups using the same 13-filter set and as a function of the same *B* quantile bins.

Normalization of diversity and divergence calculations with Rhesus macaque

To calculate human divergence with Rhesus macaque, we downloaded the

syntenic net alignments between hg19 and rheMac2 that were generated by blastz from

http://hgdownload.cse.ucsc.edu/goldenpath/hg19/vsRheMac2/syntenicNet/. We binned

the human genome into non-overlapping 100 kb bins and calculated divergence within

each bin by taking the proportion of base pair differences between human and Rhesus

macaque. Gaps between human and Rhesus macaque, positions lacking alignment in-

formation, and positions that did not pass the 13-filter set described in "Filtering and

ascertainment scheme" were ignored in the divergence estimate. Additionally, a sepa-

rate set of divergence estimates were also made using the additional set of filtering cri-

teria that removed those regions under gBGC or in recombination hotspots and were

used for normalizing diversity in those measurements that controlled for gBGC and

hotspots.

When normalizing diversity by divergence, only 100 kb bins that had at least 10

kb of divergence information were used (21,100 bins total for 13-filter set, 20,935 bins

total for the 13-filter set plus the additional gBGC and hotspot filters). Bins with less than

10 kb of divergence information were ignored. To make estimates comparable, in those

measurements of diversity that did not normalize by divergence, diversity was still calcu-

lated using the same set of 100 kb bins that had at least 10 kb for estimating diver-

gence.

Calculations of population differentiation (F_{ST}) and linear regression

F_{ST} calculations were performed as a function of B and recombination rate be-

tween every pair of non-admixed phase 3 TGP populations not belonging to the same

continental group (150 pairs total). We followed the recommendations in Bhatia et al.

[56] to limit biases in F_{ST} due to 1) type of estimator used, 2) averaging over SNPs, and

3) SNP ascertainment. Specifically, we 1) used the Hudson-based F_{ST} estimator [127],

2) used a ratio of averages for combining F_{ST} estimated across different SNPs, and 3)

ascertained SNPs based on being polymorphic in an outgroup (i.e., the KhoeSan). For

ascertaining SNPs in the KhoeSan, we also performed filtering according to the filtering

scheme described under "Filtering and ascertainment scheme." For a position to be

considered polymorphic in the KhoeSan, at least one alternate allele and one reference

allele had to be called across the 13 genomes we utilized (see "Data"). These criteria

left 3,497,105 total sites in the genome in the phase 3 dataset for F_{ST} to be estimated

across.

F_{ST} was calculated across either 2% quantile bins of B or 2% quantile bins of

recombination rate in order to perform simple linear regression, with either B or

recombination rate acting as an explanatory variable using the linear model $F_{ST} = \beta_0 +$

$\beta_1 X + \varepsilon$ (where X represents either B or recombination rate [ρ]). This was done for all

pairwise comparisons of populations between a specific pair of continental groups (25

pairs total) or across all pairwise comparisons using all continental groups (150 pairs

total). Recombination rate was scaled to be between 0 and 1 (the minimum and maxi-

mum observed recombination rate was 0.0 cM/Mb and 126.88 cM/Mb, respectively) to

aid in the comparison of the regression coefficient with B. Additionally, the mean of the

bounds defining each quantile bin was used when defining the explanatory variables for

the regression. Linear regression, robust linear regression, and simple correlation were

performed using the lm(), rlm(), and cor() functions, respectively, in the R programming

language (www. r-project.org). To generate standard errors of the mean, this same pro-

cedure was performed on F_{ST} results generated from each of 1,000 bootstrapped

iterations of the data.

F_{ST} was also calculated across bins defined by both B and recombination rate

(ρ). For this step, bins where first defined into 2% quantile bins of either variable. Each

2% bin was further defined into 4% quantile bins of the other variable, generating 1,250

separate bins in which F_{ST} was estimated for both B and recombination rate

simultaneosly. Multiple linear regression and robust regression was performed for the

model $F_{ST} = \beta_0 + \beta_1 B + \beta_2 \rho + \beta_3 B\rho + \varepsilon$ with the lm() and rlm() functions in R. As with the

simple linear regression step, recombination rate was scaled to be between 0 and 1 and

the mean of the bounds defining each quantile bin was used when defining the explana-

tory variables.

Bootstrapping

Diversity Estimates. To control for the structure of linkage disequilibrium and correla-

tion between SNPs along the genome, we partitioned the human genome into non-

overlapping 100 kb bins (these bins were identical to the 100 kb bins used for estimat-

ing divergence) and calculated mean pairwise diversity (π) or heterozygosity within each

bin. We also normalized the diversity estimates by divergence within each bin. We then

bootstrapped individual genomes by sampling, with replacement, the 100 kb bins until

the number of sampled bins equaled the number of bins used for calculating the diversi-

ty point estimates (i.e., 21,100 bins or 20,935 bins total, depending on whether filters for

gBGC and hotspots were applied). 1,000 total bootstrap iterations were completed and

standard errors of the mean were calculated by taking the standard deviation from the

resulting bootstrap distribution.

F_{ST}. For bootstrapping F_{ST}, the human genome was partitioned into non-overlapping

100 kb bins and were sampled with replacement until 28,823 bins were selected (the

total number of non-overlapping 100 kb bins in the human autosomes). F_{ST} was then

calculated genome-wide for the bootstrapped genome as a function of either B and/or

recombination rate for every pairwise comparison of non-admixed phase 3 TGP popula-

tions not belonging to the same continental group. 1,000 total bootstrap iterations were

completed and standard errors of the mean were calculated by taking the standard de-

viation from the F_{ST} distribution calculated from all 1,000 iterations.

Acknowledgements

969   We thank Lawrence Uricchio, Dominic Tong, Melissa Spear, and Nicolas Strauli for

970   helpful comments on the manuscript. The computations in this paper were run on the

971   QB3 Shared Cluster at the University of California, San Francisco.

Author contributions

974 Conceived and designed the experiments: RT SAZ RDH.

975 Performed the experiments: RT SAZ.

976 Analyzed the data: RT SAZ.

977 Wrote the paper: RT RDH.

978 Wrote S2 Appendix: SAZ.

References

1.   Cutter AD, Payseur BA. Genomic signatures of selection at linked sites: unifying
the disparity among species. Nat Rev Genet. Nature Publishing Group; 2013;14:
262–74. doi:10.1038/nrg3425
2.   Ellegren H, Galtier N. Determinants of genetic diversity. Nat Rev Genet. 2016;17:
422–433. doi:10.1038/nrg.2016.58
3.   Sabeti PC, Reich DE, Higgins JM, Levine HZP, Richter DJ, Schaffner SF, et al.
Detecting recent positive selection in the human genome from haplotype
structure. Nature. Nature Publishing Group; 2002;419: 832–837.
doi:10.1038/nature01027.1.
4.   Williamson SH, Hernandez R, Fledel-Alon a, Zhu L, Nielsen R, Bustamante CD.
Simultaneous inference of selection and population growth from patterns of
variation in the human genome. Proc Natl Acad Sci U S A. National Academy of
Sciences; 2005;102: 7882–7887. doi:10.1073/pnas.0502300102
5.   Boyko AR, Williamson SH, Indap AR, Degenhardt JD, Hernandez RD, Lohmueller
KE, et al. Assessing the evolutionary impact of amino acid mutations in the
human genome. PLoS Genet. 2008;4: e1000083.
doi:10.1371/journal.pgen.1000083
6.   McVicker G, Gordon D, Davis C, Green P. Widespread genomic signatures of
natural selection in hominid evolution. PLoS Genet. 2009;5: e1000471.
doi:10.1371/journal.pgen.1000471
7.   Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Inferring the
joint demographic history of multiple populations from multidimensional SNP
frequency data. PLoS Genet. 2009;5. doi:10.1371/journal.pgen.1000695
8.   Li H, Durbin R. Inference of human population history from individual whole-
genome sequences. Nature. 2011;475: 493–496. doi:10.1038/nature10231
9.   Auton A, Abecasis GR, Altshuler DM, Durbin RM, Abecasis GR, Bentley DR, et al.
A global reference for human genetic variation. Nature. 2015;526: 68–74.
doi:10.1038/nature15393
10. Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. A thousand fly
genomes: An expanded drosophila genome nexus. Mol Biol Evol. Oxford

University Press; 2016;33: 3308–3313. doi:10.1093/molbev/msw195

11. Gibbs RA, Taylor JF, Van Tassell CP, Barendse W, Eversole KA, Gill CA, et al.
Genome-Wide Survey of SNP Variation Uncovers the Genetic Structure of Cattle
Breeds [Internet]. NIH Public Access; 2009 Apr 24 pp. 528–532.
doi:10.1126/science.1167936

12. Marsden CD, Ortega-Del Vecchyo D, O'Brien DP, Taylor JF, Ramirez O, Vilà C,
et al. Bottlenecks and selective sweeps during domestication have increased
deleterious genetic variation in dogs. Proc Natl Acad Sci. 2016;113: 152–157.
doi:10.1073/pnas.1512501113

13. Caicedo AL, Williamson SH, Hernandez RD, Boyko A, Fledel-Alon A, York TL, et
al. Genome-wide patterns of nucleotide polymorphism in domesticated rice. PLoS
Genet. Sinauer Associates; 2007;3: 1745–1756.
doi:10.1371/journal.pgen.0030163

14. Begun DJ, Aquadro CF. African and North American populations of Drosophila
melanogaster are very different at the DNA level. Nature. 1993;365: 548–550.
doi:10.1038/365548a0

15. Haddrill PR, Thornton KR, Charlesworth B, Andolfatto P. Multilocus patterns of
nucleotide variability and the demographic and selection history of Drosophila
melanogaster populations. Genome Res. Cold Spring Harbor Laboratory Press;
2005;15: 790–799. doi:10.1101/gr.3541005

16. Ometto L, Glinka S, De Lorenzo D, Stephan W. Inferring the effects of
demography and selection on Drosophila melanogaster populations from a
chromosome-wide scan of DNA variation. Mol Biol Evol. Nature Publishing Group;
2005;22: 2119–2130. doi:10.1093/molbev/msi207

17. Hernandez RD, Hubisz MJ, Wheeler DA, Smith DG, Ferguson B, Rogers J, et al.
Demographic histories and patterns of linkage disequilibrium in Chinese and
Indian rhesus macaques. Science. 2007;316: 240–3.
doi:10.1126/science.1140462

18. Ramachandran S, Deshpande O, Roseman CC, Rosenberg N a, Feldman MW,
Cavalli-Sforza LL. Support from the relationship of genetic and geographic
distance in human populations for a serial founder effect originating in Africa. Proc
Natl Acad Sci U S A. 2005;102: 15942–15947. doi:10.1073/pnas.0507611102

19. Henn BM, Cavalli-Sforza LL, Feldman MW. The great human expansion. Proc
Natl Acad Sci U S A. 2012;109: 17758–64. doi:10.1073/pnas.1212380109

20. Charlesworth B. Fundamental concepts in genetics: effective population size and
patterns of molecular evolution and variation. Nat Rev Genet. 2009;10: 195–205.
doi:10.1038/nrg2526

21. Charlesworth D. Balancing selection and its effects on sequences in nearby
genome regions. PLoS Genet. Benjamin/Cummings; 2006;2: 379–384.
doi:10.1371/journal.pgen.0020064

22. Maynar Smith J, Haigh J. The hitch-hiking effect of a favourable gene. Genet Res.
1974;23: 23–35. doi:10.1017/S0016672308009579

23. Charlesworth B, Morgan MT, Charlesworth D. The effect of deleterious mutations
on neutral molecular variation. Genetics. 1993;134: 1289–303. Available:
http://www.ncbi.nlm.nih.gov/pubmed/8375663

24. Kim Y, Stephan W. Joint effects of genetic hitchhiking and background selection

on neutral variation. Genetics. 2000;155: 1415–1427. doi:drosophila-
melanogaster; deleterious mutations; recombination; polymorphism; rates;
population; evolution; variance; linkage; locus

25. Begun DJ, Aquadro CF. Levels of naturally occurring DNA polymorphism
correlate with recombination rates in D. melanogaster. Nature. Nature Publishing
Group; 1992;356: 519–520. doi:10.1038/356519a0

26. Charlesworth B. Background selection and patterns of genetic diversity in
Drosophila melanogaster. Genet Res. 1996;68: 131–149.
doi:10.1017/S0016672300034029

27. Andolfatto P. Hitchhiking effects of recurrent beneficial amino acid substitutions in
the Drosophila melanogaster genome. Genome Res. Cold Spring Harbor
Laboratory Press; 2007;17: 1755–1762. doi:10.1101/gr.6691007

28. Sella G, Petrov DA, Przeworski M, Andolfatto P. Pervasive natural selection in the
Drosophila genome? [Internet]. Nachman MW, editor. PLoS Genetics. Plenum;
2009. p. e1000495. doi:10.1371/journal.pgen.1000495

29. Comeron JM. Background Selection as Baseline for Nucleotide Variation across
the Drosophila Genome. Begun DJ, editor. PLoS Genet. 2014;10: e1004434.
doi:10.1371/journal.pgen.1004434

30. Elyashiv E, Sattath S, Hu TT, Strutsovsky A, McVicker G, Andolfatto P, et al. A
Genomic Map of the Effects of Linked Selection in Drosophila. Barton NH, editor.
PLoS Genet. Public Library of Science; 2016;12: e1006130.
doi:10.1371/journal.pgen.1006130

31. Flowers JM, Molina J, Rubinstein S, Huang P, Schaal BA, Purugganan MD.
Natural selection in gene-dense regions shapes the genomic pattern of
polymorphism in wild and domesticated rice. Mol Biol Evol. Oxford University
Press; 2012;29: 675–687. doi:10.1093/molbev/msr225

32. Xu X, Liu X, Ge S, Jensen JJDJJDJ, Hu F, Li X, et al. Resequencing 50
accessions of cultivated and wild rice yields markers for identifying agronomically
important genes. Nat Biotechnol. 2012;30: 105–11. doi:10.1038/nbt.2050

33. Andersen EC, Gerke JP, Shapiro JA, Crissman JR, Ghosh R, Bloom JS, et al.
Chromosome-scale selective sweeps shape Caenorhabditis elegans genomic
diversity. Nat Genet. 2012;44: 285–90. doi:10.1038/ng.1050

34. Cutter AD, Payseur BA. Selection at linked sites in the partial selfer
Caenorhabditis elegans. Mol Biol Evol. 2003;20: 665–673.
doi:10.1093/molbev/msg072

35. Reed FA, Akey JM, Aquadro CF. Fitting background-selection predictions to
levels of nucleotide variation and divergence along the human autosomes.
Genome Res. 2005;15: 1211–1221. doi:10.1101/gr.3413205

36. Voight BF, Kudaravalli S, Wen X, Pritchard JK. A map of recent positive selection
in the human genome. Hurst L, editor. PLoS Biol. Public Library of Science;
2006;4: 0446–0458. doi:10.1371/journal.pbio.0040072

37. Cai JJ, Macpherson JM, Sella G, Petrov DA. Pervasive hitchhiking at coding and
regulatory sites in humans. PLoS Genet. 2009;5: e1000336.
doi:10.1371/journal.pgen.1000336

38. Hernandez RD, Kelley JL, Elyashiv E, Melton SC, Auton A, McVean G, et al.
Classic Selective Sweeps Were Rare in Recent Human Evolution. Science (80-).

1103 2011;331: 920–924. doi:10.1126/science.1198878

1104 39. Lohmueller KE, Albrechtsen A, Li Y, Kim SY, Korneliussen T, Vinckenbosch N, et
1105  al. Natural selection affects multiple aspects of genetic variation at putatively
1106  neutral sites across the human genome. PLoS Genet. 2011;7: e1002326.
1107  doi:10.1371/journal.pgen.1002326

1108 40. Alves I, Šrámková Hanulová A, Foll M, Excoffier L. Genomic data reveal a
1109  complex making of humans [Internet]. Schierup MH, editor. PLoS Genetics. Public
1110  Library of Science; 2012. p. e1002837. doi:10.1371/journal.pgen.1002837

1111 41. Granka JM, Henn BM, Gignoux CR, Kidd JM, Bustamante CD, Feldman MW.
1112  Limited evidence for classic selective sweeps in African populations. Genetics.
1113  2012;192: 1049–1064. doi:10.1534/genetics.112.144071

1114 42. Enard D, Messer PW, Petrov DA. Genome-wide signals of positive selection in
1115  human evolution. Genome Res. Cold Spring Harbor Laboratory Press; 2014;24:
1116  885–895. doi:10.1101/gr.164822.113

1117 43. Bank C, Ewing GB, Ferrer-Admettla A, Foll M, Jensen JD. Thinking too positive?
1118  Revisiting current methods of population genetic selection inference. Trends
1119  Genet. Elsevier Ltd; 2014;30: 540–546. doi:10.1016/j.tig.2014.09.010

1120 44. Corbett-Detig RB, Hartl DL, Sackton TB. Natural Selection Constrains Neutral
1121  Diversity across A Wide Range of Species. Barton NH, editor. PLoS Biol.
1122  Springer; 2015;13: e1002112. doi:10.1371/journal.pbio.1002112

1123 45. Zeng K. A coalescent model of background selection with recombination,
1124  demography and variation in selection coefficients. Heredity (Edinb). Nature
1125  Publishing Group; 2013;110: 363–71. doi:10.1038/hdy.2012.102

1126 46. Nicolaisen LE, Desai MM. Distortions in Genealogies due to Purifying Selection
1127  and Recombination. Genetics. 2013;195: 221–230.
1128  doi:10.1534/genetics.113.152983

1129 47. Pennings PS, Kryazhimskiy S, Wakeley J. Loss and Recovery of Genetic
1130  Diversity in Adapting Populations of HIV. Fraser C, editor. PLoS Genet. Public
1131  Library of Science; 2014;10: e1004000. doi:10.1371/journal.pgen.1004000

1132 48. Brandvain Y, Wright SI. The Limits of Natural Selection in a Nonequilibrium World.
1133  Trends Genet. Elsevier Ltd; 2016;32: 201–210. doi:10.1016/j.tig.2016.01.004

1134 49. Koch E, Novembre J. A Temporal Perspective on the Interplay of Demography
1135  and Selection on Deleterious Variation in Humans. G3 (Bethesda). 2017;7:
1136  g3.117.039651. doi:10.1534/g3.117.039651

1137 50. Lohmueller KE, Indap AR, Schmidt S, Boyko AR, Hernandez RD, Hubisz MJ, et
1138  al. Proportionally more deleterious genetic variation in European than in African
1139  populations. Nature. 2008;451: 994–997. doi:10.1038/nature06611

1140 51. Simons YB, Turchin MC, Pritchard JK, Sella G. The deleterious mutation load is
1141  insensitive to recent population history. Nat Genet. 2014;46: 220–4.
1142  doi:10.1038/ng.2896

1143 52. Do R, Balick D, Li H, Adzhubei I, Sunyaev S, Reich D. No evidence that selection
1144  has been less effective at removing deleterious mutations in Europeans than in
1145  Africans. Nat Genet. 2015;47: 126–131. doi:10.1038/ng.3186

1146 53. Nordborg M, Charlesworth B, Charlesworth D. The effect of recombination on
1147  background selection. Genet Res. 1996;67: 159–74.
1148  doi:10.1017/S0016672300033619

54. Moreno-Estrada A, Gignoux CR, Fernández-López JC, Zakharia F, Sikora M,
Contreras A V, et al. The genetics of Mexico recapitulates Native American
substructure and affects biomedical traits. Science (80-). 2014;344: 1280–5.
doi:10.1126/science.1251688
55. Hey J. On the number of new world founders: A population genetic portrait of the
peopling of the Americas. Clark AG, editor. PLoS Biol. Springer; 2005;3: 0965–
0975. doi:10.1371/journal.pbio.0030193
56. Bhatia G, Patterson N, Sankararaman S, Price AL. Estimating and interpreting
FST: The impact of rare variants. Genome Res. 2013;23: 1514–1521.
doi:10.1101/gr.154831.113
57. Keinan A, Reich D. Human population differentiation is strongly correlated with
local recombination rate. Begun DJ, editor. PLoS Genet. Public Library of
Science; 2010;6: e1000886. doi:10.1371/journal.pgen.1000886
58. Willing EM, Dreyer C, van Oosterhout C. Estimates of genetic differentiation
measured by fst do not necessarily require large sample sizes when using many
snp markers. PLoS One. 2012;7: 1–7. doi:10.1371/journal.pone.0042649
59. Messer PW, Petrov DA. Frequent adaptation and the McDonald-Kreitman test.
Proc Natl Acad Sci U S A. 2013;110: 8615–20. doi:10.1073/pnas.1220835110
60. Ewing GB, Jensen JD. The consequences of not accounting for background
selection in demographic inference. Mol Ecol. Wiley-Blackwell; 2016;25: 135–141.
doi:10.1111/mec.13390
61. Gravel S, Henn BM, Gutenkunst RN, Indap AR, Marth GT, Clark AG, et al.
Demographic history and rare allele sharing among human populations. Proc Natl
Acad Sci. 2011;108: 11983–11988. doi:10.1073/pnas.1019276108
62. Tennessen JA, Bigham AW, O'Connor TD, Fu W, Kenny EE, Gravel S, et al.
Evolution and Functional Impact of Rare Coding Variation from Deep Sequencing
of Human Exomes. Science (80-). 2012;337: 64–69.
doi:10.1126/science.1219240
63. Phung TN, Huber CD, Lohmueller KE. Determining the Effect of Natural Selection
on Linked Neutral Divergence across Species. Akey JM, editor. PLoS Genet. W.
H. Freeman; 2016;12: e1006199. doi:10.1371/journal.pgen.1006199
64. Burgess R, Yang Z. Estimation of hominoid ancestral population sizes under
Bayesian coalescent models incorporating mutation rate variation and sequencing
errors. Mol Biol Evol. Oxford University Press; 2008;25: 1979–1994.
doi:10.1093/molbev/msn148
65. Mouse Genome Sequencing Consortium MGS, Waterston RH, Lindblad-Toh K,
Birney E, Rogers J, Abril JF, et al. Initial sequencing and comparative analysis of
the mouse genome. Nature. 2002;420: 520–62. doi:10.1038/nature01262
66. Charlesworth B, Nordborg M, Charlesworth D. The effects of local selection,
balanced polymorphism and background selection on equilibrium patterns of
genetic diversity in subdivided populations. Genet Res. 1997;70: 155–174.
doi:10.1017/S0016672397002954
67. Hu XS, He F. Background selection and population differentiation. J Theor Biol.
2005;235: 207–219. doi:10.1016/j.jtbi.2005.01.004
68. Coop G, Pickrell JK, Novembre J, Kudaravalli S, Li J, Absher D, et al. The role of
geography in human adaptation. Schierup MH, editor. PLoS Genet. R Foundation

for Statistical Computing; 2009;5: e1000500. doi:10.1371/journal.pgen.1000500

69.   Bhatia G, Patterson N, Pasaniuc B, Zaitlen N, Genovese G, Pollack S. Genome-
Wide Comparison of African-Ancestry Populations from CARe and Other Cohorts
Reveals Signals of Natural Selection. 2011;89: 368–381.
doi:10.1016/j.ajhg.2011.07.025.

70.   Eyre-Walker A, Keightley PD. Estimating the rate of adaptive molecular evolution
in the presence of slightly deleterious mutations and population size change. Mol
Biol Evol. 2009;26: 2097–2108. doi:10.1093/molbev/msp119

71.   Galtier N. Adaptive Protein Evolution in Animals and the Effective Population Size
Hypothesis. Schierup MH, editor. PLoS Genet. Oxford University Press; 2016;12:
e1005774. doi:10.1371/journal.pgen.1005774

72.   Garud NR, Messer PW, Buzbas EO, Petrov DA. Recent selective sweeps in North
American Drosophila melanogaster show signatures of soft sweeps. Copenhaver
GP, editor. PLoS Genet. Wiley; 2015;11: e1005004.
doi:10.1371/journal.pgen.1005004

73.   Szpiech ZA, Hernandez RD. Selscan: An efficient multithreaded program to
perform EHH-based scans for positive selection. Mol Biol Evol. 2014;31: 2824–
2827. doi:10.1093/molbev/msu211

74.   Fu W, Gittelman RM, Bamshad MJ, Akey JM. Characteristics of neutral and
deleterious protein-coding variation among individuals and populations. Am J
Hum Genet. Springer-Verlag, New York; 2014;95: 421–436.
doi:10.1016/j.ajhg.2014.09.006

75.   Henn BM, Botigué LR, Peischl S, Dupanloup I, Lipatov M, Maples BK, et al.
Distance from sub-Saharan Africa predicts mutational load in diverse human
genomes. Proc Natl Acad Sci. National Academy of Sciences; 2015;113:
201510805. doi:10.1073/pnas.1510805112

76.   Casals F, Hodgkinson A, Hussin J, Idaghdour Y, Bruat V, de Maillard T, et al.
Whole-Exome Sequencing Reveals a Rapid Change in the Frequency of Rare
Functional Variants in a Founding Population of Humans. Williams SM, editor.
PLoS Genet. Public Library of Science; 2013;9: e1003815.
doi:10.1371/journal.pgen.1003815

77.   Lim ET, Würtz P, Havulinna AS, Palta P, Tukiainen T, Rehnström K, et al.
Distribution and Medical Impact of Loss-of-Function Variants in the Finnish
Founder Population. Cutler D, editor. PLoS Genet. Public Library of Science;
2014;10: e1004494. doi:10.1371/journal.pgen.1004494

78.   Cao J, Schneeberger K, Ossowski S, Günther T, Bender S, Fitz J, et al. Whole-
genome sequencing of multiple Arabidopsis thaliana populations. Nat Genet.
2011;43: 956–963. doi:10.1038/ng.911

79.   Renaut S, Rieseberg LH. The accumulation of deleterious mutations as a
consequence of domestication and improvement in sunflowers and other
compositae crops. Mol Biol Evol. Oxford University Press; 2015;32: 2273–2283.
doi:10.1093/molbev/msv106

80.   Balick DJ, Do R, Cassa CA, Reich D, Sunyaev SR. Dominance of Deleterious
Alleles Controls the Response to a Population Bottleneck. Coop G, editor. PLoS
Genet. Cambridge University Press; 2015;11: e1005436.
doi:10.1371/journal.pgen.1005436

81. Simons YB, Sella G. The impact of recent population history on the deleterious
mutation load in humans and close evolutionary relatives [Internet]. Current
Opinion in Genetics and Development. 2016. pp. 150–158.
doi:10.1016/j.gde.2016.09.006
82. Gravel S. When is selection effective? Genetics. 2016;203: 451–462.
doi:10.1534/genetics.115.184630
83. Charlesworth B. The Effects of Deleterious Mutations on Evolution at Linked
Sites. Genetics. 2012;190: 5–22. doi:10.1534/genetics.111.134288
84. Coventry A, Bull-Otterson LM, Liu X, Clark AG, Maxwell TJ, Crosby J, et al. Deep
resequencing reveals excess rare recent variants consistent with explosive
population growth. Nat Commun. Nature Publishing Group; 2010;1: 131.
doi:10.1038/ncomms1130
85. Keinan A, Clark AG. Recent explosive human population growth has resulted in
an excess of rare genetic variants. Science (80-). 2012;336: 740–3.
doi:10.1126/science.1217283
86. Nelson MR, Wegmann D, Ehm MG, Kessner D, St. Jean P, Verzilli C, et al. An
Abundance of Rare Functional Variants in 202 Drug Target Genes Sequenced in
14,002 People. Science (80-). 2012;337: 100–104. doi:10.1126/science.1217876
87. Gazave E, Ma L, Chang D, Coventry A, Gao F, Muzny D, et al. Neutral genomic
regions refine models of recent rapid human population growth. Proc Natl Acad
Sci U S A. 2014;111: 757–62. doi:10.1073/pnas.1310398110
88. Maher MC, Uricchio LH, Torgerson DG, Hernandez RD. Population Genetics of
Rare Variants and Complex Diseases. Hum Hered. NIH Public Access; 2012;74:
118–128. doi:10.1159/000346826
89. Fu W, O'Connor TD, Jun G, Kang HM, Abecasis G, Leal SM, et al. Analysis of
6,515 exomes reveals the recent origin of most human protein-coding variants.
Nature. NIH Public Access; 2013;493: 216–220. doi:10.1038/nature11690
90. Fay JC, Wu CI. A human population bottleneck can account for the discordance
between patterns of mitochondrial versus nuclear DNA variation. Mol Biol Evol.
1999;16: 1003–1005. doi:10.1093/oxfordjournals.molbev.a026175
91. Wall JD, Andolfatto P, Przeworski M. Testing Models of Selection and
Demography in Drosophila simulans. Genetics. 2002;162. Available:
http://www.genetics.org/content/162/1/203
92. Pool JE, Nielsen R. Population size changes reshape genomic patterns of
diversity. Evolution (N Y). 2007;61: 3001–3006. doi:10.1111/j.1558-
5646.2007.00238.x
93. Gottipati S, Arbiza L, Siepel A, Clark AG, Keinan A. Analyses of X-linked and
autosomal genetic variation in population-scale whole genome sequencing. Nat
Genet. NIH Public Access; 2011;43: 741–743. doi:10.1038/ng.877
94. Arbiza L, Gottipati S, Siepel A, Keinan A. Contrasting X-linked and autosomal
diversity across 14 human populations. Am J Hum Genet. 2014;94: 827–844.
doi:10.1016/j.ajhg.2014.04.011
95. Wilson Sayres MA, Lohmueller KE, Nielsen R. Natural Selection Reduced
Diversity on Human Y Chromosomes. PLoS Genet. 2014;10.
doi:10.1371/journal.pgen.1004064
96. Hammer MF, Woerner AE, Mendez FL, Watkins JC, Cox MP, Wall JD. The ratio

of human X chromosome to autosome diversity is positively correlated with
genetic distance from genes. Nat Genet. 2010;42: 830–1. doi:10.1038/ng.651

97. Leffler EM, Bullaughey K, Matute DR, Meyer WK, Ségurel L, Venkat A, et al.
Revisiting an Old Riddle: What Determines Genetic Diversity Levels within
Species? PLoS Biol. Public Library of Science; 2012;10: e1001388.
doi:10.1371/journal.pbio.1001388

98. Vucetich JA, Waite TA, Nunney L. Fluctuating Population Size and the Ratio of
Effective to Census Population Size Published by : Society for the Study of
Evolution Stable URL : http://www.jstor.org/stable/2411022. Source Evol. 2008;51:
2017–2021. doi:10.2307/2411022

99. Coop G. Does linked selection explain the narrow range of genetic diversity
across species? [Internet]. bioRxiv. 2016. doi:10.1101/042598

100. Lewontin RC. The Genetic Basis of Evolutionary Change [Internet]. The Genetic
Basis of Evolutionary Change. New York and London: Columbia University Press;
1974. doi:10.1136/ard.2010.140574

101. Nam K, Munch K, Mailund T, Nater A, Greminger MP, Krützen M, et al. Evidence
that the rate of strong selective sweeps increases with population size in the great
apes. Proc Natl Acad Sci. National Academy of Sciences; 2016;114: 1–6.
doi:10.1073/pnas.1605660114

102. Prado-Martinez J, Sudmant PH, Kidd JM, Li H, Kelley JL, Lorente-Galdos B, et al.
Great ape genetic diversity and population history. Nature. Nature Research;
2013;499: 471–475. doi:10.1038/nature12228

103. McManus KF, Kelley JL, Song S, Veeramah KR, Woerner AE, Stevison LS, et al.
Inference of gorilla demographic and selective history from whole-genome
sequence data. Mol Biol Evol. Cambridge University Press, Cambridge; 2015;32:
600–612. doi:10.1093/molbev/msu394

104. Nater A, Greminger MP, Arora N, Van Schaik CP, Goossens B, Singleton I, et al.
Reconstructing the demographic history of orang-utans using Approximate
Bayesian Computation. Mol Ecol. 2015;24: 310–327. doi:10.1111/mec.13027

105. Kuhlwilm M, de Manuel M, Nater A, Greminger MP, Kr?tzen M, Marques-Bonet T.
Evolution and demography of the great apes. Curr Opin Genet Dev. 2016;41:
124–129. doi:10.1016/j.gde.2016.09.005

106. Uricchio LH, Torres R, Witte JS, Hernandez RD. Population genetic simulations of
complex phenotypes with implications for rare variant association tests. Genet
Epidemiol. 2015;39: 35–44. doi:10.1002/gepi.21866

107. Uricchio LH, Zaitlen NA, Ye CJ, Witte JS, Hernandez RD. Selection and explosive
growth alter genetic architecture and hamper the detection of causal rare variants.
Genome Res. Cold Spring Harbor Laboratory Press; 2016;26: 863–873.
doi:10.1101/gr.202440.115

108. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The
variant call format and VCFtools. Bioinformatics. 2011;27: 2156–2158.
doi:10.1093/bioinformatics/btr330

109. Kidd J, Sharpton T, Bobo D, Norman P, Martin A, Carpenter M, et al. Exome
capture from saliva produces high quality genomic and metagenomic data. BMC
Genomics. BioMed Central; 2014;15: 262. doi:10.1186/1471-2164-15-262

110. Kim HL, Ratan A, Perry GH, Montenegro A, Miller W, Schuster SC. Khoisan

hunter-gatherers have been the largest population throughout most of modern-
human demographic history. Nat Commun. 2014;5: 5692.
doi:10.1038/ncomms6692

111. Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral
substitution rates on mammalian phylogenies. Genome Res. Cold Spring Harbor
Laboratory Press; 2010;20: 110–121. doi:10.1101/gr.097857.109

112. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, et al.
Evolutionarily conserved elements in vertebrate, insect, worm, and yeast
genomes. Genome Res. 2005;15: 1034–1050. doi:10.1101/gr.3715005

113. Consortium ENCODEP, Bernstein BE, Birney E, Dunham I, Green ED, Gunter C,
et al. An integrated encyclopedia of DNA elements in the human genome. Nature.
2012;489: 57–74. doi:10.1038/nature11247

114. Bailey JA, Yavor AM, Massa HF, Trask BJ, Eichler EE. Segmental duplications:
organization and impact within the current human genome project assembly.
Genome Res. 2001;11: 1005–17. doi:10.1101/gr.187101

115. Capra JA, Hubisz MJ, Kostka D, Pollard KS, Siepel A. A Model-Based Analysis of
GC-Biased Gene Conversion in the Human and Chimpanzee Genomes. Coop G,
editor. PLoS Genet. Public Library of Science; 2013;9: e1003684.
doi:10.1371/journal.pgen.1003684

116. Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, Gibbs RA, et al. A second
generation human haplotype map of over 3.1 million SNPs. Nature. 2007;449:
851–61. doi:10.1038/nature06258

117. Pratto F, Brick K, Khil P, Smagulova F, Petukhova G V, Camerini-Otero RD. DNA
recombination. Recombination initiation maps of individual human genomes.
Science. 2014;346: 1256442. doi:10.1126/science.1256442

118. Wang K, Li M, Hakonarson H. ANNOVAR: Functional annotation of genetic
variants from high-throughput sequencing data. Nucleic Acids Res. Oxford
University Press; 2010;38: e164–e164. doi:10.1093/nar/gkq603

119. Hernandez RD, Williamson SH, Zhu L, Bustamante CD. Context-dependent
mutation rates may cause spurious signatures of a fixation bias favoring higher
GC-content in humans. Mol Biol Evol. 2007;24: 2196–2202.
doi:10.1093/molbev/msm149

120. Schrider DR, Shanku AG, Kern AD. Effects of linked selective sweeps on
demographic inference and model selection. Genetics. 2016;204: 1207–1223.
doi:10.1534/genetics.116.190223

121. Palamara PF, Francioli LC, Wilton PR, Genovese G, Gusev A, Finucane HK, et
al. Leveraging Distant Relatedness to Quantify Human Mutation and Gene-
Conversion Rates. Am J Hum Genet. 2015;97: 775–789.
doi:10.1016/j.ajhg.2015.10.006

122. Hernandez RD. A flexible forward simulator for populations subject to selection
and demography. Bioinformatics. 2008;24: 2786–2787.
doi:10.1093/bioinformatics/btn522

123. Torgerson DG, Boyko AR, Hernandez RD, Indap A, Hu X, White TJ, et al.
Evolutionary processes acting on candidate cis-regulatory regions in humans
inferred from patterns of polymorphism and divergence. PLoS Genet. 2009;5.
doi:10.1371/journal.pgen.1000592

124. Maples BK, Gravel S, Kenny EE, Bustamante CD. RFMix: A discriminative
modeling approach for rapid and robust local-ancestry inference. Am J Hum
Genet. The American Society of Human Genetics; 2013;93: 278–288.
doi:10.1016/j.ajhg.2013.06.020
125. O'Connell J, Gurdasani D, Delaneau O, Pirastu N, Ulivi S, Cocca M, et al. A
General Approach for Haplotype Phasing across the Full Spectrum of
Relatedness. PLoS Genet. Public Library of Science; 2014;10: e1004234.
doi:10.1371/journal.pgen.1004234
126. Mao X, Bigham AW, Mei R, Gutierrez G, Weiss KM, Brutsaert TD, et al. A
genomewide admixture mapping panel for Hispanic/Latino populations. Am J
Hum Genet. 2007;80: 1171–1178. doi:10.1086/518564
127. Hudson RR, Slatkin M, Maddison WP. Estimation of levels of gene flow from DNA
sequence data. Genetics. 1992;589: 583–589. Available:
http://www.genetics.org/content/132/2/583.short

1393
1394

Supporting information

S1 Supporting information.

Fig S1. Inference models inferred from TGP CG weak BGS neutral regions and coding four-fold degenerate sites. Solid lines are the inference results from running dadi on 53 YRI (African), 64 CEU (European), and 62 CHS (East Asian) TGP CG samples (projected down to 106 chromosomes during inference procedure) across neutral regions in the weakest 1% BGS bin ($B \geq 0.994$). Broken lines represent the inference results using the same CG samples, but with sequence data only from coding four-fold degenerate synonymous sites.

Fig S2. Diversity for TGP non-admixed populations while controlling for GC-biased gene conversion and recombination hotspots. (A) Normalized diversity (π/divergence) measured across the strongest 1% BGS bin. (B) Normalized diversity measured across the weakest 1% BGS bin. (C) Relative diversity: the ratio of normalized diversity for the strongest 1% BGS bin to normalized diversity for the weakest 1%

BGS bin (π/π_0). Error bars represent ±1 SEM calculated from 1,000 bootstrapped da-

tasets.

Fig S3. Diversity for TGP non-admixed populations without normalizing by diver-

gence with Rhesus macaque. (A) Diversity (π) measured across the strongest 1%

BGS bin. (B) Diversity measured across the weakest 1% BGS bin. (C) Relative diversi-

ty: the ratio of diversity for the strongest 1% BGS bin to diversity for the weakest 1%

BGS bin (π/π_0). Error bars represent ±1 SEM calculated from 1,000 bootstrapped da-

tasets.

Fig S4. Diversity for TGP continental groups while controlling for GC-biased gene

conversion and recombination hotspots. (A) Normalized diversity (π/divergence)

measured across the strongest 1%, 5%, 10% and 25% BGS bins and the weakest 1%

BGS bin (as classified by B). (B) Relative diversity (π/π_0) for the strongest 1%, 5%, 10%,

and 25% BGS bins. Error bars represent ±1 SEM calculated from 1,000 bootstrapped

datasets.

Fig S5. Diversity for TGP continental groups without normalizing by divergence

with Rhesus macaque. (A) Diversity (π) measured across the strongest 1%, 5%, 10%

and 25% BGS bins and the weakest 1% BGS bin (as classified by B). (B) Relative di-

versity (π/π_0) for the strongest 1%, 5%, 10%, and 25% BGS bins. Error bars represent

±1 SEM calculated from 1,000 bootstrapped datasets.

Fig S6. F_{ST} measured across joint bins of B and recombination rate for different

TGP continental groups. The left panels of Figures S6 A-E show F_{ST} measured as a

function of 25 4% quantile recombination rate bins conditional on three 2% quantile B

bins (note log scale of x-axis for recombination rate). The right panels of Figures S6 A-E

show F_{ST} measured as a function of 25 4% quantile B bins conditional on three 2%

quantile recombination rate bins. The following continental group comparisons are

shown for each plot: (A) African vs. European, (B) African vs. East Asian, (C) European

vs. South Asian, (D) European vs. East Asian, (E) South Asian vs. East Asian. Smaller

transparent points and lines show the F_{ST} estimates and corresponding lines of best fit

(using linear regression) for each of the pairwise population comparisons within a par-

ticular pair of continental groups (25 comparisons total). Larger opaque points are mean

F_{ST} estimates across all pairwise comparisons within a particular pair of continental

groups (with bold lines showing their corresponding lines of best fit).

Fig S7. Simulations of diversity and relative diversity under BGS using a human

demographic model without migration. (A) Inferred demographic model from Com-

plete Genomics TGP data. The demographic model used for the simulations in Figure

S7 are identical to those used for Figure 6, except that migration parameters between

all populations are set to 0. (B) Simulated diversity at neutral sites across populations as

a function of time under our inferred demographic model without BGS (π_0 - dashed col-

ored lines) and with BGS (π - solid colored lines). (C) Relative diversity (π/π_0) measured

by taking the ratio of diversity with BGS (π) to diversity without BGS (π_0) at each time

point. Note that the x-axes in all three figures are on the same scale. Time is scaled us-

ing a human generation time of 25 years per generation. Simulation data was sampled

every 100 generations.

Fig S8. Simulations of diversity and relative diversity under BGS using various

values of U. Values of U are provided in the header for each set of plots. Left column

plots show results of simulations under a demographic model with migration between all

human populations. Right column plots show results of simulations under a demograph-

ic model with no migration. Colored lines represent different populations though time

and are identical to those in Figure 6 and Figure S7. The demographic model used is

also identical to that in Figure 6 (for simulations with migration) and Figure S7 (for simu-

lations without migration). Simulation data was sampled every 100 generations.

1460

1461   **S2 Appendix. Soft sweep detection and implementation in selscan v1.2.0.**

1462

Main Figure Legends

Fig 1. Normalized diversity and relative diversity for non-admixed populations of

the Thousand Genomes Project (TGP).

(A) Normalized diversity (π/divergence) measured across the strongest 1% background

selection (BGS) bin. (B) Normalized diversity measured across the weakest 1% BGS

bin. (C) Relative diversity: the ratio of normalized diversity for the strongest 1% BGS bin

to normalized diversity for the weakest 1% BGS bin (π/π_0). BGS bins were classified by

B. TGP population labels are indicated below each bar (see Table S11 in Supporting

information for population label descriptions), with African populations colored by gold

shades, European populations colored by blue shades, South Asian populations colored

by violet shades, and East Asian populations colored by green shades. Error bars rep-

resent ±1 SEM calculated from 1,000 bootstrapped datasets.

Fig 2. Normalized and relative diversity for Thousand Genomes Project (TGP)

continental groups.

(A) Normalized diversity (π/divergence) measured across the strongest 1%, 5%, 10%

and 25% background selection (BGS) bins and the weakest 1% BGS bin (as classified

by *B*). (B) Relative diversity: the ratio of normalized diversity for each strong BGS bin in

(A) to normalized diversity for the weakest 1% BGS bin (π/π_0). Error bars represent ±1

SEM calculated from 1,000 bootstrapped datasets.

Fig 3. Comparing patterns of diversity between local ancestry segments of ad-

mixed samples and continental groups.

(A) Normalized diversity (heterozygosity/divergence) and (B) Relative diversity: the ratio

of normalized diversity for each strong BGS bin in (A) to normalized diversity for the

weakest 1% BGS bin. Local ancestry segments include African, European, and Native

American ancestries. Continental groups include African, European, and East Asian

populations. Error bars represent ±1 SEM calculated from 1,000 bootstrapped datasets.

Fig 4. F_{ST} is correlated with *B* but not recombination rate.

(A) F_{ST} measured across 2% quantile bins of *B*. (B) F_{ST} measured across 2% quantile

recombination rate bins. The right panel of Figure 4B displays a narrower range of re-

combination rates to show detail. Smaller transparent points and lines show the esti-

mates and corresponding lines of best fit (using linear regression) for F_{ST} between every

pairwise population comparison for a particular pair of continental groups (25 pairwise

comparisons each). Larger opaque points and lines are mean F_{ST} estimates and lines of

best fit across all Thousand Genomes Project (TGP) population comparisons between a

particular pair of continental groups. Error bars represent ±1 SEM calculated from 1,000

bootstrapped datasets.

Fig 5. F_{ST} between African (AFR) and South Asian (SASN) populations jointly

across B and recombination rate.

(A) F_{ST} as a function of 25 recombination rate bins (4% quantile bins) conditional on

three different 2% quantile B bins (note log scale of x-axis for recombination rate). (B)

F_{ST} as a function of 25 B bins (4% quantile bins) conditional on three different 2% quan-

tile recombination rate bins. Smaller transparent points and lines show the F_{ST} esti-

mates and corresponding lines of best fit (using linear regression) for each of the pair-

wise comparisons of AFR vs. SASN Thousand Genomes Project (TGP) populations (25

comparisons total). Larger opaque points are mean F_{ST} estimates across all pairwise

comparisons of AFR vs. SASN TGP populations (with bold lines showing their corre-

sponding lines of best fit).

Fig 6. Simulations confirm that demographic events shape the impact of back-

ground selection (BGS).

(A) Inferred demographic model from Complete Genomics Thousand Genomes Project

(TGP) data showing population size changes for Africans (AFR), Europeans (EUR), and

East Asians (EASN) as a function of time that was used for the simulations of BGS. (B)

Simulated diversity at neutral sites across populations as a function of time under our

inferred demographic model without BGS (π_0 - dashed colored lines) and with BGS (π -

solid colored lines). (C) Relative diversity (π/π_0) measured by taking the ratio of diversity

with BGS (π) to diversity without BGS (π_0) at each time point. Note that the x-axes in all

three figures are on the same scale. Time is scaled using a human generation time of

25 years per generation. Simulation data was sampled every 100 generations.

1521

Figure 1

Figure 2

Figure 3

Figure 4

A

B

Figure 5

Figure 6

S1 Supporting information

Table S1

Parameters	$B \geq 0.994$	four-fold degenerate
$N_{Ancestral}$	18,449	17,118
N_{AFR}	38,874	47,537
N_{Bott}	5,946	6,408
N_{EUR0}	3,413	4,331
N_{EUR}	81,901	100,614
N_{EASN0}	1,317	1,678
N_{EASN}	206,804	266,616
$T_{AFR}+T_{Bott}+T_{EUR\_EASN}$ (kya)	552,939	413,337
$T_{Bott}+T_{EUR\_EASN}$ (kya)	149,813	198,603
T_{EUR_EASN} (kya)	48,822	69,584
r_{EUR} (%)	0.163	0.113
r_{EASN} (%)	0.259	0.182
$m_{AFR\text{-}Bott}$ (x10^{-5})	7.83	7.02
$m_{AFR\text{-}EUR}$ (x10^{-5})	0.51	0.47
$m_{AFR\text{-}EASN}$ (x10^{-5})	0.13	0.18
$m_{EUR\text{-}EASN}$ (x10^{-5})	0.98	1.14

Table S1. Inferred parameters from running dadi on TGP CG data across neutral regions in the weakest 1% BGS bin ($B \geq 0.994$) and across four-fold degenerate sites. The demographic model inferred is the Out-of-Africa demographic model of Gutenkunst et al. 2009. Time parameters, T, assume a generation time of 25 years per generation. Growth rates, r, and migration rates, m, are per generation. Parameters with subscript, "*Bott*", represent parameters inferred for the ancestral European and East Asian out-of-Africa bottleneck population. Time parameters with subscript "*EUR_EASN*" represent the European-East Asian population split.

Table S3

Henn et al. 2015 samples		
SampleID	**Number of Sites**	**Mean Depth**
HGDP00991	2,207,845	6.96118
HGDP00987	2,229,426	7.19132
HGDP01036	2,373,023	11.6072
HGDP00992	2,452,509	12.1913
HGDP01029	2,415,792	12.3526
HGDP01032	2,407,400	12.8113
Kidd et al. 2014 samples		
SampleID	**Number of Sites**	**Mean Depth**
SA1000A	547,527	2.56481
SA1025A	2,136,905	9.1239
Kim et al. 2014 samples		
SampleID	**Number of Sites**	**Mean Depth**
KB2	2,756,225	27.5951
NB1	2,599,220	28.0148
MD8	2,777,871	38.4532
NB8	2,778,198	40.1789
KB1	2,757,336	50.5629

Table S3. Number of polymorphic sites and mean depth coverage of 13 Khoisan samples used for SNP ascertainment in calculations of F_{ST}.

Table S4

	top 1% *B*	top 5% *B*	top 10% *B*	top 25% *B*	bottom 1% *B*
filters	7.59	40.42	87.86	246.59	13.1
filters + gBGC and hotspots removal	7.26	38.68	83.75	231.71	7.94

Table S4. Total number of Mb in the human genome passing the set of 13 filters described in Materials and Methods that were used for calculating pairwise genetic diversity (π) for each quantile of *B*. The bottom row is the total number Mb when including the set of filters to remove regions sensitive to GC-biased gene conversion (gBGC) or sites in recombination hotspots. Additionally, these totals only include those 100 kb regions that had a minimum of 10 kb of divergence information for Rhesus macaque (see Materials and Methods).

Table S5

Ancestry	top 1% *B*	top 5% *B*	top 10% *B*	top 25% *B*	bottom 1% *B*
African	841.97	4471.54	9720.15	27333.95	1447.04
European	815.74	4296.69	9293.04	26034.57	1366.26
Native American	497.29	2603.12	5640.13	15776.71	834.46

Table S5. Total number of Mb of homozygous ancestry that passed all filters and were used in the analyses of admixed samples in the 6 admixed TGP populations (ACB, ASW, CLM, MXL, PEL, PUR) for each quantile of *B*. Additionally, these totals only include those 100 kb regions that had a minimum of 10 kb of divergence information for Rhesus macaque (see Materials and Methods).

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

Figure S7

S2 Appendix. Soft sweep detection and implementation in selscan v1.2.0.

Detecting soft sweeps

Under the model of a soft sweep, there is an increased chance of multiple distinct haplotypes sweeping to high frequency in a population. Garud et al. [1] developed a window-based statistic ($H12$) with good power to detect this process, and here we adapt $H12$ into an integrated haplotype homozygosity framework [2–4]. We call this new statistic $iHH12$. The general principle of these statistics is to combine the top two most frequent haplotypes into a single haplotype class to avoid the reduced power that iHS has when the adaptive allele segregates on more than one haplotype background. We calculate $iHH12$ as follows.

Following the notation of Szpiech and Hernandez [5] in a sample of n chromosomes we let $\mathcal{C}$ be the set of all possible distinct haplotypes at the locus x_0. $\mathcal{C}(x_i)$ is then the set of all possible distinct haplotypes extending from locus x_0 to locus x_i. Let h_i in $\mathcal{C}(x)$ be the i^{th} most frequent haplotype. We then calculate $EHH12$ of the entire sample of haplotypes from x_0 to x_i as

$$EHH12(x_i) = \frac{\binom{n_{h_1}+n_{h_2}}{2}}{\binom{n}{2}} + \sum_{j>2}^{|\mathcal{C}(x_i)|} \frac{\binom{n_{h_j}}{2}}{\binom{n}{2}}$$

where n_{h_j} is the number of h_j haplotypes in the sample.

If $EHH12(x_i)$ is calculated repeatedly for several x_i moving farther away from x_0, we expect to observe more haplotypes and therefore we expect to observe lower haplotype homozygosity. However, the decay of homozygosity is slower in a region under selection [2–4]. Therefore, we integrate the decay of $EHH12$ as a function of genetic distance in order to summarize the pattern and make genome-wide comparisons. This integrated score is calculated as

$$iHH12 = \sum_{i=1}^{|\mathcal{D}|} \frac{1}{2}(EHH12(x_{i-1}) - EHH(x_i))g(x_{i-1}, x_i)$$
$$+ \sum_{i=1}^{|\mathcal{U}|} \frac{1}{2}(EHH12(x_{i-1}) - EHH(x_i))g(x_{i-1}, x_i)$$

where $g(x_{i-1}, x_i)$ is the genetic distance between markers x_{i-1} and x_i. $\mathcal{D}$ and $\mathcal{U}$ represent sets of markers downstream and upstream from x_0, respectively. In practice the curve is integrated until $EHH12 < 0.05$ on both sides of the focal locus. Finally, $iHH12$ is normalized genome-wide in order to account for the effects of demographic history on the distribution of haplotype homozygosity. We integrated this new statistical framework to detect soft-sweeps into `selscan` version 1.2.0 (https://github.com/szpiech/selscan) [5].

We evaluated the power of our $iHH12$ statistic implementation in `selscan` to detect hard and soft sweeps relative to iHS across a range of parameters. We simulated neutrally evolving sequences with `ms` [6] and non-neutrally evolving sequences with `mssel`, a modified version of `ms` also developed by R. Hudson

that conditions on an allele frequency trajectory. We simulated trajectories backwards in time under a selection on standing variation model with $s = 0.01$. Once an adaptive variant reached a set frequency backwards in time, the selection coefficient was set to $s = 0$ and was allowed drift neutrally until loss. We simulated 200 replicates across several sampling frequencies (0.7, 0.8, 0.9), several frequencies at which the variant become adaptive (0, 0.01, 0.02, 0.05, 0.10), and several demographic histories (Constant, African, European; [7]).

For both iHS and $iHH12$ scans, we normalized scores with respect to the neutral simulations and calculated the critical threshold for the most extreme 1% of scores. Using non-overlapping 100 kb windows across the genome, we calculated the fraction of scores in each window above this threshold. The top 1% of windows are identified as putatively under positive selection. This scheme controls the false positive rate to be no greater than 1%.

$iHH12$ has good power to detect hard and soft sweeps (Appendix Fig A1 A, C, E) and has improved power to identify both types of sweeps over iHS (Appendix Fig A1 B, D, F), particularly under realistic models of human demography.

Computing iHS and iHH12 scores in the Thousand Genomes Project (TGP)

We used `selscan` to compute both iHS and $iHH12$ scores for phase 3 TGP [8] phased whole genome sequences with a genetic map from HapMap3 [9]. Genetic map locations for sites not present in HapMap3 were linearly interpolated. The statistics were calculated for each population separately, and variants of frequency < 0.05 were filtered by `selscan`. All `selscan` runs used default parameters.

Using `selscan`'s companion program `norm`, for each population we normalized $iHH12$ scores genome-wide and normalized iHS scores in 1% frequency bins genome-wide. We identified the critical threshold representing the most extreme 1% of scores for each statistic. Then, to identify putative regions under selection, we partitioned the genome into non-overlapping 100 kb windows, and then we calculated the fraction of scores in each window above this threshold. The top 1% of windows were identified as putatively under positive selection. This scheme controlled the false positive rate to be no greater than 1%.

Appendix Figure A1

Appendix Fig A1. Power of $iHH12$ and comparison with iHS. Simulated power of $iHH12$ (A), (C), and (E) under varying parameters and comparison with iHS power (B), (D), and (F) in the same scenario. Panels (A) and (B) show results for a constant demography; panels (C) and (D) show results for an African demography; and panels (E) and (F) show results for a European demography. Non-constant demographies are from Gutenkunst et al. [7]. When the frequency at which selection begins is > 0, the sweep is considered soft. All simulations assume a selection coefficient of $s = 0.01$.

References

1. Garud NR, Messer PW, Buzbas EO, Petrov DA. Recent selective sweeps in North American Drosophila melanogaster show signatures of soft sweeps. Copenhaver GP, editor. PLoS Genet. Wiley; 2015;11: e1005004. doi:10.1371/journal.pgen.1005004

2. Sabeti PC, Reich DE, Higgins JM, Levine HZP, Richter DJ, Schaffner SF, et al. Detecting recent positive selection in the human genome from haplotype structure. Nature. Nature Publishing Group; 2002;419: 832–837. doi:10.1038/nature01027.1.

3. Voight BF, Kudaravalli S, Wen X, Pritchard JK. A map of recent positive selection in the human genome. Hurst L, editor. PLoS Biol. Public Library of Science; 2006;4: 0446–0458. doi:10.1371/journal.pbio.0040072

4. Sabeti PC, Varilly P, Fry B, Lohmueller J, Hostetter E, Cotsapas C, et al. Genome-wide detection and characterization of positive selection in human populations. Nature. Nature Publishing Group; 2007;449: 913–8. doi:10.1038/nature06250

5. Szpiech ZA, Hernandez RD. Selscan: An efficient multithreaded program to perform EHH-based scans for positive selection. Mol Biol Evol. 2014;31: 2824–2827. doi:10.1093/molbev/msu211

6. Hudson RR. Generating samples under a Wright-Fisher neutral model of genetic variation. Bioinformatics. Oxford University Press; 2002;18: 337–338. doi:10.1093/bioinformatics/18.2.337

7. Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Inferring the joint demographic history of multiple populations from multidimensional SNP frequency data. PLoS Genet. 2009;5: e1000695. doi:10.1371/journal.pgen.1000695

8. Auton A, Abecasis GR, Altshuler DM, Durbin RM, Abecasis GR, Bentley DR, et al. A global reference for human genetic variation. Nature. 2015;526: 68–74. doi:10.1038/nature15393

9. The International HapMap 3 Consortium, Altshuler DM, Gibbs RA, Peltonen L, Dermitzakis E, Schaffner SF, et al. Integrating common and rare genetic variation in diverse human populations. Nature. 2010;467: 52–8. doi:10.1038/nature09298