Local and global chromatin interactions are altered by large genomic deletions

associated with human brain development

Ying Zhang[1*], Xianglong Zhang[2,3*], Xiaowei Zhu[2,3*], Carolin Purmann[2,3], Michael S Haney[3],

Thomas Ward[2,3], Jie Yao[4], Sherman M Weissman[5#], Alexander E Urban[2,3#]

1 Department of Genetics and Genomic Sciences, Icahn School of medicine at Mount Sinai, 2

Department of Psychiatry and Behavioral Sciences, 3 Department of Genetics, Stanford

University School of Medicine, 4 Department of Cell Biology, Yale University School of Medicine,

5 Department of Genetics, Yale University

Email addresses:

Ying Zhang, zzhang006@gmail.com;

Xianglong Zhang, xzhang8@stanford.edu;

Xiaowei Zhu, xwzhu@stanford.edu;

Carolin Purmann, purmann@stanford.edu;

Michael S Haney, haney@stanford.edu;

Thomas Ward, wardtr@stanford.edu;

Jie Yao, yao.j@yale.edu;

Sherman M Weissman, sherman.weissman@yale.edu;

Alexander E Urban, aeurban@stanford.edu.

*These authors contributed equally to this work.

#To whom correspondence should be addressed.

Abstract

Background: Large copy number variants (CNVs) in the human genome are strongly

associated with common neurodevelopmental, neuropsychiatric disorders such as

schizophrenia and autism. Using Hi-C analysis of long-range chromosome interactions and

ChIP-Seq analysis of regulatory histone marks we studied the epigenomic effects of the

prominent large deletion CNV on chromosome 22q11.2 and also replicated a subset of the

findings for the large deletion CNV on chromosome 1q21.1.

Results: We found that, in addition to local and global gene expression changes, there are

pronounced and multilayered effects on chromatin states, chromosome folding and

topological domains of the chromatin, that emanate from the large CNV locus. Regulatory

histone marks are altered in the deletion proximal regions, and in opposing directions for

activating and repressing marks. There are also significant changes of histone marks

elsewhere along chromosome 22q and genome wide. Chromosome interaction patterns are

weakened within the deletion boundaries and strengthened between the deletion proximal

regions. We detected a change in the manner in which chromosome 22q folds onto itself,

namely by increasing the long-range contacts between the telomeric end and the deletion

proximal region. Further, the large CNV affects the topological domain that is spanning its

genomic region. Finally, there is a widespread and complex effect on chromosome

interactions genome-wide, i.e. involving all other autosomes, with some of the effect directly

tied to the deletion region on 22q11.2.

Conclusions: These findings suggest novel principles of how such large genomic deletions

can alter nuclear organization and affect genomic molecular activity.

Keywords: Germline CNVs, Chromatin States, Chromosome Interactions, Topological

Domains

Introduction

Two of the most exciting discoveries in human genetics of the past decade are that small- to

medium-sized Copy Number Variants (CNVs) are very common in the human genome, and

that there is a group of large CNVs that are strongly associated with brain development and

neuropsychiatric disorders such as schizophrenia and the autism spectrum disorders

(ASDs) [1,2]. These large CNVs are understood to be providing enticing points of entry to the

analysis of the strong but complex genetic and molecular (and possibly even cellular) basis

of these common disorders.

We now know that small to medium-sized CNVs, i.e. deletions or duplications of

genomic DNA sequence ranging in size from hundreds to tens of thousands of basepairs, are

present in any human genome with their numbers in the thousands [3-7]. Not very much is

known as of yet about such smaller CNVs' effects on the normal phenotype, but some

examples already exist that show that such effects could be considerable, such as the copy

number variation of the amylase gene associated with a given human population's ability to

digest high-starch food [8].

Large CNVs, typically sized from hundreds of thousands to millions of basepairs of

genomic DNA sequence, were previously known to be in strong association with often-

severe but rare congenital malformations, or found in cancer genomes. It was a striking

discovery when a series of studies [1,2] showed that there is a group of more than ten large

CNVs that are strongly associated with aberrant brain development and a resulting

neuropsychiatric phenotype such as schizophrenia, ASD or Williams Syndrome. These large

neuropsychiatric CNVs can encompass many genes and their effects across the various

molecular levels of gene activity and regulation and the clinical phenotype are complex and

only poorly understood.

On the molecular level these large neuropsychiatric CNVs have been mostly studied by

applying the research paradigm of trying to determine which single gene from within the

CNV boundaries is at the root of the large CNV's effects on brain development. Many very

interesting insights have been gained using this approach into a considerable number of

genes with these large CNVs.

However, these insights about individual genes fall far short of explaining the full

effects of the large CNVs. Also there already have been a number of transcriptome wide

studies that at least hint at certain network effects emanating from the large CNVs. Which

mechanisms mediate such transcription network effects is then the question. Furthermore

there are an increasing number of studies that show a potentially very important role of

chromatin regulation in the molecular etiology of neuropsychiatric disorder [9-15].

Against this backdrop we reasoned that it was worthwhile testing whether large CNVs

with association to brain development might cause a disruption or at least alteration of one

or several aspects of chromatin conformation, such as the distribution of regulatory

chromatin marks, the long-range direct physical interactions between distant regions on

one chromosome or between different chromosomes or the higher-order chromatin domain

structures that are defined by such marks or interactions. Such effects on these important

layers of molecular regulation of gene activity would then constitute a novel principle by

which large CNVs could transmit their presence to the machinery of cellular physiology.

Here we show in a cohort of cell lines derived from patients with 22q11 Deletion

Syndrome that indeed chromatin marks, chromatin domains and long-range chromosome

interactions are affected in several distinct ways by a large, common and disease-associated

CNV on chromosome 22q11.2. We then go on to show in a smaller set of cell lines from

different patients that at least some of the same observations can also be made with another

neuropsychiatric large CNV, on chromosome 1q21.1.

22q11 Deletion Syndrome (22q11DS) is a disorder caused by a heterozygous deletion

of 3 million basepairs containing more than 60 known genes on chromosome 22q11.2. It

occurs in at least 1 in 4,000 live births. The common phenotypes of 22q11DS include a large

spectrum of congenital anomalies, for example of the facial structures and the

cardiovascular system – and notably there is a strong association with several

neurodevelopmental psychiatric disorders, in particular schizophrenia and ASDs [2,16-19]. In

this study we are using the large CNV on 22q11.2 as a model to determine the generalizable

principles along which large CNVs of this important category can lead to changes to the

various ways in which chromatin is ordered, using unbiased, genome-wide, sequencing-

based assays for discovery.

Results

High-volume sequencing data generation for Hi-C chromosomal contact maps, ChIP-
Seq chromatin marks and RNA-Seq gene expression data

To determine possible effects of the 22q11.2 deletion on chromosomal interactions, we

generated Hi-C contact maps for 11 human lymphoblastoid cell lines (5 patient cell lines

with 22q11.2 deletion and 6 control cell lines without), with a total of 3.1 billion Hi-C

contact-reads of which 680 million read-pairs were of high quality and used for the

downstream analyses (**Supplementary Table 1**). The existence of the 3 Mbp deletion in the

patient cell lines was validated by whole genome sequencing **(Supplementary Fig. 1)**.

As a first means of quality control we included in our data generation and analysis the

cell line GM06990, which was the cell line used in the original Hi-C paper [20].

Interchromosomal contacts of GM06990 as determined by our own Hi-C data for this line

(**Supplementary Fig. 2a**) show the same patterns of chromosomal interactions across the

nucleus as in [20]; i.e. small chromosomes generally have more interactions with each other

than larger chromosomes with each other, and many more than chromosomes in the

medium size range. Specific interaction pairs, e.g. between chromosomes 17 and 19 were

also replicated in our GM06990 data. Global interchromosomal contact maps resulting from

combining all our controls and cases, respectively, again replicated these global interaction

patterns (**Supplementary Fig. 2b,c**).

To determine the effects of 22q11.2 deletion on gene expression patterns and

chromatin marks we performed RNA-Seq on 14 cell lines (**Supplementary Table 1**) and

ChIP-seq of H3K27ac and H3K27me3 for 6 cell lines and CTCF for 5 of the same cell lines,

respectively (**Supplementary Table 2**).

Normalization of Hi-C data

Many factors, such as mappability of sequencing reads, GC content, length of the restriction

enzyme fragment etc., can lead to biases in Hi-C data. Several alternative computational

approaches have been developed for the normalization of Hi-C data with the aim to remove

these biases [21-30]. While these approaches use different algorithmic principles, it has been

demonstrated that when using them on Hi-C data generated for control genomes without

large CNVs, such as GM12878, the normalization metrics will be highly correlated across the

various normalization schemes [31]. For our Hi-C data however, we needed to be certain to

use a normalization method that is not thrown off from the outset by the presence of the

heterozygous 3 Mbp deletion in 22q11.2.

We tested three commonly used different normalization algorithms that were

developed for Hi-C data, using the rationale that the normalization methods should not

change the general patterns of interaction we can see in the raw data.

We found that not all of the available normalization methods are robust for use with

Hi-C data coming from genomes with large CNVs but that the hicpipe algorithm [21] is quite

suitable for this purpose (details of the comparison between normalization methods in the

Supplementary Information and **Supplementary Fig. 3**).

Chromosomal contacts decrease within the 22q11.2 deletion boundaries and

between the deletion region and the remainder of chromosome 22q

We observed that the chromosomal contacts within the 22q11.2 deletion regions in

22q11DS lines decreased significantly compared to control cell lines (**Fig. 1a**). Also strongly

reduced in the deletion cell lines were the chromosomal contacts between the 22q11.2

deletion region with the entire remainder of chromosome 22 (**Fig. 1a; Supplementary Fig.**

4a). This dosage effect is consistent with the copy number of the 22q11.2 deletion region in

the patient cell lines, as all of the 22q11DS cell lines are heterozygously deleted for this

region. No such dosage effect on the chromosomal contacts was observed that did not

involve the 22q11.2 region. Furthermore, there was no such strong dosage effect on

intrachromosomal contacts on any of the other autosomes (for example chromosome 19 in

Fig. 1b; Supplementary Fig. 4b).

Figure 1. Effects of the 22q11.2 deletion on chromosome contacts, gene expression and epigenetic profiles. *Panels a-d: each pixel in the heat-maps represents the intra- or inter-chromosomal contact frequency in Hi-C data from 22q11.2del cell lines versus control cell lines for a 500 kbp region. Yellow dashed lines indicate the 3 Mbp deletion on chromosome 22q. (a) Fold change of cis-contacts along chromosome 22 in 22q11.2del versus control cell lines. Black boxes indicate increased contacts between the deletion-flanking regions in 22q11.2del cell lines. Blue box: the signal for increased contacts between the downstream deletion-flanking region and the telomeric end of chromosome 22q (red arrows and dashed red line indicate the corresponding chromosome folding event). (b) Lack of intra-chromosomal fold change of contacts for chromosome 19. (c) Fold change of inter-chromosomal contacts between*

chromosome 22 and chromosome 17. (d) Lack of intra-chromosomal fold change of contacts

between chromosome 18 and chromosome 19. (e) Log2 transformed fold change of gene

expression for genes on chromosome 22q in RNA-Seq data from 22q11.2del versus cell lines.

Each point represents a gene. Panels f-h: Log2 transformed fold change in ChIP-Seq signals in

22q11.2del versus control cell lines. (f) H3K27ac histone modifications. (g) H3K27me3 histone

modifications. (h) CTCF binding sites. Black dashed lines indicate the 3 Mbp deletion on

chromosome 22q in e-h.

Then we investigated whether this dosage effect of the 22q11.2 deletion on *cis*-contacts
(i.e. within the deletion boundaries and between the deletion region and elsewhere on
chromosome 22q) holds for *trans*-contacts as well (i.e. for contacts between the region
within the deletion boundaries and the rest of the genome). We found the *trans*-contacts
involving the 22q11.2 deletion in the patient cell lines and any other chromosome also
decreased compared to control cell lines (**Fig. 1c; Supplementary Fig. 4c**). No other
regions of chromosome 22q showed such an effect, and neither did any other pair of
autosomes that did not include 22q (**Fig. 1d; Supplementary Fig. 4d**).

**The 22q11.2 deletion has an effect on gene expression and epigenetic profiles in the
22q11.2 region**

Next we set out to identify the effects of the 22q11.2 deletion on gene expression and
epigenetic profiles. RNA-Seq analysis showed that all of the genes that were expressed
within the deletion boundaries in the 22q11.2 region showed decreased expression in
22q11.2DS patient lines relative to control cell lines (**Fig. 1e**). Our findings are consistent
with a previous study [32].

Differential pattern analysis of H3K27ac, H3K27me3 and CTCF by ChIP-Seq showed the

majority of binding sites for these essential chromatin marks within the 22q11.2 deletion

boundaries with decreased signals in 22q11.2DS patient lines compared to control cell lines

(**Fig. 1f,g,h**). These decreased binding signals for such a large region are specific to

chromosome 22q11.

Chromosome contacts increase across the 22q11DS breakpoint junction and

chromatin marks in its flanking regions are affected in a concerted manner

The flanking regions upstream and downstream of the 22q11.2 deletion are brought into

close proximity to each other by the formation of the deletion breakpoint junction. We

hypothesized that since Hi-C contacts between two given regions will increase with

decreasing genomic distance, the contacts between the upstream and downstream deletion-

flanking regions in 22q11DS patient cells would be markedly enhanced. We indeed found

such stronger contacts between the deletion-flanking regions in 22q11.2DS cell lines (**Fig.**

1a; Supplementary Fig. 3a). These regions of increased contact extended out to two

million base pairs both upstream and downstream of the 22q11.2 deletion boundaries.

Following this observation we wanted to examine whether there is an effect on the

chromatin marks concurrent to these increased chromosomal contact patterns. To do so we

performed enrichment analysis in our ChIP-Seq data for significantly differential signal

patterns for H3K27ac, H3K27me3 and CTCF, respectively. We found that both the upstream

and downstream flanking regions of the 22q11.2 deletion were enriched with significantly

differentially bound sites of both H3K27ac and H3K27me3 (**Fig. 2a,b**). For CTCF the

downstream deletion-flanking region was enriched with significantly differentially bound

sites (Fisher's exact test $p = 2.53\text{E-}06$) (**Fig. 2c**).

Figure 2. Distribution of significantly differentially enriched H3K27ac histone marks (a), H3K27me3 histone marks (b) and CTCF binding sites (c) along chromosome 22q. Each point represents a 500 kbp bin. Red dashed lines mark the boundaries of the 3 Mbp deletion in 22q11.2.

More specifically, within the upstream deletion-flanking region, from 18 to 18.5 Mbp, we found that 5 out of 24 sites with binding by H3K27ac and 6 out of 15 sites with binding

by H3K27me3 showed significantly differential binding (Fisher's exact test p = 0.0075 for

H3K27ac and p = 5.67E-05 for H3K27me3 respectively). Intriguingly, for H3K27ac all of the

5 sites with significant differential binding were bound less strongly while for the same

region for H3K27me3 all of the 6 sites with significant differential binding were bound

more strongly in the 22q11DS patient lines. For the downstream region, from 23Mb to

23.5Mb, 5 out of 13 sites with H3K27ac binding and 11 out of 30 sites with H3K27me3

binding showed significantly differential binding (Fisher's exact test p = 0.0004 for

H3K27ac and p = 1.26E-07 for H3K27me3 respectively). Again we observed the reciprocity

between significantly differential changes for the two different histone marks: 4 out of 5

sites for H3K27ac gave a less strong signal while all of the 11 of such sites for H3K27me3

showed a stronger signal in the 22q11.2DS patient cell lines.

Such reciprocity in signal strengths between these two histone marks is a general

feature of their principle of action and makes it much more likely that the observed changes

are of physiological relevance.

Next we investigated whether there are gene expression changes in these same

deletion-flanking regions. Based on the observed changes in histone marks we reasoned

that gene expression in these regions could be downregulated in the 22q11DS patient lines,

as H3K27ac is associated with active genes and H3K27me3 with repressed genes. We found

that among the 4 genes with detectable expression in the deletion-upstream region from

18Mbp to 18.5 Mbp, the gene *BID* showed significantly decreased expression in 22q11.2DS

patient lines while the other three remained unchanged. In the deletion-downstream

region, from 23 Mbp to 23.5 Mbp, only one gene, *IGLL5,* had detectable levels of expression

and we found it to exhibit significantly decreased expression in 22q11.2DS patient lines as

well.

Taken together this part of our study shows that the effect of the 22q11.2 deletion is

not limited to the genes within the deletion boundaries but also exerts extensive influence

on its flanking regions on the level of intrachromosomal contacts, histone modifications and

gene expression.

A/B compartments of the chromatin and the topological domain structure of the

nuclear genome are affected by the 22q11.2 deletion

Previous studies of cell lines without large CNVs, have revealed that there is one level of

organization where the chromatin is partitioned into two compartments, A and B, which

correspond to active and inactive regions, respectively [20], and which can be derived from Hi-

C data. We sought to determine whether the 22q11.2 deletion might lead to changes in

these A/B compartments. Indeed when computing A/B compartments from our Hi-C data

we observed that the eigenvectors changed signs in the 22q11.2 del region while all the

other regions remained the same between 22q11.2DS patient and control cell lines (**Fig.**

3a). The original compartment that is spanning the deletion region in control cells is being

partitioned into two shorter compartments with different eigenvectors signs, signifying a

change in the A/B compartment structure in the deletion region.

Figure 3. Change in A/B compartments and topological domain signal on chromosome 22q11. *(a) The A/B compartment spanning the 22q11 deletion in controls is partitioned into two compartments in the patient cell lines (red arrows). Red dashed lines: boundaries of 22q11.2del region. Shown is the first eigenvector for the principal component analysis of the normalized contact matrix. X-axis: position on chromosome 22q. Y-axis: value of the first eigenvector. (b) Normalized Hi-C contacts for 22q11.2del and control cell lines, displayed as a triangular heat-map that is overlayed on directionality indices (DI) and chromatin domains. The signal for the topological domain spanning the 22q11.2 deletion and its downstream region (blue triangle in controls) was broken into two shorter signals (blue triangles and black arrows in 22q11.2del). Black dashed lines: boundaries of 22q11.2del region. Regions with low mappability of reads were removed from the analysis and are shown in grey.*

Topological domains are a megabase-sized structural feature of the genome

organization that is constituted of highly self-interacting chromosome regions [33]. Using our

Hi-C and ChIP-Seq data we identified clear changes of the topological domain structure in

the deletion-downstream flanking region of 22q11.2DS cell lines while the deletion-

upstream topological domain structure was preserved (**Fig. 3b**). The topological domain

spanning the 22q11.2 deletion and its downstream region was broken into two shorter

domains across the downstream boundary of the deletion. Of note, this effect of the 22q11.2

deletion on the topological domain structure extended to 2 million base pairs downstream

of the deletion region, where we observed another topological domain breaking into two

shorter ones. Our results demonstrate that the 22q11.2 deletion can result in changing the

order of topological domains both in the deletion region proper and its downstream

regions.

Change of chromosomal *cis*-contacts involving the downstream flanking region of

22q11.2del

Based on the above findings that the genomic region downstream of the 22q11.2 deletion

was affected on different levels of molecular regulation in 22q11.2DS patient cell lines, we

next sought to investigate whether the presence of the 22q11.2 deletion also affected the

chromosomal *cis*-contacts of the downstream deletion-flanking region with any other

region on chromosome 22q. To do so, we analyzed our Hi-C data by calculating the fold

change of contacts between the 21.5-22 Mbp window, which is situated right downstream

of the 22q11.2 deletion, and all the other 1 Mbp-sized regions on chromosome 22q. As

expected, we observed that region 17-18 Mbp, i.e. the breakpoint-proximal region right

upstream of the deletion, showed the largest fold change (2.04) of contact with region 21.5-

22 Mbp between 22q11DS and control cell lines, given that these regions were brought into

close proximity to each other by the 22q11.2 deletion (**Fig. 1a**).

The second largest fold change (1.96) of *cis*-contacts involving region 21.5-22 Mbp was

for contacts with region 50-51 Mbp, i.e. towards the very telomeric end of chromosome 22q

(**Fig. 1a**). Intriguingly, we also observed strong positive correlation between region 22-22.5

Mbp and region 50-50.5 Mbp for CTCF binding (Pearson's r = 0.933, p = 0.02)

(**Supplementary Fig. 5a**) and between region 22-22.5 Mbp and region 50.5-51 Mbp for

H3K27ac enrichment (Pearson's r = 0.811, p < 0.05) (**Supplementary Fig. 5b**).

Furthermore there was weak correlation between region 22-22.5 Mbp and 50.5-51 Mbp for

H3K27me3 enrichment (Pearson's r = 0.74, p = 0.090) (**Supplementary Fig. 5c**). Given that

both region 21.5-22 Mbp and region 22-22.5 Mbp are in the downstream flanking region of

the 22q11.2 deletion, our results strongly imply that increased chromosomal contacts

between the downstream deletion-flanking region and the telomeric region 50-51 Mbp may

be associated with the differential changes of histone modifications and CTCF binding that

we found to be in correlation between these two regions.

To validate the increased chromosomal contact between the downstream deletion-

proximal region and the telomeric end of 22q we performed DNA FISH on 5 cell lines with

the 22q11.2 deletion and on 6 control cell lines, using FISH probes to loci in region 21.8-

22.5 Mbp (RP11-47L18: chr22: 21,931,796 – 22,118,344) and in region 50-51Mbp (RP11-

125K3: chr22: 50,149,996- 50287311). FISH showed that the downstream flanking region

of the 22q11.2 deletion is significantly closer to region 50-51 Mbp in nuclear space in

22q11DS cell lines than in control cell lines (p value 0.008) (**Fig. 4; Supplementary Fig. 6**).

Taken together, our results strongly indicate that the 22q11.2 deletion causes

conformational changes on several levels on chromosome 22q.

Figure 4. Examples of 3D FISH visualization of intra-chromosomal interaction changes.

The Hi-C predicted interaction regions of 21.8-22.5 Mbp and 50-51 Mbp on chromosome 22q

were visualized by 3D DNA FISH using BAC probes RP11-47L18 and RP11-125K3, labeled with

biotin (red) or digoxigenin (green). Arrow: an example for the red and green FISH probes in

close proximity in a cell carrying the 22q11.2 deletion. Scale bars are 5 μm. Red and green bars

on chromosome 22q indicate the locations of the biotin and digoxigenin FISH probes,

respectively. Black dashed lines indicate the position of the 3 Mbp deletion.

Global changes of interchromosomal contact patterns

To explore the effect of the 22q11.2 deletion on *trans*-contacts, i.e. between any non-

homologous autosomes in the nucleus of 22q11.2DS patient lines, we analyzed our Hi-C

data for significantly different *trans*-contacts between 22q11DS and control cell lines. On

the genome-wide level, we found 272 *trans*-contacts with a p-value of less than 0.0001 (**Fig.**

5a). Interestingly the majority of these chromosomal *trans*-contacts did not involve

chromosome 22q as one of the interacting partners. Notably, 56 of these interchromosomal

contact signals are among the top 5% of the strongest *trans*-contacts (**Fig. 5b**). This

enrichment of significantly different genome-wide chromosomal *trans*-contacts within the

strongest *trans*-contacts is again highly statistically significant (Fisher's exact test $p < 2.2e-$

16). We found many fewer significantly different genome-wide chromosomal *trans*-contacts

by randomly swapping the assignment of 22q11.2 deletion and control status across our Hi-

C data sets, by comparing within control cell lines and by comparing within 22q11.2del cell

lines (**Supplementary Table 3**). None of these swapping analyses achieved the same

enrichment. This indicates that a relevant amount of the significantly different genome-

wide chromosomal *trans*-contacts is not due to random chromosomal motion or to as-of-yet

unknown factors such as cell-culture variations across the lymphoblastoid cell lines. Rather,

our analysis points to a genome-wide disturbance of the network of chromosomal *trans*-

contacts that is at least in part attributable to the presence of the 22q11.2 deletion on

chromosome 22q.

Figure 5. Genome-wide inter-chromosomal contact changes (either more or less frequent) determined by Hi-C analysis in 22q11.2del versus control cell lines. *(a) Circos plot of the inter-chromosomal contacts exhibiting differential interaction in 22q11.2del versus control cell lines at significance level of 0.0001. (b) Circos plot of the inter-chromosomal contacts exhibiting differential interaction in 22q11.2del versus control cell lines at significance level of 0.0001 and showing only the top 5% strongest inter-chromosomal contacts.*

Global changes of gene expression patterns

To investigate the global effect of the 22q11.2 deletion on gene expression, we performed differential expression analysis between the 22q11.2DS and control cell lines. Of the 11,374 genes with detectable levels of expression (FPKM > 0.5), 1,610 genes showed significantly

differential expression (FDR < 0.05). Gene ontology analysis indicated that these

differentially expressed genes are enriched for genes involved in mitochondrial pathways

such as the respiratory chain (n = 32, p value = 2.96E-11). KEGG pathway analysis

demonstrated   enrichment   of   genes   involved   in   oxidative   phosphorylation

neurodegenerative diseases such as Parkinson's disease (**Fig. 6a**). Earlier studies have

noted that there are several genes related to mitochondrial function that are located within

the 22q11 deletion boundaries [34], therefore the pathways associated with this cellular

function would have a high likelihood to be affected by the change in copy number. The

enrichment for pathways related to neurodegenerative disorder is encouraging, as an

association between 22q11DS and Parkinson's Disorder has been reported previously [35-37].

This reinforces the notion that the LCL cell culture model can be of relatively more

relevance for the molecular study of 22q11DS in general.

Figure 6. Differential expression analysis and correlation between gene expression and epigenetic profiles. (a) KEGG pathway analysis for genes differentially expressed between 22q11.2del and control cell lines. (b) Enrichment analysis for chromosome 22q for differentially expressed genes. Red dashed lines mark the boundaries of 22q11.2del region. (c) Differentially expressed genes correlated with different H3K27ac and H3K27me3 signal strengths in 22q11.2del versus control cell lines. A1 represents genes significantly up-regulated in 22q11.2del cell lines while B1 represents non-significantly up-regulated genes; A2 represents genes significantly down-regulated in 22q11.2del cell lines while B2 represents non-significantly down-regulated genes. The y-axis shows the z-score transformed fold change of H3K27ac signals (left) or H3K27me3 signals (right) between 22q11.2del and control cell lines. (d) Differential H3K27ac and H3K27me3 signal strengths exhibited correlation with differential gene expression between 22q11.2del and control cell lines. A1 represents genes whose TSSs were significantly more marked by H3K27ac (left) or by H3K27me3 (right) in 22q11.2del cell lines. B1 represents genes with TSSs non-significantly more marked by H3K27ac (left) or by H3K27me3 (right). A2 represents genes whose TSSs were significantly less marked by H3K27ac (left) or by H3K27me3 (right) in 22q11.2del cell lines. B2 represents genes with TSSs non-significantly less marked by H3K27ac (left) or by H3K27me3 (right). The y-axis shows the z-score transformed fold change of genes' FPKM in 22q11.2del versus control cell lines.

We next sought to identify whether genome-wide there were entire genomic regions that are enriched for differentially expressed genes. As expected, the most significant signals for this analysis were located in the 22q11.2 region (**Fig. 6b; Supplementary Fig. 7**). No other regions genome-wide achieved FDR corrected significance (**Supplementary Fig. 7**).

Correlation between histone modification and gene expression

To examine whether the gene expression changes are consistent with the histone modification changes, we assigned the H3K27ac and H3K27me3 peaks to their nearest genes based on the distance to their transcription start sites and for each gene, we only retained the closest peak for both histone marks. We observed that significantly upregulated genes in 22q11.2DS cell lines exhibited significantly higher fold change of H3K27ac enrichment (permutation test p = 0.0140) and significantly lower fold change of H3K27me3 enrichment (permutation test p = 0.0018) than genes non-significantly upregulated in 22q11.2DS cell lines (**Fig. 6c**). Consistently, significantly downregulated genes in 22q11.2DS cell lines exhibited significantly lower fold change of H3K27ac enrichment (permutation test p = 0.0002) but significantly higher fold change of H3K27me3 enrichment (permutation test p = 0.0227) than genes non-significantly down regulated in 22q11.2DS cell lines (**Fig. 6c**).

Moreover, genes whose transcription start sites (TSSs) showed significantly upregulated binding by H3K27ac in 22q11DS cell lines exhibited significantly higher fold change of expression (permutation test p < 2.2e-16) than those with non-significantly upregulated binding, while those TSSs showing significantly downregulated binding exhibited a significantly lower fold change of expression (permutation test p = 0.0202) than those with non-significantly downregulated binding (**Fig. 6d**). Consistently, genes whose TSSs showed significantly upregulated binding by H3K27me3 in 22q11DS showed significantly lower fold change of expression (permutation test p = 0.0037) while those with significantly downregulated binding exhibited significantly higher fold change of expression (permutation test p = 0.0027) between 22q11DS and control cell lines (**Fig. 6d**).

We also calculated the correlation between gene expression and H3K27ac binding

affinity across the individuals for all the genes with TSSs bound by H3K27ac. In line with the

above results, we found significantly higher Pearson's correlation coefficients than the

coefficients obtained from permutations (Wilcoxon rank sum test p < 2.2e-16)

(**Supplementary Fig. 8a**). This difference was even more significant when we included the

differential expressed genes (absolute fold change > 2) only (Wilcoxon rank sum test p <

2.2e-16) or the differential H3K27ac bound genes only into our analysis (Wilcoxon rank

sum test p < 2.2e-16) (**Supplementary Fig. 8a**). Unsurprisingly, we also observed

significantly higher Pearson's correlation coefficients between gene expression and

H3K27me3 binding than the coefficients obtained from permutations for the differential

expressed genes (absolute fold change > 2) only (Wilcoxon rank sum test p = 9.72e-04) and

the differential H3K27me3 bound genes only (Wilcoxon rank sum test p = 3.71e-06)

(**Supplementary Fig. 8b**). Together, our results demonstrated that gene expression

changes are associated with histone modification changes in 22q11DS cell lines compared

with control cell lines.

Effects of a large CNV on chromosome 1q21.1 on chromosome folding

To explore whether large CNVs other than the one on chromosome 22q11.2 can lead to

changes in the patterns of chromosome folding, we performed Hi-C on two lymphoblastoid

cell lines with a heterozygous deletion of approximately 1.35 Mbp in size on chromosome

1q21.1 (1q21.1del). This deletion is strongly associated with the development of

schizophrenia [38-40]. As for the 22q11.2 deletion, we observed that both *cis*-contacts and

trans-contacts between the 1q21.1 deletion regions and other regions were decreased in

1q21.1del cell lines relative to control cell lines (**Fig. 7a,b**). As observed in the 22q11DS cell

lines, in the 1q21.2del cell lines there was an increase of intrachromosomal contacts

between the regions directly flanking the main CNV of 1q21.1 (**Fig. 7a**). Analysis of A/B

compartments showed that the eigenvector sign of the downstream boundary region of the

1q21.1 deletion was inverted (**Fig. 7c**), showing that also on the level of chromatin

compartmentalization the large CNV on 1q21.2 can have effects, similar to the large CNV on

22q11.2. Taken together the large deletion CNV on chromosome 1q21.1 exhibited similar

effects on chromosomal *cis*-contacts and *trans*-contacts as the large deletion CNV on

chromosome 22q11.2, which points towards our findings in 22q11DS being generalizable

across large neuropsychiatric CNVs.

Figure 7. Effect of the large deletion CNV on 1q21.1 on chromosome conformation. *(a)*

Fold change of cis-contacts of chromosome 1 in 1q21.1del versus control cell lines. Blue boxes

mark the regions of increased contacts between the upstream and downstream flanking

regions of the 1q21.1 deletion. (b) Fold change of trans-contacts between chromosome 1 and 4

 in 1q21.1del cell lines versus control cell lines. Black dashed lines mark the boundaries of the

1q21.1 deletion. Each cell in the heatmap represents the inter-chromosomal contact level

between two 500 kbp regions. (c) Change in A/B compartments on chromosome 1q21. Upper

panel: control cell lines. Lower panel: 1q21.1del cell lines. Shown is the first eigenvector for the

principal component analysis on the normalized contact matrix of chromosome 1, for

1q21.1del and control cell lines. X-axis: position on chromosome 1q. Y-axis: value of the first

eigenvector. Black arrows indicate the location of change in the eigenvector sign in the

downstream boundary region of 1q21.1, in 1q21.1del cell lines.

Discussion

Large CNVs are an important feature of the genetic architecture of several major

neurodevelopmental psychiatric disorders as well as of conditions involving aberrant

morphology of many organ systems. Their effects on the level of phenotype are complex and

the molecular mechanisms mediating these effects are very incompletely understood. While

it is still a good assumption that a considerable portion of these mechanisms are a direct

consequence of the copy number change of the genes within a given CNV's boundaries and

the resulting changes in expression levels for these genes, it also seems plausible to

investigate whether additional levels of complexity exist regarding the effects of a large CNV

across multiple layers of the control of gene activity. This plausibility stems from several

observations and lines of reasoning. Namely there are large numbers of genes affected

genome wide and far distal from the main CNV, and expression changes within CNV

boundaries do not always follow exactly the change in copy number. This leads us to

consider the basic principles that govern organizational features of the nucleus and the

chromatin such as regulatory domains being bounded by protein factors that recognize

binding sites in the DNA sequence which in turn could be affected (i.e. deleted or

duplicated) by a CNV, or long-range chromosome contacts being also influenced by the

length of the involved chromosomes (and that length being changed by large CNVs). Finally

there have been several recent reports about mutations in chromatin remodeling genes in

the context of neuropsychiatric disorders pointing to the importance of proper molecular

management on the epigenomic level in these conditions.

Here we studied the effects of the important large CNV on chromosome 22q11.2, and in

a more limited fashion the effects of the large CNV on chromosome 1q21.2, on chromatin

conformation including long-range chromosome contacts and domain formation, epigenetic

profiles and gene expression. To our knowledge, this is the first study of changes in

chromatin conformation and epigenetic profile changes caused by deletions on 22q11.2,

and 1q21.2, or any such large CNVs with strong relevance to organ development and in

particular brain development and neuropsychiatric disorders. We observed dosage effects

of the large CNVs on long-range chromosome contacts, chromatin domains and epigenetic

profiles as well as on gene expression. More specifically we found increased contacts

between upstream and downstream flanking regions of the 22q11.2 deletion in 22q11DS

cell lines in contrast with control cell lines. Interestingly, both upstream and downstream

regions of the 22q11.2 deletion were enriched with differentially binding sites of H3K27ac

and H3K27me3 while downstream regions were also enriched with differentially binding

sites of CTCF. Importantly, the gene expression changes in these regions were consistent

with the histone modification changes as less H3K27ac binding and more H3K27me3

binding reflect decreased gene expression in 22q11Ds cell lines. Based on the above

findings, it is tempting to speculate that increased contacts between upstream and

downstream flanking regions of the 22q11.2 deletion contribute to gene expression changes

via the coordinated changes of epigenetic profiles in these flanking regions.

Although we found chromosome contacts to be correlated with both the epigenetic profiles and the gene expression changes in the flanking regions of the 22q11.2 deletion, we did not observe significant correlation either between chromosome contact changes and gene expression changes or between chromosome contact changes and epigenetic profile changes (data not shown) on the genome-wide level. Moreover, we did not observe the end of chromosome 22 (50-51Mb) to be enriched with differentially expressed genes (2 out of 24 genes) despite the Hi-C finding, validated by FISH, of significantly increased contacts with the downstream flanking region of the 22q11.2 deletion in 22q11.2DS cell lines. Our results indicate that chromosome conformation changes caused by the 22q11.2 deletion contribute to gene expression changes and epigenetic profile changes between 22q11.2DS and control cell lines but not always in a clearly and linearly deterministic way.

However, we did observe significant positive correlation between H3K27ac changes and gene expression changes but negative correlation between H3K27me3 and gene expression changes on the genomic level. These findings demonstrated that the epigenetic profiles were reshaped genome-wide to cause extensive gene expression changes in 22q11.2DS cell lines.

The dosage effects of the deletion on *cis*- and *trans*-contacts and the increased chromosomal contacts between upstream and downstream deletion-flanking regions in 22q11Ds cell lines were also observed in cell lines with the 1q21.1 deletion. The similar results that were obtained on that level of observation for the 22q11.2 deletion and the 1q21.1 deletion would suggest that these changes caused by a large deletion CNV are universal for such chromosomal aberrations instead of specific to the 22q11.2 deletion.

As a cautionary note on the technical level, we demonstrated here that for genomes with a large deletion CNV the appropriate normalization methods for Hi-C data have to be

chosen with great care to avoid false findings. For instance we would have reached the

conclusion that the chromosomal contacts within the deletion regions are not decreased in

cell lines with deletion compared with control cell lines if the hiclib software package [22] had

been used for normalization of Hi-C data, instead of the hicpipe package [21].

Lastly, we also observed A/B chromatin compartment changes and changes in the

signal from topological domain analyses across the 22q11.2 deletion region. This is the first

time that such a phenomenon has been reported in this context. Earlier, Lupiáñez et al. [41],

using the 4C assay, demonstrated in a mouse model as well as in human lines from patients

with rare malformations of the limbs, that large structural changes in the genomic sequence

can affect the topological domain architecture directly on top of the sequence change, giving

further support to our Hi-C based findings in 22q11.2 and 1q21.1 deletion lines. It is

important to remember that, just as for earlier studies that were reporting on A/B

compartments and topological domains (in non-CNV genomes), our analysis resulted from

combining the sequencing data of two homologous chromosomes. Only once haplotype

phasing of the genomes in question will be available will one be able to determine whether

the chromosome 22q with large deletion or the one without large deletion, or some

combination of the two, is driving such changes in the domain signals (the deletion-carrying

domain being the main cause for the change in signal certainly being the initial hypothesis).

Whether the observed chromosomal contact changes on chromosome 22q are existent only

on one chromosome 22q (again probably the one carrying the large CNV) or emanate from

both of the two homologous chromosomes is currently unknown and will be worth

investigating when haplotype-phasing data is available.

In summary, we found dosage effects of the large deletion CNV on chromosome

22q11.2 on long-range chromosome contacts, chromatin organization, epigenetic profiles

and gene expression. Extensive changes on these levels caused by the 22q11.2 deletion are

global and seem to be rippling along the entire chromosome carrying the deletion as well as

across the entire nucleus rather than being confined to the deletion region only. Such effects

had never been shown before outside of cancer cell lines [42]. Furthermore, in contrast to the

findings in cancer cells we used a larger cohort of individual patient cell lines, with all lines

carrying only one main large CNV that is clearly and strongly associated in a causative

manner with a neurodevelopmental phenotype. This, in contrast to the two cancer cell lines

that each carried multiple large CNVs which could also have been a consequence rather

than a cause of the disease phenotype, makes it much more likely that the higher-order

effects of the large CNVs that we observed may be contributing to the molecular etiology of

the disorders in question. This point is further strengthened by another recent paper [43],

where the authors describe studying the effects of large CNVs on chromosome 16p11 on

chromosome interactions. These large CNVs are almost as strongly associated with

neurodevelopmental disorders than the large CNV on 22q11.2. While the study on the

16p11 CNVs used a somewhat smaller number of cell lines than our study and also used 4C

as a method of discovery, which is, unlike the Hi-C method used by us, not able to detect

changes in a global and unbiased fashion, it is one more independent piece of supporting

evidence for the biological validity and general relevance of the findings which we describe

here.

While we were able to show possible correlations across several pairs of the molecular

levels that we assayed in this study there are other combinations of molecular levels that

show no obvious connection in our data. We believe that this could be a function of either

the developmental time point or the cell type, or both, being removed from those where the

22q11.2 and 1q21.1 deletions most likely exert some of their strongest effects (e.g. during

embryonal development and in cells of the developing central nervous system). The clear

effects that we were able to observe in lymphoblastoid cell lines could represent the

afterglow of a molecular tragedy that played out earlier in the development of various

organs in the patients carrying these deletions. At the same time the strength of this

afterglow would hint at the strength of the effects that impacted across various levels of

molecular control and gene regulation by CNVs of this size.

<u>Methods</u>

Cell lines and data generation for this study. All cell lines were either acquired from the

Coriell cell repository (cell lines IDs starting with GM or ID) or were taken from the

Molecular Genetics of Schizophrenia (MGS) cohort (dbGaP Study Accession:

phs000167.v1.p, cell lines 52425 and 82699), and were appropriately consented. The Hi-C

assay was carried out according to the original protocol [20], with several modifications. ChIP-

Seq was performed as described in previous studies [44,45]. The RNA-seq libraries were

generated according as in [46]. All experimental procedures are described in the

Supplementary Information.

Hi-C Data analysis. All Hi-C data were produced using Illumina paired-end sequencing with

a read length 2 x 101 bp. As there might be ligation junctions in the reads, we performed

iterative mapping using bowtie2 as in Imakaev et al. 2012 [22]. Briefly, we computationally cut

all the reads to 25 bp first and mapped them to human genome (hg19). Then we extended

the non-uniquely mapped reads by 5 bp to 30 bp and mapped them again. This process was

repeated until the read length was extended to 101 bp. This iterative mapping did improve

the mapping rate (**Supplementary Table 4**). Each read end was mapped separately using

the single ends mode. Only uniquely mapping reads were used and PCR duplicate read pairs

were removed. We only included autosomes in our study. The filtered contact number is

listed in **Supplementary Table 5**.

We compared three different data normalization methods: hiclib [22], hicpipe [21] and

HiCNorm [23]. All of the three tools were run using the default parameters except for the

segment length threshold being set to 600 bp. We chose normalized metrics on hicpipe for

the following analyses, using a bin size of 40 kb for topological domain analysis and of 500

kb for the other analyses. The total number of contacts was normalized for each sample

before combining cell lines in each category (control, 22q11del and 1q21del, respectively).

Fold changes of log2 transformed mean contacts between deletion cell lines and control cell

lines were calculated by (deletion – control)/control.

To identify the differential interchromosomal contacts, we only included contacts with

at least 1 supporting read pair in each of the cell lines. Differential contacts analysis was

conducted by Student's t-test using the normalized metrics. Fisher's exact test was used to

assess the enrichment of differential contacts within the top 5% strongest contacts. We also

performed the same analysis by permuting the control and 22q11del status of the cell lines

ten times. Comparison within control cell lines and within 22q11del cell lines were

performed by randomly dividing the cell lines into two groups three times.

Identification of A and B compartments was performed as in Lieberman et al. 2009 [20]

and topological domains were identified as in Dixon et al. 2012 [33]. We modified the

algorithm for identifying topological domains such that when calculating the direction index

for a given bin we will exclude neighboring bins that are spanned by segmental duplications

as well as the bins on the opposite side of the given bin and equidistant to the given bin than

the bins spanned by segmental duplications (even if such symmetrical bins are not spanned

by segmental duplications themselves). For topological domain analysis, we pooled the raw

data of all the samples for each category to obtain sufficient Hi-C sequencing coverage

before normalization.

3D FISH. Two human DNA BACs (clones RP11-47L18 and RP11-125K3) covering two

distinct regions of chromosome 22 were labeled with biotin or digoxigenin (DIG) by nick

translation kit (Roche Applied Sciences) to make FISH probes (Roche). In-situ hybridization

was performed according to the method published by [47], with several modifications.

Experimental details and data analysis could be found in the **Supplementary Information**.

Differential expression analysis. All RNA-seq data were generated using Illumina paired-

end sequencing with read length 101 bp. Reads were mapped to hg19 and transcriptome

reference with TopHat 2 [48]. TopHat 2 was run using default parameters but with the

coverage search being turned off. The mapped reads were analyzed by Cufflinks [49].

Differential expression was estimated with Cuffdiff 2 [50]. We excluded the genes with low

expression (FPKM < 0.5) from downstream analysis.

Pathway analysis of significantly differential expressed genes was conducted with

DAVID [51] using all the expressed genes as background.

ChIP-Seq data analysis. All ChIP-Seq data were generated using Illumina paired-end

sequencing with read length 101 bp. Reads were aligned to hg19 with BWA-MEM using

default parameters [52]. Reads with low mapping quality (< 30) were removed. PCR duplicate

reads were removed using Picard (http://broadinstitute.github.io/picard). As a quality

control, we calculated the normalized strand cross-correlation coefficient (NSC) and relative

strand correlation (RSC) [53] to assess the signal-to-noise ratios. All the data showed higher

NSC than RSC (**Supplementary Fig. 9**). Replicates for the same cell line on average showed

higher correlation than datasets from different cell lines (**Supplementary Fig. 10**).

For CTCF and H3K27ac, we used MACS2 [54] to call narrow peaks with default

parameters. For H3K27me3, we used the broad peak calling in MACS2. For all peak calling,

we used the corresponding whole-cell extract input library as background. For differential

bound analysis, we used the R package DiffBind [55] with the effective library size for read

counts normalization. Then DBA_DESEQ2 method was employed to conduct the differential

bound analysis. Signal artifact blacklist regions were excluded from our analysis

(http://hgdownload.cse.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeMapabil

ity).

Enrichment analysis. For the enrichment analysis of differentially expressed genes we

divided each chromosome into 500 kbp bins. Within each bin, we calculated the total

number of expressed genes and the number of genes with significantly differential

expression between 22q11del and control cell lines. Then we conducted Fisher's exact test

to identify bins enriched with significantly differentially expressed genes against the

background of the whole genome.

For the analysis of differentially enriched sites for CTCF, H3K27ac and H3K27me3, we

also used 500 kbp bins. Log2 transformed fold changes of normalized read numbers in

binding sites between 22q11del and control cell lines were further transformed to Z-scores.

We considered binding sites with Z-score >2 or < -2 as significantly bound sites. Then

within each 500 kbp bin, we calculated the total number of binding sites and the number of

significantly differentially enriched sites between 22q11del and control cell lines. Then we

conducted Fisher's exact test to identify bins with significantly differentially enriched sites

against the background of the whole genome.

Correlation analysis of gene expression and histone modification. To estimate the

correlation between gene expression and histone modification, we assigned each binding

site of H3K27ac and H3K27me3 to its nearest ENSEMBL TSS using the R package

ChIPpeakAnno [56]. If a TSS was associated with multiple binding sites, only the nearest

binding site was retained. We then used this assignment for downstream correlation

analysis. To determine the cutoff for the distance in which binding sites are associated with

TSSs, we plotted the distribution of distances between binding sites and their assigned TSSs

(**Supplementary Fig. 11**). Based on the distribution, we set the cutoff to distance to TSS to

± 1 kbp for H3K27ac and to ± 5 kbp for H3K27me3.

To investigate the effects of histone modifications on gene expression, we divided the

genes into two categories based on the differential expression analysis: differentially

expressed genes (FDR < 0.05) (category A) and not differentially expressed genes (FDR >

0.05) (category B). Within each category, we further categorized the genes into two groups:

up-regulated expressed genes (A1, B1) and down-regulated expressed genes (A2, B2) in

22q11del cell lines relative to control cell lines. Then for the genes within each of the four

groups (A1, B1, A2, B2), we calculated the Z-score transformed fold changes of the

normalized read counts in the TSS binding sites of histone marks between 22q11del and

control cell lines. To obtain the statistical significance of the fold change differences

between A1 and B1 and between A2 and B2, we performed permutation tests with 9,999

permutations.

We also carried out the reverse analysis. TSS binding sites of histone marks were

partitioned into two categories based on the differential bound analysis: differentially

bound sites (|Z-score| > 1 for H3K27ac, |Z-score| > 2 for H3K27me3) (category A) and non-

differentially bound sites (|Z-score| < 1 for H3K27ac, |Z-score| < 2 for H3K27me3) (category

B). Within each category, we further categorized the binding sites into two groups: up-

regulated bound sites (A1, B1) and down-regulated bound sites (A2, B2) in 22q11del cell

lines. Then we calculated the fold changes of the genes' FPKM between 22q11del and

control cell lines within each group. Permutation tests were performed with 9,999

permutations to obtain statistical significance.

To estimate the correlation between gene expression and histone modification in a

direct way, we calculated Pearson's correlation coefficient between gene's FPKM and

normalized read counts in the corresponding TSS binding site for each gene across all of the

cell lines. To obtain statistical significance, we first permuted genes' FPKM across the cell

lines for each TSS 10 times to assess the background correlation levels, and then performed

the Wilcoxon rank sum test between the observed correlation coefficients and the

background correlation coefficients. We also performed the same analysis using the

differentially expressed genes only and differentially enriched binding sites only.

Correlation analysis for epigenetic marks. To assess the correlation of epigenetic marks

binding between different regions on the same chromosomes, we divided the chromosomes

into 500 kbp bins. Within each bin, we calculated the mean value of the normalized read

counts for all the binding sites of each epigenetic mark. Then we calculated the Pearson's

correlation coefficients of obtained mean values across the cell lines between any two bins

on the same chromosomes.

Data availability. Hi-C, ChIP-Seq and RNA-Seq data from this study have been submitted to

the NCBI Gene Expression Omnibus (GEO; http://www.ncbi. nlm.nih.gov/geo/) under

accession number GSE76922.

Competing interests

The authors declare that no competing interests exist.

Authors' contributions (in order of position in Authors List)

YZ contributed to study design and produced experimental data, XZ contributed to study

design, carried out data analysis and contributed to writing the paper, XZ contributed to

study design and carried out data analysis, CP produced experimental data and contributed

to data interpretation, MSH produced experimental data, TW produced experimental data,

JY produced experimental data, SMW and AEU conceived of and designed the study,

directed data production and data analysis and wrote the paper.

Acknowledgments

This research was supported by external funding from the National Institute of Mental

Health (NIMH) (grant number: MH100010), NARSAD and the March of Dimes, as well as

funds from Stanford Pediatric Research Fund (Grant number: UL1 RR025744) and

departmental funds from the Stanford Department of Psychiatry and Behavioral Sciences.

We thank Dr. Eitan Yaffe (Stanford University) for his advice on the use of hicpipe.

References

1   Kirov, G. CNVs in neuropsychiatric disorders. *Hum Mol Genet* **24**, R45-49,

doi:10.1093/hmg/ddv253 (2015).

2   Torres, F., Barbosa, M. & Maciel, P. Recurrent copy number variations as risk

factors for neurodevelopmental disorders: critical overview and analysis of

clinical implications. *J Med Genet* **53**, 73-90, doi:10.1136/jmedgenet-2015-

103366 (2016).

3   Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**,

68-74, doi:10.1038/nature15393 (2015).

4   Sudmant, P. H. *et al.* An integrated map of structural variation in 2,504

human genomes. *Nature* **526**, 75-81, doi:10.1038/nature15394 (2015).

5   Korbel, J. O. *et al.* Paired-end mapping reveals extensive structural variation

in the human genome. *Science* **318**, 420-426, doi:10.1126/science.1149504

(2007).

6   Abecasis, G. R. *et al.* A map of human genome variation from population-scale

sequencing. *Nature* **467**, 1061-1073, doi:10.1038/nature09534 (2010).

7   Mills, R. E. *et al.* Mapping copy number variation by population-scale genome

sequencing. *Nature* **470**, 59-65, doi:10.1038/nature09708 (2011).

8   Santos, J. L. *et al.* Copy number polymorphism of the salivary amylase gene:

implications in human nutrition research. *J Nutrigenet Nutrigenomics* **5**, 117-

131, doi:10.1159/000339951 (2012).

9   Gavin, D. P. & Floreani, C. Epigenetics of schizophrenia: an open and shut

case. *Int Rev Neurobiol* **115**, 155-201, doi:10.1016/B978-0-12-801311-

3.00005-6 (2014).

10   Fullard, J. F. *et al.* Understanding the genetic liability to schizophrenia

through the neuroepigenome. *Schizophr Res*,

doi:10.1016/j.schres.2016.01.039 (2016).

11   Nestler, E. J., Peña, C. J., Kundakovic, M., Mitchell, A. & Akbarian, S. Epigenetic

Basis of Mental Illness. *Neuroscientist*, doi:10.1177/1073858415608147

(2015).

12   Mitchell, A. C. *et al.* The genome in three dimensions: a new frontier in human

brain research. *Biol Psychiatry* **75**, 961-969,

doi:10.1016/j.biopsych.2013.07.015 (2014).

13   Barnard, R. A., Pomaville, M. B. & O'Roak, B. J. Mutations and Modeling of the

Chromatin Remodeler CHD8 Define an Emerging Autism Etiology. *Front*

Neurosci **9**, 477, doi:10.3389/fnins.2015.00477 (2015).

14   Vallianatos, C. N. & Iwase, S. Disrupted intricacy of histone H3K4 methylation

in neurodevelopmental disorders. *Epigenomics* **7**, 503-519,

doi:10.2217/epi.15.1 (2015).

15   Zhubi, A., Cook, E. H., Guidotti, A. & Grayson, D. R. Epigenetic mechanisms in

autism spectrum disorder. *Int Rev Neurobiol* **115**, 203-244,

doi:10.1016/B978-0-12-801311-3.00006-8 (2014).

16   Biswas, A. B. & Furniss, F. Cognitive phenotype and psychiatric disorder in

22q11.2 deletion syndrome: A review. *Res Dev Disabil* **53-54**, 242-257,

doi:10.1016/j.ridd.2016.02.010 (2016).

17   Swillen, A. & McDonald-McGinn, D. Developmental trajectories in 22q11.2

deletion. *Am J Med Genet C Semin Med Genet* **169**, 172-181,

doi:10.1002/ajmg.c.31435 (2015).

18   Schneider, M. *et al.* Psychiatric disorders from childhood to adulthood in

22q11.2 deletion syndrome: results from the International Consortium on

Brain and Behavior in 22q11.2 Deletion Syndrome. *Am J Psychiatry* **171**, 627-

639, doi:10.1176/appi.ajp.2013.13070864 (2014).

19   Urban, A. E. *et al.* High-resolution mapping of DNA copy alterations in human

chromosome 22 using high-density tiling oligonucleotide arrays. *Proc Natl*

Acad Sci U S A **103**, 4534-4539, doi:10.1073/pnas.0511340103 (2006).

20   Lieberman-Aiden, E. *et al.* Comprehensive mapping of long-range

interactions reveals folding principles of the human genome. *Science* **326**,

289-293, doi:10.1126/science.1181369 (2009).

21   Yaffe, E. & Tanay, A. Probabilistic modeling of Hi-C contact maps eliminates

systematic biases to characterize global chromosomal architecture. *Nat Genet*

43, 1059-1065, doi:10.1038/ng.947 (2011).

22   Imakaev, M. *et al.* Iterative correction of Hi-C data reveals hallmarks of

chromosome organization. *Nat Methods* **9**, 999-1003,

doi:10.1038/nmeth.2148 (2012).

23   Hu, M. *et al.* HiCNorm: removing biases in Hi-C data via Poisson regression.

Bioinformatics **28**, 3131-3133, doi:10.1093/bioinformatics/bts570 (2012).

24   Schmid, M. W., Grob, S. & Grossniklaus, U. HiCdat: a fast and easy-to-use Hi-C

data analysis tool. *BMC Bioinformatics* **16**, 277, doi:10.1186/s12859-015-

0678-x (2015).

25   Servant, N. *et al.* HiC-Pro: an optimized and flexible pipeline for Hi-C data

processing. *Genome Biol* **16**, 259, doi:10.1186/s13059-015-0831-x (2015).

26   Heinz, S. *et al.* Simple combinations of lineage-determining transcription

factors prime cis-regulatory elements required for macrophage and B cell

identities. *Mol Cell* **38**, 576-589, doi:10.1016/j.molcel.2010.05.004 (2010).

27   Li, W., Gong, K., Li, Q., Alber, F. & Zhou, X. J. Hi-Corrector: a fast, scalable and

memory-efficient package for normalizing large-scale Hi-C data.

Bioinformatics **31**, 960-962, doi:10.1093/bioinformatics/btu747 (2015).

28   Servant, N. *et al.* HiTC: exploration of high-throughput 'C' experiments.

Bioinformatics **28**, 2843-2844, doi:10.1093/bioinformatics/bts521 (2012).

29   Shavit, Y. & Lio', P. Combining a wavelet change point and the Bayes factor for

analysing chromosomal interaction data. *Mol Biosyst* **10**, 1576-1585,

doi:10.1039/c4mb00142g (2014).

30   Sauria, M. E., Phillips-Cremins, J. E., Corces, V. G. & Taylor, J. HiFive: a tool

suite for easy and efficient HiC and 5C data analysis. *Genome Biol* **16**, 237,

doi:10.1186/s13059-015-0806-y (2015).

31   Rao, S. S. *et al.* A 3D map of the human genome at kilobase resolution reveals

principles of chromatin looping. *Cell* **159**, 1665-1680,

doi:10.1016/j.cell.2014.11.021 (2014).

32   Rees, E. *et al.* Evidence that duplications of 22q11.2 protect against

schizophrenia. *Mol Psychiatry* **19**, 37-40, doi:10.1038/mp.2013.156 (2014).

33   Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by

analysis of chromatin interactions. *Nature* **485**, 376-380,

doi:10.1038/nature11082 (2012).

34   Maynard, T. M. *et al.* Mitochondrial localization and function of a subset of

22q11 deletion syndrome candidate genes. *Mol Cell Neurosci* **39**, 439-451,

doi:10.1016/j.mcn.2008.07.027 (2008).

35   Butcher, N. J. *et al.* Association between early-onset Parkinson disease and

22q11.2 deletion syndrome: identification of a novel genetic form of

Parkinson disease and its clinical implications. *JAMA Neurol* **70**, 1359-1366,

doi:10.1001/jamaneurol.2013.3646 (2013).

36   Boot, E. *et al.* Movement disorders and other motor abnormalities in adults

with 22q11.2 deletion syndrome. *Am J Med Genet A* **167A**, 639-645,

doi:10.1002/ajmg.a.36928 (2015).

37   Mok, K. Y. *et al.* Deletions at 22q11.2 in idiopathic Parkinson's disease: a

combined analysis of genome-wide association data. *Lancet Neurol* **15**, 585-

596, doi:10.1016/S1474-4422(16)00071-5 (2016).

38   Stefansson, H. *et al.* Large recurrent microdeletions associated with

schizophrenia. *Nature* **455**, 232-236, doi:10.1038/nature07229 (2008).

39   Consortium, I. S. Rare chromosomal deletions and duplications increase risk

of schizophrenia. *Nature* **455**, 237-241, doi:10.1038/nature07239 (2008).

40   Levinson, D. F. *et al.* Copy number variants in schizophrenia: confirmation of

five previous findings and new evidence for 3q29 microdeletions and VIPR2

duplications. *Am J Psychiatry* **168**, 302-316,

doi:10.1176/appi.ajp.2010.10060876 (2011).

41   Lupiáñez, D. G. *et al.* Disruptions of topological chromatin domains cause

pathogenic rewiring of gene-enhancer interactions. *Cell* **161**, 1012-1025,

doi:10.1016/j.cell.2015.04.004 (2015).

42 Taberlay, P. C. *et al.* Three-dimensional disorganization of the cancer genome

occurs coincident with long-range genetic and epigenetic alterations. *Genome*

Res **26**, 719-731, doi:10.1101/gr.201517.115 (2016).

43 Loviglio, M. N. *et al.* Chromosomal contacts connect loci associated with

autism, BMI and head circumference phenotypes. *Mol Psychiatry*,

doi:10.1038/mp.2016.84 (2016).

44 Kasowski, M. *et al.* Extensive variation in chromatin states across humans.

Science **342**, 750-752, doi:10.1126/science.1242510 (2013).

45 Raha, D., Hong, M. & Snyder, M. ChIP-Seq: a method for global identification

of regulatory elements in the genome. *Curr Protoc Mol Biol* **Chapter 21**, Unit

21.19.21-14, doi:10.1002/0471142727.mb2119s91 (2010).

46 Abyzov, A. *et al.* Somatic copy number mosaicism in human skin revealed by

induced pluripotent stem cells. *Nature* **492**, 438-442,

doi:10.1038/nature11629 (2012).

47 Yao, J., Fetter, R. D., Hu, P., Betzig, E. & Tjian, R. Subnuclear segregation of

genes and core promoter factors in myogenesis. *Genes Dev* **25**, 569-580,

doi:10.1101/gad.2021411 (2011).

48 Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence

of insertions, deletions and gene fusions. *Genome Biol* **14**, R36,

doi:10.1186/gb-2013-14-4-r36 (2013).

49 Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals

unannotated transcripts and isoform switching during cell differentiation.

Nat Biotechnol **28**, 511-515, doi:10.1038/nbt.1621 (2010).

50   Trapnell, C. *et al.* Differential analysis of gene regulation at transcript

resolution with RNA-seq. *Nat Biotechnol* **31**, 46-53, doi:10.1038/nbt.2450

(2013).

51   Huang, d. W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment

tools: paths toward the comprehensive functional analysis of large gene lists.

Nucleic Acids Res **37**, 1-13, doi:10.1093/nar/gkn923 (2009).

52   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-

Wheeler transform. *Bioinformatics* **25**, 1754-1760,

doi:10.1093/bioinformatics/btp324 (2009).

53   Landt, S. G. *et al.* ChIP-seq guidelines and practices of the ENCODE and

modENCODE consortia. *Genome Res* **22**, 1813-1831,

doi:10.1101/gr.136184.111 (2012).

54   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**,

R137, doi:10.1186/gb-2008-9-9-r137 (2008).

55   Stark, R. & Brown, G.

http://bioconductor.org/packages/release/bioc/vignettes/DiffBind/inst/do

c/DiffBind.pdf, 2011.

56   Zhu, L. J. *et al.* ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq

and ChIP-chip data. *BMC Bioinformatics* **11**, 237, doi:10.1186/1471-2105-11-

237 (2010).