Molecular Analysis of Parasites in the Choreocolacaceae (Rhodophyta) Reveals a

Reduced *Harveyella mirabilis* Plastid Genome and Supports the Transfer of Genera

to the Rhodomelaceae (Rhodophyta)

Eric D. Salomaki[2]

Department of Biological Sciences, University of Rhode Island, Kingston, RI, USA.

Gary W. Saunders

Centre for Environmental and Molecular Algal Research, Department of Biology,

University of New Brunswick, Fredericton, NB, E3B 5A3, Canada.

Christopher E. Lane

Department of Biological Sciences, University of Rhode Island, Kingston, RI, USA.

[1]Received.

[2]Author for correspondence: email eric.salomaki@gmail.com.

Running Title: *Alloparasite plastid from Harveyella mirabilis*

Abstract:

Parasitism is a life strategy that has repeatedly evolved within the

Florideophyceae. Until recently, the accepted paradigm of red algal parasite evolution

was that parasites arise by first infecting a close relative and, either through host jumping

or diversification, adapt to infect more distant relatives. The terms adelphoparasite and

alloparasite have been used to distinguish parasites that are closely related to their hosts

from those more distantly related to their hosts, respectively. Phylogenetic studies have

cast doubt on the utility of these terms as data show that even alloparasites predominately

infect with the same family. All adelphoparasites that have been investigated have lost a

native plastid and instead hijack and incorporate a copy of their hosts' plastid when

packaging spores. In contrast, a highly reduced plastid that has lost all genes involved

with photosynthesis was sequenced from the alloparasite *Choreocolax polysiphoniae*,

which indicates that it did not pass through an adelphoparasite stage. In this study we

investigate whether other species in the Choreocolacaceae, a family of alloparasites, also

retains its native plastid, as well as test the hypothesis that alloparasites can arise and

subsequently speciate to form monophyletic clades that infect a range of hosts. We

present the plastid genome for *Harveyella mirabilis* which, similar to that of *C.*

polysiphoniae, has lost genes involved in photosynthesis. The *H. mirabilis* plastid shares

more synteny with free-living red algal plastids than that of *C. polysiphoniae*

Phylogenetic analysis identifies a well-supported monophyletic clade of parasites in the

Choreocolacaceae, which retain their own plastid genomes, within the Rhodomelaceae.

We therefore transfer genera in the Choreocolacaceae to the Rhodomelaceae.

Keywords: Alloparasite, Adelphoparasite, *Choreocolax*, *Leachiella*

INTRODUCTION:

Since the 1845, when *Microcolax botryocarpa* Hooker & Harvey, became the first formally described parasitic red alga, the known biodiversity of red algal parasites has been steadily increasing (Reinsch 1875, Setchell 1914, 1923, Goff 1982, Preuss et al. 2017). Red algal parasites exclusively infect other rhodophytes and are predominately unpigmented, appearing as galls or irregular growths on their free-living red algal hosts. Recent counts have identified 121 distinct species of red algal parasites distributed across eight different orders of the Florideophyceae (Salomaki and Lane 2014, Blouin and Lane 2015, Preuss et al. 2017). Florideophyceae seem to be prone to adopting parasitism, in large part due to their ability to form direct cell-to-cell fusions between adjacent, non daughter cells (Wetherbee and Quirk 1982, Goff and Coleman 1985). With few exceptions [e.g. *Choreonema thuretii* (Broadwater and LaPointe 1997)], red algal parasites leverage their ability to form cell fusions as a means of infecting their host (Goff and Coleman 1985, Zuccarello et al. 2004).

Two main hypotheses have been proposed for the origin of red algal parasites. Setchell (1918) initially proposed that parasites arose as spores from their host, which had mutated to no longer be capable of photosynthesis. Sturch (1926) later proposed that parasites evolved from epiphytes that penetrated and, over time, became nutritionally reliant on the host. Feldmann and Feldmann (1958) later added to Sturch's hypothesis by suggesting that epiphytes that are closely related to their hosts are more likely to succeed in forming secondary pit connections and therefore increase the likelihood of successfully establishing a parasitic relationship.

Despite their diminutive habit, parasitic red algae share morphological

characteristics with other close relatives, and thus were assigned to tribes or families at

the time of their initial discovery (Reinsch 1875, Feldmann and Feldmann 1958).

Historically parasites that have infected close relatives have been considered

'adelphoparasites', while those more distantly related to their hosts are called

'alloparasites' (Feldmann and Feldmann 1958, Goff et al. 1996). Approximately 80% of

the described red algal parasite diversity is considered to be adelphoparasitic, while the

remaining 20% is alloparasitic (Goff 1982, Salomaki and Lane 2014, Blouin and Lane

2015). Based initially on morphological observations, and later coupled with molecular

data, it was proposed that parasites evolve sympatrically with their hosts, as

adelphoparasites, and over time diversify or adapt to infect new and more distantly

related hosts, becoming alloparasites (Feldmann and Feldmann 1958, Goff et al. 1997).

Sturch (1926) initially described the family Choreocolacaceae as a family in the

Gigartinales, which consisted of morphologically reduced parasites lacking chlorophyll

including members of the genera *Choreocolax, Harveyella,* and *Holmsella*. The

Choreocolacaceae was the subject of a thorough phylogenetic analysis of alloparasites to

confirm whether the family was truly a monophyletic clade of parasitic red algae

(Zuccarello et al. 2004). This study supported previous morphological observations

(Fredericq and Hommersand 1990) confirming that the genus *Holmsella* was a member

of the Gracilariaceae and questioned the legitimacy of recognizing a family of red algal

parasites.

In the few other cases where molecular tools have been applied to assess

evolutionary histories of red algal parasites, data suggests that red algal parasites arise

though independent evolutionary events (Goff et al. 1996, 1997, Kurihara et al. 2010). In

addition to phylogenetic analyses, molecular tools have also been applied to investigate

the parasite-host dynamics throughout parasite development. Analyses of the

adelphoparasites *Gardneriella tuberifera* Kylin*, Gracilariophila oryzoides* Setchell & H.

L. Wilson demonstrated that, although the parasites maintain a native mitochondrion,

they have lost their native plastid and instead 'hijack' a host plastid when packaging their

own spores (Goff and Coleman 1995). To date, all red algal parasites examined maintain

a fully functional mitochondrion (Salomaki and Lane 2016). All adelphoparasites that

have been investigated have lost their native plastid (Goff and Coleman 1995; Salomaki

and Lane 2014; Salomaki and Lane unpublished). In contrast, a highly reduced native

plastid was sequenced from the alloparasite *Choreocolax polysiphoniae* Reinsch, which

has lost genes involved in photosynthesis, yet maintains functions including fatty acid

and amino acid biosynthesis (Salomaki et al. 2015). The lack of plastids in

adelphoparasites, in combination with finding a native plastid in the alloparasite *C.*

polysiphoniae, demonstrates that not all parasites pass through an adelphoparasite stage

and that there are multiple paths to parasitism in red algae.

In their study examining relationships in the Choreocolacaceae, Zuccarello *et al*.

(2004) found that *Holmsella pachyderma* and *Holmsella australis* form a monophyletic

clade within the Gracilariaceae. Additionally they identify that parasite genera

Choreocolax, *Harveyella*, and *Leachiella* are members of the Rhodomelaceae, though

have fairly low support for their relationships to each other. Using molecular data we

investigate the relationships of *Choreocolax*, *Harveyella*, and *Leachiella*, and test the

hypothesis that alloparasites can arise and subsequently speciate, forming monophyletic

clades that infect a range of hosts. Furthermore, we set out to determine if another

alloparasite besides *C. polysiphoniae* retains its own native plastid.

MATERIALS AND METHODS

Sample Collection and DNA Extraction

Choreocolax polysiphoniae, found on its host, *Vertebrata lanosa* (Linneaus) T.A.

Christensen, was collected at Beavertail State Park, Jamestown, RI, USA. *Harveyella*

mirabilis (Reinsch) F.Schmitz & Reinke, on its host, *Odonthalia washingtoniensis* Kylin,

was collected from Cattle Point, Friday Harbor, WA, USA (48.452428, -122.962774),

and *Leachiella pacifica* Kugrens, on *Polysiphonia hendryi* N.L. Gardner, was collected

off the dock at Friday Harbor Laboratories, Friday Harbor WA, USA (48.545066, -

123.012268). Individual parasite galls from *Harveyella mirabilis* and *Leachiella pacifica*

were excised from their respective hosts and collected in a 1.5 mL microcentrifuge tube.

The parasite tissue was hand-ground using a Corning Axygen® PES-15-B-SI disposable

tissue grinder pestle in a 1.5 mL microcentrifuge tube while submerged in 100µL of

DNA extraction buffer (Saunders 1993). DNA was extracted from specimens using a

standard phenol/chloroform extraction with all ratios adjusted for an initial buffer volume

of 100µL (Saunders 1993). Additionally, lyophilized DNA from specimen GWS021225

(Saunders 2014) was acquired from the Saunders lab and rehydrated in 20µL of 5mM

Tris/HCl pH 8.5.

Molecular analyses

A 664 bp fragment at the 5-prime end of the mitochondrial cytochrome oxidase 1

gene (COI-5P), which has been used extensively for barcoding red algal species

(Saunders 2005), was PCR amplified using the GWSFn (Le Gall and Saunders 2010) and

GWSRx (Saunders and Moore 2013) primer pair according to protocols in Saunders and

Moore (2013). All PCR reactions were completed with Ex Taq ^{TM} DNA Polymerase

(Takara Bio USA, Mountainview, CA, USA) following the manufacturer's

recommendations. The BigDye Terminator v3.1 Cycle Sequencing Kit (PE Applied

Biosystems [ABI], Foster City, CA, USA) was used for the sequencing of PCR products,

and reactions were analyzed using an ABI Prism 3130XL genetic analyzer. Sequences

were trimmed and assembled using Geneious v. 9.1.5. The consensus sequences resulting

from the assembly were BLAST searched against the NCBI GenBank database and the

Barcode Of Life Database (boldsystems.org) to find similar publicly available sequences.

In addition to PCR amplification of the COI-5P, genomic DNA was amplified

from all specimens using the illustra Single Cell GenomiPhi DNA Amplification kit (GE

Healthcare Life Sciences, Pittsburgh, Pa) according to manufacturer protocols. Whole-

genome amplification of an individual *C. polysiphoniae* parasite was completed using the

illustra GenomiPhi Hy DNA Amplification Kit according to manufacturer protocols. The

COI-5P was PCR amplified and sequenced, as previously described, from the whole-

genome amplified DNA to screen for contamination resulting from the amplification

process. Libraries for Illumina sequencing were constructed from the amplified DNA on

the Apollo 324 robot using the PrepX ILM DNA Library Kit (Wafergen Biosystems,

Freemont, CA, USA). The *Harveyella mirabilis, Leachiella pacifica,* and GWS021225

libraries were multiplexed and sequenced on an Illumina MiSeq paired-end 250 x 250

basepair run. The *C. polysiphoniae* library was sequenced on separate full-cell Illumina

MiSeq paired-end 300 x 300 basepair run. The sequencing effort resulted in 29,355,470

paired-end reads for *C. polysiphoniae,* 7,923,094 paired end reads for *Harveyella*

mirabilis, 7,502,360 paired-end reads for *Leachiella pacifica*, and 8,869,740 paired end

reads for GWS021225. For all datasets, sequences with PHRED scores <30 were

removed and the remaining reads were trimmed of adapter sequences. Additionally,

fifteen 5' and five 3' nucleotides were trimmed from the remaining reads and all reads

under 100 nucleotides were removed from the dataset. All trimming was completed using

CLC Genomics Workbench v. 9.5.2 (CLC Bio-Qiagen, Aarhus, Denmark) and the

remaining reads were assembled using default parameters in CLC Genomics Workbench

v. 9.5.2.

Phylogenetic Analysis

A total of 823 Rhodomelaceae COI-5P sequences available on GenBank were

downloaded into Geneious v. 9.1.5 {FormattingCitation} and aligned with the data

generate by PCR for *Harveyella mirabilis* and *Leachiella pacifica.* This alignment was

subjected to maximum likelihood analysis using RAxML with the GTR+I+G model with

1,000 bootstrap replicates. Based upon the results of the preliminary tree, 18S, 28S, COI-

5P, and *rbc*L data were compiled from GenBank for thirteen of the closest relatives to a

weakly supported clade containing *Choreocolax polysiphoniae* Reinsch, *Harveyella*

mirabilis, Leachiella pacifica, and GWS021225 (Table 1). The longest sequence for each

gene from this compiled dataset was used as a query against local BLAST databases

created from the *C. polysiphoniae, H. mirabilis, L. pacifica,* and GWS021225 assemblies,

(Table 1). The 18S and 28S regions were extracted from the ribosomal cistrons identified

in each assembly based upon its alignment to the query gene.

182   Each dataset was aligned using he MAFFT plugin in Geneious v. 9.1.5 and

subsequently concatenated using Sequence Matrix v. 1.7.8 {FormattingCitation}. The

concatenated dataset was then subjected to phylogenetic analysis using maximum

likelihood, implemented in RAxML v. 8.2.2 {FormattingCitation} and Bayesian

Inference using MrBayes v. 3.2.2 {FormattingCitation}. For the maximum likelihood

analysis the GTR+I+G model was used and bootstrap support values were calculated

using 1,000 bootstrap replicates. For the Bayesian inference analysis, two Metropolis-

coupled Markov chain Monte Carlo (MCMCMC) runs consisting of one cold chain and

three hot chains were preformed. Each run was sampled every 100 generations for

1,010,000 generations. After confirming the runs converged by checking to ensure that

the average standard deviation of split frequencies was below 0.01, the trees were

merged. The resulting tree and posterior probabilities were calculated from the 20,202

trees generated.

Plastid Genome Annotation

197   A 90,654 bp contig was identified as the plastid genome of *Harveyella mirabilis*

from the previously described assembled Illumina MiSeq data. Open reading frame

(ORF) prediction on the *H. mirabilis* plastid was done in Geneious Pro v6.1 and the

resulting ORFs were manually annotated using GenBank and Pfam (Finn et al. 2010,

2015) databases. Functional annotations were assigned from the UniProt (The UniProt

Consortium 2017) and KEGG databases (Kanehisa et al. 2016). Genes found in red algal

plastid genomes that were missing from the *H. mirabilis* plastid were searched for using

BLAST, against the plastid sequence and the genomic assemblies to verify their absence

and check for evidence of transfer to another genetic compartment. The plastid genome

sequence was submitted to the tRNAscan-SE online server v1.21 (Schattner et al. 2005)

for identification of tRNA sequences and to MFannot

(http://megasun.bch.umontreal.ca/cgi-bin/mfannot/mfannotInterface.pl) to identify rRNA

sequences and confirm manual gene annotations.

Plastid Genome Comparative Analysis

The circular plastid genomes of the free-living florideophytes, *Calliarthron*

tuberculosum (KC153978), *Ceramium cimbricum* (KR025491), *Ceramium japonicum*

(KX284719), *Chondrus crispus* (HF562234), *Dasya binghamiae* (KX247284), *Laurencia*

sp. JFC0032 (LN833431), *Grateloupia taiwanensis* (KC894740), *Gracilaria*

tenuistipitata (AY673996), *Vertebrata lanosa* (KP308097), and the alloparasites

Choreocolax polysiphoniae (KP308096) and *Harveyella mirabilis* (XXXXXXXX) were

arranged so their sequences began with the *fts*H gene. Whole genome alignment was

completed using the default settings for the progessiveMauve algorithm in the Mauve

v2.3.1 (Darling et al. 2004).

RESULTS

Identification of a Cryptic Parasite

A maximum likelihood phylogenetic analysis of 823 Rhodomelaceae COI-5P

sequences was completed using RAxML. This analysis provided weak support (bootstrap

support of 23) for a clade containing the alloparasites *Choreocolax polysiphoniae,*

Harveyella mirabilis, Leachiella pacifica, as well as 2 sequences labeled *Rhodomela*

sp1Cal voucher GWS021225 (KM254767) and *Rhodomela* sp1Cal voucher GWS021347

(KM254267) (Supplemental Figure 1). Interestingly, these two GenBank sequences had

bootstrap support of 100 as being sister to the sequence of *Leachiella pacifica* generated

in this study (data not shown). Based upon these results, the specimens were reexamined

and parasite galls were found on the host, which was subsequently identified as

Polysiphonia paniculata based upon *rbc*L sequence data.

Monophyletic Alloparasite Clade

Although identification of parasites on the samples GWS021225 and

GWS021347 was a direct result of resolving a clade of parasite sequences, bootstrap

support for a monophyletic clade of alloparasites was weak and a subsequent Bayesian

phylogeny failed to recover the same clade (data not shown). Based upon this original

analysis, COI-5P data from a reduced dataset of the parasites and 28 of their closest

relatives was subjected to maximum likelihood analysis, which again recovered a poorly

supported monophyletic alloparasite clade (Supplemental Figure 2). To resolve the issue

of low statistical support, thirteen taxa that were continually allied to the alloparasite

clade were selected based upon availability of published comparative data. Phylogenetic

analysis was conducted on a concatenated dataset comprised of 18S, 28S, COI-5P and

*rbc*L (Table 1) from the 13 free-living species, and the alloparasites *C. polysiphoniae, H.*

mirabilis, L. pacifica, and the parasite on GWS021225 (from here on called *Leachiella*

sp.). With the additional data beyond the COI-5P, both maximum likelihood and

Bayesian inference recovered a monophyletic clade of red algal alloparasites with 75%

bootstrap support and 0.99 posterior probability (Figure 1). *Choreocolax polysiphoniae*

was the earliest branching parasite followed by *Harveyella mirabilis*, with *Leachiella*

pacifica and *Leachiella* sp. being recovered as sister taxa with 100% bootstrap

support/1.0 posterior probability (Figure 1).

To exclude long-branch attraction as a contributor to the monophyly of the clade

of alloparasites, maximum likelihood analyses were conducted individually on the 18S

and 28S gene trees, as well as the concatenated dataset with both species of *Leachiella*

(the longest branches) removed. All analyses recovered a monophyletic clade of

alloparasites, however the branching patterns differed between them. In the 18S gene

tree, the clade had 82% bootstrap support, however *H. mirabilis* was the earliest

branching parasite followed by *L. pacifica, L.* sp., and *C. polysiphoniae*. The 28S gene

tree recovered a monophyletic parasite clade with the same branching pattern observed in

Figure 1, but with only 65% bootstrap support. The concatenated analysis with the two

species of *Leachiella* removed also recovered *C. polysiphoniae* and *H. mirabilis* as sister

species with 70% bootstrap support.

Taxonomic Considerations

Choreocolax polysiphoniae was initially described by Reinsch (1875) from the

Atlantic coast of North America as parasitic on *Polysiphonia fastigiata* (now *Vertebrata*

lanosa). Specimens used the phylogenetic analysis presented here (Figure 1) were

collected at Beavertail State Park in Jamestown, RI, USA infecting *Vertebrata lanosa* and

are a strong match to the type description (Reinsch 1875). As the type collection cannot

be located, we formally lectotypify *C. polysiphoniae* on image #49 accompanying the

description in Reinsch (1875). Based upon the molecular analyses here, which resolved a

monophyletic clade of parasitic red algae within the Rhodomelaceae (Figure 1), we

formally transfer *Choreocolax* to the Rhodomelaceae and recognize Choreocolacaceae as

a synonym of this family. To adhere to the principle of monophyly, the genus *Harveyella*,

based on the type and only species *H. mirabilis* and included in our analyses (Figure 1), is

also transferred to the Rhodomelaceae.

Harveyella mirabilis Plastid Genome

282   The plastid genome of *Harveyella mirabilis* was assembled as a 90,654 kb

circular molecule with 322x coverage. The plastid genome has an overall AT content of

76.5% and contains 84 protein coding genes, 3 rRNAs, and 23 tRNAs (Figure 2). Similar

to the *Choreocolax polysiphoniae* plastid (Salomaki et al. 2015), all genes related to

photosynthesis have been lost with the exception of *pet*F which has been demonstrated to

be involved in electron transport in other metabolic pathways (Happe and Naber 1993,

Jacobs et al. 2009). Genes involved in transcription/translation and fatty acid, amino acid,

protein, isoprene biosynthesis remain conserved. As in the *C. polysiphoniae* plastid, *glt*B

appears to be a vanishing pseudogene. BLAST similarity searches are able to find

conserved homology, however the presence of stop-codons throughout the region

suggests that the gene is likely no longer capable of being completely translated.

Plastid Genome Comparisons

A whole genome MAUVE alignment of the *H. mirabilis* with *C. polysiphoniae* and nine

representative free-living Florideophyceae plastid genomes identified 13 locally collinear

blocks in the *H. mirabilis* genome that aligned with the free-living plastids (Figure 3).

There were no rearrangements or inversions in the *H. mirabilis* plastid genome when

compared to photosynthetic Rhodomelaceae taxa. When aligning *H. mirabilis* to the *C.*

polysiphoniae plastid genome, 11 locally collinear blocks are identified, and several

genome inversions and rearrangements are evident (Figure 4). Additionally, gene content

varies slightly between the two parasite plastid genomes. *Harveyella mirabilis* retains

*arg*B, *car*A, *inf*B, *rnz*, *rpl*9, *rpl*17, *rpl*24, *rpl*32, *rpl*33, *rpl*34, *rpo*Z, *rps*18, *rps*20, *ycf*21,

which have all been lost in *C. polysiphoniae* (Table 2), while *C. polysiphoniae* maintains

copies of *dna*B and *fab*H which have both been lost in the *H. mirabilis* plastid genome.

DISCUSSION

Origin of Parasites

With 121 described parasites occurring across eight Florideophyeceae orders

(Salomaki and Lane 2014, Blouin and Lane 2015, Preuss et al. 2017), red algae appear

more able to transition from autotrophy to parasitic lifestyles than any other eukaryotic

lineage. The terms adelphoparasite and alloparasite have traditionally been used to

describe parasites that infect hosts within their tribe/family, or in different tribes/families,

respectively. The use of these terms has been questioned as molecular data have revealed

that alloparasites, like adelphoparasites, infect close relatives rather than distantly related

species (Zuccarello et al. 2004, Kurihara et al. 2010). Phylogenetic analysis using data

presented here place *Choreocolax polysiphoniae*, *Harveyella mirabilis*, and two species

of *Leachiella* firmly within the same family as their hosts further supporting

abandonment of these terms for differentiating two types of red algal parasites.

The Demise of Exclusively Parasitic Families

Sturch (1926) initially described the Choreocolacaceae as a family of

holoparasites, containing the genera *Choreocolax, Harveyella,* and *Holmsella*. However,

more recent morphological investigation considered that the genus *Holmsella* was related

to the parasites *Gelidiocolax* and *Pterocladiophila*, and it was moved to the family

Pterocladiophilaceae in the Gracilariales (Fredericq and Hommersand 1990). Their

observations were subsequently supported by molecular data generated with the specific

aim of testing the phylogenetic affinities of parasites that Sturch had assigned to the

Choreocolacaceae. This work demonstrated that *Holmsella australis* and *Holmsella*

pachyderma formed a well supported monophyletic clade within the Gracilariaceae

(Zuccarello et al. 2004). Their molecular data also indicated that *Choreocolax* and

Harveyella belong in the Ceramiales, leading the authors to question whether the

Choreocolacaceae should continue to be recognized (Zuccarello et al. 2004). However,

their use of 18S sequence data was insufficient to resolve the issue of a monophyletic

clade for these parasites. Other authors have also noted that species in *Choreocolax,*

Harveyella, and *Leachiella* have features aligning them to the Ceramiales, but again,

taxonomic affinities among the parasite species and within this order remained uncertain

(Goff and Cole 1975, Kugrens 1982, Fredericq and Hommersand 1990). Our data

confirm the findings of Zuccarello, Moon, and Goff (2004), placing *Choreocolax,*

Harveyella and *Leachiella*, within the Rhodomelaceae. Furthermore, the phylogeny

utilizing additional molecular markers provides strong support for a monophyletic clade

containing *Choreocolax*, *Harveyella* and *Leachiella* (Figure 1), and supports the placing

the Choreocolacaceae in synonymy with the Rhodomelaceae.

Cryptic Species

Until recently, the dogma behind red algal parasite evolution was the notion that

parasites arise sympatrically as adelphoparasites and over time evolve and adapt to infect

more distant hosts becoming alloparasites (Goff et al. 1996, 1997). Based on molecular

data demonstrating that alloparasites also infect members of their own family (Zuccarello

et al. 2004, Kurihara et al. 2010), we now consider that these terms are not suitable for

distinguishing parasites. The recognition of multiple monophyletic parasite clades, the

Holmsella clade (Zuccarello et al. 2004), and the Rhodomelaceae clade uncovered here

containing *Choreocolax, Harveyella,* and *Leachiella* (Figure 1), supports the idea of a

sympatric origin of alloparasites with subsequent speciation as parasites adapt to their

hosts, without passing through an adelphoparasite-like stage.

Previously, *Leachiella pacifica* was described from multiple hosts including

members of the genus *Polysiphonia* and *Pterosiphonia* (Kugrens 1982, Zuccarello and

West 1994). In their study on *L. pacifica* host specificity, Zuccarello and West (1994)

found that parasite spores isolated from one host genus were unable to infect members of

the other host genus. Kugrens initially designated the *Leachiella pacifica* type specimen

as a parasite infecting *Polysiphonia* spp. from Cattle Point, Friday Harbor, WA, USA

(Kugrens 1982). Based upon this, here we are considering the specimen collected on

Polysiphonia hendryi from Friday Harbor, WA, USA as true *Leachiella pacifica*.

Subsequently, parasites infecting *Polysiphonia paniculata* (and several other species)

have also been attributed to *L. pacifica* (Zuccarello and West 1994, Zuccarello et al.

2004). Molecular data here identify two distinct species of *Leachiella* that infect separate

hosts but are otherwise difficult to distinguish. Due to the highly reduced morphology of

these parasites, it seems likely that host identity may be the easiest way to identify

parasites in the genus *Leachiella* to species.

Some parasitic red algae are reportedly capable of infecting a range of hosts [e.g.,

Asterocolax – see Goff, Ashen, and Moon (1997)], however molecular data are

contradicting that notion. Rather, it appears that red algal parasite species have higher

host specificity than previously believed, and instead, we have underestimated the

amount of parasite diversity as a result of their reduced morphology. Morphology based

studies report *Leachiella pacifica* infecting at least seven unique host species (Zuccarello

and West 1994), further molecular analyses will surely uncover additional species in the

genus *Leachiella*.

Parasite Plastids

The link between plastid origin in red algal parasites and their evolutionary

relationship to their hosts may be central to our understanding of red algal parasite

evolution. The *Harveyella mirabilis* plastid genome (Figure 2) represents the second red

algal parasite demonstrated to retain a reduced native plastid. Similar to *Choreocolax*

polysiphoniae, the *H. mirabilis* plastid genome remains conserved for functions including

amino acid, fatty acid, and protein biosynthesis, but has lost genes involved in building

the light harvesting apparatus, photosystems I and II, and other photosynthesis related

genes. Identifying a monophyletic clade of parasites (Figure 1) in which the two earliest

branching members identified so far retain plastids, strongly suggests that plastids are

also retained in species of *Leachiella*. Preliminary data generated in the Lane lab supports

that hypothesis, though work remains to completely sequence plastids from *Leachiella*

species.

The *H. mirabilis* plastid gene order, with the exception of the missing

photosynthesis genes, is conserved when compared to plastids of free-living

Rhodomelaceae species (Figure 3). In contrast, we find that the *C. polysiphoniae* plastid

has undergone greater gene loss than the *H. mirabilis* plastid, and a substantial amount of

genome reorganization (Figure 4). *Harveyella mirabilis* retains 13 genes that have been

lost from the *C. polysiphoniae* plastid (Table 2); *arg*B and *car*A, which are involved in

arginine biosynthesis processes, *rpo*Z, which promotes RNA polymerase assembly, nine

ribosomal protein genes, and an uncharacterized hypothetical protein. In contrast, *C.*

polysiphoniae retains copies of *dna*B, which is involved in DNA replication, and *fab*H,

which is involved in fatty acid biosynthesis, both of which have been lost in *H. mirabilis*.

Analysis of additional plastid genomes in this clade will provide greater insights into

patterns of plastid genome evolution in red algal parasites.

Two hypotheses have been proposed for the origin of red algal parasites. Setchell

(1918) suggested that a mutation in a spore causes parasites to arise sympatrically, while

Sturch (1926) postulated that parasitic red algae start out as epiphytes that over time

become endophytes and increasingly rely on the host for nutrition. It seems plausible that

Setchell's origin hypothesis could explain the rise of the so-called adelphoparasites,

which comprise the majority of known red algal parasite biodiversity. By evolving from

their hosts, adelphoparasites could more easily incorporate a copy of a genetically similar

plastid as their own. Replacing a functionally reduced or non-functional native plastid

with a host-derived plastid seems to be an easy mechanism for survival. Unfortunately,

reliance on the host plastid also starts the newly evolved parasite down a path towards

inevitable extinction. In order for a non-photosynthetic parasite to survive it still must

retain compatibility for other plastid functions including amino acid and fatty acid

biosynthesis. Because the host-derived plastid is newly acquired during each new

infection, the host plastid experiences one set of evolutionary pressures while the parasite

evolves and accumulates mutations of its own. Eventually the parasite will inevitably lose

the ability to communicate with the host plastid as the parasite and host increasingly

become genetically distinct. This leaves the parasite with two possibilities for survival,

either to find another host with a compatible plastid or go extinct. Alternatively, the

success of an alloparasite may be explained by Sturch's hypothesis. By evolving from a

closely related epiphyte that is able to create secondary pit connections, the parasite may

retain its own plastid and therefore enable its longevity and the opportunity to even

speciate as the parasite adapts to new hosts. Therefore, what were previously viewed as

competing hypotheses to explain the evolution of red algal parasites, may each explain

how different types of parasites arise.

Although organisms have transitioned from photosynthesis to other methods of

nutrient acquisition numerous times across the tree of life, support for total plastid loss is

rare (Gornik et al. 2015) with most cases resulting in highly reduced plastids retained for

core functions (e.g. isoprenoid and fatty acid biosynthesis) (Wolfe et al. 1992, Knauf and

Hachtel 2002, de Koning and Keeling 2006) or in extreme cases the retention of the

apicoplast in Apicomplexans . The non-photosynthetic red algal parasites traditionally

considered adelphoparasites, including *Gracilariophila oryzoides* and *Gardneriella*

tuberifera, still require a copy of the host plastid for functions that, to date, remain

undetermined. Interestingly, in the *Gracilariophila oryzoides* genome and transcriptome,

photosynthesis related nuclear-encoded plastid-targeted genes remain conserved while

those same genes are absent from the *Choreocolax polysiphoniae* transcriptome

(Salomaki and Lane, unpublished).

Data presented here further highlight the need to abandon the notion that

taxonomy defines red algal parasites. By placing the family Choreocolacaceae in

synonymy with the Rhodomelaceae we are making steps to remove the artificial

appearance of parasites infecting hosts in different families. Furthermore, we recognize

that a major distinction between types of red algal parasites is the origin of the parasite

plastids. While some parasites, including *Harveyella mirabilis* and *Choreocolax*

polysiphoniae, retain a native plastid that evolves in concert with the parasite, others

including *Gracilariophila oryzoides* and *Gardneriella tuberifera*, incorporate a copy of a

host plastid when packaging spores, but will inevitably evolve and become incompatible

with the host plastid and blink out of existence. We predict that investigations of parasites

that have traditionally referred to as alloparasites, like *Holmsella* and *Pterocladiophila,*

will also provide evidence of plastid retention and monophyletic clades of parasites.

ACKNOWLEDGEMENTS

The authors thank Amanda Savoie for help identifying parasite sequence data in the UNB

database, Jillian Freese for collecting specimens of *Harveyella mirabilis* and *Leachiella*

pacifica, and Kristina Terpis for her support in the lab. Funding for this work was

provided to by a Phycological Society of America Grant-in-Aid of Research to ES and by

grant #1257472 from the National Science Foundation awarded to CL. This research is

based in part upon work conducted using the Rhode Island Genomics and Sequencing

Center which is supported in part by the National Science Foundation (MRI Grant No.

DBI-0215393 and EPSCoR Grant Nos. 0554548 & EPS-1004057), the US Department of

Agriculture (Grant Nos. 2002-34438-12688 and 2003-34438-13111), and the University

of Rhode Island.

LITERATURE CITED

Blouin, N. & Lane, C. 2015. Red algae provide fertile ground for exploring parasite

    evolution. *Perspect. Phycol.* 3:11–9.

Broadwater, S.T. & Lapointe, E.A. 1997. Parasitic interactions and vegetative

    ultrastructure of Choreonema thuretii (Corallinales, Rhodophyta). *J. Phycol.*

    33:396–407.

Darling, A.C.E., Mau, B., Blattner, F.R. & Perna, N.T. 2004. Mauve : Multiple alignment

    of conserved genomic sequence with rearrangements. *Genome Res.* 14:1394–403.

de Koning, A.P. & Keeling, P.J. 2006. The complete plastid genome sequence of the

    parasitic green alga Helicosporidium sp. is highly reduced and structured. *BMC Biol.*

    4.

Feldmann, G. & Feldmann, J. 1958. Recherches sur quelques Floridees parasites. *Rev.*

478 *Gen. Bot.* 65:49–128.

479 Finn, R.D., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Mistry, J., Mitchell, A.L., Potter,

480   S.C. et al. 2015. The Pfam protein families database: towards a more sustainable

481   future. *Nucleic Acids Res.* 44:D279–85.

482 Finn, R.D., Mistry, J., Tate, J., Coggill, P., Heger, A., Pollington, J.E., Gavin, O.L. et al.

483   2010. The Pfam protein families database. *Nucleic Acids Res.* 38:D211-22.

484 Goff, L.J. 1982. The biology of parasitic red algae. *Prog. Phycol. Res.* 1:289–369.

485 Goff, L.J., Ashen, J. & Moon, D. 1997. The evolution of parasites from their hosts: A

486   case study in the parasitic red alga. *Evolution.* 51:1068–78.

487 Goff, L.J. & Cole, K. 1975. The biology of Harveyella mirabilis (Cryptonemiales,

488   Rhodophyceae). II . Carposporophyte development as related to the taxonomic

489   affiliation of the parasitic red alga, Harveyella mirabilis. *Phycologia.* 14:227–38.

490 Goff, L.J. & Coleman, A.W. 1985. The role of secondary pit connections in red algal

491   parasitism. *J. Phycol.* 21:483–508.

492 Goff, L.J. & Coleman, A.W. 1995. Fate of parasite and host organelle DNA during

493   cellular transformation of red algae by their parasites. *Plant Cell.* 7:1899–911.

494 Goff, L.J., Moon, D.A., Nyvall, P., Stache, B., Mangin, K. & Zuccarello, G. 1996. The

495   evolution of parasitism in the red algae: Molecular comparisons of adelphoparasites

496   and their hosts. *J. Phycol.* 32:297–312.

497 Gornik, S.G., Febrimarsa, Cassin, A.M., MacRae, J.I., Ramaprasad, A., Rchiad, Z.,

498   McConville, M.J. et al. 2015. Endosymbiosis undone by stepwise elimination of the

499   plastid in a parasitic dinoflagellate. *Proc. Natl. Acad. Sci.* 112:201423400.

500 Happe, T. & Naber, J.D. 1993. Isolation, characterization and N-terminal amino acid

sequence of hydrogenase from the green alga Chlamydomonas reinhardtii. *Eur. J.*

Biochem. 214:475–81.

Jacobs, J., Pudollek, S., Hemschemeier, A. & Happe, T. 2009. A novel, anaerobically

induced ferredoxin in Chlamydomonas reinhardtii. *FEBS Lett.* 583:325–9.

Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M. & Tanabe, M. 2016. KEGG as a

reference resource for gene and protein annotation. *Nucleic Acids Res.* 44:D457–62.

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S.

et al. 2012. Geneious Basic: an integrated and extendable desktop software platform

for the organization and analysis of sequence data. *Bioinformatics*. 28:1647–9.

Knauf, U. & Hachtel, W. 2002. The genes encoding subunits of ATP synthase are

conserved in the reduced plastid genome of the heterotrophic alga Prototheca

wickerhamii. *Mol. Genet. Genomics*. 267:492–7.

Kugrens, P. 1982. Leachiella pacifica, Gen. et Sp. nov., a new parasitic red alga rom

Washington and California. *Am. J. Bot.* 123:306–19.

Kurihara, A., Abe, T., Tani, M. & Sherwood, A.R. 2010. Molecular phylogeny and

evolution of red algal parasites: A case study of Benzaitenia, Janczewskia, and

Ululania (Ceramiales). *J. Phycol.* 46:580–90.

Le Gall, L. & Saunders, G.W. 2010. DNA barcoding is a powerful tool to uncover algal

diversity: A case study of the phyllophoraceae (Gigartinales, Rhodophyta) in the

Canadian flora1. *J. Phycol.* 46:374–89.

Preuss, M., Nelson, W.A. & Zuccarello, G.C. 2017. Red algal parasites: a synopsis of

described species, their hosts, distinguishing characters and areas for continued

research. *Bot. Mar.* 60:1–13.

Reinsch, P.F. 1875. Contributiones ad Algologiam et Fungologiam. Lipsiae, Nuremberg.

61 pp.

Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget,

B. et al. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model

choice across a large model space. *Syst. Biol.* 61:539–42.

Salomaki, E.D. & Lane, C.E. 2014. Are all red algal parasites cut from the same cloth?

Acta Soc. Bot. Pol. 83:369–75.

Salomaki, E.D. & Lane, C.E. 2016. Red Algal Mitochondrial Genomes are More

Complete than Previously Reported. *Genome Biol. Evol.* 48:evw267.

Salomaki, E.D., Nickles, K.R. & Lane, C.E. 2015. The ghost plastid of Choroeocolax

polysiphoniae. *J. Phycol.* 51:217–21.

Saunders, G.W. 1993. Gel purification of red algal genomic DNA: An inexpensive and

rapid method for the isolation of polymerase chain reaction-friendly DNA. *J.*

Phycol. 29:251–4.

Saunders, G.W. 2005. Applying DNA barcoding to red macroalgae: a preliminary

appraisal holds promise for future applications. *Philos. Trans. R. Soc. Lond. B. Biol.*

Sci. 360:1879–88.

Saunders, G.W. 2014. Long distance kelp rafting impacts seaweed biogeography in the

Northeast Pacific: the kelp conveyor hypothesis. *J. Phycol.* 50:968–74.

Saunders, G.W. & Moore, T.E. 2013. Refinements for the amplification and sequencing

of red algal DNA barcode and RedToL phylogenetic markers: a summary of current

primers, profiles and strategies. *Algae.* 28:31–43.

Schattner, P., Brooks, A.N. & Lowe, T.M. 2005. The tRNAscan-SE, snoscan and

snoGPS web servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res.*

33:686–9.

Setchell, W.A. 1914. Parasitic Florideae I. *Univ. Calif. Publ. Bot.* 6:1–34.

Setchell, W.A. 1918. Parasitism Among the Red Algae. *Proc. Am. Philisophical Soc.*

57:155–72.

Setchell, W.A. 1923. Parasitic Florideae II. *Univ. Calif. Publ. Bot.* 10:393–401.

Stamatakis, A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics.* 30:1312–3.

Sturch, H.H. 1926. Choreocolax Polysiphoniae, Reinsch. *Ann. Bot.* XL.

The UniProt Consortium. 2017. UniProt: the Universal Protein knowledgebase. *Nucleic*

Acids Res. 45:D158–69.

Vaidya, G., Lohman, D.J. & Meier, R. 2011. SequenceMatrix: Concatenation software

for the fast assembly of multi-gene datasets with character set and codon

information. *Cladistics.* 27:171–80.

Wetherbee, R. & Quirk, H. 1982. The Fine Structure and Cytology of the Association

Between the Parasitic Red Alga Holmsella australis and Its Red Algal Host

Gracilaria furcellata. *Protoplasma.* 165:153–65.

Wolfe, K.H., Morden, C.W. & Palmer, J.D. 1992. Function and evolution of a minimal

plastid genome from a nonphotosynthetic parasitic plant. *Proc. Natl. Acad. Sci. U. S.*

A. 89:10648–52.

Zuccarello, G.C., Moon, D. & Goff, L.J. 2004. A phylogenetic study of parasitic genera

placed in the family Choreocolacaceae (Rhodophyta). *J. Phycol.* 40:937–45.

Zuccarello, G.C. & West, J.A. 1994. Genus and race specificity in the red algal parasite

Leachiella pacifica (Choreocolacaceae, Rhodophyta). *Phycologia*. 33:213–8.

Table 1. GenBank Sequence accession numbers for taxa used in phylogenetic analyses
(Figure 2).
575

	18S	28S	COI-5P	*rbc*L
Amansia fimbrifolia	HM582913	HQ422079	HM582889	-
Amansia glomerata	HM582909	HQ422225	HQ422913	-
Bostrychia arbuscula	-	AY920894	KM502795	AY920845
Bostrychia intricata	-	KM502850	KM502801	AY920846
Bostrychia vaga	-	KM502842	KM502791	KM502817
Choreocolax polysiphoniae	This Study	This Study	This Study	-
Harveyella mirabilis	This Study	This Study	This Study	-
Heterodasya mucronata	-	-	KC567673	KF367797
Leachiella pacifica	This Study	This Study	This Study	-
Leachiella sp. GWS021225	This Study	This Study	KM254767	-
Neorhodomela larix	AY617140	-	KM254241	GQ252553
Neorhodomela oregona	-	-	KM254304	GQ252556
Odonthalia dentata	-	JX572172	JX571960	KU564463
Odonthalia floccosa	AY617141	-	KM254276	GQ252492
Osmundaria obtusiloba	-	HQ422003	HQ422818	-
Rhodomela confervoides	AY617145	KX145642	KX258842	KX146197
Rhodomela lycopodioides	-	This Study	HM916516	KU564489

576
577

578   Table 2. *Harveyella mirabilis* plastid genes and their function, which have been lost from
579   the *Choreocolax polysiphoniae* plastid genome.
580

Gene Name	Biological Process and Molecular Function
*arg*B	Amino acid biosynthesis; ATP binding, acetylglutamate kinase activity
*car*A	Pyrimidine metabolism; ATP binding, carbamoyl-phosphate synthase (glutamine-hydrolyzing) activity
*inf*B	Translation; Translation initiation, GTP binding
*rpl*9	Translation; structural constituent of the 50S ribosome
*rpl*17	Translation; structural constituent of the 50S ribosome
*rpl*24	Translation; structural constituent of the 50S ribosome
*rpl*32	Translation; structural constituent of the 50S ribosome
*rpl*33	Translation; structural constituent of the 50S ribosome
*rpl*34	Translation; structural constituent of the 50S ribosome
*rpo*Z	Transcription; DNA binding
*rps*18	Translation; structural constituent of the 30S ribosome
*rps*20	Translation; structural constituent of the 30S ribosome
*ycf*21	Uncharacterized hypothetical protein

Figure 1. Maximum likelihood phylogeny based on concatenated 18S, 28S, COI-5P, and rbcL data as shown

in Table 1, of the parasites *Choreocolax polysiphoniae, Harveyella mirabilis, Leachiella pacifica*, an undescribed

species of *Leachiella* found on *Polysiphonia paniculata*, and thirteen close relatives. This phylogenetic analysis

supports a monophyletic clade of parasites arising within the Rhodomelaceae. Support values shown as Bayesian

posterior probability/maximum likelihood bootstrap.

Figure 2. The plastid genome of the parasitic red alga, *Harveyella mirabilis* is 90,654 basepairs and contains

84 protein coding genes (Green), the 5S, 16S, and 23S rRNAs (Red), and 23 tRNAs (Pink). All genes involved

with photosynthetic functions, except *pet*F, have been lost. The *fts*H gene is truncated but may still be transcribed,

however *glt*B is a non-functional pseudogene (Yellow).

Figure 3. Mauve alignment of the parasites *Choreocolax polysiphoniae* (top) and *Harveyella mirabilis* (second from top) with all published Rhodomelaceae plastid genomes as well as those from select representatives of other Florideophyceae families (*Dasya binghamiae*, Dasyaceae; *Chondrus crispus*, Gigartinaceae; *Calliarthron tuberculosum*, Corallinaceae; *Grateloupia taiwanensis*, Halymeniaceae; *Gracilaria tenuistipitata*, Gracilariaceae). This alignment identifies 16 locally collinear blocks (LCBs) among the selected plastid genomes and demonstrates that even with the loss of photosynthesis genes overall synteny is shared between the parasite *H. mirabilis* and other plastid genomes, while *C. polysiphoniae* has undergone several genome rearrangements.

Figure 4. Mauve alignment of the plastid genomes from the parasites *Harveyella mirabilis* (top) and

Choreocolax polysiphoniae (bottom). This alignment identifies 11 locally collinear blocks (LCBs) among

these parasite plastid genomes, highlighting the high level of genome fragmentation and rearrangements

evident in the two parasite plastids