Functional redundancy of variant and canonical histone

H3 lysine 9 modification in *Drosophila*

Taylor J.R. Penke[*], Daniel J. McKay[*,†,**,††], Brian D. Strahl[*,‡,§,]

A. Gregory Matera[*,†,§,**,††], and Robert J. Duronio[*,†,§,**,††,1]

[*]Curriculum in Genetics and Molecular Biology

[†]Integrative Program for Biological and Genome Sciences

[‡]Department of Biochemistry and Biophysics

[§]Lineberger Comprehensive Cancer Center

[**]Department of Genetics,

[††]Department of Biology,

The University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599

[1]To whom correspondence should be addressed

Robert J. Duronio

Integrative Program for Biological and Genome Sciences, CB#7100

University of North Carolina

Chapel Hill, NC 27599

Voice: (919) 962-4568

FAX: (919) 962-4574

Email: duronio@med.unc.edu

ABSTRACT

Histone post-translational modifications (PTMs) and differential incorporation of variant and canonical histones into chromatin are central modes of epigenetic regulation. Despite similar protein sequences, histone variants are enriched for different suites of PTMs compared to their canonical counterparts. For example, variant histone H3.3 occurs primarily in transcribed regions and is enriched for "active" histone PTMs like Lys9 acetylation (H3.3K9ac), whereas the canonical histone H3 is enriched for Lys9 methylation (H3K9me), which is found in transcriptionally silent heterochromatin. To determine the functions of K9 modification on variant versus canonical H3, we compared the phenotypes caused by engineering $H3.3^{K9R}$ and $H3^{K9R}$ mutant genotypes in *Drosophila melanogaster*. Whereas most $H3.3^{K9R}$ and a small number of $H3^{K9R}$ mutant animals are capable of completing development and do not have substantially altered protein coding transcriptomes, all $H3.3^{K9R}$ $H3^{K9R}$ combined mutants die soon after embryogenesis and display decreased expression of genes enriched for K9ac. These data suggest that the role of K9ac in gene activation during development can be provided by either H3 or H3.3. Conversely, we found that H3.3K9 is methylated at telomeric transposons, and this mark contributes to repressive chromatin architecture, supporting a role for H3.3 in heterochromatin that is distinct from that of H3. Thus, our genetic and molecular analyses demonstrate that K9 modification of variant and canonical H3 have overlapping roles in development and transcriptional regulation, though to differing extents in euchromatin and heterochromatin.

INTRODUCTION

DNA interacts with histones and other proteins to establish chromatin environments that

affect all DNA-dependent processes. The establishment of chromatin environments is

accomplished through multiple mechanisms that collectively comprise the bulk of

epigenetic regulation found in eukaryotes. In particular, post-translational modification

(PTM) of histones influences DNA/histone interactions and also provides binding sites

for recruitment of chromatin modulators that influence gene expression, DNA replication

and repair, and chromosome segregation during cell division (Wallrath et al. 2014). In

addition to histone PTMs, epigenetic regulation is modulated by the type of histone

protein deposited onto DNA. There are two major categories of histone proteins: the

canonical histones and the closely related histone variants (Talbert and Henikoff 2010,

2017). These two histone categories are distinguished by the timing of their expression

during the cell cycle and their mechanism of deposition onto DNA. Canonical histones

are encoded by multiple genes (e.g., ~55 in humans and ~500 in flies), organized into

clusters that are highly expressed during S-phase of the cell cycle, and are deposited

onto DNA by the histone chaperone CAF-1 in a replication-coupled manner (Marzluff et

al. 2002; Tagami et al. 2004; Verreault et al. 1996). In contrast, variant histones are

typically encoded by one or two genes, are expressed throughout the cell cycle, and

can be deposited onto DNA independently of replication by histone chaperones other

than CAF-1 (Henikoff and Ahmad 2005; Tagami et al. 2004; Szenker et al. 2011).

Variant histones are often deposited at specific genomic locations and have functions

that can differ from canonical histones. For example, two histone H2A variants, H2AX

and H2A.Z, play critical roles in DNA repair (Scully and Xie 2013; Price and Andrea

2014), and the histone H3 variant CENP-A localizes to centromeres and is essential for

kinetochore formation (Blower and Karpen 2001; Henikoff and Ahmad 2005; Mellone

and Allshire 2003).

The major histone H3 variant in animal genomes is H3.3, which in both mice and

Drosophila is encoded by two different genes (*H3.3A* and *H3.3B*) that produce identical

proteins. Variant histone H3.3 differs from canonical H3.2 and H3.1 by only four or five

amino acids, respectively (Szenker et al. 2011). In each case, three of these different

amino acids are located in the globular domain of H3.3 and are necessary and sufficient

for interaction with the replication-independent chaperones HIRA and ATRX-DAXX

(Tagami et al. 2004; Goldberg et al. 2010; Ahmad and Henikoff 2002; Lewis et al.

2010). In H3.2, the only replication-dependent histone in *Drosophila*, the fourth amino

acid difference occurs at position 31 in the unstructured N-terminal tail (Szenker et al.

2011). Histones H3.2 and H3.1 (collectively hereafter referred to as H3) along with H3.3

are some of the most conserved proteins in all eukaryotes (Malik and Henikoff 2003).

The conservation of amino acid differences between H3 and H3.3 during evolution

strongly suggests that these proteins perform distinct functions. Indeed, H3.3 and H3

are deposited in different genomic regions in a variety of species (Mito et al. 2005;

Schwartz and Ahmad 2005; Tamura et al. 2009; Jin et al. 2011; Kraushaar et al. 2013;

Allis and Wiggins 1984). H3.3 is also enriched for different histone PTMs than H3 (Hake

et al. 2006; McKittrick et al. 2004), and H3.3 containing nucleosomes can be less stable

than those with H3 (Jin and Felsenfeld 2007; Xu et al. 2010). Although the epigenetic

PTM signature on variant and canonical H3 histones is distinct, the degree to which

particular histone PTMs found on both H3 and H3.3 can compensate for one another is

not fully understood. Here, we explore the common and distinct functions of variant and

canonical H3K9 function during *Drosophila* development.

H3.3 is associated with transcriptionally active regions of the genome with high

nucleosome turnover, consistent with H3.3 being enriched in "activating" histone PTMs

and depleted in "repressing" histone PTMs (Hake et al. 2006; McKittrick et al. 2004).

One of the histone PTMs enriched on H3.3 relative to H3 is acetylation of lysine nine

(K9ac), a mark associated with accessible chromatin (Hake et al. 2006; McKittrick et al.

2004). Previous studies have identified K9ac at promoters of genes and in regions of

high transcriptional activity (Kharchenko et al. 2011; Bernstein et al. 2005; Liang et al.

2004; Roh et al. 2005). Additionally, mutation of H3K9 acetyltransferases results in

compromised transcriptional output, suggesting K9ac contributes to or is a

consequence of gene expression activation (Wang et al. 1998; Georgakopoulos and

Thireos 1992; Kuo et al. 1998). Importantly, H3K9 acetyltransferases target other

histone residues and have non-histone substrates as well (Glozak et al. 2005; Spange

et al. 2009), indicating that one cannot deduce the function of K9ac solely by mutation

of H3K9 acetyltransferases. For example, whereas mutation of the H3K9

acetyltransferase Rtt109 in budding yeast results in sensitivity to DNA-damaging

agents, H3K9R mutants, which cannot be acetylated by Rtt109, are insensitive to DNA-

damaging agents (Fillingham et al. 2008). Direct investigation of K9ac function in vivo

therefore requires mutation of H3K9 itself. Previously, we used a *Drosophila* histone

gene replacement platform (McKay et al. 2015) to generate a canonical $H3^{K9R}$ mutant,

and found no significant changes in gene expression at regions of the genome enriched

in K9ac (Penke et al. 2016). This observation raises the possibility that H3.3K9ac

functions in gene regulation and can compensate for the absence of H3K9ac.

H3.3 is also found at transcriptionally inactive, heterochromatic regions of the

genome (Goldberg et al. 2010; Lewis et al. 2010; Wong et al. 2010). Heterochromatin is

enriched in H3K9 di- and tri-methylation (me2/me3), modifications that recruit

Heterochromatin Protein 1 (HP1) and are essential for heterochromatin function

(Bannister et al. 2001; Lachner et al. 2001; Nakayama et al. 2001; Penke et al. 2016).

DNA within heterochromatin is composed of repeated sequence elements, many of

which are transcriptionally silent and consist of immobile transposons or transposon

remnants. Using H3.3 mutants it was recently demonstrated that H3.3 is essential for

repression of endogenous retroviral elements and that H3.3 can be methylated at lysine

nine (Elsässer et al. 2015). H3.3K9me3 is also important for heterochromatin formation

at mouse telomeres (Udugama et al. 2015). These studies did not assess the

contribution of canonical H3K9 because strategies for mutating all replication-dependent

H3 genes in mammalian cells have not been developed. We recently showed in

Drosophila that mutation of canonical H3K9 causes defects in heterochromatin

formation and transposon repression (Penke et al. 2016), similar to phenotypes

observed in *C. elegans* in the absence of H3K9 methyltransferases (Zeller et al. 2016).

In addition, we detected low levels of K9me2/me3 in $H3^{K9R}$ mutants. Combined, these

data suggest methylated H3.3K9 plays a role in heterochromatin formation and can

compensate for the absence of canonical H3K9. However, the extent of functional

overlap between variant and canonical H3K9 and the intriguing possibility that identical

modifications on variant or canonical histones have distinct functions has yet to be fully

investigated.

In order to better understand the functions of H3 and H3.3 and to compare the

functions of the variant and canonical H3K9 residues, we used CRISPR-Cas9 to

generate a variant K9R substitution mutation ($H3.3^{K9R}$) in *Drosophila* and combined this

with our previously described canonical $H3^{K9R}$ mutant (Penke et al. 2016). By comparing

the individual mutant phenotypes of $H3.3^{K9R}$ and $H3^{K9R}$ to the combined $H3.3^{K9R}$ $H3^{K9R}$

mutants using a variety of genomic and cell biological assays, we demonstrate that

variant and canonical versions of H3K9 can compensate for each other, although to

substantially different extents in euchromatin versus heterochromatin. H3K9 plays a

more substantial role than H3.3K9 in heterochromatin formation and in the repression of

transposons, whereas they compensate for each other in controlling euchromatic gene

expression, particularly in regions enriched in the activating modification, K9ac.

MATERIALS AND METHODS

Generation of K9R mutant genotypes: Variant $H3.3^{K9R}$ mutants generated by the

cross scheme illustrated in Supplementary Figure 1A were selected by the absence of

GFP fluorescence and/or the presence of straight wings. 1st instar larvae from the

variant and canonical $H3.3^{K9R}$ $H3.2^{K9R}$ cross described in Supplementary Figure 1B

were selected based on the presence of GFP fluorescence. Only larvae that receive the

H3^{HWT} or H3^{K9R} transgene will survive embryogenesis, as this transgene provides the

only source of canonical histone genes. In Table 2, rows one and two indicate progeny

from the cross *yw ; H3.3^{2x1} / CyO, twiGFP x yw ; Df(2L)BSC110 / CyO, twiGFP*. Rows

three and four indicate progeny from the cross *H3.3B^{K9R} ; H3.3^{2x1} / CyO, twiGFP x*

H3.3B^{K9R} ; Df(2L)BSC110 / CyO, twiGFP. In these crosses, the expected ratio of

heterozygous to homozygous *H3.3A^{2x1}* animals is 2:1, as *CyO, twiGFP/CyO, twiGFP*

animals do not eclose as adults.

CRISPR-Cas9 Mutagenesis and Transgene Integration: A single gRNA targeting

H3.3B near the K9 residue was inserted into pCFD3 and co-injected with a 2 kb

homologous repair template containing the H3.3BK9R substitution (Supplementary

Experimental Methods). Constructs were injected into embryos expressing Cas9 from

the *nanos* promoter (*nanos-cas9*; Kondo and Ueda 2013). Recovered *H3.3B^{K9R}* alleles

were subsequently crossed into *H3.3A* null backgrounds (*H3.3A^{2x1}* over deficiency

Df(2L)BSC110). Independent *H3.3B^{K9R}* CRISPR alleles were used to generate trans-

heterozygous animals for all experiments. To generate *H3.3B* rescue constructs, a 5 kb

genomic sequence containing the entire wild-type *H3.3B* transcription unit was PCR

amplified from genomic DNA of *nanos-cas9* flies and cloned into pATTB

(Supplementary Experimental Methods). Gibson assembly (Gibson et al. 2009) using

primers containing K9R or K9Q substitutions was used to generate mutated versions of

H3.3B, and all three constructs were integrated into the 86FB *attP* landing site by

ΦC31-mediated recombination.

Immunofluorescence: Salivary gland preparations stained using anti-H3K9me2, anti-

H3K9me3, anti-H3K9ac, or anti-HP1a were performed as previously described (Cai et

al. 2010). Antibody sources and concentrations are included in Supplementary

Experimental Methods. 1st instar larval brains were prepared similar to imaginal wing

disc preparations described in Estella et al. (2008).

Western Blots: ImageJ densitometry analysis was used to determine K9me2, K9ac, or

H3 band intensity (See Supplementary Experimental Methods). Histone modification

signal was normalized to corresponding H3 loading control signal. Normalized signal

from different titrations of the same genotype were averaged and consequent values

were set relative to WT value. This process was completed for two biological replicates

for both K9me2 and K9ac.

Sample Preparation and Sequence Data Analysis: FAIRE-seq and RNA-seq

samples were prepared from wandering 3rd instar imaginal wing discs as previously

described (McKay and Lieb 2013). Sequencing reads were aligned to the dm6 (6.04)

reference genome using Bowtie2 (FAIRE) and Tophat (RNA) default parameters

(Langmead and Salzberg 2012; Trapnell et al. 2014). FAIRE peaks were called with

MACS2 using a shift size of 110bp and a stringency cutoff of 0.01 (Zhang et al. 2008).

Transcripts were assembled with Cufflinks (Trapnell et al. 2014). Bedtools was used to

determine read coverage at peaks and transcripts (Quinlan and Hall 2010) and DESeq2

was used to determine statistical significance ($p < 0.05$) (Love et al. 2014). The following

modENCODE 3rd instar larval ChIP-seq data sets were used: K9me2=GSE47260, and

K9me3=GSE47258. K9ac ChIP-seq data from imaginal wings discs was generated by

Pérez-Lluch et al. (GSM1363590, 2015).

Chromatin state analysis was performed using data from Kharchenko et al

(2010), which assigns small regions of the genome into one of nine different chromatin

state. FAIRE peaks were classified as one or more chromatin states based on overlap

with regions defined by Kharchenko et al. (2010). Of all the peaks in a particular

chromatin state, we determined the percentage of peaks that had significantly different

FAIRE signal in mutant compared to WT samples. RNA chromatin state analysis was

performed in a similar fashion.

See Supplemental Experimental Procedures for a detailed description of the

methods. Strains are available upon request. Sequencing data are available at GEO

under accession number GSE106192.

RESULTS

H3.3^{K9R} mutant animals are viable but sterile

In order to investigate the role of H3.3K9 in *Drosophila* development and compare it to

the role of H3K9, we first generated an H3.3^{K9R} animal by introducing a K9R substitution

at the endogenous *H3.3B* locus using CRISPR/Cas9 and then combining recovered

H3.3B^{K9R} mutant alleles with a previously generated *H3.3A* null allele (H3.3A/B

combined genotype denoted hereafter as *H3.3^{K9R}*; see Table 1, Table S1, and Figure

S1A for histone genotype nomenclature) (Sakai et al. 2009). These *H3.3^{K9R}* mutants,

which contain the full complement of endogenous canonical *H3* genes, eclose as adults

at the expected Mendelian ratios (Table 2) and appear morphologically normal.

Therefore, canonical H3 can provide all of the H3K9 function during *Drosophila*

development. This result is consistent with a previous study finding that flies without any

H3.3 protein could be propagated as a stock if canonical H3.2 was expressed from a

transgene using the *H3.3B* promoter (Hödl and Basler 2012). Our results are also in

line with a previous report in which *H3.3A* and *H3.3B* null animals containing an

H3.3A^{K9R} transgene were viable (Sakai et al. 2009). However, whereas these *H3.3A^{K9R}*

transgenic animals were fertile (Sakai et al. 2009), we found that animals with an

endogenous *H3.3B^(K9R)* mutation and the same *H3.3A* null allele used by Sakai et al.

(2009) were sterile. The sterility of our *H3.3^(K9R)* animals was rescued in both males and

females by a transgene containing the wild-type *H3.3B* gene ectopically integrated into

the genome, suggesting that the relative abundance of H3.3^(K9R) causes sterility (Figure

S2). We conclude that H3.3K9 plays an essential role during gametogenesis and

speculate that different amounts of H3.3^(K9R) histones from *H3.3A* or *H3.3B* promoters

may account for the differences between our observations and those of Sakai et al.

(2009).

262 H3.3K9 and H3K9 have overlapping functions during development

We previously observed that canonical *H3^(K9R)* mutants could complete development,

although 98% of these mutant animals died during larval or pupal stages (Penke et al.

2016). We considered the possibility that *H3^(K9R)* mutant animals progressed to late larval

or pupal stages of development because of compensation by H3.3K9. We therefore

tested if the *H3.3^(K9R)* genotype would advance the *H3^(K9R)* mutant stage of lethality by

observing the development of animals in which the *H3.3^(K9R)* and *H3^(K9R)* mutant

genotypes were combined (Table 1, Table S1, and Figure S1B). The *H3^(K9R)* genotype

was generated using our previously described histone replacement platform (McKay et

al. 2015; Penke et al. 2016). Briefly, the endogenous array of ~100 canonical histone

gene clusters was deleted and replaced with an ectopically located transgene encoding

a BAC-based, tandem array of 12 canonical histone gene clusters in which the *H3*

genes contain a K9R mutation (Figure S1B). A 12x tandem array of wild-type, canonical

histone genes (denoted <u>h</u>istone <u>w</u>ild <u>t</u>ype or *H3^(HWT)*, Figure S1B), which fully rescues

deletion of the endogenous histone gene array, was used as a control (McKay et al.

2015). Similar to the *H3.3^{K9R}* mutants, *H3^{HWT}* animals with the *H3.3^{K9R}* mutant genotype

(denoted hereafter as *H3.3^{K9R} H3^{HWT}*; see Table 1 and Figure S1B) were viable (Table

3). However, only 34.6% of *H3.3^{K9R} H3^{HWT}* progeny eclosed as adults (Table 3)

compared to essentially 100% of the *H3.3^{K9R}* genotype that contained the full

complement of endogenous, wild-type *H3* genes (Table 2). This result suggests that in

the presence of fewer total canonical *H3* gene copies, the *H3.3^{K9R}* mutation is more

detrimental. Importantly, animals with the *H3.3^{K9R} H3^{K9R}* combined mutant genotype

containing both the variant and canonical K9R mutation were 100% inviable, dying with

high penetrance at the 1^{st} instar larval stage, much earlier than the majority of *H3^{K9R}*

mutants. These results demonstrate that H3.3K9 can partially compensate for the

absence of H3K9, indicating that H3.3K9 and H3K9 have some redundant functions.

H3K9 PTMs are lost in animals lacking H3.3K9 and H3K9

We previously found that K9me2/me3 signal in *H3^{K9R}* mutant animals is substantially

reduced but not absent. Thus, a possible reason why *H3.3^{K9R} H3^{K9R}* mutants have a

more severe developmental defect than *H3^{K9R}* mutants is complete loss of K9me

throughout the genome. We therefore assessed K9me2/me3 levels in *H3.3^{K9R}* and

H3.3^{K9R} H3^{K9R} mutants by immunofluorescence. We first assessed K9me2/me3 levels in

salivary gland polytene chromosomes of *H3.3^{K9R}* mutants, with the expectation that if

H3.3K9 is methylated the signal will be reduced relative to controls. The salivary gland

is a highly polyploid tissue (>1000C) and the alignment of chromatids in the polytene

chromosomes results in easily visible structures that provide information about levels

and genomic locations of histone PTMs using immunofluorescence. $H3.3^{K9R}$ mutants

had lower levels of both K9me2 and K9me3 compared to wild-type controls at the

largely heterochromatic chromocenter, demonstrating that H3.3K9 is

normally methylated in the pericentric heterochromatin of otherwise wild-type animals

(Figure 1A, B). In support of this result, western blot analysis of salivary glands

demonstrated that K9me2 levels were decreased in $H3.3^{K9R}$ mutants compared to wild-

type controls (Figure 1D, E).

Because $H3.3^{K9R}$ mutants exhibited reduced K9me2/me3 signal at the

chromocenter, we next used immunofluorescence to examine localization of HP1a,

which binds K9me2/me3. In line with reduced K9me2/me3 signal, HP1a signal at the

chromocenter of $H3.3^{K9R}$ mutants was reduced compared to wild-type controls

(Figure1A, C). HP1a and H3.3 also both localize to telomeres (Goldberg et al. 2010;

Lewis et al. 2010). We found that HP1a localizes to telomeres in $H3.3^{K9R}$ mutants

(Figure 1A), as it does in $H3^{K9R}$ mutants (Penke et al. 2016). These results are

consistent with previous observations that HP1 recruitment to telomeres requires

telomere binding proteins (Raffa et al. 2011; Vedelek et al. 2015; Badugu et al. 2003)

and not the H3K9 methyltransferase Su(var)3-9 (Perrini et al. 2004), suggesting that

H3K9me is not required for HP1 recruitment to telomeres.

Because $H3.3^{K9R}$ $H3^{K9R}$ combined mutants do not develop to the 3[rd] instar larval

stage, we examined K9me2 levels in 1[st] instar larval brains. $H3^{K9R}$ mutants (with wild-

type variant histones) and $H3.3^{K9R}$ $H3^{HWT}$ mutants (with a 12x transgenic complement of

wild-type canonical histone genes) each exhibited reduced K9me2 levels by

immunofluorescence compared to $H3^{HWT}$ controls, consistent with the polytene

chromosome data (Figure 2A). In contrast, the *H3.3^{K9R} H3^{K9R}* variant and canonical

combined mutant brains had undetectable levels of K9me2 in the vast majority of cells

(Figure 2A). These results provide further evidence that H3.3K9 is methylated and that

the total amount of K9me is derived from both H3.3 and H3.

Interestingly, a small number of cells in the *H3.3^{K9R} H3^{K9R}* 1st instar mutant brains

retained low levels of K9me2 signal at the chromocenter (arrowheads, Figure 2). Cells

with residual K9me2 express ELAV, a pan-neuronal marker, and lack expression of

Deadpan and Prospero, markers of proliferating neuroblasts and ganglion mother cells,

respectively (circles, Figure S3). These data indicate that cells with K9me2 positive

chromocenters in *H3.3^{K9R} H3^{K9R}* mutant 1st instar larval brains are differentiated

neurons. We suspect that the K9me2 signal in these cells reflects maternally provided

wild-type H3 protein remaining in the genomes of quiescent neurons that differentiated

prior to having their maternal H3 fully replaced by zygotically expressed H3K9R mutant

histones. A corollary to this conclusion is that the proliferating neuroblasts and their

GMC daughters likely have progressed through a sufficient number of S phases such

that replacement of maternal H3 with zygotic H3K9R eliminates detectable K9me2

signal.

We also found that levels of H3K9 acetylation were reduced in both the *H3.3^{K9R}*

mutant and the *H3^{K9R}* mutant relative to controls, as determined both by

immunofluorescence of salivary gland polytene chromosomes (Figure 3A, B) and by

western blots of salivary gland extracts (Figure 3C). Because a substantial amount of

K9ac is placed on H3.3, we considered the possibility that lack of K9ac was responsible

for the fertility defects of *H3.3^{K9R}* mutants and the early lethality of *H3.3^{K9R} H3^{K9R}*

mutants. To address this question, we integrated either an H3.3B^{K9}, an H3.3B^{K9R}, or an

H3.3B^{K9Q} transgene into the same genomic position in order to determine if a K9Q

acetyl mimic could restore fertility to *H3.3^{K9R}* mutants. Animals with only an *H3.3B^{K9R}*

mutation at the endogenous locus (i.e., containing a wild-type *H3.3A* gene), and

carrying either an H3.3B^{K9R} or H3.3B^{K9Q} transgene were sterile, precluding us from

constructing the genotype to test if these transgenes could rescue the sterility of *H3.3^{K9R}*

mutant adults (Figure S2). This result suggests that both the H3.3B^{K9R} and H3.3B^{K9Q}

transgenes acted dominantly to compromise fertility. Furthermore, these data imply that

H3.3B^{K9R} and H3.3B^{K9Q} histones are incorporated into chromatin.

H3.3K9 regulates chromatin organization at the chromocenter, telomeres, and

transposons

We next asked if the reduction of K9me2/me3 in *H3.3^{K9R}* mutants affected chromatin

organization by cytological examination of salivary gland polytene chromosomes using

DAPI staining of DNA. As we found previously in *H3^{K9R}* mutants (Penke et al. 2016), in

some *H3.3^{K9R}* mutants polytene chromosome spreads the chromocenter appeared

abnormal and not fully condensed (Figure 1F). The cause of this phenotype is unclear

but may reflect altered chromatin organization or defects in the under-replication of

salivary gland pericentric heterochromatin (Belyaeva et al. 1998; Zhimulev et al. 2003).

Based on their cytology, we binned chromocenters into three categories: "organized",

"moderately organized", and "disorganized" (Figure 1F). We categorized chromocenters

from four genotypes: wild-type (WT; i.e., with the endogenous canonical histone genes),

an *H3.3A* null mutant (*H3.3A^{Null}*), an *H3.3B* K9R substitution mutant (*H3.3B^{K9R}*), and the

H3.3B^{K9R} ; H3.3A^{Null} double mutant in which all H3.3 contains the K9R substitution

(*H3.3^{K9R}*) (Table 1, Table S1, and Figure S1A). Whereas the majority of wild-type

chromocenters were organized (60% organized vs 40% moderately organized), both the

H3.3B^{K9R} and the *H3.3A^{Null}* single mutants had increased percentages of moderately

organized and disorganized chromocenters (Figure 1F). For example, ~22% of

chromocenters in the various *H3.3* mutants were disorganized compared to less than

1% of wild-type chromocenters. These results indicate that H3.3 contributes to

chromocenter structure. Interestingly, the *H3.3B^{K9R}* ; *H3.3A^{Null}* double mutant had the

same proportion of moderately organized and disorganized chromocenters as either

single mutant. This result suggests that either reducing *H3.3* gene dose (i.e., the

H3.3A^{Null} allele) or expressing K9R mutant H3.3 histones (i.e., the *H3.3B^{K9R}* mutation),

can prevent normal H3.3 function at pericentric heterochromatin.

Given the disrupted chromocenter structure in *H3.3^{K9R}* mutants, we next

examined chromatin structure genome wide by performing Formaldehyde Assisted

Isolation of Regulatory Elements followed by whole genome sequencing (FAIRE-seq).

FAIRE-seq provides a measure of local nucleosome occupancy across the genome,

revealing regions of "open" chromatin that are relatively depleted of nucleosomes

(Simon et al. 2013). We previously found using this technique that regions of

heterochromatin enriched in K9me, particularly pericentromeric heterochromatin, were

more open in canonical *H3^{K9R}* mutants relative to *H3^{HWT}* controls (Penke et al. 2016). To

determine if variant *H3.3^{K9R}* mutants had a similar phenotype we performed FAIRE-seq

in triplicate on imaginal wing discs from wandering 3rd instar larvae in *WT, H3.3A^{Null}*,

H3.3B^{K9R}, and *H3.3B^{K9R}* ; *H3.3A^{Null} (H3.3^{K9R})* double mutant genotypes. Sequencing

reads were aligned to the genome and peaks were called on each of the three

replicates and combined into a merged peak set. Called peaks were consistent across

replicates and read coverage across peaks was highly correlated (R$\geq$ 0.96) (Figure

S4A, B). Additionally, wild-type FAIRE data was consistent with previously generated

data from wing discs (McKay and Lieb 2013) (Figure S4D). *H3.3A^{Null}, H3.3B^{K9R}, and*

H3.3^{K9R} mutants each had a similar percentage of peaks with significantly altered FAIRE

signal when compared to wild-type: 8.8%, 6.5%, and 7.9% respectively (Figures 4A-C).

Moreover, significantly changed peaks across the three mutants exhibited a high degree

of overlap. Of the 2,660 significantly changed peaks across all mutants, 21% were

shared among all three and 52% by at least two mutants (Figure S5A). FAIRE signal at

significantly changed peaks also displayed similar fold changes in mutants compared to

wild-type and were not exacerbated in the double mutant compared to either single

mutant (Figure S5B). These data suggest H3.3A and H3.3BK9 both function to regulate

chromatin architecture.

We next asked if the changes in FAIRE signal we observed in *H3.3* mutants were

characterized by a particular chromatin signature. We assigned each called FAIRE peak

to one of nine different chromatin states characterized by different combinations of

histone PTMs as defined by Kharchenko et al. (2010). We then calculated the

percentage of FAIRE peaks that changed between an *H3.3* mutant and wild-type within

each chromatin state. Regions of K9me2/me3 showed the highest percentage of

changes in FAIRE signal in *H3.3A^{Null}*, *H3.3B^{K9R}*, and the *H3.3^{K9R}* mutant compared to

wild-type, supporting the idea that H3.3K9 is methylated and plays a necessary role in

regulating chromatin architecture (Figure 4D). Changes in FAIRE signal were also more

likely to occur in regions of H3K36me3, a mark that is enriched along gene bodies that

are themselves enriched for H3.3 (Bannister et al. 2005; Szenker et al. 2011). Finally,

we used modENCODE K9me2 and K9me3 ChIP-seq data to complement the chromatin

state analysis. Of the FAIRE peaks significantly increased or decreased in $H3.3^{K9R}$

mutants compared to wild-type, 76.4% and 49.0% respectively overlapped a K9me2 or

K9me3 peak (Figure 4F). These results demonstrate that altered FAIRE signal in

$H3.3^{K9R}$ mutants occurred in regions normally occupied by K9me.

We also observed increased FAIRE signal at telomeres in all three $H3.3$ mutant

genotypes, particularly on chromosomes 2R and 3L (Figure S5C), suggesting that H3.3

regulates telomeric chromatin architecture. In *Drosophila,* telomeres are composed of

retrotransposons enriched in K9me2/me3 (Levis et al. 1993; Cenci et al. 2005). H3.3

plays a similar role in the mouse, in which H3.3 null mutant embryonic stem cells exhibit

an increase in transcripts from transposons (Elsässer et al. 2015) and telomeres

(Udugama et al. 2015). Additionally, we previously observed transposon activation and

mobilization in canonical $H3^{K9R}$ mutants (Penke et al. 2016). For these reasons, we

examined FAIRE signal at transposons in our $H3.3$ mutants using the piPipes pipeline,

which avoids ambiguity in aligning reads to repetitive transposons by mapping to

transposon families (Han et al. 2015). Both $H3.3A^{Null}$ and $H3.3B^{K9R}$ mutants resulted in

significantly increased FAIRE signal at transposons, and $H3.3^{K9R}$ mutants had on

average even higher increased FAIRE signal at transposons (Figure 5A, B). Moreover,

FAIRE signal at some telomeric transposons, particularly TART-B, was increased in

$H3.3$ mutants (Figure 5C). However, the extent of increase in $H3.3^{K9R}$ mutants was not

as severe as previously observed for $H3^{K9R}$ mutants (Penke et al. 2016) (Figure 5B).

These results support a role for H3.3K9 in chromatin-mediated transposon repression,

though to a lesser extent than H3K9.

H3.3K9 and H3K9 functions overlap in regions of K9ac and partially in regions of

K9me

To investigate the cause of lethality when both variant and canonical H3 histones

contain the K9R mutation, we performed RNA-seq of 1st instar larvae from four

genotypes: *H3^{HWT}*, *H3^{K9R}*, *H3.3^{K9R} H3^{HWT}*, and *H3.3^{K9R} H3^{K9R}* (Table 1, Table S1).

Larvae of the correct genotype were identified by GFP fluorescence (see Materials and

Methods). RNA sequencing reads were aligned to the genome using Tophat, transcript

assembly was performed by Cufflinks, and DESeq2 was used for statistical analysis

(Trapnell et al. 2014; Love et al. 2015). Each genotype was verified by examination of

RNA-seq reads mapping to the K9 codon of variant and canonical histones. Correlation

analysis demonstrated transcript abundance across all assembled transcripts was

highly similar among replicates, and was also similar to previously generated data from

wild-type 1st instar larvae (Figure S6) (Graveley et al. 2011). Additionally, histone

expression was similar across all genotypes, suggesting that variation in histone levels

do not underlie observed phenotypes (Figure S7A). In line with our previous analysis of

H3^{K9R} RNA-seq data from imaginal wing discs (Penke et al. 2016), the majority of

significantly changed transcripts in *H3^{K9R}* 1st instar samples was increased compared to

H3^{HWT} (247 increased vs 41 decreased), supporting a role for H3K9me in gene silencing

(Figure 6A). *H3.3^{K9R} H3^{HWT}* samples had a similar number of significantly changed

transcripts, and again most transcripts showed increased signal compared to *H3^{HWT}*

(203 vs 126), though fold changes were smaller than $H3^{K9R}$ mutants (Figure 6B). By

contrast, the $H3.3^{K9R} H3^{K9R}$ combined mutant genotype caused a much more

pronounced effect on gene expression compared to either the $H3.3^{K9R} H3^{HWT}$ or the

$H3^{K9R}$ mutant genotypes (Figure 6C); 869 transcripts exhibited increased RNA signal

and 1036 transcripts were decreased compared to $H3^{HWT}$ samples. The number of

decreased transcripts in $H3.3^{K9R} H3^{K9R}$ animals compared to $H3^{HWT}$ was therefore about

ten-fold higher than either the variant or canonical K9R mutant alone. Thus, similar to

our viability analysis (Table 3), these RNA-seq results demonstrated that variant and

canonical versions of H3K9 compensate for each other in the regulation of gene

expression.

Because we observed increases in FAIRE signal at transposons in $H3.3^{K9R}$

mutants from wing disc samples, we examined RNA levels of transposon families in 1st

instar larvae. Similar to our previous RNA-seq observations from $H3^{K9R}$ mutant wing

discs (Penke et al. 2016), RNA signal at transposons in $H3^{K9R}$ 1st instar larvae was

increased relative to the $H3^{HWT}$ control (Figure S7B, C). Although on average

transposon levels were only slightly higher in $H3.3^{K9R} H3^{HWT}$ mutants compared to

$H3^{HWT}$, transposon levels in $H3.3^{K9R} H3^{K9R}$ combined mutants were significantly higher

than either $H3.3^{K9R} H3^{HWT}$ or $H3^{K9R}$ mutants alone (Figure S7B, C). Moreover, telomeric

transposons are generally increased in all K9R mutants compared to $H3^{HWT}$ controls

(Figure S7D). Together these results support an overlapping role for H3.3K9 and H3K9

in regulating gene expression and transposon repression.

We next examined chromatin signatures of significantly altered transcripts to

explore the mechanism of the observed gene expression changes. All transcripts were

assigned to one or more chromatin states based on their overlap with genomic regions

defined by Kharchenko et al. (2010). We then determined the percentage of transcripts

within a given chromatin state that were either increased or decreased in K9R mutants

relative to $H3^{HWT}$ controls (Figure 7 A-C). Transcripts in regions of K9me2/me3

(chromatin state 7 and 8) were the most likely to have significantly increased RNA

levels in mutants compared to $H3^{HWT}$. Although $H3.3^{K9R}$ $H3^{K9R}$ combined mutants had

the highest percentage of chromatin state 7 transcripts that were significantly increased

(~26%), $H3^{K9R}$ mutants also displayed a high percentage (~13%) of change within

chromatin state 7 (Figure 7A, D, Figure S8A). These results suggest that H3.3K9

contributes to gene repression in regions of K9me2/me3 but cannot completely

compensate for the absence of H3K9.

In contrast to upregulated transcripts, very few transcripts were significantly

decreased in $H3.3^{K9R}$ $H3^{HWT}$ or $H3^{K9R}$ mutants. However, the $H3.3^{K9R}$ $H3^{K9R}$ combined

mutant displayed numerous significant decreases in transcript abundance. Interestingly,

transcripts in chromatin state 1, characterized by K9ac and lack of K9me, were most

likely to be decreased (Figure 7B, E). Several other chromatin states showed elevated

transcript changes, particularly in the $H3.3^{K9R}$ $H3^{K9R}$ combined mutant; however, in this

analysis transcripts can be assigned to more than one chromatin state. Indeed, many

transcripts in chromatin state 1 also overlap other chromatin states. We therefore

performed a supplementary analysis that examined only transcripts that overlap a single

chromatin state. This analysis demonstrated that transcripts solely in chromatin state 1

were much more likely to change in K9R mutants than those in other chromatin states

(Figure S8A). Similar results were obtained using imaginal wing disc K9ac ChIP data

from Pérez-Lluch et al. (2015). Whereas few transcripts that overlapped K9ac were

significantly altered in either single mutant (68 in $H3^{K9R}$ and 116 in $H3.3^{K9R}\ H3^{HWT}$),

1195 K9ac associated transcripts exhibited changed expression levels in $H3.3^{K9R}\ H3^{K9R}$

combined mutants (Figure 8). These data suggest that in regions of K9ac, H3.3 and H3

can completely compensate for each other. Additionally, these data provide evidence

that K9ac facilitates gene expression.

DISCUSSION

Overlapping and distinct developmental functions of H3 and H3.3

In this study, we determined the distinct and overlapping roles that lysine 9 of variant

and canonical histone H3 play in gene expression and heterochromatin function during

Drosophila development. Our developmental genetic analyses demonstrate that H3.3K9

is necessary for fertility but not viability in *Drosophila*. In addition, we find that some

euchromatic functions of H3K9 can be provided by either variant H3.3 or canonical H3,

whereas H3.3K9 cannot completely compensate for H3K9 in some regions of

heterochromatin as discussed below.

Several studies from multiple species have investigated the developmental

functions of H3.3 and H3. In mice, single mutation of either *H3.3A* or *H3.3B* results in

reduced viability and compromised fertility (Bush et al. 2013; Couldrey et al. 1999).

Similarly, *Drosophila H3.3A* and *H3.3B* double mutants appear at lower than expected

Mendelian ratios and are sterile (Sakai et al. 2009). H3.3 in *Tetrahymena thermophila*

is also important for sexual reproduction, although it is not required for viability or

maintenance of nucleosome density (Cui et al. 2006). Both *Tetrahymena* and

Drosophila H3.3 and H3 can compensate for one another. In *Tetrahymena*, canonical

H3 is dispensable if H3.3 is overexpressed (Cui et al. 2006). Similarly in *Drosophila*,

transgenic expression of H3 can rescue both the semi-lethality (Sakai et al. 2009) and

infertility (Hödl and Basler 2012) of *H3.3* mutants, indicating some functional

redundancy between the two histones. Indeed, when expressed equivalently,

Drosophila H3.3 can provide all of the developmental functions of H3 (Hödl and Basler

2012). Moreover, H3.3 is the sole H3 protein in *S. pombe* and *S. cerevisiae* yeast

(Malik and Henikoff 2003).

H3.3K9 functions in heterochromatin

We find that under endogenous expression conditions, H3.3K9 functions in

heterochromatin, including pericentromeric and telomeric regions of the genome. We

detected H3.3K9 methylation in pericentromeric heterochromatin, congruous with

previous data demonstrating that H3.3 in *Drosophila* is deposited at the chromocenter of

polytene chromosomes in a replication-dependent manner (Schwartz and Ahmad

2005). We also observed that *H3.3^{K9R}* mutants exhibited an abnormal chromocenter

structure in polytene chromosomes. Moreover, we provide evidence that H3.3K9 is

required for maintenance of telomeric chromatin architecture and repression of certain

telomeric transcripts, indicating that replication-coupled expression of H3 cannot provide

these particular H3K9 functions. These findings in *Drosophila* are consistent with

studies in mouse embryonic stem cells showing that H3.3 is localized to telomeres, is

methylated at K9, and functions in repression of telomeric repeat-containing RNAs

(Goldberg et al. 2010; Udugama et al. 2015). Conversely, the genetic data we

presented here and previously (Penke et al. 2016) indicate that H3K9 is essential for

repression of transposon-derived transcripts in pericentric heterochromatin, and H3.3K9

cannot compensate for the lack of H3K9 at these regions of the genome. The role of

H3.3K9 in telomere structure and function may be independent of HP1, as HP1 is

recruited to telomeres via the terminin complex independently of H3K9me (Raffa et al.

2011; Vedelek et al. 2015; Badugu et al. 2003).

K9ac regulates euchromatic gene expression

Previous studies that mapped histone modifications across the genome identified K9ac

as a characteristic of transcriptionally active regions (Kharchenko et al. 2011; Bernstein

et al. 2005; Liang et al. 2004; Roh et al. 2005). Moreover, mutation of H3K9

acetyltransferases results in compromised transcriptional activity (Wang et al. 1998;

Georgakopoulos and Thireos 1992; Kuo et al. 1998). However, H3K9 acetyltransferases

have non-histone substrates in addition to H3K9, and decreased transcriptional output

may be the result of pleiotropic effects (Glozak et al. 2005; Spange et al. 2009;

Fillingham et al. 2008). Our study provides evidence that K9ac, rather than non-histone

targets of H3K9 acetyltransferases, contributes to activating transcription, as *H3.3^{K9R}*

and *H3^{K9R}* mutants exhibit reduced gene expression in regions normally enriched for

K9ac. Importantly, these K9ac rich regions with reduced gene expression are not

normally enriched in K9me2 or me3, indicating the observed phenotype is not due to

changes in K9me2 or me3 and likely results from loss of K9ac. This change in gene

expression was accompanied by a fully penetrant lethality early in larval development of

H3.3^{K9R} H3^{K9R} combined mutant animals, raising the possibility that gene expression

control via acetylation of H3K9 is critical for the completion of animal development.

These data are also consistent with previous studies in *C. elegans* demonstrating that

H3K9 methylation is not essential for viability (Towbin et al. 2012; Zeller et al. 2016).

Overlapping and distinct genomic functions of H3K9 and H3.3K9

Functional overlap of H3K9 and H3.3K9 appears to vary at different regions of the

genome. Whereas H3.3K9 and H3K9 can perform similar functions in euchromatic

regions of the genome and can fully compensate for one another, our RNA-seq data

demonstrate H3.3K9 can only partially compensate for H3K9 in regions of

heterochromatin. Partial compensation by H3.3K9 in regions of K9me2/me3 is in line

with previous studies showing H3.3 is found at heterochromatin (Goldberg et al. 2010;

Lewis et al. 2010; Wong et al. 2010) and plays a role in transposon repression (Elsässer

et al. 2015). In the genotypes we analyzed, mRNA encoding variant and canonical H3

are expressed from their native promoters. Thus, disparity in functional overlap might be

due to differences in modes of expression and deposition and thus total amounts of

variant and canonical H3 histones in particular regions of the genome. For instance, H3

is normally enriched in heterochromatin compared to H3.3 (Ahmad and Henikoff 2002),

which may cause $H3^{K9R}$ mutations to be more detrimental in these regions. However,

H3.3 may be able to provide all H3 function when highly expressed in a replication-

dependent manner, as a transgenic histone gene array in which the *H3.2* coding region

was replaced by *H3.3* is nearly fully functional in larval imaginal discs (Hödl and Basler

2012). Thus, differences in expression and/or deposition into chromatin may be the only

basis for functional differences between H3.3 and H3.2 that we observed.

Heterochromatin may be particularly sensitive to incorporation of non-modifiable

K9 residues. H3K9 methylation serves as a binding site for the protein HP1, which can

in turn recruit H3K9 methyltransferases (Elgin and Reuter 2013; Grewal and Jia 2007).

Methylation of a neighboring nucleosome can restart the cycle and initiate propagation

of a heterochromatic configuration along the chromosome. Introduction of even a small

number of H3K9R containing nucleosomes may therefore disrupt this cycle and prevent

proper heterochromatin formation and gene repression. Incorporation of H3.3B^{K9R}

histones into regions of heterochromatin may dominantly affect chromatin structure,

resulting in the observed phenotypes at pericentromeres and telomeres in H3.3^{K9R}

mutants. In contrast, incorporation of low amounts of H3K9R histones in euchromatin

may not reduce K9ac levels sufficiently to disrupt gene expression. Finally, amino acid

differences in variant and canonical H3 may direct distinct histone modification states on

different histone types by influencing the binding of chromatin modifying enzymes

(Jacob et al. 2014). Different histone modification states on H3.3 and H3 may underlie

variation in compensation at different genomic regions.

In sum, our data investigating H3.3K9 and H3K9 function provide evidence that

K9ac activates gene expression and advance our understanding of the overlapping and

distinct functional roles of variant and canonical histones.

Figure 1: K9me2/me3 and HP1a signal is decreased in H3.3^{K9R} mutants. A) 3rd

instar larval salivary gland polytene chromosome spreads from wild-type (left) and

H3.3^{K9R} mutants (right) stained with anti-K9me2, anti-K9me3, anti-HP1a, and DAPI to

mark DNA. Right panel for each genotype shows enlarged chromocenter indicated by

white boxes. Bottom panel shows magnified view of telomere indicated by yellow boxes.

Scale bar = 20 microns (whole polytene) 5 microns (chromocenter/telomere). B, C)

Immunofluorescent signal of K9me2 (B) or HP1a (C) at chromocenters in wild-type (WT)

and H3.3^{K9R} mutants (a.u. = arbitrary units). Values were normalized to area of the

chromocenter and set relative to the average WT value from matched slides (see

Supplementary Experimental Methods). Significance was determined using t-test (*

$p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$). D) Western blot of K9me2 from salivary glands with

H3 used as loading control. E) K9me2 signal was quantified by densitometry and

normalized to corresponding H3 loading control band. Normalized values were set

relative to WT normalized signal. Error bars represent standard error of the mean from

two independent biological replicates (see Materials and Methods). F) Quantification of

chromocenter organization from *WT, H3.3^{K9R}, H3.3A^{Null}, H3.3B^{K9R}* mutants.

Figure 2: K9me2/me3 signal is diminished in K9R mutants. A) 1st instar larval brains

stained with anti-K9me2 and DAPI to mark DNA from *H3^{HWT}, H3^{K9R}, H3.3^{K9R} H3^{HWT}*, and

H3.3^{K9R} H3^{K9R} animals. Left panel shows max projection of 2 micrometer confocal

sections through the entire brain. Right panel shows a magnified, single confocal

section from the area indicated by the white boxes. Arrowheads indicate cells with

residual K9me2 signal in *H3.3^{K9R} H3^{K9R}* animals. Scale bar = 50 microns (whole brain)

10 microns (enlarged image).

Figure 3: K9ac signal is decreased in H3.3^{K9R} mutants. A,B) Polytene chromosome

spreads from wild-type (*WT*) and *H3.3^{K9R}* mutants (A) or *H3^{HWT}* and *H3^{K9R}* mutants (B)

stained with anti-K9me2, anti-K9ac, anti-HP1a, and DAPI to mark DNA. Scale bar = 20

microns. C) Western blot of K9ac from salivary glands with H3 used as loading control.

K9ac signal was quantified by densitometry and normalized to corresponding H3

loading control band. Normalized values were set relative to WT normalized signal.

Error bars represent standard error of the mean from two independent biological

replicates (see Materials and Methods).

Figure 4: H3.3K9 regulates chromatin architecture in regions of K9me. A-C)

Mutant: WT ratio of *H3.3A^{null}* (A), *H3.3B^{K9R}* (B), or *H3.3^{K9R}* (C) FAIRE signal from 3rd

instar imaginal wing discs at 19,738 FAIRE peaks called by MACS2. Red dots indicate

significantly different peaks (p<0.05), and insets indicate the number of significantly

increased (top) or decreased (bottom) peaks. Average counts signify average

normalized reads that overlap a peak in mutant and WT samples. D) Percentage of

peaks in a particular chromatin state that have significantly different FAIRE signal in

mutants versus WT (top). Bottom panel shows a summary of histone modifications or

proteins that define a chromatin state and the number of FAIRE peaks assigned to a

given chromatin state. E) Boxplot of FAIRE enrichment over input at 126 transposon

families (* indicates p < 0.05 and *** indicates p <0.0005). F) Plot from C showing only

those peaks that overlap an K9me2 or K9me3 peak from modENCODE ChIP-seq data.

Figure 5: Imaginal wing disc FAIRE signal of *H3.3* mutants is increased at

telomeres and transposons. A) Boxplot of average FAIRE enrichment determined by

piPipes pipeline across 126 transposon families (Han et al. 2015). Genomic DNA from

Drosophila embryos used as input control. B) Boxplots in A shown alongside FAIRE

enrichment for $H3^{HWT}$ and $H3^{K9R}$ mutants from a separate experiment (Penke et al.

2016). C) FAIRE enrichment of *H3.3* and $H3^{K9R}$ mutants at telomeric transposons. Error

bars indicate standard deviation from three replicates for each genotype. Statistical

significance determined by paired t-test (p<0.05 *, p<0.005 **, p<0.0005 ***, n.s. = not

significant).

Figure 6: H3.3K9 and H3K9 redundantly regulate gene expression. A-C) Mutant:

WT ratio of $H3^{K9R}$ (A), $H3.3^{K9R}$ (B), or $H3.3^{K9R}$ $H3^{K9R}$ (C) RNA signal from 1st instar

larvae at 10,253 transcripts assembled by Cufflinks. The Y axis indicates the $\log_2$

transformation of mutant/control signal between the genotypes being compared

(indicated at the top of each plot). Red dots indicate significantly different transcripts

(p<0.05) and insets signify the number of significantly increased (top) or decreased

(bottom) transcripts. Average coverage signifies the average number of normalized

reads that overlap a transcript in mutant and $H3^{HWT}$ samples.

Figure 7: H3.3K9 and H3K9 redundancy differs in heterochromatin and

euchromatin. A, B) Percentage of transcripts in a chromatin state that have

significantly increased (A) or decreased (B) RNA signal in mutants versus $H3^{HWT}$. C)

Table indicates the number of transcripts that overlap a particular chromatin state. D, E)

Heatmaps showing fold change of K9R mutants over $H3^{HWT}$ at chromatin state 7

regions (D) and chromatin state 1 regions (E). Each row indicates a transcript that

overlaps the indicated chromatin state.

Figure 8: K9ac associated transcripts are altered in $H3.3^{K9R}$ $H3^{K9R}$ double mutants.

MA plot showing fold change of normalized RNA signal in $H3^{K9R}$ (A), $H3.3^{K9R}$ $H3^{HWT}$ (B),

and $H3.3^{K9R}$ $H3^{K9R}$ (C) mutants versus $H3^{HWT}$ at all transcripts from merged

transcriptome. Average coverage on X-axis represents the mean expression level of a

transcript. Transcripts that overlap an K9ac peak called from ChIP-seq data

(GSM1363590 ; Pérez-Lluch et al. 2015) are shown in the left panel while those that do

not are shown in the right panel. Significance (shown in red) was determined using

DESeq2 (Love et al. 2015) and an adjusted p value cutoff of 0.05.

Acknowledgement

We thank Bhawana Bariar for assistance generating the gRNA construct, Jeff Sekelsky

for the pBlueSurf construct, and Kami Ahmad for providing the $H3.3A^{2x1}$ flies. This work

was supported by 5T32GM007092-39 and F31GM115194 to T.J.R.P and

R01DA036897 to D.J.M, B.D.S, A.G.M, and R.J.D.

REFERENCES

Ahmad K, Henikoff S. 2002. The histone variant H3.3 marks active chromatin by

replication-independent nucleosome assembly. *Mol Cell* **9**: 1191–1200.

Allis CD, Wiggins JC. 1984. Histone rearrangements accompany nuclear differentiation

and dedifferentiation in Tetrahymena. *Dev Biol* **101**: 282–294.

Badugu RK, Shareef MM, Kellum R. 2003. Novel Drosophila Heterochromatin Protein I

(HP1)/Origin Recognition Complex-associated Protein (HOAP) Repeat Motif in

HP1/HOAP Interactions and Chromocenter Associations. *J Biol Chem* **278**: 34491–

34498.

Bannister a J, Zegerman P, Partridge JF, Miska E a, Thomas JO, Allshire RC,

Kouzarides T. 2001. Selective recognition of methylated lysine 9 on histone H3 by

the HP1 chromo domain. *Nature* **410**: 120–4.

Bannister AJ, Schneider R, Myers FA, Thorne AW, Crane-Robinson C, Kouzarides T.

2005. Spatial distribution of di- and tri-methyl lysine 36 of histone H3 at active

genes. *J Biol Chem* **280**: 17732–17736.

Belyaeva ES, Zhimulev IF, Volkova EI, Alekseyenko a a, Moshkin YM, Koryakov DE.

1998. Su(UR)ES: a gene suppressing DNA underreplication in intercalary and

pericentric heterochromatin of Drosophila melanogaster polytene chromosomes.

Proc Natl Acad Sci U S A **95**: 7532–7.

Bernstein BE, Kamal M, Lindblad-Toh K, Bekiranov S, Bailey DK, Huebert DJ,

McMahon S, Karlsson EK, Kulbokas EJ, Gingeras TR, et al. 2005. Genomic maps

and comparative analysis of histone modifications in human and mouse. *Cell* **120**:

169–181.

Blower MD, Karpen GH. 2001. The role of Drosophila CID in kinetochore formation, cell-

cycle progression and heterochromatin interactions. *Nat Cell Biol* **3**: 730–9.

Bush KM, Yuen BT, Barrilleaux BL, Riggs JW, O'Geen H, Cotterman RF, Knoepfler PS.

2013. Endogenous mammalian histone H3.3 exhibits chromatin-related functions

during development. *Epigenetics Chromatin* **6**: 7.

Cai W, Jin Y, Girton J, Johansen J, Johansen KM. 2010. Preparation of Drosophila

Polytene Chromosome Squashes for Antibody Labeling. *Jove* 2–5.

Cenci G, Ciapponi L, Gatti M. 2005. The mechanism of telomere protection: A

comparison between Drosophila and humans. *Chromosoma* **114**: 135–145.

Couldrey C, Carlton MB, Nolan PM, Colledge WH, Evans MJ. 1999. A retroviral gene

trap insertion into the histone 3.3A gene causes partial neonatal lethality, stunted

growth, neuromuscular deficits and male sub-fertility in transgenic mice. *Hum Mol*

Genet **8**: 2489–2495.

Cui B, Liu Y, Gorovsky MA. 2006. Deposition and function of histone H3 variants in

Tetrahymena thermophila. *Mol Cell Biol* **26**: 7719–30.

Elgin SCR, Reuter G. 2013. Position-effect variegation, heterochromatin formation, and

gene silencing in Drosophila. *Cold Spring Harb Perspect Biol* **5**.

Elsässer SJ, Noh K-M, Diaz N, Allis CD, Banaszynski L a. 2015. Histone H3.3 is

required for endogenous retroviral element silencing in embryonic stem cells.

Nature **522**: 240–246.

Estella C, McKay DJ, Mann RS. 2008. Molecular Integration of Wingless,

Decapentaplegic, and Autoregulatory Inputs into Distalless during Drosophila Leg

Development. *Dev Cell* **14**: 86–96.

Fillingham J, Recht J, Silva AC, Suter B, Emili A, Stagljar I, Krogan NJ, Allis CD, Keogh

M-C, Greenblatt JF. 2008. Chaperone control of the activity and specificity of the

histone H3 acetyltransferase Rtt109. *Mol Cell Biol* **28**: 4342–53.

Georgakopoulos T, Thireos G. 1992. Two distinct yeast transcriptional activators require

the function of the GCN5 protein to promote normal levels of transcription. *EMBO J*

11: 4145–52.

Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison C a, Smith HO, Iii CAH,

America N. 2009. Enzymatic assembly of DNA molecules up to several hundred

kilobases. *Nat Methods* **6**: 343–5.

Glozak MA, Sengupta N, Zhang X, Seto E. 2005. Acetylation and deacetylation of non-

histone proteins. *Gene* **363**: 15–23.

Goldberg AD, Banaszynski LA, Noh K, Lewis PW, Elsaesser SJ, Stadler S, Dewell S,

Law M, Guo X, Li X, et al. 2010. Distinct Factors Control Histone at Specific

Genomic Regions. *Cell* **140**: 678–691.

Graveley BR, Brooks AN, Carlson JW, Duff MO, Landolin JM, Yang L, Artieri CG, Baren

MJ Van, Boley N, Booth BW, et al. 2011. The Developmental Transcriptome of

Drosophila melanogaster. *Nature* **471**: 473–479.

Grewal SIS, Jia S. 2007. Heterochromatin revisited. *Nat Rev Genet* **8**: 35–46.

Hake SB, Garcia BA, Duncan EM, Kauer M, Dellaire G, Shabanowitz J, Bazett-jones

DP, Allis CD, Hunt DF. 2006. Expression Patterns and Post-translational

Modifications Associated with Mammalian Histone H3 Variants. *J Biol Chem* **281**:

559–568.

Han BW, Wang W, Zamore PD, Weng Z. 2015. PiPipes: A set of pipelines for piRNA

and transposon analysis via small RNA-seq, RNA-seq, degradome-and CAGE-seq,

ChIP-seq and genomic DNA sequencing. *Bioinformatics* **31**: 593–595.

Henikoff S, Ahmad K. 2005. Assembly of Variant Histones into Chromatin. *Annu Rev*

Cell Dev Biol.

Hödl M, Basler K. 2012. Transcription in the Absence of Histone H3.2 and H3K4

Methylation. *Curr Biol* 2253–2257.

Jacob Y, Bergamin E, Donoghue MT a, Mongeon V, LeBlanc C, Voigt P, Underwood

CJ, Brunzelle JS, Michaels SD, Reinberg D, et al. 2014. Selective methylation of

histone H3 variant H3.1 regulates heterochromatin replication. *Science* **343**: 1249–

53.

Jin C, Felsenfeld G. 2007. Nucleosome stability mediated by histone variants H3.3 and

H2A.Z. *Genes Dev* 1519–1529.

Jin C, Zang C, Wei G, Cui K, Peng W, Zhao K, Felsenfeld G, Heart TN. 2011.

H3.3/H2A.Z double variant-containing nucleosomes mark "nucleosome-free

regions" of active promoters and other regulatory regions in the human genome.

Nat Genet **41**: 941–945.

Kharchenko P, Alekseyenko A, Schwartz YB, Minoda A, Riddle NC, Sabo PJ, Larschan

EN, Gorchakov AA, Karpen GH, Park PJ. 2011. Comprehensive analysis of the

chromatin landscape in Drosophila melanogaster. *Nature* **471**: 480–485.

Kondo S, Ueda R. 2013. Highly Improved gene targeting by germline-specific Cas9

expression in Drosophila. *Genetics* **195**: 715–721.

Kraushaar DC, Jin W, Maunakea A, Abraham B, Ha M, Zhao K. 2013. Genome-wide

incorporation dynamics reveal distinct categories of turnover for the histone variant

H3 . 3. *Genome Biol* **14**: 1.

Kuo M-H, Zhou J, Jambeck P, Churchill MEA, Allis CD. 1998. Histone acetyltransferase

activity of yeast Gcn5p is required for the activation of target genes in vivo. *Genes*

Dev **12**: 627–639.

Lachner M, O'Carroll D, Rea S, Mechtler K, Jenuwein T. 2001. Methylation of histone

H3 lysine 9 creates a binding site for HP1 proteins. *Nature* **410**: 116–120.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat*

Methods **9**: 357–360.

Levis RW, Ganesan R, Houtchens K, Tolar LA, Sheen F miin. 1993. Transposons in

place of telomeric repeats at a Drosophila telomere. *Cell* **75**: 1083–1093.

Lewis PW, Elsaesser SJ, Noh K, Stadler SC, Allis CD. 2010. Daxx is an H3.3-specific

histone chaperone and cooperates with ATRX in replication-independent chromatin

assembly at telomeres. *PNAS* **107**: 1–6.

Liang G, Lin JCY, Wei V, Yoo C, Cheng JC, Nguyen CT, Weisenberger DJ, Egger G,

Takai D, Gonzales FA, et al. 2004. Distinct localization of histone H3 acetylation

and H3-K4 methylation to the transcription start sites in the human genome. *Proc*

Natl Acad Sci **101**: 7357–7362.

Love M, Anders S, Huber W. 2015. *Differential analysis of count data – the DESeq2*

package.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* 1–21.

Malik HS, Henikoff S. 2003. Phylogenomics of the nucleosome. *Nat Struct Biol* **10**: 882–

91.

Marzluff WF, Gongidi P, Woods KR, Jin J, Maltais LJ. 2002. The Human and Mouse

Replication-Dependent Histone Genes. *Genomics* **80**: 487–498.

McKay DJ, Klusza S, Penke TJR, Meers MP, Curry KP, McDaniel SL, Malek PY,

Cooper SW, Tatomer DC, Lieb JD, et al. 2015. Interrogating the Function of

Metazoan Histones using Engineered Gene Clusters. *Dev Cell* **32**: 373–386.

McKay DJ, Lieb JD. 2013. A Common Set of DNA Regulatory Elements Shapes

Drosophila Appendages. *Dev Cell* **27**: 306–318.

McKittrick E, Gafken PR, Ahmad K, Henikoff S. 2004. Histone H3.3 is enriched in

covalent modifications associated with active chromatin. *Proc Natl Acad Sci U S A*

101: 1525–30.

Mellone BG, Allshire RC. 2003. Stretching it: putting the CEN(P-A) in centromere. *Curr*

Opin Genet Dev **13**: 191–198.

Mito Y, Henikoff JG, Henikoff S. 2005. Genome-scale profiling of histone H3.3

replacement patterns. *Nat Genet* **37**.

Nakayama J, Rice JC, Strahl BD, Allis CD, Grewal SIS. 2001. Role of Histone H3

Lysine 9 Methylation in Epigenetic Control of Heterochromatin Assembly. *Science*

(80-) **292**: 110–113.

Penke TJR, McKay DJ, Strahl BD, Gregory Matera A, Duronio RJ. 2016. Direct

interrogation of the role of H3K9 in metazoan heterochromatin function. *Genes Dev*

30: 1866–1880.

Pérez-Lluch S, Blanco E, Tilgner H, Curado J, Ruiz-Romero M, Corominas M, Guigó R.

2015. Absence of canonical marks of active chromatin in developmentally regulated

genes. *Nat Genet* **47**: 1158–67.

Perrini B, Piacentini L, Fanti L, Altieri F, Chichiarelli S, Berloco M, Turano C, Ferraro A,

Pimpinelli S. 2004. HP1 controls telomere capping, telomere elongation, and

telomere silencing by two different mechanisms in Drosophila. *Mol Cell* **15**: 467–

476.

Price BD, Andrea ADD. 2014. Chromatin Remodeling at DNA Double Strand Breaks.

Cell **152**: 1344–1354.

Quinlan AR, Hall IM. 2010. BEDTools☐: a flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26**: 841–842.

Raffa GD, Ciapponi L, Cenci G, Gatti M. 2011. Terminin: A protein complex that

mediates epigenetic maintenance of Drosophila telomeres. *Nucleus* **2**: 383–391.

Roh T, Cuddapah S, Zhao K. 2005. Active chromatin domains are defined by

acetylation islands revealed by genome-wide mapping -- Roh et al. 19 (5): 542 --

Genes and Development. 542–552.

Sakai A, Schwartz BE, Goldstein S, Ahmad K. 2009. Transcriptional and developmental

functions of the H3.3 histone variant in Drosophila. *Curr Biol* **19**: 1816–20.

Schwartz BE, Ahmad K. 2005. Transcriptional activation triggers deposition and

removal of the histone variant H3 . 3. *Genes Dev* 804–814.

Scully R, Xie A. 2013. Double strand break repair functions of histone H2AX. *Mutat Res*

750: 5–14.

Simon JM, Giresi PG, Davis IJ, Lieb JD. 2013. A detailed protocol for formaldehyde-

assisted isolation of regulatory elements (FAIRE). *Curr Protoc Mol Biol* **Chapter 21**:

Unit21.26.

Spange S, Wagner T, Heinzel T, Krämer OH. 2009. Acetylation of non-histone proteins

872 modulates cellular signalling at multiple levels. *Int J Biochem Cell Biol* **41**: 185–198.

873 Szenker E, Ray-Gallet D, Almouzni G. 2011. The double face of the histone variant

874  H3.3. *Cell Res* **21**: 421–434.

875 Tagami H, Ray-Gallet D, Almouzni G, Nakatani Y. 2004. Histone H3.1 and H3.3

876  Complexes Mediate Nucleosome Assembly Pathways Dependent or Independent

877  of DNA Synthesis. *Cell* **116**: 51–61.

878 Talbert PB, Henikoff S. 2010. Histone variants--ancient wrap artists of the epigenome.

879  *Nat Rev Mol Cell Biol* **11**: 264–275.

880 Talbert PB, Henikoff S. 2017. Histone variants on the move: substrates for chromatin

881  dynamics. *Nat Rev Mol Cell Biol* **18**: 115–126.

882 Tamura T, Smith M, Kanno T, Dasenbrock H, Nishiyama A, Ozato K. 2009. Inducible

883  deposition of the histone variant H3.3 in interferon-stimulated genes. *J Biol Chem*

884  **284**: 12217–12225.

885 Towbin BD, González-Aguilera C, Sack R, Gaidatzis D, Kalck V, Meister P, Askjaer P,

886  Gasser SM. 2012. Step-wise methylation of histone H3K9 positions

887  heterochromatin at the nuclear periphery. *Cell* **150**: 934–47.

888 Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL,

889  Rinn JL, Pachter L. 2014. Differential gene and transcript expression analysis of

890  RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc.*

891 Udugama M, Chang FTM, Chan FL, Tang MC, Pickett H a, McGhie JDR, Mayne L,

892  Collas P, Mann JR, Wong LH. 2015. Histone variant H3.3 provides the

893  heterochromatic H3 lysine 9 tri-methylation mark at telomeres. *Nucleic Acids Res*

894  gkv847-.

895   Vedelek B, Blastyák A, Boros IM. 2015. Cross-species interaction between rapidly

896   evolving telomere-specific drosophila proteins. *PLoS One* **10**: 1–16.

897   Verreault A, Kaufman PD, Kobayashi R, Stillman B. 1996. Nucleosome assembly by a

898   complex of CAF-1 and acetylated histones H3/H4. *Cell* **87**: 95–104.

899   Wallrath LL, Vitalini MW, Elgin SCR. 2014. Fundamentals of Chromatin eds. J.L.

900   Workman and S.M. Abmayr. 529–552.

901   Wang L, Liu L, Berger SL. 1998. Critical residues for histone acetylation by Gcn5,

902   functioning in Ada and SAGA complexes, are also required for transcriptional

903   function in vivo. *Genes Dev* 640–653.

904   Wong LH, Mcghie JD, Sim M, Anderson MA, Ahn S, Hannan RD, George AJ, Morgan

905   KA, Mann JR, Choo KHA. 2010. ATRX interacts with H3.3 in maintaining telomere

906   structural integrity in pluripotent embryonic stem cells. *Genome Res* 351–360.

907   Xu M, Long C, Chen X, Huang C, Chen S, Zhu B. 2010. Partitioning of Histone H3-H4

908   Tetramers During DNA Replication-Dependent Chromatin Assembly. *Science (80-)*

909   **328**: 94–98.

910   Zeller P, Padeken J, Schendel R Van, Kalck V, Tijsterman M, Gasser SM. 2016.

911   Histone H3K9 methylation is dispensable for Caenorhabditis elegans development

912   but suppresses RNA□: DNA hybrid-associated repeat instability. *Nat Genet* 1–13.

913   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C,

914   Myers RM, Brown M, Li W, et al. 2008. Open Access Model-based Analysis of

915   ChIP-Seq (MACS). *Genome Biol* **9**.

916   Zhimulev IF, Belyaeva ES, Semeshin VF, Shloma V V, Makunin I V, Volkova EI. 2003.

917   Overexpression of the SuUR gene induces reversible modifications at pericentric,

918   telomeric and intercalary heterochromatin of Drosophila melanogaster polytene

919   chromosomes. *J Cell Sci* **116**: 169–76.

Table 1: Genotype description of H3.3 and H3 K9R mutants.

Nomenclature[a]	Canonical Histone Genotypes		Variant Histone Genotypes	
	Endogenous	Transgenic	H3.3B	H3.3A
WT	WT	-	WT	WT
H3.3B^{K9R}	WT	-	K9R	WT
H3.3A^{Null}	WT	-	WT	Δ
H3.3^{K9R}	WT	-	K9R	Δ
H3^{HWT}	Δ	WT	WT	WT
H3^{K9R}	Δ	K9R	WT	WT
H3.3^{K9R} H3^{HWT}	Δ	WT	K9R	Δ
H3.3^{K9R} H3^{K9R}	Δ	K9R	K9R	Δ

[a] Wild-type (WT), gene deletion (Δ), no transgenic histone array (-). See Supplementary Table 1 for full genotypes.

Table 2: *H3.3^{K9R}* mutants are viable but sterile.

H3.3B	H3.3A[a]	Observed	Expected	p[b]	Fertile
WT	*Δ/+*	535	535.3	n.s.	yes
WT	*Δ*	268	267.7	n.s.	yes
K9R	*Δ/+*	400	438	<0.005	yes
K9R	*Δ*	257[c]	219	<0.005	No[d]

[a] *H3.3A^{2x1}* deletion allele
[b] p-value calculated with a chi-square test
[c] The higher than expected number of observed *H3.3B^{K9R} H3.3A^{Null}* animals is presumably due to non-specific detrimental effects caused by the presence of a balancer chromosome in siblings with the *H3.3B^{K9R}* mutation and balancer-derived wild-type *H3.3A*.
[d] Both males and females.

Table 3: *H3.3^{K9R}* and *H3^{K9R}* mutations are synthetically lethal.

Genotype	Embryo Hatching				Pupate[a]				Eclose			
	Obs	No.	%	P[b]	Obs	No.	%	p	Obs	No.	%	p
H3^{HWT}	389	450	86.4	-	98	140	70.0	-	88	140	62.9	-
H3^{K9R}	370	480	77.1	<0.0005	183	285	64.2	<0.05	3	285	1.1	<0.0005
H3.3^{K9R} H3^{HWT}	350	465	75.3	<0.0005	279	462	60.4	<0.0005	160	462	34.6	<0.0005
H3.3^{K9R} H3^{K9R}	214	325	65.8	<0.0005	0	130	0.0	<0.0005	0	130	0.0	<0.0005

[a] The pupation and eclosion values have an identical number of animals analyzed for each genotype because they were obtained from the same brood of animals, while the embryo hatching values were obtained from independent experiments.
[b] p-value calculated with a chi-square test using *H3^{HWT}* observed (Obs) values as expected values.

A

WT   H3.3^K9R

Merge

DAPI

K9me2

K9me3

HP1a

DAPI   HP1a   DAPI   HP1a

B K9me2 chromocenter

***

Signal / area (a.u.)

WT   H3.3^K9R

C HP1 chromocenter

*

Signal / area (a.u.)

WT   H3.3^K9R

D

H3.3^K9R   WT   H3^HWT   H3^K9R

α-K9me2

α-H3

E K9me2

Normalized Signal

H3.3^K9R   WT   H3^HWT   H3^K9R

F

n=139   n=88   n=77   n=91

Percent chromocenters

WT   H3.3^K9R   H3.3A^Null   H3.3B^K9R

organized

moderately organized

disorganized

A H3.3A^Null vs. WT

B H3.3B^K9R vs. WT

C H3.3^K9R vs. WT

D

E

state	definition	peaks
1	H3K4me2/3, H3K9ac	5433
2	H3K36me3	691
3	H3K18ac, H3K27ac	3134
4	H3K36me	2260
5	H4K16ac, H3K36me3	534
6	H3K27me2/3	1400
7	H3K9me2/3	823
8	Low H3K9me2/3	133
9	H1, SUUR	3557

F H3K9me2/me3 overlap
H3.3^K9R vs. WT

A H3^{K9R} vs. H3^{HWT}

Fold Change (log2)

Average coverage (log2)

B H3.3^{K9R} H3^{HWT} vs. H3^{HWT}

Fold Change (log2)

Average coverage (log2)

C H3.3^{K9R} H3^{K9R} vs. H3^{HWT}

1036

Fold Change (log2)

Average coverage (log2)

A

Significantly increased

Legend:
- H3^{K9} (blue)
- H3.3^{K9R} H3^{HWT} (red)
- H3.3^{K9R} H3^{K9R} (purple)

% transcripts vs chromatin state (1–9)

B

Significantly decreased

Legend:
- H3^{K9R} (blue)
- H3.3^{K9R} H3^{HWT} (red)
- H3.3^{K9R} H3^{K9R} (purple)

% transcripts vs chromatin state (1–9)

C

state	definition	genes
1	H3K4me2/3, H3K9ac	4848
2	H3K36me3	3641
3	H3K18ac, H3K27ac	1608
4	H3K36me	1857
5	H4K16ac, H3K36me3	1066
6	H3K27me2/3	777
7	H3K9me2/3	767
8	Low H3K9me2/3	389
9	H1, SUUR	4004

D

chromatin state 7 (767 genes)

Fold change (log$_2$)

H3^{K9R} H3.3^{K9R} H3^{HWT} H3.3^{K9R} H3^{K9R}

E

chromatin state 1 (4848 genes)

Fold change (log$_2$)

H3^{K9R} H3.3^{K9R} H3^{HWT} H3.3^{K9R} H3^{K9R}

H3K9ac

Nonoverlap

A

H3^K9R vs. H3^HWT

H3^K9R vs. H3^HWT

B

H3.3^K9R H3^HWT vs. H3^HWT

H3.3^K9R H3^HWT vs. H3^HWT

C

H3.3^K9R H3^K9R vs. H3^HWT

H3.3^K9R H3^K9R vs. H3^HWT

267