Molecular characterization of breast and lung tumors by integration of multiple data types with sparse-factor analysis

Tycho Bismeijer[1], Sander Canisius[1,2], Lodewyk Wessels[1,3]

1 Division of Molecular Carcinogenesis, The Netherlands Cancer Institute, Amsterdam, The Netherlands
2 Division of Molecular Pathology, The Netherlands Cancer Institute, Amsterdam, The Netherlands
3 Faculty of EEMCS, Delft University of Technology, Delft, The Netherlands

Abstract

Effective cancer treatment is crucially dependent on the identification of the biological processes that drive a tumor. This process is complicated by the fact that multiple processes may be active simultaneously and that each tumor has a unique spectrum of active processes. While clustering has been applied extensively to subtype tumors, its discrete nature makes it inherently unsuitable to this task. In addition, the availability of multiple data types per tumor has become commonplace and provides the opportunity to comprehensively profile the processes driving a tumor.

Here we introduce Functional Sparse Factor Analysis (FuncSFA) to address these challenges. FuncSFA integrates multiple data types to define a lower dimensional space capturing the variation in a set of tumors across those data types. A tailor-made module allows the association of the identified factors to biological processes. FuncSFA is inspired by iCluster, which we improve in several key aspects. First, we increase the convergence efficiency significantly by factor rescaling and more efficient optimization techniques, especially when analyzing multiple molecular datasets that have not been matched in terms of size or to contain only concordant features. Most importantly, unlike iCluster, FuncSFA does not assign samples to discrete clusters, but identifies the dominant driver processes active in each sample. This is achieved by a regression of the factors on the full RNA expression matrix followed by a functional enrichment analysis and manual curation step.

We apply FuncSFA to the TCGA breast and lung datasets. We identify several processes common to both cancer types although with subtle differences associated with each cancer type. These include EMT and Immune processes. In the breast cancer dataset we recover the known intrinsic subtypes and identify additional processes that have not been identified before in an integrated approach. These include immune infiltration and epithelial-mesenchymal transition, and processes driven by copy number gains on the 8q chromosome arm. In lung cancer we recover the major types (adenocarcinoma v squamous cell carcinoma) and identify processes that occur in both of these types. These include EMT, two immune processes, and the activity of the NFE2L2 transcription factor. In summary, FuncSFA is a robust method to perform discovery of key driver processes in a collection of samples through unsupervised integration of multiple molecular data types and functional annotation.

Introduction

Cancer is a heterogeneous disease, both at the molecular level and in response to treatment. If we can better understand the variation between tumors, we may get a better understanding of why tumors respond differently to treatment. This could, in turn, lead to better treatment selection for patients.

To chart the variation across tumors, projects such as The Cancer Genome Atlas (TCGA) have collected a variety of molecular data from thousands of tumors [1–3]. Analyses of these data provide a better understanding of the underlying biological processes associated with the cancer. For example, recurrent copy number abberations or recurrent point mutations may reveal the drivers of carcinogenesis. Complementary to this, RNA expression or protein phosphorylation can reveal downstream changes involving many genes, even if the upstream driver of those changes is unknown. Hence, the different data types are reflections of the same biological state, yet each of them encodes information not present or only partially present in the others. Therefore, a comprehensive characterization of the molecular variation across tumors requires the integration of multiple data types.

A popular approach to characterizing tumors is clustering of RNA expression data. Examples include the PAM50 subtypes [4] in breast cancer and the consensus subtypes in colorectal cancer [5]. These approaches do not take into account multiple data types, and will therefore fail to incorporate valuable information provided by other data types.

Integrative clustering approaches such as Bayesian consensus clustering [6], patient specific data fusion [7] and iCluster [8] do take multiple data types into account. However, clusters are unsuitable models of biological processes for at least two reasons. First, a biological process can be activated in multiple contexts and multiple independent biological processes can be active simultaneously. However, as clusters cannot overlap, it becomes a challenge to represent this variation in a discrete clustering. For example, immune infiltration occurs in both ER+ and ER- negative breast tumors, but once a sample is assigned to the ER+ cluster it cannot be assigned to an immune cluster that spans all breast cancer tumors. Second, the activity of a biological process can be continuous. This cannot be captured by discrete clusters.

Paradigm [9] improves upon the abovementioned approaches by integrating multiple data types to infer activity levels of biological processes. Activity levels are assigned to tumors independently of each other, avoiding the limitation of cluster analysis. To estimate these activity levels, Paradigm leverages existing knowledge available from pathway databases. A limitation of this approach is that using existing knowledge a priori limits discovery of new biological processes. More importantly, it also limits the discovery of biological processes in new contexts (e.g. tumor types) because activity of a process in a new context might involve a set of genes that is only partially overlapping with the genes currently annotated to that process.

Here we introduce FuncSFA, a sparse-factor analysis with a tailored gene-set enrichment analysis (GSEA) [10] that integrates multiple data types to provide both a continuous characterization and a functional interpretation of the variation across tumors at the molecular level (Fig 1). The sparse-factor analysis identifies factors explaining variation in multiple data types such as RNA expression, protein expression, and DNA copy number data. Subsequently, the factors are interpreted and linked to known biology using a gene-set enrichment analysis of the factors on the RNA expression data. The interpretation obtained from the gene-set enrichment analysis is validated by comparison of the genes, epitopes and copy number aberrations in the factor to external resources. Together this not only provides insight into variation across tumors but also the biology underlying the molecular data.

A Graphical representation of Sparse Factor Analysis (SFA)

Factors
SFA
Coefficients

mRNA
Features

CNA
Features

Protein
Features

B Sparse Factor Analysis

Joint likelihood of X and Z is is maximized under sparsity constraints on B.

C Factor Regression coefficients

D Functional annotation by GSEA

E Single sample factor calculation

Fig 1. Overview of FuncSFA A: Graphical representation of Functial Sparse Factor Analysis (FuncSFA). The green circles represent the factors, and the red, blue and yellow circles at the bottom represent the observed variables, with the colors representing the data types and each circle representing an individual variable. The black lines connecting the individual variables to the factors represent the regression coefficients. B: Graphical representation of the mathematical concepts of SFA with X representing the $N \times n$ data matrix, Z the $N \times k$ obtained factor matrix and B the $k \times n$ factor coefficients. C: Graphical representation of the computations of the factor expression coefficients. The coefficients represented by the $k \times n_m$ matrix C are obtained by regressing the $N \times n_m$ RNA expression matrix, X_m, on the $N \times k$ factor matrix Z. D: The gene-set enrichment analysis designed to assign biological processes or pathways to the obtained factors based on enrichment analyses. E: Application of the factors to determine the activity of the factors (or associated biological processes) in a new sample. (N: number of samples; n: number of features; k: number of factors; n_m: number of mRNA features; Z: factor matrix; X: data matrix (concatenation of mRNA, copy number and Reverse Phase Protein Array (RPPA) data); B: Sparse factor coefficients; C: Factor regression coefficients; GSEA: Gene-set enrichment analysis).

The sparse-factor analysis is based on a reinterpretation of the mathematical framework behind the iCluster method [8, 11]. Our reinterpretation improves upon the iCluster method in several key aspects. First, through proper factor rescaling and more efficient optimization approaches, we ensure convergence of FuncSFA on multiple molecular datasets that have not been pre-matched. An example of pre-matching is in the iC10 subtypes [12], where the authors selected genes where RNA expression agrees with DNA copy number. Second, in contrast to iCluster, FuncSFA does not assign samples to discrete clusters, but identifies the dominant driver processes across multiple molecular data types and across all samples, and then represents, for each sample, the spectrum of processes active in a given sample. Taken together, FuncSFA represents, for the first time, a robust method to perform discovery of the key biological processes driving the biology of a collection of samples through unsupervised integration of multiple molecular datatypes.

We applied FuncSFA to the TCGA breast and lung cohorts and identified 10 factors in each tumor type. As breast cancer is very well characterized, it served as a positive control in the sense that we could fully identify the known intrinsic subtypes in breast cancer. Uniquely, this characterizition integrates the intrinsic subtypes and the activity of Epithelial to Mesenchymal Transition (EMT) and Immune processes, each represented by an independent factor. In Lung cancer, which has remained largely uncharacterized, we also identified an EMT and Immune factor, as well as a factor associated with the main lung subtypes—Adenocarcinoma and Squamous Cell Carcinoma. We also identified a factor which captures the activation of the transcription factor NFE2L2. Here the power of integration of multiple data types is highlighted by the fact that the activity of this factor is associated with mutations in NFE2L2 as well as its inhibitor KEAP. We expect that the identified factors not only provide a more complete characterization of the biological processes active in the different samples, but will also provide a starting point for the development of better treatment strategies.

Results

FuncSFA

Sparse factor analysis

The basic idea behind sparse-factor analysis is the following. We assume that each tumor type is characterized by a set of key driving factors that give rise to molecular phenotypes. These driving factors cannot be directly measured, but are indirectly observed through noisy measurements of multiple molecular data types such as mRNA expression, copy number abberations and protein expression and modification. The challenge is to employ all these measured data types simultaneously to identify this unobserved structure in the data.

Sparse factor analysis addresses this challenge by modeling the unobserved structure by a relatively small number of continuous factors— the green circles in Fig 1A. These factors, in turn, explain the observed molecular data in a linear regression that links each molecular data type to the factors. The red, blue and yellow circles at the bottom of Fig 1A represent the observed variables, with the colors representing the data types and each circle representing an individual variable. For example, in ERBB2 (also known as HER2) positive breast cancer, the ERBB2 factor would represent the activation of the ERBB2 pathway while the measured copy number, protein and mRNA expression changes will be modeled as appropriately chosen regression functions of the ERBB2 factor.

The factors are identified by maximizing the joint likelihood of the the measured data and the factors, selecting factors and regression coefficients explaining the measured data. This is accomplished in a methodology that is akin to probabilistic principal component analysis (PCA) [13]. We employ elastic net regularization [14] to create parsimonious models by enforcing sparsity on the regression coefficients associated with the factors, as introduced by iCluster [8]. In Fig 1A the regression coefficients are represented by the black lines connecting the individual variables to the factors, while Fig 1B depicts the mathematical operations associated with the sparse-factor analysis. Consequently, a factor is defined by the subset of molecular variables that contribute to that factor as well as the regression coefficients that model the degree to which each molecular variable contributes to the factor.

Sparsity of regression coefficients induced by the elastic net penalty improves data integration by preventing one data type (especially mRNA expression) from dominating the analysis. Sparse weights are also biologically plausible as a biological process might involve the altered expression of thousands of genes but is unlikely to be driven by all genes.

A very important feature of FuncSFA is that, unlike iCluster, we do not force a sample assignment to discrete clusters, but represent the level of activity of each of the driving factors in a given sample. Consequently, each sample can be a 'member' of multiple factors, and this 'membership' can also vary in strength.

Gene-set enrichment analysis

While sparse-factor analysis efficiently identifies the hidden driving factors, it remains challenging to directly attach biological interpretation to the identified factors. In some cases, such as ERBB2 pathway activation, this may be straightforward, but in many other cases this remains very challenging. One of our important contributions include the development of a gene-set enrichment analysis tailored to the results of sparse-factor analysis.

The first step of the gene-set enrichment analysis is to regress the factors on the gene expression matrix (Fig 1C). This may sound counter-intuitive, as the whole purpose of the SFA was to pinpoint the genes contributing to the factor, hence revealing the underlying drivers. However, the motivation for this is three-fold. First, in the initial SFA the mRNA expression values of all genes may not be included for a number of reasons, including complexity reduction and to remove genes with a high signal to noise ratio. Consequently, the SFA coefficients may be incomplete. Second, the sparsity of the coefficients introduces zeros in the coefficients. Altough these zeros are beneficial for human interpreation and biological plausability, they complicate ranking of genes, an essential part of the gene-set enrichment analysis. Third, RNA expression remains a data type that captures most of the variation in the cell, and can, as such be quite informative regarding the activity of biological processes. So, having gone through the process of identifying driving factors that are robust, in the sense that they are common to all data types, we employ the regression of the factors to the complete RNA expression data set to identify all genes with expression patterns that show association with the factors. This association is captured in the regression coefficients (Matrix C in Fig 1C). We then normalize each coefficient by the gene variance to obtain the 'factor expression coefficients'.

The second step of the gene-set enrichment analysis is to rank the genes based on the the factor expression coefficients and compute the enrichment statistic for every gene-set factor pair (Fig 1D) [10]. We determined statistical significance of the enrichments by a sample permutation test [10]. The enrichment results provide input for a manual curation and verification process to identify the most likely biological

process that gives rise to the identified factors. 150

Single-sample analysis
151

Finally, in order to determine the activity of the identified factors in a new unseen 152
sample, one simply solves the same equation with the new samples while keeping the 153
sparse factor coefficients fixed, as depicted in Fig 1E. Importantly, this can be done 154
with the gene expression data only, allowing easy translation of the factors to other 155
datasets. 156

Application to breast and lung cancer
157

We have applied FuncSFA to the breast cancer [1] and lung cancer [2,3] data sets 158
from TCGA. We chose these two cancer types for the following reasons. Breast 159
cancer is arguably the most exhaustively subtyped type of cancer, and hence serves 160
as a very good positive control for FuncSFA [1,12,15,16]. Lung cancer has not been 161
studied so extensively and is at the moment the cancer type resulting in the highest 162
number of deaths per year, so there is a large unmet need for the discovery of new 163
biomarkers which could open up new treatment modalities. From a technical 164
perspective, both cancer types have a large number of samples which ensures that 165
the parameter estimates produced by FuncSFA are reliable. 166

    For both datasets, we used three data types. First, we included DNA copy number 167
at 162 (breast) or 213 (lung) frequently aberrated loci from SNP6 arrays as identified 168
by RUBIC [17]. This DNA copy number data set is clearly important as it captures 169
many copy number events that may have a functional role in oncogenesis. Second, 170
we included protein expression and modification by RPPA with 195 (breast) or 216 171
(lung) protein epitopes. These measurements capture the activity of key signaling 172
events in pathways which play a central role in many cancer types, including those 173
under study here. Third, we included RNA expression of the 1000 most variable 174
genes as measured by RNA-seq. We selected the 1000 most variable genes to 175
reduce noise in the model and to reduce the complexity of the model. We included 176
RNA expression as it is arguably the most comprehensive data type, capturing the 177
downstream effects of many upstream events such as signaling pathway activity or 178
activation of cancer genes by, for example, genetic events. Although FuncSFA allows 179
the inclusion of many data types, we have not included mutation data. The binary 180
nature of this data does not directly fit the factor analysis model we employed (see 181
Discussion and Methods). 182

    For both breast and lung cancer, we employed the abovementioned data types 183
and applied FuncSFA. We specifically set out to find the ten strongest factors. We 184
chose this number of factors based on the following argumentation. First, when 185
increasing the number of factors upto at least to twenty, one can discover more 186
detailed processes showing activity (provided that a sufficiently large sample size is 187
employed); and, as the number of factors increases, the strongest factors remain the 188
same (S2 Fig). Second, most subtyping approaches that have been applied to date 189
revealed ten or fewer subtypes: five intrinsic subtypes in breast cancer [18,19] ten 190
IC10 subtypes in breast cancer [12], four consensus subtypes in colorectal cancer [5], 191
four in squamous cell carcinoma [20] and three in lung adenocarcinoma [21]). 192
Therefore, our assumption was that ten factors would be sufficient to capture the 193
strongest (known) factors, while leaving room for the discovery of new biological 194
processes without running the risk of compromising the robustness of the factors 195
being discovered by having a too large factor-to-sample-size ratio. 196

Breast Cancer

We applied FuncSFA to the breast cancer data employing 10 factors based on the arguments given above. We performed functional annotation of the factors, as outlined above and in the methods section, employing, amongst others, the coefficients obtained from the sparse-factor analysis, as well as the gene-set enrichment for pathways and biological processes depicted in S4 Fig and S1 Table. This resulted in the following 10 factors: ER (Estrogen receptor), Normal-like, Basal, HER2, Luminal Proliferative, 8q-gained, Technical-RNA, Technical-RPPA, EMT (Epithelial to mesenchymal transition) and Immune. The strongest sparse-factor analysis coefficients (from the B matrix in Fig 1B) for the three data types are represented in Fig 2. We will first provide some general observations of the results and then provide a detailed description and analysis of each factor.

From the coefficients depicted in Fig 2, we make the following global observations. First, the ER factor is strongly associated with both mRNA and protein expression of ESR1, GATA3, PGR and AR, as expected. Second, the EMT factor shows strong association with THBS2 and COL11A1 expression—the genes identified by Anastassiou and colleagues [22] (see also the more detailed description of the EMT factor below). In addition, the EMT factor shows association with many collagens. Third, the HER2 factor shows the expected strong association with ERBB2 copy number gain and protein upregulation, GRB7 copy number gain as well as EGFR protein upregulation. Fourth, the Basal factor shows strong association with the RNA expression of basal keratins and finally, 8q-gained shows concordant copy number and expression changes for a number of genes located on Chromosome 8q, including SGK3, MYC and TP52INP1.

Fig 3A depicts, per factor, for all ten factors, the amount of explained variation for the three data types: gene expression, copy number and RPPA. Most factors explain variation in each of the data types, with the largest proportion of the variation explained by a given factor mostly being the variation in gene expression. There are a few exceptions. The HER2, Luminal proliferative and 8q-gained factors, explain more variation in copy number data than in RNA expression data. For HER2 and 8q-gained samples this is not surprising as they are clearly copy number driven (Fig 2 and 4). Most strikingly, the Techical-RPPA factor explains a very large proportion of the variation in the RPPA data. However, as the name indicates this factor most likely captures technical variation in the RPPA data, as this variation is not reflected in the other data types, and as the gene-set enrichment analysis does not reveal a clear functional enrichment associated with this factor. Similarly, the Technical-RNA factor explains technical variation in the RNAseq data. The ER factor is unique in the sense that it is the single factor that explains most variation in RNA expression and RPPA data (apart from the Technical-RPPA factor). This is not unexpected as ER signaling arguably plays the most dominant role in breast cancer. In the remainder of this section, we will discuss each of the identified factors in greater detail.

Intrinsic subtypes are covered by factors

The intrinsic subtypes proposed by Perou and Sørlie represent one of the earliest and most widely used subtypings of breast cancer [16, 18, 19]. This discrete classification of breast cancer in five subtypes (Her2-Enriched, Basal, Luminal A, Luminal B and Normal-like) is typically performed by applying a nearest centroid classifier to the RNA expression profile of the sample to be subtyped. To perform this classification based on mRNA expression, Parker and colleagues [4] developed a a 50 gene signature, the so-called PAM50. As the PAM50 subtyping is widely acknowledged as a gold standard in breast cancer subtyping, we set out to check whether the variation captured by the

Fig 2. The strongest sparse-factor analysis coefficients for the breast cancer data set for each of the three data types and all ten factors.

Fig 3. Sparse-factor analysis on the TCGA breast cancer data set. A: Explained variation per data type and factor. B: The top-left panel shows a t-SNE map of the tumors with the different colors showing PAM50 subtypes. The remaining panels show the samples in the same positions as the PAM50 map, but colored according to the value of the represented factor in each sample.

Fig 4. Copy number and factors in the TCGA breast cancer dataset. Normalized coefficients representing the contribution of DNA copy number abberations to the factors. Specifically, the coefficients represent the contribution of recurrently gained (left) or lost (right) copy number regions identified by RUBIC to the factors represented in the rows. Recurrently aberrated copy number regions are annotated with chromosomal bands or putative driver genes in the region.

PAM50 subtyping is also recapitulated by the FuncSFA factors. To this end we applied the PAM50 subtyping to the breast cancer cohort and compared the PAM50 subtyping to the variation captured by the FuncSFA factors. Fig 3B.1 depicts a t-SNE map of the breast cancer samples with the PAM50 subtype assignment indicated by the colors. Four FuncSFA factors capture the variation in the PAM50 subtypes.

First, the ER factor is associated with both mRNA and protein expression of ESR1, AR and PR (Fig 2), is enriched for gene signatures of ESR1 expression and the luminal subtype of breast cancer, which, in turn, is characterized by ESR1 overexpression (S4 Fig and S1 Table). This suggests that this factor represents the continuous variation in ESR1 expression in breast cancer, which is typically dichotomized to define ER+ and ER- tumors. This is confirmed by comparing the PAM50 subtypes with the ER factor. Indeed, the ER factor is strongly associated with the ER+ (Luminal A, Luminal B and Normal-like) and ER- (Basal, HER2-enriched) subtypes. Specifically, classification into the ER+ and ER- classes based on the ER factor results in an AUC of 0.98. The strong association between the ER factor and the ER+/ER- PAM50 subtyping is also strikingly visible in Fig 3B.2, where the same sample positions as in Fig 3B.1 are maintained whilst the samples are colored according to the value of the ER factor.

Second, the HER2 factor shows large SFA coefficients for ERBB2 protein expression and copy number gain (Fig 2), enrichment for signatures of the HER2-enriched PAM50 subtype and ERBB2 amplification signatures in the gene-set enrichment analysis. Importantly, this factor is strongly associated with the amplicon on Chromosome 17 harboring ERBB2 and GRB7, as evidenced by the large coefficients identified for this amplicon in the factor analysis (Fig 2 and 4). Taken together, this suggest that the HER2-factor indeed identifies the HER2+ tumors. This is confirmed by the strong association of this factor with the HER2-enriched PAM50 subtype (Fig 3B.3, AUC=0.96) and by the high weights of the HER2/GRB7 locus on DNA copy number (Fig 4).

Third, the Luminal-Proliferative factor shows enrichment for signatures representing the cell cycle, proliferation and high-grade tumors in the gene-set enrichment analysis. This factor shows strong association with the PAM50 Luminal A and Luminal B subtypes (AUC of 0.78), with the factor being predominantly low in Luminal A and high in Luminal B (Fig 3B.4). Notice that, in contrast to the ER and HER2 factors which are continuous but show bimodality, this factor is more continuous in nature (S3 Fig). In addition, the copy number weights associated with this factor shown in Fig 4, show the 8q and 17q gains and 8p losses characteristic of some Luminal B tumors [12].

Fourth, gene-set enrichment analysis and ROC-curves show that the Normal-like factor is associated with PAM50 normal-like tumors (AUC=0.92) and tumors with a lobular pathology (AUC=0.84). Finally, the Basal factor is associated with expression of the basal keratins KRT5, KRT6B, KRT14 and KRT17 (Fig 2) and appears only within the triple-negative tumors (S6 Fig). The Basal factor is very different from the Basal subtype from the intrinsic subtypes. The latter is mostly present in triple negative samples, and is not directly related to basal cells as the basal factor we have identified here based on the expression of basal keratins. Taken together, we can conclude that the ER, HER2, Luminal Proliferative and Normal-like factors collective capture most of the variation in the well-known breast cancer intrinsic subtypes.

EMT factor

The gene-set enrichment analysis revealed an association between the EMT factor and EMT and the extracellular matrix (S1 Table and S4 Fig). EMT is a process frequently associated with cancer [23] and it involves multiple regulators, including SNAI1, SNAI2, TWIST1 and ZEB1 as well as their targets [24]. Only SNAI1 is on the RPPA array, and its protein expression is not associated with this factor. More generally, quite a number of gene expression signatures have been developed to detect various forms of EMT (For example, see [25]). Based on the gene-set enrichment analysis, the strongest association of the EMT factor is with a consensus EMT signature proposed by Anastassiou and colleagues [22]. They compiled a pan-cancer EMT signature from multiple public data sets by comparing metastatic with non-metastatic tumors [26]. As an EMT-like expression profile can be associated with stromal contamination in a tumor sample, Anastassiou and colleagues profiled human tumors from a PDX model on a microarray with species-specific probes. This enabled the removal of the mouse stromal signal and revealed that the signature is tumor specific. A strong correlation between the EMT factor and the sum of the two most important genes in this signature (Fig 5A, ρ=0.89) confirms the interpretation of this factor as capturing the specific type of EMT modeled by the Anastassiou signature. While the claudin-low and metaplastic subtypes of breast cancer have been assiocated with EMT [27], larger studies failed to confirm the presence of the claudin-low subtype [1]. Possibly, this factor captures the biology leading the discovery of these subtypes. The t-SNE map of Fig 3B.5 reveals that this factor is equally strongly represented in all intrinsic subtypes.

Immune factor

The Immune factor shows enrichment for Interferon-Alpha Response and other immune related signatures (S1 Table and S1 Table). In order to shed further light on this factor we employed publicly available Cibersort scores [28] to estimate the immune cell fractions in all the breast cancer samples, based on their RNA expression profiles [29]. Briefly, Cibersort employs RNA expression profiles of 22 pure immune

Fig 5. Additional factors add detail over well-known subtypes of breast cancer. A: Scatterplot of the EMT factor versus the sum of the RNA expression of COL11A1 and THBS2 (CPM: count per million). B: Pearson correlation (ρ) between the factors and cell type fractions. Only significant correlations (p<0.05, |ρ|>0.2) are shown.

cell types to perform an in silico decomposition of the RNA expression profile of a sample. The resulting output provides an estimate of the relative abundance of the different immune cell types in the sample being decomposed. In other words, for every immune cell type, we obtain a profile across all the breast cancer samples representing the estimated fraction of that cell type in each sample. We then computed the Pearson correlation (ρ) of these immune cell type profiles with each of the factors. Fig 5B depicts the significant correlations, i.e. only factor and immune cell profile correlations where p<0.05 and |ρ|>0.2. The Technical RPPA, ER, Normal-Like and Immune factor all show more than one significant correlation with an immune cell type. Whereas the ER factor (resting memory DC4 T cells, resting mast cells) and the normal-like factor (γδ T-cells, resting mast cells, naive B cells) both display a more indolent immune profile, the Immune factor clearly shows high positive correlation with active immune cell types and negative correlation with resting immune cells. While the immune factor is quite uniformly present in all breast cancer samples, there is an enrichment for very large values in the Basal-like samples which corresponds with previous reports of CD274 (PD-L1) expression patterns [30], and Cibersort inferred immune infiltration patters [31]. Recently, four distinct consensus immune signatures were identified [32], relating to T-cell and/or B-cells, interferons, TGFB1, and core serum response.

Taken together, we have identified 10 factors in breast cancer. A number of these factors (ER, Normal-like, HER2, Basal and Luminal-Proliferative) capture the subtype variation previously described in the Intrinsic subtypes. This serves as a positive control of FuncSFA as it illustrates that it can capture known biological variation. More importantly, FuncSFA also captured additional variation not represented in the Intrinsic subtypes. Specifically, we identified the 8q-gained factor which is associated with gains of the q arm of Chromosome 8 and loss of the Chromosome 16q23-24

region, as is clear from the DNA copy number weights in Fig 4. In addition, we ₃₄₉ identified an EMT and Immune factor that may lead to a better understanding of the ₃₅₀ processes playing an important role in breast cancer, while also serving as a starting ₃₅₁ point for the development of new treatment approaches. ₃₅₂

Lung Cancer ₃₅₃

Having established the utility of FuncSFA on the breast cancer data, we moved on to ₃₅₄ assess FuncSFA on a less well-characterized tumor type. To this end we applied ₃₅₅ FuncSFA to the lung tumor data from the TCGA in order to further assess its ₃₅₆ usefulness in uncovering new and clinically relevant biology. We merged the TCGA ₃₅₇ lung adenocarcinoma and lung squamous cell carcinoma datasets to obtain a data set ₃₅₈ that is comparable in size to the breast cancer set. We employed the same data types ₃₅₉ (gene expression, RPPA and copy number) as we employed in the breast cancer ₃₆₀ analysis. As the dataset is of comparable size to the breast cancer dataset, and as ₃₆₁ Wilkerson and collegues defined a total of 7 lung cancer subtypes [20,21] we set out ₃₆₂ to identify 10 factors. This allows the capturing of known variation with some room for ₃₆₃ the discovery of novel subtypes. ₃₆₄

FuncSFA identified 10 lung cancer factors: Adenocarcinoma, Mitochondria, DNA ₃₆₅ Replication, Immune, Infiltrating B-cells, NFE2L2, EMT (Epithelial to mesenchymal ₃₆₆ transition), Translation, BSCC (Basal Squamous Cell Carcinoma) and 8p11-gained. ₃₆₇ S5 Fig and S2 Table list the factors and the processes and pathways that were ₃₆₈ significantly enriched in each of the factors based on the gene-set enrichment ₃₆₉ analysis. The strongest SFA coefficients (the B matrix in Fig 1) for the three data types ₃₇₀ are represented in Fig 6. As before we will first provide some general observations of ₃₇₁ the results and then provide a detailed description and analysis of each factor. ₃₇₂

The coefficients depicted in Fig 6 reveal the following. First, as before the EMT ₃₇₃ factor shows strong association with the Anastassiou signature genes (THBS2 and ₃₇₄ COL11A1) as well as a number of collagens. Second, the Mitochondrial factor has ₃₇₅ large SFA coefficients for genes encoded on the mitochondrial DNA while the ₃₇₆ Translation factor shows strong association with genes encoding ribosomal proteins. ₃₇₇ Third, the BSCC and 8p11-gained factors show large RPPA coefficients, with a ₃₇₈ concentration of lowly expressed proteins in the PI3K pathway. However, they do ₃₇₉ differ in terms of their RNA expression coefficients, and we will provide more ₃₈₀ elaborate descriptions of these factors below. Fourth, the Immune factor is ₃₈₁ characterized by large RNA expression coefficients of immunoglobulins. Finally, the ₃₈₂ Adenocarcinoma factor shows association with the RNA expression of a number of ₃₈₃ keratins, negative coefficients for both copy number and RNA expression of SOX2 ₃₈₄ and positive association with both RNA expression and copy number of NKX2-1 (also ₃₈₅ known as TTF1). ₃₈₆

Fig 7A depicts, per factor, for all ten factors, the amount of explained variation for ₃₈₇ the three data types: gene expression, copy number and RPPA. As in breast cancer, ₃₈₈ most factors explain variation in each of the data types, with the largest proportion of ₃₈₉ the variation explained by a given factor mostly being the variation in RNA expression ₃₉₀ with the exceptions being the 8p11-gained and BSCC factors where the largest ₃₉₁ portion of the variance explained by these factors is the variation in the RPPA data. It ₃₉₂ is also noteworthy that these two factors explain a significantly larger portion (≈0.32) ₃₉₃ of the variation in the RPPA data, compared to the factors that explain the largest ₃₉₄ portion of the RNA expression variation (≈0.19). The factors are negatively correlated ₃₉₅ and have coefficients of similar magnitude and equal sign. In other words, in a given ₃₉₆ sample the sum of these factors is small. (S8 Fig). ₃₉₇

In the remainder of this section, we will first discuss the Wilkerson subtypes that ₃₉₈

Fig 6. The strongest SFA coefficients for the lung cancer data set for each of the three data types and all ten factors.

Fig 7. Sparse-factor analysis on the TCGA lung cancer dataset. A: Explained variance per data type and factor. B: B.1 shows the t-SNE map of all lung cancer samples with red denoting the Adenocarcinoma samples and blue the Squamous Cell Carcinomas. With the samples in the same positions as in B.1, B.2 depicts the subtyping as proposed by Wilkerson and collegues [20, 21], B.3 depicts the values of the Adeno factor per sample, B.4 the values of the EMT factor, B.5 the values of the Immune factor and B.6 the value of the NFE2L2 factor. (LPr: Luminal Proliferative, TechR: Technical-RNA, TechP: Technical-RPPA, Imm: Immune, NLi: Normal-like, Bas: Basal, 8q: 8q-gained)

have been proposed for lung cancer. Then we will discuss each of the identified factors in greater detail.

Fig 8. Mutations and immune infiltration in lung cancer. A: Mann-Whithney U statistic of the factor values between sample with and without a mutation in a gene divided by the product of the number of samples in each group. This statistic can be interpreted as an AUC [33]. Only significant (p<0.05) values are shown. B: Pearson correlation (ρ) between the factors and cell type fractions. Only significant correlations (p<0.05, |ρ|>0.2) are shown. C: Mutations in genes of the NFE2L2 pathway and STK11 (black: sample is mutated in this gene).

The Wilkerson molecular subtypes of lung cancer

The molecular subtypes of lung adenocarcinoma and lung squamous carcinoma were defined by Wilkerson and colleagues [20, 21], and also found to be present on the larger TCGA datasets [2, 3]. Fig 7B.2 depicts the Wilkerson subtypes on top of the t-SNE plot derived from the SFA factors. Here every tumor is colored according to its subtype membership. Globally, all SFA factors are associated with these subtypes (Kruskal-Wallis test, $p < 10^{-6}$ for every factor). However, in contrast to breast cancer, it was not possible to perform a one-to-one mapping from the SFA factors to the

Fig 9. Copy number and factors in the TCGA lung cancer dataset. Normalized coefficients representing the contribution of DNA copy number abberations to the factors. Specifically, the coefficients represent the contribution of recurrently gained (left) or lost (right) copy number regions identified by RUBIC to the factors represented in the rows. Recurrently aberrated copy number regions are annotated with chromosomal bands or putative driver genes in the region.

Wilkerson subtypes. This can most likely be explained by the fact that none of the Wilkerson subtypes have (yet) been associated with a driver event, unlike breast cancer where ESR1 and ERBB2 have been identified as strong drivers giving rise to specific subtypes of breast cancer. Rather, it seems that the Wilkerson subtypes are driven by a complex interplay of multiple, heterogeneous biological processes. Under these circumstances, we would not expect to find factors that represent the subtypes directly, but that the factors should capture the underlying biology.

Adenocarinoma and squamus cell carcinoma were subtyped seperately, so we have a set of subtypes for each. The adenocarcinoma subtypes are termed 'terminal respiratory unit' (TRU), 'proximal-inflammatory' (PI) and 'proximal-proliferative' (PP). The TRU subtype is characterized by highly expressed asthma, excretion and surfactant genes, the PP subtype with the overexpression of defense response genes (chemokines) and the PI subtype with high expression of DNA-repair genes [21]. The Wilkerson squamous cell carcinoma subtypes are termed 'Basal', 'Classical', 'Secretory' and 'Primitive'. The Basal subtype has been associated with cell adhesion and epidermal development. Wilkerson and colleagues reported that the classical subtype is related to xenobiotic detoxification, that the Secretory subtype shows association with the expression of immune related genes, NKX2-1 (TTF1), MUC1 and surfactant genes. It is interesting to note that although the secretory subtype is a squamous cell carcinoma subtype, it shows intermediate scores in the Adenocarcinoma factor, which is consistent with the fact that NKX2-1 is both expressed in the Secretory subtype and highly expressed in adenocarcinomas. Finally, the Primitive subtype has previously been associated with proliferation and DNA processing and repair.

Adenocarcinoma and squamous cell carcinoma

The differences between adenocarcinoma and squamous cell carcinoma are captured by the Adenocarcinoma factor, which is high in adenocarcinomas and low in squamous cell carcinomas. Interestingly, in the primitive subtype of squamous cell carcinoma, tumors with high and low values for this factor exist. In addition, copy number is dominated by a particularly strong association with the gain of Chromosome 3q26-29, implying that the adenocarcinomas show absence of gains while squamous cell carcinomas carry this gain (Fig 9). We also observe the known associations between this factor and mutations (Fig 8A). As expected, the Adenocarcinoma factor is high in all the Wilkerson adenocarcinoma subtypes, as compared to the squamous cell carcinoma subtypes. More specifically, the Adenocarcinoma factor is very low in the Basal and Classical subtypes but higher in the Secretory and Primitive subtypes.

Two factors strongly associated with RPPA: BSCC and 8p11 gained

The largest fraction of the variance explained by the BSCC and 8p11-gained factors is associated with the RPPA data. In fact, the amount of variation explained in the RPPA data by these two factors is the highest across all factors and all data types. The 18p11-gained factor is associated with increased gain of the genes NDS3, LETM2 and, FGFR1, which are located on the 8p11 chromosomal band (Fig 9]), and shows high scores in the PI subtype. As expected, the BSCC factor shows, in general, higher scores within the Basal subtype as compared to other squamous cell carcinoma subtypes (Fig 8). Specifically, the BSCC factor shows intermediately strong association with the basal subtype of squamous cell carcinoma (AUC=0.59).

DNA replication factor

The DNA replication factor captures two biological processes. Pathway analysis reveals a positive association of this factor with pathways involved in DNA replication, implying that genes annotated to these pathways are enriched in samples with high levels of this factor. According to the Cibersort analysis, this factor is positively correlated with signatures representing activated memory CD4 T-cells and negatively correlation with signatures representing resting mast and resting dendritic cells (Fig 8B). Conversely, this factor shows negative association with the protein expression of the transcription factor NKX2-1. As expected, NKX2-1 targets (SCGB3A2 and AQP4) are amongst the top-ranking genes in terms of the magnitude of their negative RNA expression coefficients. Other genes with very large negative RNA expression coefficients include the surfactant proteins (SFTPA1, SFTPA2, SFTPB, SFTPC, SFTPD) which are co-expressed with NKX2-1 in the alveolar cells of the lung [34].

The DNA replication factor has low scores in tumors of the TRU subtype, which is consistent with the association of this subtype with high expression of surfactant proteins. Compared to the TRU subtype, the DNA-replication factor shows high scores in the PI subtype, but not at the same level as in the Primitive and Classical squamous cell carcinoma subtypes. Since the Secretory subtype shows association with surfactant proteins, the DNA-replication factor shows, as expected, low scores for the tumors of this subtype (Fig 8).

Immune factors: Immune and Infiltrating B-cells

FuncSFA identifies two immune related factors: Immune and Infiltrating B-cells (IB-cells). Pathway analysis of the Immune factor shows enrichment for interferon

associated signatures and T-cell pathways. This association is confirmed by high RNA expression coefficients of immunoglobulins, members of the histocompatibility complex (HLA) and members of the complement system (Fig 6). According to the Cibersort analysis, the Immune factor correlates with the signatures representing T-follicular helper cells and activated dendritic cells while it negatively correlates with signatures of γδ T-cells and M1 macrophages (Fig 8B). In addition, this factor is associated with an increased loss of 9p23-21. The Infiltrating B-cells factor also shows strong association with interferon associated signatures and T-cell pathways in the gene-set enrichment analysis. However, this association is weaker than for the Immune factor. When considering the RNA expression coefficients of this factor we see large coefficients for immunoglobulins, but in contrast to the Immune factor, neither the HLA nor the complement system is represented. This factor has a strong positive correlation with plasma B-cells (Fig 8B) and is associated with a higher mutation rate of STK11 (Fig 8A).

As the PP subtype shows overexpression of defense response genes (chemokines), we observe, consistent with this finding, that the infiltrating B-cell and Immune factors score highly in this subtype (Fig 7B.5 and Fig 7B.7). In contrast, the Infiltrating B-cell factor scores lowly in the TRU and PI subtypes. The Immune factor shows high scores in in the TRU, Classical and Secretory subtypes. Interestingly, the Basal subtype shows high levels of the Infiltrating B-cell factor, but no distinctive pattern for the Immune factor.

Mitochrondial, Translation and EMT factors

We identified a number of factors with a clear functional profile, but where additional research is required to understand their role and relevance in lung cancer. First, the Mitochondrial factor associated with genes encoded on the mitochondrial DNA and GAPDH protein expression (Fig 6). Second, the Translation factor which shows enrichment for translation pathways and signatures of the cell cycle (S4 Fig). This association is also strongly supported by the large coefficients of genes encoding ribosomal proteins (Fig 6). Finally, as in breast cancer we also identified an EMT factor in lung. As in breast, the EMT lung factor is strongly correlated with the Anastassiou signature factor across the samples (Pearson correlation ρ=0.73). Regarding the Wilkerson subtypes, the TRU subtype shows low scores for the EMT and Translation factors while the Translation factor is high in the Primitive subtype.

NFE2L2

The NFE2L2 factor shows NFE2L2 targets on the gene-set enrichment analysis, suggesting this factor captures the activation of NFE2L2. This conclusion is confirmed by a strong association of this factor with mutations in NFE2L2 and in its inhibitor KEAP1 (Fig 8A). Interestingly, the factor is depleted for EGFR mutations, altough EGFR activates the NFE2L2 pathway by inhibiting KEAP1 [35]. Still, these three genes in the NFE2L2 pathway show a mutually exclusive mutation pattern (Fig 8C), suggesting that they independently activate the NFE2L2 pathway effecting the common dowstream gene expression changes captured by this factor. So, this factor could be a readout of NFE2L2 pathway activation, which in some conditions has been suggested to increase resistance to EGFR inhibitors

In addition to the clear NFE2L2 association, this factor also shows large values for the coefficients for genes related to xenobiotic detoxification, such as CES1, CYP4F3 and CYP4F11. Hence, it is not surprising that the NFE2L2 factor shows high scores in the Classical subtype, which is also associated with this process.

In summary, we have identified factors that do show overall association, but not very clear one-to-one correspondence with the previously defined Wilkerson subtypes. Our factors do capture the major lung cancer subtypes (Adenocarcinoma and Squamous cell carcinoma), and identified processes that occur in both of these types. These include epithelial to mesenchymal transition, two immune processes, and, most interestingly, a factor that reports the activity of the NFE2L2 pathway. The latter may have interesting therapeutic consequences.

Discussion

In both breast and lung cancer the sparse-factor analysis was able to find known biology back. Additionally, we find factors that capture biology that traditional clustering methods have not been able to find, such as the Immune, EMT, and infiltrating B-cells factors. Our results illustrate the need for continuous factors as opposed to discrete clusters. For example, the Luminal-Proliferative factor shows that the distinction between luminal A and B is not discrete. Also the need for multiple factors per sample is illustrated by our results. For example, the Immune and EMT factors in breast cancer are active across subtypes. Finally, our results show that a complete molecular characterization requires multiple data types. Although all non-technical factors are represented in the RNA expression, the 8q-gained and Luminal Proliferative factors in breast cancer are mostly copy number driven. Also, the protein expression contributes to key factors such as the ER and HER2 factors.

Interestingly we have found the same cancer cell specific EMT factor in both lung and breast cancer. This does not necessarily mean that tumours with high levels of this factor have a large proportion of mesenchymal tumor cells. More likely, this factor represents the activation of a transcriptional program that may eventually lead to a transition to the mesenchymal phenotype. Therefore, the tumors with high levels of this factor can have a higher propensity to undergo EMT. This can potentially be resolved by applying the method pan-cancer, which could give an indication how far along the EMT transition the cells in these tumors really are.

Several factors could be implicated in a tumors' response to treatment. The NFE2L2 factor might be indicative of DMF treatment, which targets the NFE2L2 pathway and can inhibit carcinogenesis [36]. Because the factor measures activation of the transcription factor through its downstream targets, it could be more predictive than only looking at mutations. This hypothesis could be tested on patient derived xenograft models. The Immune factor in breast cancer and the Immune, DNA replication and B-cell infiltration factors in lung cancer are possibly predictive for response to immune checkpoint inhibitors. If the immune system is already highly active, a clinically relevant response might be easier to achieve. These hypotheses can be tested by obtaining RNA expression data from pre-treatment biopsies of patients treated with immune checkpoint inhibitors.

In the lung cancer data set we found two highly correlated factors. In principle, the algorithm should find almost orthogonal factors. Orthogonal factors explain the most variance, but the sparsity penalties prevent full orthogonality. There might be two explanations of the two highly correlated factors. First, the optimal sparsity penalty might be different per factor, where one biological process involves thousands of genes, and another only a few. Second, there might be non-linear interactions between genes such that a linear model cannot explain the activity of a biological process with single factor. Two factors involving the same genes could model a non-linear biological process together.

In this study we used only the samples which have all data types available. With

increasing samples sizes and an increasing number of data types this leads to ignoring larger parts of the available data. The EM framework quite naturally allows for missing data, so this might be an interesting future extension.

In addition to the data types we used here, other molecular data types that could be included are DNA methylation and miRNA expression data. As the mutation data includes driver events it would be desirable to include it as well. This would require transforming the mutation data such that it has approximately Gaussian error. One way to do this is by smoothing mutations over an interaction network [37]. The inclusion of higher level tumor phenotypes, such as features obtained from MR imaging or pathology slides, could guide the method towards finding biological processes that lead to clinically relevant differences.

We have shown the method is able to find biologically processes that are active across otherwise very different tumors, such the ER+ and ER- subtypes in breast cancer. Applying this method to a pan-cancer cohort might find biological processes that are activated in a large number of tumor types and provide insight into how tumors of different origin relate to each other.

In summary, we have shown FuncSFA is able to integrate at least three data types. It finds factors as opposed to clusters. Our results illustrate the advantage of factors over clusters with several biologically or clinically relevant examples. The identified factors represent the heterogeneity both within and between cancer types, and represent the activation of biological processes in a patient specific manner. Considering the more complete and fine-grained characterization they allow, these factors could benefit the personalized treatment of cancer.

Methods

The FuncSFA method consists of two steps. In the first step, a sparse-factor analysis integrates multiple data types into a small number of factors. In the second step, the factors are linked to existing knowledge of biology by doing a gene-set enrichment analysis.

Sparse-Factor Analysis Model

The data of a single data type i with N samples and n_i features is represented in a $n_i \times N$ data matrix X_i. The data matrices of t data types are stacked together into a $n \times N$ data matrix $X = [X_1^T, \ldots, X_t^T]^T$. This data matrix is factorized into a $n \times k$ coefficient matrix B and $k \times N$ factor matrix Z, where the number of factors k is much smaller than the total number of features n.

We want the coefficients B to be sparse for three reasons. First, a factor which involves a small number of genes is more interpretable than a factor involving almost all genes. Second, the biological processes we hope the factors will represent involve only a fraction of all genes. Third, by imposing higher sparsity on a single data type we can prevent that data type from dominating the factorization.

To achieve this sparse factorization of multiple data types we used the model described previously by Shen and colleagues. [8]. We briefly summarize this probabilistic model here. A sample vector $\mathbf{x}$ with n features is explained by factors $\mathbf{z}$ and coefficients B with Gaussian error

$$\mathbf{x} = B\mathbf{z} + \mathbf{e}, \qquad \mathbf{e} \sim \mathcal{N}(0, \Psi) \text{ i.i.d}$$

where the residual variance Ψ is zero off the diagonal and equal for features of the

same data type. When the factors are taken to be normally distributed $$\mathbf{z} \sim \mathcal{N}(0, I) \text{ i.i.d}$$

the complete data log-likelihood is:

$$\ell(B, \Psi; X, Z) = -\frac{1}{2}\text{trace}((X - BZ)^T \Psi^{-1}(X - BZ)) - \frac{n}{2}\log(\det(\Psi)) - \frac{1}{2}\text{trace}(ZZ^T)$$

where $X = [\mathbf{x}_1, ..., \mathbf{x}_N]$ and $Z = [\mathbf{z}_1, ..., \mathbf{z}_N]$. The coefficients are made sparse by a elastic net penalty:

$$\ell_p(B, \Psi; X, Z) = \ell(B, \Psi; X, Z) - \sum_{i=1}^{n} \lambda_{i,1}|(\mathbf{b}_i)|_1 - \sum_{i=1}^{n} \lambda_{i,2}|(\mathbf{b}_i)|_2^2$$

where $B = [\mathbf{b}_1^T, ..., \mathbf{b}_n^T]^T$. The penalties $\lambda_{1,i}$ and $\lambda_{2,i}$ are kept the same for all features in a data type.

Sparse-Factor Analysis Algorithm

To optimize the penalized complete data log-likelihood we use an iterative algorithm, improving over the iCluster2 algorithm [11] at several key points. It consists of the following steps.

1. Initialize the estimated coefficients $\hat{B}$ and residual variance $\hat{\Psi}$ by the loadings of a principal component analysis (PCA).

2. Calculate the expectation $E[Z|X]$ and covariance $E[ZZ^T|X]$ of the factors given the current $\hat{B}$ and $\hat{\Psi}$. In contrast to iCluster2 $E[Z|X]$ and $E[ZZ^T|X]$ are then scaled to have unit variance per factor. This prevents the regularization penalty from forcing large factor values to compensate for small coefficients.

3. Estimation of coefficients is done differently from iCluster2. We use the coordinate descent scheme for solving elastic net as used in glmnet [38]. A coefficient of a single factor i and a single feature j is updated given the current $E[Z|X]$, $\hat{B}$ and $\hat{\Psi}$:

$$\hat{B}_{[i,j]} \leftarrow \frac{\text{S}(\hat{B}_{[i,j]} + E[Z|X]_{[j,\cdot]}X_{[i,\cdot]}^T - \hat{B}_{[i,\cdot]}\,\text{E}[ZZ^T|X]_{[\cdot,j]}, \lambda_{i,1})}{1 + \lambda_{i,2}}$$

where S is the soft-tresholding operator $\text{S}(b, l_1) = \text{sign}(b)(|b| - l_1)_+$. In a single iteration of the main algorithm the coefficients are updated once in sequential order.

4. Residual variance is estimated per data type:

$$\hat{\Psi}_i \leftarrow \frac{1}{n_i N}(\text{trace}(B_i E[Z|X]X_i) - \text{trace}(X_i X_i^T))$$

for data type i.

5. Repeat from step 2 until convergence, with a maximum of 5.000 iterations. We consider the algorithm converged when the average absolute change in reconstruction error over the last 10 iterations is smaller than 10^{-6}.

On the TCGA breast cancer dataset, this algorithm converged faster and to better solutions than the iCluster2 (S1 Fig). The software is available from `https://github.com/NKI-CCB/funcsfa`.

Parameter Selection ₆₅₁

To find optimal parameters $\lambda_{i,1}$, $\lambda_{i,2}$ we reparameterize. ₆₅₂

$$\lambda_{i,1} = \alpha l_{\mathrm{datatype(i)}}$$
$$\lambda_{i,2} = 10^{-6} + (1 - \alpha)l_{\mathrm{datatype(i)}}$$

where $\mathrm{datatype(i)}$ gives the datatype of feature i. We then do a grid search for the l ₆₅₃
penalties giving the highest Bayesian information criterion (BIC), keeping α fixed at ₆₅₄
0.9. The effective number of parameters of the penalized coefficients, used for ₆₅₅
calculating the BIC, is calculated using the method from Tibshirani and Taylor [39]. ₆₅₆

Pathway Analysis ₆₅₇

We employ a gene-set enrichment analysis tailored to the results of sparse-factor ₆₅₈
analysis. For this we use gene-set enrichment analysis [10] with a modified gene ₆₅₉
ranking method. Gene ranking is done as follows. First the factors Z are regressed on ₆₆₀
the RNA expression data containing all genes. Second we get a correlation-like ₆₆₁
coefficients by dividing the regression coefficients of a gene by the standard deviation ₆₆₂
of that gene. The genes are then ranked per factor based on their correlation-like ₆₆₃
coefficient. We made available the implementation of this methnod in the ggsea ₆₆₄
package: https://github.com/NKI-CCB/ggsea. ₆₆₅

Preprocessing ₆₆₆

RNA expression data from RNA sequencing was preprocessed using ₆₆₇
limma-voom [40]. We selected the 1000 genes with the highest median absolute ₆₆₈
deviation. The mean variance-trend in RNA sequencing data was taken into account ₆₆₉
by multiplying the RNA expression with the precision weights obtained from voom. ₆₇₀
DNA copy number was sampled at regions with recurrent copy number aberrations. ₆₇₁
Recurrently aberrated copy number regions were taken from the RUBIC paper [17] for ₆₇₂
breast cancer or obtained by running RUBIC for the lung cancer. A sample by region ₆₇₃
matrix was obtained by taking the median copy number of all segments of sample ₆₇₄
overlapping with a region. Protein expression data from RPPA was used directly as ₆₇₅
processed by TCGA. To remove differences in variance between data types we ₆₇₆
divided every datatype by its standard deviation. ₆₇₇

Additional methods ₆₇₈

t-SNE [41] was used to summarize the factors in two non-linear dimensions, yielding a ₆₇₉
high-level map the factors and other tumor variables can be projected on. These ₆₈₀
t-SNE maps were calculated using scikit-learn [42]. ₆₈₁

To make coefficients comparable between features, we divide the coefficients of ₆₈₂
feature by the standard deviation of the feature data. ₆₈₃

The explained variance of a factor is calculated by subtracting the summed square ₆₈₄
error of the model with all factors from the model excluding the factor dividing by the ₆₈₅
total sum of squares of the data: ₆₈₆

$$\mathrm{explained\ variance}_{i,t} = \frac{||X_t - B_{t[\cdot,-i]}Z_{[-i,\cdot]}||_F^2 - ||X_t - B_t Z)||_F^2}{||X_t||_F^2}$$

for factor i and data type t where $B_{t[\cdot,-i]}$ and $Z_{[-i,\cdot]}$ are the coefficients and factors ₆₈₇
excluding the factor i. The squared Frobenius norm $|| \cdot ||_F^2$ is the sum of squares of the ₆₈₈
elements of a matrix. ₆₈₉

Supporting information

S1 Table.   Gene set enrichment of all 10 factors in breast cancer.

S2 Table.   Gene set enrichment of all 10 factors in lung cancer.

S1 Fig.   Convergence of iCluster, iCluster2 and sparse-factor analysis.

S2 Fig.   Correlation between the factors of the best solution with a number of factors and the best solution with one factor more.

S3 Fig.   Histograms of factor values.

S4 Fig.   Heatmap of GSEA Normalized Enrichment Statistic (Breast)

S5 Fig.   Heatmap of GSEA Normalized Enrichment Statistic (Lung)

S6 Fig.   t-SNE maps of breast cancer

S7 Fig.   t-SNE maps of lung cancer

S8 Fig.   Scatterplot of coefficients and values of RPPA technical factors in lung.

Acknowledgments

This research is part of the STW Perspectief programme Population Imaging Genetics (ImaGene) and supported by the Dutch Technology Foundation STW, which is part of the Netherlands Organisation for Scientific Research (NWO), and partly funded by Ministry of Economic Affairs.

References

1. The Cancer Genome Atlas Research Network. Comprehensive molecular portraits of human breast tumours. Nature. 2012;490(7418):61–70. doi:10.1038/nature11412.

2. The Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. Nature. 2014;511(7511):543–550. doi:10.1038/nature13385.

3. The Cancer Genome Atlas Research Network. Comprehensive genomic characterization of squamous cell lung cancers. Nature. 2012;489(7417):519–525. doi:10.1038/nature11404.

4. Parker JS, Mullins M, Cheang MCU, Leung S, Voduc D, Vickery T, et al. Supervised risk predictor of breast cancer based on intrinsic subtypes. Journal of clinical oncology: official journal of the American Society of Clinical Oncology. 2009;27(8):1160–1167. doi:10.1200/JCO.2008.18.1370.

5. Guinney J, Dienstmann R, Wang X, Reyniès A, Schlicker A, Soneson C, et al. The consensus molecular subtypes of colorectal cancer. Nature Medicine. 2015;21(11):1350–1356. doi:10.1038/nm.3967.

6. Lock EF, Dunson DB. Bayesian consensus clustering. Bioinformatics. 2013;29(20):2610–2616. doi:10.1093/bioinformatics/btt425.

7. Yuan Y, Savage RS, Markowetz F. Patient-Specific Data Fusion Defines Prognostic Cancer Subtypes. PLoS Computational Biology. 2011;7(10):e1002227. doi:10.1371/journal.pcbi.1002227.

8. Shen R, Olshen AB, Ladanyi M. Integrative clustering of multiple genomic data types using a joint latent variable model with application to breast and lung cancer subtype analysis. Bioinformatics. 2009;25(22):2906–2912. doi:10.1093/bioinformatics/btp543.

9. Vaske CJ, Benz SC, Sanborn JZ, Earl D, Szeto C, Zhu J, et al. Inference of patient-specific pathway activities from multi-dimensional cancer genomics data using PARADIGM. Bioinformatics. 2010;26(12):i237–i245. doi:10.1093/bioinformatics/btq182.

10. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proceedings of the National Academy of Sciences of the United States of America. 2005;102(43):15545–15550. doi:10.1073/pnas.0506580102.

11. Shen R, Wang S, Mo Q. Sparse integrative clustering of multiple omics data sets. The Annals of Applied Statistics. 2013;7(1):269–294. doi:10.1214/12-AOAS578.

12. Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. Nature. 2012;486(7403):346–352. doi:10.1038/nature10983.

13. Tipping ME, Bishop CM. Probabilistic principal component analysis. Journal of the Royal Statistical Society, Series B. 1999;61(3):611–621. doi:10.1111/1467-9868.00196.

14. Zou H, Hastie T. Regularization and variable selection via the elastic net. Journal of the Royal Statistical Society: Series B (Statistical Methodology). 2005;67(2):301–320. doi:10.1111/j.1467-9868.2005.00503.x.

15. Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, et al. Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. The Journal of Clinical Investigation. 2011;121(7):2750–2767. doi:10.1172/JCI45014.

16. Russnes HG, Lingjærde OC, Børresen-Dale AL, Caldas C. Breast Cancer Molecular Stratification. The American Journal of Pathology. 2017;doi:10.1016/j.ajpath.2017.04.022.

17. Dyk E, Hoogstraat M, Hoeve J, Reinders MJT, Wessels LFA. RUBIC identifies driver genes by detecting recurrent DNA copy number breaks. Nature Communications. 2016;7:12159. doi:10.1038/ncomms12159.

18. Perou CM, Sørlie T, Eisen MB, Rijn Mvd, Jeffrey SS, Rees CA, et al. Molecular portraits of human breast tumours. Nature. 2000;406(6797):747–752. doi:10.1038/35021093.

19. Sørlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. Proceedings of the National Academy of Sciences. 2001;98(19):10869–10874. doi:10.1073/pnas.191367098.

20. Wilkerson MD, Yin X, Hoadley KA, Liu Y, Hayward MC, Cabanski CR, et al. Lung squamous cell carcinoma mRNA expression subtypes are reproducible, clinically-important and correspond to different normal cell types. Clinical Cancer Research. 2010; p. clincanres.0199.2010. doi:10.1158/1078-0432.CCR-10-0199.

21. Wilkerson MD, Yin X, Walter V, Zhao N, Cabanski CR, Hayward MC, et al. Differential Pathogenesis of Lung Adenocarcinoma Subtypes Involving Sequence Mutations, Copy Number, Chromosomal Instability, and Methylation. PLOS ONE. 2012;7(5):e36530. doi:10.1371/journal.pone.0036530.

22. Anastassiou D, Rumjantseva V, Cheng W, Huang J, Canoll PD, Yamashiro DJ, et al. Human cancer cells express Slug-based epithelial-mesenchymal transition gene expression signature obtained in vivo. BMC cancer. 2011;11(1):529. doi:10.1186/1471-2407-11-529.

23. Thiery JP. Epithelial-mesenchymal transitions in tumour progression. Nature reviews Cancer. 2002;2(6):442–454. doi:10.1038/nrc822.

24. Lamouille S, Xu J, Derynck R. Molecular mechanisms of epithelial–mesenchymal transition. Nature reviews Molecular cell biology. 2014;15(3):178–196. doi:10.1038/nrm3758.

25. Loboda A, Nebozhyn MV, Watters JW, Buser CA, Shaw PM, Huang PS, et al. EMT is the dominant program in human colon cancer., EMT is the dominant program in human colon cancer. BMC medical genomics, BMC Medical Genomics. 2011;4, 4:9, 9–9. doi:10.1186/1755-8794-4-9, 10.1186/1755-8794-4-9.

26. Kim H, Watkinson J, Varadan V, Anastassiou D. Multi-cancer computational analysis reveals invasion-associated variant of desmoplastic reaction involving INHBA, THBS2 and COL11A1. BMC Medical Genomics. 2010;3:51. doi:10.1186/1755-8794-3-51.

27. Taube JH, Herschkowitz JI, Komurov K, Zhou AY, Gupta S, Yang J, et al. Core epithelial-to-mesenchymal transition interactome gene-expression signature is associated with claudin-low and metaplastic breast cancer subtypes. Proceedings of the National Academy of Sciences. 2010;107(35):15449–15454. doi:10.1073/pnas.1004900107.

28. Charoentong P, Finotello F, Angelova M, Mayer C, Efremova M, Rieder D, et al. Pan-cancer Immunogenomic Analyses Reveal Genotype-Immunophenotype Relationships and Predictors of Response to Checkpoint Blockade. Cell Reports. 2017;18(1):248–262. doi:10.1016/j.celrep.2016.12.019.

29. Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, et al. Robust enumeration of cell subsets from tissue expression profiles. Nature Methods. 2015;12(5):453–457. doi:10.1038/nmeth.3337.

30. Ali HR, Glont SE, Blows FM, Provenzano E, Dawson SJ, Liu B, et al. PD-L1 protein expression in breast cancer is rare, enriched in basal-like tumours and associated with infiltrating lymphocytes. Annals of Oncology. 2015;doi:10.1093/annonc/mdv192.

31. Ali HR, Chlon L, Pharoah PDP, Markowetz F, Caldas C. Patterns of Immune Infiltration in Breast Cancer and Their Clinical Implications: A Gene-Expression-Based Retrospective Study. PLOS Medicine. 2016;13(12):e1002194. doi:10.1371/journal.pmed.1002194.

32. Amara D, Wolf DM, van 't Veer L, Esserman L, Campbell M, Yau C. Co-expression modules identified from published immune signatures reveal five distinct immune subtypes in breast cancer. Breast Cancer Research and Treatment. 2017;161(1):41–50. doi:10.1007/s10549-016-4041-3.

33. Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. Radiology. 1982;143(1):29–36. doi:10.1148/radiology.143.1.7063747.

34. Stahlman MT, Gray ME, Whitsett JA. Expression of thyroid transcription factor-1 (TTF-1) in fetal and neonatal human lung. Journal of Histochemistry & Cytochemistry. 1996;44(7):673–678.

35. Yamadori T, Ishii Y, Homma S, Morishima Y, Kurishima K, Itoh K, et al. Molecular mechanisms for the regulation of Nrf2-mediated cell proliferation in non-small-cell lung cancers. Oncogene. 2012;31(45):4768–4777. doi:10.1038/onc.2011.628.

36. Sporn MB, Liby KT. NRF2 and cancer: the good, the bad and the importance of context. Nature Reviews Cancer. 2012;12(8):564–571. doi:10.1038/nrc3278.

37. Hofree M, Shen JP, Carter H, Gross A, Ideker T. Network-based stratification of tumor mutations. Nature methods. 2013;10(11):1108–1115. doi:10.1038/nmeth.2651.

38. Friedman J, Hastie T, Tibshirani R. Regularization paths for generalized linear models via coordinate descent. Journal of statistical software. 2010;33(1):1–22. doi:10.18637/jss.v033.i01.

39. Tibshirani RJ, Taylor J. Degrees of freedom in lasso problems. The Annals of Statistics. 2012;40(2):1198–1232. doi:10.1214/12-AOS1003.

40. Law CW, Chen Y, Shi W, Smyth GK. voom: precision weights unlock linear model analysis tools for RNA-seq read counts. Genome Biology. 2014;15(2):R29. doi:10.1186/gb-2014-15-2-r29.

41. Maaten L, Hinton G. Visualizing Data using t-SNE. Journal of Machine Learning Research. 2008;9(Nov):2579–2605.

42. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn: Machine Learning in Python. Journal of Machine Learning Research. 2011;12:2825−2830.