Nutrient stoichiometry shapes microbial coevolution

Megan L. Larsen[1], Steven W. Wilhelm[2], Jay T. Lennon[*1]

[1]Department of Biology, Indiana University, Bloomington, IN 47405

[2]Department of Microbiology, University of Tennessee, Knoxville, TN 37996

*Correspondence to: lennonj@indiana.edu.

Running title: Nutrient stoichiometry and coevolution

Keywords: bacteria, phage, resources, resistance, arms race, networks

Article type: Letter

Abstract word count:

(original manuscript: 150; revised manuscript: 150)

Main text word count: (original manuscript: 4817; revised manuscript: 5369)

Reference count: (original manuscript: 50; revised manuscript: 52)

Figure count: 5

Table count: 0

Author contributions: MLL, SWW, and JTL designed study; MLL performed research; MLL and JTL analyzed data; MLL, SWW, and JTL wrote paper.

Available code and data. Data and code are available in the public GitHub repository

https://github.com/LennonLab/eco-evo-stoich

ABSTRACT

Coevolution is a force contributing to the generation and maintenance of biodiversity. It

28   is influenced by environmental conditions including the scarcity of essential resources, which

can drive the evolution of defense and virulence traits. We conducted a long-term chemostat

30   experiment where the marine cyanobacterium *Synechococcus* was challenged with a lytic

phage under nitrogen (N) or phosphorus (P) limitation. This manipulation of nutrient

32   stoichiometry altered the stability of host-parasite interactions and the underlying mode of

coevolution. By assessing infectivity with >18,000 pairwise challenges, we documented

34   directional selection for increased phage resistance, consistent with arms-race dynamics

while phage infectivity fluctuated through time, as expected when coevolution is driven by

36   negative frequency-dependent selection. The resulting infection networks were 50 % less

modular under N- versus P-limitation reflecting host-range contraction and asymmetric

38   coevolutionary trajectories. Nutrient stoichiometry affects eco-evolutionary feedbacks in

ways that may alter the dynamics and functioning of environmental and host-associated

40   microbial communities.

 INTRODUCTION

In order to grow and reproduce, organisms must assimilate resources from the environment to meet their nutritional and energetic demands. Ecological stoichiometry is a

theoretical framework that explicitly considers the mass balance of materials and energy in the environment and the individuals that incorporate them (Sterner & Elser 2002). Of the

approximately 25 elements contained in biomass, nitrogen (N) and phosphorus (P) are two of the most limiting and essential nutrients. N and P are needed for the synthesis of major

macromolecules, including nucleic acids, ribosomes, proteins, and cellular membranes that collectively influence an organism's performance. However, the degree to which N and P are

regulated differs among major groups of taxa. For example, the biomass stoichiometry of primary producers tends to be flexible, reflecting the supply of nutrients in their environment

(Sterner & Elser 2002). In contrast, the biomass stoichiometry of consumer populations (e.g., viruses, invertebrates, mammals) is homeostatically regulated and therefore tends to remain

constant even when the nutrient content of their resources fluctuates (Sterner & Elser 2002). Such differences can create a nutritional imbalance between primary producers and consumers,

which has profound consequences for a wide range of ecological processes including resource competition, host-parasite dynamics, and ecosystem functioning (Smith & Holt 1996; Hall 2009;

Aalto *et al.* 2015).

        Nutrient stoichiometry can also influence evolutionary processes. Nutrient limitation

66   often leads to a reduction in population size, which can diminish the efficiency of selection and result in the accumulation of deleterious mutations through genetic drift (Kimura 1962).

68   However, many populations have adaptations that allow individuals to contend with absolute and relative resource scarcity. For example, the disproportionate use of nucleotides that vary in N

70   content reduces stoichiometric mismatch between an organism and its environment (Elser *et al.*

2011). Similarly, natural selection can operate on the material costs of gene expression in ways

72   that lead to the sparing of elements that are normally used in highly expressed proteins (Bragg &

Wagner 2009). These genetic responses to nutrient limitation may give rise to the evolution of

74   organismal stoichiometry. For example, long-term carbon limitation led to increased N and P

content of experimentally evolved bacterial populations (Turner *et al.* 2017). However,

76   evolutionary responses to nutrient stoichiometry may depend on the identity of the limiting

nutrient (Gresham *et al.* 2008). While N limitation can select for stress-related or catabolic genes

78   with lower guanine-cytosine content (Acquisti *et al.* 2009), P limitation can favor the

replacement of phospholipids with sulfolipids in the cell membranes of certain microorganisms

80   (Van Mooy *et al.* 2006).

       When considering multiple interacting species, the combined effects of nutrient

82   stoichiometry may give rise to eco-evolutionary feedbacks. Such feedbacks occur when

ecological interactions affect evolutionary processes, which in turn modify species interactions

84   and ecological dynamics (Haloin & Strauss 2008). For example, the rapid evolution of functional

traits can produce diminished oscillations, longer periods of cycling, and phase-shifted

86   population densities between hosts and their parasites, a phenomenon referred to as "cryptic

dynamics" (Yoshida *et al.* 2007). Theory suggests that cryptic dynamics can arise when nutrient

88   stoichiometry alters the stability of antagonistic species interactions (Yamamichi *et al.* 2015),

which may ultimately intensify the modes of coevolution, namely arms-race dynamics and

90   negative frequency-dependent selection (Aalto *et al.* 2015). However, links among nutrient

stoichiometry, eco-evolutionary feedbacks, and coevolution remain to be tested.

Major advances in evolutionary ecology have been made through experimental studies of microbial communities. In particular, bacteria and phage are ideal for studying eco-evolutionary

feedbacks owing to their large population sizes, rapid growth rates, and experimental tractability. Moreover, bacteria and phage dynamics are critical for understanding the structure and function

of microbial food webs, especially in aquatic ecosystems (Suttle 2007). For example, *Synechococcus* is a diverse and widely distributed group of primary producers in the world ocean

with an estimated global abundance of 10^{26} cells (Flombaum *et al.* 2013). While *Synechococcus* must contend with both N- and P-limitation, it is also subject to a high degree of phage-induced

mortality, which may lead to nutrient-dependent coevolutionary dynamics. These coevolutionary processes can create feedbacks that not only influence population dynamics, but also ecosystem

functioning, including the turnover of N and P (Lennon & Martiny 2008).

In this study, we tested how nutrient stoichiometry affects host-parasite eco-evolutionary

dynamics in a simplified community where a single genotype of marine *Synechococcus* was infected with a phage strain in continuous culture using chemostats that were supplied with either

N- or P-limited media. We isolated hundreds of host and phage strains, which were challenged against one another to document phenotypic changes in resistance and infectivity over the course

of the experiment. With this information, we conducted network analyses that allowed us to quantify the degree of nestedness and modularity, which can serve as signatures of shed light on

the underlying modes of coevolution, namely arms race dynamics and negative frequency dependent selection (Weitz *et al.* 2013). Our findings reveal that nutrient stoichiometry is a

bottom-up force that regulates eco-evolutionary feedbacks in ways that alter coevolutionary processes.

Strains and media — We evaluated the effects of nutrient stoichiometry on the eco-
evolutionary dynamics of the marine cyanobacterium *Synechococcus* WH7803 and a lytic T4-

118   like phage belonging to the Myoviridae family of phage (S-RIM8). Changes to nutrient supply
can alter the equilibrium density of microbial populations in continuous culture (i.e.,

120   chemostats), which in turn, may affect cyanobacteria-phage contact rates (Rhee 1978).
Therefore, we adjusted both the concentrations and ratios of N and P in modified AN artificial

122   seawater medium (Tables S1-S3) to induce N- or P-limitation while maintaining similar
equilibrium densities of *Synechococcus* between the stoichiometry treatments prior to phage

124   addition. Specifically, the N-limited medium had an N : P ratio of 10 : 1 (KNO_3; N = 220 µM)
while the P-limited medium had a N : P ratio of 40 : 1 (K_2HPO_4; P = 11 µM). Growth assays

126   confirmed that *Synechococcus* was limited by N under low N : P supply and by P under high N :
P supply (Fig. S1).

   Nutrient limitation assay — We tested for nutrient limitation in *Synechococcus* by measuring

130   growth rates on cells that had been maintained under low (10 : 1) and high (40 : 1) supply ratio
of N and P. We acclimated *Synechococcus* semi-continuously under constant light (20 µE m^{-1} s^{-1})

132   at 25 °C for three serial transfers in either N- or P-limited medium (n = 10). We then inoculated
each replicate cell line into fresh medium that doubled the concentration of N (N-limited = 440

134   µM, P-limited = 880 µM) or P (N-limited = 44 µM, P-limited = 22 µM). Next, we monitored
Synechococcus population densities for seven days *via* autofluorescence (ex: 550 nm, em: 570

136   nm) with a Biotek Synergy Mx plate reader (Winooski, VT, USA). With these data, we
estimated the maximum growth rate for each culture using a modified Gompertz equation

138   (Lennon et al. 2007) and calculated the percent change in growth rate with additional N or P as

       compared to the control, which contained no additional N or P.

       Chemostat experiment — We supplied ten chemostats, each with a 40 mL operating volume,

142   with N-limited or P-limited medium at a dilution rate of 1 d^{-1} (Tables S1-S3). The chemostats

       were maintained in a Percival growth chamber (Perry, IA, USA) at 25 °C on a 14:10 light : dark

144   cycle under 20 μE m^{-2} s^{-1} and homogenized with magnetic stir bars following inoculation with a

       single-colony strain of *Synechococcus* WH7803. We allowed the *Synechococcus* populations to

146   equilibrate in the chemostats prior to initiating the "phage-amended" treatment by introducing an

       aliquot of a plaque-purified S-RIM8 to three randomly chosen chemostats in each nutrient

148   treatment to achieve a multiplicity of infection (phage : cyanobacteria ratio) of approximately 10.

       To document the potential influence of stoichiometry on population dynamics and population

150   size, we maintained "no phage" chemostats with only *Synechococcus* in both stoichiometry

       treatments (Fig. S2).

       Community dynamics — We tracked *Synechococcus* and phage densities in each chemostat via

154   epifluorescent microscopy every other day for 172 days. *Synechococcus* populations were

       enumerated by concentrating samples from a chemostat onto 0.22-μm nominal pore-size black

156   polycarbonate filters. After removing cellular material (0.2 μm filtration) and extracellular DNA

       with DNase I, we concentrated samples onto 0.02 μm Anodisc filters to estimate the abundance

158   of free phage particles. Each phage-containing filter was then stained with SYBR Green I

       (Noble & Fuhrman 1998). We estimated population densities after capturing ten images from

160   each filter with a CY3 filter set (ex: 550 nm, em: 570 nm) for *Synechococcus* or a FITC filter set

(ex: 497 nm, em: 520 nm) for phage using a Zeiss microscope and image processing modules

162   found in the base package of the Axiovision imaging software (Release 4.5 SP1).

We evaluated the influence of nutrient stoichiometry on community dynamics in three

164   ways. First, we used repeated measures (RM) ANOVA to test for the effects of nutrient

stoichiometry and time on *Synechococcus* and phage densities. Second, because cryptic

166   dynamics can alter the temporal synchrony between hosts and parasites, we performed cross-

correlation analyses for each chemostat using Auto Regressive Moving Average (ARMA)

168   procedures (Lennon & Martiny 2008). In addition to determining whether populations in a

chemostat positively or negatively covaried (i.e., in phase vs. out of phase, respectively) we used

170   the cross-correlation analyses to test if the duration of time which viruses lagged behind host

densities was altered by nutrient stoichiometry. Last, we estimated the effects of nutrient

172   stoichiometry on the stability of microbial populations by calculating the inverse of the

coefficient of variation (CV) over time.

Coevolutionary dynamics — To test for the effects of nutrient stoichiometry on phenotypic

176   coevolution, we tracked changes in infection patterns between *Synechococcus* and its phage over

time. We isolated multiple (3-5) *Synechococcus* strains from each chemostat using serial dilution

178   techniques six days prior to phage addition (day -6) and at days 9, 23, 72, 129, 148, and 166 after

phage addition. We performed 10-fold dilutions of chemostat samples into 24-well plates

180   containing sterile AN media that matched the nutrient supply ratio of the chemostat environment.

When a dilute suspension of *Synechococcus* 7803 was incubated without shaking, single-

182   colonies formed in the bottom of the wells (25 °C on a 14:10 light : dark cycle under 20 μE m^{-2} s^{-1}). We picked these colonies with a Pasteur pipette and passaged them twice in 50 mL

184   Erlenmeyer flasks using the same medium prior to harvesting. We then used epifluorescence

microscopy to screen the cultures to make sure they did not contain free phage particles. Finally,

186   these *Synechococcus* strains were concentrated *via* centrifugation, preserved in glycerol (10 %

final concentration), and stored at -80 °C until reanimation for use in challenge assays, which are

188   described below. We also isolated multiple (3-5) phage strains from the phage-amended

chemostats on days 23, 72, 129, 148, and 166 through double plaque purification. This process

190   involved adding 40 µL of 0.2 µm-filtered chemostat sample to a soft agar (0.8 % agar) overlay

containing the ancestral *Synechococcus* WH7803 as the host. We then incubated the serially

192   diluted plates (25 °C on a 14:10 light : dark cycle under 20 $\mu E\ m^{-2}\ s^{-1}$), picked plaques that

formed on the lawns of *Synechococcus*, and transferred to a log-phase batch culture of

194   *Synechococcus* for propagation. The resulting phage lysates were syringe-filtered (1 µm) and

preserved in glycerol (10 % final concentration) at -80 °C.

196   With the isolated chemostat strains, we quantified host resistance and phage infectivity

using challenge assays. Each pair-wise challenge with strains isolated across chemostats within

198   the same nutrient treatment was carried out in triplicate by adding 20 µL of a phage stock ($\sim10^7$

particles mL^{-1}) to 200 µL of a dilute *Synechococcus* strain ($\sim10^6$ cells mL^{-1}) in 96-well plates.

200   Challenge assays were performed using the same medium from which the host strain was

originally isolated (i.e., low vs. high N : P). Turbid cultures of *Synechococcus* WH7803 appear

202   bright pink owing to the intracellular photosynthetic pigment phycoerythrin. When infected by

SRIM8, however, cells are lysed and cultures become clear (Lennon & Martiny 2008). Based on

204   this, we scored each *Synechococcus* strain as sensitive (and the phage strain as infective) if there

was a lack of growth after a two-week incubation under continuous light (20 $\mu E\ m^{-2}\ s^{-1}$) at 25 °C

206   compared to control wells (n = 3) that contained heat-killed phage. To contend with the loss of a

replicate chemostat in the N-limited treatment, we randomly selected *Synechococcus* strains

208   across the two P-limited replicate chemostats to ensure that an identical number of strains were

tested at each time point. In total, there were 18,050 pairwise challenges between chemostat-

210   isolated strains of *Synechococcus* and phage that resulted in a bipartite infection matrix, which

we used for all analyses related to phenotypic coevolution.

212   To quantify the effect of nutrient stoichiometry on coevolutionary dynamics, first, we

used RM-ANOVA to test for trends in average resistance and average infectivity over the course

214   of the chemostat experiment. Second, we used the infection matrix to perform time-shift

analyses, which involved calculating the proportion of successful infections that occurred

216   between hosts that were challenged against past, contemporary, and future phage strains (Gaba &

Ebert 2009). We statistically analyzed the time-shift data using RM-ANOVA with nutrient

218   treatment, phage treatment, and time as fixed effects, while the chemostat replicate identifier was

treated as a random effect with a corARMA covariance matrix (Koskella 2014). To visualize the

220   data, contemporary challenges were centered at a time-shift of zero, while interactions with past

phage were represented in negative space and future interactions were represented in positive

222   space. Last, to gain insight into potential mechanisms underlying stoichiometrically driven

coevolution, we used community network analyses to calculate the connectance, nestedness, and

224   modularity for each chemostat infection matrix. Connectance was calculated as the number of

interactions divided by network size. We calculated nestedness using the NODF metric, which

226   ranges from 0 (non-nested) to 1 (perfectly nested) and normalizes for matrix size. We used the

LP-BRIM algorithm to find the partition that maximizes Barber's modularity (Qb), which ranges

228   from 0 (all interactions are between modules) to 1 (all interactions are within modules). The

network statistics were calculated using 100,000 random Bernoulli simulations in the BiWeb

230   package for MATLAB (Flores *et al.* 2011; Flores *et al.* 2016), http://github.com/tpoisot/BiWeb).

We then tested for differences in connectance, nestedness, and modularity between the nutrient

232   treatments using *t*-tests.

234   **RESULTS**

Nutrient limitation of *Synechococcus* — Prior to initiating the chemostat evolution trial, we

236   conducted growth assays to test for N- and P-limitation of *Synechococcus*. After a period of

acclimation in semi-continuous culture, the growth rates of N-limited *Synechococcus* increased

238   by 87 % when additional N was supplied, but only by 35 % with the addition of P (*t*-test, t_8 =

4.55, $P < 0.001$). In contrast, the growth rate of P-limited *Synechococcus* increased by 45 % with

240   the addition of P, but only by 25 % with the addition of N (*t*-test, t_8 = -3.36, $P = 0.005$). Results

from this experiment suggest that while there may have been some degree of co-limitation, the

242   two media sources used for subsequent evolution trials induced N- and P- limitation for

Synechococcus (Fig. S1).

Stoichiometry altered community dynamics — In the chemostat experiment, nutrient

246   stoichiometry significantly affected the population dynamics of *Synechococcus* (RM-ANOVA;

time x stoichiometry, $F_{62,\ 245}$ = 2.43, $P < 0.0001$) and its phage (RM-ANOVA; time x

248   stoichiometry, $F_{58,\ 230}$ = 2.59, $P < 0.0001$). Under N-limitation, *Synechococcus* rapidly declined

and reached its minimum mean density 35 days (range = 23 – 44) following phage addition (2 x

250      $10^5 \pm 7$ x 10^4 cells mL^{-1}, mean $\pm$ SEM, Fig. 1a). This decrease in host abundance corresponded

with a peak in mean maximum density 25 days (range = 16 – 30) following phage addition (9 x

252    $10^8 \pm 1 \times 10^8$ particles mL^{-1}, mean $\pm$ SEM). Over the next ~50 days, the host densities slowly

recovered, and entered a second phase of decline near day 100.

254   *Synechococcus* and phage densities were significantly different and less dynamic under

P-limitation (Fig. 1b). Hosts declined more slowly in P-limited chemostats following phage

256   addition than in N-limited chemostats (*t*-test, $t_4 = -4.00$, $P = 0.02$) and reached their mean

minimum density 78 (range $= 63 - 93$) days following phage addition ($4 \times 10^5 \pm 6 \times 10^4$ cells

258   mL^{-1}, mean $\pm$ SEM). Though hosts recovered, they did not reach pre-phage-addition abundances

until the end of the experiment. Phage densities under P-limitation remained high and relatively

260   constant ($2 \times 10^8 \pm 9 \times 10^7$ particles mL^{-1}, mean $\pm$ SEM) throughout the duration of the

experiment. In the no-phage control chemostats, N- and P- limited *Synechococcus* densities

262   remained constant over time (Fig. S2).

Nutrient stoichiometry also affected the temporal coherence and stability of

264   *Synechococcus* and phage populations. After pre-whitening the time-series data using ARMA

procedures, phage densities in N-limited chemostats were negatively correlated with host

266   densities with lags ranging from zero to five days ($r = -0.27$ to -0.42, $P = 0.009 - 0.063$, Fig. S3).

In contrast, under P-limitation, there was no correlation between phage and *Synechococcus* at

268   any time lag ($r = -0.20$ to 0.22, $P \geq 0.23$). Last, we found that cyanobacterial and phage densities

were significantly more stable over time under P- than N-limited conditions (*Synechococcus*: *t*-

270   test, $t_4 = 3.56$, $P = 0.024$; *t*-test, phage: $t_4 = 3.83$, $P = 0.019$). See Table S4 for summary statistics.

272   **Stoichiometry altered coevolution** — Within nine days of phage addition, >50 % of the

isolated *Synechococcus* strains were resistant to the ancestral phage in both the N- and P-limited

274   chemostats (Fig. 2). Average resistance continued to increase over time (RM-ANOVA, $F_{6, 591} =$

14.2, $P < 0.0001$), but was not affected by stoichiometry ($F_{1, 4} = 2.08$, $P = 0.22$). The vast

276   majority (97 %) of *Synechococcus* strains that were resistant to the ancestral phage were also

resistant to all of the phage strains subsequently isolated from the chemostats. Phage-resistant

278   *Synechococcus* did not revert to the sensitive phenotype when grown in the absence of phage

consistent with the view that resistance is a heritable trait.

280   Phage infectivity also evolved, but in contrast to the host, was significantly affected by

nutrient stoichiometry (Fig. 2; RM-ANOVA, stoichiometry x time: $F_{1, 601} = 7.53$, $P = 0.006$). In

282   the no-phage control chemostats, *Synechococcus* strains remained susceptible to the ancestral

phage. However, nearly all (~ 96 %) of the derived phage from the phage-amended chemostats

284   lost the ability to infect some of the *Synechococcus* hosts from the P-limited (n = 51) and N-

limited (n = 22) no-phage chemostats. Despite this, we found evidence of host-range expansion.

286   For example, a phage strain from a P-limited chemostat (day 129) was able to infect a

Synechococcus strain (day 166) that was resistant to the ancestral phage. In addition, three phage

288   strains from N-limited chemostats and two phage strains from P-limited chemostats (day 166)

were able to infect phage-resistant *Synechococcus* isolated from other chemostats earlier in the

290   study.

Time-shift experiments indicated that nutrient stoichiometry altered coevolutionary

292   dynamics (Figs. 3 and 4). When hosts isolated from the phage-amended chemostats were

challenged against current and past phage, interaction strengths were generally weak owing to

294   the evolution of host resistance and phage infectivity (Fig. 3 and Fig 4 a, b; RM-ANOVA,

stoichiometry x phage isolation time x bacterial isolation time, $F_{1,597} = 24.67$, $P < 0.0001$). Hosts

296   were more susceptible when challenged against future phage, but only for *Synechococcus* strains

that were isolated earlier in the chemostat experiment (days -6, 9, and 23). The effect of nutrient

298 stoichiometry was more evident in time-shift experiments where naïve hosts from the no-phage

control chemostats were challenged against phage isolated from the phage-amended chemostats

300 (stoichiometry x phage isolation time, $F_{1, 282} = 7.25$, $P = 0.0075$). Under these conditions,

infectivity was not uniformly high across the time-shifts (Fig 3 and Fig. 4 c, d), as would be

302 expected from purely arms-race dynamics. As a result, infectivity was 70 % greater on naïve

hosts isolated from N-limited chemostats (0.61 ± 0.289) compared to naïve hosts isolated from

304 P-limited chemostats (0.36 ± 0.332).

Network analyses lent further support that nutrient stoichiometry affected host-phage

306 coevolution. Patterns of infectivity were more nested and modular than expected by chance ($P <$

0.05), which is consistent with arms race dynamics and negative frequency-dependent selection,

308 respectively. The infection networks from N- and P-limited environments had similar degrees of

connectance ($t_4 = 1.47$, $P = 0.22$) and nestedness (NODF, $t_4 = -0.37$, $P = 0.72$). However, P-

310 limited networks were 50 % more modular than N-limited networks (Fig. 5; $t_4 = -3.59$, $P = 0.02$).

See Table S6 for summary statistics.

DISCUSSION

314 The supply ratio of nitrogen (N) and phosphorus (P) had strong eco-evolutionary effects

on marine *Synechococcus* and its phage. Communities were more stable under P-limitation than

316 N-limitation, which likely was influenced by the nutritional flexibility and constraints of the host

and phage populations, respectively. In addition, host-phage dynamics were influenced by the

318 rapid invasion of resistant cyanobacteria which was followed by the appearance of evolved

phage with altered host-ranges. Using time-shift assays and network analyses, we detected strong

320 signatures of arms-race dynamics in both nutrient treatments. However, under P-limitation, we

also observed patterns of increased network modularity consistent with negative frequency-

322   dependent selection, which may reflect the underlying costs of maintaining defense and

virulence traits in different nutrient environments. Our findings generate testable predictions

324   regarding the mechanisms of local adaptation and coevolution for microorganisms among

regions of the world ocean that are limited to varying degrees by N and P (Galbraith & Martiny

326   2015).

328   **Stoichiometry altered community dynamics** — Consistent with predictions from the theory of

ecological stoichiometry (Sterner & Elser 2002), we documented that *Synechococcus* and phage

330   dynamics were highly sensitive to the N : P supply ratio. Under N-limitation, host-phage

dynamics were less stable and out-of-phase, compared to P-limited chemostats (Fig. 1; Table

332   S4). Historically, ecologists have attempted to explain such variation in populations and

communities while assuming a single currency of resource. However, systems can exhibit a

334   much wider range of behaviors when resource stoichiometry is explicitly considered owing to

feedbacks that arise when interacting species vary in the degree to which their elemental

336   composition is homeoestatically regulated (Andersen *et al.* 2004). For example, the molar N : P

ratio of marine *Synechococcus* converges upon the canonical Redfield value of 16 : 1 in some

338   environments (Garcia *et al.* 2016), but in other instances can exceed 100 : 1 (Bertilsson *et al.*

2003) reflecting the extreme plasticity of its biomass stoichiometry. In contrast, phage are

340   thought to have a fixed elemental composition owing to their relatively simple structure, which

minimally consists of genetic material (DNA or RNA) protected by a proteinaceous capsid

342   (Jover *et al.* 2014). Biophysical models predict that the T4-like phage used in our study has a

high P demand reflected by its low N : P ratio of approximately 7 : 1 (Jover *et al.* 2014). As a

344   consequence, phage productivity is largely dependent on the P content of their hosts. For

example, host lysis was delayed by 18 h leading to an 80 % reduction in phage burst size when

346   P-limited *Synechococcus* WH7803 was infected with a myovirus (Wilson *et al.* 1996). Consistent

with these reports, early in our experiment (day -6), phage infectivity was 50 % lower on P-

348   limited vs. N-limited hosts (cf. Fig. 4A and 4B). It is possible, however, for phage to overcome

the nutritional constraints of their microbial hosts. Many phage contain auxiliary metabolic genes

350   that modify a host's resource allocation (Monier *et al.* 2017). For example, phage-encoded genes

for phosphorus acquisition (*pstS* and *phoA*) were upregulated when phosphorus-starved marine

352   *Prochlorococcus* was infected by phage P-SSM2 (Zeng & Chisholm 2012). While the myovirus

used in our study (S-RIM8) does not contain phosphate-binding protein or alkaline phosphatase

354   genes, it does contain *phoH*. The exact function of *phoH* in marine microorganisms has not been

demonstrated, but it is thought to encode a phosphate-starvation inducible protein that

356   hydrolyzes ATP, thus liberating energy that can be used in P uptake (Goldsmith *et al.* 2011).

Such findings suggest that nutrient stoichiometry can profoundly shape host-parasite dynamics

358   (Aalto *et al.* 2015) and provide an ecological explanation for the patterns observed in our study

(Fig. 1).

Stoichiometry altered coevolution — Our results indicate that evolutionary processes were also

362   important in creating feedbacks that influenced host-phage dynamics. Such feedbacks arise when

rapid changes in host or parasite traits generate patterns that are not predicted from traditional

364   ecological theory (Yoshida *et al.* 2007; Frickel *et al.* 2016). These so-called cryptic dynamics

require the coexistence of multiple host genotypes, for example, through trade-offs in

366   competitive ability and parasite defense (Yoshida *et al.* 2007). While there are many intracellular

and extracellular mechanisms that bacteria can employ to resist phage infection, in

368 *Synechococcus*, mutations in genes encoding for cell-surface receptors reduce or entirely

eliminate attachment, thus precluding entry of the virus into the host (Stoddard *et al.* 2007;

370 Marston *et al.* 2012). Selection for this phenotype could help explain the recovery of host

densities in our chemostats (Fig. 1). However, resistance mutations in *Synechococcus* are often

372 accompanied by a reduction in growth rate, the magnitude of which can vary depending on the

identity of the host and virus (Lennon 2007). These fitness costs have important consequences

374 for understanding community stability. Without a reduction in growth rate, resistant

Synechococcus would outcompete the sensitive host and drive the phage population extinct.

376 Moreover, fitness costs establish a trade-off that satisfies the requirement for cryptic dynamics to

emerge (Yoshida *et al.* 2007). In sum, our data support the view that nutrient stoichiometry

378 altered host-phage dynamics and stability via eco-evolutionary feedbacks (Lennon & Martiny

2008).

380   Nutrient stoichiometry had strong, yet importantly, asymmetric effects on microbial

coevolution in our study. These effects were examined in light of the two primary modes by

382 which antagonistic coevolution is thought to occur. The first is through arms-race dynamics

involving gene-for-gene specificity, where directional selection leads to sweeps that are

384 characterized by the escalation of resistance and infectivity. Arms-race dynamics were originally

described for coevolving populations of plants and pathogens, but since then have been

386 commonly reported in studies of bacteria and phage (Dennehy 2012; Koskella & Brockhurst

2014). The second mode of antagonistic coevolution involves negative frequency-dependent

388 selection where parasites evolve to infect common hosts, which in turn favors rare host alleles.

Negative frequency-dependent selection is often associated with infections that require matching

390 alleles and is well documented in invertebrate systems (Decaestecker *et al.* 2007), but has also

been described in some studies of bacteria and phage (Hall *et al.* 2011). While arms-race

392   dynamics and negative frequency-dependent selection are often viewed as occupying different

ends of the coevolutionary spectrum, they are not mutually exclusive (Agrawal & Lively 2002).

394   One powerful way to discern modes of coevolution is through the use of time-shift

analyses. This approach involves determining the success of infections for combinations of hosts

396   and parasites that are isolated from different time points in a controlled experiment or other

longitudinal type of study (Gaba & Ebert 2009). When applied to challenge assays among host

398   and phage isolates from our chemostats, time-shift analyses indicated that coevolutionary

dynamics were significantly affected by nutrient stoichiometry (Fig. 3). The effect of nutrient

400   stoichiometry was relatively weak when examining host strains isolated from phage-amended

chemostats, which could reflect a mismatch between the time-shift interval and the rates of

402   coevolution (Gandon *et al.* 2008). In contrast, the effect of nutrient stoichiometry was driven

largely by interactions between evolved viruses that were challenged against *Synechococcus*

404   isolated from no-phage control chemostats (Fig. 4 c, d). We expected that these naïve hosts

would be uniformly and highly susceptible to all phage from the phage-amended chemostats.

406   Instead, we found that infectivity fluctuated over time and was higher on N-limited vs. P-limited

Synechococcus, reflecting expansion and contraction of the phage host-range (Fig. 4 c, d).

408   Together, these findings are consistent with observations of asymmetric coevolution that have

been attributed to the genetic constraints of viral populations (Lenski & Levin 1985). Asymmetry

410   may also arise from the costs associated with being a generalist phage (Koskella & Brockhurst

2014), which may be dependent on the resource environment (e.g., nutrient stoichiometry). Such

412   explanations have been used to explain host-phage coevolution where arms race dynamics give

way to negative frequency dependent selection (Hall *et al.* 2011), in some cases depending on

414   resource availability (Pascua *et al.* 2014).

    The effects of nutrient stoichiometry on coevolution were further supported by results

416   from our network analyses. With this approach, one can estimate the degree of nestedness that

exists among pairs of hosts and phage relative to randomized data. Often, infection data from

418   bacteria-phage systems are highly nested (Flores *et al.* 2011), which means they contain ordered

subsets of phenotypes whereby hosts from later time points are resistant to earlier phages, and

420   phage from later time points are able to infect earlier hosts (Weitz *et al.* 2013). In our study,

infection networks were significantly nested, a pattern that arises from arms-race dynamics.

422   However, the degree of nestedness was not affected by nutrient stoichiometry. Infection

networks can also exhibit modularity, which is more consistent with negative frequency-

424   dependent selection (Weitz *et al.* 2013). Modules reflect dense clusters of interacting host and

phage compared to other strain combinations found in a bipartite infection matrix. Although less

426   commonly documented, modularity has been observed in large-scale oceanic surveys of bacteria

and phage, but has been attributed to local adaptation and the phylogenetic breadth of the hosts

428   (Flores *et al.* 2013). However, theory suggests that modularity can emerge *via* coevolution

between two populations at the local scale. Using a "relaxed lock and key" model,

430   simultaneously nested and modular structures arose under simple chemostat conditions that

assumed gene matching between phage tail-fibers and host receptors (Beckett & Williams 2013).

432   Our results agree with these predictions, but suggest that network properties are affected by host

nutrition. Specifically, we found that host-phage interactions were 50 % more modular under P-

434   limitation than N-limitation, suggesting that nutrient stoichiometry constrains host-phage

interactions leading to increased specialization. Such findings are consistent with the view that

436 resources can influence the modes of coevolution based in part on the fitness costs associated

with host defense and infection strategies (Pascua *et al.* 2014)

438   Mechanistically, there are many ways that nutrient stoichiometry could shape bacteria-

phage coevolution. For example, the "dangerous nutrients" hypothesis predicts that trajectories

440 of coevolution can be influenced when receptors used by nutrient-limited hosts also serve as the

targets of phage adsorption (Menge & Weitz 2009). However, it does not appear that

442 myoviruses, including the strain used in this study, attach to the protein receptors of

cyanobacteria that are used for nutrient transport. Instead, evidence from whole-genome

444 sequencing of the host used in our study suggests that phage-resistant *Synechococcus* accumulate

mutations in hypervariable genomic islands that encode for lipopolysaccharide (LPS) (Marston

446 *et al.* 2012), a major component of the outer membrane in Gram-negative bacteria. The structural

complexity of LPS is affected by nutrient limitation (Brelles-Marino & Boiardi 1997) and such

448 changes in molecular structure of the cell membrane can interfere with phage adsorption leading

to resistance (Leon & Bastias 2015). Nevertheless, modification of tail-fibers allow phage to

450 overcome resistance of marine cyanobacteria in some instances (Schwartz & Lindell 2017) while

the acquisition of host-like P-assimilation genes may aid in the successful infection of nutrient-

452 limited hosts (Kelly *et al.* 2013).

454 **Conclusions —** Most organisms live in environments where they are limited by the relative or

absolute amount of one or more essential resources. It is well established that such variation in

456 nutrient stoichiometry can regulate ecological phenomena ranging from species interactions to

ecosystem-level processes. We demonstrated that nutrient stoichiometry also affects the

458 evolutionary dynamics of microbial communities likely through differences in the expansions

and contraction of virus host-ranges under N- vs. P-limited conditions. Identifying the targets of

460   selection in contrasting nutrient environments will help elucidate the genetic mechanisms of

coevolution and the trade-offs associated with defense and virulence traits. While our study

462   offers promising avenues to better understand bacteria-phage interactions in the aquatic

environments, nutrient stoichiometry is also important in determining the nutrition, health, and

464   disease susceptibility for non-microbial hosts and their microbiomes(Smith & Holt 1996), which

has implications for understanding the persistence, emergence, and evolution of infectious

466   diseases. In the oceans, phage represent a significant source of mortality for marine

cyanobacteria like *Synechococcus*, which play a central role in the regulation of biogeochemical

468   processes, including energy flow, carbon sequestration, and the cycling of nitrogen (N) and

phosphorus (P). Our findings suggest that the coevolutionary process was differentially affected

470   by the availability of N and P in ways that could influence the ecology and evolution of one of

the most abundant and functionally important groups of microorganisms on Earth.

Acknowledgments

474   We acknowledge technical support from R Morrison, M Carroll, and BK Lehmkuhl; ancestral

strains from MF Marston; discussion with CM Lively; feedback from BK Whitaker, KJ Locey,

476   WR Shoemaker, NI Wisnoski, V Kuo, ME Muscarella, DA Schwartz, RZ Moger-Reischer, JM

Palange on earlier versions of the manuscript. Financial support was provided by the National

478   Science Foundation (0851143, 0851113) and the US Army Research Office Grant W911NF-14-

1-0411. We dedicate this work to the memory of VH Smith.

REFERENCES

1.

484 Aalto, S.L., Decaestecker, E. & Pulkkinen, K. (2015). A three-way perspective of stoichiometric changes on host-parasite interactions. *Trends Parasitol.*, 31, 333-340.

486 2.

Acquisti, C., Elser, J.J. & Kumar, S. (2009). Ecological Nitrogen Limitation Shapes the DNA
488 Composition of Plant Genomes. *Mol. Biol. Evol.*, 26, 953-956.

3.

490 Agrawal, A. & Lively, C.M. (2002). Infection genetics: gene-for-gene versus matching-alleles models and all points in between. *Evol. Ecol. Res.*, 4, 79-90.

492 4.

Andersen, T., Elser, J.J. & Hessen, D.O. (2004). Stoichiometry and population dynamics. *Ecol.*
494 *Lett.*, 7, 884-900.

5.

496 Beckett, S.J. & Williams, H.T.P. (2013). Coevolutionary diversification creates nested-modular structure in phage-bacteria interaction networks. *Interface Focus*, 3: 20130033.

498 6.

Bertilsson, S., Berglund, O., Karl, D.M. & Chisholm, S.W. (2003). Elemental composition of
500 marine *Prochlorococcus* and *Synechococcus*: Implications for the ecological stoichiometry of the sea. *Limnol Oceanogr*, 48, 1721-1731.

502 7.

Bragg, J.G. & Wagner, A. (2009). Protein material costs: single atoms can make an evolutionary
504 difference. *Trends Genet.*, 25, 5-8.

8.

506 Brelles-Marino, G. & Boiardi, J.L. (1997). Mn^{2+} deficiency affects growth energetics of *Rhizobium etli*. *Microbiol. Res.*, 152, 367-371.

508 9.

Decaestecker, E., Gaba, S., Raeymaekers, J.A.M., Stoks, R., Van Kerckhoven, L., Ebert, D. *et al.*
510 (2007). Host-parasite 'Red Queen' dynamics archived in pond sediment. *Nature*, 450, 870-U816.

512 10.

Dennehy, J.J. (2012). What can phages tell us about host-pathogen coevolution? *Int. J. Evol. Biol.*,
514 2012, 396165.

11.

516 Elser, J.J., Acquisti, C. & Kumar, S. (2011). Stoichiogenomics: the evolutionary ecology of macromolecular elemental composition. *Trends Ecol. Evol.*, 26, 38-44.

518 12.

Flombaum, P., Gallegos, J.L., Gordillo, R.A., Rincon, J., Zabala, L.L., Jiao, N.A.Z. *et al.* (2013).
520 Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proc. Natl Acad. Sci. U S A*, 110, 9824-9829.

522 13.

Flores, C.O., Meyer, J.R., Valverde, S., Farr, L. & Weitz, J.S. (2011). Statistical structure of host-
524 phage interactions. *Proc. Natl Acad. Sci. U S A*, 108, E288-E297.

14.

526 Flores, C.O., Poisot, T., Valverde, S. & Weitz, J.S. (2016). BiMat: a MATLAB package to facilitate the analysis of bipartite networks. *Methods Ecol. Evol.*, 7, 127-132.

528 15.
Flores, C.O., Valverde, S. & Weitz, J.S. (2013). Multi-scale structure and geographic drivers of
530 cross-infection within marine bacteria and phages. *ISME J.*, 7, 520-532.

532 16. Frickel, J., Sieber, M. & Becks, L. (2016). Eco-evolutionary dynamics in a coevolving host-virus system. *Ecol. Lett.*, 19, 450-459.

534
17.
536 Gaba, S. & Ebert, D. (2009). Time-shift experiments as a tool to study antagonistic coevolution. *Trends Ecol. Evol.*, 24, 226-232.

538 18.
Galbraith, E.D. & Martiny, A.C. (2015). A simple nutrient-dependence mechanism for predicting
540 the stoichiometry of marine ecosystems. *Proc. Natl Acad. Sci. U S A*, 112, 8199-8204.
19.
542 Gandon, S., Buckling, A., Decaestecker, E. & Days, T. (2008). Host-parasite coevolution and patterns of adaptation across time and space. *J. Evol. Biol.* 21, 1861-1866.

544
20.
546 Garcia, N.S., Bonachela, J.A. & Martiny, A.C. (2016). Interactions between growth-dependent changes in cell size, nutrient supply and cellular elemental stoichiometry of marine
548 *Synechococcus*. *ISME J.*, 10, 2715-2724.
21.
550 Goldsmith, D.B., Crosti, G., Dwivedi, B., McDaniel, L.D., Varsani, A., Suttle, C.A. *et al.* (2011). Development of *phoH* as a novel signature gene for assessing marine phage diversity. *Appl*
552 *Environ Microbiol*, 77, 7730-7739.
22.
554 Gresham, D., Desai, M.M., Tucker, C.M., Jenq, H.T., Pai, D.A., Ward, A. *et al.* (2008). The repertoire and dynamics of evolutionary adaptations to controlled nutrient-limited
556 environments in yeast. *PLoS Genet.*, 4.
23.
558 Hall, A.R., Scanlan, P.D., Morgan, A.D. & Buckling, A. (2011). Host-parasite coevolutionary arms races give way to fluctuating selection. *Ecol. Lett.*, 14, 635-642.
560 24.
Hall, S.R. (2009). Stoichiometrically explicit food webs: feedbacks between resource supply,
562 elemental constraints, and species diversity. *Annu. Rev. Ecol. Evol. Syst.*, 40. 503-528.
25.
564 Haloin, J.R. & Strauss, S.Y. (2008). Interplay between ecological communities and evolution. *Ann. N.Y. Acad. Sci.*, 1133, 87-125.
566 26.
Jover, L.F., Effler, T.C., Buchan, A., Wilhelm, S.W. & Weitz, J.S. (2014). The elemental
568 composition of virus particles: implications for marine biogeochemical cycles. *Nat. Rev. Microbiol.*, 12, 519-528.
570 27.

572 Kelly, L., Ding, H.M., Huang, K.H., Osburne, M.S. & Chisholm, S.W. (2013). Genetic diversity in cultured and wild marine cyanomyoviruses reveals phosphorus stress as a strong selective agent. *ISME J.*, 7, 1827-1841.

574 28.
Kimura, M. (1962). On the probability of fixation of mutant genes in a population. *Genetics*, 47,
576 713-719.

29.
578 Koskella, B. (2014). Bacteria-phage interactions across time and space: merging local adaptation and time-shift experiments to understand phage evolution. *Am. Nat.*, 184, S9-S21.

580 30.
Koskella, B. & Brockhurst, M.A. (2014). Bacteria-phage coevolution as a driver of ecological and
582 evolutionary processes in microbial communities. *FEMS Microbiol. Rev.*, 38, 916-931.

31.
584 Lennon, J.T. (2007). Is there a cost of viral resistance in marine cyanobacteria? *ISME J.*, 1, 300-312.

586 32.
Lennon, J.T. & Martiny, J.B.H. (2008). Rapid evolution buffers ecosystem impacts of viruses in a
588 microbial food web. *Ecol. Lett.*, 11, 1178-1188.

33.
590 Lenski, R.E. & Levin, B.R. (1985). Contraints on the coevolution of bacteria and virulent phage - a model, some experiments, and predictions for natural communities. *Am. Nat.* 125, 585-
592 602.

34.
594 Leon, M. & Bastias, R. (2015). Virulence reduction in bacteriophage resistant bacteria. *Front. Microbiol.*, 6.

596 35.
Marston, M.F., Pierciey, F.J., Shepard, A., Gearin, G., Qi, J., Yandava, C. *et al.* (2012). Rapid
598 diversification of coevolving marine *Synechococcus* and a virus. *Proc. Natl Acad. Sci. U S A,*, 109, 4544-4549.

600 36.
Menge, D.N.L. & Weitz, J.S. (2009). Dangerous nutrients: Evolution of phytoplankton resource
602 uptake subject to virus attack. *J Theor Bio*, 257, 104-115.

37.
604 Monier, A., Chambouvet, A., Milner, D.S., Attah, V., Terrado, R., Lovejoy, C. *et al.* (2017). Host-derived viral transporter protein for nitrogen uptake in infected marine
606 phytoplankton. *Proc. Natl Acad. Sci. U S A*, 114: E7489–E7498.

38.
608 Noble, R.T. & Fuhrman J.A. (1998). Use of SYBR Green 1 for rapid epiluorescence count of marine viruses and bacteria. *Aquat. Microb. Ecol.* 14, 113-118.

610 39.
Pascua, L.L., Hall, A.R., Best, A., Morgan, A.D., Boots, M. & Buckling, A. (2014). Higher
612 resources decrease fluctuating selection during host-parasite coevolution. *Ecol. Lett.*, 17, 1380-1388.

614 40.
Rhee, G.Y. (1978). Effects of N:P atomic ratios and nitrate limitation on algal growth cell
616 composition, and nitrate uptake. *Limnol. Oceanogr.*, 23, 10-25.

41.

618   Schwartz, D.A. & Lindell, D. (2017). Genetic hurdles limit the arms race between *Prochlorococcus* and the T7-like podoviruses infecting them. *ISME J.*, 11, 1836-1851.

620   42.

Smith, V.H. & Holt, R.D. (1996). Resource competition and within-host disease dynamics. *Trends Ecol. Evol.*, 11, 386-389.

43.

624   Sterner, R.W. & Elser, J.J. (2002). *Ecological Stoichiometry: The Biology of Elements from Molecules to the Biosphere*. Princeton University Press.

626   44.

Stoddard, L.I., Martiny, J.B.H. & Marston, M.F. (2007). Selection and characterization of cyanophage resistance in marine *Synechococcu*s strains. *Appl. Environ. Microbiol.*, 73, 5516-5522.

630   45.

Suttle, C.A. (2007). Marine viruses - major players in the global ecosystem. *Nat. Rev. Microbiol.*, 5, 801-812.

46.

634   Turner, C.B., Wade, B.D., Meyer, J.R., Sommerfeld, B.A. & Lenski, R.E. (2017). Evolution of organismal stoichiomery in a long-term experiment with *Escherichia coli*. *R. Soc. Open Sci.*, DOI: 10.1098/rsos.170497

47.

638   Van Mooy, B.A.S., Rocap, G., Fredricks, H.F., Evans, C.T. & Devol, A.H. (2006). Sulfolipids dramatically decrease phosphorus demand by picocyanobacteria in oligotrophic marine environments. *Proc. Natl Acad. Sci. U S A*, 103, 8607-8612.

48.

642   Weitz, J.S., Poisot, T., Meyer, J.R., Flores, C.O., Valverde, S., Sullivan, M.B. *et al.* (2013). Phage-bacteria infection networks. *Trends Microbiol.*, 21, 82-91.

644   49.

Wilson, W.H., Carr, N.G. & Mann, N.H. (1996). The effect of phosphate status on the kinetics of cyanophage infection in the oceanic cyanobacterium *Synechococcus* sp WH7803. *J. Phycol.*, 32, 506-516.

648   50.

Yamamichi, M., Meunier, C.L., Peace, A., Prater, C. & Rua, M.A. (2015). Rapid evolution of a consumer stoichiometric trait destabilizes consumer-producer dynamics. *Oikos*, 124, 960-969.

652   51.

Yoshida, T., Ellner, S.P., Jones, L.E., Bohannan, B.J.M., Lenski, R.E. & Hairston, N.G. (2007). Cryptic population dynamics: Rapid evolution masks trophic interactions. *PLoS. Biol.*, 5, 1868-1879.

656   52.

Zeng, Q. & Chisholm, S.W. (2012). Marine viruses exploit their host's two-component regulatory system in response to resource limitation. *Curr Biol*, 24.

FIGURE CAPTIONS

Fig. 1. Microbial community dynamics were affected by nutrient stoichiometry. *Synechococcus*

664   and phage densities were tracked in replicate (n = 3) chemostats receiving nitrogen (N)- or

phosphorus (P)-limited media. Vertical lines at day 0 indicate time of phage amendment. See

666   Fig. S2 for *Synechococcus* dynamics in the no-phage control chemostats. Data are represented as

mean ± SEM.

Fig. 2 Average phage-resistance of *Synechococcus* strains increased monotonically over time

670   under both nitrogen (N)- and phosphorus (P)-limitation (left). Resistance was calculated as the

proportion of unsuccessful infections between all phage strains from the experiment when

672   challenged against *Synechococcus* strains that had been isolated at a specific time point and is

presented as the chemostat mean ± SEM. Average infectivity of phage over time (right) was

674   calculated as the proportion of successful infections between all *Synechococcus* strains across the

experiment when challenged against phage strains that had been isolated (on the ancestral host)

676   at a specific time point and is presented as the chemostat mean ± SEM.

678   **Fig. 3** Phenotypic coevolution between hosts (*Synechococcus*) and phage was affected by

nutrient stoichiometry. We calculated infectivity based on the proportion of successful infections

680   between *Synechococcus* strains and phage strains that were isolated from chemostats at different

time points. Infectivity is proportional to the width of the edges (lines) connecting nodes

682   (symbols). Black squares correspond to phage isolated from the phage-amended chemostats,

white circles correspond to *Synechococcu*s isolated from phage-amended chemostats, and grey

684   circles correspond to naive *Synechococcus* isolated from no-phage control chemostats. The

absence of a line indicates that *Synechococcus* isolates were resistant to a particular phage

686   challenge.

688   **Fig. 4.** Time-shift analysis of host-phage infectivity reveals the effects of stoichiometry on

coevolution. Contemporary interactions (i.e., those between a host and phage strain isolated at

690   the same time point) are centered at time zero (grey vertical line) along the time-shift

(horizontal) axis. Interactions with past phage are shifted to the left (negative values) and

692   interactions with future phage are shifted to the right (positive values). Each black line

corresponds with the mean infectivity for *Synechococcus* isolated from a specific time point as

694   indicated by the open circle containing the isolation day (-6, 9, 23, 72, 100, 129, or 166). When

comparing challenges between hosts and phage from phage-amended chemostats (a, b),

696   infectivity was weak for hosts that were isolated after day 23 or when challenged against phage

from the past owing to the evolution of resistance. Such findings are consistent with arms-race

698   dynamics where directional selection gives rise to escalating host resistance. We also challenged

phage against naive hosts from the no-phage control chemostats (c, d). From this, we found that

700   infectivity was significantly higher under nitrogen (N)-limitation than phosphorus (P)-limitation,

but overall, was lower than what would be expected under arms-race dynamics. Instead, the

702   fluctuations in infectivity with respect to time-shift are consistent with negative frequency-

dependent selection and reflect asymmetry in the coevolution between *Synechococcus* and

704   phage.

706   **Fig. 5.** Host-phage infection networks based on interactions between *Synechococcus* and phage

isolates that coevolved under nitrogen (N)- and phosphorus (P)-limitation. Networks were

708   significantly nested, consistent with expectations of host-phage systems coevolving under arms

race dynamics. However, the degree of nestedness was not affect by stoichiometry. Left panel:

710   Networks were also significantly modular when compared to randomized networks, but the

degree of modularity was significantly greater in P-limited networks. Right panel: Infection

712   networks reflecting the modular structure of host-phage interactions within each chemostat. Each

grouping of colored cells within a grey-colored background panel corresponds to a calculated

714   module within the interaction network of N-limited (a – c) or P-limited (d – f) chemostats. That

is, colored cells correspond to infectious interactions that could not be regrouped into a

716   calculated module based on the network algorithm.

Fig. 1

Fig. 2

Fig. 3

740   **Fig. 5**