1   **Title:** Genetic basis of body color and spotting pattern in redheaded pine sawfly larvae
2   (*Neodiprion lecontei*)
3

4   **Authors:** Catherine R. Linnen[*], Claire T. O'Quin[*], Taylor Shackleford[*], Connor R.
5   Sears[*†], Carita Lindstedt[‡]
6

7   **Running title:** Genetic basis of larval color
8
9   **Keywords:** pigmentation, carotenoids, melanin, genetic architecture, convergent
10   evolution, evolutionary genetics
11

12   **Corresponding author:** Catherine R. Linnen, 204E Thomas Hunt Morgan Building,
13   Lexington, KY 40506, 859-323-3160, Catherine.linnen@uky.edu

[*] Department of Biology, University of Kentucky, Lexington, KY, 40506
[†] Current affiliation: Department of Biological Sciences, University of Cincinnati,
Cincinnati, OH, 45221
[‡] Centre of Excellence in Biological Interactions, Department of Biological and
Environmental Sciences, University of Jyväskylä, Jyväskylä, Finland, FI-40014

14   ABSTRACT
15
Pigmentation has emerged as a premier model for understanding the genetic basis of
phenotypic evolution, and a growing catalog of color loci is starting to reveal biases in
the mutations, genes, and genetic architectures underlying color variation in the wild.
However, existing studies have sampled a limited subset of taxa, color traits, and
developmental stages. To expand our sample of color loci, we performed quantitative
trait locus (QTL) mapping analyses on two types of larval pigmentation traits that vary
among populations of the redheaded pine sawfly (*Neodiprion lecontei*): carotenoid-based
yellow body color and melanin-based spotting pattern. For both traits, our QTL models
explained a substantial proportion of phenotypic variation and suggested a genetic
architecture that is neither monogenic nor highly polygenic. Additionally, we used our
linkage map to anchor the current *N. lecontei* genome assembly. With these data, we
identified promising candidate genes underlying: (1) a loss of yellow pigmentation in
Mid-Atlantic/northeastern populations (*Cameo2* and *apoLTP-II/I*), and (2) a pronounced
reduction in black spotting in Great-Lakes populations (*yellow, TH, Dat*). Several of
these genes also contribute to color variation in other wild and domesticated taxa.
Overall, our findings are consistent with the hypothesis that predictable genes of large-
effect contribute to color evolution in nature.

33
34   INTRODUCTION
35
36   Over the last century, color phenotypes have played a central role in our efforts to
understand how evolutionary processes shape phenotypic variation in natural populations
(Gerould 1921; Sumner 1926; Fisher and Ford 1947; Haldane 1948; Cain and Sheppard
1954; Kettlewell 1955). More recently, as technological advances have enabled us to link
genotypes and phenotypes in non-model organisms (Davey *et al.* 2011; Gaj *et al.* 2013;
Goodwin *et al.* 2016; Huang *et al.* 2016), we've begun to decipher the genetic and
developmental mechanisms underlying naturally occurring color variation as well (True
2003; Protas and Patel 2008; Wittkopp and Beldade 2009; Manceau *et al.* 2010; Nadeau
and Jiggins 2010; Kronforst *et al.* 2012). With a growing catalog of color loci, we are
starting to assess how ecology, evolution, and development interact to bias the genetic
architectures, genes, and mutations underlying the remarkable diversity of color
phenotypes in nature (Hoekstra and Coyne 2007; Stern and Orgogozo 2008, 2009; Kopp
2009; Manceau *et al.* 2010; Streisfeld and Rausher 2011; Martin and Orgogozo 2013;
Dittmar *et al.* 2016; Massey and Wittkopp 2016; Martin and Courtier-Orgogozo 2017).
To make robust inferences about color evolution, however, we require genetic data from
diverse traits, taxa, developmental stages, and evolutionary timescales.
52   Two long-term goals of research on the genetic underpinnings of pigment
variation are: (1) to evaluate the importance of large-effect loci to phenotypic evolution
(Orr and Coyne 1992; Mackay *et al.* 2009; Rockman 2012; Remington 2015; Dittmar *et
al.* 2016), and (2) to determine the extent to which the independent evolution of similar
phenotypes (phenotypic convergence) is attributable to the same genes and mutations
(genetic convergence) (Arendt and Reznick 2008; Gompel and Prud'homme 2009;
Christin *et al.* 2010; Manceau *et al.* 2010; Elmer and Meyer 2011; Conte *et al.* 2012;
Rosenblum *et al.* 2014). Addressing these questions will require a large, unbiased sample

of pigmentation loci. At present, most of what we know about the genetic basis of color
variation in animals comes from a handful of taxa and from melanin-based color
variation in adult life stages (True 2003; Protas and Patel 2008; Wittkopp and Beldade
2009; Nadeau and Jiggins 2010; Kronforst *et al.* 2012; Sugumaran and Barek 2016).
Another important source of bias in our existing sample of pigmentation loci stems from
a tendency to focus on candidate genes and discrete pigmentation phenotypes (Kopp
2009; Manceau *et al.* 2010; Rockman 2012; but see O'Quin *et al.* 2013; Albertson *et al.*
2014; Signor *et al.* 2016; Yassin *et al.* 2016). Here, we describe an unbiased, genome-
wide analysis of multiple, continuously varying color traits in larvae from the order
Hymenoptera, a diverse group of insects that is absent from our current catalog of color
loci (Martin and Orgogozo 2013).
More specifically, our study focuses on pine sawflies in the genus *Neodiprion*.
Several factors make *Neodiprion* an especially promising and tractable system for
investigating the genetic basis of color variation. First, there is extensive variation in
many different types of color traits, and these vary both within and between *Neodiprion*
species (Figures 1-2). Second, previous phylogenetic and demographic studies of this
genus (Linnen and Farrell 2007, 2008a; b; Bagley *et al.* 2017) enable us to infer
directions of trait change and identify instances of phenotypic convergence. Third,
because many different species can be reared and crossed in the lab (Knerer and Atwood
1972, 1973; Kraemer and Coppel 1983; Bendall *et al.* 2017), unbiased genetic mapping
approaches are feasible in *Neodiprion*. Fourth, a growing list of genomic resources for
Neodiprion—including an annotated genome and a methylome for *N. lecontei* (Vertacnik
et al. 2016; Glastad *et al.* 2017)— facilitate identification of causal genes and mutations.
And finally, an extensive natural history literature (Coppel and Benjamin 1965; Knerer
and Atwood 1973) provides insights into the ecological functions of color variation in
pine sawflies, which we review briefly for context.
Under natural conditions, pine sawfly larvae are attacked by a diverse assemblage
of arthropod and vertebrate predators, by a large community of parasitoid wasps and
flies, and by fungal, bacterial, and viral pathogens (Coppel and Benjamin 1965; Wilson *et*
al. 1992; Codella and Raffa 1993). When threatened, larvae adopt a characteristic "U-
bend" posture and regurgitate a resinous defensive fluid (Figure 1), which is an effective
repellant against many different predators and parasitoids (Eisner *et al.* 1974; Codella and
Raffa 1995; Lindstedt *et al.* 2006, 2011). Although most *Neodiprion* species are
chemically defended, larvae vary from a green striped morph that is cryptic against a
background of pine foliage to highly conspicuous aposematic morphs with dark spots or
stripes overlaid on a bright yellow or white background (Figure 1). Thus, larval color is
likely to confer protection against predators either via preventing detection (crypsis) or
advertising unpalatability (aposematism) (Ruxton *et al.* 2004; Lindstedt *et al.* 2011).
Beyond selection for crypsis or aposematism, there are a myriad of abiotic and
biotic selection pressures that could act on *Neodiprion* larval color. For example,
coloration plays diverse ecological roles in insects, including thermoregulation,
protection against UV damage, desiccation tolerance, and resistance to abrasion (True
2003; Lindstedt *et al.* 2009; Wittkopp and Beldade 2009). Color alleles may also have
pleiotropic effects on other traits, such as behavior, immune function,
diapause/photoperiodism, fertility, and developmental timing (True 2003; Wittkopp and
Beldade 2009; Heath *et al.* 2013; Lindstedt *et al.* 2016). Temporal and spatial variation in

these diverse selection pressures likely contribute to the abundant color variation in the
genus *Neodiprion*.
As a first step to understanding the proximate mechanisms that generate color
variation in pine sawflies, we conducted a quantitative trait locus (QTL) mapping study
of larval body color and larval spotting pattern in the redheaded pine sawfly, *Neodiprion*
lecontei. This species is widespread across eastern North America, where it feeds on
multiple pine species. Throughout most of this range, larvae have several rows of dark
black spots overlaid on a bright, yellow body (*e.g.*, center image in Figure 1). However,
previous field surveys and demographic analyses indicate that there has been a loss of
yellow pigmentation in some populations in the Mid-Atlantic/northeastern United States
and a pronounced reduction in spotting in populations living in the Great-Lakes region of
the U.S. and Canada (Bagley *et al.* 2017; Figure 2). To describe genetic architectures and
identify candidate genes underlying these two reduced-pigmentation phenotypes, we
crossed white-bodied, dark-spotted individuals from a Virginia population to yellow-
bodied, light-spotted individuals from a Michigan population (Figure 2). After mapping
larval body-color and spotting-pattern traits in recombinant F_2 haploid males, we used
our linkage map to anchor the current *N. lecontei* genome assembly and identified
candidate color genes within our QTL intervals. We conclude by comparing these initial
findings on color genetics in pine sawflies with published studies from other insect taxa.
125
126   MATERIALS AND METHODS
127
Cross Design
To investigate the genetic basis of sawfly color traits, we crossed *Neodiprion*
lecontei females from a white-bodied, dark-spotted population (collected from Valley
View, VA; 37°54'47"N, 79°53'46"W) to *N. lecontei* males from a yellow-bodied, light-
spotted population (collected from Bitely, MI; 43°47'46"N, 85°44'24"W). Both
populations were collected from the field in 2012 and reared on *Pinus banksiana* (jack
pine) for at least two generations in the lab via standard rearing protocols (described in
more detail in Harper *et al.* 2016; Bendall *et al.* 2017). Our mapping families were
derived from four grandparental pairs, which produced 10 F_1 females. Like most
hymenopterans, *N. lecontei* adults reproduce via arrhenotokous haplodiploidy, in which
unfertilized eggs develop into haploid males and fertilized eggs develop into diploid
females (Heimpel and de Boer 2008; Harper *et al.* 2016). Therefore, to produce an F_2
haploid generation, we allowed virgin F_1 females to lay eggs and reared their haploid
male progeny on *P. banksiana* foliage until they reached a suitable size for phenotyping.
142
Color phenotyping
N. lecontei larvae pass through five (males) or six (females) feeding instars and a
single non-feeding instar, which are distinguishable on the basis of color pattern and size
(Benjamin 1955; Coppel and Benjamin 1965; Wilson *et al.* 1992). For phenotyping, we
chose only mature feeding larvae, which have an orange-red head capsule with a black
ring around each eye and up to four pairs of rows of black spots (Wilson *et al.* 1992).
Immediately after phenotyping, we preserved each larva in 100% ethanol for molecular
work. In total, we generated color-phenotype data for 30 individuals from the VA

population (mixed sex), 30 individuals from the MI population (mixed sex), 47 F_1
females, and 429 F_2 males (progeny of 10 virgin F_1 females).
153
Larval body color
We quantified larval body color in two ways: spectrometry and digital
photography. First, we immobilized larvae with CO_2 and recorded 15 reflectance
spectra—five measurements from each of three body regions: the dorsum, lateral side,
and ventrum—using a USB2000 spectrophotometer with a PX-2 Pulsed Xenon light
source and SpectraSuite software (Ocean Optics, Largo, FL). We then used the program
CLR: Colour Analysis Programs v1.05 (Montgomerie 2008) to trim the raw reflectance
data to 300-700nm, compute 1-nm bins, and to compute the following summary statistics
(described in more detail in the CLR documentation): B1, B2, B3, S1R, S1G, S1B, S1U,
S1Y, S1V, S3, S5a, S5b, S5c, S6, S7, S8, S9, H1, H3, H4a, H4b, and H4c. This
procedure produced 15 estimates for each summary statistic for each larva, which we
then averaged to obtain a single value per statistic per larva. To reduce the 22 color
summary statistics to a smaller number of independent variables, we performed a
principal component analysis (PCA). To ensure that each variable was normally
distributed, we performed a normal-quantile transformation prior to performing the PCA.
Based on examination of the resulting scree plot, we retained the first two principal
components (PC1 and PC2), which together accounted for 73.7% of the variance in the
spectral data. Based on factor loadings, PC1 (PVE: 48.2%) corresponds to saturation
(S1G, S1Y, S8) and brightness (B1, B2), while PC2 (PVE: 25.5%) corresponds to
saturation (S5A, S5B, S5C, S6) and hue (H4c) (Table S1). Unless otherwise noted, the
PCA and all other statistical analyses were performed in R (Version 3.1.3 or 3.2.3).
Because it provides objective and information-rich data, spectrometry is generally
considered to be the gold standard for color quantification (Endler 1990; Andersson and
Prager 2006). However, despite efforts to keep background lighting and reflectance probe
positioning as consistent as possible across samples, measurement of small patches of
yellow on rounded larval bodies proved challenging. Given the potential for variation in
reflectance probe position to introduce substantial noise into our data (Montgomerie
2006), we used digital photography as a complementary method for quantifying larval
body color (Stevens *et al.* 2007). For this analysis, we photographed CO_2-immobilized
larvae (dorsal and lateral surfaces) with a Canon EOS Rebel t3i camera equipped with an
Achromat S 1.0X FWD 63mm lens and mounted on a copy stand. All photographs were
taken in a dark, windowless room, and larvae were illuminated by two SLS CL-150 copy
lamp lights, each with a 23-Watt (100W equivalent) soft white compact fluorescent light
bulb. We then used Adobe Photoshop CC 2014 or 2015 (Adobe Systems Incorporated,
San Jose, CA) to ascertain the amount of yellow present, following O'Quin *et al.* (2013).
First, we converted each digital image (lateral surface) to CMYK color mode. Next, we
selected the eye-dropper tool (set to a size of 5x5 pixels) as the color sampler tool, which
we used to sample three different body locations: the body just behind the head and
parallel to the eye, the first proleg, and the anal proleg. For each of the three regions, this
procedure yielded an estimate of the proportion of the selected area that was yellow. We
then averaged the three measurements to produce a single final measurement of yellow
pigmentation (hereafter referred to as "yellow").
196

Larval spotting pattern

On each side, *N. lecontei* larvae have up to four rows of up to 11 spots extending from the mesothorax to the 9th abdominal segment (Wilson *et al.* 1992). We quantified the extent of this black spotting in two ways: the number of spots present (spot number) and the proportion of body area covered by spots (spot area). To quantify spot number, we simply counted the number of spots in digital images of the dorsal and lateral surfaces (one side only). To quantify larval spotting area, we used Adobe Photoshop's quick-selection tool to measure the area of the larval body (minus the head capsule) and the area of each row of lateral black spots. To control for differences in larval size, we divided the summed area of all lateral black spots by the area of the larval body. We also used the larval images and Adobe Photoshop to calculate the area of the head capsule, which we used as a covariate in some analyses to control for larval size (see below). We used a custom Perl script to process Photoshop measurement output files in bulk (written by John Terbot II; available in File S3).

Statistical analysis of phenotypic data

In total, we produced three measures of body color (PC1, PC2, and yellow) and two measures of spotting pattern (spot number and spot area). To determine whether mean phenotypic values for the five color traits differed between the two populations and among the three generations of our cross, we performed Welch's two-tailed *t*-tests, followed by Bonferroni correction for multiple comparisons. To determine the extent to which these five traits co-varied in the F$_2$ males, we calculated Pearson's correlation coefficients (*r*) and associated *P*-values, followed by Bonferroni correction for multiple comparisons. To determine which covariates to include in our QTL models, we performed ANOVAs to evaluate the relationship between each phenotype in the F$_2$ males (429 total) and their F$_1$ mothers (10 total) and head capsule sizes (a proxy for larval size/developmental stage). These and all other statistical analyses were performed in R version 3.3.2 (R Core Team 2013)

Genotyping

We extracted DNA from ethanol-preserved larvae using a modified CTAB method (Chen *et al.* 2010) and prepared barcoded and indexed double-digest RAD (ddRAD) libraries using methods described elsewhere (Peterson *et al.* 2012; Bagley *et al.* 2017). We prepared a total of 10 indexed libraries: one consisting of the eight grandparents and 10 F$_1$ females (18 adults total), and the remaining nine consisting of F$_2$ haploid male larvae (~48 barcoded males per library). After verifying library quality using a Bioanalyzer 2100 (Agilent, Santa Clara, CA), we sent all 10 libraries to the University of Illinois Urbana-Champaign Roy J. Carver Biotechnology Center (Urbana, IL), where the libraries were pooled and sequenced using 100-bp single-end reads on two Illumina HiSeq2500 lanes. In total, we generated 400,621,900 reads.

We de-multiplexed and quality-filtered raw reads using the protocol described in Bagley *et al.* 2017. We then used Samtools v0.1.19 (Li *et al.* 2009) to map our reads to our *N. lecontei* reference genome (Vertacnik *et al.* 2016) and STACKS v1.37 (Catchen *et al.* 2013) to extract loci from our reference alignment and to call SNPs. We called SNPs in two different ways. First, for QTL mapping analyses, our goal was to recover fixed differences between the grandparental lines. To do so, we first called SNPs in our eight

grandparents and 10 F_1 mothers. For these 18 individuals, we required that SNPs had a
minimum of 7x coverage and no more than 12% missing data. We then compiled a list of
SNPs that represented fixed differences and, as an additional quality check, confirmed
that all F_1 females were heterozygous at these loci. We then used STACKS to call SNPs
in the F_2 haploid males, requiring that each SNP had a minimum of 5x coverage, no more
than 10% missing data, and was present in the curated list produced from the
grandparents. Filtering in STACKs produced 559 SNPs genotyped in 429 F_2 males.
250   Second, to maximize the number of SNPs available for genome scaffolding, we
performed an additional STACKS run using only the F_2 haploid males, requiring that
each SNP had a minimum of 4x coverage. By removing the requirement that SNPs were
called in all grandparents, we could recover many more SNPs. We then filtered the data
in VCFtools v0.1.14 (Danecek *et al.* 2011) to remove individuals with a mean depth of
coverage less than one, retaining 408 F_2 males. After removing low-coverage individuals,
we used VCFtools to remove sites with a minor allele frequency (MAF) less than 0.05,
sites with >5 heterozygotes (in a sample of haploid males, high heterozygosity is a clear
indication of pervasive genotyping error), and sites with more than 50% missing data.
259
260   **Linkage map construction and genome scaffolding**
261   To construct a linkage map for interval mapping, we started with 559 SNPs
scored in 429 F_2 males. After an additional round of filtering in R/qtl (Broman and Sen
2009), we removed 11 haploid males that had >50% missing data, for a total of 418 F_2
males. Additionally, after removing SNPs that were genotyped in <70% of individuals,
had identical genotypes to other SNPs, and had distorted segregation ratios (at $\alpha < 0.05$,
after Bonferroni correction for multiple testing), we retained 503 SNPs. To assign these
markers to linkage groups, we then used the "formLinkageGroups" function, requiring a
minimum logarithm of odds (LOD) score of 6.0 and a maximum recombination
frequency of 0.35. To order markers on linkage groups, we used the "orderMarkers"
function, with the Kosambi mapping function to allow for crossovers. Following this
initial ordering, we performed rippling on each linkage group to check whether switching
marker order could improve LOD scores.
273   Our initial map included 503 SNPs spread across 358 scaffolds (out of 4523
scaffolds; Vertacnik *et al.* 2016). To increase the number of scaffolds and bases that we
could place on our linkage groups, we performed additional linkage mapping analyses
with a larger SNP dataset that was called as described above. We then constructed a
linkage map for each of our four grandparental families (N = 54, 73, 120, and 161). For
each grandparental family, we first performed additional data filtering in R/qtl to remove
duplicate SNPs and SNPs with distorted segregation ratios, retaining between 2,049 and
3,155 SNPs per family. We then used the "formLinkageGroups" command with variable
LOD thresholds (range: 5-15) and a maximum recombination frequency of 0.35. Because
SNPs were not coded according to grandparent of origin, many alleles were "switched".
We therefore performed an iterative process of linkage group formation, visualization of
pairwise recombination fractions and LOD scores ("plotRF" command), and allele
switching ("switchAlleles" command) until we obtained seven linkage groups (the
number of *N. lecontei* chromosomes; Smith 1942; Maxwell 1958; Sohi and Ennis 1981)
and a recombination/LOD plot indicative of linkage within, but not between, linkage
groups. Because allele ordering and examination of alternative SNP orders for these

larger panels of SNPs was prohibitively slow in R/qtl, we used the more efficient
MSTmap algorithm, implemented in R/ASMap v0.4-7 (Taylor and Butler 2017), to order
our markers along their assigned linkage groups.
Finally, to order and orient our genome scaffolds along linkage groups
(chromosomes), we used ALLMAPS (Tang *et al.* 2015) to combine information from our
five maps (initial map with all individuals, but limited markers; plus four additional
maps, each with more markers, but fewer individuals). Because maps constructed from
larger families are likely to be more accurate than those constructed from small families,
we weighted the maps according to their sample sizes.
298
Interval mapping analysis
After linkage map construction, we used R/qtl to map QTL for our five color
traits. Based on analyses of phenotypic covariates (Table S2), we included F_1 mothers
and head-capsule size as covariates in our analyses of PC1, spot number, and spot area;
F_1 mothers as covariates in our analysis of PC2; and no covariates in in our analysis of
yellow. For each trait, we performed interval mapping using multiple imputation
mapping. We first used the "sim.geno" function with a step size of 0 (*i.e.*, genotypes only
drawn at marker locations) and 64 replicates. We then used the "stepwiseqtl" command
to detect QTL and select the multiple QTL model that optimized the penalized LOD
score (Broman and Sen 2009; Manichaikul *et al.* 2009). To obtain penalties for the
penalized LOD scores, we used the "scantwo" function to perform 1,000 permutations
under a two-dimensional, two-QTL model that allows for interactions between QTL and
the "calc.penalties" function to calculate penalties from these permutation results, using a
significance threshold of $\alpha = 0.05$. Finally, for each QTL retained in the final model, we
calculated a 1.5-LOD support interval.
314
Candidate gene analysis
As a first step to moving from QTL intervals to causal loci, we compiled a list of
candidate color genes and determined their location in the *N. lecontei* genome. For larval
spotting, we included genes in the melanin synthesis pathway and genes that have been
implicated in pigmentation patterning (Wittkopp *et al.* 2003; Protas and Patel 2008;
Wittkopp and Beldade 2009; Sugumaran and Barek 2016). For larval body color, we
included genes implicated in the transport, deposition, and processing of carotenoid
pigments derived from the diet (Palm *et al.* 2012; Yokoyama *et al.* 2013; Tsuchida and
Sakudoh 2015; Toews *et al.* 2017). Although several pigments can produce yellow
coloration in insects (*e.g.,* melanins, pterins, ommochromes, and carotenoids), we
focused on carotenoids because a heated pyridine test (McGraw *et al.* 2005) was
consistent with carotenoid-based coloration in *N. lecontei* larvae (Figure S1).
Once we had compiled a list of candidate genes, we searched for these genes by
name in the *N. lecontei* v1.0 genome assembly and NCBI annotation release 100
(Vertacnik *et al.* 2016). To find missing genes and as an additional quality measure, we
obtained FASTA files corresponding to each candidate protein and/or gene from NCBI
(using *Apis, Drosophila melanogaster,* or *Bombyx mori* sequences, depending on
availability). We then used the i5k Workspace@NAL (Poelchau *et al.* 2014) BLAST
(Altschul *et al.* 1990) web application to conduct tblastn (for protein sequences) or tblastx
(for gene sequences) searches against the *N. lecontei* v1.0 genome assembly, using

default search settings. After identifying the top hit for each candidate gene/protein, we
then used the WebApollo (Lee *et al.* 2013) JBrowse (Skinner *et al.* 2009) *N. lecontei*
genome browser to identify the corresponding predicted protein coding genes (from
NCBI annotation release 100) in the *N. lecontei* genome.
We took additional steps to identify genes in the *yellow* gene family, all of which
contain a major royal jelly protein (MRJP) domain (Ferguson *et al.* 2011). First, we used
the search string "major royal jelly protein *Neodiprion*" to search the NCBI database for
all predicted *yellow*-like and *yellow-MRJP*-like *N. lecontei* genes. We then downloaded
FASTA files for the putative *yellow* gene sequences (26 total). Next, we used the
Hymenoptera Genome Database (Elsik *et al.* 2016) to conduct a blastx search of our *N.*
lecontei gene sequence queries against the *Apis mellifera* v4.5 genome NCBI RefSeq
annotation release 103. Finally, we recorded the top *A. mellifera* hit for each putative *N.*
lecontei yellow gene.
348
Data availability
Short-read DNA sequences will be made available via the NCBI SRA (Bioproject
PRJNA#######, Biosample numbers SAMN########-SAMN########). The linkage-
group anchored assembly will be submitted to NCBI and i5k to update the existing *N.*
lecontei genome assembly and annotations (Vertacnik *et al.* 2016). File S3 contains the
custom script used to process the raw spot-area data. File S4 contains phenotypic data
from all generations. File S5 contains the input file for R/qtl. Files S6 and S7 contain the
genotype data and linkage maps used in the scaffolding analysis, respectively.
357
358   RESULTS AND DISCUSSION
359
Variation in larval color traits
Lab-reared larvae derived from the two founding populations differed
significantly from one another for all color phenotypes (Figures 2, 3; Table S3). Because
all larvae were reared on the same host under the same laboratory conditions (*i.e.*,
minimal environmental variance), these results suggest that genetic variance contributes
to variance in these larval color traits. Crosses between the VA and MI lines produced
diploid F_1 female larvae that were intermediate in both body color and spotting pattern
(Figures 2, 3). With the exception of PC2, F_1 larvae differed significantly from both
grandparents for all color traits (Table S3), indicating a lack of complete dominance for
these traits. In contrast, the yellow, MI-like hue/chroma (PC2) appears to be completely
dominant to the white, VA-like hue/chroma. We can make additional inferences about
trait dominance by comparing the phenotypes of diploid F_1 female larvae (dominance
effects present) to those of the haploid F_2 male larvae (dominance effects absent). For all
three body-color measures (yellow, PC1, and PC2), average F_1 larval colors were more
MI-like (yellow) than those of F_2 larvae. In contrast, compared to the F_2 larvae, spotting
phenotypes (number and area) of F_1 larvae are more similar to the VA population (dark-
spotted). Taken together, these phenotypic data suggest that the more heavily pigmented
body-color and spotting-pattern phenotypes (*i.e.,* large values for yellow, spot number,
and spot area; small values for PC1 and PC2) are partially dominant to the less pigmented
phenotypes.

Larval color variation observed in the F_2 males spanned—and even exceeded—
the range of variation observed in the grandparental populations and F_1 females (Figure
3). The observation that grandparental body-color and spotting-pattern phenotypes are
recapitulated in the F_2 males suggests that both types of traits are controlled by a
relatively small number of loci. There are multiple, non-mutually exclusive explanations
for the transgressive color phenotypes in our haploid F_2 larvae, including: variation in the
grandparental lines, reduced developmental stability in hybrids, epistasis, unmasking of
recessive alleles in haploid males, and the complementary action of additive alleles from
the two grandparental lines (Rieseberg *et al.* 1999).
We also observed significant correlations between many of the color traits (Figure
S2). Encouragingly, estimates of body color derived from digital images (yellow)
correlated significantly with both estimates derived from spectrometry (PC1 and PC2),
suggesting that these independent data sources describe the same underlying larval color
phenotype (*i.e.,* amount of yellow pigment). Similarly, we observed strong correlations
between the number and area of larval spots, indicating that both measures reliably
characterize the extent of melanic spotting on the larval body. We also found that larvae
with yellower bodies (small values for PC2) tended to be less heavily spotted (Figure S2).
This observation suggests that body color and spotting pattern do not evolve completely
independently of one another. Nevertheless, the correlation between these traits was weak
and we observed many different combinations of spotting and pigmentation in the
recombinant F_2 males (Figure 2).
401
402 Linkage mapping and genome scaffolding
Our 503 SNP markers were spread across seven linkage groups (LGs), which
matches the number of *N. lecontei* chromosomes (Smith 1941; Maxwell 1958; Sohi and
Ennis 1981). The total map length was 1169 cM, with an average marker spacing of 2.4
cM and maximum marker spacing of 24.3 cM (Table S4; Figure 4). Together, these
results indicate that this linkage map is of sufficient quality and coverage for interval
mapping. Additionally, with an estimated genome size of 340 Mb (estimated via flow
cytometry; C. Linnen, personal observation), these mapping results yield a recombination
density estimate of 3.43 cM/Mb. This recombination rate is lower than that observed in
social hymenopterans, which have among the highest rates of recombination in
eukaryotes (Wilfert *et al.* 2007). Nevertheless, this rate is on par with that reported in
other (non-eusocial) hymenopterans, which lends support to the hypothesis that elevated
recombination rates in eusocial hymenopteran species is a derived trait and possibly an
adaptation to a social lifestyle (Gadau *et al.* 2000; Schmid-Hempel 2000; Crozier and
Fjerdingstad 2001).
Linkage maps estimated for the four grandparental families, each of which
contained >2000 markers, ranged in length from 1072 cM to 3064 cM (Table S4). This
variation in map length is likely attributable to both decreased mapping accuracy in
smaller families and decreased genotyping accuracy in these less-stringently filtered SNP
datasets. Nevertheless, our scaffolding analysis revealed that marker ordering was highly
consistent across linkage maps (Figures S3-S9). Additionally, via including more SNPs,
we were able to more than triple the number of mapped scaffolds (from 358 to 1005) and
increase the percentage of mapped bases from 41.2% to 78.9% (Tables S5-S6).

Anchored genome scaffolds, coupled with existing *N. lecontei* gene annotations, are a
valuable resource for identification of candidate genes within QTL.
427
Genetic architectures of larval body color and larval spotting pattern
Combining the results from our three measures of larval body-color (yellow, PC1,
and PC2), our interval mapping analyses detected a total of six distinct QTL regions and
one QTL x QTL interaction (Table 1; Figures 4,5A). Of these, the largest-effect QTL
were on LGs 3 and 5. We also recovered a significant interaction between these QTL for
yellow (Figure 5B). Compared to analyses of larval body color, analyses of larval
spotting pattern yielded fewer distinct QTL (four), but more QTL x QTL interactions
(three) (Table 1; Figures 4, 5C, 5D). Both spotting measures (spot number and spot area)
recovered two large-effect QTL on LG 2. These spotting QTL also overlapped with QTL
with small-to-modest effects on body color (Table 1; Figure 4). Co-localization of
spotting-pattern and body-color QTL suggests that the phenotypic correlations we
observed between body color and spotting pattern in F_2 males (Figure S2) are caused by
underlying genetic correlations via either pleiotropy or physical linkage.
These genetic mapping results provide us with our first glimpse into the genetic
architecture of naturally occurring color variation in pine sawflies. For both larval body
color and larval spotting pattern, we recovered a relatively small number of loci, some of
which had a substantial effect on the color phenotype (Table 1). Additionally, our
multiple-QTL models explained a considerable percentage of the observed variation in
larval color: up to 86% for larval body color (yellow) and up to 73% for larval spotting
pattern (spot number) (Table 2). Although limited statistical power precluded us from
quantifying the number of QTL of very small effect, our results suggest that the
proportion of variation explained by small, isolated QTL is small. It is possible, however,
that our large-effect QTL comprise many linked QTL of individually small effect (*e.g.,*
Stam and Laurie 1996; McGregor *et al.* 2007; Bickel *et al.* 2011; Linnen *et al.* 2013).
Distinguishing between oligogenic architectures (a small number of moderate to large-
effect loci) and polygenic architectures (a large number of small-effect loci) will require
fine-mapping and functionally validating genes and mutations for both color traits.
Nevertheless, for both types of color traits, our results clearly rule out both a monogenic
architecture and an architecture in which there are many unlinked, small-effect mutations.
The latter architecture is also highly unlikely if local color adaptation in pine sawflies
proceeded in the face of gene flow (Griswold 2006; Yeaman and Whitlock 2011), as
previous demographic analyses seem to indicate is the case (Bagley *et al.* 2017).
To date, QTL-mapping and candidate-gene studies of color traits have yielded
many examples of relatively simple genetic architectures in which one or a small number
of mutations have very large phenotypic effects (Rockman 2012; Martin and Orgogozo
2013). However, the relevance of these studies to broader patterns in phenotypic
evolution remains unclear due to biases that stem from a tendency to work on discrete
phenotypes and specific candidate genes. These biases can be minimized by focusing on
continuously varying color traits and employing an unbiased mapping approach as we
have done here (see also O'Quin *et al.* 2013; Albertson *et al.* 2014; Signor *et al.* 2016;
Yassin *et al.* 2016). Experimental biases aside, some have also argued that color itself—
but not color pattern—is atypically genetically simple (i.e., more likely to be mono- or
oligogenic than other traits) (Charlesworth *et al.* 1982; Rockman 2012). In our own data,

we see no obvious differences between the architectures of body color and spotting
pattern. If anything, the architecture of spotting appears slightly "simpler" (fewer QTL
with larger individual effects) than that of body color (Table 1). That said, more spotting
variation remains unexplained (Table 2), which could be attributable to a plethora of
undetected small-effect loci, and thus a more complex underlying architecture. A more
definitive test of the "color/pattern" hypothesis will therefore require fine-mapping our
QTL, with the prediction that the spotting QTL will "fractionate" to a greater extent (i.e.,
break into more QTL of individually smaller effect) than the body-color QTL. A rigorous
evaluation of the color/pattern hypothesis will also require data on the genetic
architecture of color and pattern traits that have evolved independently in other taxa. To
this end, ample variation in both body-color and color-pattern traits make *Neodiprion*
sawflies an ideal system for determining whether there are predictable differences in the
genetic architecture of different types of color traits.
484
Candidate genes for larval color traits
In total, we identified 61 candidate genes with known or suspected roles in
melanin-based or carotenoid-based pigmentation and for which we were able to identify
putative homologs in the *N. lecontei* genome (Table S7). Of these, 26 appeared to belong
to the *yellow* gene family. Notably, this number is equivalent to the number of *yellow*-
like/*yellow-MRJP*-like genes found in the genome of the jewel wasp, *Nasonia vitripennis,*
which boasts the highest reported number of *yellow*-like genes of any insect to date
(Werren *et al.* 2010). Additionally, 13 of these genes (*yellow-e, yellow-e3,* four *yellow-g,*
yellow-h, yellow-x, and five *MJRPs*) were found in tandem array along three adjacent
scaffolds (548, 170, and 36; ~1 Mb total) on LG 2. This genomic organization is
consistent with a conserved clustering of *yellow-h, -e3, -e, -g2,* and –*g* observed across
Apis, Tribolium, Bombyx, Drosophila, and *Nasonia* (Drapeau *et al.* 2006; Werren *et al.*
2010; Ferguson *et al.* 2011). Like *Nasonia* and *Apis*, this cluster also contains *MJRPs*;
like *Heliconius*, this cluster contains a *yellow-x* gene. Overall, we placed 57 of our 61
color genes on our LG-anchored assembly. With these data, we identified candidate
genes for all but two QTL (Yellow-1 on LG 1 and SpotNum-3 on LG 6) (Table S7).
501
Candidate genes for larval body color
Both of our largest-effect body-color QTL regions contained promising candidate
genes with known or suspected roles in carotenoid-based pigmentation (Table S7). First,
within the LG-3 QTL region (Yellow-4, PC1-1, PC2-3; Table 1, Figure 4), we found a
predicted protein-coding region in scaffold 518 with a high degree of similarity to the
Bombyx mori Cameo2 scavenger receptor protein (e-value: 1×10^{-18}; bitscore: 92.8).
Cameo2 encodes a transmembrane protein belonging to the CD36 family that has been
implicated in the selective transport of the carotenoid lutein from the hemolymph to
specific tissues (Sakudoh *et al.* 2010; Tsuchida and Sakudoh 2015). In the silkworm
Bombyx mori, Cameo2 is responsible for the "C mutant" phenotype, which is
characterized by a combination of yellow hemolymph and white cocoons that arises as a
consequence of disrupted transport of lutein from the hemolymph to the middle silk gland
(Sakudoh *et al.* 2010; Tsuchida and Sakudoh 2015).
Second, within the LG-5 QTL region (Yellow-5, Yellow-6, PC1-2, PC2-4; Table
1, Figure 4), we recovered a predicted protein coding gene in scaffold 164 with a high

degree of similarity to the ApoLTP-1 and ApoLTP-2 protein subunits (encoded by the gene *apoLTP-II/I*) of the *Bombyx mori* lipid transfer particle (LTP) lipoprotein (e-value: 0; bitscore: 391). LTP is one of two major lipoproteins present in insect hemolymph and appears to be involved in the transport of hydrophobic lipids (including carotenoids) from the gut to the other major lipoprotein, lipophorin, which then transports lipids to target tissues (Tsuchida *et al.* 1998; Palm *et al.* 2012; Yokoyama *et al.* 2013). Based on these observations, we hypothesize that loss-of-function mutations in both *Cameo2* and *apoLTP-II/I* contribute to the loss of yellow pigmentation in white-bodied *N. lecontei* larvae.

Candidate genes for larval spotting pattern

Our two largest-effect spotting-pattern QTL also yielded promising candidate genes—this time in the well-characterized melanin biosynthesis pathway (Table S7). In both LG-2 QTL regions, we found protein-coding genes that belong to the *yellow* gene family. The first LG-2 QTL region (SpotNum-1 and SpotArea-1; Table 1, Figure 4) contained 2 *yellow*-like genes that were most similar to *Apis yellow-x1* (e-value: 1.2×10^{-160}; bitscore: 471.47 and e-value: 3.9×10^{-151}; bitscore: 448.36). At present, there is little known about the function of *yellow-x* genes, which appear to be highly divergent from other *yellow* gene families (Ferguson *et al.* 2011). The second LG-2 QTL region (SpotNum-2 and SpotArea-2; Table 1, Figure 4) contained a cluster of 13 *yellow* genes, including *yellow-e*. In two different mutant strains of *B. mori*, mutations in *yellow-e* produced a truncated gene product that results in increased reddish-brown pigmentation in the head cuticle and anal plate compared to wildtype strains (Ito *et al.* 2010). In wildtype larvae, *yellow-e* is most highly expressed in the integument of the head and the tail (Ito *et al.* 2010). Based on these observations, one possible mechanism for the reduced spotting observed in the light-spotted MI population is an increase in *yellow-e* expression.

The second LG-2 spotting QTL region also contained a predicted protein that was highly similar to tyrosine hydroxylase (TH) (e-score: 6×10^{-123}, bitscore: 406). TH catalyzes the hydroxylation of tyrosine to 3,4-dihydroxyphenylalanine (DOPA), a precursor to melanin-based pigments (Wright 1987). Work in the swallowtail butterfly *Papilio xuthus* and the armyworm *Pseudaletia separata* demonstrates that TH and another enzyme, dopa decarboxylase (DDC), are expressed in larval epithelial cells containing black pigment (Futahashi and Fujiwara 2005; Ninomiya and Hayakawa 2007). Furthermore, inhibition of either enzyme prevented the formation of melanin-based larval pigmentation patterns (Futahashi and Fujiwara 2005). Thus, a reduction in the regional expression of *TH* (aka *pale*) is another plausible mechanism underlying reduced spotting in the light-spotted MI population.

We also found candidate genes in some of our QTL of more modest effect. As noted above, the Yellow-2 and Yellow-3 QTL overlap with the two large-effect spotting QTL (Table 1, Figures 4,5). One possible explanation for this observation is that in addition to impacting spotting phenotypes, genes in the melanin biosynthesis pathway also impact overall levels of melanin throughout the integument and therefore overall body color. Finally, the SpotNum-4 QTL contained a gene encoding dopamine N-acetyl transferase (*Dat*). Dopamine N-acetyl transferase, which catalyzes the reaction between dopamine and N-acetyl dopamine and ultimately leads to the production of a colorless

563 pigment, is responsible for a difference in pupal case color between two *Drosophila*
564 species (Ahmed-Braimah and Sweigart 2015).
565
Comparison with color loci in other taxa
Identification of candidate genes within our QTL peaks provides an opportunity
to compare our results with those from different insect taxa. For melanin-based traits, the
bulk of existing genetic studies involve *Drosophila* fruit flies (Massey and Wittkopp
2016), with a handful of additional studies in wild and domesticated lepidopterans (*e.g.,*
Ito *et al.* 2010; Hof *et al.* 2016; Nadeau *et al.* 2016). A recent synthesis of *Drosophila*
studies suggests that *cis*-regulatory changes in a handful of genes controlling melanin
biosynthesis (*yellow, ebony, tan, Dat*) and patterning (*bab1, bab2, omb, Dll,* and *wg*)
explain much of the intra- and interspecific variation in pigmentation (Massey and
Wittkopp 2016). Although our data cannot speak to the contribution of *cis*-regulatory
versus protein-coding changes, we do recover some of the same melanin biosynthesis
genes (*yellow* and *Dat*). By contrast, all of the *Drosophila* melanin patterning genes fell
outside of our spotting-pattern QTL intervals (Table S7). While we cannot rule these
genes out completely, our results suggest that none of them plays a major role in this
intraspecific comparison. Whether the same is true of patterning differences within and
between other *Neodiprion* species remains to be seen.
Compared to melanin-based pigmentation, much less is known about the genetic
underpinnings of variation in carotenoid-based pigmentation, which is widespread in
nature (Heath *et al.* 2013; Toews *et al.* 2017). Nevertheless, studies in the domesticated
silkworm (Sakudoh *et al.* 2010; Tsuchida and Sakudoh 2015), salmonid fish (Sundvold *et*
al. 2011), and scallops (Liu *et al.* 2015) collectively suggest that scavenger receptor
genes such as our candidate *Cameo2* may be a common and taxonomically widespread
source carotenoid-based color variation (Toews *et al.* 2017). Beyond evaluating patterns
of gene reuse, extensive intra- and interspecific variation in larval body color across the
genus *Neodiprion* (Figure 1) has the potential to provide novel insights into the molecular
mechanisms underlying carotenoid-based pigmentation.
592
Summary and Conclusions
Our study, which focuses on naturally occurring color variation in an under-
sampled life stage (larva), taxon (Hymenoptera), and pigment type (carotenoids),
represents a valuable addition to the invertebrate pigmentation literature. Our results also
have several implications for both hymenopteran evolution and color evolution. First, we
provide the first recombination-rate estimate for the Eusymphyta, the sister group to all
remaining Hymenoptera (*e.g.,* bees, wasps, and ants) (Peters *et al.* 2017). Our estimate
suggests that the exceptionally high recombination rates observed in some social
hymenopterans represents a derived state, supporting the hypothesis that increased
recombination is an adaptation for increasing genetic diversity in a colony setting (Gadau
et al. 2000; Schmid-Hempel 2000; Crozier and Fjerdingstad 2001; Wilfert *et al.* 2007).
Second, for both larval body color and spotting pattern, we can conclude that
intraspecific variation is due neither to a single Mendelian locus nor to a large number of
unlinked, small-effect loci. It remains to be seen, however, whether color, color pattern,
and other types of phenotypes differ predictably in their genetic architectures (Rockman
2012; Dittmar *et al.* 2016). Third, our QTL intervals contain several key players in

melanin-based and carotenoid-based pigmentation in other taxa, including *yellow, Dat,* and *Cameo2* (Massey and Wittkopp 2016; Toews *et al.* 2017). Although functional testing is needed to establish causal links, we can rule out a major role for several core melanin-patterning genes in this intraspecific comparison. Finally, although much work remains—including fine-mapping QTL, functional analysis of candidate genes, and genetic dissection of diverse color traits in additional *Neodiprion* populations and species—our rapid progression from phenotypic variation to strong candidate genes demonstrates the tremendous potential of this system for addressing fundamental questions about the genetic basis of color variation in natural populations.

ACKNOWLEDGEMENTS

For assistance with collecting and maintaining sawfly colonies, we thank past members of the Linnen laboratory, especially Robin Bagley, Adam Leonberger, Kim Vertacnik, Mary Collins, and Aubrey Mojesky. For assistance with phenotypic data collection, we thank Aubrey Mojesky, Mary Collins, and Ismaeel Siddiqqi. For advice on linkage mapping in R/qtl, we thank Karl Broman and Kelly O'Quin. For constructive comments on earlier versions of this manuscript, we thank Danielle Herrig, Emily Bendall, Kim Vertacnik, Katie Peichel, and two anonymous reviewers. Funding for this research was provided by the National Science Foundation (DEB-1257739; CRL), the United States Department of Agriculture National Institute of Food and Agriculture (2016-67014-2475; CRL), the Academy of Finland (project no. 257581; CL), and the University of Kentucky (Ribble summer undergraduate research fellowship, honors program independent research grant, and summer research and creativity fellowship to TS).

REFERENCES

Ahmed-Braimah Y. H., Sweigart A. L., 2015 A single gene causes an interspecific difference in pigmentation in *Drosophila*. Genetics 200: 331–342.

Albertson R. C., Powder K. E., Hu Y., Coyle K. P., Roberts R. B., *et al.*, 2014 Genetic basis of continuous variation in the levels and modular inheritance of pigmentation in cichlid fishes. Mol. Ecol. 23: 5135–5150.

Altschul S. F., Gish W., Miller W., Myers E. W., Lipman D. J., 1990 Basic Local Alignment Search Tool. J. Mol. Biol.: 403–410.

Andersson S., Prager M., 2006 Quantifying Colors. In: Hill GE, McGraw KJ (Eds.), *Bird Coloration Volume 1: Mechanisms and Measurements*, Harvard University Press, Cambridge, MA, pp. 41–89.

Arendt J., Reznick D., 2008 Convergence and parallelism reconsidered: what have we learned about the genetics of adaptation? Trends Ecol. Evol. 23: 26–32.

Bagley R. K., Sousa V. C., Niemiller M. L., Linnen C. R., 2017 History, geography and host use shape genomewide patterns of genetic variation in the redheaded pine sawfly (*Neodiprion lecontei*). Mol. Ecol. 26: 1022–1044.

Bendall E. E., Vertacnik K. L., Linnen C. R., 2017 Oviposition traits generate extrinsic postzygotic isolation between two pine sawfly species. BMC Evol. Biol. 17: 26.

Benjamin D. M., 1955 The biology and ecology of the red-headed pine sawfly. USDA Tech. Bull. 118: 55p.

Bickel R. D., Kopp A., Nuzhdin S. V, 2011 Composite effects of polymorphisms near
multiple regulatory elements create a major-effect QTL. PLoS Genet. 7: e1001275.
Broman K. W., Sen S., 2009 *A Guide to QTL Mapping with R/qtl*. Springer, New York.
Cain A. J., Sheppard P. M., 1954 Natural selection in *Cepaea*. Genetics 39: 89–116.
Catchen J., Hohenlohe P. A., Bassham S., Amores A., Cresko W. A., 2013 Stacks: An
analysis tool set for population genomics. Mol. Ecol. 22: 3124–3140.
Charlesworth B., Lande R., Slatkin M., 1982 A Neo-Darwinian commentary on
macroevolution. Evolution 36: 474–498.
Chen H., Rangasamy M., Tan S. Y., Wang H., Siegfried B. D., 2010 Evaluation of five
methods for total DNA extraction from western corn rootworm beetles. PLoS One 5.
Christin P. A., Weinreich D. M., Besnard G., 2010 Causes and evolutionary significance
of genetic convergence. Trends Genet. 26: 400–405.
Codella S. G., Raffa K. F., 1993 Defense strategies of folivorous sawflies. In: Wagner M,
Raffa KF (Eds.), *Sawfly Life History Adaptations to Woody Plants,* Academic Press
Inc., San Diego, pp. 261–294.
Codella S. G., Raffa K. F., 1995 Host plant influence on chemical defense in conifer
sawflies (Hymenoptera: Diprionidae). Oecologia 104: 1–11.
Conte G. L., Arnegard M. E., Peichel C. L., Schluter D., 2012 The probability of genetic
parallelism and convergence in natural populations. Proc. R. Soc. B 279: 5039–47.
Coppel H. C., Benjamin D. M., 1965 Bionomics of Nearctic pine-feeding diprionids.
Annu. Rev. Entomol. 10: 69–96.
Crozier R. H., Fjerdingstad E. J., 2001 Polyandry in social Hymenoptera: disunity in
diversity? Ann. Zool. Fennici 38: 267–285.
Danecek P., Auton A., Abecasis G., Albers C. A., Banks E., *et al.*, 2011 The variant call
format and VCFtools. Bioinformatics 27: 2156–2158.
Davey J. W., Hohenlohe P. A., Etter P. D., Boone J. Q., Catchen J. M., *et al.*, 2011
Genome-wide genetic marker discovery and genotyping using next-generation
sequencing. Nat. Rev. Genet. 12: 499–510.
Dittmar E. L., Oakley C. G., Conner J. K., Gould B., Schemske D., 2016 Factors
influencing the effect size distribution of adaptive substitutions. Proc. R. Soc. B 283:
20153065.
Drapeau M. D., Albert S., Kucharski R., Prusko C., Maleszka R., 2006 Evolution of the
Yellow / Major Royal Jelly Protein family and the emergence of social behavior in
honey bees. Genome Res.: 1385–1394.
Eisner T., Johnesse J. S., Carrel J., Hendry L. B., Meinwald J., 1974 Defensive use by an
insect of a plant resin. Science. 184: 996–999.
Elmer K. R., Meyer A., 2011 Adaptation in the age of ecological genomics: insights from
parallelism and convergence. Trends Ecol. Evol. 26: 298–306.
Elsik C. G., Tayal A., Diesh C. M., Unni D. R., Emery M. L., *et al.*, 2016 Hymenoptera
Genome Database: Integrating genome annotations in HymenopteraMine. Nucleic
Acids Res. 44: D793–D800.
Endler J. A., 1990 On the measurement and classification of colour in studies of animal
colour patterns. Biol. J. Linn. Soc. 41: 315–352.
Ferguson L. C., Green J., Surridge A., Jiggins C. D., 2011 Evolution of the insect yellow
gene family. Mol. Biol. Evol. 28: 257–272.
Fisher R., Ford E., 1947 The spread of a gene in natural conditions. Heredity. 1: 143–174.

Futahashi R., Fujiwara H., 2005 Melanin-synthesis enzymes coregulate stage-specific larval cuticular markings in the swallowtail butterfly, *Papilio xuthus*. Dev. Genes Evol. 215: 519–529.

Gadau J., Page R. E., Werren J. H., Schmid-Hempel P., 2000 Genome organization and social evolution in Hymenoptera. Naturwissenschaften 87: 87–9.

Gaj T., Gersbach C. A., Barbas C. F., 2013 ZFN, TALEN, and CRISPR/Cas-based methods for genome engineering. Trends Biotechnol. 31: 397–405.

Gerould J. H., 1921 Blue-green caterpillars: The origin and ecology of a mutuation in hemolymph color in *Colias (Eurymus) philodice*. J. Exp. Zool. Part A Ecol. Genet. Physiol. 34: 384–415.

Glastad K. M., Arsenault S. V., Vertacnik K. L., Geib S. M., Kay S., *et al.*, 2017 Variation in DNA methylation is not consistently reflected by sociality in Hymenoptera. Genome Biol. Evol. 9: 1687–1698.

Gompel N., Prud'homme B., 2009 The causes of repeated genetic evolution. Dev. Biol. 332: 36–47.

Goodwin S., Mcpherson J. D., Mccombie W. R., 2016 Coming of age : ten years of next-generation sequencing technologies. Nat. Rev. Genet. 17: 333–351.

Griswold C. K., 2006 Gene flow's effect on the genetic architecture of a local adaptation and its consequences for QTL analyses. Heredity. 96: 445–453.

Haldane J., 1948 The theory of a cline. J. Genet. 48: 277–284.

Harper K. E., Bagley R. K., Thompson K. L., Linnen C. R., 2016 Complementary sex determination, inbreeding depression and inbreeding avoidance in a gregarious sawfly. Heredity. 117: 326–335.

Heath J. J., Cipollini D. F., Stireman J. O., 2013 The role of carotenoids and their derivatives in mediating interactions between insects and their environment. Arthropod. Plant. Interact. 7: 1–20.

Heimpel G. E., Boer J. G. de, 2008 Sex determination in the Hymenoptera. Annu. Rev. Entomol. 53: 209–30.

Hoekstra H. E., Coyne J. A., 2007 The locus of evolution: Evo devo and the genetics of adaptation. Evolution. 61: 995–1016.

Hof A. E. van't, Campagne P., Rigden D. J., Yung C. J., Lingley J., *et al.*, 2016 The industrial melanism mutation in British peppered moths is a transposable element. Nature 534: 102–105.

Huang Y., Liu Z., Rong Y. S., 2016 Genome editing: from *Drosophila* to non-model insects and beyond. J. Genet. Genomics 43: 263–272.

Ito K., Katsuma S., Yamamoto K., Kadono-Okuda K., Mita K., *et al.*, 2010 Yellow-e determines the color pattern of larval head and tail spots of the silkworm B*ombyx mori*. J. Biol. Chem. 285: 5624–5629.

Kettlewell H. B. D., 1955 Selection experiments on industrial melanism in the Lepidoptera. Heredity. 9: 323–342.

Knerer G., Atwood C. E., 1972 Evolutionary trends in subsocial sawflies belonging to *Neodiprion abietis* complex (Hymenoptera: Tenthredinoidea). Am. Zool. 12: 407–418.

Knerer G., Atwood C. E., 1973 Diprionid sawflies: polymorphism and speciation. Science. 179: 1090–1099.

Kopp A., 2009 Metamodels and phylogenetic replication: a systematic approach to the

evolution of developmental pathways. Evolution. 63: 2771–2789.

Kraemer M. E., Coppel H. C., 1983 Hybridization of jack pine feeding sawflies
(Diproinidae: *Neodiprion*). Forestry Research Notes. University of Wisconsin,
Madison, WI.

Kronforst M. R., Barsh G. S., Kopp A., Mallet J., Monteiro A., *et al.*, 2012 Unraveling
the thread of nature's tapestry: The genetics of diversity and convergence in animal
pigmentation. Pigment Cell Melanoma Res. 25: 411–433.

Lee E., Helt G. A., Reese J. T., Munoz-Torres M. C., Childers C. P., *et al.*, 2013 Web
Apollo: a web-based genomic annotation editing platform. Genome Biol 14: R93.

Li H., Handsaker B., Wysoker A., Fennell T., Ruan J., *et al.*, 2009 The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.

Lindstedt C., Carita L., Mappes J., Johanna M., Päivinen J., *et al.*, 2006 Effects of group
size and pine defence chemicals on Diprionid sawfly survival against ant predation.
Oecologia 150: 519–26.

Lindstedt C., Lindstrom L., Mappes J., 2009 Thermoregulation constrains effective
warning signal expression. 63: 469–478.

Lindstedt C., Huttunen H., Kakko M., Mappes J., 2011 Disengtangling the evolution of
weak warning signals: High detection risk and low production costs of chemical
defences in gregarious pine sawfly larvae. Evol. Ecol. 25: 1029–1046.

Lindstedt C., Schroderus E., Lindström L., Mappes T., Mappes J., 2016 Evolutionary
constraints of warning signals: A genetic trade-off between the efficacy of larval and
adult warning coloration can maintain variation in signal expression. Evolution. 70:
2562–2572.

Linnen C. R., Farrell B. D., 2007 Mitonuclear discordance is caused by rampant
mitochondrial introgression in *Neodiprion* (Hymenoptera: Diprionidae) sawflies.
Evolution. 61: 1417–1438.

Linnen C. R., Farrell B. D., 2008a Phylogenetic analysis of nuclear and mitochondrial
genes reveals evolutionary relationships and mitochondrial introgression in the
sertifer species group of the genus *Neodiprion* (Hymenoptera: Diprionidae). Mol.
Phylogenet. Evol. 48: 240–257.

Linnen C. R., Farrell B. D., 2008b Comparison of methods for species-tree inference in
the sawfly genus *Neodiprion* (Hymenoptera: Diprionidae). Syst. Biol. 57: 876–90.

Linnen C. R., Poh Y.-P., Peterson B. K., Barrett R. D. H., Larson J. G., *et al.*, 2013
Adaptive evolution of multiple traits through multiple mutations at a single gene.
Science. 339: 1312–6.

Liu H., Zheng H., Zhang H., Deng L., Liu W., *et al.*, 2015 A de novo transcriptome of the
noble scallop, *Chlamys nobilis*, focusing on mining transcripts for carotenoid-based
coloration. BMC Genomics 16: 1–13.

Mackay T. F. C., Stone E. A., Ayroles J. F., 2009 The genetics of quantitative traits:
challenges and prospects. Nat. Rev. Genet. 10: 565–577.

Manceau M., Domingues V. S., Linnen C. R., Rosenblum E. B., Hoekstra H. E., 2010
Convergence in pigmentation at multiple levels: mutations, genes and function.
Philos. Trans. R. Soc. Lond. B. Biol. Sci. 365: 2439–2450.

Manichaikul A., Moon J. Y., Sen Ś., Yandell B. S., Broman K. W., 2009 A model
selection approach for the identification of quantitative trait loci in experimental
crosses, allowing epistasis. Genetics 181: 1077–1086.

Martin A., Orgogozo V., 2013 The loci of repeated evolution: a catalog of genetic
hotspots of phenotypic variation. Evolution. 67: 1235–50.
Martin A., Courtier-Orgogozo V., 2017 Morphological evolution repeatedly caused by
mutations in signaling ligand genes. In: *Diversity and Evolution of Butterfly Wing*
Patterns, Springer, Singapore, pp. 59–87.
Massey J. H., Wittkopp P. J., 2016 The genetic basis of pigmentation differences within
and between *Drosophila* species. In: *Current topics in developmental biology*,
Elsevier, pp. 27–61.
Maxwell D. E., 1958 Sawfly cytology with emphasis upon Diprionidae (Hymenoptera,
Symphyta). Proc 10th Int Congr Entomol. 2: 961–978.
McGraw K. J., Hudon J., Hill G. E., Parker R. S., 2005 A simple and inexpensive
chemical test for behavioral ecologists to determine the presence of carotenoid
pigments in animal tissues. Behav. Ecol. Sociobiol. 57: 391–397.
McGregor A. P., Orgogozo V., Delon I., Zanet J., Srinivasan D. G., *et al.*, 2007
Morphological evolution through multiple cis-regulatory mutations at a single gene.
Nature 448: 587–590.
Montgomerie R., 2006 Analyzing Colors. In: Hill GE, McGraw KJ (Eds.), *Bird*
Coloration Volume 1: Mechanisms and Measurement, Harvard University Press,
Cambridge, MA, pp. 90–147.
Montgomerie R., 2008 CLR, version 1.05.
Nadeau N. J., Jiggins C. D., 2010 A golden age for evolutionary genetics? Genomic
studies of adaptation in natural populations. Trends Genet. 26: 484–492.
Nadeau N. J., Pardo-Diaz C., Whibley A., Supple M. A., Saenko S. V., *et al.*, 2016 The
gene *cortex* controls mimicry and crypsis in butterflies and moths. Nature 534: 106–
110.
Ninomiya Y., Hayakawa Y., 2007 Insect cytokine, growth-blocking peptide, is a primary
regulator of melanin-synthesis enzymes in armyworm larval cuticle. FEBS J. 274:
1768–1777.
O'Quin C. T., Drilea A. C., Conte M. a, Kocher T. D., 2013 Mapping of pigmentation
QTL on an anchored genome assembly of the cichlid fish, *Metriaclima zebra*. BMC
Genomics 14: 287.
Orr H. A., Coyne J. A., 1992 The genetics of adaptation: a reassessment. Am. Nat. 140:
725–742.
Palm W., Sampaio J. L., Brankatschk M., Carvalho M., Mahmoud A., *et al.*, 2012
Lipoproteins in *Drosophila melanogaster*: assembly, function, and influence on
tissue lipid composition. PLoS Genet. 8.
Peters R. S., Krogmann L., Mayer C., Donath A., Gunkel S., *et al.*, 2017 Evolutionary
history of the Hymenoptera. Curr. Biol. 27: 1013–1018.
Peterson B. K., Weber J. N., Kay E. H., Fisher H. S., Hoekstra H. E., 2012 Double-digest
RADseq: an inexpensive method for de novo SNP discovery and genotyping in
model and non-model species. PLoS One 7: e37135.
Poelchau M., Childers C., Moore G., Tsavatapalli V., Evans J., *et al.*, 2014 The i5k
Workspace@NAL: enabling genomic data access, visualization and curation of
arthropod genomes. Nucleic Acids Res. 43: D714–D719.
Protas M. E., Patel N. H., 2008 Evolution of coloration patterns. Annu. Rev. Cell Dev.
Biol. 24: 425–446.

Remington D. L., 2015 Alleles versus mutations: Understanding the evolution of genetic
architecture requires a molecular perspective on allelic origins. Evolution. 69: 3025–
3038.
Rieseberg L. H., Archer M. a, Wayne R. K., 1999 Transgressive segregation, adaptation
and speciation. Heredity. 83: 363–372.
Rockman M. V., 2012 The QTN program and the alleles that matter for evolution: All
that's gold does not glitter. Evolution. 66: 1–17.
Rosenblum E. B., Parent C. E., Brandt E. E., 2014 The molecular basis of phenotypic
convergence. Annu. Rev. Ecol. Evol. Syst. 45: 203–226.
Ruxton G. D., Sherratt T. N., Speed M. P., 2004 *Avoiding Attack: The Evolutionary*
Ecology of Crypsis, Warning Signals and Mimicry. Oxford University Press, New
York.
Sakudoh T., Iizuka T., Narukawa J., Sezutsu H., Kobayashi I., *et al.*, 2010 A CD36-
related transmembrane protein is coordinated with an intracellular lipid-binding
protein in selective carotenoid transport for cocoon coloration. J. Biol. Chem. 285:
7739–7751.
Schmid-Hempel P., 2000 Mating, parasites and other trials of life in social insects.
Microbes Infect. 2: 515–520.
Signor S. A., Liu Y., Rebeiz M., Kopp A., 2016 Genetic convergence in the evolution of
male-specific color patterns in *Drosophila*. Curr. Biol. 26: 2423–2433.
Skinner M. E., Uzilov A. V, Stein L. D., Mungall C. J., Holmes I. H., 2009 JBrowse: a
next-generation genome browser. Genome Res 19: 1630–1638.
Smith S. G., 1941 A new form of spruce sawfly identified by means of its cytology and
parthenogenesis. Sci. Agric. 21: 245–305.
Sohi S. S., Ennis T. J., 1981 Chromosomal characterization of cell lines of *Neodiprion*
lecontei (Hymenoptera: Diprionidae). Proc. Entomol. Soc. Ontario 112: 45–48.
Stam L. F., Laurie C. C., 1996 Molecular dissection of a major gene effect on a
quantitative trait: the level of alcohol dehydrogenase expression in *Drosophila*
melanogaster. Evolution. 1564: 1559–1564.
Stern D. L., Orgogozo V., 2008 The loci of evolution: how predictable is genetic
evolution? Evolution. 62: 2155–2177.
Stern D. L., Orgogozo V., 2009 Is genetic evolution predictable? Science. 323: 746–751.
Stevens M., Párraga C. A., Cuthill I. C., Partridge J. C., Troscianko T. S., 2007 Using
digital photography to study animal coloration. Biol. J. Linn. Soc. 90: 211–237.
Streisfeld M. A., Rausher M. D., 2011 Population genetics, pleiotropy, and the
preferential fixation of mutations during adaptive evolution. Evolution. 65: 629–642.
Sugumaran M., Barek H., 2016 Critical analysis of the melanogenic pathway in insects
and higher animals. Int. J. Mol. Sci. 17: 1–24.
Sumner F. B., 1926 An analysis of geographic variation in mice of the *Peromyscus*
polionotus group from Florida and Alabama. J. Mammal. 7: 149–184.
Sundvold H., Helgeland H., Baranski M., Omholt S. W., Våge D. I., 2011
Characterisation of a novel paralog of scavenger receptor class B member I
(SCARB1) in Atlantic salmon (*Salmo salar*). BMC Genet. 12.
Tang H., Zhang X., Miao C., Zhang J., Ming R., *et al.*, 2015 ALLMAPS: robust scaffold
ordering based on multiple maps. Genome Biol. 16: 3.
Taylor J., Butler D., 2017 R Package ASMap: efficient genetic linkage map construction

and diagnosis. J. Stat. Softw. 79: 1–29.

R Core Team, 2013 R: A language and environment for statistical computing.

Toews D. P. L., Hofmeister N. R., Taylor S. A., 2017 The evolution and genetics of
carotenoid processing in animals. Trends Genet. 33: 171–182.

True J. R., 2003 Insect melanism: The molecules matter. Trends Ecol. Evol. 18: 640–647.

Tsuchida K., Arai M., Tanaka Y., Ishihara R., Ryan R. O., *et al.*, 1998 Lipid transfer
particle catalyzes transfer of carotenoids between lipophorins of *Bombyx mori*.
Insect Biochem. Mol. Biol. 28: 927–934.

Tsuchida K., Sakudoh T., 2015 Recent progress in molecular genetic studies on the
carotenoid transport system using cocoon-color mutants of the silkworm. Arch.
Biochem. Biophys. 572: 151–157.

Vertacnik K. L., Geib S., Linnen C. R., 2016 *Neodiprion lecontei* genome assembly
Nlec1.0. NCBI/GenBank. http://dx.doi.org/10.15482/USDA.ADC/1235565.

Werren J. H., Richards S., Desjardins C. A., Niehuis O., Gadau J., *et al.*, 2010 Functional
and evolutionary insights from the genomes of three parasitoid *Nasonia* species.
Science. 327: 343-348.

Wilfert L., Gadau J., Schmid-Hempel P., 2007 Variation in genomic recombination rates
among animal taxa and the case of social insects. Heredity. 98: 189–197.

Wilson L. F., Wilkinson R. C., Averill R. C., 1992 *Redheaded Pine Sawfly: Its Ecology*
and Management. USDA, Washington DC.

Wittkopp P. J., Carroll S. B., Kopp A., 2003 Evolution in black and white: Genetic
control of pigment patterns in *Drosophila*. Trends Genet. 19: 495–504.

Wittkopp P. J., Beldade P., 2009 Development and evolution of insect pigmentation:
Genetic mechanisms and the potential consequences of pleiotropy. Semin. Cell Dev.
Biol. 20: 65–71.

Wright T. R. F., 1987 The genetics of biogenic amine metabolism, sclerotization, and
melanization in *Drosophila melanogaster*. Adv. Genet. 24: 127–222.

Yassin A., Delaney E. K., Reddiex A. J., Seher T. D., Bastide H., *et al.*, 2016 The *pdm3*
locus is a hotspot for recurrent evolution of female-limited color dimorphism in
Drosophila. Curr. Biol. 26: 2412–2422.

Yeaman S., Whitlock M. C., 2011 The genetic architecture of adaptation under
migration-selection balance. Evolution. 65: 1897–1911.

Yokoyama H., Yokoyama T., Yuasa M., Fujimoto H., Sakudoh T., *et al.*, 2013 Lipid
transfer particle from the silkworm, *Bombyx mori*, is a novel member of the
apoB/large lipid transfer protein family. J. Lipid Res. 54: 2379–90.

922
923
Figure 1. Interspecific variation in *Neodiprion* larval color. Top row (left to right):
Neodiprion nigroscutum, *N. rugifrons*, *N. virginianus*. Middle row (left to right): *N.*
pinetum, *N. lecontei*, *N. merkeli*. Bottom row (left to right): *N. pratti, N. compar*, *N.*
swainei. Larvae in the first and last columns are exhibiting a defensive "U-bend" posture
(a resinous regurgitant is visible in *N. virginianus*, top right). *N. pratti* photo is by K.
Vertacnik, all others are by R. Bagley.
930
931

932
Figure 2. Intraspecific variation in *Neodiprion lecontei* larval color and cross design.
We crossed white, dark-spotted diploid females from Virginia to yellow, light-spotted
haploid males from Michigan. This produced haploid males with the VA genotype and
phenotype (not shown) and diploid females (F_1) with intermediate spotting and color.
Virgin F_1 females produced recombinant haploid males (F_2) with a wide range of body
color and spotting pattern (a representative sample is shown).
939

940
941
942
Figure 3. Larval color variation among *N. lecontei* populations and across
generations. For larval body color (A-C), higher scores for yellow (A) and lower scores
for PC1 (B) and PC2 (C) indicate higher levels of yellow pigment. For larval spotting
pattern (D-E), higher spot numbers (D) and spot area scores (E) indicate more melanic
spotting. For all traits, boxes represent interquartile ranges (median ± 2 s.d.), with outliers
indicated as points, and letters denote significant differences after correction for multiple
comparisons (see Table S3 for full statistical results).
950

951
952 **Figure 4.** *N. lecontei* **linkage map with larval-color QTL and 1.5-LOD support**
953 **intervals.**
954

956 **Figure 5. LOD profiles and interaction plots for larval color traits.** Interval mapping
957 analyses recovered QTL for larval body color (Yellow, PC1, PC2) on LGs 1, 2, 3, and 5
958 (A) and a single QTL x QTL interaction (B). QTL for larval spotting pattern (spot area
959 and spot number) were on LGs 2 and 6 (C), with three QTL x QTL interactions (D). QTL
960 names are as in Table 1.

Table 1. QTL locations and effect sizes for larval body color (yellow, PC1, and PC2) and larval spotting pattern (spot number and spot area).

Trait type	Trait	QTL name	LG[4]	Position (interval)[5]	Marker[6]	LOD	PVE	Effect size (SE)[7]	% diff.[8]
Body color	Yellow	Yellow-1	1	105.9 (84.4-112.5)	5744	8.29	1.32	-0.025 (0.0043)	8.06
Body color	Yellow	Yellow-2	2	28.2 (16.2-55.1)	19162	2.92	0.45	-0.014 (0.0043)	4.63
Body color	Yellow	Yellow-3	2	179.9 (140.8-194.0)	15279	3.51	0.54	-0.016 (0.0043)	5.13
Body color	Yellow	Yellow-4	3	181.9 (179.8-181.9)	1882	71.23	16.27	-0.084 (0.0043)	27.19
Body color	Yellow	Yellow-5	5	140.6 (139.9-149.6)	3222	71.17	16.25	-0.16 (0.0081)	51.64
Body color	Yellow	Yellow-6	5	149.6 (140.6-154.8)	19661	4.37	0.68	-0.035 (0.0082)	11.43
Body color	Yellow		Interaction	Yellow-3 x Yellow-5		12.45	2.03	0.061 (0.0086)	19.58
Body color	PC1	PC1-1	3	181.9 (160.4-181.9)	1882	4.31	3.56	1.15 (0.26)	15.51
Body color	PC1	PC1-2	5	140.6 (137.4-149.6)	3222	14.75	12.92	2.17 (0.26)	29.30
Body color	PC2	PC2-1	2	55.1 (46.8-63.9)	21	11.37	6.42	1.18 (0.16)	23.46
Body color	PC2	PC2-2	2	179.8 (165.2-189.9)	2143	7.24	3.99	0.91 (0.16)	17.99
Body color	PC2	PC2-3	3	181.9 (179.8-181.9)	1882	14.60	8.39	1.33 (0.16)	26.39
Body color	PC2	PC2-4	5	140.6 (139.9-149.6)	3222	33.13	21.13	2.09 (0.16)	41.45
Spotting pattern	Spot number	SpotNum-1	2	55.1 (48.0-57.6)	9282	58.46	23.52	6.31 (0.36)	33.06
Spotting pattern	Spot number	SpotNum-2	2	179.9 (179.8-187.5)	15279	78.19	35.45	7.81 (0.35)	40.96
Spotting pattern	Spot number	SpotNum-3	6	55.7 (51.1-64.9)	9040	4.84	1.43	-0.58 (0.59)	3.05
Spotting pattern	Spot number	SpotNum-4	6	64.9 (55.7-82.2)	7485	6.75	2.02	-1.00 (0.59)	5.26
Spotting pattern	Spot number		Interaction	SpotNum-1 x SpotNum-4		6.15	1.84	3.45 (0.71)	18.09
Spotting pattern	Spot number		Interaction	SpotNum-2 x SpotNum-3		4.64	1.38	-3.13 (0.71)	16.39
Spotting pattern	Spot area	SpotArea-1	2	55.1 (48.0-57.6)	9282	28.62	12.99	0.029 (0.0025)	34.86
Spotting pattern	Spot area	SpotArea-2	2	179.9 (169.1-187.5)	15279	63.73	35.47	0.048 (0.0024)	57.04
Spotting pattern	Spot area		Interaction	Spot-1 x Spot-2		1.76	0.69	0.014 (0.0049)	16.49

[4] Linkage group number

[5] Position in cM (1.5-LOD support intervals)

[6] Marker closest to QTL peak

[7] Effect sizes calculated as the difference in the phenotypic averages of among F_2 males carrying a VA allele and F_2 males carrying a MI allele (± standard error).

[8] Effect sizes calculated as a percentage of the difference between average trait values for the two grandparental lines (VA and MI).

Table 2. Summary of full multiple-QTL models for five larval-color traits.

Trait type	Trait	Covariates	# QTL	# Interactions	LOD	PVE
Body color	Yellow	None	6	1	182.03	85.83
Body color	PC1	Size, Mother	2	0	27.14	25.48
Body color	PC2	Mother	4	0	65.84	50.92
Spotting pattern	Spot number	Size, Mother	4	2	122.18	73.23
Spotting pattern	Spot area	Size, Mother	2	1	94.77	63.93