Title:

Does floral herbivory reduce pollination-mediated fitness in shelter rewarding Royal Irises?

Authors:

Mahua Ghara (1*), Christina Ewerhardy (2),Gil Yardeni (1, 3), Mor Matzliach (1), Yuval Sapir

(1)

(1) The Botanical Garden, School of Plant Science and Food Security, Tel Aviv University,

Tel Aviv 69978, Israel

(2) Albrecht von Haller Institute of Plant Sciences, University of Göttingen, DE-37073

Göttingen, Germany

(3) Current address: Department of Botany and Biodiversity Research, University of Vienna,

A-1030 Vienna, Austria

* Author for correspondence; mail: mahuaghara@gmail.com

Author Contributions: MG conceived and designed the experiments. YS supervised research.

MG, CE, GY and MM performed the experiments. MG and YS analyzed the data. YS wrote the

manuscript with contributions of all other authors.

Abstract:

Florivory, the damage to flowers by herbivores, can make flowers less attractive to pollinators.

Even when pollinated, flower consumed by florivores may fail to produce fruit or will produce

lower seed set. Despite the widespread evidence of florivory across ecosystems and plant taxa,

only a few studies tested experimentally the interaction of florivory and pollination by

manipulative study. We studied the effect of two levels of florivory on both pollinator visitation

and reproductive success over two flowering seasons in three *Iris* species. We hypothesized that

florivory will reduce pollen deposition due to reduced attractiveness to pollinators, and that

fruiting probability and seed set will depend on the extent of florivory. We performed artificial

florivory treatments, representing high, low, and no florivory (control) in two experiments. In the

first experiment, each of the three floral units of the same flower was subject to either low, high

or no artificial florivory, after which we counted the number of pollen grains present on the

stigma. In the second experiment, three flowers of the same plant were treated and were further

recorded for fruit and seed production. Surprisingly, and against our hypotheses, in all three

species no significant effect was found among and between florivory treatments and control,

neither in pollen grain deposited nor in fruit and seed set. The results undermine the assumption

that flower herbivory is necessarily antagonistic interaction and suggests that florivores are not

strong selection agents on floral reproductive biology in the *Oncocyclus* irises.

Key words: herbivory, *Iris* section *Oncocyclus*, floral herbivory, florivory, mutualist-antagonist

interaction, natural selection, pollen limitation.

Introduction

Flowers of animal-pollinated plants are the major mean of plants to advertise and attract pollinators. Floral traits serve as signals to the pollinators, usually fit to the best-efficient pollinator (Fenster et al., 2004). Flowers advertise through visual signals, fragrance cues, and even by acoustic signature, in order to stand out of the canopy or other flowering species (Schiestl and Johnson, 2013). Consequently, flower traits that increase attraction are selected by pollinators (Harder and Johnson, 2009). For example, floral size, contributing to the visibility of the flower, is positively selected by pollinators (Sletvold et al., 2010, Sletvold et al., 2016, Campbell et al., 1991, Conner and Rush, 1996, Harder and Johnson, 2009, Lavi and Sapir, 2015). However, flowers are costly organs that require investment in resources for production and maintenance. Other selection agents also act on floral traits, either in concert with, or in contrast to the selection exerted by pollinators (Strauss and Whittall, 2006). For example, floral size is under contrasting selection regimes, positive by pollinators and negative by drought and water loss (Galen, 2000, Carroll et al., 2001). Color polymorphism is also thought to be maintained by the combined effect of mutualists (pollinators) and antagonists (Carlson and Holsinger, 2010), or by opposing selection directions by herbivores and pathogens (Frey, 2004). Although many studies have examined floral adaptations to pollinators, the role of non-pollinator selection agents in shaping floral evolution and plant reproductive success is still underappreciated.

Florivory, the damage of flowers by herbivores, is widespread across plant taxa and ecosystems (Gonzáles et al., 2016, Burgess, 1991). Florivores comprises of various taxonomic orders of animals that consume the entire flower (or the buds) or floral parts (bracts, sepals, petals, nectaries, stamens, pistils or pollen). Florivory may affect plant fitness directly or indirectly, by reducing fruiting or seed-set, and can consequently affect population dynamics (Louda and Potvin, 1995). Florivory reduces fitness directly when reproductive parts are consumed, whereas indirect effect can be through effects on pollinator behavior and hence reduce pollination services. An increasing number of studies suggest that florivory can decrease pollinator visitation rate and pollination success (reviewed in Gonzáles et al., 2016, but see Zhu et al., 2017). Methods for studying the effect of florivory on pollination primarily focus on the effect of florivore presence or the extant of florivory on pollinator visits, using natural encounters and manipulative studies. For example, Kirk et al.(1995) placed black spots on flowers to simulate presence of florivore beetle and found that bees were likely to avoid flowers

with mimicry of florivore presence. Although artificial florivory performed in the field can

provide an estimate to the direct effect of florivory on pollination, few studies implement this

method. Moreover, while behavior of pollinators and/or measure of maternal fitness (measured

as the number of visits or seed set, respectively) are often used to estimate the effect of florivory

on pollination (e.g., Cardel and Koptur, 2010), direct evidence for pollination success, which is

pollen deposition on the stigma, has rarely been incorporated in florivory studies. Although

pollinator limitation and pollen limitation supply the ground for pollinator-mediated selection

(Bartkowska and Johnston, 2015, Lavi and Sapir, 2015, Sletvold et al., 2017), it is unclear

whether it affects florivore-mediated selection. Therefore, controlled manipulative experiments

that measure pollen deposition and fitness, as well as incorporating pollen limitation, are needed

to estimate directly the interaction of florivory and pollination success.

In a survey of the literature, we found only a small number of studies that used a controlled

experimental florivory that mimic florivory by artificially manipulating or removing parts of the

petals and tested for effect on pollination. Of these, seven studies found negative effect on

pollination success , one study found mixed effects that depended on flower morph (Carper et al.,

2016), and one study found no effect (Tsuji et al., 2016). Thus, it remains unclear whether

florivory affects plant fitness indirectly through inducing negative effect on pollinators. As an

example, McCall (2008) found that both natural and artificial petal damage indeed reduced

fitness, and while it deterred pollinator activity, the effect was a result of petal physical damage

rather than reduced pollinators activity, as pollen addition experiment did not recover fruit-set.

Hence, for fully understanding the multiple effects of florivory on fitness and the interplay

between direct and indirect effects it requires further experiments in diverse plant species.

We studied the effect of florivory on pollination success in three species of the Royal Irises

(*Iris* section *Oncocyclus*) in their natural habitats in Israel (Figure 1 A–C). The Royal Irises

comprise of about 30 species across the Middle East, most are narrow endemic (Rix, 1997,

Mathew, 1989). This group was well studied as a system for pollination by shelter reward to

male *Eucera* bees (Figure 1 D; Sapir et al., 2005, Sapir et al., 2006, Watts et al., 2013, Monty et

al., 2006, Vereecken et al., 2013, Lavi and Sapir, 2015). Floral organs in irises are organized in

three units consisting of a tunnel, in which the male *Eucera* bees seek shelter for the night or at

low temperatures. A stigma and anther are located within each tunnel and pollen is deposited on

stigmas by bees that move among flowers as they seek shelter (Sapir et al., 2005).

The Royal Irises are eaten by various florivores, from snails and true bugs to grasshoppers,

birds and goats, and the intensity of damage ranges from a few superficial scratches or poke

marks, up to >90% of damage to floral tissue (M. Ghara and Y. Sapir, per. observations; Figure 1

E–G). From our observations, it appears that all flower parts are potentially eaten, either bracts,

petals, the petaloid style, anthers or ovaries. While damage to the reproductive organs may

obviously reduce fitness directly, we asked whether florivory affects fitness indirectly through

reduced attraction to pollinators. In order to estimate the effect of florivory on pollinators'

visitation and pollination success we manipulated the flowers to simulate two levels of florivory,

i.e., high (more than 50% damage) and low (10-30% damage), and compared to control flowers

without damage. We asked the following questions: (1) Does florivory affect pollinator

visitation? (2) Does florivory affect fruiting probability and seed set? We used pollen deposition

on stigma as a surrogate for estimating pollinator visitation and used fruit- and seed-set to

estimate overall effect of florivory on (maternal) fitness. Accounting for two measures of

pollination success simultaneously is likely to reveal an indication of the possible effect of

florivory on pollination-mediated fitness in the royal irises.

Methods

Study species and sites

We conducted the florivory experiment in two consecutive years on three species of the Royal

Irises, namely *Iris atropurpurea* Baker, *I. petrana* Dinsm., and *I. lortetii* Barbey ex Boiss.

Experiments were conducted in the natural environment at the largest population for each of the

species. *Iris atropurpurea* was studied in Netanya Iris Reserve (32.28°N, 34.84°E, alt. 35 m),

located on stable coastal sand dunes in Mediterranean climate and consisting of mostly low shrub

vegetation. Population size is estimated >1,000 plants (Yardeni et al., 2016). Flowering season,

and hence experiment time, is earlier compared to other species of the Royal Irises, starting as

early as mid-January, and peaks in February. Experiments in *I. atropurpurea* in Netanya were

conducted between February 12[th] and March 2[nd], 2016, and between February 19[th] and March

7[th], 2017. *Iris petrana* was studied in Yeruham Iris Reserve (31.02°N, 34.97°E, alt. 560 m), a

large population (estimated >10,000 plants) growing on sandy loess hills over Neogene

sandstone in arid climate. Vegetation is sparse desert shrubs, mostly *Retama raetam* (Forssk.)

Webb and *Anabasis articulata* (Forssk.) Moq. Flowering season is in March, and experiments in

I. petrana were conducted between March 5[th] and 14[th], 2016, and between March 19[th] and April

2^{nd}, 2017. The shift in dates in 2017 resulted due to about two weeks delay in flowering period in

2017. *Iris lortetii* was studied in two sub-populations near Malkiya in the upper Galilee (central

coordinates: 33.09°N, 35.52°E, alt. 620 m). Populations of *I. lortetii* are sparse and relatively

small, thus, two sub-populations at a distance of 3 km of each other were pooled to achieve a

sufficient sample size. Plants are growing on Eocene limestone in mesic Mediterranean climate

and vegetation is open woodland dominated by *Quercus calliprinos* Webb and *Pistacia atlantica*

Desf. trees, accompanied by dense herbaceous vegetation. *Iris lortetii* is the late blooming

species among the Israeli species of the Royal Irises; experiments were conducted between

March 29^{th} and April 7^{th}, 2016. The experiments were not conducted on *Iris lortetii* in 2017

because of high herbivory in 2016 and therefore a potential decrease in sample size.

Flowers of the Royal Irises grow singly on a stem, but the plant (genet) comprises of one to

hundreds of stems (ramets) in a well-defined patch. Plants for the experiments were randomly

selected in a dense part of the population in Netanya, or along transects in Yeruham. In Malkiya

the plants are sparse and plants in all genets located were used. The three experiments described

below were conducted simultaneously in time with only a single experiment conducted in each

genet to avoid the joint effect of several treatments.

Pollinator and pollen limitation

To assess the extent of pollinator and pollen limitations we took the approach of Lavi and

Sapir (2015). In Netanya (*I. atropurpurea*) we supplemented 54 flowers in 2016 and 30 flowers

in 2017 with additional pollen. In Yeruham (*I. petrana*) we supplemented additional pollen for

22 flowers in 2016 and 45 in 2017. Pollinator and pollen limitation were not quantified in *I.*

lortetii.

Flowers were randomly chosen and were pollinated using a mixture of pollen collected from

>20 flowers located at least 10 meters away from the supplemented flowers. The flowers were

left open for natural pollination. At the end of the season, the fruits of all flowers were collected

and seeds were counted after full ripening. Fraction of fruits produced and number of seeds were

compared with flowers of the control treatment in the experiment testing for florivory effect on

maternal fitness (see below).

Florivory manipulations – effect on pollination

Flowers of the Royal Irises comprise of three identical units, each bearing one upright and one

downright petals (standard and fall, respectively), and one petaloid style, curved above the fall to

create a tunnel where the reproductive organs reside (Figure 1). The three stigmas, located at the

top of the entrance of each of the pollination tunnels, are merged in the base of the flower to one

style. A previous study showed that pollinating one style is sufficient to produce seeds in all

three carpels in the ovary (Watts et al., 2013). To study the effect of florivory on pollen

deposition, a single flower in a genet was selected and each flower unit (i.e., pollination tunnel)

was given one of three treatments as follows. High damage – both lower petal and its opposite

upper petal were manually damaged up to 50% or more of the petal's area, using a puncher of 6-

8 mm in diameter. Low – 10-30% of the petals' area were removed using puncher. Control –

petals were not damaged and left un-manipulated (Figure 2). To control for the possible effect of

the contact between the metal hole-puncher and the flower the puncher was rubbed on the petal

surface in the control treatments. In addition, because damaging the petals required holding a

layer of tissue paper against the puncher, we also gently rubbed tissue paper under the surface of

the petal in the control treatment.

Flowers were treated in the first day after anthesis and left open for two consecutive evenings

following treatments to enable pollinators to visit naturally. In the morning of the third day, the

stigmas of the three pollination units were collected in separate vials containing 1 ml of 70%

ethanol. To prevent naturally occurring florivores from damaging the flowers, the stem of the

treated flower was coated with a layer of double sided sticky paper tape, as well as a layer of

Petroleum jelly (Vaseline). Occasionally we found insects trapped on the Vaseline layer, and in

some rare cases, we found florivores that passed this barrier. Flowers found to be damaged

naturally (mostly by flying insects, snails, or mammals) were discarded in order to account for

the effect of controlled artificial florivory only.

Collected stigmas were brought to the laboratory for pollen counting and kept in room

temperature. Pollen grains were stained using a drop of basic fucshin (Calberla's stain). Stigma

was then dissected in a drop 70% aqueous glycerol (Dafni et al., 2005), mounted on microscope

glass slide and the pollen present on the stigma was counted under dissecting microscope (WILD

Heerbrugg Switzerland M5-72558).

Florivory manipulations – effect on maternal fitness

To study the effect of florivory on seed set, three buds of the same genet, roughly of the same

developmental stage (i.e., before emerging from bracts) were selected and bagged to avoid bud

florivory. Upon anthesis, each of the three flowers was randomly assigned to one of the florivory

treatments described above, i.e., high, low, or control. To control for the effect of visual

attraction mediated by flower size (Lavi and Sapir, 2015) we measured flower length as a

surrogate for display size of each flower before manipulation. Flower length was measured from

the bottom of the lower petal to the top end of the upper one. The flowers were left open to

enable naturally occurring pollination. At the end of the season, approximately three weeks after

the end of the flowering in each site, the fruits were collected and brought to the laboratory. In

2016, fruits of *I. atropurpurea, I. petrana* and *I. lortetii* were collected on March 21st, April 2nd

and April 27th, respectively. In 2017, fruits were collected on March 12th and 22nd for *I.*

atropurpurea and on April 13th and 27th for *I. petrana*. Fruits were kept in paper bags in room

temperature until seed maturation. Fitness was recorded as presence or absence of a fruit

(binomial data), and as the number of viable seeds (count data).

Data analyses

The data was analyzed in R (R Development Core Team, 2014) using R-studio interface. To

test for differences between control and supplemental pollination treatments, we used

generalized linear models (GLMs) with year and treatment effects nested within site. For fruit-

set, we used GLM with binomial distribution errors, and for number of seeds we used quasi-

Poisson distribution errors due to over-dispersion of the data. Pollen and pollinator limitation

indices were calculated as *PLI=1–O/S*, where *O* is fruit-set or seed-set in open flowers, and *S* is

fruit-set or seed-set in supplementary pollination treatment. Number of pollen grains deposited

on stigmas was analyzed using GLM with year and treatment effects nested within species (note

that in *I. lortetii* experiment was done only in 2016). As in number of seeds, we used quasi-

Poisson distribution errors due to over-dispersion of the data. Because the three treated floral

units were on the same flower, for analysis of pollen deposition these were considered as non-

independent. Thus, we initially tested models that included also the effect of flower ("block"

effect). We found this effect to be non-significant (not shown), hence it was removed from the

analysis. For analyses using fruit or seeds as explained variables, we incorporated flower size as

a covariate. For seeds as a response variable, we used only the subset of flowers that set a fruit.

In order to account for non-normal distribution, we used GLM with binomial distribution errors

for fruits and GLM with quasi-Poisson distribution errors for seeds.

Results

Pollinator and pollen limitation

We found a significant effect of site and a significant effect of treatment, nested within year

and within site, on both fruit-set and seed-set (Table 1). The effect of year was not significant. In

Netanya, flowers supplemented with pollen produced significantly higher fruit-set than control

flowers, open to natural pollinators only (48.1% versus 5.5% in 2016 and 60% versus 15% in

2017; P<0.001 in both years; Figure 3 A&B). Seed-set in Netanya in the two years was also

significantly higher in flowers supplemented with excess pollen, compared to open flowers

pollinated only naturally (mean number of seeds 11.5 versus 1.1 in 2016 and 10.3 versus 1.7 in

2017, P<0.005 in both years; Figure 3 C&D). Both pollinator limitation and pollen limitation

were extremely high in Netanya in both years (PLI=0.88 for fruit-set and PLI=0.91 for seed-set

in 2016; PLI=0.75 for fruit-set and PLI=0.84 for seed-set in 2017).

In Yeruham, fruit-set was similar between supplementary and open flowers in 2016 (40.9%

versus 41.7%, respectively, P=0.96; Figure 3A). In 2017, fruit-set was significantly higher in

flowers supplemented with pollen, compared to not-treated control flowers (73.3% versus

34.8%, respectively, P<0.001; Figure 3 B). Likewise, seed-set was not significantly different

between supplementary and open flowers in 2016 (mean number of seeds 7.1 versus 9.4,

respectively, P=0.675) but differ between treatments in 2017 (mean number of seeds 15.6 versus

5.8, respectively, P=0.003; Figure 3 C&D). Pollinator limitation and pollen limitation were

below zero in Yeruham in 2016 (PLI= −0.02 for fruit-set and PLI= −0.32 for seed-set), but

relatively high in 2017 (PLI= 0.53 for fruit-set and PLI= 0.64 for seed-set).

Pollen deposition on stigma

We found significant effect of year and species on number of pollen grains deposited on

stigmas of flowers ($F_{2,691}$=4.38, P=0.013 for species and $F_{2,691}$=24.96, P<0.001 for year effect,

nested within species). Nonetheless, treatment effect (nested within year and within species) was

not significant ($F_{2,691}$=0.63, P=0.706). In *Iris atropurpurea*, 49 (26.3%) stigmas did not receive

pollen at all in 2016 and 8 (5.3%) in 2017. Flowering units with high artificial florivory revealed

lower number of pollen grains on stigma in 2016 and lower in 2017(Figure 4), but these

differences were small and far from significance ($F_{2,183}$=0.67, P=0.515 in 2016 and $F_{2,135}$=0.16,

P=0.855 in 2017). In *I. petrana*, all stigmas received pollen grains in 2016, but in 2017, 51 (34%)

did not receive any pollen grain. Number of pollen grains deposited on the stigmas was an order

of magnitude larger than in *I. atropurpurea* in 2016 (Figure 4 A), but similar in 2017 (Figure 4

B). Nonetheless, as in *I. atropurpurea*, no significant treatment effect was found in both years

($F_{2,195}=0.53$, P=0.588 in 2016 and $F_{2,144}=1.89$, P=0.155 in 2017). In *Iris lortetii*, as in *I. petrana*

in 2016, all stigmas received pollen and stigmas in units treated by high artificial florivory

received pollen grains in a similar level as the control, untreated units, both higher than medium

artificial florivory treatment (Figure 4 A). However, this difference was not significant as well

($F_{2,30}=0.322$, P=0.727).

Fruit and seed sets

In *Iris atropurpurea*, 80 flowers were included in the final analyses in 2016, because 16 of the

treated flowers were either not found or damaged. Of these, only 13 flowers set fruits, indicating

extreme pollinator limitation and lack of pollinator visitations. Treatment did not affect either

fruit-set or number of seeds ($F_{2,75}=0.69$, P=0.501, and $F_{2,8}=0.26$, P=0.774, respectively; Figure 5

A & C). In both analyses flower size did not affect significantly (P=0.266 and P=0.554 for fruits

and seeds, respectively). In 2017, 124 flowers were included in the experiment, of which four

were damaged. Of the remaining 120 flowers, 23flowers set fruits. As in 2016, no effect of the

treatment was found, neither on fruit-set, nor on number of seeds ($F_{2,116}=0.37$, P=0.695, and

$F_{2,19}=0.95$, P=0.404, respectively; Figure 5 B & D). Similar to 2016, flower size as covariate did

not affect fruit-set or seed-set (P=0.188 and P=0.857, respectively).

In *I. petrana* population in Yeruham, 132 flowers were treated in 2016, but 29 flowers of all

treatments were eaten by goats that entered the reserve illegally and ate wilting flowers and

young fruits in the pre-dispersal stage. Of the remaining 103 treated flowers, 31 flowers set

fruits. Treatment effect on fruit-set was not significant ($F_{2,98}=1.89$, P=0.156 ;Figure 5 A).

Interestingly, flower length (before treatment) significantly affected fruit set ($F_{1,98}=5.09$,

P=0.026), but with no significant interaction with treatment ($F_{2,96}=0.26$, P=0.770). Although

control flowers produced almost twice fraction of fruits compared to florivory treatment (34%

vs. 18%), this difference was not significant when controlled for flower size. Number of seeds

was not affected by treatment ($F_{2,98}=1.89$, P=0.156, n=31; Figure 5C). Flower size did not affect

seed-set ($F_{1,26}=1.23$, P=0.278). In 2017, only six flowers were eaten or not found, out of 188

flowers treated. As in 2016, no effect of the treatment was found, neither on fruit-set, nor on

number of seeds ($F_{2,133}=1.64$, P=0.199, and $F_{2,29}=0.72$, P=0.495, respectively; Figure 5 B & D).

As opposed to 2016, flower size as covariate did not affect fruit-set but did affect seed-set

(P=0.472 and P<0.001, respectively). No interaction was found between flower size and

florivory treatment in their effect on seed-set (P=0.463).

In the two sites of *Iris lortetii*, 61 out of 62 flowers treated (12 in Avivim and 50 in Malkiya)

were found at the end of the season and included in the analyses. No significant difference was

found among treatments ($F_{2,57}$=0.21, P=0.811; Figure 4 A). Flower size affected fruit-set

($F_{1,57}$=4.42, P=0.40). Number of seeds was not significantly affected by treatment ($F_{2,12}$=0.37,

P=0.698; Figure 4 C), and neither by flower size ($F_{1,12}$=0.92, P=0.356).

Discussion

Florivory, namely the damage herbivores cause to floral organs, can affect fitness either

directly by consuming pollen or ovules or physiological costs, or indirectly, by reducing plant

attraction signal for the pollinators (Burgess, 1991, McCall and Irwin, 2006). Here we tested for

both direct and indirect effects of florivory on fitness by executing artificial florivory and

measuring both fitness and pollination. Our results do not support the hypothesis that florivory

affects pollination success in the Royal Irises either directly or indirectly. Instead, we show that

artificial damage to reproductive tissues in three species of the royal irises did not significantly

reduce either pollen deposition or fitness of manipulated flowers, compared to un-treated ones

(Figure 4 and Figure 5).

While numerous studies were concerned with the effect of florivory on plant fitness,

controlled, artificial florivory was rarely performed. Most studies examined flowers that were

naturally attacked by florivores (e.g., Meindl et al., 2013, Ruane et al., 2014, Eliyahu et al.,

2015) or used experimental florivore removal or prevention (e.g., Krupnick et al., 1999, Theis

and Adler, 2012, Althoff et al., 2013). Only a few studies implemented methods similar to ours,

using cutting flowers to simulate florivory. These studies revealed mixed results. For example,

Sõber et al. (2010) experimentally showed a correlation between extent of florivory and

pollinator visitations at both population and plant level. On the other hand, Tsuji et al. (2016)

found no evidence for pollinators discrimination against experimentally damaged flowers.

Interestingly, mixed results can be found within the same system: Carper et al.(2016) found

differences between heterostylous morphs in pollinator responses to artificial damage, and found

no effect on fitness. Our study adds to the puzzle by providing yet another piece of evidence that

florivory itself does not deter pollinators, nor reduces fitness.

Negative effect of florivory on pollination may act in two avenues. One possible effect of

florivory on pollinator behavior is the deterring of pollinators from eaten flowers. This is

achieved by either avoidance flowers where florivore is visually detected (Kirk et al., 2005) or

by a change in volatile compounds (Kessler et al., 2013). Another possible effect of florivory on

pollination is mediated by the effect of the overall advertisement size of the flower and reducing

visual signaling for the pollinators (Sánchez-Lafuente, 2007). This may reduce number of visits

and lead to pollinator limitation or pollen limitation, which in turn reduces fruit-set and seed-set,

respectively (Sapir et al., 2015). This study was conducted in natural populations of which at

least one (Netanya) indeed experience strong pollinator limitation. It is likely that the effect of

pollen limitation in this population obscures the effect of florivory (McCall, 2010). Thus, we

propose that selection mediated by pollinator/pollen limitation is stronger than selection pressure

exerted by florivory. A previous study that tested for pollen limitation in two *Iris* species

showed, indeed, that pollinators limitation provides conditions for pollinator-mediated selection

(Lavi and Sapir, 2015). However, in Yeruham we found no evidence for pollen limitation, still

no effect of florivory was found, which contradicts this hypothesis. Our mixed results suggest

that while pollinators may be a selection agent on flower traits (as in Lavi & Sapir, 2015),

florivores do not act as selection agents because pollen limitation balances their effect (cf. Jogesh

et al., 2016), but this connection cannot be generalized beyond our specific system.

Pollinators are thought to be the major selection agent on floral traits through their positive

effect on fitness; this, however, was challenged by observations on contrasting effects of abiotic

conditions or antagonistic biotic interactions (Herrera, 1996, reviewed in Strauss and Whittall,

2006). Strauss and Whittall (2006) proposed two scenarios of such mutualistic-antagonistic

effect, in which overall selection acts as either directional or stabilizing on floral traits. Based on

our results, we propose a third scenario, where florivory affects fitness at the same level as the

mean effect of pollinators and regardless floral trait. In this case, pollinator-mediated selection

will govern trait evolution, but the presence of florivory reduces or diminishes effect size (Figure

6). We speculate that florivory may indeed not affect the direction of selection but the intensity

of it. Thus, we suggest that testing for the net selection mediated by pollinators should control

experimentally for the effect of florivores. In a follow-up experiment, we intend to test for net

pollinator-mediated selection on floral color in the Royal Irises. Floral pigment concentration is

expected to deter florivores and attract pollinators (de Jager and Ellis, 2014, McCall et al., 2013),

and given the results of the current study we assume that florivory may not necessarily reduce

fitness; instead, it is expected that a weak directional pollinator-mediated selection on floral color

will be detected after controlling for florivory.

While few previous studies have presented only slight (or no) effect of florivory on

pollination, it may be further hypothesized that the Royal Irises are unique and thus do not

represent a general rule relevant to other species. The Royal Irises present a unique pollination

syndrome in which pollination is performed by night sheltering male bees (Sapir et al., 2005,

Watts et al., 2013). Previous studies suggest that flower display itself is not necessarily the major

attractant for these pollinators (Lavi and Sapir, 2015). While our florivory-like manipulations

were performed on petals, it is likely that these manipulations did not affect pollinator choice

because the shelter itself (the pollination tunnel) was not damaged. Because we have observed

florivory of stigma and anthers (M. Ghara pers. observations), we suspect that florivory may

affect male and female fitness through consumption of reproductive organs, but we have yet to

assess the cost of florivory on reproductive organs. Nonetheless, while natural florivory is

widespread in all species and most populations of the Royal Irises in Israel, the estimated

proportion of flowers of which pollination tunnels were eaten is rather small (M. Ghara,

manuscript in preparation). Thus, our manipulation on petals accurately mimics natural florivory

and our conclusions on the role of florivory in selection on flowers of the Royal Irises are valid.

Finally, our study provides an application to conservation. Of the three species studied, two

(*Iris atropurpurea* and *I. lortetii*) are rare and endangered species (Sapir, 2016a, Sapir, 2016b).

Understanding the relative contribution of biotic interactions to population dynamics may shed

light on the factors affecting species survival in a way that may contribute for evidence-based

management. The study presented here suggests that reduced mutualistic interactions, namely,

pollination services, rather than antagonistic florivory, threatens the maintenance of positive

population growth in the Royal Irises.

Acknowledgement

We thank many students, volunteers and friends for assisting in field work in various

capacities, especially Aakash Jog, Anil MG, Danya Ayehlet Cooper, Gali Paltin Ben Adiva,

Gavriella Shani, Hagai Shemesh, Ilana Sichel, Ilan Etam, Ilya Kutuzov, Kineret Manevich,

Maithili Jog, Maeva Relet, Mor Marzan, Yuval Shtrool, Valery Zayt, Yael Avidan, Yonatan

Ben-Ami and Zion Awaskar. Prof. D. Eisikowitch provided lab support and fruitful discussions

and ideas. The study was partially supported by the Israeli Nature-Parks Authority and a grant

from Israel Science Foundation #336/16 to YS. MG was supported by Israel Planning and

Budgeting Commission (PBC) post-doctoral fellowship.

References

Althoff DM, Xiao W, Sumoski S, Segraves KA. 2013. Florivore impacts on plant reproductive
success and pollinator mortality in an obligate pollination mutualism. *Oecologia,* **173**:
1345-1354.

Bartkowska MP, Johnston MO. 2015. Pollen limitation and its influence on natural selection
through seed set. *Journal of Evolutionary Biology,* **28**: 2097-2105.

Botto-Mahan C, Ramírez PA, Ossa CG, Medel R, Ojeda-Camacho M, González AV. 2011.
Floral herbivory affects female reproductive success and pollinator visitation in the
perennial herb *Alstroemeria ligtu* (Alstroemeriaceae). *International Journal of Plant*
Sciences, **172**: 1130-1136.

Burgess K. 1991. *Florivory: The ecology of flower feeding insects and their host plants,* PhD,
Harvard University.

Campbell DR, Waser NM, Price MV, Lynch EA, Mitchell RJ. 1991. Components of
phenotypic selection: pollen export and flower corolla width in *Ipomopsis aggregata*.
Evolution, **45**: 1458-1467.

Cardel YJ, Koptur S. 2010. Effects of Florivory on the Pollination of Flowers: An
Experimental Field Study with a Perennial Plant. *International Journal of Plant Sciences,*
171: 283-292.

Carlson JE, Holsinger KE. 2010. Natural selection on inflorescence color polymorphisms in
wild Protea populations: The role of pollinators, seed predators, and intertrait
correlations. *American Journal of Botany,* **97**: 934-944.

Carper AL, Adler LS, Irwin RE. 2016. Effects of florivory on plant-pollinator interactions:
Implications for male and female components of plant reproduction. *American Journal of*
Botany, **103**: 1061-1070.

Carroll AB, Pallardy SG, Galen C. 2001. Drought stress, plant water status, and floral trait
expression in fireweed, *Epilobium angustifolium* (Onagraceae). *American Journal of*
Botany, **88**: 438-446.

Conner JK, Rush S. 1996. Effects of flower size and number on pollinator visitation to wild
radish, *Raphanus raphanistrum. Oecologia,* **105**: 509-516.

Dafni A, Kevan PG, Husband BC. 2005. Practical Pollination Biology. Ontario: Enviroquest.

de Jager ML, Ellis AG. 2014. Floral polymorphism and the fitness implications of attracting
pollinating and florivorous insects. *Annals of Botany,* **113**: 213-222.

Eliyahu D, McCall AC, Lauck M, Trakhtenbrot A. 2015. Florivory and nectar-robbing
perforations in flowers of pointleaf manzanita *Arctostaphylos pungens* (Ericaceae) and
their effects on plant reproductive success. *Arthropod-Plant Interactions,* **9**: 613-622.

Fenster CB, Armbuster WS, Wilson P, Dudash MR, Thomson JD. 2004. Pollination
syndromes and floral specialization. *Annual Review of Ecology, Evolution and*
Systematics, **35**: 375-403.

Frey FM. 2004. Opposing natural selection from herbivores and pathogens may maintain floral-
color variation in *Claytonia virginica* (Portulacacea). *Evolution,* **58**: 2426-2437.

Galen C. 2000. High and dry: drought stress, sex-allocation trade-offs, and selection on flower
size in the alpine wildflower *Polemonium viscosum* (Polemoniaceae). *The American*
Naturalist, **156**: 72-83.

Gonzáles WL, Suárez LH, Gianoli E. 2016. Genetic variation in the reduction of attractive
floral traits of an annual tarweed in response to drought and apical damage. *Journal of*
Plant Ecology, **9**: 629-635.

Harder LD, Johnson SD. 2009. Darwin's beautiful contrivances: evolutionary and functional
evidence for floral adaptation. *New Phytologist,* **183**: 530-545.

Herrera CM. 1996. Floral traits and plant adaptation to insect pollinators: a devil's advocate
approach. In: Lloyd DG, Barrett SCH, eds. *Floral biology: studies on floral evolution in*
animal-pollinated plants. New York: Chapman & Hall.

Jogesh T, Overson RP, Raguso RA, Skogen KA. 2016. Herbivory as an important selective
force in the evolution of floral traits and pollinator shifts. *AoB Plants*.

Kessler D, Diezel C, Clark DG, Colquhoun TA, Baldwin IT. 2013. Petunia flowers solve the
defence/apparency dilemma of pollinator attraction by deploying complex floral blends.
Ecology Letters, **16**: 299-306.

Kirk H, Vrieling K, Klinkhamer PGL. 2005. Maternal effects and heterosis influence the
fitness of plant hybrids. *The New Phytologist,* **166**: 685-694.

Kirk WDJ, Ali M, Breadmore KN. 1995. The effects of pollen beetles on the foraging
behaviour of honey bees. *Journal of Apicultural Research,* **34**: 15-22.

Krupnick GA, Weis AE, Campbell DR. 1999. The consequences of floral herbivory for
pollinator service to *Isomeris arborea. Ecology,* **80**: 125-134.

Lavi R, Sapir Y. 2015. Are pollinators the agents of selection for the extreme large size and
dark color in Oncocyclus irises? *New Phytologist,* **205**: 369-377.

Louda SM, Potvin MA. 1995. Effect of Inflorescence-Feeding Insects on the Demography and
Lifetime of a Native Plant. *Ecology,* **76**: 229-245.

Mathew B. 1989. *The Iris.* London: Batsford.

McCall AC. 2008. Florivory affects pollinator visitation and female fitness in *Nemophila*
menziesii. Oecologia, **155**: 729-737.

McCall AC. 2010. Does dose-dependent petal damage affect pollen limitation in an annual
plant? *Botany,* **88**: 601-606.

McCall AC, Irwin RE. 2006. Florivory: the intersection of pollination and herbivory. *Ecology*
Letters, **9**: 1351-1365.

McCall AC, Murphy SJ, Venner C, Brown M. 2013. Florivores prefer white versus pink petal
color morphs in wild radish, *Raphanus sativus. Oecologia,* **172**: 189-195.

Meindl GA, Bain DJ, Ashman T-L. 2013. Edaphic factors and plant–insect interactions: direct
and indirect effects of serpentine soil on florivores and pollinators. *Oecologia,* **173**: 1355-
1366.

Monty A, Saad L, Mahy Gg. 2006. Bimodal pollination system in rare endemic Oncocyclus
irises (Iridaceae) of Lebanon. *Canadian Journal of Botany,* **84**: 1327–1338.

Muola A, Weber D, Malm LE, Egan PA, Glinwood R, Parachnowitsch AL, Stenberg JA.
2017. Direct and Pollinator-Mediated Effects of Herbivory on Strawberry and the
Potential for Improved Resistance. *Frontiers in Plant Science,* **8**: 823.

Pohl N, Carvallo G, Botto-Mahan C, Medel R. 2006. Nonadditive effects of flower damage
and hummingbird pollination on the fecundity of Mimulus luteus. *Oecologia,* **149**: 648-
655.

R Development Core Team. 2014. R: A Language and Environment for Statistical Computing.
Vienna: R Foundation for Statistical Computing.

Rix M. 1997. Section *Oncocyclus* (Siemssen) Baker. In: The species group of the British Iris
Society, ed. *A Guide to Species Irises*: Cambridge University Press.

Ruane LG, Rotzin AT, Congleton PH. 2014. Floral display size, conspecific density and
florivory affect fruit set in natural populations of *Phlox hirsuta*, an endangered species.
Annals of Botany, **113**: 887-893.

Sánchez-Lafuente AM. 2007. Corolla Herbivory, Pollination Success and Fruit Predation in
Complex Flowers: An Experimental Study with *Linaria lilacina* (Scrophulariaceae).
Annals of Botany, **99**: 355-364.

Sapir Y. 2016a. *Iris atropurpurea. The IUCN Red List of Threatened Species 2016*:
e.T13161450A18611400.

Sapir Y. 2016b. *Iris lortetii. The IUCN Red List of Threatened Species 2016*:
e.T13161743A18612235.

Sapir Y, Dorchin A, Mandelik Y. 2015. Indicators of Pollinator Decline and Pollen Limitation.
In: Armon RH, Hänninen O, eds. *Environmental Indicators*: Springer Netherlands.

Sapir Y, Shmida A, Ne'eman G. 2005. Pollination of the *Oncocyclus* irises (*Iris*: Iridaceae) by
night-sheltering male bees. *Plant Biology,* **7**: 417-424.

Sapir Y, Shmida A, Ne'eman G. 2006. Morning floral heat as a reward to the pollinators of the
Oncocyclus irises. *Oecologia,* **147**: 53-59.

Schiestl FP, Johnson SD. 2013. Pollinator-mediated evolution of floral signals. *Trends in
Ecology and Evolution,* **28**: 307-315.

Sletvold N, Grindeland JM, Ågren J. 2010. Pollinator-mediated selection on floral display,
spur length and flowering phenology in the deceptive orchid *Dactylorhiza lapponica.*
New Phytologist, **188**: 385–392.

Sletvold N, Tye M, Ågren J. 2016. Resource- and pollinator-mediated selection on floral traits.
Functional Ecology: doi: 10.1111/1365-2435.12757.

Sletvold N, Tye M, Ågren J. 2017. Resource- and pollinator-mediated selection on floral traits.
Functional Ecology, **31**: 135-141.

Sõber V, Moora M, Teder T. 2010. Florivores decrease pollinator visitation in a self-
incompatible plant. *Basic and Applied Ecology,* **11**: 669-675.

Strauss SY, Whittall JB. 2006. Non-pollinator agents of selection on floral traits. In: Harder
LD, Barrett SCH, eds. *Ecology and Evolution of Flowers*. Oxford Oxford University
Press.

Theis N, Adler LS. 2012. Advertising to the enemy: enhanced floral fragrance increases beetle
attraction and reduces plant reproduction. *Ecology,* **93**: 430-435.

Tsuji K, Dhami MK, Cross DJR, Rice CP, Romano NH, Fukami T. 2016. Florivory and
pollinator visitation: a cautionary tale. *AoB Plants,* **8**.

Vereecken NJ, Dorchin A, Dafni A, Hötling S, Schulz S, Watts S. 2013. A pollinators' eye
view of a shelter mimicry system. *Annals of Botany,* **111**: 1155-1165.

Watts S, Sapir Y, Segal B, Dafni A. 2013. The endangered *Iris atropurpurea* (Iridaceae) in
Israel: honey-bees, night-sheltering male bees and female solitary bees as pollinators.
Annals of Botany, **111**: 395-407.

Yardeni G, Tessler N, Imbert E, Sapir Y. 2016. Reproductive isolation between populations of
Iris atropurpurea is associated with ecological differentiation. *Annals of Botany,* **118**:
971-982.

Zhu Y-R, Yang M, Vamosi JC, Armbruster WS, Wan T, Gong Y-B. 2017. Feeding the
enemy: loss of nectar and nectaries to herbivores reduces tepal damage and increases
pollinator attraction in *Iris bulleyana*. *Biology Letters,* **13**: 20170271.

Tables

Table 1 – Analysis of variance table for the effects of site (Netanya and Yeruham), year (2016

and 2017) and treatment (supplementary or open pollination) on fruit-set and seed-set.

Source	DF	SS	F	Significance
Fruit-set				
Site	1	3.05	15.68	P<0.001
Site/year	2	0.72	1.84	P=0.161
Site/year/treatment	2	10.6	27.13	P<0.001
Residuals	309	60.2		
Seed-set				
Site	1	1063	5.52	P=0.794
Site/year	2	172	0.45	P=0.641
Site/year/treatment	2	5303	13.77	P<0.001
Residuals	305	58724		

Figure legends

Figure 1 –(A) *Iris atropurpurea* in Netanya; (B) *Iris petrana* in Yeruham; (C) *Iris lortetii* in Malkiya; (D) Male *Eucera* bee, the specific pollinator of the royal irises, sheltering within a pollination tunnel of *Iris petrana*; (E-G) Natural florivory in flowers of *Iris atropurpurea* (E), *I. petrana* (F) and *I. lortetii* (G).

Figure 2 –Artificial florivory manipulations exemplified in *Iris atropurpurea*. (A) Within flower manipulation – each floral unit treated as either high florivory (>50% petal cut), low florivory (10-30% petal removed by hole puncher), or control (no treatment). These treatments were used for testing the effect on pollination. (B–D) Flowers used for testing the effect of florivory on fitness. (B) Flower treated as high florivory. (C) Flower treated as low florivory. (D) Control flower.

Figure 3 – Fruit-set and seed-set in pollen limitation experiment in *Iris atropurpurea* in Netanya, and *I. petrana* in Yeruham. Bars are fractions ± standard errors. (A) Fraction of flowers that produced fruits in 2016 (B) and in 2017. (C) Mean number of seeds in 2016 (D) and in 2017. Open – flowers open to natural pollination. Supp. – flowers received supplementary pollen artificially.

Figure 4–Mean number of pollen grains counted on stigma (± standard errors) as a function of florivory treatment in three species.(A) Pollen grains in 2016 experiment; (B) Pollen grains in 2017 experiment. Bars are fractions ± standard errors.

Figure 5–Fitness as a function of artificial florivory manipulations. (A) Fruit-set (fraction of flowers that set fruits) in three species in 2017 and in 2017 (B). (C) Seed-set (mean number of seeds in a fruit) in 2016 and in 2017 (D). Bars are means ± standard errors.

Figure 6–Hypothetical floral trait evolution as a function of selection by both mutualists (pollinators) and antagonists (florivores). (A) When pollinators and florivores exert selection in the same direction, the concert selection favors the same trait optimum. (B) When pollinators and florivores exert opposing selection on a trait, an intermediate trait optimum is favored. A and B

adapted from figure 7.1 in Strauss &Whittall (2006). (C) When florivores have no preference, or

their effect is similar to mean fitness derived from pollinators' effect, trait will have optimum

fitness selected by pollinators like in A.

Figure 1

A   B   C   D

E   F   G

Figure 2

A   B

570   C   D

     Figure 3

573   Figure 4

575   Figure 5

577   Figure 6