1

Global connectivity of the frontoparietal cognitive control network is related to

depression symptoms in the general population

Douglas H. Schultz[1], Takuya Ito[1], Levi I. Solomyak[1], Richard H. Chen[1], Ravi D.

Mill[1], Alan Anticevic[2], & Michael W. Cole[1]

[1]Center for Molecular and Behavioral Neuroscience,

Rutgers University – Newark

[2]Department of Psychiatry,

Yale University

Corresponding author: Douglas H. Schultz

Center for Molecular and Behavioral Neuroscience

197 University Ave, Suite 212

Newark, NJ 07102

dhs95@newark.rutgers.edu

KEYWORDS: fMRI, depression, frontoparietal network, functional connectivity

ABSTRACT

We all vary in our mental health, even among people not meeting diagnostic

criteria for mental illness. Understanding this individual variability may reveal

factors driving the risk for mental illness, as well as factors driving sub-clinical

problems that still adversely affect quality of life. To better understand the large-

scale brain network mechanisms underlying this variability we examined the

relationship between mental health symptoms and resting-state functional

connectivity patterns in cognitive control systems. One such system is the

frontoparietal cognitive control network (FPN). Changes in FPN connectivity may

impact mental health by disrupting the ability to regulate symptoms in a goal-

directed manner. Here we test the hypothesis that FPN dysconnectivity relates to

mental health symptoms even among individuals who do not meet formal

diagnostic criteria but may exhibit meaningful symptom variation. We found that

depression symptoms severity negatively correlated with between-network global

connectivity (BGC) of the FPN. This suggests that decreased connectivity

between the FPN and the rest of the brain is related to increased depression

symptoms in the general population. These findings complement previous clinical

studies to support the hypothesis that global FPN connectivity contributes to the

regulation of mental health symptoms across both health and disease.

AUTHOR SUMMARY

Understanding how large-scale network interactions in the brain contribute to (or

serve a protective role against) mental health symptoms is an important step

toward developing more effective mental health treatments. Here we test the

hypothesis that cognitive control networks play an important role in mental health

by being highly connected to other brain networks and able to serve as a

feedback mechanism capable of regulating symptoms in a goal-directed manner.

We found that the more well-connected the frontoparietal cognitive control

network was to other networks in the brain the less depression symptoms were

reported by participants. These results contribute to our understanding of how

brain network interactions are related to mental health symptoms, even in

individuals who have not been diagnosed with a disorder.

INTRODUCTION

People vary in their degree of mental health. Indeed, people who do not meet formal criteria for mental illness as defined by our current diagnostic systems (American Psychiatric Association, 2013; World Health Organization, 1992) may still experience a number of symptoms associated with that disorder (World Health Organization, 2017). Here we use this natural variability to better understand neural factors potentially contributing to day-to-day experiences of poor mental health, as well as (prodromal) factors that may elevate the risk for severe mental illness. We hypothesized that the variability observed in mental health symptoms among individuals is related to the function of the frontoparietal cognitive control network (FPN), based on our previously-developed theoretical proposal that the FPN plays a domain-general protective role against mental health symptoms (Cole et al., 2014).

The proposed theoretical framework suggests that alterations in FPN function may play a common role in multiple mental disorders by disrupting a domain-general cognitive control feedback mechanism that can regulate symptoms when they are experienced (Cole et al., 2014). The FPN is a candidate network for this function because it is a flexible hub, meaning it has a high degree of connectivity across the brain (Cole et al., 2010; Power et al., 2011) and can rapidly modify functional connections according to current goals (Cole et al., 2013). There is strong evidence that these FPN functions are domain general (Chein and Schneider, 2005; Cole et al., 2013; Dosenbach et al., 2007), such that individual differences in the general ability to regulate cognition can

influence symptoms. Finally, alterations in FPN functional connectivity (FC) have

been identified in a number of mental disorders including: depression Kaiser et

al., 2015), anxiety (Sylvester et al., 2012), schizophrenia (Baker et al., 2014; Cole

et al., 2011; Fornito et al., 2011; Yang et al., 2016), attention deficit hyperactivity

disorder (Li et al., 2014; Park et al., 2016), and eating disorders (Boehm et al.,

2014; Cowdrey et al., 2014). Consistent with most of these studies, we focus

here on FC measured using functional magnetic resonance imaging (fMRI),

calculated as the temporal relationship in the blood oxygenation level dependent

(BOLD) signal between brain regions (Biswal et al., 1995) while participants rest

in the scanner.

Consistent with the flexible hub framework, a number of studies using

different measures have provided converging evidence that the FPN is especially

well connected to the rest of the brain (Buckner et al., 2009; Cole et al., 2010).

Both of these studies calculated a summary statistic reflecting the degree of

connectivity across the whole brain. However, these estimates can be influenced

by the relative size of different networks. For example, nodes of a larger network

will have a larger overall number of strong connections than nodes of a smaller

network simply because, by definition, within-network connections are stronger

on average than between-network connections (Power et al., 2013; Wig et al.,

2011). Therefore, we estimated how well each region of the brain was connected

to the rest of the brain using between-network global connectivity (BGC) (Ito et

al., 2017), a measure not influenced by network size.

Particularly important for our specific test of the flexible hub framework

here, patients diagnosed with major depression exhibit differences in FC patterns

throughout the brain, including FPN functional connections (Brakowski et al.,

2017). Specifically, connectivity between regions of the FPN is decreased in

depressed individuals (Alexopoulos et al., 2012), as well as in undiagnosed

individuals experiencing depression symptoms (Wei et al., 2014). However, Wei

and colleagues (2014) looked at FC with specific seed regions, not global

connectivity, in their sample. Another study found that global brain connectivity

was decreased in the medial prefrontal cortex and the dorsolateral prefrontal

cortex (dlPFC) portions of the FPN in depressed patients (Murrough et al., 2016).

Depression has been shown to generally disrupt FC in the dlPFC as well as the

default mode network (DMN) (for review see: Gong and He, 2015). Decreases in

within-network FPN connectivity have also been observed in a group of

individuals reporting depression symptoms in the absence of a clinical diagnosis

(Hwang et al., 2015). Researchers have also attempted to subdivide depression

into various types based on FC patterns. Decreases in FPN connectivity were

most pronounced in one particular subtype of depression associated with

increased symptoms of fatigue and decreased symptoms of anxiety (Drysdale et

al., 2016). Furthermore, activation in the FPN and changes in FPN FC have been

observed when participants are actively attempting to regulate their emotions

(Banks et al., 2007; Buhle et al., 2014; Kohn et al., 2014) and when participants

are in threatening situations (Balderston et al., 2017). Decreased global brain

connectivity, the mean FC value for each region's connections, has been

observed in the prefrontal cortex in depression patients (Abdallah et al., 2016).

This decrease in global brain connectivity was rescued 24 hours following

ketamine treatment. These previous results are broadly consistent with our

hypothesis, yet the extension of results to test whether FPN BGC is related to

mental health symptoms among healthy individuals would provide important new

evidence for the general nature of FPN's role in regulating mental health.

Consistent with our previously-developed theoretical framework (Cole et

al., 2014) along with observed FPN FC alterations in patients with major

depression, we hypothesized that individual differences in depression symptoms

in undiagnosed individuals would be correlated with BGC in the FPN. Support for

our hypothesis would provide important evidence for a potentially general role of

global FPN intrinsic FC in facilitating the regulation of mental health symptoms.

METHODS

Participants

Data were collected at the Rutgers University Brain Imaging Center

(RUBIC). The participants were recruited from the Rutgers University-Newark

campus and surrounding community. All participants provided informed consent

and all procedures were approved by the Rutgers University-Newark Institutional

Review Board. We collected data from 106 participants. Technical error or

equipment malfunction during the scanning session resulted in removing six

participants from the study. Four participants were removed from the study

because they did not complete the Center for Epidemiological Studies

Depression Scale (CESD) during a behavior-only session separate from the MRI

session. We also collected demographic information and asked participants 11

questions asking what hand they used for various activities including writing,

throwing, using a scissors, holding their toothbrush, striking a match, opening a

box, kicking, using a knife, using a spoon, which hand was placed on top while

using a broom, and which eye they used in situations where they would only

being using one eye. They replied to each question by answering always right,

usually right, no preference, usually left, or always left. Answers were scored in

the following manner: always right (2), usually right (1), no preference (0), usually

left (-1), always left (-2). We calculated a laterality quotient (LQ) by summing

these scores and dividing by the maximum score of 22. A LQ score of -100

indicates a strong left hand preference, 0 indicates no hand preference, and 100

indicates a strong right hand preference. All participants self reported as right

handed. However, one participant demonstrated a LQ indicating they had a left

hand preference. The participant with a left hand preference has been included in

all analyses.

Studies have proposed that the CESD can be broken down into between

one and four factors (Cole et al., 2004; Herrero and Meneses, 2006). In addition

to calculating the raw CESD score for each participant we also calculated three

factor scores: somatic symptoms, negative affect, and anhedonia, based on a

recent study (Carleton et al., 2013). Participants also completed several

measures of flexible cognition during the behavior-only session. These measures

included Raven's progressive matrices (Bilker et al., 2012), Cattell's culture fair

test (Cattell and Horn, 1978), and Duncan's goal neglect task (Dumontheil et al.,

2011). The final sample consisted of 96 participants (See Table 1).

Table 1. Demographic information

N	96
Age	$M = 22.06,\ SD = 3.84$
Gender	
Male	42 (43.8%)
Female	54 (56.2%)
Handedness (LQ)	$M = 74.67,\ SD = 25.2$
Education (Highest level)	
High school	14 (14.6%)
Some college	50 (52.1%)
Bachelor's degree	29 (30.2%)
Graduate degree	3 (3.1%)
Cognitive measures	(Percent correct)
Raven	$M = 52.29,\ SD = 15.69$
Cattell	$M = 55.20,\ SD = 12.03$
Duncan	$M = 76.19,\ SD = 18.34$
CESD (raw score)	$M = 17.41,\ SD = 9.26$

MRI Parameters

Multiband whole-brain echo-planar imaging (EPI) acquisition was collected using a 32-channel head coil on a 3T Siemens Trio MRI scanner with the following parameters: TR = 785 ms, TE = 34.8 ms, flip angle = 55°, Bandwidth 1924/Hz/Px, in-plane FoV read = 208 mm, 72 slices, 2.0 mm isotropic voxels, with a multiband acceleration factor of 8. Whole-brain high-resolution T1-weighted and T2-weighted anatomical scans with 0.8 mm isotropic voxels were also collected. Spin echo field maps were collected in both the anterior to posterior direction and the posterior to anterior direction consistent with the Human Connectome Project preprocessing pipelines (Glasser et al., 2013). The resting-state fMRI scan was 14 minutes (1070 TRs) in duration.

fMRI Preprocessing

Functional imaging data were preprocessed using the Human Connectome Project minimal preprocessing pipeline version 3.5.0. Preprocessing consisted of anatomical restructuring and segmentation, EPI reconstruction, segmentation, and spatial normalization to a standard template, intensity normalization, and motion correction (Glasser et al., 2013). All further processing was conducted in CIFTI 64k greyordinate standard space. The data were parcellated into 360 regions described previously (Glasser et al., 2016), taking the average time series of the vertices within a parcel. At this point all subsequent data analysis was conducted in MATLAB R2014b (The Mathworks). We performed nuisance regression using 12 motion parameters, ventricle and

white matter timeseries (as well as their first derivatives), and global signal. We

also performed motion scrubbing (Power et al., 2012) based on framewise

displacement. Framewise displacement was calculated as the amount of head

movement for each frame relative to the previous in terms of Euclidean distance.

We next applied a low pass temporal filter (0.3 Hz) to the framewise

displacement vector in order to reduce the effect of respiration on our framewise

displacement measure (Siegel et al., 2016). The framewise displacement

threshold for motion scrubbing was set at 0.3 mm. Motion scrubbing consisted of

removing the flagged frame from the timeseries as well as one frame prior and

the two frames following. FC was estimated by calculating the Pearson

correlation of the BOLD timeseries between each pair of the parcels defined by

Glasser and colleagues (Glasser et al., 2016).

Network Assignment and Analysis

The network assignment of each of the parcels was completed on an

independent dataset, the Human Connectome Project (100 unrelated) (Van

Essen et al., 2013). Briefly, each parcel was assigned to a network using the

Generalized Louvain method for community detection. This process was

conducted using resting-state data. We identified 12 functional networks (Spronk

et al., 2017) (See Figure 1). The functional network topology findings replicate

the major features of several previously published network partitions (Gordon et

al., 2016; Power et al., 2011; Yeo et al., 2011).

Cortical partition

- Visual 1
- Visual 2
- Somatomotor
- Cingulo-Opercular
- Dorsal Attention
- Language
- Frontoparietal
- Auditory
- Default Mode
- Posterior Multimodal
- Ventral Multimodal
- Orbito-Affective

Figure 1 - Network assignment. Resting-state fMRI data from an independent

dataset (HCP: 100 unrelated) was used to assign each parcel to a network using

a community detection algorithm. This resulted in 12 total networks. Color

indicates the network assignment for each parcel.

We were interested in the relationship between the frequency at which

participants experienced depression symptoms and the degree of between-

network global connectivity (BGC). Specifically, we were interested in a measure

that would estimate the strength of FC for each region of a network to all of the

other regions in different networks. BGC was calculated for each region

individually and defined as the mean FC for all out-of-network connections. Out-

of-network connections were defined as all connections from a source region to

target regions outside the source region's network. This process was completed

for all regions until we had a BGC value for each region in the brain. Then we

calculated the mean BGC value for each of the functional networks to summarize

effects at the network level.

239   More formally, BGC was defined for each region as:

$$BGC_i = \frac{\sum_{j \notin C} W_{ij}}{N_{total} - N_C}$$

240   where BGC_i corresponds to the out-of-network weighted degree of region i in

241   network C, $j \notin C$ corresponds to all regions not in network C, W_{ij} corresponds to

242   the FC estimate between regions i and j, N_{total} corresponds to the total number

243   of regions, and N_C corresponds to the total number of regions in network C.

244   We tested the relationship between BGC and depression symptoms in all

brain networks. Due to the large number of statistical tests we report the false-

discovery rate (FDR)-corrected *p*-values for primary analyses (Benjamini and

Hochberg, 1995). Our main hypothesis was that BGC in the FPN was correlated

with depression symptoms. Because this was a primary hypothesis based on our

previous publication (Cole et al., 2014) we report the uncorrected p-value.

Follow-up and control analyses were not independent of the primary tests and

are therefore not corrected.

RESULTS

Depression Symptom Scores

255   We hypothesized that the frequency of experiencing depression

symptoms is related to individual differences in the between-network global

connectivity of the FPN. This would be consistent with the involvement of FPN in

domain-general cognitive regulation, broadly construed to include emotion (Cole

et al., 2014). We used a standard measure of depression symptoms, the CESD,

to measure the frequency of depression symptoms. The CESD consists of 20

questions asking how frequently participants have experienced symptoms related

to depression over the last seven days. CESD scores ($M = 17.41, SD = 9.26$)

varied from a minimum of 0 to a maximum of 43 (out of a possible 60) (Figure 2).

CESD scores from 16 to 24 indicate mild levels of depression symptoms while

scores above 24 indicate moderate to severe levels of depression. In our sample

48.9% of participants demonstrated at least mild levels of depression, while 25%

of participants exhibited at least moderate levels of depression. The

characteristics of our sample were consistent with previous studies using the

CESD with undiagnosed young adults (Gress-Smith et al., 2015; Van Dam and

Earleywine, 2011). However, the scores were not normally distributed based on a

Kolomogorov–Smirnov test ($p < 0.023$). We therefore used the Box-Cox power

transformation (Box and Cox, 1964), a common approach to correct for non-

normality. After the transformation the data no longer deviated from a normal

distribution ($p = 0.17$). The transformed data were used for all subsequent

analyses. CESD scores were not correlated with any of the flexible cognition

measures (*smallest p = 0.28*).

Figure 2 – Histogram of raw CESD scores.

We also calculated CESD sub-scores for somatic, negative affect, and

anhedonia factors. After Box-Cox transformation none of the factor score

distributions significantly deviated from a normal distribution (*somatic p = 0.4,*

negative affect p = 0.19, anhedonia p = 0.056). All three of the factors were

significantly correlated with the overall CESD scores (*somatic r = 0.77,*

uncorrected p < 0.001; negative affect r = 0.80, uncorrected p< 0.001; anhedonia

r = 0.52, uncorrected p < 0.001, See Table 2). The three factors were also

correlated with each other (*correlation between somatic and negative affect r =*

0.55, correlation between somatic and anhedonia r = 0.18, correlation between

negative affect and anhedonia r = 0.26, See Table 1). Although all of the factors

were highly related to the overall CESD score, they appear to have contributed

unique variance as the highest correlation between any two factors was only

accounting for 30% of the variance (based on R^2 values).

Table 2. Correlation between the CESD and CESD factors

	Overall CESD	Somatic	Negative Affect	Anhedonia
Overall CESD	-----	-----	-----	-----
Somatic	0.77	-----	-----	-----
Neg. Affect	0.80	0.55	-----	-----
Anhedonia	0.52	0.18	0.26	-----

Between-Network Global Connectivity

We calculated BGC values using resting-state fMRI data for each

functionally defined cortical parcel (Figure 3). DMN regions tended to have lower

BGC values, consistent with the conceptually-similar participation coefficient

results reported by Power et al. (2011). In contrast, the FPN had high levels of

BGC in lateral prefrontal portions of the network ($M = 0.002, SD = 0.02$) relative

to the mean of regions outside of the FPN ($M = -0.0081, SD = 0.013; t(95) =$

$5.08, p < 0.0001$). However, FPN had lower BGC values in parietal ($M = -0.0258,$

$SD = 0.0190$), inferior frontal ($M = -0.0177, SD = 0.0209$) and temporal lobe

regions of the network ($M = -0.0420, SD = 0.0221$) relative to the mean of

regions outside of the FPN (*largest p < 0.0001*). Sensory networks exhibited two

patterns. The visual network had some of the lowest BGC values in the brain with

the primary visual cortex region showing the least BGC in the network ($M = -$

0.0742, SD = 0.0348). In contrast, the auditory network (*M = 0.0212, SD =*

0.0171) and portions of the sensorimotor network surrounding the central sulcus

(*M = 0.0117, SD = 0.0249*) both showed a high degree of BGC. The

sensorimotor regions closest to the central sulcus showed more moderate BGC

scores (*M = -0.0139, SD = 0.0289*).

Figure 3 – Between network global connectivity (BGC) across all cortical

regions. Between network global connectivity for region A is defined as the

mean FC (Pearson correlation) between that region A and all other regions

outside of region A's network. Warm values indicate positive values and cool

values indicate negative values.

Correlation Between BGC and Depression Symptoms

We next tested for a relationship between BGC and depression

symptoms. This involved creating a mean BGC score for each functional network

and testing for correlation between those values and the depression symptom

measure (See Table 3). Consistent with our a priori hypothesis, we found that

FPN BGC was significantly correlated with depression symptoms ($r = -0.247$, $p =$

0.015) (Figure 4A). Unexpectedly, we found that DMN and language network

BGC showed a similar magnitude effect (DMN $r = -0.241$, uncorrected $p = 0.018$;

language $r = -0.268$, uncorrected $p = 0.008$) (Figure 4B & 4C). These findings

suggest that greater FC between the FPN and the rest of the brain (outside of the

FPN) is related to less frequent depression symptoms. Less depression

symptoms are also related to greater BGC in the DMN and language networks.

We report an uncorrected p-value for the test of our a priori hypothesis that FPN

BGC would be related to depression symptoms. We also examined the

relationship between depression symptoms and BGC in the other 11 brain

networks. These p-values do not survive FDR correction for multiple

comparisons, but because the magnitude of the effect observed in the DMN and

language network was similar to that in the FPN we continue to report

subsequent uncorrected statistics for these networks. Although it should be noted

that the DMN and language network results should be interpreted with caution as

they were not a priori hypotheses and do not survive multiple comparison

correction.

To address whether these findings were specific to this particular network

partition we conducted a similar analysis using a different network partition

scheme (Ito et al., 2017). This network partition includes a FPN and DMN, but

lacks a network analogous to the language network. Using this network partition

we found a similar negative correlation between BGC and depression symptoms

in the FPN (*r = -0.262, p = 0.009, FDR adjusted for multiple comparisons p =*

0.044) and DMN (*r = -0.250, p = 0.014, FDR adjusted p = 0.049*). The results for

the FPN and DMN both survive correction for multiple comparisons with this

network partition.

We conducted a similar analysis, correlating BGC and depression

symptoms on the region level, to examine if there were particular brain regions

within these networks driving the network level results. None of the individual

brain regions showed a significant correlation after correcting for multiple

comparisons using FDR. This suggests that the network level results are not

being driven by particularly strong relationships in smaller number of brain

regions within these networks.

Table 3. Correlation between BGC and CESD scores

Network	Pearson's r	uncorrected p-value
Visual 1	0.015	0.884
Visual 2	0.020	0.850
Somatomotor	-0.177	0.085
Cingulo-Opercular	-0.143	0.164
Language	-0.268	0.008
Default Mode	-0.241	0.018
Frontoparietal	**-0.247**	**0.015***
Auditory	-0.099	0.340
Post. Multimodal	-0.110	0.284
Dorsal Attention	-0.113	0.275
Ventral Multimodal	0.116	0.261
Orbito-Affective	-0.078	0.448

Figure 4 – CESD scores and BGC in the FPN, DMN, and language networks are negatively correlated. A) BGC in the FPN is plotted on the y-axis and CESD scores are plotted on the x-axis. B) BGC in the DMN is plotted on the y-axis and CESD scores are plotted on the x-axis. C) BGC in the language network is plotted on the y-axis and CESD scores are plotted on the x-axis.

In order to verify that these correlations were not dependent on the Box-

Cox transformation we used to normalize the depression symptom data we also

ran non-parametric Spearman's rank correlations, which do not require normally

distributed data. The results of testing our a priori hypothesis that BGC in the

FPN would be related to depression symptoms were unchanged. Specifically, the

BGC-depression rank correlation was consistent with significant Pearson

correlations (FPN $rho = -0.227, p = 0.026$). The Spearman rank correlations also

reflected similar results to the Pearson correlation in the DMN ($rho = -0.220,$

uncorrected p = 0.031) and the language network (*rho = -0.285, uncorrected p =*

0.005).

To ensure that our observed correlation between depression symptoms

and BGC in the FPN was not being driven by other factors we removed the

variance in the CESD scores that could be accounted for by age and gender

factors with a regression model and used the residuals to rerun the correlation.

We still found a significant relationship between BGC in the FPN and depression

symptoms (*r = -0.220, p = 0.031*). This result suggests that the relationship

between BGC in the FPN and depression scores is not being influenced by age

or gender differences in our sample.

Next, we tested if the relationships between BGC and overall depression

scores were driven by specific depression factors. We tested for correlation

between the BGC values and three factors derived from the CESD questionnaire:

somatic symptoms, negative affect, and anhedonia symptoms (Carleton et al.,

2013). The three factors were correlated with one another so we also used a

partial correlation approach to explore the relationship between the unique

variance contributed by each factor and BGC in the FPN, DMN and language

networks. The partial correlation approach allows us to examine the relationship

between BGC and each of the factors while accounting for the variance

contributed by the other factors. After accounting for the variance contributed by

other factors we found that none of the factors were significantly correlated with

BGC in the FPN (*largest rho = -0.110, p = 0.292*). We also found that the unique

variance from each factor was not correlated with BGC in the DMN (*largest rho =*

-0.137, p = 0.188). BGC in the language network was not significantly correlated

with the unique variance contributed by any of the factors (*largest rho = -0.185, p*

= 0.075). These results suggest that the relationships we observed between the

overall CESD scores and BGC were driven by a combination of all factors, and

not specific features of depression.

Observed BGC-depression Correlations were not Dependent on

Connections Between FPN, DMN, and language network

The magnitude and direction of the BGC-depression correlations were

similar for the FPN, DMN, and language network. One possible explanation for

these results is that the observed BGC-depression effects for these networks

were driven primarily by the FC between them. We tested this possibility by

recalculating BGC for each network by removing connections between the FPN,

DMN, and language network from the calculation. This approach minimizes the

possibility that connections between these networks drove the original

correlations between depression symptoms and BGC The magnitude and

direction of the correlations between depression symptoms and BGC in the FPN

(*r = -0.226, p = 0.027*), the DMN (*r = -0.236, uncorrected p = 0.021*), and the

language network (*r = -0.201, uncorrected p = 0.049*) were similar when we

excluded the connections between these networks from the BGC calculation.

This suggests the original BGC-depression correlations were not dependent on

connections between the FPN, DMN, and language network.

In order to further explore the possibility that the observed relationship

between depression symptoms and BGC in the FPN might be primarily driven by

FPN connections to the DMN and language network we calculated the

correlation between the mean FPN to DMN, and FPN to language network FC

with depression symptoms. There was not a significant correlation between

depression symptoms and the mean connectivity between the FPN and DMN ($r =$

-0.016, uncorrected p = 0.881) or the mean connectivity between the FPN and

language network ($r = -0.146$, *uncorrected p = 0.156*). This suggests that the

relationship between FPN BGC and depression symptoms is not driven by

connections to the DMN and language networks, which both showed a negative

correlation between BGC and depression symptoms as well.

Within-Network Connectivity does not Correlate with Depression

Symptoms

We focused on between-network connectivity, yet the degree of

connectivity within a functional network might also be related to depression

symptoms. For example, a high degree of FC within a network may reflect more

homogeneous network activity. In this case the network would be tightly coupled

and all of the component regions would likely be serving a very similar function.

However, a lower degree of FC within a network may reflect more heterogeneous

processing. In this case the different regions of a functional network may all be

involved in the same general process, but they may be contributing in different

ways. These differences in within-network functionality may, in turn, relate to

depression symptoms.

Consistent with our choice to focus primarily on between-network effects,

within-network FC in the FPN was not significantly correlated with depression

symptoms ($r = 0.066, p = 0.526$). Within-network FC in the DMN was also not

significantly correlated with depression symptoms ($r = 0.143$, *uncorrected p =*

0.164). The language network showed a negative correlation between

depression symptoms and within-network FC ($r = -0.219$, *uncorrected p = 0.032*).

We observed a significant difference in the DMN correlation coefficients

comparing BGC ($r = -0.241$) and within-network connectivity ($r = 0.143$) with

depression symptoms ($z = 2.66, p = 0.008$). It will be important for future studies

to test this effect for replication, given the non-significance of the DMN within-

network FC-depression effect relative to a correlation of 0.

DISCUSSION

Based on convergent evidence across a variety of mental health

conditions we predicted that individual differences in FPN BGC would be

correlated with symptoms associated with depression (Cole et al., 2014). This is

consistent with extensive evidence that the FPN is a domain-general cognitive

control system regulating general goal pursuit processes (Cole and Schneider,

2007; Duncan, 2010; Schneider, 2003), including regulation of mental illness

symptoms (Cole et al., 2014). We sought to expand the general relevance of this

framework by including symptoms experienced in everyday life, which express

dimensionally across health and disease. This involved investigating the

relationship between the mood symptoms, one of the most commonly

experienced set of symptoms in the general population (Centers for Disease

Control and Prevention (CDC), 2010), and FPN global connectivity properties.

We further ensured the broad relevance of the findings by focusing on how a

general population who had not been diagnosed with a mental disorder varied in

their depression symptoms and primarily experienced them in the normal (i.e.,

sub-clinical) range.

481   As predicted, we found a significant relationship between how well-

connected the FPN was to the rest of the brain and the frequency of depression

symptoms in adults who had not been diagnosed with depression. We used BGC

to estimate the degree of brain-wide connectivity for each brain region. The

results suggest that individuals who report fewer depression symptoms are

characterized by a FPN that is more connected to the rest of the brain. These

results support our previously developed theoretical framework, suggesting

natural variance in FPN global connectivity influences each individual's ability to

regulate mood symptoms in everyday life. Extending this framework to test if

FPN function relates to natural variation in other symptom domains will be critical

as well as examining if such effects persist when an individual crosses a

threshold necessary for a formal diagnosis.

Between-network global connectivity identifies how well each brain region is connected to other brain networks

We hypothesized that individuals exhibiting greater FC between the FPN and the rest of the brain would report fewer depression symptoms. We used BGC, a measure that calculates the mean FC between each brain region and all other out-of-network brain regions (Ito et al., 2017), to evaluate this hypothesis. BGC reduces the potential bias of other graph centrality measures, which can be inflated in regions assigned to large networks (Power et al., 2013). Previous methods using global brain connectivity and degree, which both include within network connections, result in DMN regions showing high connectivity with the rest of the brain (Buckner et al., 2009; Cole et al., 2010; Liang et al., 2013). However, we found that BGC was relatively low in the DMN. This is consistent with DMN results from studies using participation coefficient (Power et al., 2013, 2011). While our measure of BGC showed similarity to results obtained using participation coefficient (Power et al., 2011), there were some differences. Some of these differences may be due to the differences between BGC and participation coefficient. Participation coefficient is generally calculated by first thresholding a graph and then examining the distribution of edges between networks. A high value indicates that the edges are relatively evenly distributed to other networks and lower values indicate that the edges tend to preferentially connect nodes in a smaller number of other networks. BGC does not require thresholding a graph prior to calculation and simply measures the mean FC weight for each region to all other out of network regions. These results suggest

that previous findings identifying the DMN as highly connected to the rest of the

brain are largely driven by high FC within the DMN, and do not necessarily reflect

greater connectivity between the DMN and nodes in other functional networks.

The results we observed with BGC were similar to previous attempts to

identify hubs and the most well connected regions in the brain. BGC was high in

the lateral prefrontal cortex, the motor and tactile cortex, the auditory cortex, and

higher order visual regions. Lateral prefrontal regions and higher order visual

areas show a greater degree of FC (Buckner et al., 2009) and a higher global

brain connectivity (Cole et al., 2010). Higher global brain connectivity has also

been reported in the lateral prefrontal cortex, higher order visual regions, auditory

cortex, and somatosensory cortex (Liang et al., 2013). We observed that BGC

was consistent with previous attempts to classify the degree of connectivity in

many brain regions.

There were some discrepancies between previous methods and BGC. We

found that BGC was quite low in lower visual regions in contrast to higher

connectivity estimates calculated by others (Cole et al., 2010; Liang et al., 2013).

The differences observed between BGC and other measures of connectivity

strength may be driven by BGC not considering the relatively strong local

connections within the visual network. In fact, primary and secondary visual

cortex show high local connectivity relative to distant connectivity strength

(Sepulcre et al., 2010). It will be important for future studies to identify whether

differences observed in primary sensory cortices are driven by an imbalance

between within and between-network connectivity.

BGC in the FPN is negatively correlated with depression symptoms

BGC in the FPN is negatively correlated with depression symptoms

We found that BGC in the FPN showed a significant negative correlation

with depression symptoms. This suggests that individuals exhibiting greater

connectivity of the FPN with the rest of the brain experience fewer symptoms of

depression. The correlation between BGC in the FPN and depression symptoms

supports our hypothesis that a well-connected FPN may serve a protective role

against depression symptoms and possibly mental health symptoms in general

(Cole et al., 2014).

Decreases in FPN FC have been reported in individuals diagnosed with

major depression (Alexopoulos et al., 2012; Murrough et al., 2016; Veer, 2010).

Differences in FC patterns can also be used to divide depression into distinct

subtypes (Drysdale et al., 2016). We build on these findings by observing similar

results in a variable sample of undiagnosed individuals that likely includes

primarily mentally healthy, but also some mentally unhealthy, participants.

Another study examining FC in a group of undiagnosed individuals with higher

levels of depression symptoms found that connectivity between the superior

parietal lobule and the dlPFC portion of the FPN was decreased (Wei et al.,

2014). Undiagnosed participants experiencing more depression symptoms have

also been reported to have reduced FC between dlPFC and the supramarginal

gyrus, insula, operculum, precuneus and parahippocampal gyrus (Hwang et al.,

2015). Previous studies have also found that anti-depressant and

electroconvulsive therapy treatments can modulate prefrontal cortex FC in

depression patients (Perrin et al., 2012; Wang et al., 2015). It is possible that

future therapeutic interventions could target and attempt to alter aspects of

network communication. Our findings suggest that greater global connectivity of

the FPN is related to reduced depression symptoms. Our data regarding global

connectivity differences is consistent with a recent study that found reduced

global brain connectivity in the lateral PFC in depression patients (Abdallah et al.,

2016). Here we extend the relationship between global connectivity in the FPN to

include samples that have not been diagnosed with a clinical disorder and further

refine the global connectivity findings by suggesting that the between network

connections are particularly important. Another recent study has suggested that

the FPN can be split into two subsystems, one more involved in introceptive

processes, and another more involved in the regulation of perceptual attention

(Dixon et al., 2018). Future studies should examine if FC relationships with

depression are stronger in the introceptive FPN system that may be more

involved in rumination symptoms.

Unexpectedly, we also found a negative correlation between BGC in the

DMN and depression symptoms. This result did not survive multiple comparison

correction, but because it was of a similar magnitude as our predicted

relationship in the FPN we have explored it. It is important to note that the results

in the DMN should be interpreted with caution and future studies should attempt

to replicate these results. DMN activity has been linked to rumination symptoms

in depression (Hamilton et al., 2011). This finding may reflect an inability for

individuals experiencing more depression symptoms to disengage the DMN in

situations when attentional or cognitive resources need to be allocated (Sheline

et al., 2009). Depression patients also exhibit decreased FC between the DMN

and executive networks (Abbott et al., 2013; Manoliu et al., 2014). These findings

provide further support for the interpretation that depressed individuals may have

a difficult time disengaging the DMN because the connections from cognitive

control networks to the DMN are decreased.

A number of studies have suggested that FC within portions of the DMN is

increased in depression patients (Greicius et al., 2007; Li et al., 2013). Although

we did not find a significant correlation between within-network FC in the DMN

and depression symptoms, we did see a trend toward a positive correlation. Our

lack of a correlation between within-network FC in the DMN and depression

symptoms could be due to methodological differences in using a population of

individuals who have not been diagnosed with a disorder versus comparing a

control group to a group diagnosed with depression. Additionally, we considered

the entire DMN rather than using a seed correlation approach. It will be important

for future studies to investigate the relationship between within-DMN FC and

depression symptoms, verifying this effect in a larger sample and tying it to

specific neural mechanisms and specific depression symptoms.

We also found a negative correlation between BGC in the language

network and depression symptoms. Similar to the DMN, this result did not survive

multiple comparison correction, but because the magnitude of the effect was

similar to that of our predicted relationship between BGC in the FPN and

depression symptoms we have explored it. Because we did not have an a priori

prediction about this network and the result did not survive multiple comparison

correction, the interpretation of this result should be tempered. Decreases in FC

in the language network have been reported in depression patients (Buchanan et

al., 2014) and language performance deficits have also been reported in

depression patients (Baune et al., 2010). The current results might be consistent

with these observed deficits in language in depressed individuals. However,

future studies should attempt to replicate these results in an independent large

sample.

Conclusion

The current study sought to test our previously-developed framework that

suggests the FPN (along with other cognitive control networks) acts as a

protective factor against mental disease via its widespread FC with other

networks (Cole et al., 2014). We identified a negative correlation between

depression symptoms and a measure of between-network, global FC in the FPN

in a sample of individuals from the general population who had not been

diagnosed with depression. These results suggest the human brain's global

network architecture is critical for maintaining mental health even in undiagnosed

individuals, supporting the possibility that FPN maintains a goal-directed

feedback loop to regulate symptoms as they arise. It will be important for future

research to characterize the exact mechanisms by which FPN influences

symptoms, and to assess the possibility of enhancing FPN FC in the interest of

reducing symptoms and potentially preventing the onset of mental illness.

Acknowledgements

This work was supported by the National Institute of Health (MH096801, MH109520, and AG055556). The content is solely the responsibility of the authors and does not necessarily represent the official views of any of the funding agencies.

References

Abbott, C.C., Lemke, N.T., Gopal, S., Thoma, R.J., Bustillo, J., Calhoun, V.D., Turner, J.A., 2013. Electroconvulsive Therapy Response in Major Depressive Disorder: A Pilot Functional Network Connectivity Resting State fMRI Investigation. Front. Psychiatry 4. https://doi.org/10.3389/fpsyt.2013.00010

Abdallah, C.G., Averill, L.A., Collins, K.A., Geha, P., Schwartz, J., Averill, C., DeWilde, K.E., Wong, E., Anticevic, A., Tang, C.Y., others, 2016. Ketamine treatment and global brain connectivity in major depression. Neuropsychopharmacology.

Alexopoulos, G.S., Hoptman, M.J., Kanellopoulos, D., Murphy, C.F., Lim, K.O., Gunning, F.M., 2012. Functional connectivity in the cognitive control network and the default mode network in late-life depression. J. Affect. Disord. 139, 56–65. https://doi.org/10.1016/j.jad.2011.12.002

American Psychiatric Association, 2013. Diagnostic and statistical manual of mental disorders, 5th ed. American Psychiatric Publishing, Arlington, VA.

Baker, J.T., Holmes, A.J., Masters, G.A., Yeo, B.T.T., Krienen, F., Buckner, R.L., Öngür, D., 2014. Disruption of Cortical Association Networks in Schizophrenia and Psychotic Bipolar Disorder. JAMA Psychiatry 71, 109. https://doi.org/10.1001/jamapsychiatry.2013.3469

Balderston, N.L., Hale, E., Hsiung, A., Torrisi, S., Holroyd, T., Carver, F.W., Coppola, R., Ernst, M., Grillon, C., 2017. Threat of shock increases

excitability and connectivity of the intraparietal sulcus. eLife 6.

https://doi.org/10.7554/eLife.23608

Banks, S.J., Eddy, K.T., Angstadt, M., Nathan, P.J., Phan, K.L., 2007.

Amygdala–frontal connectivity during emotion regulation. Soc. Cogn.

Affect. Neurosci. 2, 303–312. https://doi.org/10.1093/scan/nsm029

Baune, B.T., Miller, R., McAfoose, J., Johnson, M., Quirk, F., Mitchell, D., 2010.

The role of cognitive impairment in general functioning in major

depression. Psychiatry Res. 176, 183–189.

https://doi.org/10.1016/j.psychres.2008.12.001

Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a

practical and powerful approach to multiple testing. J R Stat Soc Ser. B

Stat Methodol 289–300.

Bilker, W.B., Hansen, J.A., Brensinger, C.M., Richard, J., Gur, R.E., Gur, R.C.,

2012. Development of abbreviated nine-item forms of the Raven's

Standard Progressive Matrices Test. Assessment 354–369.

Biswal, B., Zerrin Yetkin, F., Haughton, V.M., Hyde, J.S., 1995. Functional

connectivity in the motor cortex of resting human brain using echo-planar

mri. Magn. Reson. Med. 34, 537–541.

Boehm, I., Geisler, D., King, J.A., Ritschel, F., Seidel, M., Deza Araujo, Y.,

Petermann, J., Lohmeier, H., Weiss, J., Walter, M., Roessner, V., Ehrlich,

S., 2014. Increased resting state functional connectivity in the fronto-

parietal and default mode network in anorexia nervosa. Front. Behav.

Neurosci. 8. https://doi.org/10.3389/fnbeh.2014.00346

Box, G.E., Cox, D.R., 1964. An analysis of transformations. J. R. Stat. Soc. Ser.

B Methodol. 211–252.

Brakowski, J., Spinelli, S., Dorig, N., Bosch, O.G., Manoliu, A., Holtforth, M.G.,

Seifritz, E., 2017. Resting state brain network function in major depression

- Depression symptomatology, antidepressant treatment effects, future

research. J. Psychiatr. Res. 92, 147–159.

https://doi.org/10.1016/j.jpsychires.2017.04.007

Buchanan, A., Wang, X., Gollan, J.K., 2014. Resting-state functional connectivity

in women with Major Depressive Disorder. J. Psychiatr. Res. 59, 38–44.

https://doi.org/10.1016/j.jpsychires.2014.09.002

Buckner, R.L., Sepulcre, J., Talukdar, T., Krienen, F.M., Liu, H., Hedden, T.,

Andrews-Hanna, J.R., Sperling, R.A., Johnson, K.A., 2009. Cortical Hubs

Revealed by Intrinsic Functional Connectivity: Mapping, Assessment of

Stability, and Relation to Alzheimer's Disease. J. Neurosci. 29, 1860–

1873. https://doi.org/10.1523/JNEUROSCI.5062-08.2009

Buhle, J.T., Silvers, J.A., Wager, T.D., Lopez, R., Onyemekwu, C., Kober, H.,

Weber, J., Ochsner, K.N., 2014. Cognitive Reappraisal of Emotion: A

Meta-Analysis of Human Neuroimaging Studies. Cereb. Cortex 24, 2981–

2990. https://doi.org/10.1093/cercor/bht154

Carleton, R.N., Thibodeau, M.A., Teale, M.J.N., Welch, P.G., Abrams, M.P.,

Robinson, T., Asmundson, G.J.G., 2013. The Center for Epidemiologic

Studies Depression Scale: A Review with a Theoretical and Empirical

Examination of Item Content and Factor Structure. PLoS ONE 8, e58067.

https://doi.org/10.1371/journal.pone.0058067

Cattell, R.B., Horn, J.L., 1978. A check on the theory of fluid and crystallized

intelligence with description of new subtest designs. J. Educ. Meas. 15,

139–164.

Centers for Disease Control and Prevention (CDC), 2010. Current depression

among adults---United States, 2006 and 2008. MMWR Morb. Mortal.

Wkly. Rep. 59, 1229–1235.

Chein, J.M., Schneider, W., 2005. Neuroimaging studies of practice-related

change: fMRI and meta-analytic evidence of a domain-general control

network for learning. Cogn. Brain Res. 25, 607–623.

https://doi.org/10.1016/j.cogbrainres.2005.08.013

Cole, J.C., Rabin, A.S., Smith, T.L., Kaufman, A.S., 2004. Development and

Validation of a Rasch-Derived CES-D Short Form. Psychol. Assess. 16,

360–372. https://doi.org/10.1037/1040-3590.16.4.360

Cole, M.W., Anticevic, A., Repovs, G., Barch, D., 2011. Variable Global

Dysconnectivity and Individual Differences in Schizophrenia. Biol.

Psychiatry 70, 43–50. https://doi.org/10.1016/j.biopsych.2011.02.010

Cole, M.W., Pathak, S., Schneider, W., 2010. Identifying the brain's most globally

connected regions. NeuroImage 49, 3132–3148.

https://doi.org/10.1016/j.neuroimage.2009.11.001

Cole, M.W., Repovš, G., Anticevic, A., 2014. The Frontoparietal Control System

A Central Role in Mental Health. The Neuroscientist 20, 652–664.

Cole, M.W., Reynolds, J.R., Power, J.D., Repovs, G., Anticevic, A., Braver, T.S.,

2013. Multi-task connectivity reveals flexible hubs for adaptive task

control. Nat. Neurosci. 16, 1348–1355. https://doi.org/10.1038/nn.3470

Cole, M.W., Schneider, W., 2007. The cognitive control network: Integrated

cortical regions with dissociable functions. NeuroImage 37, 343–360.

https://doi.org/10.1016/j.neuroimage.2007.03.071

Cowdrey, F.A., Filippini, N., Park, R.J., Smith, S.M., McCabe, C., 2014.

Increased resting state functional connectivity in the default mode network

in recovered anorexia nervosa: Resting State Functional Connectivity in

the DMN in Recovered AN. Hum. Brain Mapp. 35, 483–491.

https://doi.org/10.1002/hbm.22202

Dixon, M.L., De La Vega, A., Mills, C., Andrews-Hanna, J., Spreng, R.N., Cole,

M.W., Christoff, K., 2018. Heterogeneity within the frontoparietal control

network and its relationship to the default and dorsal attention networks.

Proc. Natl. Acad. Sci. 201715766.

Dosenbach, N.U., Fair, D.A., Miezin, F.M., Cohen, A.L., Wenger, K.K.,

Dosenbach, R.A., Fox, M.D., Snyder, A.Z., Vincent, J.L., Raichle, M.E.,

others, 2007. Distinct brain networks for adaptive and stable task control

in humans. Proc. Natl. Acad. Sci. 104, 11073–11078.

Drysdale, A.T., Grosenick, L., Downar, J., Dunlop, K., Mansouri, F., Meng, Y.,

Fetcho, R.N., Zebley, B., Oathes, D.J., Etkin, A., Schatzberg, A.F.,

Sudheimer, K., Keller, J., Mayberg, H.S., Gunning, F.M., Alexopoulos,

G.S., Fox, M.D., Pascual-Leone, A., Voss, H.U., Casey, B., Dubin, M.J.,

Liston, C., 2016. Resting-state connectivity biomarkers define

neurophysiological subtypes of depression. Nat. Med. 23, 28–38.

https://doi.org/10.1038/nm.4246

Dumontheil, I., Thompson, R., Duncan, J., 2011. Assembly and use of new task

rules in fronto-parietal cortex. J. Cogn. Neurosci. 23, 168–182.

Duncan, J., 2010. The multiple-demand (MD) system of the primate brain: mental

programs for intelligent behaviour. Trends Cogn. Sci. 14, 172–179.

https://doi.org/10.1016/j.tics.2010.01.004

Fornito, A., Yoon, J., Zalesky, A., Bullmore, E.T., Carter, C.S., 2011. General and

Specific Functional Connectivity Disturbances in First-Episode

Schizophrenia During Cognitive Control Performance. Biol. Psychiatry 70,

64–72. https://doi.org/10.1016/j.biopsych.2011.02.019

Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub,

E., Ugurbil, K., Andersson, J., Beckmann, C.F., Jenkinson, M., Smith,

S.M., Van Essen, D.C., 2016. A multi-modal parcellation of human

cerebral cortex. Nature 536, 171–178. https://doi.org/10.1038/nature18933

Glasser, M.F., Sotiropoulos, S.N., Wilson, J.A., Coalson, T.S., Fischl, B.,

Andersson, J.L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J.R., Van

Essen, D.C., Jenkinson, M., 2013. The minimal preprocessing pipelines

for the Human Connectome Project. NeuroImage 80, 105–124.

https://doi.org/10.1016/j.neuroimage.2013.04.127

Gong, Q., He, Y., 2015. Depression, Neuroimaging and Connectomics: A

Selective Overview. Biol. Psychiatry 77, 223–235.

https://doi.org/10.1016/j.biopsych.2014.08.009

Gordon, E.M., Laumann, T.O., Adeyemo, B., Huckins, J.F., Kelley, W.M.,

Petersen, S.E., 2016. Generation and Evaluation of a Cortical Area

Parcellation from Resting-State Correlations. Cereb. Cortex 26, 288–303.

https://doi.org/10.1093/cercor/bhu239

Greicius, M.D., Flores, B.H., Menon, V., Glover, G.H., Solvason, H.B., Kenna, H.,

Reiss, A.L., Schatzberg, A.F., 2007. Resting-state functional connectivity

in major depression: abnormally increased contributions from subgenual

cingulate cortex and thalamus. Biol. Psychiatry 62, 429–437.

Gress-Smith, J.L., Roubinov, D.S., Andreotti, C., Compas, B.E., Luecken, L.J.,

2015. Prevalence, Severity and Risk Factors for Depressive Symptoms

and Insomnia in College Undergraduates: Depressive Symptoms and

Insomnia in College. Stress Health 31, 63–70.

https://doi.org/10.1002/smi.2509

Hamilton, J.P., Furman, D.J., Chang, C., Thomason, M.E., Dennis, E., Gotlib,

I.H., 2011. Default-Mode and Task-Positive Network Activity in Major

Depressive Disorder: Implications for Adaptive and Maladaptive

Rumination. Biol. Psychiatry 70, 327–333.

https://doi.org/10.1016/j.biopsych.2011.02.003

Herrero, J., Meneses, J., 2006. Short Web-based versions of the perceived

stress (PSS) and Center for Epidemiological Studies-Depression (CESD)

Scales: a comparison to pencil and paper responses among Internet

users. Comput. Hum. Behav. 22, 830–846.

https://doi.org/10.1016/j.chb.2004.03.007

Hwang, J.W., Egorova, N., Yang, X.Q., Zhang, W.Y., Chen, J., Yang, X.Y., Hu,

L.J., Sun, S., Tu, Y., Kong, J., 2015. Subthreshold depression is

associated with impaired resting-state functional connectivity of the

cognitive control network. Transl. Psychiatry 5, e683.

https://doi.org/10.1038/tp.2015.174

Ito, T., Kulkarni, K.R., Schultz, D.H., Mill, R.D., Chen, R.H., Solomyak, L.I., Cole,

M.W., 2017. Cognitive task information is transferred between brain

regions via resting-state network topology. Nat. Commun. 8.

https://doi.org/10.1038/s41467-017-01000-w

Kaiser, R.H., Andrews-Hanna, J.R., Wager, T.D., Pizzagalli, D.A., 2015. Large-

Scale Network Dysfunction in Major Depressive Disorder: A Meta-analysis

of Resting-State Functional Connectivity. JAMA Psychiatry 72, 603.

https://doi.org/10.1001/jamapsychiatry.2015.0071

Kohn, N., Eickhoff, S.B., Scheller, M., Laird, A.R., Fox, P.T., Habel, U., 2014.

Neural network of cognitive emotion regulation — An ALE meta-analysis

and MACM analysis. NeuroImage 87, 345–355.

https://doi.org/10.1016/j.neuroimage.2013.11.001

Li, B., Liu, L., Friston, K.J., Shen, H., Wang, L., Zeng, L.-L., Hu, D., 2013. A

Treatment-Resistant Default Mode Subnetwork in Major Depression. Biol.

Psychiatry 74, 48–54. https://doi.org/10.1016/j.biopsych.2012.11.007

Li, F., He, N., Li, Y., Chen, L., Huang, X., Lui, S., Guo, L., Kemp, G.J., Gong, Q.,

2014. Intrinsic brain abnormalities in attention deficit hyperactivity

disorder: a resting-state functional MR imaging study. Radiology 272,

514–523.

Liang, X., Zou, Q., He, Y., Yang, Y., 2013. Coupling of functional connectivity and

regional cerebral blood flow reveals a physiological basis for network hubs

of the human brain. Proc. Natl. Acad. Sci. 110, 1929–1934.

https://doi.org/10.1073/pnas.1214900110

Manoliu, A., Meng, C., Brandl, F., Doll, A., Tahmasian, M., Scherr, M.,

Schwerthöffer, D., Zimmer, C., Förstl, H., Bäuml, J., Riedl, V.,

Wohlschläger, A.M., Sorg, C., 2014. Insular dysfunction within the

salience network is associated with severity of symptoms and aberrant

inter-network connectivity in major depressive disorder. Front. Hum.

Neurosci. 7. https://doi.org/10.3389/fnhum.2013.00930

Murrough, J.W., Abdallah, C.G., Anticevic, A., Collins, K.A., Geha, P., Averill,

L.A., Schwartz, J., DeWilde, K.E., Averill, C., Jia-Wei Yang, G., Wong, E.,

Tang, C.Y., Krystal, J.H., Iosifescu, D.V., Charney, D.S., 2016. Reduced

global functional connectivity of the medial prefrontal cortex in major

depressive disorder: Global Connectivity in MDD. Hum. Brain Mapp. 37,

3214–3223. https://doi.org/10.1002/hbm.23235

Park, B., Hong, J., Lee, S.-H., Park, H., 2016. Functional Connectivity of Child

and Adolescent Attention Deficit Hyperactivity Disorder Patients:

Correlation with IQ. Front. Hum. Neurosci. 10.

https://doi.org/10.3389/fnhum.2016.00565

Perrin, J.S., Merz, S., Bennett, D.M., Currie, J., Steele, D.J., Reid, I.C.,

Schwarzbauer, C., 2012. Electroconvulsive therapy reduces frontal

cortical connectivity in severe depressive disorder. Proc. Natl. Acad. Sci.

109, 5464–5468. https://doi.org/10.1073/pnas.1117206109

Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012.

Spurious but systematic correlations in functional connectivity MRI

networks arise from subject motion. NeuroImage 59, 2142–2154.

https://doi.org/10.1016/j.neuroimage.2011.10.018

Power, J.D., Cohen, A.L., Nelson, S.M., Wig, G.S., Barnes, K.A., Church, J.A.,

Vogel, A.C., Laumann, T.O., Miezin, F.M., Schlaggar, B.L., Petersen, S.E.,

2011. Functional Network Organization of the Human Brain. Neuron 72,

665–678. https://doi.org/10.1016/j.neuron.2011.09.006

Power, J.D., Schlaggar, B.L., Lessov-Schlaggar, C.N., Petersen, S.E., 2013.

Evidence for Hubs in Human Functional Brain Networks. Neuron 79, 798–

813. https://doi.org/10.1016/j.neuron.2013.07.035

Schneider, W., 2003. Controlled & automatic processing: behavior, theory, and

biological mechanisms. Cogn. Sci. 27, 525–559.

https://doi.org/10.1016/S0364-0213(03)00011-9

Sepulcre, J., Liu, H., Talukdar, T., Martincorena, I., Yeo, B.T.T., Buckner, R.L.,

2010. The Organization of Local and Distant Functional Connectivity in the

Human Brain. PLoS Comput. Biol. 6, e1000808.

https://doi.org/10.1371/journal.pcbi.1000808

Sheline, Y.I., Barch, D.M., Price, J.L., Rundle, M.M., Vaishnavi, S.N., Snyder,

A.Z., Mintun, M.A., Wang, S., Coalson, R.S., Raichle, M.E., 2009. The

default mode network and self-referential processes in depression. Proc.

Natl. Acad. Sci. 106, 1942–1947.

Siegel, J.S., Mitra, A., Laumann, T.O., Seitzman, B.A., Raichle, M., Corbetta, M.,

Snyder, A.Z., 2016. Data Quality Influences Observed Links Between

Functional Connectivity and Behavior. Cereb. Cortex.

https://doi.org/10.1093/cercor/bhw253

Spronk, M., Ji, J.L., Kulkarni, K., Repovs, G., Anticevic, A., Cole, M.W., 2017.

Mapping the human brain's cortical-subcortical functional network

organization. bioRxiv 206292.

Sylvester, C.M., Corbetta, M., Raichle, M.E., Rodebaugh, T.L., Schlaggar, B.L.,

Sheline, Y.I., Zorumski, C.F., Lenze, E.J., 2012. Functional network

dysfunction in anxiety and anxiety disorders. Trends Neurosci. 35, 527–

535. https://doi.org/10.1016/j.tins.2012.04.012

Van Dam, N.T., Earleywine, M., 2011. Validation of the Center for Epidemiologic

Studies Depression Scale—Revised (CESD-R): Pragmatic depression

assessment in the general population. Psychiatry Res. 186, 128–132.

https://doi.org/10.1016/j.psychres.2010.08.018

Van Essen, D.C., Smith, S.M., Barch, D.M., Behrens, T.E.J., Yacoub, E., Ugurbil,

K., 2013. The WU-Minn Human Connectome Project: An overview.

NeuroImage 80, 62–79. https://doi.org/10.1016/j.neuroimage.2013.05.041

Veer, I.M., 2010. Whole brain resting-state analysis reveals decreased functional

connectivity in major depression. Front. Syst. Neurosci. 4.

https://doi.org/10.3389/fnsys.2010.00041

Wang, L., Xia, M., Li, K., Zeng, Y., Su, Y., Dai, W., Zhang, Q., Jin, Z., Mitchell,

P.B., Yu, X., He, Y., Si, T., 2015. The effects of antidepressant treatment

on resting-state functional brain networks in patients with major

depressive disorder: Antidepressant Effects on Depressed Brain

Networks. Hum. Brain Mapp. 36, 768–778.

https://doi.org/10.1002/hbm.22663

Wei, X., Shen, H., Ren, J., Li, X., Xu, X., Yang, R., Lai, L., Chen, L., Hu, J., Liu,

W., Jiang, X., 2014. Altered Resting-State Connectivity in College

Students with Nonclinical Depressive Symptoms. PLoS ONE 9, e114603.

https://doi.org/10.1371/journal.pone.0114603

Wig, G.S., Schlaggar, B.L., Petersen, S.E., 2011. Concepts and principles in the

analysis of brain networks: Brain networks. Ann. N. Y. Acad. Sci. 1224,

126–146. https://doi.org/10.1111/j.1749-6632.2010.05947.x

World Health Organization, 2017. Depression and other common mental

disorders: Global health estimates. World Health Organization, Geneva.

World Health Organization, 1992. The ICD-10 classifications of mental and

behavioural disorder: Clinical descriptions and diagnostic guidelines.

World Health Organization, Geneva.

Yang, G.J., Murray, J.D., Wang, X.-J., Glahn, D.C., Pearlson, G.D., Repovs, G.,

Krystal, J.H., Anticevic, A., 2016. Functional hierarchy underlies

preferential connectivity disturbances in schizophrenia. Proc. Natl. Acad.

Sci. 113, E219–E228. https://doi.org/10.1073/pnas.1508436113

Yeo, B.T.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D.,

Hollinshead, M., Roffman, J.L., Smoller, J.W., Zollei, L., Polimeni, J.R.,

Fischl, B., Liu, H., Buckner, R.L., 2011. The organization of the human

cerebral cortex estimated by intrinsic functional connectivity. J.

Neurophysiol. 106, 1125–1165. https://doi.org/10.1152/jn.00338.2011

Figure 1 - Network assignment. Resting-state fMRI data from an independent

dataset (HCP: 100 unrelated) was used to assign each parcel to a network using

a community detection algorithm. This resulted in 12 total networks. Color

indicates the network assignment for each parcel.

Figure 2 – Histogram of raw CESD scores.

Figure 3 – Between network global connectivity (BGC) across all cortical

regions. Between network global connectivity for region A is defined as the

mean FC (Pearson correlation) between that region A and all other regions

outside of region A's network. Warm values indicate positive values and cool

values indicate negative values.

Figure 4 – CESD scores and BGC in the FPN, DMN, and language networks

are negatively correlated. A) BGC in the FPN is plotted on the y-axis and

CESD scores are plotted on the x-axis. B) BGC in the DMN is plotted on the y-

axis and CESD scores are plotted on the x-axis. C) BGC in the language network

is plotted on the y-axis and CESD scores are plotted on the x-axis.

936