1
14   Global connectivity of the frontoparietal cognitive control network is related to
15   depression symptoms in the general population
16

17   Douglas H. Schultz[1], Takuya Ito[1], Levi I. Solomyak[1], Richard H. Chen[1], Ravi D.
18   Mill[1], Alan Anticevic[2], & Michael W. Cole[1]
19
20   [1]Center for Molecular and Behavioral Neuroscience,
21   Rutgers University – Newark
22   [2]Department of Psychiatry,
23   Yale University

29   Corresponding author: Douglas H. Schultz
30   Center for Molecular and Behavioral Neuroscience
31   197 University Ave, Suite 212
32   Newark, NJ 07102
33   dhs95@newark.rutgers.edu

34
35
36

37   KEYWORDS: fMRI, depression, frontoparietal network, functional connectivity

38
39
40

ABSTRACT

We all vary in our mental health, even among people not meeting diagnostic criteria for mental illness. Understanding this individual variability may reveal factors driving the risk for mental illness, as well as factors driving sub-clinical problems that still adversely affect quality of life. To better understand the large-scale brain network mechanisms underlying this variability we examined the relationship between mental health symptoms and resting-state functional connectivity patterns in cognitive control systems. One such system is the frontoparietal cognitive control network (FPN). Changes in FPN connectivity may impact mental health by disrupting the ability to regulate symptoms in a goal-directed manner. Here we test the hypothesis that FPN dysconnectivity relates to mental health symptoms even among individuals who do not meet formal diagnostic criteria but may exhibit meaningful symptom variation. We found that depression symptoms severity negatively correlated with between-network global connectivity (BGC) of the FPN. This suggests that decreased connectivity between the FPN and the rest of the brain is related to increased depression symptoms in the general population. These findings complement previous clinical studies to support the hypothesis that global FPN connectivity contributes to the regulation of mental health symptoms across both health and disease.

AUTHOR SUMMARY

Understanding how large-scale network interactions in the brain contribute to (or serve a protective role against) mental health symptoms is an important step toward developing more effective mental health treatments. Here we test the hypothesis that cognitive control networks play an important role in mental health by being highly connected to other brain networks and able to serve as a feedback mechanism capable of regulating symptoms in a goal-directed manner. We found that the more well-connected the frontoparietal cognitive control network was to other networks in the brain the less depression symptoms were reported by participants. These results contribute to our understanding of how brain network interactions are related to mental health symptoms, even in individuals who have not been diagnosed with a disorder.

INTRODUCTION

People vary in their degree of mental health. Indeed, people who do not meet formal criteria for mental illness as defined by our current diagnostic systems (American Psychiatric Association, 2013; World Health Organization, 1992) may still experience a number of symptoms associated with that disorder (World Health Organization, 2017). Here we use this natural variability to better understand neural factors potentially contributing to day-to-day experiences of poor mental health, as well as (prodromal) factors that may elevate the risk for severe mental illness. We hypothesized that the variability observed in mental health symptoms among individuals is related to the function of the frontoparietal cognitive control network (FPN), based on our previously-developed theoretical proposal that the FPN plays a domain-general protective role against mental health symptoms (Cole et al., 2014).

The proposed theoretical framework suggests that alterations in FPN function may play a common role in multiple mental disorders by disrupting a domain-general cognitive control feedback mechanism that can regulate symptoms when they are experienced (Cole et al., 2014). The FPN is a candidate network for this function because it is a flexible hub, meaning it has a high degree of connectivity across the brain (Cole et al., 2010; Power et al., 2011) and can rapidly modify functional connections according to current goals (Cole et al., 2013). There is strong evidence that these FPN functions are domain general (Chein and Schneider, 2005; Cole et al., 2013; Dosenbach et al., 2007), such that individual differences in the general ability to regulate cognition can influence symptoms. Finally, alterations in FPN functional connectivity (FC) have been identified in a number of mental disorders including: depression (Kaiser et al., 2015), anxiety (Sylvester et al., 2012), schizophrenia (Baker et al., 2014; Cole et al., 2011; Fornito et al., 2011; Yang et al., 2016), attention deficit hyperactivity disorder (Li et al., 2014; Park et al., 2016), and eating disorders (Boehm et al., 2014; Cowdrey et al., 2014). Consistent with most of these studies, we focus here on FC measured using functional magnetic resonance imaging (fMRI), calculated as the temporal relationship in the blood oxygenation level dependent (BOLD) signal between brain regions (Biswal et al., 1995) while participants rest in the scanner.

Consistent with the flexible hub framework, a number of studies using different measures have provided converging evidence that the FPN is especially well connected to the rest of the brain (Buckner et al., 2009; Cole et al., 2010). Both of these studies calculated a summary statistic reflecting the degree of connectivity across the whole brain. However, these estimates can be influenced by the relative size of different networks. For example, nodes of a larger network will have a larger overall number of strong connections than nodes of a smaller network simply because, by definition, within-network connections are stronger on average than between-network connections (Power et al., 2013; Wig et al., 2011). Therefore, we estimated how well each region of the brain was connected to the rest of the brain using between-network global connectivity (BGC) (Ito et al., 2017), a measure not influenced by network size.

Particularly important for our specific test of the flexible hub framework
here, patients diagnosed with major depression exhibit differences in FC patterns
throughout the brain, including FPN functional connections (Brakowski et al.,
2017). Specifically, connectivity between regions of the FPN is decreased in
depressed individuals (Alexopoulos et al., 2012), as well as in undiagnosed
individuals experiencing depression symptoms (Wei et al., 2014). However, Wei
and colleagues (2014) looked at FC with specific seed regions, not global
connectivity, in their sample. Another study found that global brain connectivity
was decreased in the medial prefrontal cortex and the dorsolateral prefrontal
cortex (dlPFC) portions of the FPN in depressed patients (Murrough et al., 2016).
Decreases in within-network FPN connectivity have also been observed in a
group of individuals reporting depression symptoms in the absence of a clinical
diagnosis (Hwang et al., 2015). Researchers have also attempted to subdivide
depression into various types based on FC patterns. Decreases in FPN
connectivity were most pronounced in one particular subtype of depression
associated with increased symptoms of fatigue and decreased symptoms of
anxiety (Drysdale et al., 2016). These previous results are broadly consistent
with our hypothesis, yet the extension of results to test whether FPN BGC is
related to mental health symptoms among healthy individuals would provide
important new evidence for the general nature of FPN's role in regulating mental
health.
Consistent with our previously-developed theoretical framework (Cole et
al., 2014) along with observed FPN FC alterations in patients with major
depression, we hypothesized that individual differences in depression symptoms
in undiagnosed individuals would be correlated with BGC in the FPN. Support for
our hypothesis would provide important evidence for a potentially general role of
global FPN intrinsic FC in facilitating the regulation of mental health symptoms.
148
METHODS
Participants
Data were collected at the Rutgers University Brain Imaging Center
(RUBIC). The participants were recruited from the Rutgers University-Newark
campus and surrounding community. All participants provided informed consent
and all procedures were approved by the Rutgers University-Newark Institutional
Review Board. We collected data from 106 participants. Technical error or
equipment malfunction during the scanning session resulted in removing six
participants from the study. Four participants were removed from the study
because they did not complete the Center for Epidemiological Studies
Depression Scale (CESD) during a behavior-only session separate from the MRI
session. We also collected demographic information and asked participants 11
questions asking what hand they used for various activities including writing,
throwing, using a scissors, holding their toothbrush, striking a match, opening a
box, kicking, using a knife, using a spoon, which hand was placed on top while
using a broom, and which eye they used in situations where they would only
being using one eye. They replied to each question by answering always right,
usually right, no preference, usually left, or always left. Answers were scored in

the following manner: always right (2), usually right (1), no preference (0), usually
left (-1), always left (-2). We calculated a laterality quotient (LQ) by summing
these scores and dividing by the maximum score of 22. A LQ score of -100
indicates a strong left hand preference, 0 indicates no hand preference, and 100
indicates a strong right hand preference.
Studies have proposed that the CESD can be broken down into between
one and four factors (Cole et al., 2004; Herrero and Meneses, 2006). In addition
to calculating the raw CESD score for each participant we also calculated three
factor scores: somatic symptoms, negative affect, and anhedonia, based on a
recent study (Carleton et al., 2013). Participants also completed several
measures of flexible cognition during the behavior-only session. These measures
included Raven's progressive matrices (Bilker et al., 2012), Cattell's culture fair
test (Cattell and Horn, 1978), and Duncan's goal neglect task (Dumontheil et al.,
2011). The final sample consisted of 96 participants (See Table 1).
181

Table 1. Demographic information

N	96
Age	*M* = 22.06, *SD* = 3.84
Gender	
Male	42 (43.8%)
Female	54 (56.2%)
Handedness (LQ)	*M* = 74.67, *SD* = 25.2
Education (Highest level)	
High school	14 (14.6%)
Some college	50 (52.1%)
Bachelor's degree	29 (30.2%)
Graduate degree	3 (3.1%)
Cognitive measures	(Percent correct)
Raven	*M* = 52.29, *SD* = 15.69
Cattell	*M* = 55.20, *SD* = 12.03
Duncan	*M* = 76.19, *SD* = 18.34
CESD (raw score)	*M* = 17.41, *SD* = 9.26

182
183

MRI Parameters
Multiband whole-brain echo-planar imaging (EPI) acquisition was collected
using a 32-channel head coil on a 3T Siemens Trio MRI scanner with the
following parameters: TR = 785 ms, TE = 34.8 ms, flip angle = 55°, Bandwidth
1924/Hz/Px, in-plane FoV read = 208 mm, 72 slices, 2.0 mm isotropic voxels,
with a multiband acceleration factor of 8. Whole-brain high-resolution T1-
weighted and T2-weighted anatomical scans with 0.8 mm isotropic voxels were
also collected. Spin echo field maps were collected in both the anterior to
posterior direction and the posterior to anterior direction consistent with the
Human Connectome Project preprocessing pipelines (Glasser et al., 2013). The
resting-state fMRI scan was 14 minutes (1070 TRs) in duration.
195

196
fMRI Preprocessing

Functional imaging data were preprocessed using the Human Connectome Project minimal preprocessing pipeline version 3.5.0. Preprocessing consisted of anatomical restructuring and segmentation, EPI reconstruction, segmentation, and spatial normalization to a standard template, intensity normalization, and motion correction (Glasser et al., 2013). All further processing was conducted in CIFTI 64k greyordinate standard space. The data were parcellated into 360 regions described previously (Glasser et al., 2016), taking the average time series of the vertices within a parcel. At this point all subsequent data analysis was conducted in MATLAB R2014b (The Mathworks). We performed nuisance regression using 12 motion parameters, ventricle and white matter timeseries (as well as their first derivatives), and global signal. We also performed motion scrubbing (Power et al., 2012) based on framewise displacement. Framewise displacement was calculated as the amount of head movement for each frame relative to the previous in terms of Euclidean distance. We next applied a low pass temporal filter (0.3 Hz) to the framewise displacement vector in order to reduce the effect of respiration on our framewise displacement measure (Siegel et al., 2016). The framewise displacement threshold for motion scrubbing was set at 0.3 mm. Motion scrubbing consisted of removing the flagged frame from the timeseries as well as one frame prior and the two frames following. FC was estimated by calculating the Pearson correlation of the BOLD timeseries between each pair of the parcels defined by Glasser and colleagues (Glasser et al., 2016).

Network Assignment and Analysis

The network assignment of each of the parcels was completed on an independent dataset, the Human Connectome Project (100 unrelated) (Van Essen et al., 2013). Briefly, each parcel was assigned to a network using the Generalized Louvain method for community detection. This process was conducted using resting-state data. We identified 12 functional networks (Spronk et al., 2017) (See Figure 1). The functional network topology findings replicate the major features of several previously published network partitions (Gordon et al., 2016; Power et al., 2011; Yeo et al., 2011).

Cortical partition

Visual 1
Visual 2
Somatomotor
Cingulo-Opercular
Dorsal Attention
Language
Frontoparietal
Auditory
Default Mode
Posterior Multimodal
Ventral Multimodal
Orbito-Affective

231
232 **Figure 1 - Network assignment.** Resting-state fMRI data from an independent dataset (HCP:
233 100 unrelated) was used to assign each parcel to a network using a community detection
234 algorithm. This resulted in 12 total networks. Color indicates the network assignment for each
235 parcel.
236
We were interested in the relationship between the frequency at which
participants experienced depression symptoms and the degree of between-
network global connectivity (BGC). Specifically, we were interested in a measure
that would estimate the strength of FC for each region of a network to all of the
other regions in different networks. BGC was calculated for each region
individually and defined as the mean FC for all out-of-network connections. Out-
of-network connections were defined as all connections from a source region to
target regions outside the source region's network. This process was completed
for all regions until we had a BGC value for each region in the brain. Then we
calculated the mean BGC value for each of the functional networks to summarize
effects at the network level.
More formally, BGC was defined for each region as:

$$BGC_i = \frac{\sum_{j \notin C} W_{ij}}{N_{total} - N_C}$$

where BGC_i corresponds to the out-of-network weighted degree of region i in
network C, $j \notin C$ corresponds to all regions not in network C, W_{ij} corresponds to
the FC estimate between regions i and j, N_{total} corresponds to the total number
of regions, and N_C corresponds to the total number of regions in network C.
We tested the relationship between BGC and depression symptoms in all
brain networks. Due to the large number of statistical tests we report the false-
discovery rate (FDR)-corrected *p*-values for primary analyses (Benjamini and
Hochberg, 1995). Our main hypothesis was that BGC in the FPN was correlated
with depression symptoms. Because this was a primary hypothesis based on our
previous publication (Cole et al., 2014) we report the uncorrected p-value.

Follow-up and control analyses were not independent of the primary tests and
are therefore not corrected.
261
262 **RESULTS**
263
Depression Symptom Scores
We hypothesized that the frequency of experiencing depression
symptoms is related to individual differences in the between-network global
connectivity of the FPN. This would be consistent with the involvement of FPN in
domain-general cognitive regulation, broadly construed to include emotion (Cole
et al., 2014). We used a standard measure of depression symptoms, the CESD,
to measure the frequency of depression symptoms. The CESD consists of 20
questions asking how frequently participants have experienced symptoms related
to depression over the last seven days. CESD scores ($M = 17.41, SD = 9.26$)
varied from a minimum of 0 to a maximum of 43 (out of a possible 60). The
characteristics of our sample were consistent with previous studies using the
CESD with undiagnosed young adults (Gress-Smith et al., 2015; Van Dam and
Earleywine, 2011). However, the scores were not normally distributed based on a
Kolomogorov–Smirnov test ($p < 0.023$). We therefore used the Box-Cox power
transformation (Box and Cox, 1964), a common approach to correct for non-
normality. After the transformation the data no longer deviated from a normal
distribution ($p = 0.17$). The transformed data were used for all subsequent
analyses. CESD scores were not correlated with any of the flexible cognition
measures (*smallest p = 0.28*).
We also calculated CESD sub-scores for somatic, negative affect, and
anhedonia factors. After Box-Cox transformation none of the factor score
distributions significantly deviated from a normal distribution (*somatic p = 0.4,*
negative affect p = 0.19, anhedonia p = 0.056). All three of the factors were
significantly correlated with the overall CESD scores (*somatic r = 0.77,*
uncorrected p < 0.001; negative affect r = 0.80, uncorrected p< 0.001; anhedonia
r = 0.52, uncorrected p < 0.001, See Table 2). The three factors were also
correlated with each other (*correlation between somatic and negative affect r =*
0.55, correlation between somatic and anhedonia r = 0.18, correlation between
negative affect and anhedonia r = 0.26, See Table 1). Although all of the factors
were highly related to the overall CESD score, they appear to have contributed
unique variance as the highest correlation between any two factors was only
accounting for 30% of the variance (based on R^2 values).

Table 2. Correlation between the CESD and CESD factors

	Overall CESD	Somatic	Negative Affect	Anhedonia
Overall CESD	-----	-----	-----	-----
Somatic	0.77	-----	-----	-----
Negative Affect	0.80	0.55	-----	-----
Anhedonia	0.52	0.18	0.26	-----

305
306
Between-Network Global Connectivity
We calculated BGC values using resting-state fMRI data for each
functionally defined cortical parcel (Figure 2). DMN regions tended to have lower
BGC values, consistent with the conceptually-similar participation coefficient
results reported by Power et al. (2011). In contrast, the FPN had high levels of
BGC in lateral prefrontal portions of the network (*M = 0.002, SD = 0.02*) relative
to the mean of regions outside of the FPN (*M = -0.0081, SD = 0.013; t(95) =*
5.08, p < 0.0001). However, FPN had lower BGC values in parietal (*M = -0.0258,*
SD = 0.0190), inferior frontal (*M = -0.0177, SD = 0.0209*) and temporal lobe
regions of the network (*M = -0.0420, SD = 0.0221*) relative to the mean of
regions outside of the FPN (*largest p < 0.0001*). Sensory networks exhibited two
patterns. The visual network had some of the lowest BGC values in the brain with
the primary visual cortex region showing the least BGC in the network (*M = -*
0.0742, SD = 0.0348). In contrast, the auditory network (*M = 0.0212, SD =*
0.0171) and portions of the sensorimotor network surrounding the central sulcus
(*M = 0.0117, SD = 0.0249*) both showed a high degree of BGC. The
sensorimotor regions closest to the central sulcus showed more moderate BGC
scores (*M = -0.0139, SD = 0.0289*).
325

Figure 2 – Between network global connectivity (BGC) across all cortical regions. Between network global connectivity for region A is defined as the mean FC (Pearson correlation) between that region A and all other regions outside of region A's network. Warm values indicate positive values and cool values indicate negative values.

327
328
Correlation Between BGC and Depression Symptoms
We next tested for a relationship between BGC and depression
symptoms. This involved creating a mean BGC score for each functional network
and testing for correlation between those values and the depression symptom
measure (See Table 3). Consistent with our a priori hypothesis, we found that
FPN BGC was significantly correlated with depression symptoms (r = *-0.247, p =*
0.015) (Figure 3A). Unexpectedly, we found that DMN and language network
BGC showed a similar magnitude effect (DMN r = *-0.241, uncorrected p = 0.018;*
language r = *-0.268, uncorrected p = 0.008*) (Figure 3B & 3C). These findings
suggest that greater FC between the FPN and the rest of the brain (outside of the
FPN) is related to less frequent depression symptoms. Less depression
symptoms are also related to greater BGC in the DMN and language networks.
We report an uncorrected p-value for the test of our a priori hypothesis that FPN
BGC would be related to depression symptoms. We also examined the
relationship between depression symptoms and BGC in the other 11 brain
networks. These p-values do not survive FDR correction for multiple
comparisons, but because the magnitude of the effect observed in the DMN and
language network was similar to that in the FPN we continue to report
subsequent uncorrected statistics for these networks.
To address whether these findings were specific to this particular network
partition we conducted a similar analysis using a different network partition
scheme (Ito et al., 2017). This network partition includes a FPN and DMN, but
lacks a network analogous to the language network. Using this network partition
we found a similar negative correlation between BGC and depression symptoms
in the FPN (r = *-0.262, p = 0.009, FDR adjusted for multiple comparisons p =*

0.044) and DMN (*r = -0.250, p = 0.014, FDR adjusted p = 0.049*). The results for
the FPN and DMN both survive correction for multiple comparisons with this
network partition.
357
358

Table 3. Correlation between BGC and CESD scores

Network	Pearson's r	uncorrected p-value
Visual 1	0.015	0.884
Visual 2	0.020	0.850
Somatomotor	-0.177	0.085
Cingulo-Opercular	-0.143	0.164
Language	-0.268	0.008
Default Mode	-0.241	0.018
Frontoparietal	**-0.247**	**0.015***
Auditory	-0.099	0.340
Post. Multimodal	-0.110	0.284
Dorsal Attention	-0.113	0.275
Ventral Multimodal	0.116	0.261
Orbito-Affective	-0.078	0.448

359
360

Figure 3 – CESD scores and BGC in the FPN, DMN, and language networks are negatively correlated. A) BGC in the FPN is plotted on the y-axis and CESD scores are plotted on the x-axis. B) BGC in the DMN is plotted on the y-axis and CESD scores are plotted on the x-axis. C) BGC in the language network is plotted on the y-axis and CESD scores are plotted on the x-axis.

362
363
In order to verify that these correlations were not dependent on the Box-
Cox transformation we used to normalize the depression symptom data we also
ran non-parametric Spearman's rank correlations, which do not require normally
distributed data. The results of testing our a priori hypothesis that BGC in the
FPN would be related to depression symptoms were unchanged. Specifically, the

BGC-depression rank correlation was consistent with significant Pearson
correlations (FPN *rho = -0.227, p = 0.026*). The Spearman rank correlations also
reflected similar results to the Pearson correlation in the DMN (*rho = -0.220,*
uncorrected p = 0.031) and the language network (*rho = -0.285, uncorrected p =*
0.005).
To ensure that our observed correlation between depression symptoms
and BGC in the FPN was not being driven by other factors we removed the
variance in the CESD scores that could be accounted for by age and gender
factors with a regression model and used the residuals to rerun the correlation.
We still found a significant relationship between BGC in the FPN and depression
symptoms (*r = -0.220, p = 0.031*). This result suggests that the relationship
between BGC in the FPN and depression scores is not being influenced by age
or gender differences in our sample.
Next, we tested if the relationships between BGC and overall depression
scores were driven by specific depression factors. We tested for correlation
between the BGC values and three factors derived from the CESD questionnaire:
somatic symptoms, negative affect, and anhedonia symptoms (Carleton et al.,
2013). In the FPN, BGC was only significantly correlated with anhedonia
symptoms (*r = -0.253, p = 0.013*). FPN BGC was not correlated with somatic
symptoms (*r = -0.149, p = 0.149*) or negative affect (*r = -0.137, p = 0.183*). In the
DMN, BGC was only marginally significantly correlated with somatic symptoms (*r*
= -0.189, uncorrected p = 0.064) and anhedonia symptoms (*r = -0.192,*
uncorrected p = 0.060), and not significantly correlated with negative affect (*r = -*
0.119, uncorrected p = 0.247). In the language network, BGC was correlated with
somatic symptoms (*r = -0.257, uncorrected p = 0.011*) and negative affect (*r = -*
0.227, uncorrected p = 0.026), but it was not significantly correlated with
anhedonia (*r = -0.167, uncorrected p = 0.105*).
396
Observed BGC-depression Correlations were not Dependent on
Connections Between FPN, DMN, and language network
The magnitude and direction of the BGC-depression correlations were
similar for the FPN, DMN, and language network. One possible explanation for
these results is that the observed BGC-depression effects for these networks
were driven primarily by the FC between them. We tested this possibility by
recalculating BGC for each network by removing connections between the FPN,
DMN, and language network from the calculation. This approach minimizes the
possibility that connections between these networks drove the original
correlations between depression symptoms and BGC The magnitude and
direction of the correlations between depression symptoms and BGC in the FPN
(*r = -0.226, p = 0.027*), the DMN (*r = -0.236, uncorrected p = 0.021*), and the
language network (*r = -0.201, uncorrected p = 0.049*) were similar when we
excluded the connections between these networks from the BGC calculation.
This suggests the original BGC-depression correlations were not dependent on
connections between the FPN, DMN, and language network.
In order to further explore the possibility that the observed relationship
between depression symptoms and BGC in the FPN might be primarily driven by

FPN connections to the DMN and language network we calculated the correlation between the mean FPN to DMN, and FPN to language network FC with depression symptoms. There was not a significant correlation between depression symptoms and the mean connectivity between the FPN and DMN ($r = -0.016$, uncorrected $p = 0.881$) or the mean connectivity between the FPN and language network ($r = -0.146$, uncorrected $p = 0.156$). This suggests that the relationship between FPN BGC and depression symptoms is not driven by connections to the DMN and language networks, which both showed a negative correlation between BGC and depression symptoms as well.

Within-Network Connectivity does not Correlate with Depression Symptoms

We focused on between-network connectivity, yet the degree of connectivity within a functional network might also be related to depression symptoms. For example, a high degree of FC within a network may reflect more homogeneous network activity. In this case the network would be tightly coupled and all of the component regions would likely be serving a very similar function. However, a lower degree of FC within a network may reflect more heterogeneous processing. In this case the different regions of a functional network may all be involved in the same general process, but they may be contributing in different ways. These differences in within-network functionality may, in turn, relate to depression symptoms.

Consistent with our choice to focus primarily on between-network effects, within-network FC in the FPN was not significantly correlated with depression symptoms ($r = 0.066$, $p = 0.526$). Within-network FC in the DMN was also not significantly correlated with depression symptoms ($r = 0.143$, uncorrected $p = 0.164$). The language network showed a negative correlation between depression symptoms and within-network FC ($r = -0.219$, uncorrected $p = 0.032$). We observed a significant difference in the DMN correlation coefficients comparing BGC ($r = -0.241$) and within-network connectivity ($r = 0.143$) with depression symptoms ($z = 2.66$, $p = 0.008$). It will be important for future studies to test this effect for replication, given the non-significance of the DMN within-network FC-depression effect relative to a correlation of 0.

DISCUSSION

Based on convergent evidence across a variety of mental health conditions we predicted that individual differences in FPN BGC would be correlated with symptoms associated with depression (Cole et al., 2014). This is consistent with extensive evidence that the FPN is a domain-general cognitive control system regulating general goal pursuit processes (Cole and Schneider, 2007; Duncan, 2010; Schneider, 2003), including regulation of mental illness symptoms (Cole et al., 2014). We sought to expand the general relevance of this framework by including symptoms experienced in everyday life, which express dimensionally across health and disease. This involved investigating the relationship between the mood symptoms, one of the most commonly experienced set of symptoms in the general population (Centers for Disease

Control and Prevention (CDC), 2010), and FPN global connectivity properties.
We further ensured the broad relevance of the findings by focusing on how a
general population who had not been diagnosed with a mental disorder varied in
their depression symptoms and primarily experienced them in the normal (i.e.,
sub-clinical) range.
As predicted, we found a significant relationship between how well-
connected the FPN was to the rest of the brain and the frequency of depression
symptoms in adults who had not been diagnosed with depression. We used BGC
to estimate the degree of brain-wide connectivity for each brain region. The
results suggest that individuals who report fewer depression symptoms are
characterized by a FPN that is more connected to the rest of the brain. These
results support our previously developed theoretical framework, suggesting
natural variance in FPN global connectivity influences each individual's ability to
regulate mood symptoms in everyday life. Extending this framework to test if
FPN function relates to natural variation in other symptom domains will be critical
as well as examining if such effects persist when an individual crosses a
threshold necessary for a formal diagnosis.
478
Between-network global connectivity identifies how well each brain region
is connected to other brain networks
We hypothesized that individuals exhibiting greater FC between the FPN
and the rest of the brain would report fewer depression symptoms. We used
BGC, a measure that calculates the mean FC between each brain region and all
other out-of-network brain regions (Ito et al., 2017), to evaluate this hypothesis.
BGC reduces the potential bias of other graph centrality measures, which can be
inflated in regions assigned to large networks (Power et al., 2013). Previous
methods using global brain connectivity and degree, which both include within
network connections, result in DMN regions showing high connectivity with the
rest of the brain (Buckner et al., 2009; Cole et al., 2010; Liang et al., 2013).
However, we found that BGC was relatively low in the DMN. This is consistent
with DMN results from studies using participation coefficient (the proportion of
between-network vs. within-network connectivity) (Power et al., 2011, 2013).
These results suggest that previous findings identifying the DMN as highly
connected to the rest of the brain are largely driven by high FC within the DMN,
and do not necessarily reflect greater connectivity between the DMN and nodes
in other functional networks.
The results we observed with BGC were similar to previous attempts to
identify hubs and the most well connected regions in the brain. BGC was high in
the lateral prefrontal cortex, the motor and tactile cortex, the auditory cortex, and
higher order visual regions. Lateral prefrontal regions and higher order visual
areas show a greater degree of FC (Buckner et al., 2009) and a higher global
brain connectivity (Cole et al., 2010). Higher global brain connectivity has also
been reported in the lateral prefrontal cortex, higher order visual regions, auditory
cortex, and somatosensory cortex (Liang et al., 2013). We observed that BGC
was consistent with previous attempts to classify the degree of connectivity in
many brain regions.

There were some discrepancies between previous methods and BGC. We
found that BGC was quite low in lower visual regions in contrast to higher
connectivity estimates calculated by others (Cole et al., 2010; Liang et al., 2013).
The differences observed between BGC and other measures of connectivity
strength may be driven by BGC not considering the relatively strong local
connections within the visual network. In fact, primary and secondary visual
cortex show high local connectivity relative to distant connectivity strength
(Sepulcre et al., 2010). It will be important for future studies to identify whether
differences observed in primary sensory cortices are driven by an imbalance
between within and between-network connectivity.
517
BGC in the FPN is negatively correlated with depression symptoms
We found that BGC in the FPN showed a significant negative correlation
with depression symptoms. This suggests that individuals exhibiting greater
connectivity of the FPN with the rest of the brain experience fewer symptoms of
depression. The correlation between BGC in the FPN and depression symptoms
supports our hypothesis that a well-connected FPN may serve a protective role
against depression symptoms and possibly mental health symptoms in general
(Cole et al., 2014).
Decreases in FPN FC have been reported in individuals diagnosed with
major depression (Alexopoulos et al., 2012; Murrough et al., 2016; Veer, 2010).
Differences in FC patterns can also be used to divide depression into distinct
subtypes (Drysdale et al., 2016). We build on these findings by observing similar
results in a variable sample of undiagnosed individuals that likely includes
primarily mentally healthy, but also some mentally unhealthy, participants.
Another study examining FC in a group of undiagnosed individuals with higher
levels of depression symptoms found that connectivity between the superior
parietal lobule and the dlPFC portion of the FPN was decreased (Wei et al.,
2014). Undiagnosed participants experiencing more depression symptoms have
also been reported to have reduced FC between dlPFC and the supramarginal
gyrus, insula, operculum, precuneus and parahippocampal gyrus (Hwang et al.,
2015). Our findings suggest that greater global connectivity of the FPN is related
to reduced depression symptoms. Our data regarding global connectivity
differences is consistent with a recent study that found reduced global brain
connectivity in the lateral PFC in depression patients (Abdallah et al., 2016).
Unexpectedly, we also found a negative correlation between BGC in the
DMN and depression symptoms. This result did not survive multiple comparison
correction, but because it was of a similar magnitude as our predicted
relationship in the FPN we have explored it. It is important to note that the results
in the DMN should be interpreted with caution and future studies should attempt
to replicate these results. DMN activity has been linked to rumination symptoms
in depression (Hamilton et al., 2011). This finding may reflect an inability for
individuals experiencing more depression symptoms to disengage the DMN in
situations when attentional or cognitive resources need to be allocated (Sheline
et al., 2009). Depression patients also exhibit decreased FC between the DMN
and executive networks (Abbott et al., 2013; Manoliu et al., 2014). These findings

provide further support for the interpretation that depressed individuals may have
a difficult time disengaging the DMN because the connections from cognitive
control networks to the DMN are decreased.
A number of studies have suggested that FC within portions of the DMN is
increased in depression patients (Greicius et al., 2007; Li et al., 2013). Although
we did not find a significant correlation between within-network FC in the DMN
and depression symptoms, we did see a trend toward a positive correlation. Our
lack of a correlation between within-network FC in the DMN and depression
symptoms could be due to methodological differences in using a population of
individuals who have not been diagnosed with a disorder versus comparing a
control group to a group diagnosed with depression. Additionally, we considered
the entire DMN rather than using a seed correlation approach. It will be important
for future studies to investigate the relationship between within-DMN FC and
depression symptoms, verifying this effect in a larger sample and tying it to
specific neural mechanisms and specific depression symptoms.
We also found a negative correlation between BGC in the language
network and depression symptoms. Similar to the DMN, this result did not survive
multiple comparison correction, but because the magnitude of the effect was
similar to that of our predicted relationship between BGC in the FPN and
depression symptoms we have explored it. Because we did not have an a priori
prediction about this network and the result did not survive multiple comparison
correction, the interpretation of this result should be tempered. Decreases in FC
in the language network have been reported in depression patients (Buchanan et
al., 2014) and language performance deficits have also been reported in
depression patients (Baune et al., 2010). The current results might be consistent
with these observed deficits in language in depressed individuals. However,
future studies should attempt to replicate these results in an independent large
sample.
581
Conclusion
The current study sought to test our previously-developed framework that
suggests the FPN (along with other cognitive control networks) acts as a
protective factor against mental disease via its widespread FC with other
networks (Cole et al., 2014). We identified a negative correlation between
depression symptoms and a measure of between-network, global FC in the FPN
in a sample of individuals from the general population who had not been
diagnosed with depression. These results suggest the human brain's global
network architecture is critical for maintaining mental health even in undiagnosed
individuals, supporting the possibility that FPN maintains a goal-directed
feedback loop to regulate symptoms as they arise. It will be important for future
research to characterize the exact mechanisms by which FPN influences
symptoms, and to assess the possibility of enhancing FPN FC in the interest of
reducing symptoms and potentially preventing the onset of mental illness.
596
597
598

599 **Acknowledgements**
600   This work was supported by the National Institute of Health (MH096801,
601 MH109520, and AG055556). The content is solely the responsibility of the
602 authors and does not necessarily represent the official views of any of the
603 funding agencies.

References

Abbott, C.C., Lemke, N.T., Gopal, S., Thoma, R.J., Bustillo, J., Calhoun, V.D., Turner, J.A., 2013. Electroconvulsive Therapy Response in Major Depressive Disorder: A Pilot Functional Network Connectivity Resting State fMRI Investigation. Front. Psychiatry 4. https://doi.org/10.3389/fpsyt.2013.00010

Abdallah, C.G., Averill, L.A., Collins, K.A., Geha, P., Schwartz, J., Averill, C., DeWilde, K.E., Wong, E., Anticevic, A., Tang, C.Y., others, 2016. Ketamine treatment and global brain connectivity in major depression. Neuropsychopharmacology.

Alexopoulos, G.S., Hoptman, M.J., Kanellopoulos, D., Murphy, C.F., Lim, K.O., Gunning, F.M., 2012. Functional connectivity in the cognitive control network and the default mode network in late-life depression. J. Affect. Disord. 139, 56–65. https://doi.org/10.1016/j.jad.2011.12.002

American Psychiatric Association, 2013. Diagnostic and statistical manual of mental disorders, 5th ed. American Psychiatric Publishing, Arlington, VA.

Baker, J.T., Holmes, A.J., Masters, G.A., Yeo, B.T.T., Krienen, F., Buckner, R.L., Öngür, D., 2014. Disruption of Cortical Association Networks in Schizophrenia and Psychotic Bipolar Disorder. JAMA Psychiatry 71, 109. https://doi.org/10.1001/jamapsychiatry.2013.3469

Baune, B.T., Miller, R., McAfoose, J., Johnson, M., Quirk, F., Mitchell, D., 2010. The role of cognitive impairment in general functioning in major depression. Psychiatry Res. 176, 183–189. https://doi.org/10.1016/j.psychres.2008.12.001

Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J R Stat Soc Ser. B Stat Methodol 289–300.

Bilker, W.B., Hansen, J.A., Brensinger, C.M., Richard, J., Gur, R.E., Gur, R.C., 2012. Development of abbreviated nine-item forms of the Raven's Standard Progressive Matrices Test. Assessment 354–369.

Biswal, B., Zerrin Yetkin, F., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar mri. Magn. Reson. Med. 34, 537–541.

Boehm, I., Geisler, D., King, J.A., Ritschel, F., Seidel, M., Deza Araujo, Y., Petermann, J., Lohmeier, H., Weiss, J., Walter, M., Roessner, V., Ehrlich, S., 2014. Increased resting state functional connectivity in the fronto-parietal and default mode network in anorexia nervosa. Front. Behav. Neurosci. 8. https://doi.org/10.3389/fnbeh.2014.00346

Box, G.E., Cox, D.R., 1964. An analysis of transformations. J. R. Stat. Soc. Ser. B Methodol. 211–252.

Brakowski, J., Spinelli, S., Dorig, N., Bosch, O.G., Manoliu, A., Holtforth, M.G., Seifritz, E., 2017. Resting state brain network function in major depression - Depression symptomatology, antidepressant treatment effects, future research. J. Psychiatr. Res. 92, 147–159. https://doi.org/10.1016/j.jpsychires.2017.04.007

Buchanan, A., Wang, X., Gollan, J.K., 2014. Resting-state functional connectivity
in women with Major Depressive Disorder. J. Psychiatr. Res. 59, 38–44.
https://doi.org/10.1016/j.jpsychires.2014.09.002
Buckner, R.L., Sepulcre, J., Talukdar, T., Krienen, F.M., Liu, H., Hedden, T.,
Andrews-Hanna, J.R., Sperling, R.A., Johnson, K.A., 2009. Cortical Hubs
Revealed by Intrinsic Functional Connectivity: Mapping, Assessment of
Stability, and Relation to Alzheimer's Disease. J. Neurosci. 29, 1860–
1873. https://doi.org/10.1523/JNEUROSCI.5062-08.2009
Carleton, R.N., Thibodeau, M.A., Teale, M.J.N., Welch, P.G., Abrams, M.P.,
Robinson, T., Asmundson, G.J.G., 2013. The Center for Epidemiologic
Studies Depression Scale: A Review with a Theoretical and Empirical
Examination of Item Content and Factor Structure. PLoS ONE 8, e58067.
https://doi.org/10.1371/journal.pone.0058067
Cattell, R.B., Horn, J.L., 1978. A check on the theory of fluid and crystallized
intelligence with description of new subtest designs. J. Educ. Meas. 15,
139–164.
Centers for Disease Control and Prevention (CDC), 2010. Current depression
among adults---United States, 2006 and 2008. MMWR Morb. Mortal.
Wkly. Rep. 59, 1229–1235.
Chein, J.M., Schneider, W., 2005. Neuroimaging studies of practice-related
change: fMRI and meta-analytic evidence of a domain-general control
network for learning. Cogn. Brain Res. 25, 607–623.
https://doi.org/10.1016/j.cogbrainres.2005.08.013
Cole, J.C., Rabin, A.S., Smith, T.L., Kaufman, A.S., 2004. Development and
Validation of a Rasch-Derived CES-D Short Form. Psychol. Assess. 16,
360–372. https://doi.org/10.1037/1040-3590.16.4.360
Cole, M.W., Anticevic, A., Repovs, G., Barch, D., 2011. Variable Global
Dysconnectivity and Individual Differences in Schizophrenia. Biol.
Psychiatry 70, 43–50. https://doi.org/10.1016/j.biopsych.2011.02.010
Cole, M.W., Pathak, S., Schneider, W., 2010. Identifying the brain's most globally
connected regions. NeuroImage 49, 3132–3148.
https://doi.org/10.1016/j.neuroimage.2009.11.001
Cole, M.W., Repovš, G., Anticevic, A., 2014. The Frontoparietal Control System
A Central Role in Mental Health. The Neuroscientist 20, 652–664.
Cole, M.W., Reynolds, J.R., Power, J.D., Repovs, G., Anticevic, A., Braver, T.S.,
2013. Multi-task connectivity reveals flexible hubs for adaptive task
control. Nat. Neurosci. 16, 1348–1355. https://doi.org/10.1038/nn.3470
Cole, M.W., Schneider, W., 2007. The cognitive control network: Integrated
cortical regions with dissociable functions. NeuroImage 37, 343–360.
https://doi.org/10.1016/j.neuroimage.2007.03.071
Cowdrey, F.A., Filippini, N., Park, R.J., Smith, S.M., McCabe, C., 2014.
Increased resting state functional connectivity in the default mode network
in recovered anorexia nervosa: Resting State Functional Connectivity in
the DMN in Recovered AN. Hum. Brain Mapp. 35, 483–491.
https://doi.org/10.1002/hbm.22202

Dosenbach, N.U., Fair, D.A., Miezin, F.M., Cohen, A.L., Wenger, K.K., Dosenbach, R.A., Fox, M.D., Snyder, A.Z., Vincent, J.L., Raichle, M.E., others, 2007. Distinct brain networks for adaptive and stable task control in humans. Proc. Natl. Acad. Sci. 104, 11073–11078.

Drysdale, A.T., Grosenick, L., Downar, J., Dunlop, K., Mansouri, F., Meng, Y., Fetcho, R.N., Zebley, B., Oathes, D.J., Etkin, A., Schatzberg, A.F., Sudheimer, K., Keller, J., Mayberg, H.S., Gunning, F.M., Alexopoulos, G.S., Fox, M.D., Pascual-Leone, A., Voss, H.U., Casey, B., Dubin, M.J., Liston, C., 2016. Resting-state connectivity biomarkers define neurophysiological subtypes of depression. Nat. Med. 23, 28–38. https://doi.org/10.1038/nm.4246

Dumontheil, I., Thompson, R., Duncan, J., 2011. Assembly and use of new task rules in fronto-parietal cortex. J. Cogn. Neurosci. 23, 168–182.

Duncan, J., 2010. The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. Trends Cogn. Sci. 14, 172–179. https://doi.org/10.1016/j.tics.2010.01.004

Fornito, A., Yoon, J., Zalesky, A., Bullmore, E.T., Carter, C.S., 2011. General and Specific Functional Connectivity Disturbances in First-Episode Schizophrenia During Cognitive Control Performance. Biol. Psychiatry 70, 64–72. https://doi.org/10.1016/j.biopsych.2011.02.019

Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C.F., Jenkinson, M., Smith, S.M., Van Essen, D.C., 2016. A multi-modal parcellation of human cerebral cortex. Nature 536, 171–178. https://doi.org/10.1038/nature18933

Glasser, M.F., Sotiropoulos, S.N., Wilson, J.A., Coalson, T.S., Fischl, B., Andersson, J.L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J.R., Van Essen, D.C., Jenkinson, M., 2013. The minimal preprocessing pipelines for the Human Connectome Project. NeuroImage 80, 105–124. https://doi.org/10.1016/j.neuroimage.2013.04.127

Gordon, E.M., Laumann, T.O., Adeyemo, B., Huckins, J.F., Kelley, W.M., Petersen, S.E., 2016. Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. Cereb. Cortex 26, 288–303. https://doi.org/10.1093/cercor/bhu239

Greicius, M.D., Flores, B.H., Menon, V., Glover, G.H., Solvason, H.B., Kenna, H., Reiss, A.L., Schatzberg, A.F., 2007. Resting-state functional connectivity in major depression: abnormally increased contributions from subgenual cingulate cortex and thalamus. Biol. Psychiatry 62, 429–437.

Gress-Smith, J.L., Roubinov, D.S., Andreotti, C., Compas, B.E., Luecken, L.J., 2015. Prevalence, Severity and Risk Factors for Depressive Symptoms and Insomnia in College Undergraduates: Depressive Symptoms and Insomnia in College. Stress Health 31, 63–70. https://doi.org/10.1002/smi.2509

Hamilton, J.P., Furman, D.J., Chang, C., Thomason, M.E., Dennis, E., Gotlib, I.H., 2011. Default-Mode and Task-Positive Network Activity in Major Depressive Disorder: Implications for Adaptive and Maladaptive

Rumination. Biol. Psychiatry 70, 327–333.
https://doi.org/10.1016/j.biopsych.2011.02.003

Herrero, J., Meneses, J., 2006. Short Web-based versions of the perceived
stress (PSS) and Center for Epidemiological Studies-Depression (CESD)
Scales: a comparison to pencil and paper responses among Internet
users. Comput. Hum. Behav. 22, 830–846.
https://doi.org/10.1016/j.chb.2004.03.007

Hwang, J.W., Egorova, N., Yang, X.Q., Zhang, W.Y., Chen, J., Yang, X.Y., Hu,
L.J., Sun, S., Tu, Y., Kong, J., 2015. Subthreshold depression is
associated with impaired resting-state functional connectivity of the
cognitive control network. Transl. Psychiatry 5, e683.
https://doi.org/10.1038/tp.2015.174

Ito, T., Kulkarni, K.R., Schultz, D.H., Mill, R.D., Chen, R.H., Solomyak, L.I., Cole,
M.W., 2017. Cognitive task information is transferred between brain
regions via resting-state network topology. Nat. Commun. 8.
https://doi.org/10.1038/s41467-017-01000-w

Kaiser, R.H., Andrews-Hanna, J.R., Wager, T.D., Pizzagalli, D.A., 2015. Large-
Scale Network Dysfunction in Major Depressive Disorder: A Meta-analysis
of Resting-State Functional Connectivity. JAMA Psychiatry 72, 603.
https://doi.org/10.1001/jamapsychiatry.2015.0071

Li, B., Liu, L., Friston, K.J., Shen, H., Wang, L., Zeng, L.-L., Hu, D., 2013. A
Treatment-Resistant Default Mode Subnetwork in Major Depression. Biol.
Psychiatry 74, 48–54. https://doi.org/10.1016/j.biopsych.2012.11.007

Li, F., He, N., Li, Y., Chen, L., Huang, X., Lui, S., Guo, L., Kemp, G.J., Gong, Q.,
2014. Intrinsic brain abnormalities in attention deficit hyperactivity
disorder: a resting-state functional MR imaging study. Radiology 272,
514–523.

Liang, X., Zou, Q., He, Y., Yang, Y., 2013. Coupling of functional connectivity and
regional cerebral blood flow reveals a physiological basis for network hubs
of the human brain. Proc. Natl. Acad. Sci. 110, 1929–1934.
https://doi.org/10.1073/pnas.1214900110

Manoliu, A., Meng, C., Brandl, F., Doll, A., Tahmasian, M., Scherr, M.,
Schwerthöffer, D., Zimmer, C., Förstl, H., Bäuml, J., Riedl, V.,
Wohlschläger, A.M., Sorg, C., 2014. Insular dysfunction within the
salience network is associated with severity of symptoms and aberrant
inter-network connectivity in major depressive disorder. Front. Hum.
Neurosci. 7. https://doi.org/10.3389/fnhum.2013.00930

Murrough, J.W., Abdallah, C.G., Anticevic, A., Collins, K.A., Geha, P., Averill,
L.A., Schwartz, J., DeWilde, K.E., Averill, C., Jia-Wei Yang, G., Wong, E.,
Tang, C.Y., Krystal, J.H., Iosifescu, D.V., Charney, D.S., 2016. Reduced
global functional connectivity of the medial prefrontal cortex in major
depressive disorder: Global Connectivity in MDD. Hum. Brain Mapp. 37,
3214–3223. https://doi.org/10.1002/hbm.23235

Park, B., Hong, J., Lee, S.-H., Park, H., 2016. Functional Connectivity of Child
and Adolescent Attention Deficit Hyperactivity Disorder Patients:

Correlation with IQ. Front. Hum. Neurosci. 10.
https://doi.org/10.3389/fnhum.2016.00565

Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012.
Spurious but systematic correlations in functional connectivity MRI
networks arise from subject motion. NeuroImage 59, 2142–2154.
https://doi.org/10.1016/j.neuroimage.2011.10.018

Power, J.D., Cohen, A.L., Nelson, S.M., Wig, G.S., Barnes, K.A., Church, J.A.,
Vogel, A.C., Laumann, T.O., Miezin, F.M., Schlaggar, B.L., Petersen, S.E.,
2011. Functional Network Organization of the Human Brain. Neuron 72,
665–678. https://doi.org/10.1016/j.neuron.2011.09.006

Power, J.D., Schlaggar, B.L., Lessov-Schlaggar, C.N., Petersen, S.E., 2013.
Evidence for Hubs in Human Functional Brain Networks. Neuron 79, 798–
813. https://doi.org/10.1016/j.neuron.2013.07.035

Schneider, W., 2003. Controlled & automatic processing: behavior, theory, and
biological mechanisms. Cogn. Sci. 27, 525–559.
https://doi.org/10.1016/S0364-0213(03)00011-9

Sepulcre, J., Liu, H., Talukdar, T., Martincorena, I., Yeo, B.T.T., Buckner, R.L.,
2010. The Organization of Local and Distant Functional Connectivity in the
Human Brain. PLoS Comput. Biol. 6, e1000808.
https://doi.org/10.1371/journal.pcbi.1000808

Sheline, Y.I., Barch, D.M., Price, J.L., Rundle, M.M., Vaishnavi, S.N., Snyder,
A.Z., Mintun, M.A., Wang, S., Coalson, R.S., Raichle, M.E., 2009. The
default mode network and self-referential processes in depression. Proc.
Natl. Acad. Sci. 106, 1942–1947.

Siegel, J.S., Mitra, A., Laumann, T.O., Seitzman, B.A., Raichle, M., Corbetta, M.,
Snyder, A.Z., 2016. Data Quality Influences Observed Links Between
Functional Connectivity and Behavior. Cereb. Cortex.
https://doi.org/10.1093/cercor/bhw253

Spronk, M., Ji, J.L., Kulkarni, K., Repovs, G., Anticevic, A., Cole, M.W., 2017.
Mapping the human brain's cortical-subcortical functional network
organization. bioRxiv 206292.

Sylvester, C.M., Corbetta, M., Raichle, M.E., Rodebaugh, T.L., Schlaggar, B.L.,
Sheline, Y.I., Zorumski, C.F., Lenze, E.J., 2012. Functional network
dysfunction in anxiety and anxiety disorders. Trends Neurosci. 35, 527–
535. https://doi.org/10.1016/j.tins.2012.04.012

Van Dam, N.T., Earleywine, M., 2011. Validation of the Center for Epidemiologic
Studies Depression Scale—Revised (CESD-R): Pragmatic depression
assessment in the general population. Psychiatry Res. 186, 128–132.
https://doi.org/10.1016/j.psychres.2010.08.018

Van Essen, D.C., Smith, S.M., Barch, D.M., Behrens, T.E.J., Yacoub, E., Ugurbil,
K., 2013. The WU-Minn Human Connectome Project: An overview.
NeuroImage 80, 62–79. https://doi.org/10.1016/j.neuroimage.2013.05.041

Veer, I.M., 2010. Whole brain resting-state analysis reveals decreased functional
connectivity in major depression. Front. Syst. Neurosci. 4.
https://doi.org/10.3389/fnsys.2010.00041

Wei, X., Shen, H., Ren, J., Li, X., Xu, X., Yang, R., Lai, L., Chen, L., Hu, J., Liu,
W., Jiang, X., 2014. Altered Resting-State Connectivity in College
Students with Nonclinical Depressive Symptoms. PLoS ONE 9, e114603.
https://doi.org/10.1371/journal.pone.0114603
Wig, G.S., Schlaggar, B.L., Petersen, S.E., 2011. Concepts and principles in the
analysis of brain networks: Brain networks. Ann. N. Y. Acad. Sci. 1224,
126–146. https://doi.org/10.1111/j.1749-6632.2010.05947.x
World Health Organization, 2017. Depression and other common mental
disorders: Global health estimates. World Health Organization, Geneva.
World Health Organization, 1992. The ICD-10 classifications of mental and
behavioural disorder: Clinical descriptions and diagnostic guidelines.
World Health Organization, Geneva.
Yang, G.J., Murray, J.D., Wang, X.-J., Glahn, D.C., Pearlson, G.D., Repovs, G.,
Krystal, J.H., Anticevic, A., 2016. Functional hierarchy underlies
preferential connectivity disturbances in schizophrenia. Proc. Natl. Acad.
Sci. 113, E219–E228. https://doi.org/10.1073/pnas.1508436113
Yeo, B.T.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D.,
Hollinshead, M., Roffman, J.L., Smoller, J.W., Zollei, L., Polimeni, J.R.,
Fischl, B., Liu, H., Buckner, R.L., 2011. The organization of the human
cerebral cortex estimated by intrinsic functional connectivity. J.
Neurophysiol. 106, 1125–1165. https://doi.org/10.1152/jn.00338.2011
851