Task-related effective connectivity reveals that the cortical rich club gates cortex-wide communication

Abbreviated Title

The cortical rich club gates network communication

Authors

Mario Senden[1,2*], Niels Reuter[3,4*], Martijn P. van den Heuvel[5], Rainer Goebel[1,2,6], Gustavo Deco[7,8,9,10], Matthieu Gilson[7]

* These authors contributed equally to the paper

1) Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience, Maastricht University, 6201BC Maastricht, The Netherlands

2) Maastricht Brain Imaging Centre, Faculty of Psychology and Neuroscience, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands

3) Institute of Systems Neuroscience and Institute of Clinical Neuroscience & Medical Psychology, Medical Faculty, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

4) Institute of Neuroscience and Medicine (INM-1, INM-7), Research Centre Jülich, Jülich, Germany

5) Brain Center Rudolf Magnus, Department of Psychiatry, University Medical Center Utrecht, Brain Center Rudolf Magnus, 3508 GA Utrecht, The Netherlands

6) Department of Neuroimaging and Neuromodeling, Netherlands Institute for Neuroscience, an Institute of the Royal Netherlands Academy of Arts and Sciences (KNAW), 1105BA Amsterdam, The Netherlands

7) Center for Brain and Cognition, Computational Neuroscience Group, Department of Information and Communication Technologies, Universitat Pompeu Fabra, Roc Boronat 138, Barcelona, 08018, Spain

8) Institució Catalana de la Recerca i Estudis Avançats (ICREA), Universitat Pompeu Fabra, Passeig Lluís Companys 23, Barcelona, 08010, Spain

9) Department of Neuropsychology, Max Planck Institute for Human Cognitive and Brain Sciences, 04103 Leipzig, Germany

10) School of Psychological Sciences, Monash University, Melbourne, Clayton VIC 3800, Australia

Address for Correspondence

Mario Senden, Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience, Maastricht University, Oxfordlaan 55, 6200 MD Maastricht, P.O. Box 616, The Netherlands, Phone number: +31 43 38 82071;

Email: mario.senden@maastrichtuniversity.nl

Abstract

Higher cognition may require the globally coordinated integration of specialized brain regions into functional networks. A collection of cortical hubs – referred to as the rich club – has been hypothesized to support task-specific functional integration. In the present paper, we use a whole-cortex model to estimate directed interactions between 68 cortical regions from fMRI activity for four different tasks (reflecting different cognitive domains) and resting state. We analyze the state-dependent input and output effective connectivity of the rich club and relate these to whole-cortex dynamics and network reconfigurations. We find that the cortical rich club exhibits an increase in outgoing effective connectivity during task performance as compared to rest while incoming connectivity remains constant. Increased outgoing connectivity targets a sparse set of peripheral regions with specific regions strongly overlapping between tasks. At the same time, community detection analyses reveal massive reorganizations of interactions among peripheral regions, including those serving as target of increased rich cub output. This suggests that while peripheral regions can play a role in several tasks, their interplay might nonetheless be task-specific. Furthermore, we observe that whole-cortex dynamics are faster during task as compared to rest. The decoupling effects usually accompanying faster dynamics appear to be counteracted by the increased rich club outgoing effective connectivity. Together our findings speak to a gating mechanism of the rich club that supports fast-paced information exchange among relevant peripheral regions in a task-specific and goal-directed fashion, while constantly listening to the whole network.

Introduction

The brain's structural connectivity profile has been shown to contain a set of densely interconnected hub regions (Colizza, Flammini, Serrano, & Vespignani, 2006;van den Heuvel & Sporns, 2011; Zamora-López, Zhou, & Kurths, 2009). These regions, collectively termed the rich club, have been hypothesized to form a central high-capacity backbone for brain communication (van den Heuvel, Kahn, Goñi, & Sporns, 2012). As such they could play a crucial role in the integration of segregated brain regions into transient functional networks assumed to underlie higher cognition (Baars, 2005; Deco, Jirsa, & McIntosh, 2011; Deco, Van Hartevelt, Fernandes, Stevner, & Kringelbach, 2017; Dehaene & Naccache, 2001; Ghosh et al., 2008).

Several lines of research support this notion. Neuroimaging studies have shown that echoes of resting- and task related functional networks (Braga, Sharp, Leeson, Wise, & Leech, 2013; Leech, Braga, & Sharp, 2012) are present in the blood oxygen-level dependent (BOLD) time-series observed within cortical hub regions, indicative of the rich club serving as a central relay for cortical communication. Further support comes from the observation that selective disruption of connectivity among cortical hubs observed in schizophrenia is associated with reduced global communication capacity (Martijn P. van den Heuvel et al., 2013). Simulation studies have shown that cortical hubs, and specifically the rich club, may allow the brain to sustain a large functional repertoire characterized by diverse configurations of peripheral regions, i.e. regions of lower structural degree, around a stable high-degree core (Deco, Senden, & Jirsa, 2012; Senden, Deco, De Reus, Goebel, & van den Heuvel, 2014). Furthermore, there is evidence that rich club regions harmonize peripheral regions by engaging in infraslow oscillations which they exhibit exclusively during task performance (Senden, Reuter, van den Heuvel, Goebel, & Deco, 2017). Finally, the presence of hubs has been shown to increase network controllability (Liu, Slotine, & Barabási, 2011) by providing peripheral regions with the means to exhibit control over the network (Gu et al., 2015; Liu et al., 2011).

In light of these findings it is reasonable to conceive of the rich club as a central workspace of information integration wherein peripheral brain regions compete for control of the system, as recently proposed by Shanahan (2012). In this context, the rich club might act as a gate furthering communication among a winning set of task-relevant peripheral regions while intercepting signals from competing regions. During rest, the gate should largely be closed and the rich club should thus receive more input from peripheral regions than sending

output to these regions. During task performance, the gate would then be opened with the rich club increasing its output. This increase should not result in broad communication between all peripheral regions but rather in the establishment of a specific community with the rich club relaying information between a sparse, task-relevant, set of peripheral regions.

Figure 1: Brain regions are classified as rich club or peripheral regions. Communication occurs within each set of regions (black arrows) as well as between regions of different sets. *Panel A)* During rest, the rich club receives a fair amount of input from the periphery (purple arrows) but provides only little output back to peripheral regions (red arrows). *Panels B & C)* During task performance, the rich cub increases the output it projects to the periphery. The specific peripheral targets of increased rich club output may depend on which task is currently being performed.

The present study aims to address these hypotheses using an integrative approach combining empirical functional magnetic resonance imaging (fMRI) data with large-scale computational modeling. Specifically, we employ a recently developed estimation procedure for whole-cortex effective connectivity in a noise-diffusion network model that reproduces the empirical spatiotemporal covariance of BOLD fluctuations (Gilson et al., 2016). The latter have been shown to convey information about the behavioral conditions of subjects (Mitra et al., 2015). Our whole-brain model captures changes in this spatiotemporal FC and interprets them in terms of local activity and network connectivity (Gilson et al., 2017). In contrast to previous modeling studies this procedure does not rely on identification of an optimal working point given fixed structural connectivity (Deco et al., 2013; Messé, Rudrauf, Benali, Marrelec, & Honey, 2014) but optimizes the strengths of interactions between brain regions in addition to their individual level of activity. The resulting estimates of directed effective connectivity allow us to study input/output relations of the rich club for rest and various task conditions in order to evaluate our gating hypothesis under an information propagation perspective. We optimized connectivity for a collection of fMRI measurements during which subjects engaged in a range of tasks tapping into different cognitive domains (visual working memory, response inhibition, mental rotation, verbal reasoning) and whose associated functional connectivity profiles are minimally overlapping (Smith et al., 2009).

Materials and Methods

Whole-Cortex Dynamic Model to Fit Empirical BOLD Covariances

The model comprised $N = 68$ cortical regions of interest (ROI; Desikan et al., 2006), whose local fluctuating activity (an amalgamation of internal processing and external input) is shaped by the recurrent effective connectivity C. Following Gilson et al. (2016; https://github.com/MatthieuGilson/optimization_linear_EC_Sigma.git), the model aims to reproduce BOLD covariances with and without time shift, which are calculated as:

$$\hat{Q}_{ij}^{\tau} = \frac{\sum\limits_{t}^{T-\tau} x_i^t x_j^{t+\tau}}{T - \tau} \tag{1}$$

where s_i^t are the centered BOLD time series for each region $1 \leq i \leq N$ with time indexed by $1 \leq t \leq T$ (*T*=192 time points separated by a TR=2 seconds).

The local dynamics correspond to an Ornstein-Uhlenbeck process, where the nodal activity variable x_i decays exponentially with time constant τ_x and is affected by the rest of the network via $\frac{dx_i}{dt} = \frac{-x_i}{\tau_x} + \sum\limits_{j \neq i} C_{ij} x_j + dB_i$. Here Each ROI experiences local fluctuations formally described by a Wiener process dB_i (equivalent to white Gaussian noise) with a diagonal covariance matrix Σ. The between-region effective connectivity is embodied by the matrix C, whose skeleton is determined by a generic matrix of structural connectivity (see below). The advantage of this model is its analytical tractability, which allows for a quick calculation of its network covariance pattern Q^0 and Q^1. In essence, the model thus decomposes functional connectivity (FC) into two sets of parameters: effective connectivity and local variability. Those can be seen as a fingerprint of brain dynamics, which captures the propagation of fluctuating BOLD activity across ROIs.

Figure 2: Schematic diagram of the model. *Panel A)* shows example BOLD time series (green) of two populations in the network. Dark blue arrows indicate temporal covariance with zero lag, whereas light blue arrows indicate covariance with time series *j* (light green) shifted by one TR (i.e. 2 seconds) with respect to time series *i* (dark green). *Panel B)* shows the logarithm of the autocovariances for each node as a function of time shift exemplary for rest and the n-back task. *Panel C)* shows a schematic diagram of the noise diffusion model: neural populations (gray circles) exhibit local, noisy, fluctuations (purple) as well as recurrent feedback (red arrows). *Panel D)* shows a schematic representation of a single iteration within the Lyapunov optimization procedure. Model covariances ($\hat{Q}^0$ and $\hat{Q}^1$) given current values of the network parameters EC and Σ are evaluated. From the comparison between model covariances with their empirical counterparts, the desired parameter updates for both parameters are calculated.

Estimation Procedure for Effective Connectivity and Local Excitability

We iteratively tune model parameters C and Σ such that model covariance matrices Q^0 and Q^1 best reproduce the empirical FC $\hat{Q}^0$ and $\hat{Q}^1$. We summarize the essential steps of the procedure described in Gilson et al. (2016) that iteratively optimizes the network parameters C

and Σ. For each state (rest & tasks), we calibrate the model by calculating the time constant τ_x associated with the exponential decay of the autocovariance function $\hat{Q}_{ii}^{\tau}$ averaged over all regions: $\tau_x = \frac{1}{\frac{1}{N}\sum_i \log(\hat{Q}_{ii}^0) - \log(\hat{Q}_{ii}^1)}$. This gives the diagonal elements of the Jacobian of the dynamic system: $J_{ij} = -\delta_{ij}\tau_x + C_{ij}$, where δ_{ij} is the Kronecker delta. Starting from no recurrent connectivity ($C = 0$), the model FC matrices Q^0 and Q^1 are calculated by solving the Lyapunov consistency equation $JQ^0 + Q^0 J^T + \Sigma = 0$ using the Bartels-Stewart algorithm and then evaluating $Q^1 = Q^0 \text{expm}(J^T)$, where expm denotes the matrix exponential. The desired Jacobian update is the matrix $\Delta J^T = (Q^0)^{-1}[\Delta Q^0 + \Delta Q^1 \text{expm}(\text{-}J^T \tau)]$, which involves the FC error between the empirical and model matrices: $\Delta Q^0 = \hat{Q}^0 - Q^0$ and $\Delta Q^{\tau} = \hat{Q}^{\tau} - Q^{\tau}$. This yields the connectivity update $\Delta C_{ij} = \eta_C \Delta J_{ij}$ for existing connections and the local variance update $\Delta \Sigma_{ii} = -\eta_{\Sigma}(J\Delta Q^0 + \Delta Q^0 J^T)_{ii}$ for the diagonal elements only. We use $\eta_C = 0.0001$, $\eta_{\Sigma} = 0.1$, and impose non-negativity for the weights in C and the diagonal elements of Σ.

Compared to the classical approach of dynamic causal modelling (DCM) for the estimation of effective connectivity (Friston, 2011), our network model relies on simpler assumptions to obtain a maximum-likelihood estimate of the whole-cortex connectivity. The optimization tunes the model parameters to reproduce the empirical cross-covariances between ROIs, which are canonically related to the cross spectral density used in recent studies that apply DCM to resting state fMRI data (Frässle et al., 2017; Friston, Kahan, Biswal, & Razi, 2014). Moreover, the proposed model relies on an exponential approximation of BOLD autocovariance (locally over a few TRs) and discards very slow-frequency fluctuations below 0.1 Hz. Despite these differences, we still borrow the term "effective connectivity" from DCM literature as our connectivity estimates relate to directional interactions between ROIs in the brain network. This model-based approach has been successfully applied to identify changes in the cortical coordination between rest and movie viewing (Gilson et al., 2017). In essence, our approach aims to bridge the gap between whole-brain modeling where the connectivity is fixed and taken from DTI (Deco et al., 2013; Messé et al., 2014) and studies focusing on a few cortical areas only to test hypotheses on specific brain subsystems (Goebel, Roebroeck, Kim, & Formisano, 2003; He, 2011).

Structural Connectivity (SC)

We obtained high-quality diffusion-weighted MRI data of 215 subjects from the Q3 release of the human connectome project (HCP; Glasser et al., 2013; Van Essen et al., 2012). Combined diffusion tensor and generalized q-sampling imaging (GQI) were used to fit

complex fiber architecture, with white matter pathways reconstructed by means of streamline tractography (Romme, de Reus, Ophoff, Kahn, & van den Heuvel, 2017; Yeh, Wedeen, & Tseng, 2010)., The cortex was parcellated into 68 cortical regions according to the widely used Desikan-Killany (DK) atlas (Desikan et al., 2006; Klein & Tourville, 2012) forming the regions of the reconstructed anatomical connectome map. Details on these processing steps can be found in de Reus and van den Heuvel (2014). A weighted group-average structural connectivity matrix was generated by averaging streamlines over subjects and keeping only those entries which had positive values for at least 60% of subjects (de Reus & van den Heuvel, 2013). Finally, since Gaussian resampling (or taking the logarithm) of the data has previously been shown to enhance correspondence between diffusion tractography and in vivo animal tract-tracing measurements of anatomical connectivity (Martijn P. van den Heuvel et al., 2015), the data was resampled with a mean of 0.5 and a standard deviation of 0.15 (Honey et al., 2009).

Functional MRI Data

We used resting and task-state BOLD time-series data from fourteen healthy subjects (8 females, mean age = 28.76) previously described by Senden *et al.* (2017). Briefly, data comes from five functional runs consisting of a resting-state measurement (eyes closed), four individual task measurements including a visual n-back (n=2) task (Kirchner, 1958), the Eriksen flanker task (Eriksen & Eriksen, 1974), a mental rotation task (Shepard & Metzler, 1971), and a verbal odd-man-out task (Flowers & Robertson, 1985). All runs comprise 192 data points with tasks being continuously performed during this period. For the n-back and flanker task, stimuli were presented at a rate of 0.5 Hz; for the mental rotation and odd-man out tasks they were presented at a rate of 0.25 Hz. The data were acquired using a Siemens Prisma Fit (upgraded Tim Trio) scanner and a 64-channel head coil. In addition to 3D-motion correction and high-pass filtering with a frequency cutoff of .01 Hz, signals were cleaned by performing wavelet despiking (Patel & Bullmore, 2015) and regressing out a global noise signal given by the first principal component of signals observed within the cerebrospinal fluid of the ventricles. Next, voxels were uniquely assigned to one of the 68 cortical ROIs specified by the DK atlas and an average BOLD time-series was computed for each region as the mean time-series over all voxels of that region (preprocessed BOLD time series per ROI and state are publicly available at datadryad.org [*doi not yet available*]). Finally, empirical τ-lag FC matrices were given by the covariances of the z-normalized BOLD time series according to equation (1).

Louvain Community Detection Method

We identified ROI communities within the effective connectivity that maximize the modularity of possible partitions of the network (Newman, 2006), using the implementation developed by Bondel et al. (2008). Modularity measures the excess of connections between ROIs compared to the expected values estimated from the sum of incoming and outgoing weights for the nodes (targets and sources, respectively). In practice, the method stochastically determines local communities and then iteratively aggregates ROIs to maximize the modularity at each step, until reaching a minimum. The process was repeated 20 times for each task to average over the initial random clustering. In the end, the reported value is the participation index for each pair of ROIs to belong to the same community.

The overlap between communities is evaluated using the binary matrices M obtained from each repetition of the Louvain algorithm, where the element M_{ij} is 1 if the ROIs i and j belong to the same community and 0 otherwise. The overlap measure for two matrices M^1 and M^2 is given by $\sum_{ij} M_{ij}^1 M_{ij}^2 \Big/ \sqrt{\sum_{ij} M_{ij}^1 \sum_{ij} M_{ij}^2}$, giving a number between 0 (no overlap) and 1 (identical matrices).

Statistical Analyses

In order to assess to what extent a set of brain regions (such as the rich club) collectively exhibits significant differences from the cortex as a whole, we used a clustered-based bootstrapping procedure. That is, we averaged the measure of interest across regions falling within a previously defined cluster of size *N* (such as the rich club forming a theoretically motivated a-priori cluster). Subsequently, we repeatedly (10,000 times) sampled *N* regions from the entire set of cortical regions with replacement and calculated the average within each sample.

In order to assess differences observed for tasks with respect to rest, we employed a blocked bootstrapping procedure. Null distributions for each task were created from running connectivity optimization for 1,000 randomly resampled functional rest and task samples, calculating the difference (task - rest) for effective connectivity, and obtaining average changes in the measure of interest. Samples were created by first randomly drawing from the subject pool with replacement and then randomly placing one of each subject's two states in the *rest* and the other in the *task* sample repeatedly until each sample comprised 14 subjects. Connectivity and local variability (sigma) were then optimized in each of these samples.

We compared the overlap of the communities corresponding the estimated ECs with that for random communities generated by partitioning the ROI indices into 4 to 7 groups (same range as the original communities). We repeated the process 1000 times to obtain a Null distribution of overlap measures. The difference between distributions was assessed using Welch's t-test.

All p-values reported in the results have been corrected for multiple comparisons according to the false discovery rate (FDR) controlling procedure.

Results

Rich club

We followed the procedure outlined in Senden *et al.* (2017) to identify rich club regions in the SC. Specifically, we binarized the SC matrix by setting its non-zero entries to one. From this binary matrix rich club coefficients were calculated as the fraction of the number of existing connections between regions with degree larger than k to the possible number of connections among these regions (Colizza et al., 2006; Martijn P. van den Heuvel & Sporns, 2011; Zhou & Mondragon, 2004). Next, the statistical significance of rich club coefficients for each degree k was determined by calculating the rich club coefficients for a set of 1000 degree-preserving rewired adjacency matrices (Maslov & Sneppen, 2002) and identifying the first k for which the rich club coefficient of the binarized SC was larger than the 95[th] percentile of the rich club coefficients corresponding to the rewired matrices. In our data, this was the case for $k = 21$. Seven candidate rich club regions were subsequently identified as those whose degree exceeded this cutoff. These included the bilateral precuneus, the bilateral superior frontal cortex, the bilateral superior parietal cortex, and the right insula. To ensure that these regions were not only individually rich but indeed also formed part of a dense club we evaluated the contribution of each region to the internal density of the set. To that end, we calculated internal density of the full set as well as of subsets resulting from leaving each candidate region out once. This procedure revealed that removal of the right insula lead to an increase in internal density by 21.33%. Since this exceeded the 95[th] percentile of observed density changes (± 8.31%), the right insula was not considered a rich club region in this study. The six remaining regions have been identified as rich cub hub regions in a multitude of studies (e.g. Daianu et al., 2015; Dennis et al., 2013; M. P. van den Heuvel, Scholtens, Feldman Barrett, Hilgetag, & de Reus, 2015; Martijn P. van den Heuvel & Sporns, 2011)

Local variability

We first examined the local variability (Σ) exhibited by rich club and peripheral regions for resting and the four task states. Local variability exhibited by rich club regions was low compared to peripheral regions irrespective of state. Specifically, mean local variability within the rich club was 0.35 (95% CI [0.10, 0.59]), 0.40 (95% CI [0.19, 0.60]), 0.34 (95% CI [0.11, 0.57]), 0.39 (95% CI [0.24, 0.45]), and 0.38 (95% CI [0.16, 0.60]) for rest, the n-back task, the flanker task, the mental rotation task, and the odd-man out task, respectively. In contrast, local variability exhibited by peripheral regions was 1.36 (95% CI [1.14, 1.58]), 1.50 (95% CI [1.24, 1.77]), 1.43 (95% CI [1.19, 1.67]), 1.46 (95% CI [1.21, 1.70]), 1.41 (95% CI [1.16, 1.66]). A cluster-based bootstrapping test revealed that local variability exhibited by the rich club was significantly lower than that exhibited by peripheral regions in each state (figure 3a; $p \ll 0.01$ for all comparisons). Note that these results do not imply that the model predicts no fluctuations in rich club activity as measured by the variance given by the diagonal of the model FC matrix ($\hat{Q}^0$). Mean variance for rich club regions was 1.00 (95% CI [0.78, 1.22]), 0.73 (95% CI [0.55, 0.91]), 0.80 (95% CI [0.63, 0.96]), 0.75 (95% CI [0.51, 0.99]), 0.77 (95% CI [0.61, 0.92]), for the five states, respectively. Mean variance of peripheral regions was 1.13 (95% CI [1.02, 1.25]), 0.93 (95% CI [0.83, 1.04]), 0.98 (95% CI [0.87, 1.09]), 0.91 (95% CI [0.80, 1.03]), and 0.93 (95% CI [0.82, 1.04]), for the five states respectively. Our clustered-based bootstrap procedures showed that differences between rich club and peripheral regions were not significant for any of the five states (figure 3b). Our model results thus reveal that fluctuating activity observed for the rich club is largely the result of network contributions from peripheral regions and not due to local variability.

Figure 3: Cluster-based bootstrap analyses. This figure shows the region enclosed by the 2.5th and 97.5th percentile of sampled cluster averages (blue) for each task. Averages observed for the a priori rich club cluster falling inside this region (red) were considered non-significant while those falling outside this range (green) were considered significant. *Panel A)* shows results for local variability Σ. Local variability exhibited by the rich club was significantly lower than that exhibited by peripheral regions for rest and each task. *Panel B)* shows results for model variance $diag(\hat{Q}^0)$. Variance observed for the rich club did not differ significantly from that observed for peripheral regions for either rest or task states. *Panel C)* shows results for the input-output ratio φ. The ratio exhibited by the

rich club was significantly larger than that exhibited by peripheral regions for rest but not for any of the tasks.

Input-output ratio

We examined to what extent the rich club gates the input it receives from peripheral regions by computing the input-to-output ratio of total incoming/outgoing effective connectivity. We focus on this ratio rather than absolute input and output strength since the rich club is likely to exceed peripheral regions in both simply by virtue of its high structural degree. The ratio provides an indication in how far the received input is passed on and is less dependent on structural degree. During rest, the total input the rich club receives from peripheral regions was roughly twice ($\varphi_{rc} = 2.12$, 95% CI [0.33, 3.90]) the output it sends to the periphery. In contrast, the ratio exhibited by peripheral regions was only 1.14, 95% CI [1.03, 1.26]. During task performance, the rich club presented with lower input-output ratios. Specifically, the ratio exhibited by the rich club was 1.15 (95% CI [0.42, 1.88]), 1.71 (95% CI [0.52, 2.90]), 0.97 (95% CI [0.53, 1.41]), and 1.48 (95% CI [0.36, 2.60]) for the n-back, flanker, mental rotation, and odd-man out task respectively. The corresponding ratios observed for peripheral regions were 1.24 (95% CI [1.05, 1.44]), 1.18 (95% CI [1.03, 1.32]), 1.20 (95% CI [1.04, 1.36]), and 1.19 (95% CI [1.03, 1.34]). To assess whether input-output ratios exhibited by the rich club were significantly different from those exhibited by the cortex as a whole for each task, we again employed the cluster-based bootstrapping procedure. In line with our expectation, the input-output ratio was significantly larger for the rich club compared to peripheral regions during rest ($p = 0.02$) but not during any of the task states (figure 3c).

Engaging in tasks is accompanied by increased rich club output

We follow up on the previous results by investigating to what extent decreases in the input-output ratio are due to the rich club receiving a decreased amount of input from the periphery or due to the rich club projecting an increased amount of output to the periphery during task states as compared to rest. To that end, we calculated the difference between effective connectivity observed during rest and during each of the task states and calculated the average change in input (output) received from (sent to) the periphery across the rich club. We assessed the significance of these changes per task by performing one-sided blocked bootstrap tests.

Input to the rich club from the periphery slightly increased (rather than decreased) for all tasks compared to rest. Specifically, average changes in input were 1.57 (95% CI [0.73,

2.41]), 1.45 (95% CI [0.67, 2.23]), 1.82 (95% CI [0.96, 2.67]), and 0.95 (95% CI [0.38, 1.52]) for the n-back, flanker, mental rotation, and odd-man out task, respectively. There could thus no significant decreases in input be observed. Increases in output from the rich club, on the other hand, were significant for all tasks except the flanker task ($p = 0.04$, $p = 0.09$, $p = 0.04$, and $p = 0.04$, for n-back, flanker, mental rotation, and odd-man out tasks). Specifically, change in output was 5.30 (95% CI [3.96, 6.91]), 5.71 (95% CI [4.72, 6.70]), and 3.73 (95% CI [1.97, 5.50]), for the n-back, mental rotation, and odd-man out task, respectively while it was 2.26 (95% CI [0.74, 3.79]) for the flanker task.

After establishing that the rich club is more conducive to whole-cortex communication during task performance compared to rest, we identified the specific peripheral targets of increased output from the rich club. These are defined as those peripheral regions which receive an increase of output exceeding the 95[th] percentile obtained from a blocked bootstrap procedure on differences in effective connectivity between tasks and rest from at least one rich club region. Figure 4 shows an overview of the thusly identified target regions per task. For the n-back task, the rich club displayed significantly increased output to 18 out of 62 peripheral regions. For the flanker task, it displayed significantly increased output to 11 peripheral regions. For the mental rotation task, it displayed significantly increased output to 18 peripheral regions. Finally, for the odd-man out task, it displayed significantly increased output to 17 peripheral regions. The increase in output strength was thus sparsely distributed. At the same time, the targets of rich club output for the different tasks were strongly overlapping. Specifically, the n-back task had 5 regions in common with the flanker task, 12 with the mental rotation task, and 10 with the odd-man out task. The flanker task had 9 and 8 regions in common with the mental rotation and odd-man out tasks, respectively. Finally, the mental rotation task had 13 regions in common with the odd-man out task.

Figure 4: Targets of increased rich club output. This figure shows peripheral regions receiving increased output from at least one rich club region for each task as compared to rest. Colors indicate the percent of rich club regions sending increased output to a particular peripheral region with warmer

colors reflecting a larger percentage. Targets of rich club output for the different tasks are largely overlapping. These regions include the bilateral cuneus cortex, the bilateral postcentral gyrus, the left rostral anterior cingulate cortex, the left insula, the right paracentral lobule, the left paracentral lobule, the left precentral gyrus, the left posterior-cingulate cortex, the right fusiform gyrus, the right lingual gyrus, and the right pericalcarine cortex. These regions are generally involved in processes common to each task such as visual perception, attention, and conflict monitoring (see discussion). At the same time, each task also presents with a set of regions serving as targets for increased rich club output uniquely for that task.

The pairwise analytical probability of obtaining the observed or more extreme degrees of overlap given the total number of regions selected per task constitutes a p-value for the Null hypothesis that the observed degree of overlap occurred by chance. FDR corrected p-values for each task pair are presented in table 1. As can be appreciated from this table, the degree of overlap was significant for all task pairs with the exception of the [n-back, flanker] pair. This indicates that many targets for the increased output of the rich club might be task-general. Indeed, the bilateral postcentral gyrus, the left rostral anterior cingulate cortex, the left insula, and the right paracentral lobule are targets of increased rich club output for all tasks. If regions which are target of increased output from the rich club for a majority of tasks are included, this set extends to include the bilateral cuneus cortex, the left paracentral lobule, the left precentral gyrus, the left posterior-cingulate cortex, the right fusiform gyrus, the right lingual gyrus, and the right pericalcarine cortex.

In addition to these task-general target regions for rich club output, each task also presents with a specific set of target regions. For the n-back task these regions include the bilaretal caudal middle frontal gyri, the bilateral lateral occipital cortex, the bilateral pars opercularis, the right pars triangularis, and the right rostral middle frontal gyrus. For the flanker task target regions include the left isthmus-cingulate cortex, the left medial orbitofrontal cortex, and the supramarginal gyrus. For the mental rotation task target regions include the bilateral lateral occipital cortex, the left fusiform gyrus, the left lingual gyrus, and the left medial orbitofrontal cortex. Finally, for the odd-man out task target regions include the bilateral inferior parietal cortex, the right insula, and the right posterior-cingulate cortex.

Task-specific network communication

The final set of analyses focuses on task-related network communication as well as dynamics as compared to rest. In terms of communication, target regions of increased rich club output likely constitute the set of specialized peripheral regions whose integration into

functional networks underlies task performance. As such, for each task we would expect higher levels of communication and hence larger empirical covariance between these target regions as compared to the remaining peripheral regions (i.e. those not forming targets of increased rich club output). Indeed, we find that average covariance (excluding variance) among target peripheral regions was 0.44 (95% CI [0.340, 0.47]), 0.42 (95% CI [0.35 0.47]), 0.48 (95% CI [0.43, 0.53]), and 0.48 (95% CI [0.44, 0.52]), for the n-back, flanker, mental rotation, and odd-man out task, respectively. For comparison, average covariance among non-target peripheral regions was 0.30 (95% CI [0.29, 0.31]), 0.34 (95% CI [0.32 0.35]), 0.28 (95% CI [0.27, 0.29]), and 0.30 (95% CI [0.29, 0.31]), for the four tasks, respectively (see figure 6a). To assess whether average covariance among target peripheral regions was significantly higher than among non-target peripheral for each task, we again employed the cluster-based bootstrapping procedure with the set of target regions identified in each task as cluster and generating a Null distribution by sampling new clusters of equal size from all peripheral regions. In line with our expectation, average covariance was significantly larger for target compared to non-target peripheral regions for the n-back ($p = 0.05$), mental rotation ($p = 0.04$), and odd-man out ($p = 0.04$) but not the flanker task ($p = 0.189$).

Figure 5: Changes in BOLD properties and model estimates with respect to rest. *Panel A)* shows changes in empirically measured region-specific and network time constants (τ_x and τ_c, respectively) from rest to task averaged across tasks. Decreases for both constants reveal that cortical dynamics are generally faster during task performance as compared to rest. *Panel B)* shows changes in empirically observed variance, model variance, and local variability parameter Σ. Red indicate rich club regions, blue indicates peripheral regions serving as target for increased rich club in at least one task, and green indicates the remaining regions. *Panel C)* shows changes in both incoming and outgoing effective connectivity for the three groups of regions (color coding as before).

At the same time, variance (and to a lesser extent covariance) decreased homogeneously across the cortex for task states compared to rest (see figures 5b and 6b). Specifically, average empirically observed variance across all cortical regions was 0.15 (95% CI [0.13, 0.16]), 0.12 (95% CI [0.11 0.13]), 0.13 (95% CI [0.11, 0.14]), 0.12 (95% CI [0.11, 0.13]), and 0.12 (95% CI [0.11, 0.13]), for rest and the four tasks, respectively. Differences in variance between task and rest were significant according to a blocked bootstrap test for the n-back (p = 0.04), mental rotation (p = 0.04), and odd-man out (p = 0.04) tasks while it marginally failed to reach significance for the flanker task (p = 0.053). This global reduction in variance can be explained by decreases in the measured time constants τ_x for individual regions in all tasks (see figure 5a). These decreased nodal time constants lead to faster network dynamics (as measured by τ_c, the negative inverse of the dominating eigenvalue of the Jacobian) and weaken recurrent network interactions. Non-significant increases in local variability Σ and effective connectivity (figure 5b & c), in conjunction with significant, highly specific, increases in outgoing EC from the rich club (figure 6c), partially offset these decoupling effects and are likely necessary to sustain meaningful network communication in light of faster network dynamics during task performance.

Figure 6: Comparison between whole-cortex connectivity observables and estimates for the four tasks. *Panel A)* displays empirical covariance matrices. The two horizontal and vertical lines demarcate rich club regions (lower left corner) from peripheral regions serving as target in a specific

task and those, in turn, from the remaining peripheral regions. *Panel B)* shows the change in covariance with respect to rest. *Panel C)* shows changes in EC with respect to rest.

Finally, we examined whether the changes observed in output connections across tasks have a homogeneous effect over the whole network. The goal was to test whether the network could be simply summarized into three groups (rich club, target and non-target regions) that clearly appeared in figure 6a-c. To do so, we performed community detection (see Methods) to partition the cortex into strongly connected subnetworks. This method compensates for the dense connectivity of rich club regions such that they do not belong to the same community. The outcome shown in figure 7a reveals reorganization of the communities for each task, both for target and non-target peripheral regions. Interestingly, the overlap between the task-communities displayed in figure 7b is at the same level as when shuffling 20-50% of the ROIs from a given community (undistinguishable using Welch t-test with $p>0.5$). In comparison, shuffling 10-20% of the ROIs results in much more overlap and shuffling all much less (random), both with $p\sim0$. This indicates that even non-significant changes in effective connectivity collectively induce a profound reconfiguration of communication at the network level, with an intermediate level of complexity.

Moving one step further, we examine the contribution of the three groups (rich club, target peripheral or non-target peripheral) in figure 7c, by shuffling 6 regions (same number as rich club) within each group (yellow violin plots) or irrespective of the groups (in gray). Shuffling non-target regions reduced the overlap much more than shuffling target regions (with $p \ll 0.01$), and the latter than the rich club (with $p \ll 0.01$). This suggests that the task-specific EC rewiring between the rich club and the target regions mediates a dramatic community reconfiguration for the whole network, especially within peripheral regions.

Figure 7: Overlap between task-related communities. *Panel A)* shows community participation indices between all pairs of cortical regions. Rich club regions are indicated in red (RC), peripheral regions serving as target in all tasks in blue (COM), peripheral regions serving as target in at least one task in pink (OTH), all remaining peripheral regions in black (REM). The plotted participation indices correspond to averages over 20 repetitions of the stochastic detection algorithm per task; the order of the ROIs is the same for the four matrices. *Panel B)* compares the overlap between the communities for pairs of tasks (original) with the overlap between a given community partition and its shuffled version by permuting a portion of the ROIs (as indicated in the x-labels; random corresponds to 100%). *Panel C)* compares the overlap as in B with group-specific shuffling: within RC, within target and within non-targets (yellow distributions). In addition, the grey distribution corresponds to shuffling the same number of regions as for the yellow distribution on the left, but chosen at random from the full set of 68 regions.

Discussion

In line with the hypothesis that the rich club might serve as a gated relay for whole-cortex communication, we found that during rest the rich club impedes transmission of activation between peripheral brain regions as total outgoing effective connectivity from the rich club was half of its total incoming effective connectivity. During task performance, on the other hand, a larger proportion of the input received by the rich club was passed on. The decrease in the input-output ratio observed for the rich club was not due to increases in local fluctuations as these remained consistently low irrespective of state. Neither was it due to decreases in input strength, the rich club always listens to the entire cortex. Instead, the change in input-output ratio was due to increased output strength from the rich club to the

remainder of the cortex. Furthermore, outgoing effective connectivity did not increase for all regions, but only for a specific set of brain regions comprising between 18 and 29 percent of the periphery. These target peripheral regions showed strong coupling among each other (as measured by their empirical covariance) indicative of their functional relevance in their respective task. A subset of these peripheral regions served as targets of increased output from the rich club for all tasks while others were targeted specifically during individual tasks. The former subset included the cuneus cortex and the fusiform and lingual gyri which are involved in visual processing such as object recognition (e.g. Haxby, Gobbini, Furey, & Ishai, 2001; Ishai, Ungerleider, & Martin, 1999; Machielsen, Rombouts, & Barkhof, 2000; Mangun, Buonocore, & Girelli, 1998; Okada, Tanaka, Nakai, Nishizawa, & Inui, 2000; Vanni, Tanskanen, Seppä, Uutela, & Hari, 2001). It further included the posterior and anterior cingulate cortices which are involved in regulating the focus of attention and error/conflict monitoring, respectively (e.g. Bush, Luu, & Posner, 2000; Hampson, Driesen, Skudlarski, Gore, & Constable, 2006). Finally, it included the postcentral gyrus and the insula which are involved in sensing touch and motor control, respectively (e.g. Anderson, Jenkins, Brooks, & Hawken, 1994; Fink, Frackowiak, Pietrzyk, & Passingham, 1997). All of these functions pertain to each of our tasks as stimuli were presented visually, tasks were cognitively demanding, and motor responses were required.

Those regions serving as target for increased rich club output specifically for individual tasks have generally been associated with functions directly relevant for the task at hand. For the visual n-back task, regions included the caudal middle frontal gyri and the pars triangularis which have been implicated in visuospatial and semantic working memory as well as selection among competing items in memory (e.g. Carlson et al., 1998; Gabrieli, Poldrack, & Desmond, 1998; Leung, Gore, & Goldman-Rakic, 2002; Thompson-Schill, D'Esposito, & Kan, 1999). Other regions to which rich club output was increased during the n-back task were the lateral occipital cortex and the pars opercularis which have been associated with object recognition and selective response suppression (e.g. Forstmann, van den Wildenberg, & Ridderinkhof, 2008; Grill-Spector, Kourtzi, & Kanwisher, 2001). For the flanker task, target regions included the medial orbitofrontal cortex and the supramarginal gyrus; both related to decision making and evaluating decision outcomes (e.g. Kringelbach, 2005; Schoenbaum, Takahashi, Liu, & McDannald, 2011; Silani, Lamm, Ruff, & Singer, 2013). For the mental rotation task, regions included the lateral occipital cortex, the fusiform gyrus, and the lingual gyrus which are involved in encoding visual stimuli and object recognition (e.g. Grill-Spector et al., 2001; Haxby et al., 2001; Ishai et al., 1999; Machielsen et al., 2000;

Okada et al., 2000) as well as the medial orbitofrontal cortex which is involved in decision making (e.g. Kringelbach, 2005). Finally, for the verbal odd-man out task, regions included the inferior parietal cortex implicated in word comprehension (e.g. Price, 1998) as well as the right posterior-cingulate cortex which is involved in regulating the focus of attention (e.g. Hampson et al., 2006).

In terms of whole-cortex network dynamics, we observed that BOLD variance decreased for task states as compared to rest and that these changes were due to faster cortical dynamics. This is in line with findings of He (2011) who showed that time-lagged autocorrelation of fMRI signals decreases during task performance suggestive of decreased long-range memory and more efficient online information processing. The fact that faster dynamics occur in spite of increases in recurrent effective connectivity further indicates that rest is closer to criticality than tasks. The cortex is thus optimally positioned in terms of its dynamics to allow for massive reorganization of functional interactions to meet task demands. This is also evident from our community detection analyses which revealed massive reconfigurations between tasks. Interestingly, while reconfigurations occurred among rich cub, target peripheral, and non-target peripheral regions, they exhibit a hierarchical structure with non-target regions exhibiting most pronounced reconfigurations and rich cub regions least pronounced reconfigurations. This may indicate that not only brain regions but also some communication channels (notably among rich club but also among target and to a lesser extent among non-target peripheral regions) may be of importance for all tasks. At the same time, the fact that reconfigurations do occur among target peripheral regions suggests that while these regions may play a role in several tasks, their concrete interplay might nevertheless be highly task-specific.

Taken together, our results provide further support for the notion that the rich club serves as a central workspace wherein peripheral brain regions compete for the establishment of communication channels among them. During rest, slower, temporally dependent network dynamics reflect exploration of the brain's functional repertoire. With impending task demands, network dynamics become faster in order to process information effectively. At the same time, the rich cub increases its outgoing effective connectivity to establish communication among task-relevant peripheral regions and prevent global decoupling (a consequence of faster dynamics). With a transition to task performance, the rich club thus supports the interplay of a set of task-relevant peripheral regions as required for higher cognition.

Acknowledgements

Authors MS and RG were supported by the European Research Council under the European Union's Seventh Framework Programme (ERC-2010-AdG, ERC grant agreement no. 269853) and by the European Union's Horizon 2020 research and innovation programme under grant agreement n. 720270 (HBP SGA1). Author GD was supported by the ERC Advanced Grant: DYSTRUCTURE (n. 295129), by the Spanish Research Project PSI2016-75688-P (AEI/FEDER), and by the European Union's Horizon 2020 research and innovation programme under grant agreement n. 720270 (HBP SGA1). Author MG was supported by a Marie Sklodowska-Curie Action grant (H2020-MSCA-656547), by European Research Council Advanced Grant DYSTRUCTURE (Grant 295129), and by the European Union via the Human Brain Project (grant FP7-FET-ICT-604102). Author MPvdH was supported by a VIDI grant of The Netherlands Organization for Scientific Research (NWO 452-16-015), by a MQ Fellowship and by NWO ALW open (ALWOP.179). The authors declare no competing financial interests.

References

Anderson, T., Jenkins, I., Brooks, D., & Hawken, M. (1994). Cortical control of saccades and fixation in man. *Brain*. Retrieved from http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.458.5181&rep=rep1&type=pdf

Baars, B. J. (2005). Global workspace theory of consciousness: toward a cognitive neuroscience of human experience. *The Boundaries of Consciousness: Neurobiology and Neuropathology*, *150*, 45–53. Retrieved from http://books.google.com/books?hl=en&lr=&id=muaMA6PeSLcC&oi=fnd&pg=PA45&dq=global+workspace+Baars&ots=LGu0ZUWHQG&sig=Z4hRg-_YHLEnuP86jEvFU9VdQzE

Blondel, V. D., Guillaume, J.-L., Lambiotte, R., & Lefebvre, E. (2008). Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment*, *2008*(10), P10008. Retrieved from http://iopscience.iop.org/1742-5468/2008/10/P10008

Braga, R. M., Sharp, D. J., Leeson, C., Wise, R. J. S., & Leech, R. (2013). Echoes of the brain within default mode, association, and heteromodal cortices. *The Journal of Neuroscience*, *33*(35), 14031–14039. https://doi.org/10.1523/JNEUROSCI.0570-13.2013

Bush, G., Luu, P., & Posner, M. I. (2000). Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Sciences*, *4*(6), 215–222. https://doi.org/10.1016/S1364-6613(00)01483-2

Carlson, S., Martinkauppi, S., Rämä, P., Salli, E., Korvenoja, A., & Aronen, H. J. (1998). Distribution of cortical activation during visuospatial n-back tasks as revealed by functional magnetic resonance imaging. *Cerebral Cortex*, *8*(8), 743–752. https://doi.org/10.1093/cercor/8.8.743

Colizza, V., Flammini, A., Serrano, M. A., & Vespignani, A. (2006). Detecting rich-club ordering in complex networks. *Nature Physics*, *2*(2), 110–115. Retrieved from http://www.nature.com/nphys/journal/vaop/ncurrent/full/nphys209.html

Daianu, M., Jahanshad, N., Nir, T. M., Jack, C. R., Weiner, M. W., Bernstein, M. A., & Thompson, P. M. (2015). Rich club analysis in the Alzheimer's disease connectome reveals a relatively undisturbed structural core network. *Human Brain Mapping*, *36*(8), 3087–3103. https://doi.org/10.1002/hbm.22830

de Reus, M. A., & van den Heuvel, M. P. (2013). Estimating false positives and negatives in brain networks. *NeuroImage*, *70*, 402–409. https://doi.org/10.1016/j.neuroimage.2012.12.066

de Reus, M. A., & van den Heuvel, M. P. (2014). Simulated rich club lesioning in brain networks: a scaffold for communication and integration? *Frontiers in Human Neuroscience*, *8*. https://doi.org/10.3389/fnhum.2014.00647

Deco, G., Jirsa, V. K., & McIntosh, A. R. (2011). Emerging concepts for the dynamical organization of resting-state activity in the brain. *Nature Reviews. Neuroscience*, *12*(1), 43–56. https://doi.org/10.1038/nrn2961

Deco, G., Ponce-Alvarez, A., Mantini, D., Romani, G. L., Hagmann, P., & Corbetta, M. (2013). Resting-state functional connectivity emerges from structurally and dynamically shaped slow linear fluctuations. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *33*(27), 11239–52. https://doi.org/10.1523/JNEUROSCI.1091-13.2013

Deco, G., Senden, M., & Jirsa, V. (2012). How anatomy shapes dynamics: a semi-analytical study of the brain at rest by a simple spin model. *Frontiers in Computational Neuroscience*, *6*, 68. https://doi.org/10.3389/fncom.2012.00068

Deco, G., Van Hartevelt, T. J., Fernandes, H. M., Stevner, A., & Kringelbach, M. L. (2017). The most relevant human brain regions for functional connectivity: Evidence for a dynamical workspace of binding nodes from whole-brain computational modelling. *NeuroImage*, *146*, 197–210. https://doi.org/10.1016/j.neuroimage.2016.10.047

Dehaene, S., & Naccache, L. (2001). Towards a cognitive neuroscience of consciousness: basic evidence and a workspace framework. *Cognition*, *79*(1), 1–37. Retrieved from http://www.sciencedirect.com/science/article/pii/S0010027700001232

Dennis, E. L., Jahanshad, N., Toga, A. W., McMahon, K. L., De Zubicaray, G. I., Hickie, I., … Thompson, P. M. (2013). Development of the "rich club" in brain connectivity networks from 438 adolescents & adults aged 12 to 30. In *Proceedings - International Symposium on Biomedical Imaging* (pp. 624–627). NIH Public Access. https://doi.org/10.1109/ISBI.2013.6556552

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., … Killiany, R. J. (2006). An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*, *31*(3), 968–80. https://doi.org/10.1016/j.neuroimage.2006.01.021

Eriksen, B. A., & Eriksen, C. W. (1974). Effects of noise letters upon the identification of a target letter in a nonsearch task. *Perception & Psychophysics*, *16*(1), 143–149. Retrieved from http://link.springer.com/article/10.3758/BF03203267

Fink, G. R., Frackowiak, R. S. J., Pietrzyk, U., & Passingham, R. E. (1997). Multiple Nonprimary Motor Areas in the Human Cortex. *Journal of Neurophysiology*, *77*(4). Retrieved from http://jn.physiology.org/content/77/4/2164.short

Flowers, K. A., & Robertson, C. (1985). The effect of Parkinson's disease on the ability to maintain a mental set. *Journal of Neurology, Neurosurgery & Psychiatry*, *48*(6), 517–529. Retrieved from http://jnnp.bmj.com/content/48/6/517.short

Forstmann, B. U., van den Wildenberg, W. P. M., & Ridderinkhof, K. R. (2008). Neural Mechanisms, Temporal Dynamics, and Individual Differences in Interference Control. *Journal of Cognitive Neuroscience*, *20*(10), 1854–1865. https://doi.org/10.1162/jocn.2008.20122

Frässle, S., Lomakina, E. I., Razi, A., Friston, K. J., Buhmann, J. M., & Stephan, K. E. (2017). Regression DCM for fMRI. *NeuroImage*, *155*, 406–421. https://doi.org/10.1016/j.neuroimage.2017.02.090

Friston, K. J. (2011). Functional and Effective Connectivity: A Review. *Brain Connectivity*, *1*(1), 13–36. https://doi.org/10.1089/brain.2011.0008

Friston, K. J., Kahan, J., Biswal, B., & Razi, A. (2014). A DCM for resting state fMRI. *NeuroImage*, *94*, 396–407. https://doi.org/10.1016/j.neuroimage.2013.12.009

Gabrieli, J. D., Poldrack, R. A., & Desmond, J. E. (1998). The role of left prefrontal cortex in language and memory. *Proceedings of the National Academy of Sciences of the United States of America*, *95*(3), 906–13. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/9448258

Ghosh, A., Rho, Y., McIntosh, A. R., K\ötter, R., Jirsa, V. K., K\textbackslashötter, R., … Jirsa, V. K. (2008). Noise during rest enables the exploration of the brain's dynamic repertoire. *PLoS Comput Biol*, *4*(10), e1000196.

Gilson, M., Deco, G., Friston, K., Hagmann, P., Mantini, D., Betti, V., … Corbetta, M. (2017). Effective connectivity inferred from fMRI transition dynamics during movie viewing points to a balanced reconfiguration of cortical interactions. *bioRxiv*. Retrieved from http://www.biorxiv.org/content/early/2017/02/20/110015

Gilson, M., Moreno-Bote, R., Ponce-Alvarez, A., Ritter, P., Deco, G., & Grady, C. (2016). Estimation of Directed Effective Connectivity from fMRI Functional Connectivity Hints at Asymmetries of Cortical Connectome. *PLOS Computational Biology*, *12*(3), e1004762. https://doi.org/10.1371/journal.pcbi.1004762

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., … others. (2013). The minimal preprocessing pipelines for the Human Connectome Project. *Neuroimage*, *80*, 105–124. Retrieved from http://www.sciencedirect.com/science/article/pii/S1053811913005053

Goebel, R., Roebroeck, A., Kim, D.-S., & Formisano, E. (2003). Investigating directed cortical interactions in time-resolved fMRI data using vector autoregressive modeling and Granger causality mapping. *Magnetic Resonance Imaging*, *21*(10), 1251–1261. https://doi.org/10.1016/j.mri.2003.08.026

Grill-Spector, K., Kourtzi, Z., & Kanwisher, N. (2001). The lateral occipital complex and its role in object recognition. *Vision Research*, *41*(10–11), 1409–22. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/11322983

Gu, S., Pasqualetti, F., Cieslak, M., Telesford, Q. K., Alfred, B. Y., Kahn, A. E., … others. (2015). Controllability of structural brain networks. *Nature Communications*, *6*. Retrieved from http://www.nature.com/ncomms/2015/151001/ncomms9414/full/ncomms9414.html

Hampson, M., Driesen, N. R., Skudlarski, P., Gore, J. C., & Constable, R. T. (2006). Brain connectivity related to working memory performance. *Journal of Neuroscience*, *26*(51), 13338.

Haxby, J., Gobbini, M., Furey, M., & Ishai, A. (2001). Distributed and overlapping representations of faces and objects in ventral temporal cortex. Retrieved from http://science.sciencemag.org/content/293/5539/2425.short

He, B. J. (2011). Scale-Free Properties of the Functional Magnetic Resonance Imaging Signal during Rest and Task. *Journal of Neuroscience*, *31*(39). Retrieved from http://www.jneurosci.org/content/31/39/13786.short

Honey, C. J., Sporns, O., Cammoun, L., Gigandet, X., Thiran, J. P., Meuli, R., & Hagmann, P. (2009). Predicting human resting-state functional connectivity from structural connectivity. *Proceedings of the National Academy of Sciences*, *106*(6), 2035.

Ishai, A., Ungerleider, L., & Martin, A. (1999). Distributed representation of objects in the human ventral visual pathway. *Proceedings of the*. Retrieved from http://www.pnas.org/content/96/16/9379.short

Kirchner, W. K. (1958). Age differences in short-term retention of rapidly changing information. *Journal of Experimental Psychology*, *55*(4), 352. Retrieved from http://psycnet.apa.org/journals/xge/55/4/352/

Klein, A., & Tourville, J. (2012). 101 Labeled Brain Images and a Consistent Human Cortical Labeling Protocol. *Frontiers in Neuroscience*, *6*, 171. https://doi.org/10.3389/fnins.2012.00171

Kringelbach, M. L. (2005). The human orbitofrontal cortex: linking reward to hedonic experience. *Nature Reviews Neuroscience*, *6*(9), 691–702.

https://doi.org/10.1038/nrn1747

Leech, R., Braga, R., & Sharp, D. J. (2012). Echoes of the brain within the posterior cingulate cortex. *The Journal of Neuroscience*, *32*(1), 215–222. https://doi.org/10.1523/JNEUROSCI.3689-11.2012

Leung, H.-C., Gore, J. C., & Goldman-Rakic, P. S. (2002). Sustained Mnemonic Response in the Human Middle Frontal Gyrus during On-Line Storage of Spatial Memoranda. *Journal of Cognitive Neuroscience*, *14*(4), 659–671. https://doi.org/10.1162/08989290260045882

Liu, Y.-Y., Slotine, J.-J., & Barabási, A.-L. (2011). Controllability of complex networks. *Nature*, *473*(7346), 167–173. https://doi.org/10.1038/nature10011

Machielsen, W., Rombouts, S., & Barkhof, F. (2000). FMRI of visual encoding: reproducibility of activation. *Human Brain*. Retrieved from http://onlinelibrary.wiley.com/doi/10.1002/(SICI)1097-0193(200003)9:3%3C156::AID-HBM4%3E3.0.CO;2-Q/full

Mangun, G., Buonocore, M., & Girelli, M. (1998). ERP and fMRI measures of visual spatial selective attention. *Human Brain*. Retrieved from https://www.researchgate.net/profile/George_Mangun/publication/13498445_ERP_and_fMRI_Measures_of_Visual_Spatial_Selective_Attention/links/02e7e539358ba7b050000000.pdf

Maslov, S., & Sneppen, K. (2002). Specificity and stability in topology of protein networks. *Science*, *296*(5569), 910–913. https://doi.org/10.1126/science.1065103

Messé, A., Rudrauf, D., Benali, H., Marrelec, G., & Honey, C. (2014). Relating Structure and Function in the Human Brain: Relative Contributions of Anatomy, Stationary Dynamics, and Non-stationarities. *PLoS Computational Biology*, *10*(3), e1003530. https://doi.org/10.1371/journal.pcbi.1003530

Mitra, A., Snyder, A. Z., Tagliazucchi, E., Laufs, H., Raichle, M. E., Zwarts, M., … Milham, M. (2015). Propagated infra-slow intrinsic brain activity reorganizes across wake and slow wave sleep. *eLife*, *4*, 5263–5264. https://doi.org/10.7554/eLife.10781

Newman, M. E. J. (2006). Modularity and community structure in networks. *Proceedings of the National Academy of Sciences*, *103*(23), 8577–8582. Retrieved from http://www.pnas.org/content/103/23/8577.short

Okada, T., Tanaka, S., Nakai, T., Nishizawa, S., & Inui, T. (2000). Naming of animals and tools: a functional magnetic resonance imaging study of categorical differences in the human brain areas commonly used for naming visually. *Neuroscience*. Retrieved from http://www.sciencedirect.com/science/article/pii/S0304394000016128

Patel, A. X., & Bullmore, E. T. (2015). A wavelet-based estimator of the degrees of freedom in denoised fMRI time series for probabilistic testing of functional connectivity and brain graphs. *NeuroImage*. Retrieved from http://www.sciencedirect.com/science/article/pii/S1053811915003523

Price, C. J. (1998). The functional anatomy of word comprehension and production. *Trends in Cognitive Sciences*, *2*(8), 281–288. https://doi.org/10.1016/S1364-6613(98)01201-7

Romme, I. A. C., de Reus, M. A., Ophoff, R. A., Kahn, R. S., & van den Heuvel, M. P. (2017). Connectome Disconnectivity and Cortical Gene Expression in Patients With Schizophrenia. *Biological Psychiatry*, *81*(6), 495–502. https://doi.org/10.1016/j.biopsych.2016.07.012

Schoenbaum, G., Takahashi, Y., Liu, T.-L., & McDannald, M. A. (2011). Does the

orbitofrontal cortex signal value? *Annals of the New York Academy of Sciences*, *1239*(1), 87–99. https://doi.org/10.1111/j.1749-6632.2011.06210.x

Senden, M., Deco, G., De Reus, M. A. M. A., Goebel, R., & van den Heuvel, M. P. (2014). Rich club organization supports a diverse set of functional network configurations. *NeuroImage*, *96*, 174–182. https://doi.org/10.1016/j.neuroimage.2014.03.066

Senden, M., Reuter, N., van den Heuvel, M. P., Goebel, R., & Deco, G. (2017). Cortical rich club regions can organize state-dependent functional network formation by engaging in oscillatory behavior. *NeuroImage*, *146*, 561–574. https://doi.org/10.1016/j.neuroimage.2016.10.044

Shanahan, M. (2012). The brain's connective core and its role in animal cognition. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, *367*(1603), 2704–2714. https://doi.org/10.1098/rstb.2012.0128

Shepard, R. N., & Metzler, J. (1971). Mental rotation of three-dimensional objects. Retrieved from https://books.google.nl/books?hl=en&lr=&id=DcJAqyCK6T8C&oi=fnd&pg=PA254&dq=shepard&ots=q3xg0-lbIF&sig=xgM8A3yyvBB0dTjkgLxn-CQjzSo

Silani, G., Lamm, C., Ruff, C., & Singer, T. (2013). Right supramarginal gyrus is crucial to overcome emotional egocentricity bias in social judgments. *Journal of Neuroscience*. Retrieved from http://www.jneurosci.org/content/33/39/15466.short

Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., … others. (2009). Correspondence of the brain's functional architecture during activation and rest. *Proceedings of the National Academy of Sciences*, *106*(31), 13040–13045. Retrieved from http://www.pnas.org/content/106/31/13040.short

Thompson-Schill, S. L., D'Esposito, M., & Kan, I. P. (1999). Effects of Repetition and Competition on Activity in Left Prefrontal Cortex during Word Generation. *Neuron*, *23*(3), 513–522. https://doi.org/10.1016/S0896-6273(00)80804-1

van den Heuvel, M. P., de Reus, M. A., Feldman Barrett, L., Scholtens, L. H., Coopmans, F. M. T., Schmidt, R., … Li, L. (2015). Comparison of diffusion tractography and tract-tracing measures of connectivity strength in rhesus macaque connectome. *Human Brain Mapping*, *36*(8), 3064–3075. https://doi.org/10.1002/hbm.22828

van den Heuvel, M. P., Kahn, R. S., Goñi, J., & Sporns, O. (2012). High-cost, high-capacity backbone for global brain communication. *Proceedings of the National Academy of Sciences*, *109*(28), 11372–11377. Retrieved from http://www.pnas.org/content/109/28/11372.short

van den Heuvel, M. P., Scholtens, L. H., Feldman Barrett, L., Hilgetag, C. C., & de Reus, M. A. (2015). Bridging Cytoarchitectonics and Connectomics in Human Cerebral Cortex. *Journal of Neuroscience*, *35*(41), 13943–13948. https://doi.org/10.1523/JNEUROSCI.2630-15.2015

van den Heuvel, M. P., & Sporns, O. (2011). Rich-club organization of the human connectome. *The Journal of Neuroscience*, *31*(44), 15775–15786. Retrieved from http://www.jneurosci.org/content/31/44/15775.short

van den Heuvel, M. P., Sporns, O., Collin, G., Scheewe, T., Mandl, R. C. W., Cahn, W., … Kahn, R. S. (2013). Abnormal Rich Club Organization and Functional Brain Dynamics in Schizophrenia. *JAMA Psychiatry*, *70*(8), 783. https://doi.org/10.1001/jamapsychiatry.2013.1328

Van Essen, D. C., Ugurbil, K., Auerbach, E., Barch, D., Behrens, T. E. J., Bucholz, R., …

others. (2012). The Human Connectome Project: a data acquisition perspective. *Neuroimage*, *62*(4), 2222–2231. Retrieved from http://www.sciencedirect.com/science/article/pii/S1053811912001954

Vanni, S., Tanskanen, T., Seppä, M., Uutela, K., & Hari, R. (2001). Coinciding early activation of the human primary visual cortex and anteromedial cuneus. *Proceedings of the National Academy of Sciences of the United States of America*, *98*(5), 2776–80. https://doi.org/10.1073/pnas.041600898

Yeh, F.-C., Wedeen, V. J., & Tseng, W.-Y. I. (2010). Generalized-sampling imaging. *Medical Imaging, IEEE Transactions on*, *29*(9), 1626–1635. Retrieved from http://ieeexplore.ieee.org/xpls/abs_all.jsp?arnumber=5432996

Zamora-López, G., Zhou, C., & Kurths, J. (2009). Graph analysis of cortical networks reveals complex anatomical communication substrate. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, *19*(1), 15117. Retrieved from http://scitation.aip.org/content/aip/journal/chaos/19/1/10.1063/1.3089559

Zhou, S., & Mondragon, R. J. (2004). The rich-club phenomenon in the Internet topology. *IEEE Communications Letters*, *8*(3), 180–182. https://doi.org/10.1109/LCOMM.2004.823426