Title: Cued reactivation during slow-wave sleep induces connectivity changes related to memory stabilization.

All authors: Ruud M.W.J. Berkers* (1,2), Matthias Ekman* (3), Eelco V. van Dongen (1), Atsuko Takashima (3,4), Marcus Barth (3,5), Ken A. Paller (6), Guillen Fernandez (1).
      *Shared first authors

Affiliations:
Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Centre, Nijmegen, the Netherlands
Max Planck Institute for Human Cognitive & Brain Sciences, Leipzig, Germany
Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands
Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands
Centre for Advanced Imaging, The University of Queensland, Brisbane, Australia
Northwestern University, Evanston, Illinois, United States

Author contributions: R.M.W.J.B., M.E., E.V.vD., A.T., M.B., K.A.P. & G.F. conceived and designed the study, E.V.vD., A.T. & M.B. performed research, M.E. contributed analytic tools, R.M.W.J.B., M.E. & E.V.vD. analyzed data, R.M.W.J.B. wrote the paper, and all authors provided feedback to the final manuscript.

Corresponding author: Ruud M.W.J. Berkers, MSc.
                                Max Planck Institute for Human Cognitive & Brain Sciences
                                Postfach 500355, D-04303 Leipzig, Germany
                                Email: berkers@cbs.mpg.de

Number of pages: 33
Number of figures: 5
Number of tables: 0
Number of words
   - **Abstract: 219**
   - **Introduction: 1165**
   - **Discussion: 1819**

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgement: R.M.W.J.B., E.V.vD. and G.F were supported by a grant from the European Research Council ("Neuroschema" ERC R0001075). G.F. was also supported by a grant from the NWO ("Language in Interaction"). E.V.v.D (400-06-110) and A.T. (451-06-006) were supported by grants of the Dutch Organization for Scientific Research (NWO).

Abstract

Memory reprocessing following acquisition enhances memory consolidation. Specifically, neural activity during encoding is thought to be 'replayed' during subsequent slow-wave sleep (SWS). This natural tendency of memory replay can be induced by external cueing, known as "targeted memory reactivation". Here, we analyzed data from a published study (van Dongen et al. 2012), where auditory cues reactivated learned visual object-location memories during SWS. Memory replay during sleep presumably involves a shift in connectivity across the brain. Therefore, we characterized the effects of memory reactivation on brain network connectivity using graph-theory. We found that cue presentation during SWS introduced dynamic changes in network integration of the occipital cortex, a visual region that was also active during the object retrieval task. Importantly, enhanced network integration of the occipital cortex showed a behavioural benefit and predicted overnight memory stabilization. Furthermore, occipital cortex displayed enhanced connectivity with mnemonic regions, namely the hippocampus, parahippocampal gyrus, thalamus and medial prefrontal cortex during cue versus control sound presentation. Finally, network integration of early occipital cortex during cueing in SWS was related to increased activation of the bilateral parahippocampal gyrus, a region involved in coding for spatial associative information, at the post-sleep test. Together, these results support a neural mechanism where cue-induced replay during sleep promotes memory consolidation by increased integration of task-relevant perceptual regions with mnemonic regions.

Introduction

The neural representations of recently acquired declarative memories are thought to be

selectively reactivated during ensuing rest periods, particularly during slow-wave sleep

(Diekelmann and Born 2010; Diekelmann et al. 2011). For instance, hippocampal place cells in

rats that co-activated during active behavior were found to be reactivated in concert during

ensuing slow-wave sleep (Wilson and McNaughton 1994). Later studies found that this

coordinated reactivation, also dubbed replay, extended to the perceptual cortex involved in the

initial processing of the stimulus, such as the visual cortex (Ji and Wilson 2007). Furthermore,

evidence of neural replay has also been found in the medial prefrontal cortex (Euston et al.

2007; Peyrache et al. 2009; Genzel and Battaglia 2017). These results demonstrate the

occurrence of neural memory replay in rodents, and have been shown to be behaviorally

relevant for the retention of associative memories (Dupret et al. 2010).

Systems-level consolidation theory posits that neural memory traces are initially hippocampus-

dependent, and subsequently integrate into neocortical networks (Squire et al. 1984;

McClelland et al. 1995; Nadel and Moscovitch 1997; Frankland and Bontempi 2005). The neural

trace is presumed to initially involve the hippocampus, which acts like a hub binding elements

into a coherent memory trace, and the cortical regions that were involved in the initial

processing of the stimulus at encoding (Nyberg et al. 2000; Polyn et al. 2005; Danker and

Anderson 2010). During sleep, especially when the brain is in a highly synchronous state (i.e.

slow-wave sleep stage), specific tightly coupled oscillatory patterns are observed, including

slow-oscillations, thalamic spindles, and hippocampal ripples (Siapas and Wilson 1998; Mölle et

al. 2002, 2006; Staresina et al. 2015) that have been related to memory consolidation (Schabus

et al. 2004; Marshall et al. 2006; Axmacher et al. 2008; Girardeau et al. 2009; Cox et al. 2012).

Thus, it is plausible to posit that memory reactivation during sleep facilitates information

exchange between neocortex and hippocampus, enabling the strengthening of connections

between these areas as well as amongst neocortical modules, thereby contributing to systems-

level memory consolidation (Marshall and Born 2007).

Neural memory reactivation is difficult to observe in humans, as its timing is generally

unknown. However, reactivation can also be induced using cues associated with information

learned previously. For example, presenting an odor during slow-wave sleep induces activation

of the hippocampus and stabilizes associated memory traces (Rasch et al. 2007; Diekelmann et

al. 2010, 2011). Furthermore, whereas studies have found that audio-visual-spatial associations

can be stabilized in a targeted manner using specific auditory cues (Rudoy et al. 2009; Creery et

al. 2015), these cueing effects can also affect all associations acquired within the same learning

context (Oudiette et al. 2013). Van Dongen and colleagues (2012) performed targeted memory

reactivation in the magnetic resonance (MR) scanner using auditory cues that were paired with

specific object-location associations. Even though no consistent effect of cueing on memory

stabilization was found, individual differences in the effect of cueing positively correlated with

activity in the hippocampus, thalamus and cerebellum during slow-wave sleep. Given repeated

earlier findings of memory improvement from auditory cues presented during slow-wave sleep

(Oudiette et al. 2013; Ong et al. 2016; Schouten et al. 2017), it is likely that the noisy scanner

environment contributed to excessive variability here, with some individuals effectively tuning

out all auditory input, or alternatively that cueing affected all associations acquired within the

same learning context (Oudiette et al. 2013).

Here, we re-analyzed data acquired by van Dongen and colleagues (2012) using a principled

graph theoretical analysis approach. Triggering reactivation of memory traces with auditory

cues, could induce a process akin to spontaneous replay, which likely involves whole-brain

connectivity changes facilitated by synchronous brain states during SWS. Conventional analysis

methods are often unable to capture the coordinated whole-brain connectivity changes

expected during memory replay. Thus, a network perspective can be leveraged using methods

developed within the framework of graph theory (Bullmore and Sporns 2009). The participation

coefficient is a metric that captures the extent to which a region integrates and distributes

information as a result of the number and positioning of their contacts in the network (Power

et al. 2013). Specifically, it quantifies network integration by looking at the importance of a

given node for interactions between subnetworks. The participation coefficient can be used to

index the integration of a given voxel (node) in the wider brain (network) during cued

reactivation of visuospatial associations during SWS.

If a cue induces memory reactivation akin to replay, and this indeed plays a role in memory

consolidation involving a hippocampal-neocortical trace shift, then one should expect a

coordinated neural activation of multiple regions, including the sensory features of the cued

memory (in this case visual object features in the occipital cortex and the spatial layout in the

parahippocampal gyrus), as well as the key mnemonic regions (such as the hippocampus,

thalamus, and medial prefrontal cortex). This increased integration of these and other regions

would be expected to be beneficial for memory stabilization across sleep and result in an

increased involvement of the neocortex in the memory trace.

Materials and Methods

The Materials and Methods have been described extensively in the original report (van Dongen

et al. 2012). Here, we summarize relevant details about the experimental setup, supplemented

with details about the data analysis performed for this report.

Participants

This re-analysis includes data from all the 22 participants included in the original report (for

complete description, see van Dongen et al., 2012). Briefly, 56 participants were initially

recruited from the online research participant system of the Radboud University Nijmegen (age

range 18–27; 14 males). Twenty-two participants in the sample reached a sufficient amount of

slow-wave sleep in the scanner to allow at least 80 % of the cueing protocol to be completed

whilst in the slow-wave sleep stage. Furthermore, these participants did not display micro-

arousals in response to sound presentations, nor did they report explicit knowledge of the

sounds having been presented while they slept. The experiment was conducted in accordance

with national legislation for the protection of human volunteers in nonclinical research settings

and the Helsinki Declaration and approved by the local ethics committee. Participants were

compensated for participation with course credits or a monetary fee.

Procedures

The experiment started between 7 and 8 PM, and participants went to sleep between 10 PM

and midnight (see figure 1). Participants were placed in the MR scanner, and the sound volume

was individually calibrated to a level at which the participant could distinguish individual sounds

while the scanner was operating. Participants learned 50 object-location associations, each

object-location pair being presented together with a unique sound associated with the object,

inside the scanner. When all associations were learned to criterion (all objects placed within 4

cm from the correct location on two subsequent rounds), a pre-test was performed on all

associations (Test 1). Next, participants were prepared outside the scanner for

polysomnographic recordings, and placed in the scanner again for a 2 hr. rest/sleep period,

while EEG was monitored online and fMRI data was continuously acquired. When participants

had entered stable slow-wave sleep for a period of time as visible on the ongoing

polysomnographic EEG, sounds were presented to the participants using an MR-compatible

headphone. Specifically, half of the sounds that were associated with specific object-location

associations during learning were presented as 'cue sounds' during sleep (25 sounds presented

twice, for a total of 50 cue sound presentations). Moreover, control sounds were presented

that had not been previously associated with learning materials (5 sounds presented five times,

for a total of 25 control sound presentations). After a 2 hr sleep, participants were awoken

before they performed the post-test with all object-location associations (Test 2).

Object-location test

The object-location task was performed in three stages: learning, Test 1 and Test 2. Here, we

focus on the behavioral data from Test 1 and Test 2 for further analysis. Participants learned

the location of 50 object pictures. For each participant, 50 objects were randomly assigned to

50 screen locations (screen size: 47 × 35 cm, resolution: 1024 × 768, viewing distance: 60 cm).

In the learning phase, participants first passively viewed all 50 objects in their respective screen

locations (duration: 3s presentation followed by a 1s inter-stimulus interval), paired with

hearing the object related sound (e.g. a cat's 'meow' when the object was a cat, sound duration:

500 ms). Next, the learning phase continued with several iterative rounds of active learning,

where an object was presented at the center of the screen simultaneously with the auditory

stimulus. Participants were then required to place the object in its original location using an

MR-compatible joystick, and to press a button to confirm the object placement (self-paced

timing). Feedback was given in every round of learning by displaying the object in the correct

location for 3s. Objects were presented in a random order, and with every new round those

objects were excluded that had been placed within 4 cm from the correct location on the two

preceding rounds. Upon reaching criterion performance on all objects, participants performed

the pre-sleep test (Test 1) for all objects without feedback. An identical post-sleep test was

performed after sleep (Test 2). During tests, object placement was followed by an inter-

stimulus period of fixation (duration jittered between 3 and 5s).

MRI Data Acquisition

Participants were scanned using a reduced-noise EPI sequence with sinusoidal gradients to

avoid acoustic resonances of the scanner (Schmitter et al. 2008). Functional (T2*) images were

acquired with whole-brain coverage (28 axial slices, ascending slice acquisition, TR = 2,511 ms,

TE = 38 ms, 90° flip angle, matrix = 64 x 64, bandwidth = 1,502 Hz per voxel, slice thickness = 3.5

mm, slice gap = 15%, field of view (FOV) = 244 mm). Structural (T1) images were acquired with a

MPRAGE sequence (176 sagittal slices, TR = 2,250 ms, TE = 2.95 ms, 15° flip angle, matrix = 256

x 256, slice thickness = 1.0 mm, FOV = 256 mm).

MRI Data Preprocessing

The fMRI data acquired during Test 1, the sleeping period and Test 2 were preprocessed using

standard routines implemented in SPM8 (www.fil.ion.ucl.ac.uk). The first five volumes of each

functional EPI run were discarded, and an outlier algorithm was used to check for corrupted

slices from each image and replaced using between-volume interpolation. The functional

images were then realigned, and coregistered to the structural image, spatially normalized to

MNI-space (resampled at voxel size 2 x 2 x 2 mm), and smoothed using a Gaussian kernel (8 mm

FWHM). The structural images were segmented into gray matter, white matter, cerebrospinal

fluid, and residual compartments (outside brain and skull) using the unified segmentation

algorithm as implemented in SPM8. These compartments were used as masks to extract the

mean intensity level across the whole time-series, and entered as compartment regressors to

account for effects related to non-specific signal fluctuations.

Network analysis during cueing

We employed a graph theoretic framework (Bullmore and Sporns 2009; Ekman et al. 2012) to

analyze connectivity patterns during the presentation of sound cues in the scanned sleep

period. The aim was to measure dynamic changes in the whole-brain network properties during

memory reactivation, and specifically changes in integration of certain brain regions with the

whole brain network. The metric of interest was defined as the participation coefficient

(Guimera et al. 2005; Power et al. 2013; Backus et al. 2016), which quantifies for each node (i.e.

voxel) the diversity of its inter-modular connections (Rubinov and Sporns 2010), specified as the

amount of connections with nodes in other submodules relative to the total amount of

connections. It therefore is a measure of the importance of a given node for global inter-

modular integration across the brain.

A high-pass filter with a cutoff period of 128s was applied to the functional time series. Sound

presentation during sleep was modelled using a separate regressor for each sound presentation

(to obtain trial-specific beta-estimates for the 50 cue sound presentations and 25 control sound

presentations) with a duration of 5 seconds each (the duration of the individual sound

presentations). The design matrix also included six regressors of no interest to account for head

movement, and the three compartment regressors to account for non-specific signal

fluctuations. The resulting beta estimates were concatenated separately for cue sound and

control sound trials. This procedure resulted in two separate beta time-series (i.e., cue sounds

and control sounds) for all voxels in the brain. Next, voxel-wise correlation coefficients of beta

time-series were computed to quantify pairwise functional connectivity (Rissman et al. 2004)

for each condition separately (i.e., cue sounds and control sounds). The connectivity matrices

were thresholded by preserving only significant connections ($p < 0.05$ FDR corrected) and

setting all other connections, as well as all negative correlations to zero. Following the

procedure used by Power and co-workers (2011), modular network structure was first derived

across all sound presentations using a connectivity matrix based on the regions defined by the

AAL (Automated Anatomical Labeling) atlas (Tzourio-Mazoyer et al. 2002). Here, beta time-

series were averaged across voxels of the region, and a 116 x 116 region-by-region connectivity

matrix was constructed. After thresholding this connectivity matrix (edges > 0, $p < 0.05$ FDR

corrected), the matrix was parcellated into subnetworks using modularity detection according

to the Louvain method (Blondel et al. 2008). Next, each voxel in the whole-brain voxel-wise

connectivity matrix was assigned to a module, allowing the calculation of their participation

coefficient ranging from 0 (provincial hub: connections are only present within its module) and

1 (connector hub: connections are only present with other modules; Power et al., 2013). The

resulting participant coefficient maps for cue sounds and control sounds were contrasted on an

individual subject level and this contrast was subsequently tested at the second level using a

one-sample t-test.

Psychophysiological interaction (PPI) analysis during cueing

The previous analysis determined which of the regions increased in global inter-modular

network integration. To determine which regions in other modules of the brain were connected

with the 'hub' (the region found to display increased network integration during cueing), we

performed a follow-up seed-based functional connectivity analysis. This psychophysiological

interaction (PPI) analysis served to determine which brain regions were more connected with

the hub during the presentation of cue versus control sounds. Therefore, we extracted the

physiological signal time series from the seed region found in the network analysis (cluster-

forming threshold at Z > 2.33) to serve as a physiological regressor. This was entered into a

general linear model (GLM) along with the psychological variable (cue versus control sounds)

and the psychophysiological interaction regressor. Small-volume correction was used for the

bilateral parahippocampal gyri, as this region was activated by the object-location retrieval task

(van Dongen et al. 2012), and thalamus and hippocampus, as these are smaller structures

hypothesized to be involved in memory reactivation. Here, anatomical masks implemented in

the IBASPM 71 Atlas of the WFU Pick Atlas Tool in SPM were used (Iturria-Medina et al. 2007).

Activity analysis during pre- and post-sleep test

We next assessed whether network integration of visual cortex during cueing in slow-wave

sleep results in increased involvement of cortical associative regions at retrieval. For this

reason, we contrasted the neural activations observed during the pre-sleep and post-sleep test,

and related this contrast to the participation coefficient of the occipital cortex as observed in

the graph theory analysis (see Results-section). The fMRI data for Test 1 and Test 2 were

analyzed in one GLM. Similar to van Dongen and colleagues (2012), this model included

regressors for cued and uncued associations, which were modeled as delta functions to be

convolved with the canonical hemodynamic response function (HRF), along with temporal

derivatives provided by SPM8. The design matrix included six regressors of no interest per test

session to account for head movement using the realignment parameters, as well as

compartment regressors accounting for non-specific signal fluctuations. Furthermore, a high-

pass filter with a cutoff period of 128s was implemented to remove low-frequency fluctuations

from the time series. Parameter images were generated on the first level (for each individual

participant) based on a contrast between all trials of Test 1 and all trials of Test 2, to assess

changes in retrieval activity from the pre- to the post-sleep test. We collapsed across cued and

uncued associations because network participation of the occipital cortex was not specifically

related to only the cued associations in the learned set (see 'Results'). These contrast images

were entered into a one-sample t-test on the second level, with the participation coefficient of

the occipital cortex during cue versus control sound presentation entered as a covariate. Small-

volume correction was used for the bilateral parahippocampal gyri, as this region was activated

by the object-location retrieval task and cue-sound presentation (van Dongen et al. 2012). Here,

an anatomical mask of the parahippocampal gyrus implemented in the IBASPM 71 Atlas of the

WFU Pick Atlas Tool in SPM was used (Iturria-Medina et al. 2007).

Results

Network changes following cueing

We probed changes in network dynamics in response to the presentation of cue sounds versus

control sounds. The presentation of cue sounds was hypothesized to induce reactivation of the

corresponding memory trace, which would presumably be reflected in increased integration of

areas that represent sensory features of the cued memory representation, e.g. occipital cortex.

Indeed, we found a specific increase in the participation coefficient of early visual (occipital)

cortex (peak MNI coordinates: x,y,z = [4, -70, 6], z_{21} = 3.12, cluster-forming threshold at Z>2.33,

corrected for the whole brain at p < 0.05 using Gaussian Random Field Theory), indicating that

occipital cortex displays an increased inter-modular connectivity when presented with cue

sounds during deep sleep. The region observed overlaps with regions that were activated in

response to the object-location task during Test 1 (see figure 2b). This overlap may be indicative

of replay of the original visuo-spatial neural memory trace in response to auditory cues

presented during sleep, in the absence of any external visual input.

To determine what brain regions this area in the occipital cortex preferentially connects to

when recruited in the network by the auditory cues, we performed a psychophysiological

connectivity analysis. The seed was defined as the region in cortex that displayed an increase in

participation coefficient during cueing (cluster-forming threshold at $z > 2.33$). The task contrast

was formed by the presentation of cue sounds versus control sounds. The occipital cortex was

found to be connected to the left hippocampus (peak MNI coordinates: x,y,z = [-32, -16, -18],

z_{21} = 3.36),   right hippocampus (peak MNI coordinates: x,y,z = [22, -2, -22], z_{21} = 2.98), left

parahippocampal gyrus (peak MNI coordinates: x,y,z = [-26, -22, -26], z_{21} = 3.10), left thalamus

(peak MNI coordinates: x,y,z = [-4, -2, -0], z_{21} = 3.11), and right thalamus (peak MNI coordinates:

x,y,z = [2, -16, -12], z_{21} = 2.67), and a medial prefrontal region extending from premotor cortex

to anterior cingulate cortex and into dorsomedial prefrontal cortex (see figure 3, all cluster-

forming threshold at $z > 2.33$, cluster-size corrected for the whole brain, or reduced search

regions based on predefined anatomical areas for the bilateral hippocampus, parahippocampal

gyrus and thalamus, thresholded at $p < 0.05$ using Gaussian Random Field Theory). Thus, during

cueing, the occipital cortex was connected to several regions where memory replay has been

shown to occur (hippocampus, medial prefrontal cortex), and that are involved in oscillatory

patterns of activity (spindles, ripples, slow-waves) that allow for systematic interactions

between brain regions (e.g., the hippocampus, medial prefrontal cortex and thalamus).

Network dynamics related to behavioral cueing effect

As expected, memory performance decreased on average (Test 1 performance: error = 2.74 ±

0.12 cm; Test 2 performance: error = 3.12 ± 0.14: Δerror = 0.37 ± 0.08; t_{21} = 4.48, p < 0.001),

both for cued (Δerror = -0.44 ± 0.11 cm, t_{21}(3.86, P = 0.001)), and uncued associations (Δerror =

−0.33 ± 0.11 cm, t_{21}(2.56, P = 0.018)). There was no effect of cueing on participants' memory

accuracy as tested after sleep ($F_{1,21}$ = 0.83; P = 0.374). Thus, there was forgetting across the

sleep/rest session, and this forgetting appeared to be unspecific across cued and uncued

object-location associations.

However, there was substantial variability in the effect of cueing on forgetting. Therefore,

different neural responses to cueing might explain individual differences in the effect of cueing.

Particularly, increased network participation of the occipital cortex could be an index of how

much the visual-spatial memory trace became reactivated in a given subject when auditory

cues were presented. Indeed, the increased participation coefficient during cue sound

presentation was found to be associated (see figure 2) with reduced overnight forgetting of

visual-spatial object locations (τ_{21} = -0.311, p = 0.045). This relationship was specifically present

for the visual-spatial locations that were associated with sound cues (τ_{21} = -0.338, p = 0.029),

but not for other locations (τ_{21} = -0.211, p = 0.174), although the difference in these correlations

did not reach significance (Williams-Hotelling test, t = 0.433, p = 0.670).

The distribution of the increase in participation coefficient followed a non-normal positively

skewed distribution, with about half of participants displaying only a minimal increase in

participation coefficient during cueing (PC < 0.1). Therefore, we performed a median split to

better characterize the group that experienced an increase in network participation of the

occipital cortex (PC+), versus the group with no increase (PC=). The PC= group displayed

significant forgetting (Δerror = 0.68 ± 0.32; t_{10} = 6.042, p < 0.001), for both cued (Δerror = 0.69 ±

0.41; t_{10} = 5.620, p < 0.001) and uncued (Δerror = 0.48 ± 0.44; t_{10} = 3.636, p = 0.005)

associations. In contrast, the PC+ group did not display any forgetting (Δerror = 0.16 ± 0.35; t_{10} =

1.551, p = 0.152), for both cued (Δerror = 0.20 ± 0.57; t_{10} = 1.186, p = 0.263) and uncued (Δerror

= 0.12 ± 0.61; t_{10} = 0.653, p = 0.529) associations. Note that we do not report the statistical test

to compare performance differences between the two groups, as this comparison would not be

statistically independent from the across-subject correlation reported earlier. This result

indicates that in the group where there was an increased network participation of occipital

cortex, overnight forgetting was reduced, but the effect was not specific to the cued items,

extending to other object-location associations encoded within the same spatial layout and

temporal learning context (see figure 4).

Overnight activity changes related to network integration during cueing

We then considered whether the increased network integration of the occipital cortex during

sleep had subsequent effects on neural activity when retrieving object-locations during the

post-sleep test. Here, we found that the participation coefficient was related to a positive

increase in retrieval-related activity from Test 1 to Test 2 in both the posterior left

parahippocampal gyrus (peak MNI coordinates: x,y,z = [-16, -32, -18], z_{21} = 4.14), and posterior

right parahippocampal gyrus (peak MNI coordinates: x,y,z = [16, -32, -18], z_{21} = 3.26, cluster-

forming threshold at Z>2.33, corrected for a reduced search volume based on anatomical masks

of the left and right parahippocampal gyrus at a threshold of p < 0.05 using Gaussian Random

Field Theory). Thus, increased network integration of the occipital cortex during cueing had a

subsequent effect on the neural regions activated during the retrieval of object-locations at

Test 2, thereby increasingly recruiting the bilateral parahippocampal gyrus.

Discussion

Here, we investigated the whole-brain neural network reorganization induced by cued

reactivation of memory traces during deep sleep, and its behavioral and neural effects on the

subsequent retrieval of those memories. For that purpose, we re-analyzed fMRI data reported

in study by van Dongen and colleagues (2012) using whole-brain connectivity measures. We

demonstrate that when participants were in a state of slow-wave sleep, absent of visual input,

and were cued with sounds associated with previously learned visuospatial information, the

occipital cortex displayed increased network integration as measured with the participation

coefficient. Furthermore, subjects who displayed an increase in cue-induced network

integration of the occipital cortex showed increased memory stabilization. This finding is

indicative of a process in which the sound cue induces a global reprocessing of the learning

episode, akin to memory replay, which is conducive to subsequent memory retention. Note

that while the occipital cortex is not directly related to mnemonic processes, previous studies

have shown that cue-triggered reactivation can lead to instantiations of previously experienced

episodes in this region (Xu et al. 2012; Ekman et al. 2017). It is likely that this reinstitution is

coordinated with other mnemonic regions. Indeed, various regions displayed increased

connectivity with the occipital cortex during the presentation of cue sounds: memory and

replay-related regions, such as the hippocampus, thalamus and medial prefrontal cortex; and

higher-order associative cortices implicated in the encoding of the memory trace, such as the

parahippocampal gyrus. These results are in line with a study by Ji & Wilson (2007) that shows

coordinated replay between the hippocampus and a perceptual region that was involved in the

initial processing of the stimulus (here: the occipital cortex). Cueing during slow-wave sleep also

showed an enduring neural effect during memory retrieval as measured during the post-sleep

retrieval test. Specifically, subjects with greater network integration of the occipital cortex

during SWS recruited the parahippocampal gyrus more at subsequent retrieval.

It should be noted that the network reorganization found during the presentation of cue

sounds was not specifically related to memory stabilization of those object-location

associations associated with the cues. The association of network participation with memory

stabilization was numerically larger for cued, versus uncued material, but the interaction was

not statistically significant. There could be various explanations for this unspecific effect. First,

the MRI scanner is a suboptimal environment for auditory presentation during sleep. Therefore,

it is conceivable that the acoustic stimuli presented during deep sleep in a noisy environment

effectively modulated ongoing brain processes in some individuals, but not others, contributing

to large inter-individual differences and lower power of demonstrating this effect than in other

studies conducted outside of the MR-scanner environment (Rudoy et al. 2009). Indeed, we

observed a non-normal positively skewed distribution of increase in participation coefficient

during cueing. There was an overall effect on memory retention for participants who showed

increased participation coefficient during cueing – but there did not appear to be a cue-specific

memory benefit even when only considering this sub group. It is also possible that the study

was underpowered to observe an interaction effect, and that a larger sample would have

rendered a significant interaction. Furthermore, it is possible that the cueing had a non-specific

effect on all material acquired within the same temporal and spatial learning context (as also

suggested by the results reported in Oudiette et al., 2013). Indeed, all object-location

associations were acquired within the same learning context; the same scanning session and

temporal learning sequence, and most importantly, all objects were located on the same two-

dimensional spatial grid. Therefore, it is plausible that the various object-locations became

integrated into a unitary map, or spatial schema (Tse et al. 2007; van Buuren et al. 2014) at

initial learning and/or during replay (Durrant et al. 2015; Hennies et al. 2016). As such, object

locations could have been stored not only with reference to the two-dimensional grid, but also

directly or indirectly in spatial reference to the other objects. As such, a sound cue could have

reactivated certain object-location associations, but also objects located within the objects'

vicinity. In fact, it has been proposed that sleep-dependent memory replay is instrumental in

the integration of memories into a cognitive schema (Lewis and Durrant 2011). Thus, while

network integration of the occipital cortex during cueing is demonstrated to be positively

related to overall memory stabilization of a set of object-location associations, the fact that an

integrated spatial schema was cued may have contributed to non-specific reactivation of all

object-locations within this set.

Bearing in mind this caveat, our results show a network reorganization in response to cues

presented during SWS that is consistent with an active model of sleep-dependent memory

consolidation. Previous research has shown that whole-brain connectivity patterns display a

general reduction in thalamo-cortical and cortico-cortical connectivity and an increase in local

clustering during slow-wave sleep (Spoormaker et al. 2010). Against this backdrop, a

hippocampal-neocortical dialogue takes place facilitating systems-level consolidation by

integrating hippocampal-dependent memories into neocortical storage sites (Buzsáki 1996;

Diekelmann and Born 2010; Mitra et al. 2016). This hippocampal-neocortical dialogue is

orchestrated through oscillatory electrophysiological patterns characteristic of slow-wave sleep

and related to memory replay (Diba and Buzsáki 2007; Davidson et al. 2009). Specifically, slow

(delta wave) oscillations propagate across the brain and to the medial temporal lobes, including

the hippocampus, exerting a global control over spiking activity (Massimini et al., 2004; Nir et

al., 2011). Mechanistically, the up-state of slow oscillations has been found to enable thalamic

sleep spindles in the sigma band, which in turn cluster high-frequency hippocampal ripples in

their troughs (Staresina et al., 2015). These hippocampal ripples then are proposed to allow for

a precisely timed exchange of mnemonic information between hippocampus and neocortex,

potentiating and strengthening the neocortical memory trace. Indeed, the importance of slow

oscillations and spindles for memory consolidation has been demonstrated with behavioral

correlations (Schabus et al., 2004; Holz et al., 2012). Moreover, intracranial recordings in

humans have found time-locked ripple events during sleep in both hippocampus and

neocortex, the latter being related to memory consolidation (Axmacher et al., 2008).

Experimental disruption of hippocampal ripples in rats impaired consolidation of hippocampal-

dependent spatial memories (Girardeau et al., 2009). Therefore, during slow-wave sleep,

hierarchically nested oscillations presumably serve to consolidate memories by tightly coupling

the neocortex (especially medial prefrontal cortex), thalamus and medial temporal lobes, which

are the regions that are displaying an increased interconnectivity during cueing here.

Thus, during slow-wave sleep, there is increased cross-talk between the hippocampus,

thalamus and neocortex. If external auditory cueing would induce a reactivation of the visuo-

spatial memory traces, one would expect that cortical representation areas corresponding to

the modality of the memory trace would be recruited to participate in this cross-talk. Indeed,

the occipital cortex displayed an increase in global network integration during cueing, and when

probing the neural connectivity changes paired with this increase, we found that this region

increases its connectivity with the regions involved in active cross-talk during slow-wave sleep,

namely the hippocampus, thalamus and medial prefrontal cortex. These regions resemble those

that are involved in the oscillatory patterns that orchestrate hippocampal-neocortical cross-talk

(Mölle et al. 2006; Staresina et al. 2015). It might at first sight be surprising that the auditory

cortex is not recruited in response to auditory cueing. However, it should be noted that a

contrast was made between two conditions in which sounds were presented. The critical

difference in the contrast is that cue sounds were associated with specific visuo-spatial

memories, but not control sounds. It remains the case that one could still expect increased

connectivity of the auditory cortex with visual, spatial, and memory regions during cueing, the

sounds being associated with the object-locations. As psychophysiological interaction analyses

were informed by results of the graph theory analyses, we did not further probe connectivity

from an anatomically defined seed in the auditory cortex here. Despite this caveat, it remains to

be noted that cueing recruited the occipital cortex to engage with mnemonic regions that have

been shown to display increased cross-talk during slow-wave sleep, and this promoted the

stabilization of the associated memory traces in neocortical storage sites.

Sleep supports memory retention in a selective manner, and information is prioritized based on

perceived future relevance at encoding (Wilhelm et al. 2011; Van Dongen et al. 2012) emotional

salience (Hu et al. 2006; Wagner et al. 2006; Payne et al. 2007) and consistency across episodes

and with prior knowledge (Tamminen et al. 2013; Durrant et al. 2015). Possibly, relevant

information is selectively prioritized for active processing during sleep, whereas non-prioritized

memory traces are forgotten, perhaps aided by homeostatic regulation of synaptic plasticity

through global downscaling (Tononi and Cirelli 2003, 2006; Genzel et al. 2014). The net result is

an increase in signal-to-noise for the prioritized memory traces in the neocortex, contributing

to memory stabilization. Indeed, this appears to be the case in the parahippocampal gyrus,

where an increase in retrieval-related activation of the posterior parahippocampal gyrus at

post-sleep test versus pre-sleep test was related to an increase in network participation of the

occipital cortex during cueing. The parahippocampal gyrus is known to be involved in

representing the local visual environment (Epstein and Kanwisher 1998) and learning spatial

layouts (Aguirre et al. 1996; Epstein et al. 1999). The increased involvement of the

parahippocampal gyrus during post-sleep retrieval is consistent with a role of the

parahippocampal gyrus in binding together low-level visual information from downstream

occipital cortex. Notably, the parahippocampal gyrus was activated already during retrieval at

the pre-sleep test, showed selective increased activation in response to cueing, and displayed

increased connectivity with the occipital cortex during cued reactivation, as originally reported

in van Dongen & colleagues (2012). The results reported here further elucidate the mechanisms

involved in targeted memory reactivation by taking a network perspective.

In sum, we show here that inducing the reactivation of object-location memories with

associated auditory cues during slow-wave sleep increases the integration of occipital cortex

with the global brain network, specifically those regions involved in memory replay, such as the

medial temporal lobe, thalamus and medial prefrontal cortex. Furthermore, global network

integration of occipital cortex during cueing is predictive of overnight memory stabilization of

learned materials and an increased involvement of parahippocampal gyrus during the post-

sleep retrieval. These findings highlight how graph theory analysis can be used to assess whole-

brain connectivity patterns during targeted memory reactivation in SWS, and contribute to a

better understanding of sleep-related memory consolidation.

References

Aguirre GK, Detre JA, Alsop DC, D'Esposito M. 1996. The parahippocampus subserves
topographical learning in man. Cereb cortex. 6:823–829.
Axmacher N, Elger CE, Fell J. 2008. Ripples in the medial temporal lobe are relevant for human
memory consolidation. Brain. 131:1806–1817.
Backus AR, Bosch SE, Ekman M, Grabovetsky AV, Doeller CF. 2016. Mnemonic convergence in
the human hippocampus. Nat Commun. 7.
Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E. 2008. Fast unfolding of communities in
large networks. J Stat Mech theory Exp. 2008:P10008.
Bullmore E, Sporns O. 2009. Complex brain networks: graph theoretical analysis of structural
and functional systems. Nat Rev Neurosci. 10:186–198.
Buzsáki G. 1996. The hippocampo-neocortical dialogue. Cereb cortex. 6:81–92.
Cox R, Hofman WF, Talamini LM. 2012. Involvement of spindles in memory consolidation is slow
wave sleep-specific. Learn Mem. 19:264–267.
Creery JD, Oudiette D, Antony JW, Paller KA. 2015. Targeted memory reactivation during sleep
depends on prior learning. Sleep. 38:755–763.
Danker JF, Anderson JR. 2010. The ghosts of brain states past: remembering reactivates the
brain regions engaged during encoding. Psychol Bull. 136:87.
Davidson TJ, Kloosterman F, Wilson MA. 2009. Hippocampal replay of extended experience.
Neuron. 63:497–507.
Diba K, Buzsáki G. 2007. Forward and reverse hippocampal place-cell sequences during ripples.
Nat Neurosci. 10:1241–1242.
Diekelmann S, Born J. 2010. The memory function of sleep. Nat Rev Neurosci. 11:114–126.
Diekelmann S, Born J, Wagner U. 2010. Sleep enhances false memories depending on general
memory performance. Behav Brain Res. 208:425–429.
Diekelmann S, Büchel C, Born J, Rasch B. 2011. Labile or stable: opposing consequences for
memory when reactivated during waking and sleep. Nat Neurosci. 14:381–386.
Dupret D, O'Neill J, Pleydell-Bouverie B, Csicsvari J. 2010. The reorganization and reactivation of
hippocampal maps predict spatial memory performance. Nat Neurosci. 13:995–1002.
Durrant SJ, Cairney SA, McDermott C, Lewis PA. 2015. Schema-conformant memories are
preferentially consolidated during REM sleep. Neurobiol Learn Mem. 122:41–50.
Ekman M, Derrfuss J, Tittgemeyer M, Fiebach CJ. 2012. Predicting errors from reconfiguration
patterns in human brain networks. Proc Natl Acad Sci. 109:16714–16719.
Ekman M, Kok P, de Lange FP. 2017. Time-compressed preplay of anticipated events in human
primary visual cortex. Nat Commun. 8.
Epstein R, Harris A, Stanley D, Kanwisher N. 1999. The parahippocampal place area: recognition,
navigation, or encoding? Neuron. 23:115–125.
Epstein R, Kanwisher N. 1998. A cortical representation of the local visual environment. Nature.
392:598–601.
Euston DR, Tatsuno M, McNaughton BL. 2007. Fast-forward playback of recent memory
sequences in prefrontal cortex during sleep. Science (80-). 318:1147–1150.
Frankland PW, Bontempi B. 2005. The organization of recent and remote memories. Nat Rev

Neurosci. 6:119–130.

Genzel L, Battaglia FP. 2017. Cortico-Hippocampal Circuits for Memory Consolidation: The Role
of the Prefrontal Cortex. In: Cognitive Neuroscience of Memory Consolidation. Springer. p.
265–281.

Genzel L, Kroes MCW, Dresler M, Battaglia FP. 2014. Light sleep versus slow wave sleep in
memory consolidation: a question of global versus local processes? Trends Neurosci.
37:10–19.

Girardeau G, Benchenane K, Wiener SI, Buzsáki G, Zugaro MB. 2009. Selective suppression of
hippocampal ripples impairs spatial memory. Nat Neurosci. 12:1222–1223.

Guimera R, Mossa S, Turtschi A, Amaral LAN. 2005. The worldwide air transportation network:
Anomalous centrality, community structure, and cities' global roles. Proc Natl Acad Sci.
102:7794–7799.

Hennies N, Ralph MAL, Kempkes M, Cousins JN, Lewis PA. 2016. Sleep spindle density predicts
the effect of prior knowledge on memory consolidation. J Neurosci. 36:3799–3810.

Hu P, Stylos-Allan M, Walker MP. 2006. Sleep facilitates consolidation of emotional declarative
memory. Psychol Sci. 17:891–898.

Iturria-Medina Y, Canales-Rodriguez EJ, Melie-Garcia L, Valdes-Hernandez PA, Martinez-Montes
E, Alemán-Gómez Y, Sánchez-Bornot JM. 2007. Characterizing brain anatomical
connections using diffusion weighted MRI and graph theory. Neuroimage. 36:645–660.

Ji D, Wilson MA. 2007. Coordinated memory replay in the visual cortex and hippocampus during
sleep. Nat Neurosci. 10:100–107.

Lewis PA, Durrant SJ. 2011. Overlapping memory replay during sleep builds cognitive schemata.
Trends Cogn Sci. 15:343–351.

Marshall L, Born J. 2007. The contribution of sleep to hippocampus-dependent memory
consolidation. Trends Cogn Sci. 11:442–450.

Marshall L, Helgadóttir H, Mölle M, Born J. 2006. Boosting slow oscillations during sleep
potentiates memory. Nature. 444:610.

McClelland JL, McNaughton BL, O'Reilly RC. 1995. Why there are complementary learning
systems in the hippocampus and neocortex: insights from the successes and failures of
connectionist models of learning and memory. Psychol Rev. 102:419–457.

Mitra A, Snyder AZ, Hacker CD, Pahwa M, Tagliazucchi E, Laufs H, Leuthardt EC, Raichle ME.
2016. Human cortical–hippocampal dialogue in wake and slow-wave sleep. Proc Natl Acad
Sci. 201607289.

Mölle M, Marshall L, Gais S, Born J. 2002. Grouping of spindle activity during slow oscillations in
human non-rapid eye movement sleep. J Neurosci. 22:10941–10947.

Mölle M, Yeshenko O, Marshall L, Sara SJ, Born J. 2006. Hippocampal sharp wave-ripples linked
to slow oscillations in rat slow-wave sleep. J Neurophysiol. 96:62–70.

Nadel L, Moscovitch M. 1997. Memory consolidation, retrograde amnesia and the hippocampal
complex. Curr Opin Neurobiol. 7:217–227.

Nyberg L, Habib R, McIntosh AR, Tulving E. 2000. Reactivation of encoding-related brain activity
during memory retrieval. Proc Natl Acad Sci. 97:11120–11124.

Ong JL, Lo JC, Chee NIYN, Santostasi G, Paller KA, Zee PC, Chee MWL. 2016. Effects of phase-
locked acoustic stimulation during a nap on EEG spectra and declarative memory
consolidation. Sleep Med. 20:88–97.

Oudiette D, Antony JW, Creery JD, Paller KA. 2013. The role of memory reactivation during
wakefulness and sleep in determining which memories endure. J Neurosci. 33:6672–6678.
Payne JD, Jackson ED, Hoscheidt S, Ryan L, Jacobs WJ, Nadel L. 2007. Stress administered prior
to encoding impairs neutral but enhances emotional long-term episodic memories. Learn
Mem. 14:861–868.
Peyrache A, Khamassi M, Benchenane K, Wiener SI, Battaglia FP. 2009. Replay of rule-learning
related neural patterns in the prefrontal cortex during sleep. Nat Neurosci. 12:919–926.
Polyn SM, Natu VS, Cohen JD, Norman KA. 2005. Category-specific cortical activity precedes
retrieval during memory search. Science (80-). 310:1963–1966.
Power JD, Cohen AL, Nelson SM, Wig GS, Barnes KA, Church JA, Vogel AC, Laumann TO, Miezin
FM, Schlaggar BL. 2011. Functional network organization of the human brain. Neuron.
72:665–678.
Power JD, Schlaggar BL, Lessov-Schlaggar CN, Petersen SE. 2013. Evidence for hubs in human
functional brain networks. Neuron. 79:798–813.
Rasch B, Büchel C, Gais S, Born J. 2007. Odor cues during slow-wave sleep prompt declarative
memory consolidation. Science (80-). 315:1426–1429.
Rissman J, Gazzaley A, D'Esposito M. 2004. Measuring functional connectivity during distinct
stages of a cognitive task. Neuroimage. 23:752–763.
Rubinov M, Sporns O. 2010. Complex network measures of brain connectivity: uses and
interpretations. Neuroimage. 52:1059–1069.
Rudoy JD, Voss JL, Westerberg CE, Paller KA. 2009. Strengthening individual memories by
reactivating them during sleep. Science (80-). 326:1079.
Schabus M, Gruber G, Parapatics S, Sauter C, Klösch G, Anderer P, Klimesch W, Saletu B,
Zeitlhofer J. 2004. Sleep spindles and their significance for declarative memory
consolidation. Sleep. 27:1479–1485.
Schmitter S, Diesch E, Amann M, Kroll A, Moayer M, Schad LR. 2008. Silent echo-planar imaging
for auditory FMRI. Magn Reson Mater Physics, Biol Med. 21:317–325.
Schouten DI, Pereira SIR, Tops M, Louzada FM. 2017. State of the art on targeted memory
reactivation: sleep your way to enhanced cognition. Sleep Med Rev. 32:123–131.
Siapas AG, Wilson MA. 1998. Coordinated interactions between hippocampal ripples and
cortical spindles during slow-wave sleep. Neuron. 21:1123–1128.
Spoormaker VI, Schröter MS, Gleiser PM, Andrade KC, Dresler M, Wehrle R, Sämann PG, Czisch
M. 2010. Development of a large-scale functional brain network during human non-rapid
eye movement sleep. J Neurosci. 30:11379–11387.
Squire LR, Cohen NJ, Nadel L. 1984. The medial temporal region and memory consolidation: A
new hypothesis. Mem Consol Psychobiol Cogn. 185–210.
Staresina BP, Bergmann TO, Bonnefond M, Van Der Meij R, Jensen O, Deuker L, Elger CE,
Axmacher N, Fell J. 2015. Hierarchical nesting of slow oscillations, spindles and ripples in
the human hippocampus during sleep. Nat Neurosci.
Tamminen J, Ralph MAL, Lewis PA. 2013. The role of sleep spindles and slow-wave activity in
integrating new information in semantic memory. J Neurosci. 33:15376–15381.
Tononi G, Cirelli C. 2003. Sleep and synaptic homeostasis: a hypothesis. Brain Res Bull. 62:143–
150.
Tononi G, Cirelli C. 2006. Sleep function and synaptic homeostasis. Sleep Med Rev. 10:49–62.

Tse D, Langston RF, Kakeyama M, Bethus I, Spooner PA, Wood ER, Witter MP, Morris RGM.
2007. Schemas and Memory Consolidation. Sci . 316:76–82.

Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer B,
Joliot M. 2002. Automated anatomical labeling of activations in SPM using a macroscopic
anatomical parcellation of the MNI MRI single-subject brain. Neuroimage. 15:273–289.

van Buuren M, Kroes MCW, Wagner IC, Genzel L, Morris RGM, Fernández G. 2014. Initial
investigation of the effects of an experimentally learned schema on spatial associative
memory in humans. J Neurosci. 34:16662–16670.

van Dongen E V, Takashima A, Barth M, Zapp J, Schad LR, Paller KA, Fernández G. 2012. Memory
stabilization with targeted reactivation during human slow-wave sleep. Proc Natl Acad Sci .
109:10575–10580.

Van Dongen E V, Thielen J-W, Takashima A, Barth M, Fernández G. 2012. Sleep supports
selective retention of associative memories based on relevance for future utilization. PLoS
One. 7:e43426.

Wagner U, Hallschmid M, Rasch B, Born J. 2006. Brief sleep after learning keeps emotional
memories alive for years. Biol Psychiatry. 60:788–790.

Wilhelm I, Diekelmann S, Molzow I, Ayoub A, Mölle M, Born J. 2011. Sleep selectively enhances
memory expected to be of future relevance. J Neurosci. 31:1563–1569.

Wilson MA, McNaughton BL. 1994. Reactivation of hippocampal ensemble memories during
sleep. Science (80-). 265:676–679.

Xu S, Jiang W, Poo M, Dan Y. 2012. Activity recall in a visual cortical ensemble. Nat Neurosci.
15:449–455.

Figures

Figure 1

Schematic depiction of the experimental procedure.

Participants learned 50 object-location associations, presented simultaneously with object-related

sounds, inside the MR-scanner. After a baseline test (Test 1), participants were set up for

polysomnographic recordings and went to sleep inside the MR-scanner. Half of the learned associations

were cued with sounds during slow-wave sleep (SWS). After awaking and a short break, participants

performed another post-sleep test (Test 2). Figure adapted with permission from the initial report (van

Dongen et al., 2012).

Replay-related changes in network integration predict behavioral memory stabilization.

A) *Participation Coefficient mapped for every voxel in the brain during the presentation of control*

sounds (not previously paired with object-location information, left panel) and cue sounds

(previously paired with object-location associations, right panel).

B) *Increased network integration was found only in the occipital cortex during the presentation of*

cue sounds versus control sounds during SWS. This region overlapped with the set of regions

activated in response to the object-location association task. Parametric maps were

superimposed onto a template brain, using the cluster-forming threshold of Z>2.33.

C) *The increase in participation coefficient of the occipital cortex during cueing predicted memory*

stabilization, indicated by reduced forgetting (expressed as the difference in error distances)

between test 1 (pre) and test 2 (post).

Figure 3

Regions displaying greater coupling with occipital cortex during cueing in SWS

Regions displaying stronger coupling with the occipital cortex for cue sounds compared to control

sounds. Parametric maps were superimposed onto a template brain, using the cluster-forming threshold

of Z>2.33. HC = Hippocampus, PHG = Parahippocampal Gyrus, mCC = Middle Cingulate Cortex, SMA =

Supplementary Motor Area, dmPFC = Dorsomedial Prefrontal Cortex.

Figure 4

680
Overnight forgetting, split by network integration in occipital cortex

The bar graph depicts the change in error distance between the pre-sleep Test 1 and the post-sleep Test

2 for a group where there was an increase in network integration (as measured with the participation

coefficient) in occipital cortex (PC+) and a group where no such increase was found (PC=). The groups

were defined using a median split on the participation coefficient values acquired from the occipital

cortex in the cue sounds > control sounds comparison. The error distance is operationalized as the

distance between the location where the object is placed at test, and the actual object location as

presented during the learning phase. A higher overnight error increase indicates a reduction in accuracy

of placement across sleep, indicative of increased forgetting. The group where cueing induced increasing

network integration in the occipital cortex (PC+) displayed reduced forgetting overall, indicating that, in

this group, cueing stabilizes memories overnight.

Figure 5

Overnight activity increases related to network integration of occipital cortex

Regions where there is a relation between on the one hand, an activity increase between the pre-sleep

and post-sleep test and, on the other hand, increased network integration of occipital cortex during

cueing in the SWS between Test1 and Test 2. Parametric maps were superimposed onto a template

brain, using the cluster-forming threshold of Z>2.33.