TITLE

Hsf1 phosphorylation generates cell-to-cell variation in Hsp90 levels and

promotes phenotypic plasticity

AUTHORS

Xu Zheng[1,*], Ali Beyzavi[2,5,*], Joanna Krakowiak[1], Nikit Patel[3], Ahmad S. Khalil[3,4,#], David

Pincus[1,#,§]

[1] Whitehead Institute for Biomedical Research, Cambridge, USA

[2] Department of Mechanical Engineering, Boston University, Boston, USA

[3] Department of Biomedical Engineering and Biological Design Center, Boston University,

Boston, USA

[4] Wyss Institute for Biologically Inspired Engineering, Harvard University, Boston, USA

[5] Present address: Koch Institute for Integrative Cancer Research, Cambridge, USA

* These authors contributed equally

[#] Correspondence: khalil@bu.edu, pincus@wi.mit.edu

[§] Lead contact: pincus@wi.mit.edu

ABSTRACT

Clonal populations of cells exhibit cell-to-cell variation in the transcription of individual genes. In

addition to this "noise" in gene expression, heterogeneity in the proteome and the proteostasis

network expands the phenotypic diversity of a population. Heat shock transcription factor (Hsf1)

regulates chaperone gene expression, thereby coupling transcriptional noise to proteostasis.

Here we show that cell-to-cell variation in Hsf1 activity is an important determinant of phenotypic

plasticity. Budding yeast cells with high Hsf1 activity were enriched for the ability to acquire

resistance to an antifungal drug, and this enrichment depended on Hsp90 – a known

"phenotypic capacitor" and canonical Hsf1 target. We show that Hsf1 phosphorylation promotes

cell-to-cell variation, and variation in Hsf1 activity – rather than absolute Hsf1 activity – promotes

antifungal resistance. We propose that Hsf1 phosphorylation enables differential tuning of the

proteostasis network in individual cells, allowing populations to access a wide range of

phenotypic states.

INTRODUCTION

Genetically identical cells grown together in the same environment nonetheless display cell-to-

cell variation in gene expression (Colman-Lerner et al., 2005; Elowitz et al., 2002; Raser and

O'Shea, 2004, 2005; Weinberger et al., 2005). While most frequently observed in

microorganisms, such as bacteria and yeast, gene expression variation is also found in

developing mammalian cells and human embryonic stem cells (Silva and Smith, 2008; Stelzer

et al., 2015). Such variation has been proposed to be the mechanistic underpinning of lineage

commitment during human development, the epithelial to mesenchymal transition in cancer

metastasis, organ regeneration in planarians, bacterial persistence in the presence of

antibiotics, and the ability of yeast cells to remain fit in fluctuating environments (Harms et al.,

2016; Newman et al., 2006; Oderberg et al., 2017; Silva and Smith, 2008; Ye and Weinberg,

2015). While differences in cell size, cell cycle position and chromatin state can partially account

for cell-to-cell variation, much of the variability has been attributed to the inherently stochastic

process of gene expression (Colman-Lerner et al., 2005; Raj and van Oudenaarden, 2008;

Raser and O'Shea, 2005). Despite the underlying stochasticity, gene expression variation itself

varies widely across the genome, with some sets of genes showing very low variation among

cells (e.g., ribosomal protein genes) and other sets of genes (e.g., stress responsive genes)

showing high levels of variation (Newman et al., 2006). Yet, individual genes within these

regulons show strong covariance, indicating the source of the variation lies in the activity of

upstream transcription factors and signaling pathways (Stewart-Ornstein et al., 2012). As such,

cell-to-cell variation may be a property that is under genetic control and can be tuned up and

down over evolution.

On top of this gene expression variation, cell-to-cell differences also exist in the state of the

proteome. Perhaps the most striking examples of proteome variation come from prion proteins,

which can exist in either soluble or self-templating amyloid conformations (Shorter and

Lindquist, 2005). Prions have been shown to have the ability to broadly reshape the proteome

by challenging chaperones and other components of the protein homeostasis (proteostasis)

machinery and even by globally altering protein translation (Serio and Lindquist, 1999; Shorter

and Lindquist, 2008). Moreover, chaperones themselves can exist in large heterotypic

complexes that differ between cells in what has been termed the "epichaperome," giving rise to

altered susceptibility of cancer cells to drugs that target the essential chaperone Hsp90 (Rodina

et al., 2016). By buffering the proteome and stabilizing near-native protein folds, Hsp90 has

been shown to mask latent genetic variation in fruit flies and plants and to enhance the ability of

yeast cells to acquire novel phenotypes such as resistance to anti-fungal drugs (Cowen and

Lindquist, 2005; Queitsch et al., 2002; Rutherford and Lindquist, 1998). In this regard, Hsp90

has been termed a "phenotypic capacitor" (Sangster et al., 2004).

Heat shock factor 1 (Hsf1) regulates the expression of many components of the proteostasis

machinery – including Hsp90 – in eukaryotes from yeast to humans. In unstressed budding

yeast cells, a different chaperone, Hsp70, binds to Hsf1 and restrains its activity. Upon heat

shock, Hsp70 dissociates from Hsf1 leaving Hsf1 free to induce expression of its target genes.

Heat shock also triggers Hsf1 hyper-phosphorylation. Although phosphorylation is a conserved

hallmark of Hsf1 activation, it is dispensable for acute Hsf1 activity during heat shock. Rather

than switching Hsf1 on, phosphorylation enables Hsf1 to sustain increased activity during

prolonged exposure to elevated temperature. Here we identify a novel role for Hsf1 – and Hsf1

phosphorylation – that may have provided a strong selective advantage during evolution. We

show that Hsf1 generates cell-to-cell variation in Hsp90 levels, which in turn contributes to the

ability of *Saccharomyces cerevisiae* to acquire resistance to the antifungal drug fluconazole. We

find that the ability of Hsf1 to become phosphorylated is a key factor in generating population

level heterogeneity in its activity. We propose that by coordinately controlling cytosolic

chaperone genes including Hsp90, Hsf1 promotes phenotypic plasticity.

RESULTS

Differential cell-to-cell variation in Hsf1 activity in response to heat shock and AZC

In addition to heat shock, Hsf1 is known to respond to a variety of chemical stressors that impair

proteostasis. In particular, the small molecule azetidine 2-carboxylic acid (AZC) is known to

strongly activate Hsf1 (Trotter et al., 2002). AZC is a proline analog that is charged onto tRNA[pro]

and incorporated into nascent proteins during translation, impairing their subsequent folding

(Fowden and Richmond, 1963). Our previous mass spectrometry data indicated that Hsf1

displays distinct phosphorylation patterns in cells that had been heat shocked compared to cells

treated with AZC (Zheng et al., 2016). To explore these differences, we monitored

phosphorylation of FLAG-tagged Hsf1 following heat shock and treatment with AZC by

observing its electrophoretic mobility over time by western blot. Heat shock induced progressive

and sustained Hsf1 phosphorylation leading to a dramatic shift in its mobility over the heat

shock time course (Figure 1A). By contrast, AZC led to only a modest shift in Hsf1 mobility

(Figure 1A). Despite these differences in phosphorylation, both heat shock and AZC robustly

induced Hsf1 transcriptional activity as measured by flow cytometry of cells expressing HSE-

YFP, a fluorescent reporter of Hsf1 transcriptional activity (Figure 1B) (Zheng et al., 2016).

While heat shock resulted in rapid HSE-YFP induction, plateauing after 1 hour, AZC induced

Hsf1 activity with delayed but sustained kinetics, ultimately leading to the same maximal output

level as heat shock after 4 hours (Figure 1B).

Along with altered activation kinetics, we observed differences in the single cell fluorescence

distributions: the populations of cells that had been heat shocked showed a broad distribution of

HSE-YFP levels, while cells treated with AZC showed narrower distributions, indicating reduced

levels of cell-to-cell variation in Hsf1 activity. To account for potential differences in cell size, we

normalized each cell's HSE-YFP fluorescence by its size as measured by side scatter (SSC)

and plotted the resulting normalized distributions over the heat shock and AZC time courses

(Figure 1C, D) (Stewart-Ornstein et al., 2012). Direct comparison of the HSE-YFP/SSC

distributions following 4 hours of heat shock or treatment with AZC reveals that despite a slightly

higher average level of Hsf1 activation in the AZC-treated cells, cellular heterogeneity is greatly

reduced compared to heat shocked cells (Figure 1E). To quantify this cell-to-cell variation in the

HSE-YFP reporter, we calculated the square of the coefficient of variation (CV^2) (Stewart-

Ornstein et al., 2012). While the CV^2 drops as the mean increases for both heat shocked and

AZC-treated cells, it is 3-fold higher in heat-shocked cells by the end of the experiment (Figure

1F). These data show that while heat shock and AZC both potently activate Hsf1, they do so

with distinctive features. Heat shock triggers rapid Hsf1 activation coupled to high levels of

phosphorylation and results in a high degree of cell-to-cell variation in the HSE-YFP reporter. By

contrast, AZC induces slow, sustained Hsf1 activation, modest Hsf1 phosphorylation and

reduced noise in the HSE-YFP reporter.

Hsf1 phosphorylation generates cell-to-cell variation during heat shock

Since increased Hsf1 phosphorylation was associated with increased cell-to-cell variation in the

HSE-YFP reporter, we wondered if Hsf1 phosphorylation was responsible for increasing noise

during heat shock. To test this, we leveraged a non-phosphorylatable mutant we had previously

generated, Hsf1$^{\Delta po4}$, in which we mutated all 152 possible sites of phosphorylation to alanine

(Zheng et al., 2016). We performed a heat shock time course and measured the HSE-YFP

reporter by flow cytometry in wild type cells and cells expressing Hsf1$^{\Delta po4}$. Indeed, after

normalizing for cell size, we observed reduced noise in the HSE-YFP reporter in the Hsf1$^{\Delta po4}$

cells compared to wild type throughout the heat shock time course (Figure 2A, B). As an

orthogonal approach to measure cell-to-cell variation in Hsf1 activity, we developed a

microfluidic-based assay to measure the HSE-YFP reporter in single cells over time in response

to heat shock using a novel platform that enables precision temperature control (Figure 2C,

Figure S1). In agreement with the flow cytometry data, cells expressing Hsf1$^{\Delta po4}$ showed lower

noise in HSE-YFP levels over the heat shock time course (Figure 2D, E). Thus, Hsf1

phosphorylation promotes cell-to-cell variation in Hsf1 activity during heat shock.

Hsp90 expression displays Hsf1-dependent cell-to-cell variation

To investigate functional roles for cell-to-cell variation in Hsf1 activity, we monitored expression

of Hsp90, an endogenous Hsf1 target gene known to influence phenotypic plasticity (Cowen

and Lindquist, 2005). Budding yeast encodes two Hsp90 paralogs, Hsc82 and Hsp82, with

Hsc82 being highly expressed under all conditions and Hsp82 showing both basal and inducible

expression in response to heat shock (Solis et al., 2016). We tagged Hsp82 with YFP and

monitored its expression in single cells expressing either wild type Hsf1 or Hsf1$^{\Delta po4}$ by

fluorescence microscopy and flow cytometry. Consistent with the HSE-YFP reporter, Hsp82-

YFP expression showed greater variability among wild type cells than Hsf1$^{\Delta po4}$ cells (Figure 3A,

B).

Prior studies have implicated Hsp90 as a "phenotypic capacitor" that promotes the ability of cells

to acquire novel phenotypes (Sangster et al., 2004). In particular, in fungi, Hsp90 has been

shown to increase the rate at which cells acquire resistance to the anti-fungal drug fluconazole

(Cowen and Lindquist, 2005). We hypothesized that the increased cell-to-cell variation in

Hsp82-YFP expression levels in wild type cells compared to Hsf1$^{\Delta po4}$ cells would translate into

increased fluconazole resistance. Indeed, wild type cells generated significantly more

fluconazole resistant colonies than Hsf1$^{\Delta po4}$ cells (Figure 3C, D).

Acquired fluconazole resistance is enriched in cells with high Hsf1 activity

Since noise in Hsp82-YFP levels and the HSE-YFP reporter correlated with phenotypic

plasticity, we wondered if individual cells with elevated HSE-YFP levels would be more likely to

become resistant to fluconazole. To test this, we used fluorescence activated cell sorting

(FACS) to isolate the tails of the HSE-YFP distribution – the 10% of cells with the highest and

lowest HSE-YFP expression levels – in wild type cells, *hsp82Δ* and *hsc82Δ* cells (Figure 3E). In

wild type cells, the tail with high HSE-YFP expression was enriched for fluconazole resistant

colonies over the low expression tail (Figure 3F). By contrast, the high expression tails of the

hsp82Δ and *hsc82Δ* cells showed no enrichment, with the *hsc82Δ* high expression tail actually

showing decreased resistance (Figure 3F). These results indicate that cells with higher levels of

Hsf1 activity have an increased ability to acquire resistance to an antifungal drug, and this ability

depends on Hsp90.

Fluconazole resistance correlates with variation in Hsf1 activity, not its magnitude

Since the high expression tail of the HSE-YFP distribution was enriched for cells able to acquire

resistance to fluconazole, we hypothesized that if we could increase the average HSE-YFP

expression, we could increase the ability of cells to acquire fluconazole resistance. To test this,

we expressed wild type Hsf1 from a promoter under the control of a synthetic, estradiol-

responsive (ER) transcription factor (Zheng et al., 2016). In this way, we could titrate the amount

of estradiol to tune the expression level of Hsf1, and thereby control the expression level of its

target genes and the HSE-YFP reporter. In parallel, we also expressed Hsf1$^{\Delta po4}$ using the same

system to determine if the deficit in cell-to-cell variation could be overcome by increasing the

absolute expression (Figure 4A). We grew the wild type and Hsf1$^{\Delta po4}$ cells in a dilution series of

estradiol, maintained growth at $OD_{600} < 0.5$ for 18 hours and measured the HSE-YFP reporter.

Estradiol induced dose-dependent induction of the HSE-YFP in both wild type and Hsf1$^{\Delta po4}$

cells, though with a steeper curve for Hsf1$^{\Delta po4}$ cells (Figure 4B, C). At all doses of estradiol, wild

type had higher CV^2 in the HSE-YFP reporter than Hsf1$^{\Delta po4}$ cells (Figure 4B, D). We tested the

ability of cells expressing wild type Hsf1 and Hsf1$^{\Delta po4}$ at low (1 nM), medium (8 nM) and high (32

nM estradiol) levels to acquire resistance to fluconazole (Figure 4E). Surprisingly, increasing

median HSE-YFP did not increase resistance: both wild type and Hsf1$^{\Delta po4}$ cells showed maximal

resistance at low Hsf1 expression (Figure 4F). However, wild type cells showed greater

resistance at all Hsf1 expression levels than Hsf1$^{\Delta po4}$ cells (Figure 4F). Thus, increased variation

in HSE-YFP levels – not increased median levels – positively correlates with fluconazole

resistance (Figure 4G-I).

DISCUSSION

Cell-to-cell variation in gene expression and the state of the proteome have been proposed to

contribute to adaptability in populations of genetically identical cells. Here we identify a single

regulatory mechanism that couples variation at the transcriptional and proteomic levels:

phosphorylation of the transcription factor Hsf1. We found that when wild type Hsf1 is activated

and concomitantly hyperphosphorylated – as is the case during heat shock – cells display high

variation in their expression of downstream Hsf1 target genes. By contrast, when Hsf1 is

activated without hyperphosphorylation following treatment with AZC, cells show more uniform

target gene expression. Moreover, removing the ability of Hsf1 to be phosphorylated reduced

cell-to-cell variation. Since the Hsf1 regulon consists of chaperones and other proteostasis

factors, variation in Hsf1 activity is likely to lead to variation in the proteostasis network and the

state of the proteome.

Functionally, we demonstrated that cell-to-cell variation in Hsf1 activity is correlated with the

ability of cells to acquire resistance to the antifungal drug fluconazole, in large part via

expression of the known phenotypic capacitor Hsp90. While wild type cells with high levels of

Hsf1 activity were enriched for the ability to acquire fluconazole resistance, cells with reduced

levels of Hsp90 lost this enrichment. This observation suggested that cells with more Hsf1

activity would have more Hsp90 and therefore show increased antifungal resistance. However,

synthetically increasing Hsf1 activity resulted in diminished fluconazole resistance. Moreover,

increased expression of non-phosphorylatable Hsf1$^{\Delta po4}$ – which showed reduced cell-to-cell

variation in its activity compared to wild type Hsf1 – could not compensate for its reduced ability

to support fluconazole resistance. Thus, antifungal resistance correlates with variation in Hsf1

activity rather than with the magnitude of Hsf1 activity.

How does Hsf1 phosphorylation lead to cell-to-cell variation and why does variation promote

fluconazole resistance? It is likely that differential activation of cell cycle, metabolic and stress

responsive kinase pathways results in differential Hsf1 phosphorylation states and leads to

distinct levels of transcriptional activity in single cells. However, the finding that variation in Hsf1

activity rather than its absolute activity drives resistance is more intriguing. Although cells with

synthetically elevated Hsf1 activity produce more Hsp90, they also overproduce the rest of the

Hsf1 regulon, which may be maladaptive. In addition, with forced Hsf1 activity, cells are unable

to dynamically regulate Hsf1 to precisely tune its activity according to need. While a high level of

Hsf1 activity – and thus Hsp90 levels – may be beneficial early in the process of acquiring

resistance to fluconazole, it may be that a subsequent reduction in Hsf1 activity promotes

proliferation. The capacity for dynamic and coordinated control over the proteostasis network

may endow cells with the plasticity required to maintain fitness in fluctuating environments. By

enabling both population-level variation and dynamic control over the proteostasis network, Hsf1

is a powerful phenotypic capacitor.

ACKNOWLEDGEMENTS

We dedicate this paper to Susan Lindquist and her scientific legacy. We are grateful to K. Reynolds and R. Ranganathan for beneficial discussions, to the Whitehead Institute FACS facility for technical assistance and to N. Azubuine and T. Nanchung for a constant supply of plates and media. This work was supported by an NIH Early Independence Award (DP5 OD017941-01 to D.P.) and a National Science Foundation CAREER Award (MCB-1350949 to A.S.K.).

AUTHOR CONTRIBUTIONS

Conceptualization, D.P. and A.S.K; Methodology, X.Z., A.B. N.P.; Investigation, X.Z., A.B., J.K., D.P., N.P.; Writing, D.P. and A.S.K; Funding Acquisition, D.P. and A.S.K.; Supervision, D.P. and A.S.K.

  FIGURE LEGENDS

Figure 1. Differential cell-to-cell variation in Hsf1 activity in response to heat shock and

AZC.

(A) Anti-FLAG Western blot showing Hsf1 phosphorylation by electrophoretic mobility shift over

time in response to heat shock at 39ºC or 4 mM AZC.

(B) Hsf1 activity quantified by the levels of the HSE-YFP transcriptional reporter measured by

flow cytometry over time in response to heat shock at 39ºC or 4 mM AZC. The reporter consists

of four repeats of the heat shock element (HSE) recognized by Hsf1 upstream of a crippled

CYC1 promoter. Each point represents the median of the fluorescence distribution of 10000

cells averaged over 3 biological replicates; error bars show the standard deviation of the

replicates.

(C) Single cell fluorescence distributions over a 4-hour heat shock time course. For each cell,

the YFP level was normalized by side scatter (HSE-YFP/SSC) to control for cell size.

(D) As in (C), but over a time course of treatment with 4 mM AZC.

(E) Overlay of the 4-hour time points from (C) and (D).

(F) Quantification of cell-to-cell variation in HSE-YFP/SSC levels over time courses in (C) and

(D) using the square of the coefficient of variation (CV^2) as a metric.

Figure 2. Hsf1 phosphorylation generates cell-to-cell variation during heat shock.

(A) Single cell fluorescence distributions over a heat shock time course in wild type cells and

cells expressing Hsf1$^{\Delta po4}$ as the only copy of Hsf1.

(B) Quantification of cell-to-cell variation in HSE-YFP/SSC levels over time courses in (A).

(C) Images of wild type cells expressing the HSE-YFP reporter growing over time in a

microfluidic device at 25ºC and shifted to 39ºC at the indicated time. Scale bar is 10 µm.

(D) Quantification of HSE-YFP levels over time in the microfluidic device in wild type cells and

cells expressing Hsf1$^{\Delta po4}$ as the only copy of Hsf1. For each cell, the total YFP fluorescence was

divided by cell area to account for cell size differences. Each gray trace represents the trajectory

of a single cell and the thick lines show population averages.

(E) Quantification of cell-to-cell variation in HSE-YFP levels over time courses in (D).

Figure 3. Cell-to-cell variation in Hsf1 activity promotes antifungal resistance in an

Hsp90-dependent manner.

(A) Spinning disc confocal images of wild type cells and Hsf1$^{\Delta po4}$ cells expressing Hsp82-YFP.

Scale bar is 10 µm.

(B) Single cell fluorescence distributions of wild type cells and Hsf1$^{\Delta po4}$ cells expressing Hsp82-

YFP as determined by flow cytometry.

(C) Appearance of fluconazole-resistant colonies in wild type cells and Hsf1$^{\Delta po4}$ cells. 10^6 cells

were plated on YPD plates supplemented with 128 µg/ml fluconazole and incubated at room

temperature for 4 days.

(D) Quantification of the number of fluconazole resistant colonies from three biological replicates

of the experiment shown in (C). Error bars show the standard deviation of the replicates (* $p <$

0.05; ** $p < 0.01$ by two-tailed T-test).

(E) Single cell fluorescence distributions of HSE-YFP/SSC in wild type and Hsp90 mutant cells.

under basal conditions. Deletion of *HSP82* leads to a modest increase in HSE-YFP levels, while

deletion of *HSC82* leads to a dramatic increase in HSE-YFP levels. The blue and red boxes

indicate the tails of the distribution with the highest and lowest expressing 10% of cells.

(F) Quantification of the number of fluconazole resistant colonies in the tails of the distributions

shown in (E) (* $p < 0.05$; ** $p < 0.01$ by two-tailed T-test)..

Figure 4. Fluconazole resistance correlates with cell-to-cell variation in Hsf1 activity.

(A) Schematic showing the estradiol-inducible system used to titrate expression of wild type

Hsf1 and Hsf1$^{\Delta po4}$. A chimeric transcription factor containing the ligand binding domain of the

human estrogen receptor drives the expression of Hsf1 as a function of the concentration of

estradiol added to the growth media.

(B) Single cell HSE-YFP/SSC fluorescence distributions in cells treated across a dose response

of estradiol, ranging from 2-128 nM in cells expressing wild type Hsf1 (gray) or Hsf1$^{\Delta po4}$ (green)

as the only copy of Hsf1. Cells were incubated in log phase for 18 hours in the presence of the

indicated dose of estradiol to achieve steady state before measuring.

(C) Quantification of the median HSE-YFP/SSC across the estradiol dose response in wild type

and Hsf1$^{\Delta po4}$ cells relative to the maximum value.

(D) Quantification of the cell-to-cell variation (CV^2) of the HSE-YFP/SSC distributions across the

estradiol dose response in wild type and Hsf1$^{\Delta po4}$ cells.

(E) Overlays of the HSE-YFP/SSC distributions from wild type and Hsf1$^{\Delta po4}$ cells treated with the

indicated concentrations of estradiol.

(F) Quantification of fluconazole resistant colonies in wild type and Hsf1$^{\Delta po4}$ cells in the presence

of the indicated doses of estradiol (* $p < 0.05$; ** $p < 0.01$ by two-tailed T-test).

(G) Fluconazole resistant wild type and Hsf1$^{\Delta po4}$ colonies plotted as a function of median Hsf1

activity as measured by the HSE-YFP reporter. Dashed line is a linear regression to all six data

points showing a negative correlation.

(H) Fluconazole resistant wild type and Hsf1$^{\Delta po4}$ colonies plotted as a function of the variation in

Hsf1 activity as measured by the CV^2 of the HSE-YFP reporter. Dashed line is a linear

regression to all six data points showing a positive correlation.

(I) Correlation coefficients (r) of the data plotted in **(G)** and **(H)**.

Figure S1. Characterization of the microfluidic heat shock device.

(A) Schematic of the microfluidic heat shock device. The assembled device consists of a glass

coverslip patterned with Pt/Ti heater and sensor wires (grey), and a multilayer PDMS device

with flow (black and blue channels) and control (red channels) layers. Cells loaded from either

port A or B are trapped in chambers (blue) that have been fabricated to the height of a single

monolayer of *S. cerevisiae* cells. W denotes waste port.

(B) Representative calibration curve for the on-chip sensor showing a linear response across

the operating temperature range.

(C) Using the on-chip heater to achieve and accurately maintain chamber temperature of 39˚C,

as measured by a high-resolution long-wave infrared thermal camera.

METHODS

Yeast strains and cell growth

Yeast cells were cultured in SDC media as described (Zheng et al., 2016).

DPY number	Description	Genotype
144	WT HSE-YFP	W303 MATa 4xHSE-Venus::LEU
118	Hsf1-3xFLAG/V5	W303 MATa hsf1Δ::KAN Hsf1-3xFLAG-V5::TRP1
416	Hsf1Δpo4 HSE-YFP	W303 MATa hsf1Δ::KAN 4xHSE-Venus::LEU2 Hsf1Δpo4::TRP1
1221	hsc82Δ HSE-YFP	W303 MATa 4xHSE-Venus::LEU2 hsc82Δ::HYG
1222	hsp82Δ HSE-YFP	W303 MATa 4xHSE-Venus::LEU2 hsp82Δ::HYG

Flow cytometry

Flow cytometry was performed as described (Zheng et al., 2016). Data were processed in

FlowJo 10. Data were left ungated and YFP fluorescence was normalized by side scatter (SSC)

for each cell.

Microfluidic heat shock time courses

Single cell heat shock experiments were performed with a custom microfluidic device that uses

microscale, on-chip heaters to enable programmable thermal perturbations. The multilayer

device was fabricated out of the silicone elastomer polydimethylsiloxane (PDMS/Sylgard 184,

Dow Corning) using soft lithographic techniques, as described previously (Duffy et al., 1998;

Thorsen et al., 2002; Unger et al., 2000; Vega et al., 2012). The device was aligned and sealed

to a pre-cleaned No. 1.5 glass coverslip (Fisher Scientific), onto which micro-scale heater and

resistance temperature detector (RTD) wires were patterned. Briefly, a thin layer of Ti (100

Angstroms) was deposited on the glass, followed by a layer of Pt (150 nm) and finally a layer of

SPR 220-7 positive photoresist (MicroChem Corp.). After transferring the heater/RTD pattern to

the photoresist using a high-resolution transparency photomask (CAD/Art Services, Inc.), the

Pt/Ti layers were etched in Aqua Regia solution to generate the final serpentine pattern of

heater and RTD wires. The photoresist was then washed off with acetone, and a layer of SiO_2

(300 nm) was deposited on the final pattern. After assembly, the RTD of each device was

calibrated and confirmed to respond linearly to changes in the input temperature, as controlled

by a hot plate, across the operating temperature range of 25°C to 45°C (CITE FIGURE?).

Additionally, the devices were characterized by applying user-defined voltages to the heater and

measuring on-chip temperatures using a Flir A655sc high-resolution long-wave infrared thermal

camera (Flir Systems, Inc.) to validate the device's ability to achieve accurate and stable heat

shock conditions (CITE FIGURE?).

To perform microfluidic experiments, we used a custom microfluidic platform that controls the

delivery of liquids to the device and the actuation of valves (Vega et al., 2012). The voltage

across the heater wire and the resistance across the RTD wire were controlled and measured,

respectively, to modulate the on-chip temperature. For each experiment, cells were inoculated

1:100 from overnight SDC cultures into 2 mL SDC and allowed to grow 3-4 hours (OD ~0.5)

before seeding the device. Cells were then loaded into the device to trap a small number of cells

in growth chambers. Cells were grown on-chip in SDC media for 4-5 hours at 25°C, ambient

conditions maintained with a Controlled Environment Microscope Incubator (Nikon Instruments,

Inc.) designed for live-cell imaging. Cells were then subjected to a heat shock by applying

voltage across the heater wire to ramp to and maintain an on-chip temperature of 39°C.

Throughout, images were collected at 15-minute intervals at 100x magnification (Plan Apo

Lambda 100X, NA 1.45) using an Eclipse Ti-E inverted microscope (Nikon Instruments, Inc.)

equipped with the "Perfect Focus System", a XYZ-motorized stage, and a Clara-E charge-

coupled device (CCD) camera (Andor Technology). Images were acquired in phase contrast

configuration and in the YFP fluorescent channel. Filters and light sources (Nikon LED and

Lumencor SPECTRA X Light Engine) were automatically controlled using the supplier's

software (NIS-Elements Advanced Research). Following each experiment, cells were

segmented using custom image analysis software written for Matlab (Mathworks, Natick, MA),

and fluorescence values for each cell, averaged over cell area, were obtained from the time

series of YFP images.

Spinning disc confocal imaging

Imaging was performed as described (Zheng et al., 2016).

Fluconazole acquired resistance assay

Wild type and Hsf1$^{\Delta po4}$ cells were grown to mid-log phase and 10^6 cells were plated on YPD

plates with 128 µg/ml fluconazole + 25 nM estradiol. Plates were incubated for 4 days at room

temperature. Colonies were counted from 3 biological replicates and mean and standard

deviation were calculated.

Fluorescence activated cell sorting

10^6 cells from the high- and low-expressing tails (top and bottom 10% of cells) of the HSE-

YFP/SSC distribution in wild type, *hsp82Δ* and *hsc82Δ* cells were sorted in the Whitehead

Institute FACS facility on a BD FACS Aria.

Estradiol dose responses

Performed as described (Zheng et al., 2016).

REFERENCES

Colman-Lerner, A., Gordon, A., Serra, E., Chin, T., Resnekov, O., Endy, D., Pesce, C.G., and Brent, R. (2005). Regulated cell-to-cell variation in a cell-fate decision system. Nature *437*, 699-706.

Cowen, L.E., and Lindquist, S. (2005). Hsp90 potentiates the rapid evolution of new traits: drug resistance in diverse fungi. Science *309*, 2185-2189.

Duffy, D.C., McDonald, J.C., Schueller, O.J., and Whitesides, G.M. (1998). Rapid Prototyping of Microfluidic Systems in Poly(dimethylsiloxane). Anal Chem *70*, 4974-4984.

Elowitz, M.B., Levine, A.J., Siggia, E.D., and Swain, P.S. (2002). Stochastic gene expression in a single cell. Science *297*, 1183-1186.

Fowden, L., and Richmond, M.H. (1963). Replacement of Proline by Azetidine-2-Carbolic Acid during Biosynthesis of Protein. Biochim Biophys Acta *71*, 459-&.

Harms, A., Maisonneuve, E., and Gerdes, K. (2016). Mechanisms of bacterial persistence during stress and antibiotic exposure. Science *354*.

Newman, J.R., Ghaemmaghami, S., Ihmels, J., Breslow, D.K., Noble, M., DeRisi, J.L., and Weissman, J.S. (2006). Single-cell proteomic analysis of S. cerevisiae reveals the architecture of biological noise. Nature *441*, 840-846.

Oderberg, I.M., Li, D.J., Scimone, M.L., Gavino, M.A., and Reddien, P.W. (2017). Landmarks in Existing Tissue at Wounds Are Utilized to Generate Pattern in Regenerating Tissue. Curr Biol *27*, 733-742.

Queitsch, C., Sangster, T.A., and Lindquist, S. (2002). Hsp90 as a capacitor of phenotypic variation. Nature *417*, 618-624.

Raj, A., and van Oudenaarden, A. (2008). Nature, nurture, or chance: stochastic gene

expression and its consequences. Cell *135*, 216-226.

Raser, J.M., and O'Shea, E.K. (2004). Control of stochasticity in eukaryotic gene expression.

Science *304*, 1811-1814.

Raser, J.M., and O'Shea, E.K. (2005). Noise in gene expression: origins, consequences, and

control. Science *309*, 2010-2013.

Rodina, A., Wang, T., Yan, P., Gomes, E.D., Dunphy, M.P., Pillarsetty, N., Koren, J.,

Gerecitano, J.F., Taldone, T., Zong, H.*, et al.* (2016). The epichaperome is an integrated

chaperome network that facilitates tumour survival. Nature *538*, 397-401.

Rutherford, S.L., and Lindquist, S. (1998). Hsp90 as a capacitor for morphological evolution.

Nature *396*, 336-342.

Sangster, T.A., Lindquist, S., and Queitsch, C. (2004). Under cover: causes, effects and

implications of Hsp90-mediated genetic capacitance. Bioessays *26*, 348-362.

Serio, T.R., and Lindquist, S.L. (1999). [PSI+]: an epigenetic modulator of translation

termination efficiency. Annu Rev Cell Dev Biol *15*, 661-703.

Shorter, J., and Lindquist, S. (2005). Prions as adaptive conduits of memory and inheritance.

Nat Rev Genet *6*, 435-450.

Shorter, J., and Lindquist, S. (2008). Hsp104, Hsp70 and Hsp40 interplay regulates formation,

growth and elimination of Sup35 prions. EMBO J *27*, 2712-2724.

Silva, J., and Smith, A. (2008). Capturing pluripotency. Cell *132*, 532-536.

Solis, E.J., Pandey, J.P., Zheng, X., Jin, D.X., Gupta, P.B., Airoldi, E.M., Pincus, D., and Denic,

V. (2016). Defining the Essential Function of Yeast Hsf1 Reveals a Compact Transcriptional

Program for Maintaining Eukaryotic Proteostasis. Mol Cell *63*, 60-71.

Stelzer, Y., Shivalila, C.S., Soldner, F., Markoulaki, S., and Jaenisch, R. (2015). Tracing

dynamic changes of DNA methylation at single-cell resolution. Cell *163*, 218-229.

Stewart-Ornstein, J., Weissman, J.S., and El-Samad, H. (2012). Cellular noise regulons underlie

fluctuations in Saccharomyces cerevisiae. Mol Cell *45*, 483-493.

Thorsen, T., Maerkl, S.J., and Quake, S.R. (2002). Microfluidic large-scale integration. Science

298, 580-584.

Trotter, E.W., Kao, C.M., Berenfeld, L., Botstein, D., Petsko, G.A., and Gray, J.V. (2002).

Misfolded proteins are competent to mediate a subset of the responses to heat shock in

Saccharomyces cerevisiae. J Biol Chem *277*, 44817-44825.

Unger, M.A., Chou, H.P., Thorsen, T., Scherer, A., and Quake, S.R. (2000). Monolithic

microfabricated valves and pumps by multilayer soft lithography. Science *288*, 113-116.

Vega, N.M., Allison, K.R., Khalil, A.S., and Collins, J.J. (2012). Signaling-mediated bacterial

persister formation. Nat Chem Biol *8*, 431-433.

Weinberger, L.S., Burnett, J.C., Toettcher, J.E., Arkin, A.P., and Schaffer, D.V. (2005).

Stochastic gene expression in a lentiviral positive-feedback loop: HIV-1 Tat fluctuations drive

phenotypic diversity. Cell *122*, 169-182.

Ye, X., and Weinberg, R.A. (2015). Epithelial-Mesenchymal Plasticity: A Central Regulator of

Cancer Progression. Trends Cell Biol *25*, 675-686.

Zheng, X., Krakowiak, J., Patel, N., Beyzavi, A., Ezike, J., Khalil, A.S., and Pincus, D. (2016).

Dynamic control of Hsf1 during heat shock by a chaperone switch and phosphorylation. Elife *5*.

Figure 1

Figure 2

Figure 3

Figure 4

Figure S1