1 **KCC1 Activation protects Mice from the Development of Experimental**
2 **Cerebral Malaria.**
3

Elinor Hortle[1], Fiona C. Brown[2], Stephen M. Jane[2,4,5], David J. Curtis[2,4], Brendan J.
McMorran[1], Simon J. Foote[1], and Gaetan Burgio[1]

Affiliations:
[1] Department of Immunology and Infectious Disease, John Curtin School of
Medical Research, Australian National University. Australian Capital Territory,
Australia
[2] Australian Centre for Blood Diseases, Central Clinical School, Monash
University, Melbourne, Australia;
[4] Clinical Hematology, Alfred Health, Melbourne, Australia;
[5] Department of Medicine, Central Clinical School, Monash University, Melbourne,
Australia

Corresponding author:
Dr Gaetan Burgio
The John Curtin School of Medical Research, The Australian National University
131 Garran Road, Canberra, ACT 2600, Australia
Gaetan.burgio@anu.edu.au
+61 2 612 59428
23
24

Abstract

Plasmodium falciparum malaria causes half a million deaths per year, with up to 9% of this mortality caused by cerebral malaria (CM). One of the major processes contributing to the development of CM is an excess of host inflammatory cytokines. Recently K+ signaling has emerged as an important mediator of the inflammatory response to infection; we therefore investigated whether mice carrying an ENU-induced activation of the electroneutral K+ channel KCC1 had an altered response to *Plasmodium berghei*. Here we show that $Kcc1^{M935K/M935K}$ mice are protected from the development of experimental cerebral malaria, and that this protection is associated with an attenuated INF-ɣ response. This is the first description of a K+ channel affecting the development of cerebral malaria.

Introduction

Plasmodium falciparum malaria is a major cause of mortality worldwide, leading to an estimated 429,000 deaths per year[1]. One of the most severe and lethal complications of *P. falciparum* infection is the sudden onset of seizures and/or coma known as cerebral malaria (CM). Its occurrence varies from region to region, with a case fatality rate as high as 9% of severe malaria cases in some areas [2,3]. The causes of CM are not well understood, but hypotheses include both the accumulation of parasitized red blood cells in the brain microvasculature, as well as imbalance in the pro- and anti- inflammatory responses to infection[4].

In recent years, potassium (K+) signaling has emerged as an important mediator of the immune response to infection. Several studies have shown *in vitro* that functional outwardly rectifying K+ channels are necessary for macrophage activation and production of TNFα[5,6], for activation of the NALP inflammasome[7], for the activation of T helper cells, and the formation of T regulatory cells [8-10]. The K+ content of the RBC also has a large effect on intra-erythrocytic *Plasmodium*. It has been shown that an outwardly directed K+ gradient is needed for normal parasite growth and maintenance of the parasite plasma membrane potential [11-13].

A mouse line expressing an activated form of KCC1, discovered from a large scale

ENU mutagenesis screen, has recently been described[14]. The induced mutation –

an M to K substitution at amino acid 935 of the protein – impairs

phosphorylation of neighboring regulatory threonines, leading to aberrant

activity of the transporter. The resulting increase in K+ efflux from the RBC

causes $Kcc1^{M935K}$ mice to display a microcytic anemia, with homozygous mutants

showing a 21% decrease in Mean corpuscular Volume (MCV), 8% decrease in

total hemoglobin, and 21% increase in number of red cells. Mutant cells are also

significantly less osmotically fragile[14] indicating a dehydration of the red blood

cells.

Here we use the $Kcc1^{M935K}$ mice line to investigate the effect of increased host K+

efflux on susceptibility to malaria infection. When $Kcc1^{M935K}$ mice were infected

with *Plasmodium berghei,* they showed a marked protection from the

development of experimental cerebral malaria (ECM), associated with a

significant decrease in IFNɣ and IL-10 during infection, suggesting K+ efflux

through KCC1 attenuates the inflammatory response to infection. This is the first

description of a cation transporter affecting the development of ECM.

Results

Kcc1[M935K] *has increased survival to P. berghei challenge*

$Kcc1^{M935K/M935K}$ mice were inoculated with *P. berghei* to determine their

resistance to infection. Cumulative survival and peripheral parasitemia were

monitored daily over the course of infection. When mice were infected with

$1x10^4$ *P. berghei* parasitized red cells, survival was significantly increased in the

mutants, with 100% of homozygotes surviving past day 10 of infection,

compared to 7% of WT females (P=0.0004; Figure 1A), and 11% of WT males

(P=$2.66x10^{-7}$; Figure1B). Significantly lower parasitemia was observed in both

$Kcc1^{M935K}$ females and males. In females, parasitemia was reduced by 63% on day

7, 48% on day 8, and 42% on day 9 post inoculation. Parasitemia in males was

similarly reduced, by 66%, 41%, and 53% respectively (Figure 1A and B). To

determine if this reduction in parasitemia was caused by impaired parasitic

invasion of $Kcc1^{M935K/M935K}$ RBCs and survive within them, we conducted TUNEL

staining of infected RBCs to detect fragmented nuclei in the parasites indicative

of maturation arrest, and an *in vivo* invasion assay as previously described[15]. No

significant differences were observed in either assay (Figure 1C and D),

suggesting the $Kcc1^{M935K}$ mutation does not affect parasite invasion or growth.

Kcc1^{M935K} *is resistant to ECM*

When infected with *P. berghei* in the experiments described above, most WT

mice died 8 to 10 days after infection. To determine if the WT mice dying from *P.*

berghei infection were succumbing to experimental cerebral malaria (ECM), mice

were injected with *P. berghei* and symptoms of ECM were scored according to

severity, from 0 (no symptoms) to 5 (death). Severe clinical symptoms were

observed in WT mice, with most dying from seizures, whereas Kcc1$^{M935K/M935K}$

remained asymptomatic for the length of the experiment (Figure 2B). One of the

key hallmarks of cerebral malaria is breakdown of the blood brain barrier.

Therefore, mice were injected intravenously with Evan's Blue to assess blood

brain barrier integrity. Infected WT mice showed an average of 14.5±2.9 grams

of dye per gram of brain tissue, which was significantly higher than the

8.1±1.4g/g observed in non-infected mice. Infected Kcc1$^{M935K/M935K}$ did not differ

significantly from non-infected mice with 9.4 ± 1.3 g/g (Figure 2A and C). These

results indicate a significant breakdown of the blood brain barrier in WT mice

that is not observed in the mutants. Together with the clinical scores, this

suggests that Kcc1$^{M935K/M935K}$ mice are resistant to the development of ECM.

Kcc1^{M935K} *has an abnormal immune response to infection*

It has been shown that depletion of CD4+ T cells, CD8+ T cells, and inflammatory

monocytes can prevent the development of ECM [16-18]. ECM resistance is also

observed in mice with impaired thymic development of CD8+ T cells[19]. Because

KCC1 is expressed ubiquitously[20], it was hypothesized that the *Kcc1^M935K*

mutation might cause alterations to some of these immune cell populations.

Therefore, the relative proportion of CD3+, CD11c+, and CD19+ cells were

measured by flow cytometry in the blood, spleen and thymus, both in non-

infected mice, and day 8 of *P. berghei* infection where the inflammatory response

is expected to be highest[21,22]. No significant differences in these cell populations

were observed in any of these tissues, before or during infection (Figure 3).

CD3+ cells were further analyzed to determine the relative proportion of CD4+

and CD8+ subsets. Again, no significant differences were observed. In WT mice

the average proportion of CD4+ cells rose with infection in the blood, spleen, and

thymus, from 41 to 62%, 37 to 75%, and 40 to 56% respectively. This trend was

also observed in *KCC1^M935K/M935K* where CD4+ cells rose from an average

proportion of 36 to 62%, 34 to 78%, and 35 to 48% respectively (Figure 3). The

average proportion of CD8+ also increased in the blood with infection, from 22 to

32% in WT mice, and from 28 to 31% in *KCC1^M935K/M935K*. Conversely, in both the

spleen and thymus the average proportion of CD8+ cells was similar between

uninfected and infected animals. Together we noted no difference in T-cell

response.

One of the major host processes known to contribute to the development of

cerebral malaria in *P. berghei* infection is an over-active cytokine-driven

inflammatory response. Both *in vivo* neutralisation of host molecules, and studies

with knock-out mice have shown that cerebral malaria can be prevented by

depletion of the pro-inflammatory cytokines IFN-γ [23,24] and TNFα [25], and can be

induced by depletion of the anti-inflammatory cytokine IL-10 [26]. CM resistance is

also observed in mice with defective T cell dependent IFN-γ production [19]. We

therefore measured plasma cytokine levels in infected mice over the course of

P. berghei infection.

As expected, WT mice showed strong increases in both IFN-γ and TNFα over the

course of infection, with IFN-γ peaking around the onset of symptoms at day 9.

IL-10 was variable over the course of infection, and IL-6 showed a smaller, late

stage response. *Kcc1^M935K* mice displayed similar trends, but differed significantly

from WT in several key points. On day 7 post infection, $Kcc1^{M935K}$ showed a 1.2-fold increase in TNFα, which was followed on day 9 by a 0.6-fold reduction in the amount of IFN-γ, and a 0.8-fold reduction in the amount of IL-6 compared to WT. On day 10 of infection $Kcc1^{M935K}$ displayed a 0.9-fold reduction in the amount of IL-10 compared to WT (Figure 4).

Discussion

This study provides the first evidence that host KCC1 plays a role in malaria resistance. It shows that over-activation of the transporter causes resistance to experimental cerebral malaria (ECM) in *P. berghei* infection. This is the first description of a mutation in a cation transporter that has an effect on ECM; other previously discovered genes have involved host cytokines, antigen presentation [27,28], or erythrocyte membrane proteins [29-31].

Whereas the vast majority of non-mutant littermates succumbed to a classical ECM, all homozygous mutant animals survived the infection and did not develop any symptoms of the syndrome. Resistance in the $Kcc1^{M935K/M935K}$ animals was accompanied with a significantly lower parasitemia than WT during the first 10 days of infection. However the mutation did not appear to have a red cell autonomous effect on either parasite invasion or survival within the RBC. A possible but untested explanation for the reduced parasitemia effect is the mutation affects the cytoadherence of infected red cells to the endothelium of blood vessels. Lower levels of sequestration in the mutants would leave more late stage parasites vulnerable to splenic clearance, and therefore result in reduced parasite burden[32]. Reduced sequestration would also be consistent with protection from cerebral malaria, as infected cells would be less likely to adhere within the microvasculature of the brain.

All of the T cell populations examined in the spleen, thymus and circulation were similar between mutant and WT mice, and none changed appreciably during the *Plasmodium berghei* infection. However, production of several cytokines previously shown to direct the ECM response [23-25] were shown to be attenuated

in the mutant animals, particularly IFN-γ and IL10. It is not clear from this study

exactly what is causing decreased INF-γ and IL-10 in *Kcc1^M935K^*. This reduced

inflammation may simply be the result of the lower parasitemia observed in

mutants. However, KCC1 is expressed on a wide range of immune cells, and this

may affect their function. Previous studies have shown that K+ efflux can alter

cellular cytokine production[5-7]; can increase assembly of the NALP

inflammasome in response to pathogen associated proteins[7]; and is essential

macrophage migration[33]. All of these may help contribute to both the attenuated

inflammatory response, and the lower parasitemia observed in *Kcc1^M935K^* mice.

Here we have shown that activation of KCC1 causes a dramatic resistance to

P. berghei by completely preventing the development of experimental cerebral

malaria (ECM). This is the first description of a mutation in a transporter that has

an effect on ECM. Previous studies have shown that pharmacological activation

of KCC channels is achiavable[34,35], therefore future research into KCC1 activation

may provide novel treatments for cerebral malaria.

Methods

Animals

Mice were bred under specific pathogens free conditions. All procedures

conformed to the National Health and Medical Research Council (NHMRC) code

of practice. All mouse procedures have been approved by the Australian National

University Animal Experimentation Ethics Committee (AEEC A2014/054). The

Kcc1^M935K^ mutation is carried on a mixed BALB/c and C57BL/6 background[14].

These two mouse strains differ in their susceptibility to *P. berghei*, and this

introduced a greater amount of variability into results than is usually observed.

Therefore, WT x WT and Kcc1^M935K/M935K^ x Kcc1^M935K/M935K^ breeding pairs were

maintained. To exclude the possibility that the resistance phenotype was due to

the mixed background, and carried by chance in mutant breeding pairs,

Kcc1^M935K^ was periodically crossed back to WT, and new WT x WT and

Kcc1^M935K/M935K^ x Kcc1^M935K/M935K^ pairs established from the progeny.

Infections

Experiments used either the rodent parasite *P. berghei ANKA*. Parasite stocks were prepared from passage through resistant SJL/J mice, as described previously[36]. Experimental mice were infected intraperitoneally at a dose of $1x10^4$ parasitised RBC. Blood stage parasitemia was determined by counting thin smears from tail blood stained in 10% Giemsa solution. A least 300 cells were counted per slide.

Histology

Thin tail smears from *P. berghei* infected mice were fixed in 100% MeOH, and stained with an APO-BrdU TUNEL assay kit according to the manufacturer's instructions (Invitrogen, Carlsbad, CA). Slides were examined on an upright epifluorescence microscope (ZIESS) 600x magnification. 10 fields of view were counted for each slide.

Evans Blue

P. berghei infected mice, and uninfected controls, were injected IV with 200µl 1% Evans Blue/PBS solution. 1hr post injection, mice were sacrificed and their brains collected and weighed. Brains were placed in 2ml 10% neutral buffered formalin at room temperature for 48hrs to extract dye. 200µl of formalin from each brain was then collected and absorbance measured at 620nm. Amount of Evans blue extracted per gram brain tissue was calculated using a standard curve ranging from 40µg/ml to 0µg/ml. Injections were carried out on the day of infection that the first mouse died.

Clinical Score

Mice were monitored three times daily, and given a score from 0 to 5 based on the type and severity of their symptoms. '0' indicated no symptoms; '1' reduced or languid movement; '2' rapid breathing and/or hunched posture; '3' ruffled fur, dehydration and/or external bleeding; '4' fitting and/or coma; '5' death. Mice were considered comatose if they were unable to right themselves after being placed on their side. The highest score recorded for each mouse on each day was used to generate daily averages.

Cytokines

Peripheral blood was taken either by cardiac puncture or mandibular bleed and centrifuged for 4 minutes at 11,000xg. Plasma was then taken into a separate tube and stored at -20°C until needed. Cytokine analysis was conducted on un-diluted plasma using a CBA Mouse Th1/Th2/Th17 Cytokine Kit to the maker's instructions (BD biosciences).

Lymphocyte Analysis

Peripheral blood was taken either by cardiac puncture or mandibular bleed, and lymphocytes were isolated on Ficoll-Paque™ according to the maker's instructions. Lymphocytes were then incubated with Fc-block in MT-FACS for 10 minutes at 4°C.

Both spleen and thymus were prepared for flow cytometry using the same method. $\frac{1}{2}$ of each organ was passed through a 70μm BD Falcon® Cell Strainer with 0.5 ml of MT-FACS buffer, and then centrifuged at 300xg for 5 minutes at 4°C. The supernatant was removed and the pellet re-suspended in 5 ml cold MT-FACS buffer. A 200μl aliquot of this suspension was then incubated with 0.8μl Fc-block.

Blood, spleen and thymus samples were then stained with CD4-PacificBlue, CD11-PE, CD8-FITC, CD25-PECy7, CD19-PERCPCy5.5 and CD3-APC-Cy7, then fixed with 300μl of MT-PBS containing 1% formalin and 1% BSA for 10 minutes at 4°C. Cells were then permeabilised with MT-PBS containing 0.1% saponin and 1%BSA at 4°C for 20 minutes. Finally, cells were stained with FoxP3-APC. Samples were acquired using a BD FACSAria™ II flow cytometer, and analysed using BD FACSDiva™ software (BD Biosciences).

TUNEL staining

Thin tail smears from *P. chabaudi* infected mice were fixed in 100% MeOH. For all TUNEL staining, an APO-BrdU TUNEL assay kit was used (Invitrogen, Carlsbad, CA). Slides were initially washed three times with 1ml of wash buffer solution before being incubated with 50μl of DNA labelling solution mix (according to manufacturer's instructions) overnight at room temperature. The next day slides were rinsed three times with 1ml rinse solution with two minutes incubation time each. Stained sections were then incubated with 100μl of BrdU-antibody for one hour in the

dark at room temperature with subsequent washes in 1%BSA/ 1x MT-PBS (three
times with a two minute soak each). Slides were then further labelled with 100μl of
nuclear yellow (Invitrogen, Carlsbad, CA) (1:5000 in 1% BSA/ 1 x MT-PBS) for one
minute in the dark. After further washes with 1ml MT-PBS (three times with a two
minute soak each) sections were affixed with Fluorescent Mounting Medium
(DakoCytomation). Once the medium dried, slides were examined on an upright
epifluorescence microscope (ZIESS) 600x magnification. 10 fields of view were
counted for each slide.

In-vivo Invasion Assay

Blood from Mutant and WT uninfected mice was collected by cardiac puncture.
1800μl of blood was collected and pooled for each genotype, then halved and stained
with either NHS-Atto 633 (1μl/100μl) or sulfobiotin-LC-NHS-Biotin (1μl/100μl of
25mg/ml in DMF). Cells were then incubated at RT for 30 minutes, and washed twice
in MTPBS. Stained cells were combined in equal proportions to achieve the following
combinations:

1) WT-Biotin + Mutant-Atto 2) WT-Atto + Mutant Biotin

Combined cells were then resuspended in 2ml MTPBS, and injected intravenously
into WT *P. berghei* infected mice. 4 mice at 1-5% parasitemia, plus 1 uninfected
control, were injected with 200μl dye combination 1; the same numbers of mice were
injected with 200μl dye combination 2. Injections were carried out when parasites
were undergoing schizogeny, at ~1am.

30 minutes post injection, 1μl tail blood was collected and stained for 30 minutes at
4°C in 50μl MTPBS containing 0.25μl CD45-APC-Cy7, 0.25μl CD71-PE-Cy5, 0.5μl
Step-PE-Cy7. Next, 400μl MTPBS containing 0.5μl Hoechst 33342 and 1μl
800μg/ml Thiazole orange was added, and cells were incubated for a further 5
minutes at 4°C. Stained cells were then centrifuged at 750xg for 3 minutes, re-
suspended in 700μl MTPBS, and analyzed on a BD Fortessa Flow Cytometer. $2x10^6$
cells were collected for each sample, and data was analysed using FlowJo (FlowJo,
LLC, Oregon, USA).

References

WHO. World Malaria Report. 198 (World Health Organization, 2016).

Grau, G. E. & Craig, A. G. Cerebral malaria pathogenesis: revisiting parasite and host contributions. *Future microbiology* **7**, 291-302, (2012).

Rockett, K. A. *et al.* Reappraisal of known malaria resistance loci in a large multicenter study. *Nat Genet* **46**, 1197-1204, (2014).

Brooks, H. M. & Hawkes, M. T. Repurposing Pharmaceuticals as Neuroprotective Agents for Cerebral Malaria. *Curr Clin Pharmacol*, (2017).

Qiu, M. R., Campbell, T. J. & Breit, S. N. A potassium ion channel is involved in cytokine production by activated human macrophages. *Clin Exp Immunol* **130**, 67-74 (2002).

Ren, J. D. *et al.* Involvement of a membrane potassium channel in heparan sulphate-induced activation of macrophages. *Immunology* **141**, 345-352, (2014).

Petrilli, V. *et al.* Activation of the NALP3 inflammasome is triggered by low intracellular potassium concentration. *Cell Death Differ* **14**, 1583-1589, (2007).

Bittner, S. *et al.* Upregulation of K2P5.1 potassium channels in multiple sclerosis. *Ann Neurol* **68**, 58-69, doi:10.1002/ana.22010 (2010).

Fellerhoff-Losch, B. *et al.* Normal human CD4(+) helper T cells express Kv1.1 voltage-gated K(+) channels, and selective Kv1.1 block in T cells induces by itself robust TNFalpha production and secretion and activation of the NFkappaB non-canonical pathway. *J Neural Transm (Vienna)* **123**, 137-157, (2016).

Wulff, H. *et al.* The voltage-gated Kv1.3 K(+) channel in effector memory T cells as new target for MS. *J Clin Invest* **111**, 1703-1713, (2003).

Staines, H. M., Ellory, J. C. & Kirk, K. Perturbation of the pump-leak balance for Na(+) and K(+) in malaria-infected erythrocytes. *Am J Physiol Cell Physiol* **280**, C1576-1587 (2001).

Brand, V. B. *et al.* Dependence of Plasmodium falciparum in vitro growth on the cation permeability of the human host erythrocyte. *Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology* **13**, 347-356, (2003).

Allen, R. J. & Kirk, K. The membrane potential of the intraerythrocytic malaria parasite Plasmodium falciparum. *J Biol Chem* **279**, 11264-11272, (2004).

Brown, F. C. *et al.* Activation of the erythroid K-Cl cotransporter Kcc1 enhances sickle cell disease pathology in a humanized mouse model. *Blood* **126**, 2863-2870, (2015).

Lelliott, P. M., McMorran, B. J., Foote, S. J. & Burgio, G. In vivo assessment of rodent Plasmodium parasitemia and merozoite invasion by flow cytometry. *J Vis Exp*, e52736, (2015).

Claser, C. *et al.* CD8+ T cells and IFN-gamma mediate the time-dependent accumulation of infected red blood cells in deep organs during experimental cerebral malaria. *PLoS One* **6**, e18720, (2011).

Hermsen, C., van de Wiel, T., Mommers, E., Sauerwein, R. & Eling, W. Depletion of CD4+ or CD8+ T-cells prevents Plasmodium berghei induced

370		cerebral malaria in end-stage disease. *Parasitology* **114 (Pt 1)**, 7-12
371		(1997).
372	18	Schumak, B. *et al.* Specific depletion of Ly6C(hi) inflammatory monocytes
373		prevents immunopathology in experimental cerebral malaria. *PLoS One*
374		**10**, e0124080, (2015).
375	19	Bongfen, S. E. *et al.* An N-Ethyl-N-Nitrosourea (ENU)-Induced Dominant
376		Negative Mutation in the JAK3 Kinase Protects against Cerebral Malaria.
377		*PloS one* **7**, e31012, (2012).
378	20	Gillen, C. M., Brill, S., Payne, J. A. & Forbush, B., 3rd. Molecular cloning and
379		functional expression of the K-Cl cotransporter from rabbit, rat, and
380		human. A new member of the cation-chloride cotransporter family. *J Biol*
381		*Chem* **271**, 16237-16244 (1996).
382	21	Hanum, P. S., Hayano, M. & Kojima, S. Cytokine and chemokine responses
383		in a cerebral malaria-susceptible or -resistant strain of mice to
384		Plasmodium berghei ANKA infection: early chemokine expression in the
385		brain. *Int Immunol* **15**, 633-640 (2003).
386	22	Lacerda-Queiroz, N. *et al.* Inflammatory changes in the central nervous
387		system are associated with behavioral impairment in Plasmodium
388		berghei (strain ANKA)-infected mice. *Exp Parasitol* **125**, 271-278, (2010).
389	23	Yanez, D. M., Manning, D. D., Cooley, A. J., Weidanz, W. P. & van der Heyde,
390		H. C. Participation of lymphocyte subpopulations in the pathogenesis of
391		experimental murine cerebral malaria. *J Immunol* **157**, 1620-1624 (1996).
392	24	Amani, V. *et al.* Involvement of IFN-gamma receptor-medicated signaling
393		in pathology and anti-malarial immunity induced by Plasmodium berghei
394		infection. *Eur J Immunol* **30**, 1646-1655 (2000).
395	25	Grau, G. E. *et al.* Tumor necrosis factor (cachectin) as an essential
396		mediator in murine cerebral malaria. *Science* **237**, 1210-1212 (1987).
397	26	Kossodo, S. *et al.* Interleukin-10 modulates susceptibility in experimental
398		cerebral malaria. *Immunology* **91**, 536-540 (1997).
399	27	Koch, O. *et al.* IFNGR1 gene promoter polymorphisms and susceptibility
400		to cerebral malaria. *J Infect Dis* **185**, 1684-1687, (2002).
401	28	Hill, A. V. *et al.* Common west African HLA antigens are associated with
402		protection from severe malaria. *Nature* **352**, 595-600, (1991).
403	29	Cortes, A., Benet, A., Cooke, B. M., Barnwell, J. W. & Reeder, J. C. Ability of
404		Plasmodium falciparum to invade Southeast Asian ovalocytes varies
405		between parasite lines. *Blood* **104**, 2961-2966, (2004).
406	30	Fischer, P. R. & Boone, P. Short report: severe malaria associated with
407		blood group. *Am J Trop Med Hyg* **58**, 122-123 (1998).
408	31	Fry, A. E. *et al.* Common variation in the ABO glycosyltransferase is
409		associated with susceptibility to severe Plasmodium falciparum malaria.
410		*Hum Mol Genet* **17**, 567-576, (2008).
411	32	Simpson, J. A., Aarons, L., Collins, W. E., Jeffery, G. M. & White, N. J.
412		Population dynamics of untreated Plasmodium falciparum malaria within
413		the adult human host during the expansion phase of the infection.
414		*Parasitology* **124**, 247-263 (2002).
415	33	Kan, X. H. *et al.* Kv1.3 potassium channel mediates macrophage migration
416		in atherosclerosis by regulating ERK activity. *Arch Biochem Biophys* **591**,
417		150-156, (2016).

34   Delpire, E. & Kahle, K. T. The KCC3 cotransporter as a therapeutic target
for peripheral neuropathy. *Expert Opin Ther Targets* **21**, 113-116, (2017).
35   Yamada, K. *et al.* Small-molecule WNK inhibition regulates cardiovascular
and renal function. *Nat Chem Biol* **12**, 896-898, (2016).
36   Lelliott, P. M., Lampkin, S., McMorran, B. J., Foote, S. J. & Burgio, G. A flow
cytometric assay to quantify invasion of red blood cells by rodent
Plasmodium parasites in vivo. *Malar J* **13**, 100, (2014).
425
426

Acknowledgement

We would like to acknowledge Shelley Lampkin and Australian Phenomics Facility (APF) for the maintenance of the mouse colonies. This study was funded by the National Health and Medical Research Council of Australia (Program Grant 490037, and Project Grants 605524 and APP1047090), the National Collaborative Research Infrastructure Strategy (NCRIS), the Education Investment Fund from the Department of Education and Training, the Australian Phenomics Network, Howard Hughes Medical Institute and the Bill and Melinda Gates Foundation.

Author Contribution Statement:

E.J.H, B.J.M, S.F.J and G.B designed and planed the experimental work. E.J.H, and F.C.B performed the research. E.J.H, S.M.J, D.J.C. B.J.M, S.F.J and G.B interpreted and analyzed the data. E.J.H and G.B performed the statistical analysis. E.J.H and G.B wrote the manuscript. All authors reviewed the manuscript.

Competing Financial Interests:

The authors declare no competing financial interests.

Figure 1: The *Kcc1^{M935K}* mutation causes resistance to *P. berghei*. (A) and **(B)** Cumulative
survival and average ± SEM parasitemia for WT and *Kcc1^{M935K/M935K}* in male and female mice
respectively. WT male n=28, WT female n=8, *Kcc1^{M935K/M935K}* male n=14, *Kcc1^{M935K/M935K}* female
n=10. Combined results of two independent experiments. *P<0.05, **P<0.01, ***P<0.001.
P values calculated using Log rank test or the student's T-test. **(C)** average ± SEM percentage of
parasites which are TUNEL positive. **(D)** Average ± SEM fold change in parasitemia of
Kcc1^{M935K/M935K} labelled cells compared to WT labelled cells injected into the same *P. berghei*
infected host (n=8)

Figure 2: The *Kcc1^{M935K}* mutation causes resistance to cerebral malaria. **(A)** Clinical score
for WT (n=37) and *Kcc1^{M935K/M935K}* (n=24) mice infected with *P. berghei* 0= no symptoms, 1=
reduced movement, 2= rapid breathing/hunched posture, 3= ruffled fur/external bleeding, 4=
fitting/coma, 5= death. **(B)** Amount of Evan's Blue dye extracted from *P. berghei* infected
Kcc1^{M935K/M935K} (n=8), WT (n=7) and uninfected (n=4) brains. **(C)** Representative brains dissected
from mice injected with Evan's Blue dye. Values are average ± SEM. *P<0.05, **P<0.01,
***P<0.001. P values calculated using the Student's T-test.

Figure 3: The *Kcc1^{M935K}* mutation does not alter immune cell populations (A) Average ±
SEM proportion of lymphocytes that are CD3+, CD11c+, and CD19+ in the blood spleen and
thymus. **(B)** Average ± SEM proportion of CD3+ cells that are CD4+, CD8+, and CD4+CD8+ in the
blood spleen and thymus. Uninfected WT (n=5), uninfected *Kcc1^{M935K/M935K}* (n=5), infected WT
(n=3), infected *Kcc1^{M935K/M935K}* (n=3).

Figure 4: The *Kcc1^{M935K}* mutation alters the inflammatory response to *P. berghei* infection.
Average ± SEM concentration of cytokines in the plasma of WT (n=3-5)and *Kcc1^{M935K/M935K}*
(n=3-5) mice during infection with *P. berghei*. **P<0.01, Significance calculated using the
Student's T test.

483   **Figure 1**

487   **Figure 2**

493   **Figure 3**

Figure 4

510