Crossmodal congruency effect scores decrease with repeat test exposure

1
2   Satinder Gill[1], Daniel Blustein[1*], Adam Wilson[2], Jon Sensinger[1,2]
3
4   [1]Institute of Biomedical Engineering, University of New Brunswick, Fredericton, NB, Canada
5   [2]Department of Electrical and Computer Engineering, University of New Brunswick,
6   Fredericton, NB, Canada
7
8   * Corresponding author
9   Email: dan.blu@unb.ca (DB)
10

11   **Abstract**
The incorporation of feedback into a person's body schema is well established. The crossmodal
congruency effect (CCE) task is used to objectively quantify incorporation without being
susceptible to experimenter biases. This visual-tactile interference task is used to calculate the
CCE score as a difference in response time for incongruent and congruent trials. Here we show
that this metric is susceptible to a learning effect that causes attenuation of the CCE score due to
repeated task exposure sessions. We demonstrate that this learning effect is persistent, even after
a 6 month hiatus in testing. Two mitigation strategies are proposed: 1. Only use CCE scores that
are taken after learning has stabilized, or 2. Use a modified CCE protocol that decreases the task
exposure time. We show that the modified and shortened CCE protocol, which may be required
to meet time or logistical constraints in laboratory or clinical settings, reduced the impact of the
learning effect on CCE results. Importantly, the CCE scores from the modified protocol were not
significantly more variable than results obtained with the original protocol. This study highlights
the importance of considering exposure time to the CCE task when designing experiments and
suggests two mitigation strategies to improve the utility of this psychophysical assessment.
26
27   *Keywords:* Crossmodal congruency; Learning effect; multisensory integration; reaction time
28

Introduction

There is an increasing interest concerning the multisensory representation of space in

humans due to converging findings from a number of different disciplines. The crossmodal

congruency task is a visual-tactile interference task that has been used to investigate

multisensory representation of space in humans (Spence, Nicholls, Gillespie & Driver 1998;

Spence, Pavani & Driver 2004; Spence, Pavani, Maravita & Holmes 2004), including those with

brain damage (Spence, Kingstone, Shore & Gazzaniga 2001). Investigation of crossmodal

selective attention has been used to demonstrate the detrimental effects of age on the ability to

ignore irrelevant sensory information when attending to relevant sensory information (Poliakoff, Ashworth, Lowe & Spence 2006). Other studies have investigated changes in the representation of peripersonal space that are elicited by the prolonged use of hand-held tools (Holmes, Calvert & Spence 2007; Holmes & Spence 2004; Maravita, Spence, Kennett & Driver 2002). Further studies used this task to extend findings regarding physical and pointing tools to virtual robotic tools using techniques from haptics and virtual reality (Sengül et al., 2012; Sengül et al., 2013). Recently it has been used in conjunction with the rubber hand illusion paradigm to investigate realistic rubber hand incorporation into body schema (Zopf, Savage & Williams 2010; Zopf, Savage & Williams 2013). The crossmodal congruency effect (CCE) assessment has become a well-established quantitative metric of feedback incorporation that extends to a variety of applications.

Typically, CCE study consists of a participant holding two foam blocks in either hand with vibrotactile targets and visual distractors embedded in the top and bottom of each foam block. A trial consists of a random and independent presentation of a single vibrotactile target paired with a single visual distractor from any of the four possible pairs of locations. Participants are instructed to make a speeded response regarding the elevation of the vibrotactile target (i.e. "up", at the index finger verses "down", at the thumb), while simultaneously ignoring visual distractors (Spence et al., 2001). Participants are typically slower at discriminating the elevation of vibrotactile targets when visual distractors are presented from an incongruent elevation (i.e. when vibrotactile target and visual distractors are presented from different elevations) as compared to a congruent elevation. The CCE score is calculated as a difference in the reaction time for congruent and incongruent trials and is used as a quantitative performance metric for multisensory representation of space. It has been demonstrated that CCE scores are typically

higher when the spatial separation between visual distractor and vibrotactile targets is low (e.g.

both locations on same block vs. locations on two different blocks held in different hands)

(Spence, Pavani & Driver 2004; Maravita, Spence & Driver 2003). CCE is a simple stereotypical

behavioral task that provides a robust performance metric of feedback incorporation.

Despite the widespread use of the crossmodal congruency task in a variety of experimental

paradigms, important questions related to task exposure times remain unanswered. For example,

in a majority of the studies the experimental protocol consists of multiple blocks of trials with

experimental sessions lasting up to approximately 60 minutes. These time-intensive experimental

sessions might result in extended learning of the task and adversely affect a participant's CCE

score. Also, there is no specific mention of a participant selection criterion based on previous

knowledge of the crossmodal congruency task. To our knowledge, there exist no studies to date

that have investigated modulation of CCE score due to repeated task exposures. If a task learning

effect exists for the CCE, results from previous studies that ignored such an effect may be

misinterpreted. Participants tested with previous CCE experience would be expected to have

lower CCE scores than participants naïve to the assessment. A learning effect that has been

ignored by the field may render inaccurate our current understanding of multisensory space

representation in humans. In this study we present evidence that CCE scores decrease with

repeated task exposure sessions and that this learning effect is persistent over time. We propose

two mitigation strategies, including a shortened CCE protocol that we compare to the established

protocol. Best practices for future CCE score use are suggested.

Materials and Methods

CCE test implementation

During CCE testing, participants sat comfortably on a chair in front of a table in a dimly illuminated room. They wore over-ear noise-canceling headphones playing Brownian noise to mask background noise. Identical vibrotactile stimulation motors were placed on the thumb and index fingertips of the participant's right hand (see the Materials and Apparatus section for hardware details). Distractor light emitting diodes (LEDs) were placed on the thumb and index finger of the participant's right hand while a fixation LED was centered between the distractor LEDs using a plastic mounting strip. Fig 1 shows the system setup used for this experiment.

At the beginning of the CCE assessment, participants were instructed to make a speeded response to vibrotactile targets that were presented randomly to either the thumb or index fingertip. Participants responded by pressing the left or right foot pedals to indicate stimulation of the thumb or index finger, respectively. Participants were explicitly instructed to respond as fast as possible, while making as few errors as possible. Participants were also informed that visual distractors would be presented simultaneously with the vibrotactile targets but that they were completely irrelevant to the vibrotactile target discrimination task. They were specifically instructed to ignore visual distractors by keeping their eyes open and fixating their vision on the central fixation LED. The fixation LED was presented 1000ms before the vibrotactile targets and visual distractors were simultaneously presented for 250ms.

The CCE test session was preceded by a practice session consisting of three blocks of 10 trials each. The first block of the practice session consisted of presenting the fixation LED and vibrotactile targets only (i.e. no visual distractors were presented) so that participants could

familiarize themselves with the vibrotactile target discrimination. The visual distractors were

presented along with vibrotactile targets in the following two blocks of the practice session. This

was followed by the experimental session during which 64 trials were presented to each

participant during each block of trials. Either four or eight blocks of trials were presented

depending upon the particular experiment. Each trial ended when the participant responded by

pressing one of the foot pedals or no response was made within 1500ms of target onset.

Participants

A total of 30 healthy participants were recruited from the local community (23 male, 7

female; aged 19 – 57 years, mean ± SD = 27.5 ± 7.8 years; 26 right hand dominant, 1 left hand

dominant, 3 reported no particular hand dominance). All participants had normal or corrected to

normal vision, no disorder of touch, were able to use both foot pedals, and were naïve to the

CCE task. Participants were informed about the general purpose of the research and were given

the opportunity to ask questions but they were not aware of the specific goals of the research.

Experiments were conducted under human ethics approval at the University of New Brunswick

(Fredericton, NB, Canada) and the U.S. Department of the Navy's Human Research Protection

Program. Written informed consent was obtained from each participant before conducting

experiments and no compensation was provided to participants.

Materials and apparatus

The experimental test platform was designed around the National Instruments (NI) myRIO

embedded hardware system to achieve millisecond timing accuracy. The three digital outputs of

the myRIO system were used to drive one 3mm green LED, termed the fixation LED, and two

3mm green LEDs named distractor LEDs. Two 308-107 Pico Vibe 8mm vibratory motors from

Precision Microdrives were used as the vibrotactile targets for the thumb and index finger. The

analog outputs of the myRIO system were used to drive STMicroelectronics L272 power

operational amplifiers that drove the vibratory motors. The amplifiers were necessary to provide

sufficient driving current to the vibratory motors. Two OnStage KSP100 keyboard sustain pedals

were interfaced to the myRIO system through digital inputs to measure a participant's speeded

responses to the vibrotactile targets. NI LabVIEW was used to develop the firmware for the

myRIO system to randomly activate the visual distractors and vibrotactile targets at a fixed

interval and measure the speeded response time.   A desktop NI LabVIEW graphical user

interface (GUI) was designed to interact with the myRIO embedded system, allowing the

experimenter to set various experimental parameters such as vibratory stimulus amplitude, the

number of trials and to specify the filename to record the timing results of speeded responses.

Experiment design

Three experiments were run to examine CCE learning effects. In Experiment 1 we assessed

the learning effect over five exposures to the CCE task. Twelve subjects each completed five

consecutive days of CCE testing using the standard 8-block protocol (Spence, Pavani & Driver

2004). The number of sessions and subjects was determined using a power analysis using

preliminary CCE results from eight pilot subjects (see Appendix). Subjects were naïve to the

CCE and testing was run at about the same time each day. Before each test session, generalized

reaction time was measured using a ruler drop task. Subjects held their thumb and first finger 4

cm apart with a vertically-aligned ruler held by the experimenter set so the bottom of the ruler

was just within the grasp of the subject. The subject was asked to grasp the ruler as quickly as

possible after they noticed it falling. The researcher dropped the ruler at an unspecified and

variable time and recorded the distance fallen as a proxy for reaction time. The 'reaction

distance' was calculated as the average result of three ruler drop trials. In Experiment 2, the

persistence of the CCE learning effect was assessed. Eight subjects from Experiment 1 returned

for an additional CCE test session after more than six months with no CCE exposure. For both

Experiments 1 and 2, all CCE sessions consisted of eight blocks of 64 trials each. Each block

lasted for approximately four minutes and a break period of two minutes was provided between

consecutive blocks.

In Experiment 3, we examined the effect of reducing the number of CCE blocks per

exposure from eight to four. 18 subjects were tested across two exposures on consecutive days;

half of the subjects (n=9) were randomly assigned to test with four blocks and the other nine

subjects tested with eight blocks for each session.

Analysis

Practice session trials were not analyzed. To calculate the CCE score from the test trials,

we first discarded trials with an incorrect response, trials with a premature response (i.e. reaction

time less than 200ms), and trials with a delayed response (i.e. reaction time greater than 1500ms)

as these conditions most likely occurred due to lapses in attention (Spence, Pavani & Driver

2004; Sengül et al., 2012). The remaining trials in each block were used to calculate the mean

congruent and mean incongruent reaction times. The CCE score for each block was calculated by

taking the difference between mean incongruent and congruent reaction times. The mean of the

block CCE scores was used to calculate the CCE score for each participant for a particular

exposure session. Selection error rates were used as a separate metric of analysis.

In Experiment 1, attenuation across sessions was evaluated using a one-way repeated-

measures ANOVA. For Experiment 2, we ran an intraclass correlation coefficient analysis [two-

way mixed model with single measurements] and calculated the standard error of measurement

[SEM = SD$\sqrt{(1\text{-}ICC)}$] to assess the test-retest reliability between the 6+ month follow-up visit

and the Day 5 visit (Weir 2005). Experiment 3 data were analyzed using paired t-tests within the

data   collected   using   each   protocol.   All   data   are   available   on   Dryad   at

doi:10.5061/dryad.150v8g3.

183   **Results**

Experiment 1: CCE score decreases over repeated exposures

We measured the CCE score of 12 subjects across five sessions using the conventional

CCE method of 8-blocks. We observed a significant effect of exposure number on CCE score

determined with a repeated-measures ANOVA with Greenhouse-Geisser correction (Fig 2)

($F(2.00,21.95)$=6.93, p = .005, partial-eta=.39). Tests of within-subjects polynomial contrasts

over exposure number indicated a significant linear trend ($F(1,11) = 8.63$, p=.013), a significant

4th order trend ($F(1,11) = 5.14$, p=.044), and a quadratic trend ($F(1,11) = 4.83$, p=.05). The

statistically-supported trends match the observed initial decrease in CCE score and subsequent

stabilization, indicating a task learning effect (Fig 2).

To verify that the change in CCE score was due to a learning effect and not attributed to

other interacting factors, we analyzed the CCE selection error rates, generalized reaction times

measured with a ruler grasp task, and overall reaction times for congruent and incongruent

stimuli. We found no significant effect of exposure number on CCE error rate (Fig 3a), as

indicated by a repeated-measures ANOVA with Greenhouse-Geisser correction

(F(1.15,12.66)=0.53, p = .51). Repeated-measures ANOVA with Greenhouse-Geisser correction

indicated a significant effect of exposure on both incongruent reaction time

(F(2.16,23.78)=13.57, p <.01) and congruent reaction time (F(1.46,16.07)=8.89, p<.01) (Fig. 3b).

To summarize, the only metrics presenting similar exposure-dependent decreases as the CCE

scores (Fig 2) were the congruent and incongruent reaction times (Fig 3b). We found no

significant effect of exposure number on generalized reaction time as measured by a ruler drop

fall-to-grasp distance, as indicated by a repeated-measures ANOVA with Greenhouse-Geisser

correction (F(1.463,14.63)=1.269, p = .29).

Experiment 2: CCE learning effect persists over time

After observing a statistically significant CCE learning effect, observed in both

incongruent and congruent reaction times but not in error rates and generalized reaction times,

we wanted to see if the learning effect persisted over longer time periods. Eight subjects who

completed Experiment 1 were re-tested between 6 and 7 months after their initial testing. Re-

visit CCE scores were similar, and even slightly lower, than the CCE scores measured on

Exposure 5, about 6 months prior (Fig 4). Similarity between Exposure 5 and the re-visit session

was indicated by a high intraclass correlation coefficient (ICC(3,1) = .71) and a low standard

error of measurement (SEM = 21.6ms) compared to baseline reliability data calculated by

comparing exposure 1 and re-visit results (ICC(3,1) = .46; SEM = 45.7ms).

Experiment 3: Modified CCE protocol reduces learning effect

We next sought to explore mitigation strategies to diminish the impact of the CCE learning

effect on research results. The persistence of the learning effect suggests that CCE scores should

only be considered after task learning has stabilized. However, in certain research or clinical

settings with subject access or time constraints, extended testing may be impractical. In

Experiment 3 we tested a modified CCE protocol designed to reduce task exposure to determine

if the shortened testing could: 1. Produce valid CCE results; and 2. Mitigate the learning effect.

The modified protocol reduced the duration of task exposure in each session from eight

blocks to four blocks. The 4-block protocol reduced the drop in CCE score from first exposure to

second exposure (Fig 5). For the 8-block protocol, mean CCE score dropped by 14.7ms from the

first exposure to the second, compared to a 7.6ms drop for the 4-block protocol. Variability in the

second session was only slightly higher for the 4-block protocol (SD = 43.8ms) compared to the

8-block protocol (SD = 39.0ms). The variability difference was more pronounced when

analyzing the first session results [4-block SD =64.9ms; 8-block SD = 47.5ms]. None of these

differences were statistically significant (paired t-tests, p>.05).

Discussion

The CCE is a well-established quantitative measure of feedback incorporation that has

important implications for objective human sensorimotor assessments (Spence et al., 1998;

Spence, Pavani & Driver 2004; Spence, Pavani, Maravita & Holmes 2004; Spence 2015). The

resulting scores have been used to quantify the rubber hand illusion (Zopf et al., 2010) and show

promise for assessing neuroprosthetic devices (Spence 2015). Many applications of the CCE task

require repeated testing – for example, testing the same subject under different sensory feedback

conditions to see which one enables the highest degree of feedback incorporation. Oftentimes in

research or clinical settings time allotted for sensorimotor assessment can be limited. This study

sought to characterize the CCE task performance through repeat testing to inform its future

implementation.

In Experiment 1, we found that with repeat testing, CCE scores decreased over the initial

exposure sessions. The results are not explained by a fatigue effect as the CCE score trend was

not matched by the overall reaction times measured with a ruler grasp task. Different motivation

levels or strategies fail to explain the CCE score trend as CCE error rate did not show a trend

matching the decreasing CCE scores. Thus, we consider the decrease in CCE score to be

evidence of a task learning effect.

In Experiment 2, we demonstrated that the learning effect is persistent as CCE scores

after six months remained near the stabilized scores observed at the end of five days of

consecutive testing. The persistence of the learning effect provides an opportunity for researchers

to only use CCE scores collected after learning has stabilized.

In certain circumstances, additional testing to arrive at stable CCE scores may not be

feasible. In Experiment 3, we showed that a reduced-duration CCE protocol with 4 testing blocks

instead of 8 reduced the impact of the learning effect across two testing sessions. Importantly,

CCE score variability did not significantly increase with the 4-block protocol.

Reducing the number of blocks per session has the added benefit of reducing fatigue,

improving the ability of subjects to concentrate over the duration of the study, and shortening the

length of the session to approximately 30 minutes (which in turn makes it viable to test a greater

number of subjects or for use in a clinical setting). Using a modified version of the CCE with

only four blocks of 64 trials has many potential benefits, with the only drawback identified being

a slight increase in the variability of results.

Through this characterization of the learning effect associated with the CCE task, we

have provided support for two distinct mitigation strategies. If extended testing is feasible,

researchers can provide enough practice so that CCE scores are only measured and compared

after task learning has stabilized. In situations where reduced testing is required, researchers can

use the 4-block CCE protocol to reduce overall task exposure and subsequently reduce the

impact of the learning effect on the results. It is important that these strategies not be mixed;

CCE scores should only be compared to other scores that were collected in the same manner.

It is not obvious what is being 'learned' through repeated task exposure. We suspect that

subjects strengthen an internal model of the foot pedal location tied to the vibration stimulus,

resulting in faster response times. Anecdotal support for this idea comes from one subject's

comment that "When the feedback is in the opposite location to the light, I sometimes find

myself thinking about which pedal to press only to realize I've already pressed the correct

pedal". Somatosensory reorganization of an individual's body schema through training may also

be implicated (Cardinali et al., 2009).

The initial session CCE scores (i.e. before learning effect stabilization) appear to be

higher than scores reported by other researchers (Spence, Pavani & Driver 2004; Zopf, Savage &

Williams 2010). One potential explanation for this observation is that other studies may have

different baseline CCE scores due to the experimental inquiry, such as studies using spatial

misalignment (Spence, Pavani & Driver 2004) or artificial hands (Zopf, Savage & Williams

2010; Marini et al., 2014). We would expect a lower degree of feedback incorporation, and thus

a lower CCE score (Maravita et al. 2002), under these conditions compared to the ideal

conditions we used with both vibratory feedback and visual distractors aligned in place on one of

the subject's hands. Additionally, different experimental setups could account for differences in

CCE scores. Others have recorded user inputs differently, e.g with a rocking heel-toe foot pedal

setup (Spence, Pavani & Driver 2004) or with finger presses (Zopf, Savage & Williams 2010).

Other variable techniques include the exclusion of trials with extensive eye movement (over 9%

omitted in one study (Spence, Pavani & Driver 2004)), or the use of no-go trials as a control

(Zopf, Savage & Williams 2010).

Another possible explanation for the observed variability in CCE scores across studies is

the different methods used to apply feedback. Different feedback modalities result in different

CCE scores (Frigs & Spence 2010; Mayer, Franco, Canive & Harrington 2009). Even when

comparing studies using vibration feedback, the method of application can vary and may result

in CCE score differences. We used small vibratory motors (0.8cm diameter vibrating surface)

but others have used larger bone conduction vibrators (1.6cm x 2.4cm vibrating surface, (Spence,

Pavani & Driver 2004)) or small speakers (0.9cm diameter vibrating surface, (Zopf, Savage &

Williams 2010)), all could result in different degrees of feedback incorporation.

Differences in subject characteristics could also explain differences in CCE scores. All of

the subjects in this study were initially naïve to the CCE task, but it is unclear if that was an

inclusion criteria in other studies. Even comparing the first session results from this study across

the two cohorts of subjects tested with 8 CCE test blocks we see differences in mean CCE. In

Experiment 1, a mean CCE score of 140ms was observed (Fig 2) compared to 115ms in

Experiment 3 (Fig 5). To summarize, there are lots of factors that could affect CCE score and it

is impossible to determine if results reported elsewhere are capturing pre-, mid- or post-learning

effect scores.

To our knowledge, this is the first study that has demonstrated the modulation of CCE

score due to task overexposure. We recognize some limitations with this study that could warrant

additional studies. Although we collected a generalized reaction time metric with a ruler grasp

task, more sophisticated tracking of mental and physical fatigue would be helpful to better

inform our understanding of the observed learning effect. Although CCE testing for individual

participants was conducted at the same time of the day for each exposure, no strict guidelines

were used to ensure that participants were in similar mental or physical states before each test

session. Nevertheless, the learning effect we have described has important implications for future

use of the CCE task. Going forward, the learning effect must be considered when measuring

CCE scores by only using scores from after learning has stabilized, or by using a modified

protocol that reduces exposure to the task.

321   **References**

Cardinali, L., Frassinetti, F., Brozzoli, C., Urquizar, C., Roy, A.C., & Farnè, A. (2009). Tool-use

induces morphological updating of the body schema. *Current Biology, 19*(12), R478–9.

doi:10.1016/j.cub.2009.05.009

Frings, C., & Spence, C. (2010). Crossmodal congruency effects based on stimulus identity.

Brain Research, 1354, 113–22. doi:10.1016/j.brainres.2010.07.058

Holmes, N.P., Calvert, G.A., & Spence, C. (2007). Tool use changes multisensory interactions in

seconds: evidence from the crossmodal congruency task. *Experimental Brain Research*,

183(4), 465–76. doi:10.1007/s00221-007-1060-7

Holmes, N.P., & Spence, C. (2004). The body schema and the multisensory representation(s) of

peripersonal space. *Cognitive Processing, 5*(2), 94–105. doi:10.1007/s10339-004-0013-3

Maravita, A., Spence, C., & Driver, J. (2003). Multisensory integration and the body schema:

close to hand and within reach. *Current Biology, 13*(13), R531–9. doi:10.1016/S0960-

9822(03)00449-4

Maravita, A., Spence, C., Kennett, S., & Driver, J. (2002). Tool-use changes multimodal spatial

interactions between vision and touch in normal humans. *Cognition, 83*(2), B25–34.

doi:10.1016/S0010-0277(02)00003-3

Marini, F., Tagliabue, C.F., Sposito, A.V., Hernandez-Arieta, A., Brugger, P., Estévez, N.,

Maravita, A. (2014). Crossmodal representation of a functional robotic hand arises after

extensive training in healthy participants. *Neuropsychologia, 53*, 178–86.

doi:10.1016/j.neuropsychologia.2013.11.017

Mayer, A.R., Franco, A.R., Canive, J., & Harrington, D.L.. (2009). The effects of stimulus

modality and frequency of stimulus presentation on cross-modal distraction. *Cerebral*

Cortex, 19(5), 993–1007. doi:10.1093/cercor/bhn148

Poliakoff, E., Ashworth, S., Lowe, C., & Spence, C. (2006). Vision and touch in ageing:

crossmodal selective attention and visuotactile spatial interactions. *Neuropsychologia,*

44(4), 507–17. doi:10.1016/j.neuropsychologia.2005.07.004

Sengül, A., Rognini, G., van Elk, M., Aspell, J.E., Bleuler, H., Blanke, O. (2013). Force

feedback facilitates multisensory integration during robotic tool use. *Experimental Brain*

Research, 227(4), 497–507. doi:10.1007/s00221-013-3526-0

Sengül, A., van Elk, M., Rognini, G., Aspell, J.E., Bleuler, H., & Blanke, O. (2012). Extending

the body to virtual tools using a robotic surgical interface: evidence from the crossmodal

congruency task. *PLoS ONE, 7*(12), e49473. doi:10.1371/journal.pone.0049473

Spence, C. (2015). The Cognitive Neuroscience of Incorporation: Body Image Adjustment and

Neuroprosthetics. In: K. Kansaku, L. Cohen, & N. Birbaumer (Eds.), *Clinical Systems*

Neuroscience (pp. 151–68). Tokyo: Springer. doi:10.1007/978-4-431-55037-2_9

Spence, C., Kingstone, A., Shore, D.I., & Gazzaniga, M.S. (2001). Representation of visuotactile

space in the split brain. *Psychological Science, 12*(1), 90–3. doi:10.1111/1467-

9280.00316

Spence, C., Nicholls, M. E., Gillespie, N., & Driver, J. (1998). Cross-modal links in exogenous

covert spatial orienting between touch, audition, and vision. *Perception & Psychophysics,*

60(4),544–57. doi:10.3758/BF03206045

Spence, C., Pavani, F., & Driver, J. (2004). Spatial constraints on visual-tactile cross-modal

distractor congruency effects. *Cognitive, Affective, & Behavorial Neuroscience, 4*(2),

148–69. doi:10.3758/CABN.4.2.148

Spence, C., Pavani, F., Maravita, A., & Holmes, N. (2004). Multisensory contributions to the 3-

D representation of visuotactile peripersonal space in humans: evidence from the

crossmodal congruency task. *Journal of Physiology - Paris, 98*(1-3), 171–89.

doi:10.1016/j.jphysparis.2004.03.008

Weir, J.P. (2005). Quantifying test-retest reliability using the intraclass correlation coefficient

and the SEM. *Journal of Strength & Conditioning Research, 19*(1), 231–40.

doi:10.1519/15184.1

Zopf, R., Savage, G., & Williams, M.A. (2010). Crossmodal congruency measures of lateral

distance effects on the rubber hand illusion. *Neuropsychologia, 48*(3), 713–25.

doi:10.1016/j.neuropsychologia.2009.10.028

Zopf, R., Savage, G., & Williams, M.A. (2013). The Crossmodal Congruency Task as a Means

to Obtain an Objective Behavioral Measure in the Rubber Hand Illusion Paradigm.

Journal of Visualized Experiments, (77), e50530. doi:10.3791/50530

Fig 1. Overall system setup. Subjects make speeded responses with foot pedals to vibratory

feedback presented on the index finger or thumb. Subject is asked to focus on the fixation LED

while ignoring the distractor LEDs. See text for details.

Fig 2. CCE score decreases with task exposure. Each data point represents the mean CCE

score for 12 subjects. Error bars show standard error.

Fig 3. Decreasing CCE score trend matched in isolated reaction times but not in CCE error

rates. a. Plot of selection error rate over repeated exposures. An error occurred when the wrong

pedal was pressed (e.g. vibration stimulus on the thumb resulted in subject pressing the pedal

indicating an index finger stimulus). b. Congruent and incongruent reaction times plotted

independently.

Fig 4. Persistent CCE learning effect after 6-month follow-up testing. Eight subjects from

Experiment 1 returned for a sixth visit about 6 months after their initial visits. Data shown are

means and standard error for those eight subjects.

Fig 5. First exposure vs. second exposure mean CCE scores for 8-block and 4-block

protocols. Means and standard error for 8-block (n=9) and 4-block (n=9) participants. Dark

colored bars represent the first exposure session and light colored bars represent the second

exposure session.

Distractor LEDs

Fixation LED

myRIO System

Vibratory Motors

Foot Pedals