

Identification of DISE-inducing shRNAs by monitoring cellular responses

Monal Patel^a and Marcus E. Peter^{a,b}

^aDepartment of Medicine/Division Hematology/Oncology and ^bDepartment of Biochemistry and Molecular Genetics, Feinberg School of Medicine, Northwestern University, Chicago, IL 60611, USA

Contact: Marcus Peter, E-mail: m-peter@northwestern.edu. Division Hematology/Oncology Feinberg School of Medicine, Northwestern University, Chicago, IL 60611, USA

Keywords:

RNAi, tumor suppressors, cancer, CRISPR, cell death, DISE

Abstract

Off-target effects (OTE) are an undesired side effect of RNA interference (RNAi) caused by partial complementarity between the targeting siRNA and mRNAs other than the gene to be silenced. The death receptor CD95 and its ligand CD95L contain multiple sequences that when expressed as either si- or shRNAs kill cancer cells through a defined OTE that targets critical survival genes. Death induced by survival gene elimination (DISE) is characterized by specific morphological changes such as elongated cell shapes, senescence-like enlarged cells, appearance of large intracellular vesicles, release of mitochondrial ROS followed by activation of caspase-2, and induction of a necrotic form of mitotic catastrophe. Using genome-wide shRNA lethality screens with eight different cancer cell lines, we recently identified 651 genes as critical for the survival of cancer cells. To determine whether the toxic shRNAs targeting these 651 genes contained shRNAs that kill cancer cell through DISE rather than by silencing their respective target genes, we tested all shRNAs in the TRC library derived from a subset of these genes targeting tumor suppressors (TS). We now report that only by monitoring the responses of cancer cells following expression of shRNAs derived from these putative TS it was possible to identify DISE-inducing shRNAs in five of the genes. These data indicate that DISE in general is not an undefined toxic response of cells caused by a random OTE but rather a specific cellular response with shared features that points at a specific biological function involving multiple genes in the genome.

Introduction

RNA interference is a widely used tool to reduce the expression of mRNAs. RNAi is initiated by double-stranded (ds)RNAs or pre-microRNAs, which are cleaved by Dicer, an RNase III enzyme, producing 21-23 nucleotide short interfering (si)RNAs or micro (mi)RNAs respectively, containing 2nt 3' overhangs^{1, 2}. The antisense (guide) strand is then loaded onto the endonuclease argonaute 2 (Ago2) in the RNA-induced silencing complex (RISC) and directs the downstream targeting events mostly through complete complementarity between positions 2–8 (the seed region) at the 5' end of the guide strand and a matching sequence (seed match) in the 3'UTR of targeted mRNAs.³⁻⁶ While in case of miRNAs, the guide strand recruits the RISC to the 3' untranslated regions (UTRs) of partially complementary mRNAs to promote translation repression or mRNA cleavage^{7, 8}, the guide strand of an siRNA is designed to be fully complementary to the target mRNA and directs the enzymatic cleavage of the mRNA by the Ago2 protein⁹⁻¹¹. RNAi can be induced by either transfecting cells with siRNAs, or by introducing short hairpin (sh)RNAs in the form of expression vectors or viruses. Apart from the intended target, the guide strands of siRNAs also recognize many mRNAs with partial complementarity in a manner similar to miRNAs, mostly involving the guide RNA seed, and studies have suggested that 3'UTR complementarity to si/shRNA seed sequences can mediate gene silencing based on an off-target effect (OTE) both through translational repression and mRNA degradation¹²⁻¹⁵. In addition, improper loading of the sense/passenger strand can also lead to OTEs¹⁶. This can be caused by imprecise cleavage of shRNAs by Dicer prior to RISC loading¹⁶. The main causes of OTE are therefore cross-reactivities of either the guide RNA or the passenger strand loaded into the RISC^{17, 18} with transcripts of undesired genes in the genome. The goal for virtually all RNAi projects is to selectively silence targeted genes with little or no OTE. In fact, most of the latest generation siRNAs are chemically modified to increase their stability, specificity, and to reduce OTE¹⁹, and shRNAs are expressed using optimized vector systems that allow preferential loading of guide strand into the RISC^{18, 20}.

We recently reported that >80% of 22 different nonoverlapping si-, Dsi- or shRNAs derived from either CD95L or CD95 killed cancer cells by activating multiple cell death pathways^{21, 22}. Activation of the CD95/Fas surface receptor upon binding to its cognate ligand (CD95L) induces apoptosis. The CD95/CD95L system is used by immune cells to eliminate virus-infected and cancer cells through the secretion of CD95 ligand (CD95L)²³. Hence, the CD95/CD95L system has a tumor suppressive function. Interestingly, what at first appeared to be cancer cell death caused by silencing the expression of these two genes, actually turned out to be initiated by a mechanism completely independent of the presence of CD95 or CD95L gene products²². We demonstrated that the toxic si- or shRNAs derived from either CD95 or CD95L killed cells in what appeared to be a combination of apoptosis, necrosis and mitotic catastrophe, mediated by the release of mitochondrial ROS, activation of caspase-2, and DNA damage. Morphologically, most cells responded by forming oddly shaped elongated cell structures, with likely stress induced large vesicles, and anaphase bridges²¹. While some cells died as early as one day after introducing the toxic shRNAs, most cells died when attempting to divide²¹. This form of cell death could not be inhibited and cancer cells had a hard time developing resistance both *in vitro* and *in vivo*^{21, 24}. We recently presented data to suggest that cells actually die through an OTE that results in the *preferential* targeting of the 3'UTRs of a set of critical survival genes²². We have therefore named this form of cell death DISE (for death induced by survival gene elimination).

The discovery of DISE raised a number of puzzling questions: Why did the cancer cells appear to respond to the toxic shRNAs in a highly similar way? Why would an OTE not result in a variety of unintended cellular responses, depending on what gene or sets of genes are affected? In this study we set out to identify novel toxic shRNAs derived from a small subset of putative tumor suppressor genes other than CD95 and CD95L. Solely by monitoring cellular responses (morphology, biochemical changes, and ability to divide) by the cancer cells we have identified

shRNAs derived from 5 putative tumor suppressive genes that can kill multiple cancer cells by an OTE in the absence of the coded protein that resembles DISE. We propose that these RNAi active sequences can be used to kill cancer cells.

Results

A subset of genes recently found to be critical for the survival of cancer cells are tumor suppressors

Previously, based on 12 shRNA-based lethality screens of 8 human cancer cell lines/cell line variants (HeLa, S3, HeLa N10, CHP-100, FU-UR-1, HEK293, A549 EGFRB, A549, H2030), we nominated 651 out of ~18,0000 genes targeted (by ~78,000 shRNAs, individually tested) as critical survival factors for cancer cells²¹. Included were all genes for which at least 3 out of 5 shRNAs (H factor = 60) reduced cell viability more than 95% in at least 9 out of 12 independent screens²¹. Most of the 651 genes had genuine survival functions and included genes coding for ribosomal proteins, cell cycle regulators or all three RAS genes (see Table S2 in²¹). However, a survival function was not immediately obvious for a number of these genes and therefore they could be sources of DISE-inducing shRNAs. To increase the chance of finding such toxic shRNAs, we decided to focus on a subset of genes most unlikely to be required for cancer cell survival: tumor suppressors (TS). To identify potential TS among the 651 genes identified as survival genes, we compared the 651 genes with a curated list of 637 putative TS genes²⁵. This analysis resulted in 17 putative TS genes (plus CD95L) for which up to 94% of the targeting shRNAs killed a number of cancer cell lines in the shRNA lethality screen (**Fig. S1**). For each of the 17 genes, tumor suppressive activities have been described for various cancers (see legend of **Fig. S1B**).

Identification of RNAi active toxic sequences derived from certain tumor suppressors

The finding that shRNAs derived from TS can kill cancer cells suggested that they may not act by reducing protein levels of their targeted genes, but by another mechanism, possibly DISE. We therefore decided to first validate the toxicity by testing five shRNAs per gene, a total of 85 shRNAs. Because we were only interested in shRNAs that killed all cancer cells, we chose three additional cell lines for this test, which were not part of the original shRNA lethality screen: HeyA8 (ovarian cancer), T89G (glioblastoma), and HCT116 (colon cancer). The latter two cell lines were chosen because we used them before to study and biochemically characterize DISE²¹. We decided on a sequential strategy: test the shRNAs on HeyA8 cells, then test the toxic ones on T98G cells and test all shRNAs that killed these two cell lines on HCT116 cells. To identify two shRNAs per gene that killed HeyA8 cells, the 85 shRNAs (in the pLKO backbone) were screened in 96 well plates targeting the 17 TSs using a Thermo Multidrop Combi and a Tecan Freedom EVO200 for infecting cells at an MOI of five. After puromycin selection the effect on growth was monitored in the IncuCyte Zoom. For each gene, the two shRNAs that caused the strongest growth reduction were identified and used for further analysis (data not shown). This resulted in the identification of 34 toxic shRNAs targeting the 17 TS.

Because we were interested in determining if these shRNAs had similar activities and elicited cellular responses similar to the DISE-inducing shRNAs derived from CD95 or CD95L, we retested these 34 toxic shRNAs at an MOI of three on HeyA8, T98G and HCT116 cells. This was done again in the IncuCyte Zoom and growth reduction (50% reduction compared to cells infected with a nontargeting shRNA at half maximal confluency) was used as an initial surrogate marker for cell death. All 34 shRNAs targeting the 17 TS were identified as toxic to HeyA8 cells, validating the original screen. (**Fig. 1A**). Not surprisingly, while shRNAs against the tested TS were toxic, when we tested 4-5 shRNAs targeting two of the most widely studied and most highly mutated TS in human cancers, p53 and PTEN, none of them qualified as toxic shRNAs using the threshold we had defined (**Fig. 1B**). This suggested that only certain TS contain RNAi active sequences that can kill cancer cells. Of the 34 shRNAs targeting the 17 TS genes, 30 shRNAs were also toxic to T98G cells targeting 15 of the TSs (**Fig. S2A**). Because we were only interested in shRNAs that kill all three cancer cells, we only tested these 30 shRNAs on HCT116

cells (**Table 1** and **Fig. S2B**). This reduced the number of shRNAs that killed all three cell lines to 26 shRNAs (targeting 13 TS).

Toxic TS derived shRNAs trigger DISE-like cell death

To determine which of the shRNAs killed the three cell lines in a fashion similar to DISE induction observed with CD95L derived shRNAs, we compared the morphological changes seen in HeyA8 and T98G cells after infection with a lentiviral shRNA with that seen in cells infected with the CD95L derived shL3. We were unable to perform a morphological analysis in HCT116 cells since the cells were too small and without clear morphological features to distinguish. In HeyA8 cells, 3-7 days after infection, we detected the typical stress-induced elongated cell shapes (**Fig. 2A**), appearance of large intracellular vesicles, and enlargement and senescence-like cell flattening (**Fig. 2B**). 8 of the 13 remaining TS genes had shRNAs that both elicited these changes (**Fig. 2** and **Table 1**). In T98G cells, for the same 8 genes (all of the 16 shRNAs), we observed that the cells attempted to divide and immediately after that they rounded up and died (data not shown).

In order to determine whether the remaining 16 shRNAs induced cell death biochemically similar to DISE, we tested whether these shRNAs caused ROS production and caspase-2 activation in HeyA8 cells - two characteristic features of DISE seen in HeyA8 cells after infection with shL3 and also observed in multiple other cell lines after introducing multiple CD95L or CD95 targeting shRNAs²¹. For 7 of the remaining TS, both shRNAs caused significant induction of ROS and caspase-2 activation (**Fig. 3A** and **B**). Our sequential analysis in three cancer cell lines allowed us to narrow down the list of potential shRNAs that killed cancer cells by DISE to seven (**Table 1**). To confirm that all shRNAs derived from these seven genes did not just result in growth reduction but actually killed cancer cells, we quantified DNA fragmentation in HeyA8 cells 8 days after lentiviral infection (**Fig. 3C**). Indeed, all 14 shRNAs caused a significant increase in subG1 DNA, suggesting that they all at various levels killed cancer cells.

Toxic shRNAs derived from five TS genes kill cancer cells through DISE

One of the most surprising properties of DISE-inducing shRNAs is that they kill cancer cells independent of targeting the mRNA they were designed to silence; instead we reported that these sequences are toxic to cells through a unique form of OTE that targets a network of critical survival genes²². We narrowed down our list of toxic shRNAs to only include shRNAs that killed cancer cells in a way that was similar to DISE in morphology and biochemistry; we now considered whether these TS genes were enriched in shRNAs that kill cancer cells in a way independent of the expression of the coding protein, which would be indicative of the toxic OTE that is DISE. We chose to study this using HAP1 cells for two reasons: 1) They are available as knock-out cells (generated by using CRISPR/Cas9 gene editing) for most human genes (that are not essential for cell survival) and 2) We recently demonstrated that DISE-inducing shRNAs derived from either CD95 or CD95L could still kill CD95 or CD95L deficient HAP1 cells (data not shown). For five of the seven genes, both shRNAs reduced growth of unmodified HAP1 cells >50% (**Table 1, Fig. S2C**); hence these five genes could be tested in HAP1 CRISPR/Cas9 modified cells. In all HAP1 cells using CRISPR/Cas9 gene editing a frame shift mutation was introduced downstream of the translational start codon. Two of the mutant clones, ARMC10 and MAPKAPK5, were validated by Western blotting to be protein knock outs (**Fig. 4B**, far right panel). Two of the genes, SOCS3 and TMEFF1, are not expressed in HAP1 cells (Transcripts Per Kilobase Million (TPM) of less than 3 in RNA Seq analysis are considered undetectable²⁶). All of the 5 CRISPR/Cas9 modified cell lines still died after the introduction of shRNAs derived from these genes (**Fig. 4A** and **4B, Table 1**). Because for two of the genes the Western blot confirmation of a complete knockout was inconclusive (MAPKAPK5- multiple bands; and

TGFBR2 - no band), an additional k.o clone was generated and tested. For MAPKAPK5 an out-of-frame deletion was introduced into exon 8 and for TGFBR2 in exon 4 (data not shown). Both clones were as sensitive to the two toxic shRNAs derived from these genes as wt cells (data not shown). The data indicate that all of the toxic shRNAs we identified derived from the five TS killed the cells through an OTE. Because the result of this OTE is cell death and because this cell death in all tested cell lines resembled DISE we conclude that these genes contain toxic sequences that can kill cancer cells by DISE. These data suggest that CD95 and CD95L are not unique and that the human genome likely contains multiple genes that contain sequences that have DISE inducing activities when expressed as small double stranded RNAs.

Discussion

RNAi has become one of the most utilized methods to study the function of genes. Countless reports of gene-specific silencing and genome-wide screens document the power of RNAi^{21, 27-30}. However, one caveat of RNAi screens is the OTE. This can be caused by cross-reactivities between the guide strand of the siRNA and the target mRNAs due to partial complementarity. In addition, OTE can be caused by unintended loading of the passenger strand into the RISC. OTE has been described to occur¹⁶ and was found to affect many genes and to only require a complementarity of 6-7 nucleotides between the targeting si/shRNA and the affected mRNAs¹²⁻¹⁴. However, studies were unable to predict which genes would be and which genes would not be affected by OTE¹³. This finding is consistent with the assumption that OTEs are truly random. When all OTEs are truly random, one would expect cells to respond in various ways depending on the mRNAs affected by the OTE. Similarly, when OTEs lead to cell death, one would assume that different forms of cell death with different morphologies and different signaling pathways would be activated.

We recently identified a general OTE that results in the death of most tested cancer cells. It was found to preferentially affect transformed cells²¹, and among them cancer stem cells³¹. Interestingly, this OTE preferentially affected genes that are critical for cancer cell survival. We named this form of cell death DISE (death induced by survival gene elimination). Cells dying by DISE, in most cases, display similar morphologies and share a number of biochemical responses suggesting that DISE is not a random occurrence but has an underlying specific biological purpose which we are currently studying.

DISE was discovered by testing a large number of si- and shRNAs derived from either CD95 or CD95L. Since it was a sequence-specific OTE, it was likely that other genes also contained sequences that when expressed as shRNAs would induce DISE. We have now confirmed that shRNAs derived from a number of TS can induce a form of cell death that resembles DISE, with the same morphology, elongated cell shapes, ROS production, activation of caspase-2, inability to properly divide followed by DNA degradation and cell death. Solely by scoring similarities between the responses of cells to different toxic shRNAs did we identify 10 shRNAs (targeting 5 TS) that all killed cancer cells in which the gene was either disabled by CRISPR/Cas9 gene editing (protein knockout confirmed for two of them) or not expressed. In contrast, none of the shRNAs designed to silence the two most highly studied TS, p53 and PTEN caused significant cell death, suggesting that it is a selective group of genes that contain toxic RNAi active sequences.

Our data do not allow us to conclude that shRNAs derived from TS are particularly prone to inducing DISE. TS were merely chosen as a group of genes that were most unlikely to be critical for the survival of cancer cells. Hence, the shRNAs derived from our lethality screens designed to target TS were expected to be enriched in shRNAs that induce cell death by an OTE. Our data now suggest that certain TS-derived shRNAs can kill cancer cells through DISE and in five cases, we provide evidence to suggest that these shRNAs killed the cells in the absence of functional protein consistent with the action of the DISE mechanism. Based on these data, we propose that DISE is a general mechanism through which toxic shRNAs derived from multiple genes kill cancer cells.

Materials and methods

Reagents

Propidium iodide (#P4864) and puromycin (#P9620) were purchased from Sigma-Aldrich. Antibodies used for western blot were: anti-MAPKAPK5 (D70A10) rabbit mAb from Cell Signaling (#7419); anti-ARMC10 (#NBP1-81127) rabbit pAb from Novus Biologicals; and goat anti-rabbit IgG human adsorbed-HRP #4010-05) was from Southern Biotech.

Cell lines

The ovarian cancer cell line HeyA8, and the colon cancer cell line HCT116 were grown in RPMI 1640 medium (Mediatech Inc), supplemented with 10% heat-inactivated FBS (Sigma-Aldrich), 1% L-glutamine (Mediatech Inc), and 1% penicillin/streptomycin (Mediatech Inc). The glioblastoma cell line T98G was grown in EMEM (ATCC#30-2003), containing 10% heat-inactivated FBS, 1% L-Glutamine, and 1% penicillin/streptomycin. The chronic myelogenous leukemia cell line HAP1 (Horizon Discovery #C631), HAP1 ARMC10 k.o. (Horizon Discovery # HZGH005198c009, 2 bp deletion in exon 2, k.o. validated by Western blotting), HAP1 TGFBR2 k.o. (Horizon Discovery # HZGHC000035c015, 13 bp deletion in exon 1, and cat# HZGHC006289c002 - 7 bp deletion in exon 4, protein not detectable by Western blotting), HAP1 TMEFF1 k.o. (Horizon Discovery # HZGHC005199c011, 2 bp deletion in exon 2, k.o. not validated by Western blotting), HAP1 SOCS3 k.o. (Horizon Discovery # HZGHC005447c010, 25 bp deletion in exon 2, protein not detectable by Western blotting), and HAP1 MAPKAPK5 k.o. (Horizon Discovery # HZGHC000217c004, 4 bp deletion in exon 2, and cat# HZGHC006287c012 - 4 bp deletion in exon 8, k.o. validated by Western blotting) cell lines, were cultured in Gibco IMDM (Life Technologies #12440053), supplemented with 10% heat-inactivated FBS, 1% L-Glutamine, and 1% penicillin/ streptomycin.

Knockdown via lentiviral shRNAs

Cells were infected with the following MISSION® Lentiviral Transduction Particles (Sigma): pLKO.1-puro Control Transduction Particle coding for a nontargeting (scrambled) shRNA (#SHC002V), shRNAs against mRNA NM_000430 (Homo sapiens *PAFAH1B1*) TRCN0000050966 (#1: TGACCATTAAACTATGGGATT) and TRCN0000050964 (#7: CGTATGGGATTACAAGAACAA), shRNAs against mRNA NM_002507 (Homo sapiens *NGFR*) TRCN0000058155 (#3: CCTCCAGAACAAGACCTCATA) and TRCN0000058157 (#5: GCCTACGGCTACTACCAGGAT), shRNAs against mRNA NM_002210 (Homo sapiens *ITGAV*) TRCN0000010768 (#4: GTGAGGTCGAAACAGGATAAA) and TRCN0000010769 (#5: CGACAGGCTCACATTCTACTT), shRNAs against mRNA NM_001935 (Homo sapiens *DPP4*) TRCN0000050773 (#1: GCCCAATTTAACGACACAGAA) and TRCN0000050776 (#7: GACTGAAGTTATACTCCTTAA), shRNAs against mRNA NM_010109 (Homo sapiens *EFNA5*) TRCN0000058218 (#1: GAGACCAACAAATAGCTGTAT) and TRCN0000058220 (#3: CGCGGCACAAACACCAAGGAT), shRNAs against mRNA NM_003692 (Homo sapiens *TMEFF1*) TRCN0000073510 (#3: CATGCCAATTCAGTGCCATA) and TRCN0000073511 (#4: GCCAATTTTCAGTGCCATACAA), shRNAs against mRNA NM_001274 (Homo sapiens *CHEK1*) TRCN0000009947 (#2: GACAGAATAGAGCCAGACATA) and TRCN0000039856 (#3: GCCACATGTCCTGATCATAT), shRNAs against mRNA NM_003738 (Homo sapiens *PTCH2*) TRCN0000033327 (#9: GCTGCATTACACCAAGGAGAA) and TRCN0000033328 (#10: CGTACTCACATCCATCAACAA), shRNAs against mRNA NM_031905 (Homo sapiens *ARMC10*) TRCN0000130777 (#3: GCACATGCTTCACAGTTACAT) and TRCN0000128466 (#5: GCTTTAGTTGATCACCATGAT), shRNAs against mRNA NM_003766 (Homo sapiens *BECN1*) TRCN0000033552 (#2: CTCAAGTTCATGCTGACGAAT) and TRCN0000033553 (#8: GCTTGGGTGTCCTCACAATTT), shRNAs against mRNA NM_001356 (Homo sapiens *DDX3X*) TRCN0000000002 (#2: CGGAGTGATTACGATGGCATT) and TRCN0000000003

(#3: CGTAGAATAGTCGAACAAGAT), shRNAs against mRNA NM_006288 (Homo sapiens *THY1*) TRCN0000057023 (#1: GCCATGAGAATACCAGCAGTT) and TRCN0000057024 (#2: CGAACCAACTTCACCAGCAA), shRNAs against mRNA NM_002634 (Homo sapiens *PHB*) TRCN0000029204 (#1: CCCAGAAATCACTGTGAAATT) and TRCN0000029208 (#5: GAGTTCACAGAAGCGGTGGAA), shRNAs against mRNA NM_003955 (Homo sapiens *SOCS3*) TRCN0000057073 (#1: CCACCTGGACTCCTATGAGAA) and TRCN0000057076 (#4: CGGCTTCTACTGGAGCGCAGT), shRNAs against mRNA NM_152625 (Homo sapiens *ZNF366*) TRCN0000020134 (#1: AGGCAGTTCAAATATAGCTTT) and TRCN0000020135 (#2: GCCACAAAGATGCCCTATAA), shRNAs against mRNA NM_003668 (Homo sapiens *MAPKAPK5*) TRCN0000000681 (#1: GCGGCACTGTCACTTGTTAAA) and TRCN0000195129 (#4: CAGTATCAATTGGACTCAGAA), shRNAs against mRNA NM_003242 (Homo sapiens *TGFBR2*) TRCN0000195606 (#4: CGACATGATAGTCACTGACAA) and TRCN0000197056 (#5: GACCTCAAGAGCTCCAATATC), shRNA targeting mRNA NM_000639 (Homo sapiens *FasLG*) TRCN0000059000 (shL3: ACTGGGCTGTACTTTGTATAT), 5 shRNAs targeting mRNA NM_000314 (Homo sapiens *PTEEN*) TRCN0000355840 (#1: GGCACAAGAGGCCCTAGATTT), TRCN0000355841 (#2: ACAGTAGAGGAGCCGTCAAAT), TRCN0000355842 (#3: GACTTAGACTTGACCTATATT), TRCN0000355843 (#4: GACGAACTGGTGTAATGATAT), TRCN0000355946 (#5: ACATTATGACACCGCCAAATT), 4 shRNAs targeting mRNA NM_000546 (Homo sapiens *TP53*) TRCN0000342334 (#1: CACCATCCACTACAACACTACAT), TRCN0000342335 (#2: CGGCGCACAGAGGAAGAGAAT), TRCN0000003754 (#3: TCAGACCTATGGAAACTACTT), TRCN0000342259 (#4: GTCCAGATGAAGCTCCCAGAA).

Infection was performed according to the manufacturer's protocol. Briefly, 50,000 cells seeded the day before on a 6-well plate were infected with each lentivirus at an MOI of 3 in presence of 8µg/mL polybrene overnight. Media was changed the next day, followed by selection with 3µg/mL puromycin 24 hours later. Cells were selected for at least 48 hours, then seeded on a 96-well plate and placed in the IncuCyte (Essen Bioscience) to measure confluence or expanded for 4 days to assess cell viability with propidium iodide staining.

ROS measurement

Intracellular ROS production was measured after 8 days of infection with lentiviral shRNAs by incubating cells with 10 µM CM-H2DCFDA (C6827; Invitrogen Molecular Probes) in media at 37°C for 30 min. CM-H2DCFDA, a cell-permeable fluorogenic probe, is cleaved by intracellular esterases forming DCFH, which in presence of ROS, gets oxidized to the fluorescent compound DCF. Following incubation, cells were washed three times with PBS, and ROS was quantified by flow cytometry.

Caspase-2 activity measurement

Intracellular caspase-2 activity was detected *in situ* using FAM-VDVAD-FMK (ImmunoChemistry Technologies, LLC) according to the manufacturer's instructions. Briefly, cells were harvested 8 days after infection with lentiviral shRNAs. The pellet was resuspended in 290 µl of medium, to which 10 µl of 30x FAM-VDVAD-FMK was added. Cells were incubated at 37°C for 1 hour, washed with PBS, and resuspended in 300 µl of medium. Cells were kept on ice protected from light and immediately analyzed by flow cytometry.

Cell death assay (propidium iodide staining)

Cells infected with lentiviral shRNAs were plated in triplicates on 12 well plates after 2 days of puromycin selection, and plates were incubated at 37°C for 4 days. The total cell pellet consisting of live and dead cells was resuspended in Nicoletti buffer (0.1% sodium citrate, pH 7.4, 0.05% Triton X-100, 50 µg/ml propidium iodide). After incubating for 2-4 hours in the dark at 4°C, percent cell death was quantified by flow cytometry.

Western blot analysis

Cells were lysed using RIPA lysis buffer (1% SDS, 1% Triton X-100, 1% deoxycholic acid) and protein concentration was determined using the DC Protein Assay kit (Bio-Rad). Equal amounts of protein (30 µg) were resolved on 10% SDS-PAGE gels and transferred to nitrocellulose membrane (Amersham Protran 0.45 µm, GE Healthcare Life Science). The membranes were blocked with 5% non-fat dry milk in 0.1% Tween-20/TBS and then incubated in primary antibodies at 4°C overnight. After washing 3 times with TBST, membranes were incubated with secondary antibodies followed by washing again. Detection was performed using the ECLTM Western Blotting Detection Reagents reagent (GE Healthcare) and developed using a chemiluminescence imager, G:BOX Chemi XT4 (Syngene). Both primary and secondary antibodies were diluted in the blocking buffer (5% milk in 0.1% Tween-20/TBS) as follows: anti-ARMC10 (1:250), anti-MAPKAPK5 (1:1000) and goat anti-rabbit IgG human adsorbed-HRP (1:5000).

Statistical analyses

Growth reduction was scored as significant when cell growth was inhibited at least 50% at the half maximal growth of shScr infected cells. Percent growth reduction values were calculated using the formula: $[(y_1 - c_1) - (y_2 - c_2)] / [(y_1 - c_1)] * 100$ where y_1 is the half maximal confluency for cells infected with shScr (i.e. if the cells grew from 5% to 100% then $y_1 = [(100 + 5) / 2]$); c_1 is the starting confluency for cells infected with shScr. STATA1C software was then used to obtain the time (t_1) for y_1 and also to obtain the value of y_2 , which is the confluency of cells infected with TS shRNAs at t_1 , and c_2 is their starting confluency. Experiments were performed in triplicates and the data were expressed as mean \pm SD. Statistical analysis was performed using Student's two-tailed t-test. A value of $p < 0.05$ was considered to be significant.

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Acknowledgments

We are grateful to Denise Scholtens for biostatistics support and to Sam Bettis at the Cellular Screening Center at the University of Chicago.

Funding

This work was funded by the NIH training grant T32CA070085 (to M.P.) and R35CA197450 (to M.E.P.).

Author contributions

M.P. performed the experiments and M.E.P. designed the experiments and wrote the manuscript. Both authors read and approved the final manuscript.

FIGURE LEGENDS

Figure 1. shRNAs derived from 17 TS genes cause growth reduction in HeyA8 cells.

(A) Percent cell confluence over time of HeyA8 cells after infection with shScr, shL3 and two shRNAs for each of the 17 TS genes. The curves for cells infected with two independent shRNAs for each TS gene and their specific ID number and respective growth reduction caused by each shRNA are shown in blue and green. Percent growth reduction values (as shown in Table 1) were calculated using STATA1C software when cells infected with shScr reached half maximal confluency as indicated by the red dotted line. (B) Percent cell confluence over time of HeyA8 cells after infection with shScr, shL3, and five shRNAs targeting PTEN (left) and four shRNAs targeting p53 (right). Percent growth reduction values are shown in **Table 1**.

Figure 2. Toxic shRNAs derived from eight TS genes induce DISE-like morphological changes in HeyA8 cells.

Representative phase-contrast images showing elongated cell shapes (A); enlarged, flattened cells and presence of intracellular granules in HeyA8 cells infected with shRNAs against eight of the 17 TS and shL3 (B). shScr treated cells are shown as control.

Figure 3. Toxic shRNAs derived from seven TS genes induce death that is biochemically similar to DISE.

Quantification of ROS production by dichlorofluorescein (DCFH) fluorescence (A), caspase-2 activity (B) and quantification of cell death with PI staining (C) in HeyA8 cells 8 days after infection with shScr, shL3, and shRNAs derived from eight of the TS genes. ID numbers are shown in **Table 1**. p-values were calculated using students t-test. * $p < 0.01$, ** $p < 0.001$, *** $p < 0.0001$. For genes in black, both shRNAs were functionally active whereas genes shown in grey, only one the two shRNAs had a significant effect. Color code for the two shRNAs per gene is the same as in Fig. 1.

Figure 4. shRNAs derived from four TS genes kill cells in the absence of the transcript and/or protein produced from the targeted gene.

Percent cell confluence over time (A) and percent nuclear PI staining (B) of HAP1 parental cells, or HAP1 knock-out cells, after infection with shScr, shL3, or one of two shRNAs each targeting the respective TS. Percent growth reduction values (as shown in Table 1) were calculated using STATA1C software when cells infected with shScr reached half maximal confluency as indicated by the red dotted line. p-values were calculated using a t-test. Western blot analyses in (B) confirm the knock-out of ARMC10 and MAPKAPK5 at the protein level. Arrowhead marks likely unspecific band. * $p < 0.01$, ** $p < 0.001$, *** $p < 0.0001$. Color code for the two shRNAs per gene is the same as in Fig. 1.

Table 1: Summary of all assays leading to the identification of DISE inducing TS-derived shRNAs.

Results of systematic analyses (left to right) of two shRNAs to each of the 17 TS in various assays.

References

1. Zamore PD, Tuschl T, Sharp PA, Bartel DP. RNAi: double-stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell* 2000; 101:25-33.
2. Bernstein PL, Herrick DJ, Prokipcak RD, Ross J. Control of c-myc mRNA half-life in vitro by a protein capable of binding to a coding region stability determinant. *Genes Dev* 1992; 6:642-54.
3. Kawamata T, Seitz H, Tomari Y. Structural determinants of miRNAs for RISC loading and slicer-independent unwinding. *Nat Struct Mol Biol* 2009; 16:953-60.
4. Matranga C, Tomari Y, Shin C, Bartel DP, Zamore PD. Passenger-strand cleavage facilitates assembly of siRNA into Ago2-containing RNAi enzyme complexes. *Cell* 2005; 123:607-20.
5. Rivas FV, Tolia NH, Song JJ, Aragon JP, Liu J, Hannon GJ, et al. Purified Argonaute2 and an siRNA form recombinant human RISC. *Nat Struct Mol Biol* 2005; 12:340-9.
6. Yoda M, Kawamata T, Paroo Z, Ye X, Iwasaki S, Liu Q, et al. ATP-dependent human RISC assembly pathways. *Nat Struct Mol Biol* 2010; 17:17-23.
7. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 2004; 116:281-97.
8. Carthew RW, Sontheimer EJ. Origins and Mechanisms of miRNAs and siRNAs. *Cell* 2009; 136:642-55.
9. Hammond SM, Bernstein E, Beach D, Hannon GJ. An RNA-directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells. *Nature* 2000; 404:293-6.
10. Elbashir SM, Lendeckel W, Tuschl T. RNA interference is mediated by 21- and 22-nucleotide RNAs. *Genes Dev* 2001; 15:188-200.
11. Nykanen A, Haley B, Zamore PD. ATP requirements and small interfering RNA structure in the RNA interference pathway. *Cell* 2001; 107:309-21.
12. Jackson AL, Burchard J, Schelter J, Chau BN, Cleary M, Lim L, et al. Widespread siRNA "off-target" transcript silencing mediated by seed region sequence complementarity. *RNA* 2006; 12:1179-87.
13. Birmingham A, Anderson EM, Reynolds A, Ilesley-Tyree D, Leake D, Fedorov Y, et al. 3' UTR seed matches, but not overall identity, are associated with RNAi off-targets. *Nat Methods* 2006; 3:199-204.
14. Lin X, Ruan X, Anderson MG, McDowell JA, Kroeger PE, Fesik SW, et al. siRNA-mediated off-target gene silencing triggered by a 7 nt complementation. *Nucleic Acids Res* 2005; 33:4527-35.
15. Lim LP, Lau NC, Garrett-Engle P, Grimson A, Schelter JM, Castle J, et al. Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature* 2005; 433:769-73.
16. Gu S, Jin L, Zhang Y, Huang Y, Zhang F, Valdmann PN, et al. The loop position of shRNAs and pre-miRNAs is critical for the accuracy of dicer processing in vivo. *Cell* 2012; 151:900-11.
17. Petri S, Meister G. siRNA design principles and off-target effects. *Methods Mol Biol* 2013; 986:59-71.
18. Fellmann C, Hoffmann T, Sridhar V, Hopfgartner B, Muhar M, Roth M, et al. An optimized microRNA backbone for effective single-copy RNAi. *Cell Rep* 2013; 5:1704-13.
19. Deleavey GF, Damha MJ. Designing chemically modified oligonucleotides for targeted gene silencing. *Chem Biol* 2012; 19:937-54.
20. Chang K, Marran K, Valentine A, Hannon GJ. Creating an miR30-based shRNA vector. *Cold Spring Harbor protocols* 2013; 2013:631-5.
21. Hadji A, Ceppi P, Murmann AE, Brockway S, Pattanayak A, Bhinder B, et al. Death induced by CD95 or CD95 ligand elimination. *Cell Reports* 2014; 10:208-22.

22. Putzbach W, Gao QQ, Patel M, van Dongen S, Haluck-Kangas A, Sarshad AA, et al. Toxic si/shRNAs that kill cancer cells by targeting survival genes *BioRxiv* 2017.
23. Krammer PH. CD95's deadly mission in the immune system. *Nature* 2000; 407:789-95.
24. Murmann AE, McMahon KM, Halluck-Kangas A, Ravindran N, Patel M, Law C, et al. Induction of DISE in ovarian cancer cells in vivo. *BioRxiv* 2017.
25. Zhao M, Sun J, Zhao Z. TSGene: a web resource for tumor suppressor genes. *Nucleic Acids Res* 2013; 41:D970-6.
26. Essletzbichler P, Konopka T, Santoro F, Chen D, Gapp BV, Kralovics R, et al. Megabase-scale deletion using CRISPR/Cas9 to generate a fully haploid human cell line. *Genome Res* 2014; 24:2059-65.
27. Wang T, Birsoy K, Hughes NW, Krupczak KM, Post Y, Wei JJ, et al. Identification and characterization of essential genes in the human genome. *Science* 2015; 350:1096-101.
28. Hart T, Brown KR, Sircoulomb F, Rottapel R, Moffat J. Measuring error rates in genomic perturbation screens: gold standards for human functional genomics. *Mol Syst Biol* 2014; 10:733.
29. Morgens DW, Deans RM, Li A, Bassik MC. Systematic comparison of CRISPR/Cas9 and RNAi screens for essential genes. *Nat Biotechnol* 2016; 34:634-6.
30. Cowley GS, Weir BA, Vazquez F, Tamayo P, Scott JA, Rusin S, et al. Parallel genome-scale loss of function screens in 216 cancer cell lines for the identification of context-specific genetic dependencies. *Scientific data* 2014; 1:140035.
31. Ceppi P, Hadji A, Kohlhapp F, Pattanayak A, Hau A, Xia L, et al. CD95 and CD95L promote and protect cancer stem cells. *Nature Commun* 2014; 5:5238.

Figure 1




Figure 2




Figure 4




Table 1

TS	shRNA ID#	TRC#	% growth reduction compared to shScr			Morphological changes similar to DISE (in both HeyA8 & T98G)	Caspase 2 activity (in HeyA8 cells)	ROS (in HeyA8 cells)	% growth reduction in HAP1 cells	% growth reduction in respective k.o. cells
			HeyA8	T98G	HCT116					
PAFAH1B1	#1	TRCN0000050966	82	57	78	yes	yes	yes	0	
	#7	TRCN0000050964	87	90	88	yes	yes	yes	29	
NGFR	#3	TRCN0000058155	67	72	27					
	#5	TRCN0000058157	72	51	86					
ITGAV	#4	TRCN0000010769	80	99	91	no				
	#5	TRCN0000010768	83	80	85	no				
DPP4	#1	TRCN0000050773	88	87	78	yes	yes	yes		
	#7	TRCN0000050776	68	58	73	yes	no	no		
EFNA5	#1	TRCN0000058218	55	63						
	#3	TRCN0000058220	57	0						
TMEFF1	#3	TRCN0000073510	82	88	89	yes	yes	yes	90	89
	#4	TRCN0000073511	94	99	91	yes	yes	yes	93	85
CHEK1	#2	TRCN0000009947	69	76	88	no				
	#3	TRCN0000039856	77	86	86	yes				
PTCH2	#9	TRCN0000033327	96	96	94	yes	yes	yes	90	
	#10	TRCN0000033328	84	83	84	yes	yes	yes	0	
ARMC10	#3	TRCN0000130777	71	95	88	yes	yes	yes	60	93
	#5	TRCN0000128466	70	86	57	yes	yes	yes	73	95
BECN1	#2	TRCN0000033552	62	39						
	#8	TRCN0000033553	80	97						
DDX3X	#2	TRCN0000000002	91	80	96	yes				
	#3	TRCN0000000003	76	92	64	no				
THY1	#1	TRCN0000057023	58	60	83	no				
	#2	TRCN0000057024	86	75	69	yes				
PHB	#1	TRCN0000029204	86	89	99	no				
	#5	TRCN0000029208	95	91	98	no				
SOCS3	#1	TRCN0000057073	57	72	55	yes	yes	yes	90	93
	#4	TRCN0000057076	87	97	88	yes	yes	yes	85	78
ZNF366	#1	TRCN0000020134	92	93	46					
	#2	TRCN0000020135	68	58	56					
MAPKAPK5	#1	TRCN000000681	99	98	94	yes	yes	yes	97	98
	#4	TRCN0000195129	82	65	71	yes	yes	yes	88	78
TGFB2	#4	TRCN0000195606	82	90	83	yes	yes	yes	91	77
	#5	TRCN0000197056	57	84	62	yes	yes	yes	95	71
PTEN	#1	TRCN0000355840	0							
	#2	TRCN0000355841	0							
	#3	TRCN0000355842	0							
	#4	TRCN0000355843	0							
	#5	TRCN0000355946	29							
P53	#1	TRCN0000342334	30							
	#2	TRCN0000342335	0							
	#3	TRCN0000003754	19							
	#4	TRCN0000342259	22							


 shRNAs were effective
 shRNAs were not effective

Patel and Peter

Identification of DISE-inducing shRNAs by monitoring cellular responses

Monal Patel^a and Marcus E. Peter^{a,b}

Supplementary data



B

The TS-D genes	Full name	TSG for Cancer	Kd lethal for # of cell lines	AVG H score
PAFAH1B1/LIS1	Platelet-activating factor acetylhydrolase 1b, regulatory subunit 1 (45kDa)	HCC [1]	12	80
NGFR	Nerve growth factor receptor	HCC [2], retinoblastoma [3], prostate cancer [4], bladder cancer [5]	11	89
ITGAV/CD51	Integrin Alpha V	Ovarian cancer [6], SCC [7]	11	76
DPP4/CD26	Dipeptidyl-peptidase 4	NSCLC [8]	10	94
EFNA5	Ephrin-A5	Glioma [9]	10	80
TMEFF1	Transmembrane protein with EGF-like and two follistatin-like domains 1	Brain cancers [10]	10	84
CHEK1	Checkpoint kinase 1	BrCa [11], Multiple cancers [12]	10	55
PTCH2	Patched 2	Basal cell carcinoma [13]	9	91
ARMC10/SVH	Armadillo repeat containing 10	Osteosarcoma [14], Leukemia [15]	9	84
BECN1/ATG6	Beclin 1	Breast cancer [16, 17, 18]	9	84
FASLG	Fas ligand	Multiple cancers [19]	9	84
DDX3X/DBX	DEAD (Asp-Glu-Ala-Asp) box helicase 3, X-linked	HCC [20, 21], cutaneous squamous cell carcinoma [21]	9	80
THY1/CD90	Thy-1 cell surface antigen	Nasopharyngeal carcinoma [22], ovarian cancer [23, 24]	9	87
PHB	Prohibitin	Prostate cancer [25], liver cancer [26]	9	84
SOCS3	Suppressor of cytokine signaling 3	Breast cancer [27]	9	82
ZNF366/DC-SCRIPT	Zinc finger protein 366	Breast cancer [28]	9	80
MAPKAPK5/MK5/PRAK	Mitogen-activated protein kinase-activated protein kinase 5	Colon cancer [29], skin cancer [30]	9	68
TGFBR2	Transforming growth factor, beta receptor II (70/80kDa)	HCC [31], breast cancer [32, 33]	9	58

Figure S1: Identification of TS genes among 651 survival genes. (A) Venn diagram showing the overlap of the 651 putative survival genes we identified in 12 genome-wide shRNAs screens with a list of 637 putative tumor suppressors (<http://bioinfo.mc.vanderbilt.edu/TSGene>). (B) A list of the 17 genes that are putative tumor suppressors and were identified in our lethality screen. The genes are ranked first according to the number of lethality screens in which these genes were found to be survival genes, and second according to the average H score. Higher counts are indicated by darker colors. HCC, hepatocellular carcinoma. FASLG is also shown for comparison. [1], ¹; [2], ²; [3], ³; [4], ⁴; [5], ⁵; [6], ⁶; [7], ⁷; [8], ⁸; [9], ⁹; [10], ¹⁰; [11], ¹¹; [12], ¹²; [13], ¹³; [14], ¹⁴; [15], ¹⁵; [16], ¹⁶; [17], ¹⁷; [18], ¹⁸; [19], ¹⁹; [20], ²⁰; [21], ²¹; [22], ²²; [23], ²³; [24], ²⁴; [25], ²⁵; [26], ²⁶; [27], ²⁷; [28], ²⁸; [29], ²⁹; [30], ³⁰; [31], ³¹; [32], ³²; [33], ³³. <http://bioinfo.mc.vanderbilt.edu/TSGene> ³⁴.






Figure S2. Toxic shRNAs cause growth reduction in T98G, HCT116 and Hap1 cells.

Percent cell confluency over time of T98G (A), HCT116 (B) and Hap1 (C) cells infected with shScr, shL3, and two shRNAs derived from each TS gene. The curves for cells infected with two independent shRNA for each TS gene and their specific ID number and respective growth reduction caused by each shRNA are shown in blue and green. Percent growth reduction values (as shown in Table 1) were calculated using STATA1C software when cells infected with shScr reached half maximal confluency as indicated by the red dotted line. Names of genes for which only one of the two shRNAs reduced growth more than 50% are shown in grey.

Supplementary references

1. Xing Z, Tang X, Gao Y, Da L, Song H, Wang S, et al. The human LIS1 is downregulated in hepatocellular carcinoma and plays a tumor suppressor function. *Biochem Biophys Res Commun* 2011; 409:193-9.
2. Yuanlong H, Haifeng J, Xiaoyin Z, Jialin S, Jie L, Li Y, et al. The inhibitory effect of p75 neurotrophin receptor on growth of human hepatocellular carcinoma cells. *Cancer Lett* 2008; 268:110-9.
3. Dimaras H, Gallie BL. The p75 NTR neurotrophin receptor is a tumor suppressor in human and murine retinoblastoma development. *Int J Cancer* 2008; 122:2023-9.
4. Khwaja F, Tabassum A, Allen J, Djakiew D. The p75(NTR) tumor suppressor induces cell cycle arrest facilitating caspase mediated apoptosis in prostate tumor cells. *Biochem Biophys Res Commun* 2006; 341:1184-92.
5. Tabassum A, Khwaja F, Djakiew D. The p75(NTR) tumor suppressor induces caspase-mediated apoptosis in bladder tumor cells. *Int J Cancer* 2003; 105:47-52.
6. Kaur S, Kenny HA, Jagadeeswaran S, Zillhardt MR, Montag AG, Kistner E, et al. β 3-integrin expression on tumor cells inhibits tumor progression, reduces metastasis, and is associated with a favorable prognosis in patients with ovarian cancer. *Am J Pathol* 2009; 175:2184-96.
7. McCarty JH, Barry M, Crowley D, Bronson RT, Lacy-Hulbert A, Hynes RO. Genetic ablation of α v integrins in epithelial cells of the eyelid skin and conjunctiva leads to squamous cell carcinoma. *Am J Pathol* 2008; 172:1740-7.
8. Wesley UV, Tiwari S, Houghton AN. Role for dipeptidyl peptidase IV in tumor suppression of human non small cell lung carcinoma cells. *Int J Cancer* 2004; 109:855-66.
9. Li JJ, Liu DP, Liu GT, Xie D. EphrinA5 acts as a tumor suppressor in glioma by negative regulation of epidermal growth factor receptor. *Oncogene* 2009; 28:1759-68.
10. Gery S, Yin D, Xie D, Black KL, Koeffler HP. TMEFF1 and brain tumors. *Oncogene* 2003; 22:2723-7.
11. Sinha S, Singh RK, Bhattacharya N, Mukherjee N, Ghosh S, Alam N, et al. Frequent alterations of LOH11CR2A, PIG8 and CHEK1 genes at chromosomal 11q24.1-24.2 region in breast carcinoma: clinical and prognostic implications. *Mol Oncol* 2011; 5:454-64.
12. Lam MH, Liu Q, Elledge SJ, Rosen JM. Chk1 is haploinsufficient for multiple functions critical to tumor suppression. *Cancer Cell* 2004; 6:45-59.
13. Smyth I, Narang MA, Evans T, Heimann C, Nakamura Y, Chenevix-Trench G, et al. Isolation and characterization of human patched 2 (PTCH2), a putative tumour suppressor gene in basal cell carcinoma and medulloblastoma on chromosome 1p32. *Human Mol Gen* 1999; 8:291-7.
14. Zhou X, Yang G, Huang R, Chen X, Hu G. SVH-B interacts directly with p53 and suppresses the transcriptional activity of p53. *FEBS Lett* 2007; 581:4943-8.
15. Curtiss NP, Bonifas JM, Lauchle JO, Balkman JD, Kratz CP, Emerling BM, et al. Isolation and analysis of candidate myeloid tumor suppressor genes from a commonly deleted segment of 7q22. *Genomics* 2005; 85:600-7.
16. Furuya N, Yu J, Byfield M, Pattingre S, Levine B. The evolutionarily conserved domain of Beclin 1 is required for Vps34 binding, autophagy and tumor suppressor function. *Autophagy* 2005; 1:46-52.
17. Liang XH, Yu J, Brown K, Levine B. Beclin 1 contains a leucine-rich nuclear export signal that is required for its autophagy and tumor suppressor function. *Cancer Res* 2001; 61:3443-9.
18. Aita VM, Liang XH, Murty VV, Pincus DL, Yu W, Cayanis E, et al. Cloning and genomic organization of beclin 1, a candidate tumor suppressor gene on chromosome 17q21. *Genomics* 1999; 59:59-65.

19. Muschen M, Warskulat U, Beckmann MW. Defining CD95 as a tumor suppressor gene. *J Mol Med (Berl)* 2000; 78:312-25.
20. Shih JW, Tsai TY, Chao CH, Wu Lee YH. Candidate tumor suppressor DDX3 RNA helicase specifically represses cap-dependent translation by acting as an eIF4E inhibitory protein. *Oncogene* 2008; 27:700-14.
21. Chao CH, Chen CM, Cheng PL, Shih JW, Tsou AP, Lee YH. DDX3, a DEAD box RNA helicase with tumor growth-suppressive property and transcriptional regulation activity of the p21waf1/cip1 promoter, is a candidate tumor suppressor. *Cancer Res* 2006; 66:6579-88.
22. Lung HL, Cheung AK, Cheng Y, Kwong FM, Lo PH, Law EW, et al. Functional characterization of THY1 as a tumor suppressor gene with antiinvasive activity in nasopharyngeal carcinoma. *Int J Cancer* 2010; 127:304-12.
23. Abeysinghe HR, Pollock SJ, Guckert NL, Veyberman Y, Keng P, Halterman M, et al. The role of the THY1 gene in human ovarian cancer suppression based on transfection studies. *Cancer Genet Cytogenet* 2004; 149:1-10.
24. Abeysinghe HR, Cao Q, Xu J, Pollock S, Veyberman Y, Guckert NL, et al. THY1 expression is associated with tumor suppression of human ovarian cancer. *Cancer Genet Cytogenet* 2003; 143:125-32.
25. Dart DA, Spencer-Dene B, Gamble SC, Waxman J, Bevan CL. Manipulating prohibitin levels provides evidence for an in vivo role in androgen regulation of prostate tumours. *Endocr Relat Cancer* 2009; 16:1157-69.
26. Ko KS, Tomasi ML, Iglesias-Ara A, French BA, French SW, Ramani K, et al. Liver-specific deletion of prohibitin 1 results in spontaneous liver injury, fibrosis, and hepatocellular carcinoma in mice. *Hepatology* 2010; 52:2096-108.
27. Barclay JL, Anderson ST, Waters MJ, Curlew JD. SOCS3 as a tumor suppressor in breast cancer cells, and its regulation by PRL. *Int J Cancer* 2009; 124:1756-66.
28. Ansems M, Hontelez S, Karthaus N, Span PN, Adema GJ. Crosstalk and DC-SCRIPT: expanding nuclear receptor modulation. *Biochim Biophys Acta* 2010; 1806:193-9.
29. Kress TR, Cannell IG, Brenkman AB, Samans B, Gaestel M, Roepman P, et al. The MK5/PRAK kinase and Myc form a negative feedback loop that is disrupted during colorectal tumorigenesis. *Mol Cell* 2011; 41:445-57.
30. Sun P, Yoshizuka N, New L, Moser BA, Li Y, Liao R, et al. PRAK is essential for ras-induced senescence and tumor suppression. *Cell* 2007; 128:295-308.
31. Riehle KJ, Campbell JS, McMahan RS, Johnson MM, Beyer RP, Bammler TK, et al. Regulation of liver regeneration and hepatocarcinogenesis by suppressor of cytokine signaling 3. *J Exp Med* 2008; 205:91-103.
32. Forrester E, Chytil A, Bieri B, Aakre M, Gorska AE, Sharif-Afshar AR, et al. Effect of conditional knockout of the type II TGF-beta receptor gene in mammary epithelia on mammary gland development and polyomavirus middle T antigen induced tumor formation and metastasis. *Cancer Res* 2005; 65:2296-302.
33. Shida N, Ikeda H, Yoshimoto T, Oshima M, Taketo MM, Miyoshi I. Estrogen-induced tumorigenesis in the pituitary gland of TGF-beta(+/-) knockout mice. *Biochim Biophys Acta* 1998; 1407:79-83.
34. Zhao M, Sun J, Zhao Z. TSGene: a web resource for tumor suppressor genes. *Nucleic Acids Res* 2013; 41:D970-6.