Loss of generalist plant species and functional diversity decreases the robustness of a seed dispersal network

Vinicius A. G. Bastazini[12], Vanderlei J. Debastiani[2], Bethânia O. Azambuja[2], Paulo R. Guimarães Jr.[3], Valério D. Pillar[4]

[1]Theoretical and Experimental Ecology Station, National Center for Scientific Research – Paul Sabatier University, 2 route du CNRS, 09200, Moulis, France

[2]Graduate Program in Ecology, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, 91501-970, Brazil.

[3]Departmento de Ecologia, Instituto de Biociências, Universidade de São Paulo, Rua do Matão, Travessa 14, no. 321, 05508-900, São Paulo, SP, Brazil.

[4]Department of Ecology, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, 91501-970, Brazil.

Abstract: Understanding cascading effects of species loss has become a major challenge for ecologists. Traditionally, the robustness of ecological networks has been evaluated based on scenarios where primary extinctions occur at random or as a function of species specialization, ignoring other important biological factors. Here, we estimated the robustness of a seed dispersal network from a grassland-forest mosaic in southern Brazil simulating distinct scenarios of woody plant species extinction, including scenarios where species are eliminated based

on their evolutionary and functional distinctiveness. Our results suggest that the network is more robust when species are eliminated based on their evolutionary uniqueness, followed by random extinctions, the extinction of the most specialist species, functional distinctiveness and, at last, when the most generalist species are sequentially eliminated. Our results provide important information for grassland-forest mosaic management, as they indicate that loss of generalist species and functional diversity makes the system more likely to collapse.

Introduction

As we face the prospect of an unpreceded anthropogenic-induced mass extinction, with species being lost at rates that are two or three orders of magnitude greater than the background rates from the geological record (Pimm et al. 2014, Ceballos et al. 2015), understanding and predicting the consequences of species extinction has become an urgent task for conservation scientists and practitioners (Vieira & Almeida-Neto 2014, Ceballos et al. 2015). Most empirical studies that have examined the magnitude of biodiversity loss have usually ignored coextinction scenarios (Dunn et al. 2009, Vieira & Almeida-Neto 2014), where the loss of one or more species may unleash a cascade of secondary extinctions that may bring entire communities and ecosystems to collapse (Jackson et al. 2001, Fowler 2010, Colwell et al. 2012, Säterberg et al. 2013, Brodie et al. 2014, Donoso et al. 2017). Although the magnitude of coextinction process is still virtually unknown, it has been suggested that these cascading effects should be more pervasive in some special types of ecological interactions, such as, parasitism and mutualism (Dunn et al. 2009, Kiers et al. 2010). The

recent and rapid rise of Network Ecology – i.e., a field of research interested in understanding the function, structure, and evolution of ecological systems, using network models and analyses (Borrett et al. 2014) - has proved itself to be a powerful tool to describe the complexity of species interactions and their interdependence (Proulx et al. 2005, Pascual and Dunne 2006, Miranda et al. 2013), including their response to disturbance and cascading effects, such as coextinctions. Understanding what determines the robustness of ecological networks, i.e., the system's tolerance to secondary extinctions, has become one of the main challenges of network ecologists that have addressed this question in a variety of mutualistic systems (Solé & Montoya 2001, Dunne et al. 2002, Memmot et al. 2004, Burgos et al. 2007, Rezende et al. 2007, Mello et al. 2011a, b, Pocock et al. 2012, Eklöf et al. 2013, Vieira et al. 2013, Astegiano et al. 2015). However, despite their importance, most studies trying to estimate the robustness of ecological networks have been traditionally based on scenarios that ignore other important ecological and evolutionary factors, such as species traits, associated to the likelihood of a species becoming extinct (e.g., Dunne et al. 2002, Memmot et al. 2004, Burgos et al. 2007, Pocock et al. 2012, but see Curtsdotter et al. 2011, Srinivasan et al. 2007, Vieira et al. 2013, Astegiano et al. 2015). In these studies, primary species extinctions are assumed to occur at random or as a consequence of species specialization (Solé & Montoya 2001, Dunne et al. 2002, Memmot et al. 2004, Burgos et al. 2007, Rezende et al. 2007, Mello et al. 2011a, b, Pocock et al. 2012).

Plants-animal seed disperser interactions are an important mutualism that shapes both, ecological and evolutionary dynamics, through demographic

processes that affect the fitness, reproduction success and the distribution of partner species (Snow 1971, Morton 1973, Herrera 1985, Jordano 2000, Correa et al. 2015). That these interactions are key drivers of biodiversity patterns is supported by the fact that only in the Tropics, up to 90% of all plant species rely on animals to disperse their seeds (Jordano 2000). Given their significance to maintain biodiversity and support ecosystem functioning, plant-disperser interactions are considered a crucial component in conservation and restoration strategies (Wunderle Jr. et al. 1997, Trakhtenbrot et al. 2005).

Seed dispersal networks also play a critical role in terrestrial ecotone zones, such as in grassland-forest ecotones, as animal seed disperser may help plant species to expand their distribution across phytophysiognomy boundaries, ultimately transforming the landscape (Jensen et al. 1986, Bossuyt et al. 1999, Fragoso et al. 2003, Azambuja 2009, Carlucci et al. 2011, Müller et al. 2012, Myster 2012). Grasslands are one of the major biomes of the planet, occupying an area equivalent to 31-43% of the land surface (Coupland 1979, White et al. 2000). Characterized by their high levels of biological diversity and endemism (White et al. 2000, Bond & Parr 2010), grasslands are responsible for maintaining important ecosystem processes and economic activities (Coupland 1979, Breymer & Van Dyne 1980, White et al. 2000). Nonetheless, their conservation has been historically neglected by environmental policies (Overbeck et al. 2007, Bond & Parr 2010) and despite their large territory and economic and environmental value, grasslands are among the most altered and threatened ecosystems of the planet (White et al. 2000, Bond & Parr 2010, Henwood et al. 2010). In addition to anthropogenic threats, temperate grasslands

and savannas suffer from a natural process of forest expansion (Schwartz et al. 1996, Bowman et al. 2001, Bond & Parr 2010, Overbeck et al. 2007, Müller et al. 2012). Although forest expansion is considered a threat to grassland conservation, this process creates a unique and useful scenario for ecological studies, as grassland-forest ecotones encompass a "mosaic of life forms" that share very distinct evolutionary origins and life histories (Luza et al. 2015) that, compete for space and other resources. Seed dispersal networks, formed by woody plant and frugivores, are likely to be an important driver of spatial and temporal dynamics in grassland-forest mosaics, as seed dispersal of woody plants may accelerate the expansion of forest ecosystems (Azambuja 2009, Carlucci et al. 2011, Müller et al. 2012, Myster 2012). Thus, grassland-forest ecotones make an interesting case scenario to study simulated coextinctions, with important consequences to the development of conservation policies. For instance, if the goal of conservation efforts is to protect grasslands, these analyses can help us to identify species that should be managed in order to maintain grasslands. On the other hand, if conservation efforts are aimed to maintain forest expansion or forest species, this sort of analyses can help us to identify species that should be of higher priority in conservation.

Here, we estimated the robustness of a seed dispersal network formed by woody plants and birds from a grassland-forest mosaic from southern Brazil. We simulated five distinct scenarios of woody plant species extinction, including scenarios where species are eliminated based on their evolutionary and functional distinctiveness. Although the role of bottom-up and top-down ecosystem regulation has been a much-debated topic for decades now (see for

instance Hanley & La Pierre 2015 and references therein), both theoretical and empirical evidence suggest that bottom-up dynamics are more prominent in shaping community dynamics in multitrophic scenarios (Goudard & Loreau 2008, Scherber et al. 2010). Moreover, recent evidence suggests that plant extinctions may be more likely to trigger animal coextinctions than vice versa (Schleuning et al. 2016). Thus, we assumed a bottom-up regulation and simulated distinct scenarios of primary woody plant extinction and their effects on bird coextinctions.

Material and Methods

Study system and seed dispersal network

We applied our framework to a seed dispersal network from a forest-grassland mosaic located in southern Brazil (Azambuja 2009, Bastazini et al. 2017). The site (between 30°25'03"S 52°21'37"W and 30°25'54"S 52°22'40"W) is characterized by gently rolling terrains, with altitude ranging from 100 to 210m and by a grassland matrix, which is manly used for cattle ranching (Azambuja 2009). Large forests patches occupy hill slopes and riparian areas and smaller shrubby and forests patches are sparsely distributed across the grassy landscape (Azambuja 2009, Bastazini et al. 2017). The average annual temperature is 16.5°C and the mean annual precipitation is 1504 mm (Instituto de Pesquisas Agronômicas 1989). For further detailed characterization of the region, please refer to Azambuja (2009).

The data was collected during a bird trapping study that lasted one year,

between July 2007 and June 2008 (Azambuja 2009, Bastazini et al. 2017). Birds were captured in mist nets, which were set up for eight consecutive days every month in grasslands near to forest patches. Captured birds were placed into fabric bags for 20 minutes, so their feces could be collected (Azambuja 2009, Bastazini et al. 2017). The seeds found in fecal samples were identified to the finest taxonomic resolution possible. Based on this information, we built a qualitative interaction matrix comprising 22 woody plant species and 12 frugivorous bird species (Fig. 1). In a recent contribution, Bastazini et al. (2017) have demonstrated that trait coupling between birds and plants is an important driver of the structure of this network. We used the phylogeny generated by Bastazini et al. (2017), based on the APG III megatree (R20100701.new; Angiosperm Phylogeny Group 2009), and the traits used in their analyses, two continuous (diaspore diameter and maximum plant height) and three categorical traits (aril presence, diaspore shape and color), to estimate the phylogenetic and functional distinctiveness index explained in detail below. Further information on the phylogeny and the functional traits we used in our analyses can be found in Bastazini et al. (2017) and their Supplementary Material 4.

Fig. 1. Bi-adjacency matrix representing the seed dispersal network from a forest-grassland mosaic in southern Brazil.

Numerical analyses

We estimated the phylogenetic distinctiveness of each plant species, i.e., how isolated a species is on the phylogenetic tree, using the Fair Proportion metric, which is defined as the sum of all edge lengths between the species and the root of the phylogenetic tree, with each edge length being divided by the number of species in the cluster it subtends (Redding et al. 2008). We repeated the same procedure in order to estimate the functional distinctiveness of each species, i.e., how distinct a species is in terms of its functional traits, based on a functional dendrogram using plant traits. The functional dendrogram was built based on a Euclidean distance matrix that was calculated based on species trait values, using the Ward's sum of squares clustering procedure (Legendre & Legendre 2012).

Secondary extinctions of woody plants were simulated based on five distinct elimination schemes. The first, based on **i)** random extinctions (using 1,000 simulations), which provides a baseline scenario to compare the effects of the other scenarios (Dunne et al. 2002, Memmot et al. 2004, Astegiano et al. 2015). The other scenarios were based on species specialization in the seed dispersal network, either with **ii)** the most generalist (most connected plants) or **iii)** the most specialist (poorly connected plants) species disappearing first; **iv)** species eliminated based on their evolutionary distinctiveness; and **v)** based on species' functional distinctiveness. For each scenario, we calculated network robustness (R), defined as the area below the Attack Tolerance Curve (ATC; Albert & Barabási 2002, Burgos et al. 2007), which in our case represents the curve of the fraction of surviving bird species as a function of the eliminated plant species. R values vary between 0 and 1, and values closer to 1 indicate higher network robustness (Burgos et al. 2007).

All numerical simulations and analyses were performed in the R environment (R Core Team 2012) and the code is available on Github (https://github.com/bastazini/Networks).

Results

Species were more variable in terms of their phylogenetic distinctiveness (range: 0.19 – 0.48) than in terms of their functional distinctiveness (range: 0.19 – 0.25; Fig. 2). There was no correlation between both measures of distinctiveness (r= -0.015; P-value = 0.94), which indicates that most unique

species in the phylogenetic tree are not the most unique species in terms of their functional traits (Fig. 2).

Fig. 2. Functional and phylogenetic distinctiveness of plant species, with the phylogeny plotted alongside, in a seed dispersal network from southern Brazil.

Considering all five scenarios, our seed dispersal network is likely to be very robust to the loss of woody plant species (mean R considering all scenarios

= 0.74 ± SD = 11.85). However, our simulations show that distinct extinction scenarios have distinct effects on network robustness (Fig. 3). The loss of generalist species is more detrimental to network robustness, as the ATC of this scenario exhibits a very abrupt decline (Fig. 3). The other scenarios show a less steep response to the loss of plant species. Network robustness was higher when species were eliminated based on their evolutionary uniqueness, followed by random extinctions, the extinction of the most specialist species and functional distinctiveness (Fig. 3).

Fig. 3. Attack Tolerance Curves under different plant extinction sequences: i) random; extinctions based on species specialization either with ii) the most generalist (most connected plants) or iii) the most specialist (poorly connected

plants) species disappearing first; iv) species evolutionary distinctiveness; v) species functional distinctiveness.

Discussion

Over the past years, there has been a growing interest in trying to understand and estimate the consequences of coextinctions in ecological networks (e.g., Solé & Montoya 2001, Dunne et al. 2002, Memmot et al. 2004, Burgos et al. 2007, Rezende et al. 2007, Mello et al. 2011a, b, Pocock et al. 2012). Insofar, most empirical evaluations have been based on scenarios where primary extinctions occur as a function of species specialization or as random events (Solé & Montoya 2001, Dunne et al. 2002, Memmot et al. 2004, Burgos et al. 2007, Pocock et al 2012, Vieira et al. 2013), which are likely to be unrealistic scenarios, as it is well known that extinctions are unlikely to occur at random and that other ecological and evolutionary factors, besides specialization, such as body size and geographical range, influence the probability of a species becoming extinct (Purvis et al. 2000, Cardillo et al. 2005, 2008, Reynolds et al. 2005). Here, we provide further developments to coextinction analyses based on ATCs, combining functional and phylogenetic information, and show that the loss of generalist species and functional diversity has a large effect on network robustness, when compared to other extinction scenarios. Vieira et al. (2013) investigated the functional and phylogenetic consequences of random pollinator extinction in several pollination networks using a different metric to quantify functional and phylogenetic uniqueness. Their results suggest that there is an uncoupled response between functional and phylogenetic loss in mutualistic systems. Our

study corroborates their findings, since our extinction scenarios based on functional and phylogenetic distinctiveness showed very different responses in terms of network robustness. We expected an uncoupled functional and phylogenetic response to species extinction, as there is no phylogenetic signal in plant traits in this seed dispersal network (Bastazini et al. 2017).

Despite this growing interest in coextinction and robustness analyses, studies focusing on seed dispersal networks are still scarce. Mello et al. (2011b) analyzed the robustness of 21 seed dispersal networks by simulating random extinctions of plants and frugivores (bats and birds). For plant-bird networks, they found that networks should be very robust to the loss of plant species, with a mean value of robustness ($R = 0.75$) close to what we presented here. Future studies are necessary to demonstrate whether seed dispersal networks are as robust as demonstrated by Mello et al. (2011b) and our own analyses. But as plant-seed disperser interactions are a system of low specialization (Herrera 1995, Muller-Landau & Hardesty 2005, Donoso et al. 2017), one should expect a high degree of robustness. The sensitiveness of dispersal networks to the primary extinction of plants or seed dispersers are still unclear. Although Mello et al. (2011b) suggest that networks should be less robust when animal species are lost, Schleuning et al. (2016) argue that plant extinctions are more detrimental than animal extinctions in mutualistic networks. In any case, an obvious further step to our study is to expand our framework and investigate the effects of the loss of consumer species.

It is well recognized that generalist species, i.e., species with larger number of connections, play an important role in ecosystem functioning and

stability (Memmot et al. 2004, Richmond et al. 2005, Poisot et al. 2013, Valiente-Banuet et al. 2015). González et al. (2010) suggest that generalist species have, at least, two important roles in pollination networks; the first obvious and intuitive role is that generalist species are able to interact with more species than specialist ones and the second role is the ability to connect otherwise unconnected sub-networks. Together, these two roles implicate that generalist species are responsible for creating a more cohesive network. This cohesive pattern is likely to increment the ability of the system to respond to perturbations, and consequently increase its stability (Bascompte et al. 2003). Our seed dispersal network comprises a very heterogeneous set of plant species, in terms of phylogenetic history (15 taxonomic families), life history and functional groups, including trees (e.g., *Scutia buxifolia*, *Cupania vernalis* and *Styrax leprosus*), shrubs (*Berberis laurina*, *Chomelia obtusa* and *Daphnopsis racemosa*), vines (*Chiococca* alba) and hemiparasite species (*Tripodanthus* spp.). Among this diverse set of taxa, five of them were responsible for approximately 50% of all pairwise interactions, showing a larger number of connections: *Scutia buxifolia*, *Schinus polygamus*, *Myrsine* spp., *Daphnopsis racemosa* and *Tripodanthus* spp. Overall, these five taxa are characterized by low functional distinctiveness, with values lower than the median (except for *Tripodanthus* spp. and *Myrsine* spp.) and, except for *Schinus polygamus,* high phylogenetic distinctiveness (with values above the median). A common trait shared by these species is the size of their fruits, which is relatively small compared to the other species. As in other trophic interactions where predators have to swallow food items whole, fruit size is an important constraint in seed dispersal networks, and as a general rule, plant with large seeds and fruits tend to attract fewer animal

species than plants with small fruits (Wheelwright 1985, Jordano 2000, Dehling et al. 2016, Donoso et al. 2017). Temporal availability and co-ocurrence with its seed dispersers is also another important factor that increments the chance of a plant being consumed and dispersed by an animal (Jordano 2000). Among these five taxa, *Myrsine* spp. is the only one that produces fruits almost year-round (Azambuja 2009, Bastazini et al. 2017). Nonetheless, these well-connected species are among the taxa with larger temporal co-ocurrence with the most bird species (Bastazini et al. 2017). Thus, small fruits and phenological patterns seem to be a possible explanation to some structural patterns and the robustness of this seed dispersal network.

At last, we would like to underline the implications of our study to conservation policies, especially in southern Brazil. The conservation of grassland-forest mosaics of southern Brazil has been the focus of much debate in the recent years (Overbeck et al., 2007, 2013, Luza et al. 2014). As in other grassland ecosystem around the World, the grasslands of southern Brazil have experienced an increase in density of woody species that is drastically changing its physiognomy (Overbeck et al 2007, Müller et al. 2012). Forest expansion seems to be influenced by nucleation processes influenced by the presence of facilitating structures, such as rocky outcrops or isolated trees established in the grassland matrix, that acts as perches for vertebrate dispersers and as safe sites for woody plants (Carlucci et al., 2011, Müller et al. 2012). Most of the generalist tree species that we found in our study seem to be important species for woody plant encroachment into grasslands of southern Brazil (Carlucci et al. 2011, Müller et al. 2012). As we found the loss of generalist species and functional

uniqueness are more detrimental to the robustness of seed dispersal networks of southern Brazil, these results could help to develop sounder strategies in species management and restoration of grassland-forest mosaics in southern Brazil.

In summary, our findings provide important information for forest and grassland management in southern Brazil, as they indicate that the sequential extinction of generalist woody plant species and functional plant diversity makes the system more likely to collapse, whereas the loss of the most "unique" species, in terms of their evolutionary history, and specialist species has a smaller effect on network robustness. Moreover, despite its simplicity, our framework stresses the importance of considering distinct extinction scenarios and can help ecologists to understand and predict cascading effects in ecological systems.

Acknowledgments

We thank Andreas Kindel, André L. Luza, Paulo I. K. L. Prado, Rodrigo S. Bergamin and Sandra C. Muller for revising and providing valuable suggestions that greatly improved this manuscript. VAGB received support from CAPES (grant #1002302), the TULIP Laboratory of Excellence (ANR-10-LABX-41 and 394 ANR-11-IDEX-002-02) and from a Region Midi-Pyrenees project (CNRS 121090). VP received support from CNPq, Brazil (grant # 307689/2014-0). PRG was supported by FAPESP (grant #2009/54422-8) and CNPq.

References

Angiosperm Phylogeny Group (2009) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Botanical Journal of the Linnean Society* 161: 105–121.

Albert R, Barabási A (2002) Statistical mechanics of complex networks. *Reviews of Modern Physics* 74: 47–74.

Astegiano J, Massol F, Vidal, MM, Cheptou, PO, Guimarães Jr, PR (2015) The Robustness of Plant-Pollinator Assemblages: Linking Plant Interaction Patterns and Sensitivity to Pollinator Loss. *PloS One* 10(2): e0117243.

Azambuja, BO (2009) *Relações entre aves dispersoras de sementes e manchas florestais em matriz campestre na Serra do Sudeste, RS*. Ph. D. Thesis, Departament of Ecology, Federal University of Rio Grande do Sul, Rio Grande do Sul, Brazil.

Bascompte J, Jordano P, Melián CJ, Olesen JM (2003) The nested assembly of plant-animal mutualistic networks. *Proceedings of the National Academy of Sciences* 100: 9383–9387.

Bastazini VAG, Ferreira PMA, Azambuja BO, Casas G, Debastiani VJ, Guimaraes Jr. PR, Pillar VD (2017) Untangling the tangled bank: a novel method for partitioning the effects of phylogenies and traits on ecological networks. *Evolutionary Biology* 44:312-324

Bond WJ, Parr CL (2010) Beyond the forest edge: ecology, diversity and conservation of the grassy biomes. *Biological Conservation* 143: 2395–2404.

Borrett SR, Moody J, Edelmann A (2014) The rise of network ecology: maps of the topic diversity and scientific collaboration. *Ecological Modelling* 293: 111-127.

Bossuyt B, Hermy M, Deckers J (1999). Migration of herbaceous plant species across ancient–recent forest ecotones in central Belgium. *Journal of Ecology* 87: 629-638.

Bowman DMJS, Walsh A, Milne DJ (2001) Forest expansion and grassland contraction within a Eucalyptus savanna matrix between 1941 and 1994 at Litchfield National Park in the Australian monsoon tropics. *Global Ecology & Biogeography* 10: 535–548.

Breymeyer AI, van Dyne GM (1980) *Grasslands, systems analysis and man*. Cambridge, UK: Cambridge University Press.

Brodie JF, Aslan CE, Rogers HS, Redford KH, Maron JL, Bronstein JL, Groves, CR (2014) Secondary extinctions of biodiversity. Trends in *Ecology & Evolution* 29: 664-672.

Burgos E, Ceva H, Perazzo RPJ, Devoto M, Medan D, Zimmermann M, Delbue AM (2007) Why nestedness in mutualistic networks? *Journal of Theoretical Biology* 249: 307–313.

Cardillo M, Mace GM, Jones KE, Bielby J, Bininda-Emonds OR, Sechrest W, Orme CDL, Purvis, A (2005) Multiple causes of high extinction risk in large mammal species. *Science* 309: 1239-1241.

Cardillo M, Mace GM, Gittleman JL, Jones KE, Bielby J, Purvis A (2008) The

predictability of extinction: biological and external correlates of decline in mammals. *Proceedings of the Royal Society of London B: Biological Sciences* 275: 1441-1448.

Carlucci MB, Duarte LDS, Pillar VD (2011) Nurse rocks influence forest expansion over native grassland in southern Brazil. *Journal of Vegetation Science* 22: 111-119.

Ceballos G, Ehrlich PR, Barnosky AD, García A, Pringle RM, Palmer, TM (2015) Accelerated modern human–induced species losses: Entering the sixth mass extinction. *Science Advances* 1: e1400253.

Colwell RK, Dunn RR, Harris NC (2012) Coextinction and persistence of dependent species in a changing world. *Annual Review in Ecology, Evolution and Systematic* 43: 183–203.

Correa SB, Costa-Pereira R, Fleming T, Goulding M, Anderson JT (2015) Neotropical fish–fruit interactions: eco-evolutionary dynamics and conservation. *Biological Reviews* 90: 1263–1278.

Coupland RT (ed) (1979) *Grasslands ecosystems of the world: analysis of grasslands and their uses*. Cambridge,UK: Cambridge University Press.

Curtsdotter A, Binzer A, Brose U, de Castro F, Ebenman B, Eklöf A, Riede JO, Thierry A, Rall, BC (2011) Robustness to secondary extinctions: comparing trait-based sequential deletions in static and dynamic food webs. *Basic and Applied Ecology* 12: 571-580.

Dehling D, Jordano P, Schaefer H, Böhning-Gaese K, Schleuning M (2016) Morphology predicts species' functional roles and their degree of specialization in plant–frugivore interactions. *Proceedings of the Royal Society of London B: Biological Sciences* 283: 20152444.

Donoso I, Schleuning M, García D, Fründ J (2017) Defaunation effects on plant recruitment depend on size matching and size trade-offs in seed-dispersal networks. *Proceedings of the Royal Society of London B: Biological Sciences* 284: 20162664.

Dunn RR, Harris NC, Colwell RK, Koh LP, Sodhi NS (2009) The sixth mass coextinction: are most endangered species parasites and mutualists? *Proceedings of the Royal Society of London B: Biological Sciences* 276: 3037-3045.

Dunne JA, Williams RJ, Martinez ND (2002) Network structure and biodiversity loss in food webs: robustness increases with connectance. *Ecology Letters* 5: 558-567.

Eklöf A, Tang S, Allesina S (2013) Secondary extinctions in food webs: a Bayesian network approach. *Methods in Ecology and Evolution* 4(8): 760-770.

Fowler M (2010).Exticntion cascades and the distribution of species interactions. *Oikos* 119: 864-873.

Fragoso JMV, Silvius KM, Correa, JA (2003) Long-distance seed dispersal by tapirs increases seed survival and aggregates tropical trees. *Ecology 84:* 1998-2006.

González AMM, Dalsgaard B, Olesen JM (2010) Centrality measures and the

importance of generalist species in pollination networks. Ecological Complexity 7(1): 36-43

Goudard A, Loreau M (2008) Nontrophic interactions, biodiversity, and ecosystem functioning: an interaction web model. *The American Naturalist* 171: 91-106

Hanley TC, La Pierre KJ (eds) (2015) *Trophic Ecology: Bottom-up and Top-down Interactions across Aquatic and Terrestrial Systems*. Cambridge,UK. Cambridge University Press.

Henwood WD (2010) Toward a strategy for the conservation and protection of the world's temperate grasslands. *Great Plains Research* 20: 121–134.

Herrera, CM (1985) Determinants of plant-animal coevolution: the case of mutualistic dispersal of seeds by vertebrates. *Oikos* 44: 132-141.

Instituto de Pesquisas Agronômicas (1989.) *Atlas Agroclimático do Estado do Rio Grande do Sul*. Porto Alegre, Brazil: Pallotti.

Jackson JB, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, Bradbury RH et al. (2001) Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293: 629-637.

Jensen TS, Nielsen, O F (1986). Rodents as seed dispersers in a heath—oak wood succession. *Oecologia* 70: 214-221.

Jordano P (2000) Fruits and frugivory. In: *Seeds: The Ecology of Regeneration in Natural Plant Communities*, ed Fenner M, pp125-165. Wallingford, UK:CABI Publishing.

Kiers ET, Palmer TM, Ives AR, Bruno JF, Bronstein JL (2010) Mutualisms in a changing world: an evolutionary perspective *Ecology Letters* 13: 1459-1474.

Legendre P, Legendre LF (2012) Numerical ecology. Amsterdam, Netherlands : Elsevier.

Luza AL, Carlucci MB, Hartz SM, Duarte LD (2014) Moving from forest vs. grassland perspectives to an integrated view towards the conservation of forest–grassland mosaics. *Natureza & Conservação* 12: 166-169.

Luza AL, Gonçalves GL, Hartz, SM (2015) Phylogenetic and morphological relationships between nonvolant small mammals reveal assembly processes at different spatial scales. *Ecology and Evolution* 5: 889-902.

Mello MAR, Marquitti FMD, Guimarães Jr PR, Kalko EKV, Jordano P, de Aguiar, MM (2011a) The missing part of seed dispersal networks: structure and robustness of bat-fruit interactions. *PLoS One* 6: e17395.

Mello MAR, Marquitti FMD, Guimarães Jr PR, Kalko EKV, Jordano P, de Aguiar, MM (2011b) The modularity of seed dispersal: differences in structure and robustness between bat–and bird–fruit networks. *Oecologia* 167: 131-140.

Memmott, J., Waser, N.M. & Price, M.V. 2004. Tolerance of pollination networks to species extinctions. *Proceedings of the Royal Society B* 271: 2605–2611.

Miranda M, Parrini F, Dalerum F (2013) A categorization of recent network approaches to analyse trophic interactions. *Methods in Ecology and Evolution* 4: 897-905.

Morton ES (1973) On the evolutionary advantages and disadvantages of fruit eating in tropical birds. *The American Naturalist* 107: 8-22.

Muller-Landau HC, Hardesty BD (2005) Seed dispersal of woody plants in tropical forests: concepts, examples and future directions. In: *Biotic interactions in the tropics: Their role in the maintenance of species diversity,* eds. Burslem DF, Pinard MA, Hartley SE, pp. 267-309. Cambridge, UK: Cambridge University Press.

Müller SC, Overbeck GE, Pfadenhauer J, Pillar VD (2012) Woody species patterns at forest–grassland boundaries in southern Brazil. *Flora-Morphology, Distribution, Functional Ecology of Plants* 207: 586-598.

Myster RW (2012) Ecotones between forest and grassland. New York, USA: Springer.

Overbeck GE, Müller SC, Fidelis A, Pfadenhauer J, Pillar VD, Blanco, CC, Boldrini, II, Both R, Forneck ED (2007). Brazil's neglected biome: The South Brazilian Campos. *Perspectives in Plant Ecology, Evolution and Systematics* 9: 101–116.

Overbeck GE, Hermann JM, Andrade BO, Boldrini II, Kiehl K, Kirmer A, Koch C, et al. (2013) Restoration ecology in Brazil–time to step out of the forest. *Natureza & Conservação* 11: 92-95

Pascual M, Dunne JA (2006) Ecological networks: linking structure to dynamics in food webs. Oxford, UK: Oxford University Press.

Pimm SL, Jenkins CN, Abell R, Brooks TM, Gittleman JL, Joppa LN, Raven PH, Roberts CM, Sexton JO (2014) The biodiversity of species and their rates of

extinction, distribution, and protection. *Science* 344: 1246752.

Pocock, M.J., Evans, D.M. & Memmott, J. 2012. The robustness and restoration of a network of ecological networks. *Science* 335: 973-977.

Poisot T, Mouquet N, Gravel D (2013) Trophic complementarity drives the biodiversity–ecosystem functioning relationship in food webs. *Ecology Letters* 16: 853-861.

Proulx SR, Promislow DE, Phillips PC (2005) Network thinking in ecology and evolution. *Trends in Ecology & Evolution* 20: 345-353.

Purvis A, Gittleman JL, Cowlishaw G, Mace GM (2000) Predicting extinction risk in declining species. *Proceedings of the Royal Society of London B: Biological Sciences* 267: 1947-1952.

R Core Team (2012) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, URL http://www.R-project.org/.

Redding DW, Hartmann K, Mimoto A, Bokal D, DeVos M, Mooers AØ (2008) Evolutionarily distinctive species often capture more phylogenetic diversity than expected. *Journal of Theoretical Biology* 251(4): 606-615.

Rezende EL, Lavabre JE, Guimarães PR, Jordano P, Bascompte J (2007) *Non-random coextinctions in phylogenetically structured mutualistic networks*. Nature 448: 925-928.

Reynolds JD, Dulvy NK, Goodwin NB, Hutchings JA (2005) Biology of extinction risk in marine fishes. *Proceedings of the Royal Society of London B: Biological*

Sciences 272: 2337-2344.

Richmond CE, Breitburg DL, Rose KA (2005) The role of environmental generalist species in ecosystem function. *Ecological Modelling* 188: 279-295.

Säterberg T, Sellman S, Ebenman B (2013) High frequency of functional extinctions in ecological networks. *Nature* 499: 468-470.

Scherber C, Eisenhauer N, Weisser WW, Schmid B, Voigt W, Fischer M, Schulze E, et al. 2010. Bottom-up effects of plant diversity on multitrophic interactions in a biodiversity experiment. *Nature* 468: 553-556.

Schleuning M, Fründ J, Schweiger O, Welk E, Albrecht J, Albrecht M, Beil MN et al. (2016). Ecological networks are more sensitive to plant than to animal extinction under climate change. *Nature Communications* 7*:* 13965.

Schwartz D, Foresta H, Mariotti X, Balesdent J, Massimba JP, Girardin C (1996) Present dynamics of the savanna-forest boundary in the Congolese Mayombe: a pedological, botanical and isotopic (13C and 14C) study. *Oecologia* 106: 516-524.

Snow DW (1971) Evolutionary aspects of fruit-eating by birds. *Ibis* 113: 194-202.

Solé RV, Montoya, M (2001) Complexity and fragility in ecological networks. *Proceedings of the Royal Society of London B: Biological Sciences* 268: 2039-2045

Srinivasan UT, Dunne JA, Harte J, Martinez ND (2007) Response of complex food webs to realistic extinction sequences. Ecology 88(3): 671-682.

Trakhtenbrot A, Nathan R, Perry G, Richardson DM (2005) The importance of long-distance dispersal in biodiversity conservation. *Diversity and Distributions* 11: 173-181.

Valiente-Banuet A, Aizen MA, Alcántara JM, Arroyo J, Cocucci A, Galetti M, García MB, et al. (2015). Beyond species loss: the extinction of ecological interactions in a changing world. *Functional Ecology* 29: 299-307.

Vieira MC, Cianciaruso MV, Almeida-Neto M (2013) Plant-Pollinator coextinctions and the loss of plant functional and phylogenetic diversity. *Plos One* e 81242.

Vieira MC, Almeida-Neto M (2014) A simple stochastic model for complex coextinctions in mutualistic networks: robustness decreases with connectance. *Ecology Letters* 18: 144-152.

Wheelwright NT (1985) Fruit-size, gape width, and the diets of fruit-eating birds. *Ecology* 66: 808-818.

White R, Murray S, Rohweder M (2000) *Pilot Analysis of Global Ecosystems: Grasslands Ecosystems*. Washington, USA: World Resource Institute.

Wunderle Jr JM (1997) The role of animal seed dispersal in accelerating native forest regeneration on degraded tropical lands. *Forest Ecology and Management* 99: 223-235.