Standing Sentinel during Sleep: Continued Evaluation of Environmental Stimuli in the Absence of Consciousness

Christine Blume[1,2], Renata del Giudice[1,2], Malgorzata Wislowska[1,2],

Dominik P. J. Heib[1,2], Manuel Schabus[1,2]

[1] University of Salzburg, Department of Psychology, Laboratory for Sleep, Cognition and Consciousness Research

[2] University of Salzburg, Centre for Cognitive Neuroscience Salzburg (CCNS)

Corresponding Author:

Univ.-Prof. Dr. Manuel Schabus

University of Salzburg

Centre for Cognitive Neuroscience (CCNS)

Laboratory for "Sleep, Cognition and Consciousness Research"

Hellbrunner Str. 34

A-5020 Salzburg

Email: manuel.schabus@sbg.ac.at

Authors' email addresses:

christine.blume@sbg.ac.at

renata.delgiudice@sbg.ac.at

malgorzata.wislowska@sbg.ac.at

dominik.heib@sbg.ac.at

Funding

CB is supported by the Konrad-Adenauer-Stiftung e.V. and CB, MW and DPJH are supported by a grant from the Austrian Science Fund FWF (Y-777). CB, RdG, MW and DPJH are also supported by the Doctoral College ''Imaging the Mind'' (FWF; W1233-G17).

Abstract

While it is well-known that subject's own names (SON) or familiar voices are salient during

wakefulness, we investigate stimulus processing during sleep including N3 and REM sleep. Additionally,

we investigate how sleep EEG patterns (i.e. sleep spindles and slow oscillations [SOs]) relate to stimulus

processing. Using 256-channel EEG we studied stimulus processing by means of event-related oscillatory

responses (de-/synchronisation, ERD/ERS) and potentials (ERPs). We varied stimulus salience by

manipulating subjective (SON vs. unfamiliar name) and paralinguistic emotional relevance (familiar vs.

unfamiliar voice, FV/UFV). We show that evaluation of voice familiarity continues during all NREM

sleep stages and even REM sleep suggesting a 'sentinel processing mode' in the absence of

consciousness. Especially UFV stimuli elicit larger responses in a 1-15Hz range suggesting they are

salient. Unlike previously suggested sleep spindles and the negative slope of SOs do not uniformly inhibit

information processing but inhibition seems to be tuned to stimulus salience.

Keywords: sleep, sleep spindles, slow oscillations, high-density electroencephalography, auditory

stimulation

Introduction

Cognitive processing and task performance are well-known to vary with time of day (Dijk, Duffy, & Czeisler, 1992; Santhi et al., 2016; Wyatt, Cecco, Czeisler, & Dijk, 1999). Behaviourally, it can readily be observed with major variations in performance paralleling the sleep-wake cycle. Beyond these within-state studies that investigated wakefulness only, we lately studied cognitive processing during the fading of consciousness, which we here define as behavioural responsiveness, that is across vigilance stages from waking to light NREM sleep (Blume et al., 2016). Specifically, we compared processing of subjectively relevant vs. irrelevant stimuli (i.e. subject's own names [SONs] vs. unfamiliar names [UNs]) during wakefulness and non-rapid eye movement (NREM) sleep stages N1 and N2 during a nap and additionally varied the emotional prosody of the stimuli (i.e. stimuli spoken by an angry vs. a neutral voice [AV vs. NV]). Interestingly, we found evidence for preferential processing of salient stimuli (i.e. SONs and AV stimuli) not only during wakefulness, but also during light NREM sleep with these findings suggesting not only continued processing of external stimuli, but a 'sentinel processing mode' of the brain during states of decreased consciousness and naturally occurring unconsciousness, that is N1 and N2 sleep, respectively. Moreover, this initial preferential processing of salient stimuli seemed to be accompanied by a subsequent inhibitory sleep-protecting process during N2 sleep that was reflected by a K-complex-like response.

In the present study we sought to replicate our previous findings on the interaction between 'global brain' or vigilance states (i.e. wakefulness, N1 and N2 sleep) and stimulus characteristics and expand them to deep N3 as well as rapid-eye-movement (REM) sleep during a full night. Beyond this, we aimed at investigating the interaction between stimulus characteristics and 'local brain states' in more fine-grained analyses. In particular, we were interested in the interaction between stimuli and sleep-specific electroencephalogram (EEG) phenomena, that is sleep spindles and slow oscillations. Sleep spindles are considered the hallmark of N2 sleep albeit they also occur during sleep stage N3. They are defined as bursts of oscillatory activity in the sigma range (11-15Hz) with a characteristic waxing and waning shape

and a duration of 0.5-3s. Slow oscillations (SOs), on the other hand, are defined as large delta (0.5-3Hz)

waves with a first negative going wave that is followed by a positive going deflection. They occasionally

occur during N2 sleep already, where they may appear as isolated K-complexes, but their probability of

occurrence strongly increases with sleep depth also being a criterion for the definition of deep N3 sleep.

While it is well-established that the brain is not completely shut off from the environment during

sleep but continues to process external stimuli (e.g. Bastuji & García-Larrea, 1999; Blume et al., 2016;

Perrin, Garcia-Larrea, Mauguiere, & Bastuji, 1999; Strauss et al., 2015), studies also suggest that sleep-

specific oscillatory patterns, that is sleep spindles as well as SOs, can significantly alter stimulus

processing. Generally, it has been suggested that during spindles the thalamus acts as a sensory filter

inhibiting sensory transmission to the forebrain (Steriade, 1991). The negative or positive going slope of

SOs on the other hand has been associated with changes in the probability of synaptic release at the

cortical level, which could affect stimulus processing (Massimini & Amzica, 2001). In a combined EEG

and functional magnetic resonance imaging (fMRI) study Schabus et al. (2012) found that responses to

simple tones during NREM sleep were comparable to responses during wakefulness except for when

tones were presented during a spindle or the negative going slope of a slow oscillation thereby also

confirming previous findings (Dang-Vu et al., 2011; see De Gennaro & Ferrara, 2003 for an overview;

Massimini, Rosanova, & Mariotti, 2003). Likewise, in a study that looked at event-related potentials

(ERPs) Elton et al. (1997) suggested that sleep spindles inhibit processing of auditory stimuli and Cote,

Epps, and Campbell (2000) additionally found the effect of sleep spindles on processing to be modulated

by stimulus intensity. Specifically, they report that spindles co-occurring with more intense (i.e. louder)

stimuli seemed to inhibit processing to a greater extent than it was the case with less intense stimuli.

Regarding slow oscillatory activity on the other hand, a pioneering study by Oswald, Taylor, and

Treisman (1960) already showed that SONs evoke more K-complexes (KCs) than do unfamiliar names.

Beyond this, Massimini et al. (2003) showed that evoked somatosensory EEG potentials were strongly

modified not only by the presence but also by the phase of the slow oscillation. In summary, these

findings strongly suggest that sleep spindles and slow oscillatory activity systematically alter stimulus
processing during NREM sleep in a dynamic manner.

The aim of the present study was to investigate processing of more complex auditory stimuli (as
compared to simple tones) in relation to (i) 'global' as well as (ii) 'local' states of the brain. Complex
stimuli were first names that varied in salience on two dimensions, namely subjective relevance (SONs
vs. UNs) and familiarity or paralinguistic aspects of emotional relevance. Specifically, stimuli were
uttered by a familiar voice (FV) vs. a stranger's voice (unfamiliar voice [UFV]). Regarding the first aim,
we studied stimulus processing during all 'global brain states' across the vigilance continuum (i.e. during
wakefulness, N1, N2, N3 and REM sleep) irrespective of the 'local state'. Regarding 'local' states, we
investigated between-stimulus differences in oscillatory activity when (i) a spindle was present during
stimulus presentation, when a stimulus was presented during the (ii) positive slope of a SO, (iii) during
the negative slope and when (iv) stimulus presentation evoked a SO. Processing was studied by
comparing oscillatory brain responses evoked by stimulus presentation in each of these cases, that is
event-related synchronisation (ERS) and desynchronisation (ERD) in the delta (1-3Hz), theta (4-7Hz),
alpha (8-12Hz) and sigma (11-15 Hz) frequency range. Functionally, delta ERS has repeatedly been
linked to attentional processes and the detection of salient or motivationally relevant stimuli (for reviews
see Knyazev, 2007; Knyazev, 2012) while theta ERS has been suggested to indicate the encoding of new
information as well as working and episodic memory involvement (for a review see Klimesch, 1999;
Klimesch, Schack, & Sauseng, 2005). Alpha ERD on the other hand is thought to reflect task demands,
attentional processes and memory retrieval processes (for a review see Klimesch, 1999; Klimesch,
Doppelmayr, Russegger, Pachinger, & Schwaiger, 1998). Importantly, all these interpretations have been
established during wakefulness and it is likely that their functional roles are different during sleep. In a
previous publication, we suggested that delta and theta ERS during sleep may mirror an inhibitory sleep-
protecting response following initial processing of salient stimuli as has been suggested for sigma ERS
(Blume et al., 2016).

122   We hypothesised that oscillatory responses would mirror salience of SONs as well as FV stimuli

123 (compared to UNs and UFV) during wakefulness. Moreover, we expected responsiveness to stimuli to

124 vary with the 'global brain state', that is a decrease in responsiveness from wakefulness to N3 sleep.

125 Regarding the 'local brain state' we expected that when stimulus-presentation co-occurs with sleep

126 spindles and slow oscillations the differential brain response elicited by stimulus salience would vanish.

127 This should specifically be the case when stimulus onset coincided with the negative slope of the slow

128 oscillation or stimulus presentation largely overlapped with a sleep spindle.

130 **Results**

131 *Wakefulness*

132   Analyses in the delta band (1-3Hz) yielded a significant effect of *name* (see Fig. 1A).

133 Specifically, analyses revealed that SONs led to stronger ERS at 2Hz in a frontocentral and a parieto-

134 occipital cluster ($p = .01$ and $p = .02$, respectively) with an effect size of $d = 1.11$. This effect was also

135 visible in the ERP with SONs giving rise to a stronger P3 component than UNs ($d = .5$, see Fig 1B).

136 Analyses did not indicate a significant effect of *voice* or a *name × voice* interaction (*voice*: $ps > 0.37$;

137 *name × voice*: $ps > .35$).

NAME
WAKEFULNESS

139  **Fig. 1: Event-Related Responses during Wakefulness. (A)** Responses in the delta (1-3Hz) range. Bar plot

140 for the effect of *name* (left) and corresponding scalp plot of differences in ERS between SONs and UNs (right).

Large black dots indicate the electrodes that are part of the significant clusters at 2Hz. Error bars indicate ±1

standard error of the mean. Please note that for illustration purposes we show the effects at a representative

frequency (i.e. 2 Hz) although significant clusters may have comprised a larger frequency range (see main text). **(B)**

Event-related P3 response. Left: Grand average of the ERP elicited by SONs and UNs during wakefulness at all

electrodes that were part of the cluster (see scalp plot). The horizontal grey line represents the time window during

which the effect was significant (348 to 452ms). Right: Scalp plot of the difference in the ERPs evoked by SONs

and UNs. Large black dots indicate electrodes that were part of the marginally significant cluster and d is Cohen's d

for the significant clusters. SON = subject's own name, UN = unfamiliar name. Analyses and figures are based on data

from $n = 17$ participants.

In the theta, alpha and sigma bands (4-15Hz), the analyses yielded no significant effects (*voice*: *ps*

> 0.62; *name*: *ps* > 0.24; *name* × *voice*: *ps* > 0.4).

Sleep

Analysis of the sleep staging results revealed that the median of the total sleep time (TST) during

the experimental night was 430.5 minutes (range 300-481.5 min). Wakefulness after sleep onset (WASO)

had a median of 20 minutes (range 3.5-110 min). The total number of awakenings varied between 5 and

25 with a median of 15. SOL to N2 was characterised by a median of 20 minutes (range 10-107.5 min),

and SOL to REM had a median of 92.5 minutes (range 68.5-228 min). Regarding sleep architecture

participants had a median of 7.2% N1 sleep (range 2.7-13.7%), a median of 37% N2 sleep (range 23-

54.4%), a median of 34.2% N3 sleep (range 16.5-46.1%) and a median of 18.9% REM sleep (range 12.2-

45.1%).

"Global Brain State" Analyses

N1 sleep

During light N1 sleep, analyses yielded significant stimulus-induced differences in delta (1-3Hz)

ERS for the effect of *voice* ($p < .001$). Here, UFV stimuli elicited stronger delta ERS than FV stimuli in a

cluster that spanned large areas of the scalp with a frontal-central focus (see Fig. 2A). There were no

further significant stimulus-induced differences in the delta range (*name*: $p > .21$; *name × voice:* no

significant clusters). Analyses of responses in the theta, alpha and sigma bands (4-15Hz) also yielded a

significant effect of *voice* ($p = .003$, see Fig. 2C). Here, UFV stimuli elicited considerable ERS in the

theta through sigma frequency range in all time windows analysed (T1-T6: 4-15Hz). Analyses did not

show a significant effect of *name* ($ps > .26$), but a marginally significant *name × voice* interaction ($p =$

.055) with UFV stimuli eliciting stronger ERS irrespective of the name that was presented, i.e. SON or

UN thus confirming the main effect of *voice*. The effects of *voice* were also confirmed by ERP analyses

with a stronger positive (92-428ms) and negative component (440-996ms) for UFV as compared to FV

(see Suppl. Fig. 2A).

FAMILIARITY OF VOICE

Fig. 2: Event-related responses during NREM sleep. (A, B, C): Event-related responses in the delta range (1-3 Hz) during N1, N2 and N3. Bar plots

for the effect of *voice* (left) and corresponding scalp plots of differences in ERS between FV and UFV (right). **(D, E, F)** Event-related responses in the

theta/alpha/sigma range (4-15Hz) during N1, N2 and N3. Bar plot for the effect of *voice* during the six time windows (top) and corresponding scalp plots of

differences in ERS/ERD between FV and UFV stimuli (bottom). Large black dots indicate the electrodes that are part of the significant clusters and *d* is Cohen's *d*

for the significant clusters. Error bars indicate ±1 standard error of the mean. Please note that for illustration purposes we show the effects at representative

frequencies (i.e. 2 and 4 Hz) although significant clusters may have comprised a larger frequency range (see main text). FV = familiar voice, UFV = unfamiliar

voice. Analyses and figures are based on data from $n = 17$ participants.

N2 sleep

Analyses in the delta range yielded a significant effect of *voice* ($p < .001$) with a cluster covering

the whole scalp. Specifically, UFV stimuli elicited stronger delta ERS than FV stimuli for all frequencies

between 1 and 3Hz. The (fronto-central) topography was comparable to the N1 effect of *voice* in the delta

range (see Fig. 2A and 2B). Analyses did not yield an effect of *name* or a *name × voice* interaction (*ps >*

.18 and *no clusters*, respectively). In the theta to sigma range (4-15Hz), analyses also revealed a

significant effect of *voice* ($p = .002$). Here, again UFV stimuli elicited strong ERS between 4 and 15Hz

following about 200ms while FV stimuli elicited much less ERS (T1: 4-7 & 15Hz; T2-T6: 4-15Hz). The

topography and time course was comparable to the N1 effect (see Fig. 2D and Suppl. Fig. 1D). Besides

this, analyses showed no effect of *name* (*ps > .16*) and no *name × voice* interaction (*ps > .34*). The effects

of *voice* in oscillatory analyses were also confirmed by ERP analyses (see Suppl. Fig. 2B).

N3 sleep

During N3 sleep, analyses in the delta range (1-3Hz) revealed a significant effect of *voice* ($p <$

.001). UFV stimuli gave rise to stronger delta ERS than did FV stimuli in a cluster covering large areas of

the scalp. Again, the topography was comparable to the results obtained in N1 and N2 (see. Fig. 2A, Fig.

2B and Fig. 2C). Analyses did not reveal any stimulus-induced differences for the *name* effect (no

clusters) or the *name × voice* interaction (no clusters) in the delta range. Analyses in the theta to sigma

range (4-15Hz) revealed a significant effect of *voice* ($p = .002$; T1: 4-9 & 15Hz; T2-6: 14-15Hz). Here,

UFV stimuli elicited stronger ERS than did UFV stimuli, an effect that was especially pronounced

between about 200 and 1200ms following stimulus onset in a cluster that spanned more or less the whole

scalp. Also here, the time course and topography was comparable to the results obtained during N1 and

N2 (cf. Fig. 2D, Fig. 2E and Fig. 2F). Analyses did not yield any other significant effects (*name*: *ps > .23*;

name × voice: *ps > .31*). Analyses of ERPs confirmed the effects of *voice* (see Suppl. Fig. 2C).

 REM sleep

Analyses during REM sleep yielded a significant effect of *voice* in the delta range (1-3Hz, *p* =

.018, see Fig. 3A). As during N1-N3 FV stimuli were associated with stronger delta ERS between 1 and

2Hz than were UFV. There were no further stimulus-induced differences in delta ERS/ERD (*name*: *ps* >

.18; *name* × *voice* interaction: *ps* > .17). Analyses in the theta to sigma range (4-15Hz) yielded a

significant *voice* effect (*p* = .006, see Fig. 3B). Here, UFV stimuli elicited stronger ERS than FV stimuli

following about 200ms. The effect mainly covered the alpha through sigma range (T1: 5Hz; T2: 4-15Hz;

T3: 5-6 & 12-15Hz; T4/5: 8-15Hz; T6: 9-15Hz) and was most pronounced at central and centroparietal

electrodes. Generally, effects during REM were much less pronounced and delayed compared to the

NREM sleep stages. Analyses did not yield any further significant effects (*name*: *ps* > .37; *name* × *voice*:

ps > .32).

FAMILIARITY OF VOICE
REM SLEEP

Fig. 3: Event-related responses during REM sleep. (A) Event-related responses in the delta (1-3Hz)

range. Bar plot for the effect of *voice* (left) and corresponding scalp plot of differences in ERS between FV and UFV

(right). Large black dots indicate the electrodes that are part of the significant cluster at 2Hz. **(B)** Event-related

responses in the theta/alpha/sigma (4-15Hz) range. Bar plot for the effect of *voice* during the six time windows (top)

and corresponding scalp plots of differences in ERS/ERD between FV and UFV stimuli (bottom). Large black dots

indicate the electrodes that are part of the cluster at 12Hz and *d* is Cohen's *d* for the significant clusters. Error bars

indicate ±1 standard error of the mean. Please note that for illustration purposes we show the effects at

representative frequencies (i.e. 2 and 12 Hz) although significant clusters may have comprised a larger frequency

range (see main text). FV = familiar voice, UFV = unfamiliar voice. Analyses and figures are based on data from n =

17 participants.

"Local Brain State"- Analyses

Sleep Spindle vs. No Spindle

In both conditions, analyses of ERD/ERS revealed significant effects of *voice* in the delta range

("spindle" condition [S+]; p = .03; 1-2Hz, see Fig. 4A and "no spindle" condition [S-]: p = .028; 1-3Hz,

cf. Fig. 4B) with UFV stimuli eliciting stronger delta ERS than FV stimuli. Besides, post hoc analyses

indicated that stimulus presentation elicited more ERS in the S- compared to the S+ condition (p = .073).

There were no further effects in the delta range in either condition (S+: *name: p* = .40; *name* × *voice*: no

clusters; S-: *name*: no clusters; *name* × *voice*: *ps* > .26). In the theta to sigma range (4-15Hz) there were

also significant effects of *voice* in both the "spindle" and the "no spindle" conditions (S+: p = .033, T1:

15Hz; T2: 4 & 8-15Hz; T3/4: 4-15Hz; T5: 8-15Hz, T6: 6-15Hz, see Figs. 4C and E; S-: p = .016, T1: not

part of the cluster; T2: 4-12 Hz; T3: 4-14Hz; T4/5: 4-15Hz: T6: 4-9Hz, see Figs. 4D and F). Interestingly,

the topography and time course of the effects in the "no spindle" condition were only comparable to the

results in the "spindle" condition in the slower frequencies up to about 9Hz. While in the slower

frequencies UFV stimuli elicited stronger ERS than FV stimuli, in the faster frequencies (10-15Hz), FV

stimuli were specifically associated with a marked ERD in the "spindle" condition only (Condition

differences: p < .001, diamonds in Figs. 4E and F indicate time windows with sign. differences). Analyses

did not yield any further significant differences in the theta to sigma range (S+: *name: ps* > .12 and *name*

× *voice: ps* > .10; S-: *name*: *ps* > .22; *name* × *voice*: *ps* > .24). ERP analyses showed a significant effect of

voice that corresponded to the effects in the oscillatory analyses only in the "spindle" condition (see

Suppl. Fig. 3).

Fig. 4: Event-related responses during N2/N3 sleep depending on the presence/absence of sleep spindles. (A,

B) Event-related responses in the delta (1-3Hz) range. Bar plots for the effect of *voice* (left) and corresponding scalp

plot of differences in ERS between FV and UFV (right). **(C, D)** Event-related responses in the theta/alpha/sigma (4-

15Hz) range at 4Hz and **(E, F)** responses at 14Hz. Bar plots for the effect of *voice* during the six time windows (top)

and corresponding scalp plots of differences in ERS/ERD between FV and UFV stimuli (bottom). Diamonds in

figures E and F indicate the time windows with significant differences between S+ and S- conditions at 14 Hz. Large

black dots indicate the electrodes that are part of the clusters at 2Hz, 4Hz or 14Hz, respectively. Error bars indicate

±1 standard error of the mean and *d* is Cohen's *d* for the significant clusters. Please note that for illustration purposes we

show the effects at representative frequencies (i.e. 2, 4 and 14 Hz) although significant clusters may have comprised

a larger frequency range (see main text). FV = familiar voice, UFV = unfamiliar voice. Analyses and figures are

based on data from *n* = 10 participants.

Stimulus Presentation along Slow Oscillation Positive vs. Negative Slope

Irrespective of the slope of a SO during which a stimulus was presented, analyses yielded

significant effects of *voice* (pos. slope: $p < .001$, $d = .70$ see Fig. 5A, neg. slope: $p = .003$, $d = .67$, see

Fig. 5B) in the delta range. Specifically, like in the other conditions UFV stimuli elicited stronger ERS

than did FV stimuli between 1 and 3Hz in clusters spanning large parts of the scalp. There were no

differences in delta ERS between positive and negative SO slope ($p = .18$) and no further effects were

evident in the delta range (pos. slope: *name*: no clusters; *voice × name: $p > .1$*; neg. slope: *name*: $p = .18$;

voice × name: $p > .16$). In the theta to sigma range (4-15Hz) analyses also revealed significant effects of

voice in both conditions (pos. slope: $p < .001$, see Fig. 5C; neg. slope: $p = .008$, see Fig. 5D) with UFV

stimuli eliciting stronger ERS than FV stimuli following about 200ms in a broad frequency range

comparable to the effects in the other conditions regarding topography and time course (pos. slope: T1:

9Hz; T2-6: 4-15Hz; neg. slope: T1: not part of the cluster; T2-6: 4-15Hz). There were no differences in

the theta through sigma range between positive and negative SO slope ($ps > .72$). The effects of *voice*

were also confirmed by ERP analyses (pos. slope: see Suppl. Fig. 4B; neg. slope: see Suppl. Fig. 4C).

There were no further effects in the theta through sigma range (pos. slope: *name*: $ps > .27$ *name × voice:*

$ps > .31$; neg. slope: *name*: $ps > .40$; *name × voice*: $ps > .48$).

Fig. 5: Event-related responses during N2/N3 sleep when a stimulus was presented along the positive vs. negative slope of the SO. (A, B) Event-related responses in the delta (1-3Hz) range. Bar plots for the effect of *voice* (left) and corresponding scalp plots of differences in ERS between FV and UFV (right). (C, D) Event-related responses in the theta/alpha/sigma (4-15Hz) range. Bar plots for the effect of *voice* during the six time windows (top) and corresponding scalp plots of differences in ERS/ERD between FV and UFV stimuli (bottom). Large black dots indicate the electrodes that are part of the significant clusters and d is Cohen's d for the significant clusters. Error bars indicate ±1 standard error of the mean. Please note that for illustration purposes we show the effects at representative frequencies (i.e. 2 and 10 Hz) although significant clusters may have comprised a larger frequency range (see main text). FV = familiar voice, UFV = unfamiliar voice. Analyses and figures are based on data from n = 16 participants.

Discussion

298
In this study we show that especially processing of paralinguistic aspects of verbal stimuli such as

the familiarity of a voice is even possible during fading and in the absence of consciousness during sleep.

The findings add to existing evidence that the detection and evaluation of meaningful stimuli is still

possible in these states (e.g. Perrin et al., 1999; Portas et al., 2000). Intriguingly, we do not only find that

a differential response to familiar vs. unfamiliar voice (FV vs. UFV) stimuli persists during light NREM

sleep stages N1 and N2 thus replicating previous results (cf. Blume et al., 2016; Perrin et al., 1999), but

we extend this finding to deep N3 and intriguingly also REM sleep. Beyond this, we show that local brain

states that have been suggested to alter sensory information processing during sleep, i.e. sleep spindles

(Cote et al., 2000; Elton et al., 1997; Schabus et al., 2012) and slow oscillation down-states (Massimini et

al., 2003; Schabus et al., 2012), do at least not uniformly, that is irrespective of stimulus characteristics,

inhibit stimulus processing. Rather, their inhibitory function seems to be tuned to stimulus salience.

During wakefulness, SONs seemed to be salient when compared to UNs thus drawing more

attentional resources. This was indicated by SONs eliciting stronger delta ERS than UNs across large

areas of the scalp (Knyazev, 2007, 2012) and is well in line with the relatively larger P3 component

evident in ERP analyses as well as results from earlier studies (e.g. Berlad & Pratt, 1995; Blume et al.,

2016; Perrin et al., 1999). In an earlier study from our group, del Giudice et al. (2014) had also found

stronger alpha ERD for SONs than for UNs, which we not could not replicate here. This may ultimately

be due to methodological differences and the more conservative statistical analysis methods employed

here. Somewhat surprisingly, no differences were evident between FV and UFV stimuli when participants

were awake. However, we experience situations in which voices of varying degrees of familiarity are

present along with unfamiliar voices every day and, in comparison to the SON, familiar voices do usually

not draw attention automatically (cf. the "Cocktail Party Phenomenon"; Wood & Cowan, 1995).

Probably, this were even maladaptive as the manifold familiar voices would constantly disturb orienting

and eventually allocation of attention. Thus, the lack of a differential response evoked by stimulus
familiarity may well be considered adaptive.

During NREM sleep, that is from light N1 to deep N3 sleep, we consistently find that processing
of FV vs. UFV stimuli gives rise to a differential response in the delta to sigma frequency range, an effect
that is present in oscillatory analyses as well as ERPs. Most importantly, this provides support for the
notion that processing of auditory stimuli and especially of paralinguistic stimulus aspects such as the
familiarity of a voice is incessantly processed even in states where consciousness is absent. While this is
well in line with earlier findings during light sleep stages N1 and N2 (e.g. Blume et al., 2016; Oswald et
al., 1960; Portas et al., 2000), our results suggest that the same holds true even for deep N3 sleep. Thus,
the findings also support the notion of a 'sentinel processing mode' of the brain during sleep, which we
suggested in a previous publication (cf. Blume et al., 2016). Specifically, this mode describes the idea that
(low-level) stimulus evaluation continues even when consciousness fades during sleep and the result of
this evaluation may subsequently either trigger an inhibitory sleep-protecting response or awakening. In
detail, we here find UFV stimuli to be associated with stronger ERS in the delta range than FV stimuli
during all NREM sleep stages, an effect which was widespread across the scalp with the response being
most pronounced above frontocentral areas. Adopting the interpretation of delta oscillations during
wakefulness, the results suggest that UFV stimuli may become salient when consciousness fades
(Knyazev, 2007, 2012). In particular, the presence of unfamiliar voices could challenge the impression of
a safe environment that is necessary to 'let go of consciousness' and eventually fall and stay asleep,
rendering them salient. However, an increase in delta ERS could also be related to a sleep-specific ERP,
namely a K-complex-like response. K-complexes (KCs), whose peak frequency is in the delta range, have
been suggested to serve cortical excitation and low-level information processing as well as the subsequent
protection of sleep by neuronal silencing and they have been shown to be elicited by salient or high-
intensity stimuli (Amzica & Steriade, 1997; Bastien & Campbell, 1992; Cash et al., 2009; Laurino et al.,
2014). In line with the notion that KCs indicate ongoing cognitive processing, Vallat et al. (2017) have

recently reported a KC-like response during N2 sleep that was stronger for auditory stimuli that were

followed by an arousal or awakening. The authors concluded that this reflects stronger reactivity of the

brain to external stimuli, which in turn leads to stronger arousal. In accordance with this, ERP analyses of

our data indicated that stimulus-induced differences in the delta range indeed reflected KC-like responses

evoked by stimulus presentation with considerably larger amplitudes for UFV stimuli. In line with earlier

ideas, we suggest that this ERP reflects increased (low-level) information processing of especially salient

UFV stimuli (indexed by a larger positive wave), which is then followed by an inhibitory or sleep-

protecting 'down-state' (indexed by a larger negative wave) that is likewise scaled to stimulus salience.

Further support for this interpretation comes from analyses when stimulus presentation evoked an SO,

with evoked SOs also seeming to be sensitive to stimulus salience. Also here, UFV stimuli were

associated with stronger delta through sigma activity than FV stimuli and ERP analyses revealed that

UFV stimuli were associated with a very slight positive-going wave, which was followed by a SO down-

state that appeared much more pronounced for UFV stimuli (cf. Suppl. Fig. 4A). Besides the results

obtained in the delta range, we also find that during all NREM sleep stages UFV stimuli are associated

with stronger ERS in the theta through sigma range than FV stimuli, an effect which is most pronounced

following about 200ms after stimulus onset. Most importantly, these findings are well in line with the

delta results and they provide further convincing support for the notion that the brain is still able to

process paralinguistic stimulus aspects even when consciousness fades and is absent. On a functional

level, especially frequencies in the alpha and sigma range are thought to mirror an increase in arousal

during sleep (cf. American Academy of Sleep Medicine & Iber, 2007). This suggests that UFV stimuli

may be more arousing than FV stimuli during NREM sleep, an interpretation that, also given the observed

KC-like response, is well in line with Vallat et al.'s results. As suggested above, the presence of

unfamiliar voices may challenge the impression of an environment 'safe to sleep' and thus be arousing.

Admittedly, our findings during N2 sleep partly contrast results earlier studies, where the brain also

seemed to continue differentiating between UNs and SONs (e.g. Blume et al., 2016; Perrin et al., 1999).

The deviating findings could be due to methodological differences and/or participants sleeping during a

whole night and not just an afternoon nap (cf. Blume et al., 2016) with differences in the homeostatic and

circadian factors rendering it questionable whether a daytime nap can be considered a short night sleep

equivalent.

In summary, results obtained during wakefulness and NREM sleep suggest that familiarity of a

voice can be processed even during the fading of consciousness (N1) and in the full absence of

(behavioural) consciousness (N2 and N3). For REM sleep, a paradoxical state characterised by (i) the

return of 'altered consciousness', namely 'dreaming', (ii) enhanced brain metabolism (Maquet, 2000;

Nofzinger, Mintun, Wiseman, Kupfer, & Moore, 1997) and (iii) an increase in higher frequency EEG

power (Uchida, Maloney, & Feinberg, 1992), we also observed a relatively stronger increase in delta as

well as alpha/sigma ERS elicited by UFV compared to FV stimuli, which may indicate continued

processing and/or arousal of salient or potentially 'dangerous' UFV stimuli. This is especially interesting

because REM sleep has been suggested to reflect a 'closed loop', that is a state in which the brain is rather

occupied with intrinsic activity than processing of external stimuli (Andrillon, Poulsen, Hansen, Léger, &

Kouider, 2016; Llinás & Paré, 1991; Wehrle et al., 2007) with our results challenging this notion. At the

same time, while the oscillatory response pattern was generally similar to NREM sleep findings, REM

responses were considerably weaker and markedly delayed by approx. 400ms. This underlines the idea

that brain activity and processing of environmental stimuli during REM is qualitatively different although

not generally precluded.

Beyond investigating stimulus processing across global brain states, i.e. wakefulness and different

sleep stages, we were also interested in how stimulus presentation relates to 'local oscillatory activity',

that is sleep spindles and slow oscillations (SOs), during N2 and N3 sleep. Generally, sleep spindles

(Elton et al., 1997; Schabus et al., 2012) and the negative slope of slow oscillations (Schabus et al., 2012)

have been suggested to inhibit processing of external stimuli. Here we find that this does not seem to be

universally true but that brain responses are still tuned to stimulus salience suggesting that at least 'low-

level' processing is not precluded. More specifically, we find that when a sleep spindle overlapped with

stimulus presentation UFV stimuli still elicited responses in the delta through lower alpha (i.e. up to about

9Hz) range that were similar to those obtained when not taking 'local oscillatory activity' into account.

Intriguingly and unlike proposed earlier (Schabus et al., 2012; Steriade, 1991), this suggests that

processing of external stimuli is not or at least not uniformly inhibited by the presence of a sleep spindle,

i.e. spindles do not generally seem to act as a sensory filter at the thalamic level. Interestingly, this is well

in line with recent findings in rodents where thalamocortical sensory relay was shown to persist even

during sleep spindles (Sela, Vyazovskiy, Cirelli, Tononi, & Nir, 2016). Beyond this, above ≈9Hz the

response pattern when a spindle was present was markedly different from the general NREM (see Fig. 2)

and, most importantly, the 'no spindle' (see Fig. 4F) patterns with FV stimuli eliciting stronger ERD than

UFV stimuli in the ≈11-15Hz spindle range (see Fig. 4C). We speculate that this could reflect a relatively

stronger release of inhibition (reflected by 10-15Hz ERD) for seemingly less relevant FV stimuli by sleep

spindles. Arguably, a selective mechanism that specifically filters information that is considered

irrelevant, i.e. here FV stimuli, seems more adaptive than the uniform inhibition of all environmental

stimuli. Following the idea of a 'sentinel processing mode' of the brain during sleep, spindles just as slow

oscillations could thus reflect a sleep-protecting response that follows initial stimulus evaluation during

N2 and N3. Besides sleep spindles, previous studies suggested that also the slope of a SO during stimulus

presentation affects stimulus processing. In particular the negative slope has been found to be associated

with decreased responses in studies using somatosensory stimuli and simple tones as compared to the

positive SO slope. Surprisingly, in our study stimulus delivery during negative and positive slopes

revealed similar responses with responses in both conditions being tuned to stimulus salience.

Specifically, as during all other sleep stages UFV stimuli elicited stronger (delta to sigma) ERS than FV

stimuli. These results were supported by ERP analyses indicating that UFV stimuli induced a more

pronounced down-state that was preceded by an up-state. The findings thereby contrast earlier findings

and suggest that also the negative slope of a SO does at least not uniformly inhibit information processing

and allows continued evaluation of stimulus characteristics. Likewise, the findings also suggest that

during a positive SO slope the brain is not uniformly open to external stimulation.

In conclusion, this study shows that stimulus characteristics and especially the familiarity of a

voice continue to be evaluated during all stages of NREM sleep and thus even in the complete absence of

behavioural consciousness. Surprisingly, this is the case even during REM sleep with processing of

external seeming to be slowed and decreased though. Our findings thereby provide support for the idea of

a 'sentinel processing mode' of the brain during sleep, i.e. the continued processing of environmental

stimuli even in the absence of consciousness that may then be followed by either an inhibitory sleep-

protective response or awakening depending on the result of stimulus evaluation. Beyond this, it appears

that even 'local oscillatory activity', i.e. sleep spindles and slow oscillations are sensitive to paralinguistic

emotional stimulus characteristics. Furthermore, we provide novel evidence that even during spindles and

the negative slope of a SO the brain reacts differentially to incoming information. In a wider context, our

findings also suggest that using emotional stimuli such as familiar voices, or favourite sounds and music

may be helpful in the medical and therapeutic context when patients are in states of reduced or altered

awareness e.g. following severe brain injury.

Methods and Materials

Participants

We recruited 20 healthy individuals for the study. Three participants were excluded from the data

analysis, one dropped out after the adaptation night and two had to be excluded due to technical problems

during the acquisition. The remaining sample comprised 17 participants (three males) and had a median

age of 22.6 years (SD = 2.3 years). Prior to the study, participants gave written informed consent. Ethical

consent had been obtained from the ethics committee of the University of Salzburg and the study was in

accordance with the Declaration of Helsinki (World Medical Association (WMA), 1964). For more

details on the study sample please see supplementary material.

Experimental procedure

Participants were advised to keep a regular sleep/wake rhythm with eight hours time in bed (TIB) for

at least four days prior to their first visit at our sleep laboratory, which was verified with wrist actigraphy

(Cambridge Neurotechnology Actiwatch ©). Participants slept in the sleep laboratory of the University of

Salzburg for two nights, one adaptation night and one experimental night. The adaptation and

experimental nights were comparable except for no auditory stimulation during sleep taking place during

the adaptation night. On both nights and the following mornings participants were tested during

wakefulness resulting in four wakefulness recordings per participant. The wakefulness part comprised a

passive listening as well as an active counting condition, during which participants listened to the stimuli

presented via in-ear headphones at a volume of approximately 65 dB. For the passive condition

participants were instructed to listen attentively to the stimuli while in the active condition they were to

count the number of presentations of one specific name (i.e. the target). The passive condition always

preceded the active one. In this publication, we only present the results from the passive listening

condition, in which participants were presented with their own name (SON) as well as two unfamiliar

names (UNs) as it is the only condition that can be analysed meaningfully across 'global brain' or

vigilance stages (i.e. wakefulness, NREM and REM sleep). Moreover, each name was uttered by a

familiar and by an unfamiliar voice. The stimulus set was specific for each participant and all names of

one stimulus set were matched regarding the number of syllables and the occurrence in the general

population. During the wakefulness recording, each stimulus was presented 40 times and the interstimulus

interval (ISI) was 2000ms. Following the wakefulness recordings in the evenings, participants went to

bed for eight hours of sleep. During the experimental night, stimulation was continued and the volume

was adjusted individually so stimuli were clearly audible, but participants felt they could sleep despite the

stimulation. The auditory stimulation protocol was akin to the passive condition of the wake part,

although during the night, the stimulus onset asynchrony (SOA) was jittered between 2.8 and 7.8 s in

500ms steps. SOA was jittered specifically in the sleep protocol as this was necessary to allow for an

investigation of stimulus processing in relation to various EEG sleep phenomena (i.e. sleep spindles and

slow oscillations) independent of expectation effects. SOA was not jittered during wakefulness as this

would have rendered the tasks lengthy and probably too fatiguing. During the night each stimulus was

presented 690 times and had the same probability of occurrence as had each SOA. For more details on the

experimental procedure please see the supplementary material.

Electrophysiological data collection and reduction

For EEG acquisition we used a 256 electrode GSN HydroCel Geodesic Sensor Net (Electrical

Geodesics Inc., Eugene, Oregon, USA) and a Net Amps 400 amplifier.

Wakefulness data

EEG data were processed using the Fieldtrip toolbox (Oostenveld, Fries, Maris, & Schoffelen, 2010)

in Matlab (Mathworks, Natick, USA). First, the number of electrodes was reduced to 183 as the others

contained a lot of 'non-neural' artefacts and high-pass filtered at 0.5Hz. Subsequently, eye movement

artefacts were corrected using independent component analysis (ICA), data were segmented into 4s

epochs (symmetrically to stimulus onset) and bad intervals were removed manually during visual data

inspection. In the next step, the number of electrodes was further reduced to a final number of 173

electrodes now excluding 10 more electrodes that had initially been kept for the identification of eye and

muscular artefacts. Bad channels identified during visual data inspection were interpolated and data were

re-referenced to average reference. Subsequently, we randomly selected the same number of trials for

each stimulus to account for imbalances in the stimulus set (only one SON, but two UNs were presented).

We then applied a Morlet wavelet transformation (cycles = 3, 1-16Hz, 1Hz frequency steps) to each of the

segments, which was followed by a baseline correction (baseline interval: -600 to 0ms relative to stimulus

onset) and averaging across trials.

Sleep data

Sleep was scored semi-automatically by The Siesta Group© (Somnolyzer 24×7; cf. Anderer et al.,

2005; Anderer et al., 2004) according to standard criteria (American Academy of Sleep Medicine & Iber,

2007). Spindles were detected automatically during NREM sleep stages N2 and N3 at central leads using

the algorithm by Anderer et al. (2005). Slow oscillations (SOs) were also detected automatically on

frontal electrodes using lab-internal Matlab routines (cf. Heib et al., 2013) based on the criteria by

Massimini, Huber, Ferrarelli, Hill, and Tononi (2004). For more details on the detection of spindles and

SOs please see supplementary material. Pre-processing for the sleep data was essentially the same as for

the wakefulness data; but we refrained from an automatic eye movement correction in order to not

remove REMs. Beyond investigating processing of different stimuli across 'global brain states', that is in

each sleep stage, we also investigated stimulus processing with regard to 'local brain states', that is sleep

spindles and SOs. To this end, we compared evoked oscillatory responses elicited by different stimuli

when a spindle was present during stimulus onset (i.e. spindle offset min. 200ms after stimulus onset) or

when there was a substantial overlap between a spindle and stimulus presentation (spindle onset 0-400ms

after stimulus onset, i.e. spindle overlapping with at least half of the stimulus on average, cf. Suppl.Fig.1,

A). Moreover, we were interested in stimulus-specific differences in the evoked slow oscillatory

responses ("SO evoked"). More precisely, a SO was defined as "evoked" when the negative peak

occurred between 300 and 600ms after stimulus onset (cf. Suppl.Fig.1, B1), that is the time range when

the negative components of evoked K-complexes (i.e. N350 and N550) have been found to occur (Cote,

De Lugt, Langley, & Campbell, 1999). Beyond this, we compared stimulus processing when stimulus

onset was during the positive going slope of a SO (cf. Suppl.Fig.1, B2) to when stimulus onset coincided

with the down-state (cf. Suppl.Fig.1, B3). For more details on data collection and analysis please refer to

the supplementary material.

Event-Related Potentials

Although we focus on oscillatory activity in different frequency bands in the present manuscript,

we provide results from event-related potential (ERP) analyses in the supplementary material (and Fig.1).

Statistical Analyses

Statistical analyses were performed using the cluster-based permutation approach to correct for

multiple comparisons implemented in Fieldtrip that uses a Monte Carlo method for calculating

significance probabilities (Maris & Oostenveld, 2007). Three tests were run for the main effects of *name*

(SON vs. UNs), *voice* (FV vs. UFV) and the *name × voice* interaction with significant (or marginally

significant) interaction clusters being followed by post-hoc tests. We ran a first set of tests for the delta

range that included the dimensions electrode and frequency (1-3Hz in 1Hz frequency steps). In the delta

range, values were averaged across time (0-1000ms after stimulus onset for the WAKE condition, 0-

1200ms during SLEEP) as time resolution obtained with these low frequencies was considered

insufficient for an analysis in the time dimension. A second test was then run for the theta, alpha and

sigma ranges including the dimensions electrode, frequency (4-15Hz, 1Hz frequency steps) and time (0-

1000ms, five time windows à 200ms each in the WAKE condition, 6 time windows from 0-1200ms

during SLEEP). For the "spindle vs. no spindle" and "negative vs. positive SO slope" contrasts we

calculated averaged values for FV/UVF for each condition, which we then compared. For all permutation

tests the critical *p*-value for the *T*-statistic for dependent samples was set to 0.05 and 1000 randomisations

were used. Spatial clusters were formed only if electrodes had a minimum of two neighbouring electrodes

that were also significant. We report the Monte Carlo approximation for the estimate of *p*-values. Effects

with (one-sided) Monte Carlo $p < .05$ are denoted significant, effects with $p < .1$ are denoted marginally

significant. Critical *p*-values for post-hoc tests were adjusted for multiple comparisons using Bonferroni-

corrected *p*-values. We report Cohen's *d* (*d* = mean difference in a significant cluster/standard deviation

of the differences) as a measure of the effect size for all analyses. For more details on the statistical

analysis please see the supplementary material.

Conflict of interest

The authors declare no competing financial interests.

Acknowledgements

We thank Daniel Koerner and Adriana Michalak for their help with the data collection and pre-

processing and Vincenzo Muto for his advice on the sleep staging. We also thank Wolfgang Klimesch for

the valuable discussion about the results.

References

American Academy of Sleep Medicine, & Iber, C. (2007). *The AASM manual for the scoring of sleep and associated events: rules, terminology and technical specifications*: American Academy of Sleep Medicine.

Amzica, F., & Steriade, M. (1997). The K-complex: Its slow (<1-Hz) rhythmicity and relation to delta waves. *Neurology, 49*(4), 952-959. doi:10.1212/wnl.49.4.952

Anderer, P., Gruber, G., Parapatics, S., Woertz, M., Miazhynskaia, T., Klösch, G., Saletu, B., Zeitlhofer, J., Barbanoj, M. J., & Danker-Hopfe, H. (2005). An E-health solution for automatic sleep classification according to Rechtschaffen and Kales: validation study of the Somnolyzer 24× 7 utilizing the Siesta database. *Neuropsychobiology, 51*(3), 115-133.

Anderer, P., Saletu, B., Saletu-Zyhlarz, G. M., Gruber, G., Parapatics, S., Miazhynskaia, T., Woertz, M., Klösch, G., Zeitlhofer, J., & Dorffner, G. (2004). Recent advances in the electrophysiological evaluation of sleep. *Essentials and applications of EEG research in preclinical and clinical pharmacology. Berlin: Unipublish Verlag für Studium & Praxis OHG*, 307-339.

Andrillon, T., Poulsen, A. T., Hansen, L. K., Léger, D., & Kouider, S. (2016). Neural Markers of Responsiveness to the Environment in Human Sleep. *The Journal of Neuroscience, 36*(24), 6583-6596. doi:10.1523/jneurosci.0902-16.2016

Bastien, C., & Campbell, K. (1992). The evoked K-complex: all-or-none phenomenon? *Sleep, 15*(3), 236-245.

Bastuji, H., & García-Larrea, L. (1999). Evoked potentials as a tool for the investigation of human sleep. *Sleep Medicine Reviews, 3*(1), 23-45. doi:http://dx.doi.org/10.1016/S1087-0792(99)90012-6

Berlad, I., & Pratt, H. (1995). P300 in response to the subject's own name. *Electroencephalography and Clinical Neurophysiology, 96*(5), 472-474.

575 Blume, C., del Giudice, R., Lechinger, J., Wislowska, M., Heib, D. P. J., Hoedlmoser, K., & Schabus, M.
576   (2016). Preferential processing of emotionally and self-relevant stimuli persists in unconscious
577   N2 sleep. *Brain and Language*. doi:http://dx.doi.org/10.1016/j.bandl.2016.02.004

578 Cash, S. S., Halgren, E., Dehghani, N., Rossetti, A. O., Thesen, T., Wang, C., Devinsky, O., Kuzniecky,
579   R., Doyle, W., & Madsen, J. R. (2009). The human K-complex represents an isolated cortical
580   down-state. *Science, 324*(5930), 1084-1087.

581 Cote, K. A., De Lugt, D. R., Langley, S. D., & Campbell, K. B. (1999). Scalp topography of the auditory
582   evoked K-complex in stage 2 and slow wave sleep. *Journal of Sleep Research, 8*(4), 263-272.

583 Cote, K. A., Epps, T. M., & Campbell, K. B. (2000). The role of the spindle in human information
584   processing of high-intensity stimuli during sleep. *Journal of Sleep Research, 9*(1), 19-26.
585   doi:10.1046/j.1365-2869.2000.00188.x

586 Dang-Vu, T. T., Bonjean, M., Schabus, M., Boly, M., Darsaud, A., Desseilles, M., Degueldre, C.,
587   Balteau, E., Phillips, C., & Luxen, A. (2011). Interplay between spontaneous and induced brain
588   activity during human non-rapid eye movement sleep. *Proceedings of the National Academy of*
589   *Sciences, 108*(37), 15438-15443.

590 De Gennaro, L., & Ferrara, M. (2003). Sleep spindles: an overview. *Sleep Medicine Reviews, 7*(5), 423-
591   440.

592 del Giudice, R., Lechinger, J., Wislowska, M., Heib, D. P., Hoedlmoser, K., & Schabus, M. (2014).
593   Oscillatory brain responses to own names uttered by unfamiliar and familiar voices. *Brain*
594   *Research, 1591*, 63-73.

595 Dijk, D.-J., Duffy, J. F., & Czeisler, C. A. (1992). Circadian and sleep/wake dependent aspects of
596   subjective alertness and cognitive performance. *Journal of Sleep Research, 1*(2), 112-117.
597   doi:10.1111/j.1365-2869.1992.tb00021.x

Elton, M., Winter, O., Heslenfeld, D., Loewy, D., Campbell, K., & Kok, A. (1997). Event-related potentials to tones in the absence and presence of sleep spindles. *Journal of Sleep Research, 6*(2), 78-83.

Heib, D. P. J., Hoedlmoser, K., Anderer, P., Zeitlhofer, J., Gruber, G., Klimesch, W., & Schabus, M. (2013). Slow Oscillation Amplitudes and Up-State Lengths Relate to Memory Improvement. *PLoS ONE, 8*(12), e82049. doi:10.1371/journal.pone.0082049

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. *Brain Research Reviews, 29*(2-3), 169-195.

Klimesch, W., Doppelmayr, M., Russegger, H., Pachinger, T., & Schwaiger, J. (1998). Induced alpha band power changes in the human EEG and attention. *Neuroscience Letters, 244*(2), 73-76. doi:http://dx.doi.org/10.1016/S0304-3940(98)00122-0

Klimesch, W., Schack, B., & Sauseng, P. (2005). The functional significance of theta and upper alpha oscillations. *Experimental Psychology, 52*(2), 99-108.

Knyazev, G. G. (2007). Motivation, emotion, and their inhibitory control mirrored in brain oscillations. *Neuroscience & Biobehavioral Reviews, 31*(3), 377-395. doi:http://dx.doi.org/10.1016/j.neubiorev.2006.10.004

Knyazev, G. G. (2012). EEG delta oscillations as a correlate of basic homeostatic and motivational processes. *Neuroscience & Biobehavioral Reviews, 36*(1), 677-695. doi:http://dx.doi.org/10.1016/j.neubiorev.2011.10.002

Laurino, M., Menicucci, D., Piarulli, A., Mastorci, F., Bedini, R., Allegrini, P., & Gemignani, A. (2014). Disentangling different functional roles of evoked K-complex components: Mapping the sleeping brain while quenching sensory processing. *NeuroImage, 86*, 433-445. doi:http://dx.doi.org/10.1016/j.neuroimage.2013.10.030

Llinás, R. R., & Paré, D. (1991). Of dreaming and wakefulness. *Neuroscience, 44*(3), 521-535. doi:http://dx.doi.org/10.1016/0306-4522(91)90075-Y

Maquet, P. (2000). Functional neuroimaging of normal human sleep by positron emission tomography.
Journal of sleep research, 9(3), 207-232.

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-and MEG-data. *Journal of*
Neuroscience Methods, 164(1), 177-190.

Massimini, M., & Amzica, F. (2001). Extracellular calcium fluctuations and intracellular potentials in the
cortex during the slow sleep oscillation. *Journal of Neurophysiology, 85*(3), 1346-1350.

Massimini, M., Huber, R., Ferrarelli, F., Hill, S., & Tononi, G. (2004). The sleep slow oscillation as a
traveling wave. *The Journal of Neuroscience, 24*(31), 6862-6870.

Massimini, M., Rosanova, M., & Mariotti, M. (2003). EEG slow ($\sim$ 1 Hz) waves are associated with
nonstationarity of thalamo-cortical sensory processing in the sleeping human. *Journal of*
Neurophysiology, 89(3), 1205-1213.

Nofzinger, E. A., Mintun, M. A., Wiseman, M., Kupfer, D. J., & Moore, R. Y. (1997). Forebrain
activation in REM sleep: an FDG PET study. *Brain research, 770*(1), 192-201.

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2010). FieldTrip: open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational*
Intelligence and Neuroscience, 2011.

Oswald, I., Taylor, A. M., & Treisman, M. (1960). Discriminative responses to stimulation during human
sleep. *Brain*.

Perrin, F., Garcia-Larrea, L., Mauguiere, F., & Bastuji, H. (1999). A differential brain response to the
subject's own name persists during sleep. *Clinical Neurophysiology, 110*(12), 2153-2164.

Portas, C. M., Krakow, K., Allen, P., Josephs, O., Armony, J. L., & Frith, C. D. (2000). Auditory
processing across the sleep-wake cycle: simultaneous EEG and fMRI monitoring in humans.
Neuron, 28(3), 991-999.

Santhi, N., Lazar, A. S., McCabe, P. J., Lo, J. C., Groeger, J. A., & Dijk, D.-J. (2016). Sex differences in
the circadian regulation of sleep and waking cognition in humans. *Proceedings of the National*

Academy of Sciences of the United States of America, 113(19), E2730-E2739. doi:10.1073/pnas.1521637113

Schabus, M., Dang-Vu, T. T., Heib, D. P. J., Boly, M., Desseilles, M., Vandewalle, G., Schmidt, C., Albouy, G., Darsaud, A., & Gais, S. (2012). The fate of incoming stimuli during NREM sleep is determined by spindles and the phase of the slow oscillation. *Frontiers in Neurology, 3*.

Sela, Y., Vyazovskiy, V. V., Cirelli, C., Tononi, G., & Nir, Y. (2016). Responses in rat core auditory cortex are preserved during sleep spindle oscillations. *Sleep, 39*(5), 1069-1082.

Steriade, M. (1991). Normal and Altered States of Function. *Cerebral Cortex, 9*, 279e357.

Strauss, M., Sitt, J. D., King, J.-R., Elbaz, M., Azizi, L., Buiatti, M., Naccache, L., van Wassenhove, V., & Dehaene, S. (2015). Disruption of hierarchical predictive coding during sleep. *Proceedings of the National Academy of Sciences, 112*(11), E1353-E1362.

Uchida, S., Maloney, T., & Feinberg, I. (1992). Beta (20-28 Hz) and delta (0.3-3 Hz) EEGs oscillate reciprocally across NREM and REM sleep. *Sleep, 15*(4), 352-358.

Vallat, R., Lajnef, T., Eichenlaub, J.-B., Berthomier, C., Jerbi, K., Morlet, D., & Ruby, P. M. (2017). Increased Evoked Potentials to Arousing Auditory Stimuli during Sleep: Implication for the Understanding of Dream Recall. *Frontiers in Human Neuroscience, 11*(132). doi:10.3389/fnhum.2017.00132

Wehrle, R., Kaufmann, C., Wetter, T. C., Holsboer, F., Auer, D. P., Pollmächer, T., & Czisch, M. (2007). Functional microstates within human REM sleep: first evidence from fMRI of a thalamocortical network specific for phasic REM periods. *European Journal of Neuroscience, 25*(3), 863-871.

Wood, N., & Cowan, N. (1995). The cocktail party phenomenon revisited: how frequent are attention shifts to one's name in an irrelevant auditory channel? *Journal of Experimental Psychology: Learning, Memory, and Cognition, 21*(1), 255-260.

World Medical Association (WMA). (1964). Ethical Principles for Medical Research Involving Human Subjects. 10/13. Retrieved from http://www.wma.net/en/30publications/10policies/b3/

673   Wyatt, J. K., Cecco, A. R.-D., Czeisler, C. A., & Dijk, D.-J. (1999). Circadian temperature and melatonin

674   rhythms, sleep, and neurobehavioral function in humans living on a 20-h day. *American Journal*

675   *of Physiology - Regulatory, Integrative and Comparative Physiology, 277*(4), R1152-R1163.