Standing Sentinel during Human Sleep: Continued Evaluation of Environmental

Stimuli in the Absence of Consciousness

Christine Blume[1,2], Renata del Giudice[1,2], Malgorzata Wislowska[1,2],

Dominik P. J. Heib[1,2], Manuel Schabus[1,2]

[1] University of Salzburg, Department of Psychology, Laboratory for Sleep, Cognition and

Consciousness Research

[2] University of Salzburg, Centre for Cognitive Neuroscience Salzburg (CCNS)

Corresponding Author:

Univ.-Prof. Dr. Manuel Schabus

University of Salzburg

Centre for Cognitive Neuroscience (CCNS)

Laboratory for "Sleep, Cognition and Consciousness Research"

Hellbrunner Str. 34

A-5020 Salzburg

Email: manuel.schabus@sbg.ac.at

Authors' email addresses:

christine.blume@sbg.ac.at

renata.delgiudice@sbg.ac.at

malgorzata.wislowska@sbg.ac.at

dominik.heib@sbg.ac.at

Funding

CB is supported by the Konrad-Adenauer-Stiftung e.V. and CB, MW and DPJH are supported by a grant

from the Austrian Science Fund FWF (Y-777). CB, RdG, MW and DPJH are also supported by the

Doctoral College ''Imaging the Mind'' (FWF; W1233-G17).

Abstract

While it is a well-established finding that subject's own names (SON) or familiar voices are salient

during wakefulness, we here investigated processing of environmental stimuli during sleep including deep

N3 and REM sleep. Besides the effects of sleep depth we investigated how sleep-specific EEG patterns

(i.e. sleep spindles and slow oscillations [SOs]) relate to stimulus processing. Using 256-channel EEG we

studied processing of auditory stimuli by means of event-related oscillatory responses (de-

/synchronisation, ERD/ERS) and potentials (ERPs). We varied stimulus salience by manipulating

subjective (SON vs. unfamiliar name) and paralinguistic emotional relevance (familiar vs. unfamiliar

voice, FV/UFV). Results reveal that evaluation of voice familiarity continues during all NREM sleep

stages and even REM sleep suggesting a 'sentinel processing mode' of the human brain in the absence of

consciousness. Especially UFV stimuli elicit larger responses in a 1-15Hz range suggesting they continue

being salient. Beyond this, we show for the first time that, unlike previously suggested, sleep spindles and

the negative slope of SOs do not uniformly inhibit information processing but inhibition seems to be

tuned to stimulus salience.

Keywords: sleep, sleep spindles, slow oscillations, high-density electroencephalography, auditory

stimulation

Introduction

Cognitive processing and task performance are well-known to vary with time of day (1-3). Behaviourally, it can readily be observed with major variations in performance paralleling the sleep-wake cycle. Beyond these within-state studies that investigated wakefulness only, we lately studied cognitive processing during the fading of consciousness, which we here define as behavioural responsiveness, that is across vigilance stages from waking to light NREM sleep (4). Specifically, during a nap we compared processing of subjectively relevant vs. irrelevant stimuli (i.e. subject's own names [SONs] vs. unfamiliar names [UNs]) during wakefulness and non-rapid eye movement (NREM) sleep stages N1 and N2. Besides subjective relevance we additionally varied the emotional prosody of stimuli (i.e. stimuli spoken by an angry vs. a neutral voice [AV vs. NV]). Interestingly, we found evidence for preferential processing of salient stimuli (i.e. SONs and AV stimuli) not only during wakefulness, but also during light NREM sleep, with these findings suggesting not only continued processing of external stimuli, but a 'sentinel processing mode' of the brain during states of decreased consciousness and naturally occurring unconsciousness, that is N1 and N2 sleep, respectively. Moreover, this initial preferential processing of salient stimuli seemed to be accompanied by a subsequent inhibitory sleep-protecting process during N2 sleep that was reflected by a K-complex-like response.

In the present study we sought to replicate our previous findings on the interaction between 'longer lasting brain' or vigilance states (i.e. wakefulness, N1 and N2 sleep) and stimulus characteristics and expand them to deep N3 as well as rapid-eye-movement (REM) sleep during a full night. Beyond this, we aimed at investigating the interaction between stimulus characteristics and 'transient brain states' in more fine-grained analyses. In particular, we were interested in the interaction between stimuli and sleep-specific electroencephalogram (EEG) phenomena, that is sleep spindles and slow oscillations. Sleep spindles are considered the hallmark of N2 sleep albeit they also occur during sleep stage N3. They are defined as bursts of oscillatory activity in the sigma range (11-15Hz) with a characteristic waxing and waning shape and a duration of 0.5-3s. Slow oscillations (SOs), on the other hand, are defined as large

delta (usually < 1Hz; for criteria applied here see 5) waves with a first negative going wave that is

followed by a positive going deflection. They occasionally occur during N2 sleep already, where they

may appear as isolated K-complexes, but their probability of occurrence strongly increases with sleep

depth also being a criterion for the definition of deep N3 sleep.

While it is well-established that the brain is not completely shut off from the environment during

sleep but continues to process external stimuli (e.g. 4, 6, 7, 8), studies also suggest that sleep-specific

oscillatory patterns, that is sleep spindles as well as SOs, can significantly alter stimulus processing.

Generally, it has been suggested that during spindles the thalamus acts as a sensory filter inhibiting

sensory transmission to the forebrain (9). The negative or positive going slope of SOs on the other hand

has been associated with changes in the probability of synaptic release at the cortical level, which could

affect stimulus processing (10). In a combined EEG and functional magnetic resonance imaging (fMRI)

study Schabus, Dang-Vu (11) found that responses to simple tones during NREM sleep were comparable

to responses during wakefulness except for when tones were presented during a spindle or the negative

going slope of a slow oscillation thereby also confirming previous findings (12, see 13 for an overview,

14). Likewise, in a study that looked at event-related potentials (ERPs) Elton, Winter (15) suggested that

sleep spindles inhibit processing of auditory stimuli and Cote, Epps (16) additionally found the effect of

sleep spindles on processing to be modulated by stimulus intensity. Specifically, they report that spindles

co-occurring with more intense (i.e. louder) stimuli seemed to inhibit processing to a greater extent than it

was the case with less intense stimuli. Regarding slow oscillatory activity on the other hand, a pioneering

study by Oswald, Taylor (17) already showed that SONs evoke more K-complexes (KCs) than do

unfamiliar names. Beyond this, Massimini, Rosanova (14) showed that evoked somatosensory EEG

potentials were strongly modified not only by the presence but also by the phase of the slow oscillation.

In summary, these findings strongly suggest that sleep spindles and slow oscillatory activity

systematically alter stimulus processing during NREM sleep in a dynamic manner.

The aim of the present study was to investigate processing of more complex auditory stimuli (as

compared to simple tones) in relation to (i) 'longer lasting' as well as (ii) 'transient' states of the brain.

Complex stimuli were first names that varied in salience on two dimensions, namely subjective relevance

(SONs vs. UNs) and familiarity or paralinguistic aspects of emotional relevance. Specifically, stimuli

were uttered by a familiar voice (FV) vs. a stranger's voice (unfamiliar voice [UFV]). Regarding the first

aim, we studied stimulus processing during all 'longer lasting brain states' across the vigilance continuum

(i.e. during wakefulness, N1, N2, N3 and REM sleep) irrespective of the 'transient state'. Regarding

'transient' states, we investigated between-stimulus differences in oscillatory activity when (i) a spindle

was present during stimulus presentation, when a stimulus was presented during the (ii) positive slope of

a SO, (iii) during the negative slope and when (iv) stimulus presentation evoked a SO. Processing was

studied by comparing oscillatory brain responses evoked by stimulus presentation in each of these cases,

that is event-related synchronisation (ERS) and desynchronisation (ERD) in the delta (1-3Hz), theta (4-

7Hz), alpha (8-12Hz) and sigma (11-15 Hz) frequency range. Functionally, delta ERS has repeatedly been

linked to attentional processes and the detection of salient or motivationally relevant stimuli (for reviews

see 18, 19) while theta ERS has been suggested to indicate the encoding of new information as well as

working and episodic memory involvement (for a review see 20, 21). Alpha ERD on the other hand is

thought to reflect task demands, attentional processes and memory retrieval processes (for a review see

20, 22). Importantly, all these interpretations have been established during wakefulness and it is likely

that their functional roles are different during sleep. In a previous publication, we suggested that delta and

theta ERS during sleep may mirror an inhibitory sleep-protecting response following initial processing of

salient stimuli as has been suggested for sigma ERS (4).

We hypothesised that oscillatory responses would mirror salience of SONs as well as FV stimuli

(compared to UNs and UFV) during wakefulness. Moreover, we expected responsiveness to stimuli to

vary with the 'longer lasting brain state', that is a decrease in responsiveness from wakefulness to N3

sleep. Regarding the 'transient brain state' we expected that when stimulus-presentation co-occurs with

sleep spindles and slow oscillations the differential brain response elicited by stimulus salience would

vanish. This should specifically be the case when stimulus onset coincided with the negative slope of the

slow oscillation or stimulus presentation largely overlapped with a sleep spindle.

Results

Wakefulness

Analyses in the delta band (1-3Hz) yielded a significant effect of *name* (see Fig. 1A).

Specifically, analyses revealed that SONs led to stronger ERS at 2Hz in a frontocentral and a parieto-

occipital clusters (*Monte Carlo p* = .01 and *Monte Carlo p* = .02, respectively) with an effect size of *d* =

1.11. This effect was also visible in the ERP with SONs giving rise to a stronger P3 component than UNs

(*d* = .5, see Fig 1B). Analyses did not indicate a significant effect of *voice* or a *name × voice* interaction

(*voice*: *Monte Carlo ps* > 0.37; *name × voice*: *Monte Carlo ps* > .35).

Fig. 1: Event-Related Responses during Wakefulness. (A) Responses in the delta (1-3Hz) range. Bar plot

for the effect of *name* (left) and corresponding scalp plot of differences in ERS between SONs and UNs (right).

Large black dots indicate the electrodes that are part of the significant clusters at 2Hz. Error bars indicate ±1

standard error of the mean. Please note that for illustration purposes we show the effects at a representative

frequency (i.e. 2 Hz) although significant clusters may have comprised a larger frequency range (see main text). **(B)**

Event-related P3 response. Left: Grand average of the ERP elicited by SONs and UNs during wakefulness at all

electrodes that were part of the cluster (see scalp plot). The horizontal grey line represents the time window during which the effect was significant (348 to 452ms). Right: Scalp plot of the difference in the ERPs evoked by SONs and UNs. Large black dots indicate electrodes that were part of the marginally significant cluster and *d* is Cohen's *d* for the significant clusters. SON = subject's own name, UN = unfamiliar name. Analyses and figures are based on data from *n* = 17 participants.

In the theta, alpha and sigma bands (4-15Hz), the analyses yielded no significant effects (*voice*: *Monte Carlo ps > 0.62*; *name*: *Monte Carlo ps > 0.24*; *name × voice*: *Monte Carlo ps > 0.4*).

Sleep

Analysis of the sleep staging results revealed that the median of the total sleep time (TST) during the experimental night was 430.5 minutes (range 300-481.5 min). Wakefulness after sleep onset (WASO) had a median of 20 minutes (range 3.5-110 min). The total number of awakenings varied between 5 and 25 with a median of 15. SOL to N2 was characterised by a median of 20 minutes (range 10-107.5 min), and SOL to REM had a median of 92.5 minutes (range 68.5-228 min). Regarding sleep architecture participants had a median of 7.2% N1 sleep (range 2.7-13.7%), a median of 37% N2 sleep (range 23-54.4%), a median of 34.2% N3 sleep (range 16.5-46.1%) and a median of 18.9% REM sleep (range 12.2-45.1%).

"Longer lasting Brain State" Analyses

N1 sleep

During light N1 sleep, analyses yielded significant stimulus-induced differences in delta (1-3Hz) ERS for the effect of *voice* (*Monte Carlo p < .001*). Here, UFV stimuli elicited stronger delta ERS than FV stimuli in a cluster that spanned large areas of the scalp with a frontal-central focus (see Fig. 2A). There were no further significant stimulus-induced differences in the delta range (*name*: *p > .21*; *name ×*

voice: no significant clusters). Analyses of responses in the theta, alpha and sigma bands (4-15Hz) also

yielded a significant effect of *voice* (*Monte Carlo p* = .003, see Fig. 2C). Here, UFV stimuli elicited

considerable ERS in the theta through sigma frequency range in all time windows analysed (T1-T6: 4-

15Hz). Analyses did not show a significant effect of *name* (*Monte Carlo ps* > .26), but a marginally

significant *name × voice* interaction (*Monte Carlo p* = .055) with UFV stimuli eliciting stronger ERS

irrespective of the name that was presented, i.e. SON or UN, thus confirming the main effect of *voice*.

The effects of *voice* were also confirmed by ERP analyses with a stronger positive (92-428ms) and

negative component (440-996ms) for UFV as compared to FV (see Suppl. Fig. 2A).

FAMILIARITY OF VOICE

Fig. 2: Event-related responses during NREM sleep. (A, B, C): Event-related responses in the delta range (1-3 Hz) during N1, N2 and N3. Bar plots

for the effect of *voice* (left) and corresponding scalp plots of differences in ERS between FV and UFV (right). **(D, E, F)** Event-related responses in the

theta/alpha/sigma range (4-15Hz) during N1, N2 and N3. Bar plot for the effect of *voice* during the six time windows (top) and corresponding scalp plots of

differences in ERS/ERD between FV and UFV stimuli (bottom). Large black dots indicate the electrodes that are part of the significant clusters and *d* is Cohen's *d*

for the significant clusters. Error bars indicate ±1 standard error of the mean. Please note that for illustration purposes we show the effects at representative

frequencies (i.e. 2 and 4 Hz) although significant clusters may have comprised a larger frequency range (see main text). FV = familiar voice, UFV = unfamiliar

voice. Analyses and figures are based on data from $n = 17$ participants.

N2 sleep

Analyses in the delta range yielded a significant effect of *voice* ($p < .001$) with a cluster covering

the whole scalp. Specifically, UFV stimuli elicited stronger delta ERS than FV stimuli for all frequencies

between 1 and 3Hz. The (fronto-central) topography was comparable to the N1 effect of *voice* in the delta

range (see Fig. 2A and 2B). Analyses did not yield an effect of *name* or a *name* × *voice* interaction

(*Monte Carlo ps* > .18 and *no clusters*, respectively). In the theta to sigma range (4-15Hz), analyses also

revealed a significant effect of *voice* (*Monte Carlo p* = .002). Here, again UFV stimuli elicited strong

ERS between 4 and 15Hz following about 200ms while FV stimuli elicited much less ERS (T1: 4-7 &

15Hz; T2-T6: 4-15Hz). The topography and time course was comparable to the N1 effect (see Fig. 2D

and Suppl. Fig. 1D). Besides this, analyses showed no effect of *name* (*Monte Carlo ps* > .16) and no

name × *voice* interaction (*Monte Carlo ps* > .34). The effects of *voice* in oscillatory analyses were also

confirmed by ERP analyses (see Suppl. Fig. 2B).

N3 sleep

During N3 sleep, analyses in the delta range (1-3Hz) revealed a significant effect of *voice* (*Monte*

Carlo p < .001). UFV stimuli gave rise to stronger delta ERS than did FV stimuli in a cluster covering

large areas of the scalp. Again, the topography was comparable to the results obtained in N1 and N2 (see.

Fig. 2A, Fig. 2B and Fig. 2C). Analyses did not reveal any stimulus-induced differences for the *name*

effect (no clusters) or the *name* × *voice* interaction (no clusters) in the delta range. Analyses in the theta to

sigma range (4-15Hz) revealed a significant effect of *voice* (*Monte Carlo p* = .002; T1: 4-9 & 15Hz; T2-

6: 14-15Hz). Here, UFV stimuli elicited stronger ERS than did FV stimuli, an effect that was especially

pronounced between about 200 and 1200ms following stimulus onset in a cluster that spanned more or

less the whole scalp. Also here, the time course and topography was comparable to the results obtained

during N1 and N2 (cf. Fig. 2D, Fig. 2E and Fig. 2F). Analyses did not yield any other significant effects

(*name*: *Monte Carlo ps* > .23; *name × voice*: *Monte Carlo ps* > .31). Analyses of ERPs confirmed the

effects of *voice* (see Suppl. Fig. 2C).

REM sleep

Analyses during REM sleep yielded a significant effect of *voice* in the delta range (1-3Hz, *Monte*

Carlo p = .018, see Fig. 3A). As during N1-N3, FV stimuli were associated with stronger delta ERS

between 1 and 2Hz than were UFV. There were no further stimulus-induced differences in delta

ERS/ERD (*name*: *Monte Carlo ps* > .18; *name × voice* interaction: *Monte Carlo ps* > .17). Analyses in the

theta to sigma range (4-15Hz) yielded a significant *voice* effect (*Monte Carlo p* = .006, see Fig. 3B).

Here, UFV stimuli elicited stronger ERS than FV stimuli following about 200ms. The effect mainly

covered the alpha through sigma range (T1: 5Hz; T2: 4-15Hz; T3: 5-6 & 12-15Hz; T4/5: 8-15Hz; T6: 9-

15Hz) and was most pronounced at the central and centroparietal electrodes. Generally, effects during

REM were much less pronounced and delayed compared to the NREM sleep stages. Analyses did not

yield any further significant effects (*name*: *Monte Carlo ps* > .37; *name × voice: Monte Carlo ps* > .32).

Fig. 3: Event-related responses during REM sleep. (A) Event-related responses in the delta (1-3Hz)

range. Bar plot for the effect of *voice* (left) and corresponding scalp plot of differences in ERS between FV and UFV

(right). Large black dots indicate the electrodes that are part of the significant cluster at 2Hz. **(B)** Event-related

responses in the theta/alpha/sigma (4-15Hz) range. Bar plot for the effect of *voice* during the six time windows (top)

and corresponding scalp plots of differences in ERS/ERD between FV and UFV stimuli (bottom). Large black dots

indicate the electrodes that are part of the cluster at 12Hz and *d* is Cohen's *d* for the significant clusters. Error bars

indicate ±1 standard error of the mean. Please note that for illustration purposes we show the effects at

representative frequencies (i.e. 2 and 12 Hz) although significant clusters may have comprised a larger frequency

range (see main text). FV = familiar voice, UFV = unfamiliar voice. Analyses and figures are based on data from *n* =

17 participants.

"Transient Brain State"- Analyses

Sleep Spindle vs. No Spindle

In both conditions, analyses of ERD/ERS revealed significant effects of *voice* in the delta range

("spindle" condition [S+]: *Monte Carlo p* = .03; 1-2Hz, see Fig. 4A and "no spindle" condition [S-]:

Monte Carlo p = .028; 1-3Hz, cf. Fig. 4B) with UFV stimuli eliciting stronger delta ERS than FV stimuli.

Besides, post hoc analyses indicated that stimulus presentation elicited more ERS in the S- compared to

the S+ condition (*Monte Carlo p* = .073). There were no further effects in the delta range in either

condition (S+: *name: Monte Carlo p* = .40; *name × voice*: no clusters; S-: *name*: no clusters; *name ×*

voice: *Monte Carlo ps* > .26). In the theta to sigma range (4-15Hz) there were also significant effects of

voice in both the "spindle" and the "no spindle" conditions (S+: *Monte Carlo p* = .033, T1: 15Hz; T2: 4

& 8-15Hz; T3/4: 4-15Hz; T5: 8-15Hz, T6: 6-15Hz, see Figs. 4C and E; S-: *Monte Carlo p* = .016, T1: not

part of the cluster; T2: 4-12 Hz; T3: 4-14Hz; T4/5: 4-15Hz: T6: 4-9Hz, see Figs. 4D and F). Interestingly,

the topography and time course of the effects in the "no spindle" condition were only comparable to the

results in the "spindle" condition in the slower frequencies up to about 9Hz. While in the slower

frequencies UFV stimuli elicited stronger ERS than FV stimuli, in the faster frequencies (10-15Hz), FV

stimuli were specifically associated with a marked ERD in the "spindle" condition only (Condition

differences: *Monte Carlo p* < .001, diamonds in Figs. 4E and F indicate time windows with sign.

differences). Analyses did not yield any further significant differences in the theta to sigma range (S+: *name: Monte Carlo ps > .12* and *name × voice: Monte Carlo ps > .10*; S-: *name*: *Monte Carlo ps > .22*; *name × voice*: *Monte Carlo ps > .24*). ERP analyses showed a significant effect of *voice* that corresponded to the effects in the oscillatory analyses only in the "spindle" condition (see Suppl. Fig. 3).

Fig. 4: Event-related responses during N2/N3 sleep depending on the presence/absence of sleep spindles. (A, B) Event-related responses in the delta (1-3Hz) range. Bar plots for the effect of *voice* (left) and corresponding scalp

plot of differences in ERS between FV and UFV (right). **(C, D)** Event-related responses in the theta/alpha/sigma (4-

15Hz) range at 4Hz and **(E, F)** responses at 14Hz. Bar plots for the effect of *voice* during the six time windows (top)

and corresponding scalp plots of differences in ERS/ERD between FV and UFV stimuli (bottom). Diamonds in

figures E and F indicate the time windows with significant differences between S+ and S- conditions at 14 Hz. Large

black dots indicate the electrodes that are part of the clusters at 2Hz, 4Hz or 14Hz, respectively. Error bars indicate

±1 standard error of the mean and *d* is Cohen's *d* for the significant clusters. Please note that for illustration purposes we

show the effects at representative frequencies (i.e. 2, 4 and 14 Hz) although significant clusters may have comprised

a larger frequency range (see main text). FV = familiar voice, UFV = unfamiliar voice. Analyses and figures are

based on data from *n* = 10 participants.

Stimulus Presentation along Slow Oscillation Positive vs. Negative Slope

Irrespective of the slope of a SO during which a stimulus was presented, analyses yielded

significant effects of *voice* (pos. slope: *Monte Carlo p < .001, d = .70* see Fig. 5A, neg. slope: *Monte*

Carlo Monte Carlo p = .003, d = .67, see Fig. 5B) in the delta range. Specifically, like in the other

conditions UFV stimuli elicited stronger ERS than did FV stimuli between 1 and 3Hz in clusters spanning

large parts of the scalp. There were no differences in delta ERS between the positive and negative SO

slope (*Monte Carlo p = .18*) and no further effects were evident in the delta range (pos. slope: *name*: no

clusters; *voice × name: p > .1*; neg. slope: *name: Monte Carlo p = .18*; *voice × name: Monte Carlo p >*

.16). In the theta to sigma range (4-15Hz) analyses also revealed significant effects of *voice* in both

conditions (pos. slope: *Monte Carlo p < .001*, see Fig. 5C; neg. slope: *Monte Carlo p = .008*, see Fig. 5D)

with UFV stimuli eliciting stronger ERS than FV stimuli following about 200ms in a broad frequency

range comparable to the effects in the other conditions regarding topography and time course (pos. slope:

T1: 9Hz; T2-6: 4-15Hz; neg. slope: T1: not part of the cluster; T2-6: 4-15Hz). There were no differences

in the theta through sigma range between the positive and negative SO slope (*Monte Carlo ps > .72*). The

effects of *voice* were also confirmed by ERP analyses (pos. slope: see Suppl. Fig. 4B; neg. slope: see

Suppl. Fig. 4C). There were no further effects in the theta through sigma range (pos. slope: *name: Monte*

Carlo ps > .27 name × voice: Monte Carlo ps > .31; neg. slope: *name*: *Monte Carlo ps > .40; name ×*

voice: *Monte Carlo ps > .48*).

Fig. 5: Event-related responses during N2/N3 sleep when a stimulus was presented along the positive

vs. negative slope of the SO. (A, B) Event-related responses in the delta (1-3Hz) range. Bar plots for the effect of

voice (left) and corresponding scalp plots of differences in ERS between FV and UFV (right). **(C, D)** Event-related

responses in the theta/alpha/sigma (4-15Hz) range. Bar plots for the effect of *voice* during the six time windows

(top) and corresponding scalp plots of differences in ERS/ERD between FV and UFV stimuli (bottom). Large black

dots indicate the electrodes that are part of the significant clusters and *d* is Cohen's *d* for the significant clusters. Error

bars indicate ±1 standard error of the mean. Please note that for illustration purposes we show the effects at

representative frequencies (i.e. 2 and 10 Hz) although significant clusters may have comprised a larger frequency

range (see main text). FV = familiar voice, UFV = unfamiliar voice. Analyses and figures are based on data from *n* =

16 participants.

Discussion

In this study we show that especially processing of paralinguistic aspects of verbal stimuli such as

the familiarity of a voice is even possible during fading and in the absence of consciousness defined as

behavioural responsiveness during sleep. The findings add to existing evidence that the detection and

evaluation of meaningful stimuli is still possible in these states (e.g. 6, 23). Intriguingly, we do not only

find that a differential response to familiar vs. unfamiliar voice (FV vs. UFV) stimuli persists during light

NREM sleep stages N1 and N2 thus replicating previous results (cf. 4, 6), but we extend this finding to

deep N3 and intriguingly also REM sleep. Beyond this, we show that transient brain states, which have

been suggested to alter sensory information processing during sleep, i.e. sleep spindles (11, 15, 16) and

slow oscillation down-states (11, 14), do, at least not uniformly or irrespective of stimulus characteristics,

inhibit stimulus processing. Rather, their inhibitory function seems to be tuned to stimulus salience.

During wakefulness, SONs seemed to be salient when compared to UNs thus drawing more

attentional resources. This was indicated by SONs eliciting stronger delta ERS than UNs across large

areas of the scalp (18, 19) and is well in line with the relatively larger P3 component evident in ERP

analyses as well as results from earlier studies (e.g. 4, 6, 24). In an earlier study from our group, del

Giudice, Lechinger (25) had also found stronger alpha ERD for SONs than for UNs, which we could not

replicate here. This may ultimately be due to methodological differences and the more conservative

statistical analysis methods employed here. Somewhat surprisingly, no differences were evident between

FV and UFV stimuli when participants were awake. However, we experience situations in which voices

of varying degrees of familiarity are present along with unfamiliar voices every day and, in comparison to

the SON, familiar voices do not usually draw attention automatically (cf. the "Cocktail Party

Phenomenon"; 26). Probably, this was even maladaptive as the manifold familiar voices would constantly

disturb orienting and eventually allocation of attention. Thus, the lack of a differential response evoked by

stimulus familiarity may well be considered adaptive.

Discussion

In this study we show that especially processing of paralinguistic aspects of verbal stimuli such as the familiarity of a voice is even possible during fading and in the absence of consciousness defined as behavioural responsiveness during sleep. The findings add to existing evidence that the detection and evaluation of meaningful stimuli is still possible in these states (e.g. 6, 23). Intriguingly, we do not only find that a differential response to familiar vs. unfamiliar voice (FV vs. UFV) stimuli persists during light NREM sleep stages N1 and N2 thus replicating previous results (cf. 4, 6), but we extend this finding to deep N3 and intriguingly also REM sleep. Beyond this, we show that transient brain states, which have been suggested to alter sensory information processing during sleep, i.e. sleep spindles (11, 15, 16) and slow oscillation down-states (11, 14), do, at least not uniformly or irrespective of stimulus characteristics, inhibit stimulus processing. Rather, their inhibitory function seems to be tuned to stimulus salience.

During wakefulness, SONs seemed to be salient when compared to UNs thus drawing more attentional resources. This was indicated by SONs eliciting stronger delta ERS than UNs across large areas of the scalp (18, 19) and is well in line with the relatively larger P3 component evident in ERP analyses as well as results from earlier studies (e.g. 4, 6, 24). In an earlier study from our group, del Giudice, Lechinger (25) had also found stronger alpha ERD for SONs than for UNs, which we could not replicate here. This may ultimately be due to methodological differences and the more conservative statistical analysis methods employed here. Somewhat surprisingly, no differences were evident between FV and UFV stimuli when participants were awake. However, we experience situations in which voices of varying degrees of familiarity are present along with unfamiliar voices every day and, in comparison to the SON, familiar voices do not usually draw attention automatically (cf. the "Cocktail Party Phenomenon"; 26). Probably, this was even maladaptive as the manifold familiar voices would constantly disturb orienting and eventually allocation of attention. Thus, the lack of a differential response evoked by stimulus familiarity may well be considered adaptive.

During NREM sleep, that is from light N1 to deep N3 sleep, we consistently find that processing

of FV vs. UFV stimuli gives rise to a differential response in the delta to sigma frequency range, an effect

that is present in oscillatory analyses as well as ERPs. Most importantly, this provides support for the

notion that processing of auditory stimuli and especially of paralinguistic stimulus aspects such as the

familiarity of a voice is incessantly processed even in states where consciousness is absent. While this is

well in line with earlier findings during light sleep stages N1 and N2 (e.g. 4, 17, 23), our results suggest

that the same holds true even for deep N3 sleep. Thus, the findings also support the notion of a 'sentinel

processing mode' of the brain during sleep, which we suggested in a previous publication (cf. 4).

Specifically, this mode describes the idea that (low-level) stimulus evaluation continues even when

consciousness fades during sleep and the result of this evaluation may subsequently either trigger an

inhibitory sleep-protecting response or awakening. In detail, we here find UFV stimuli to be associated

with stronger ERS in the delta range than FV stimuli during all NREM sleep stages, an effect which was

widespread across the scalp with the response being most pronounced above frontocentral areas.

Adopting the interpretation of delta oscillations during wakefulness, the results suggest that UFV stimuli

may become salient when consciousness fades (18, 19). In particular, the presence of unfamiliar voices

could challenge the impression of a safe environment that is necessary to 'let go of consciousness' and

eventually fall and stay asleep, rendering them salient. However, an increase in delta ERS could also be

related to a sleep-specific ERP, namely a K-complex-like response. K-complexes (KCs), whose peak

frequency is in the delta range, have been suggested to serve cortical excitation and low-level information

processing as well as the subsequent protection of sleep by neuronal silencing and they have been shown

to be elicited by salient or high-intensity stimuli (27-30). In line with the notion that KCs indicate

ongoing cognitive processing, Vallat, Lajnef (31) have recently reported a KC-like response during N2

sleep that was stronger for auditory stimuli that were followed by an arousal or awakening. The authors

concluded that this reflects stronger reactivity of the brain to external stimuli, which in turn leads to

stronger arousal. In accordance with this, ERP analyses of our data indicated that stimulus-induced

differences in the delta range indeed reflected KC-like responses evoked by stimulus presentation with

considerably larger amplitudes for UFV stimuli. In line with earlier ideas, we suggest that this ERP

reflects increased (low-level) information processing of especially salient UFV stimuli (indexed by a

larger positive wave), which is then followed by an inhibitory or sleep-protecting 'down-state' (indexed

by a larger negative wave) that is likewise scaled to stimulus salience. Further support for this

interpretation comes from analyses when stimulus presentation evoked an SO, with evoked SOs also

seeming to be sensitive to stimulus salience. Also here, UFV stimuli were associated with stronger delta

through sigma activity than FV stimuli and ERP analyses revealed that UFV stimuli were associated with

a very slight positive-going wave, which was followed by a SO down-state that appeared much more

pronounced for UFV stimuli (cf. Suppl. Fig. 4A). Besides the results obtained in the delta range, we also

find that during all NREM sleep stages UFV stimuli are associated with stronger ERS in the theta through

sigma range than FV stimuli, an effect which is most pronounced following about 200ms after stimulus

onset. Most importantly, these findings are well in line with the delta results and they provide further

convincing support for the notion that the brain is still able to process paralinguistic stimulus aspects even

when consciousness fades and is absent. On a functional level, especially frequencies in the alpha and

sigma range are thought to mirror an increase in arousal during sleep (cf. 32). This suggests that UFV

stimuli may be more arousing than FV stimuli during NREM sleep, an interpretation that, also given the

observed KC-like response, is well in line with Vallat et al.'s results. As suggested above, the presence of

unfamiliar voices may challenge the impression of an environment 'safe to sleep' and thus be arousing.

Admittedly, our findings during N2 sleep partly contrast results earlier studies, where the brain also

seemed to continue differentiating between UNs and SONs (e.g. 4, 6). The deviating findings could be

due to methodological differences and/or participants sleeping during a whole night and not just an

afternoon nap (cf. 4) with differences in the homeostatic and circadian factors rendering it questionable

whether a daytime nap can be considered a short night sleep equivalent.

In summary, results obtained during wakefulness and NREM sleep suggest that familiarity of a

voice can be processed even during the fading of consciousness (N1) and in the full absence of

(behavioural) consciousness (N2 and N3). For REM sleep, a paradoxical state characterised by (i) the

return of 'altered consciousness', namely 'dreaming', (ii) enhanced brain metabolism (33, 34) and (iii) an

increase in higher frequency EEG power (35), we also observed a relatively stronger increase in delta as

well as alpha/sigma ERS elicited by UFV compared to FV stimuli, which may indicate continued

processing and/or arousal of salient or potentially 'dangerous' UFV stimuli. This is especially interesting

because REM sleep has been suggested to reflect a 'closed loop', that is a state in which the brain is rather

occupied with intrinsic activity than processing of external stimuli (36-38) with our results challenging

this notion. At the same time, while the oscillatory response pattern was generally similar to NREM sleep

findings, REM responses were considerably weaker and markedly delayed by approx. 400ms. This

underlines the idea that brain activity and processing of environmental stimuli during REM is

qualitatively different although not generally precluded.

Beyond investigating stimulus processing across longer lasting brain states, i.e. wakefulness and

different sleep stages, we were also interested in how stimulus presentation relates to 'transient oscillatory

activity', that is sleep spindles and slow oscillations (SOs), during N2 and N3 sleep. Generally, sleep

spindles (11, 15) and the negative slope of slow oscillations (11) have been suggested to inhibit

processing of external stimuli. Here we find that this does not seem to be universally true but that brain

responses are still tuned to stimulus salience suggesting that at least 'low-level' processing is not

precluded. More specifically, we find that when a sleep spindle overlapped with stimulus presentation

UFV stimuli still elicited responses in the delta through lower alpha (i.e. up to about 9Hz) range that were

similar to those obtained when not taking 'transient oscillatory activity' into account. Intriguingly and

unlike proposed earlier (9, 11), this suggests that processing of external stimuli is not or at least not

uniformly inhibited by the presence of a sleep spindle, i.e. spindles do not generally seem to act as a

sensory filter at the thalamic level. Interestingly, this is well in line with recent findings in rodents where

thalamocortical sensory relay was shown to persist even during sleep spindles (39). Beyond this, above

≈9Hz the response pattern when a spindle was present was markedly different from the general NREM

(see Fig. 2) and, most importantly, the 'no spindle' (see Fig. 4F) patterns with FV stimuli eliciting

stronger ERD than UFV stimuli in the ≈11-15Hz spindle range (see Fig. 4C). We speculate that this could

reflect a relatively stronger release of inhibition (reflected by 10-15Hz ERD) for seemingly less relevant

FV stimuli by sleep spindles. Arguably, a selective mechanism that specifically filters information that is

considered irrelevant, i.e. here FV stimuli, seems more adaptive than the uniform inhibition of all

environmental stimuli. Following the idea of a 'sentinel processing mode' of the brain during sleep,

spindles just as slow oscillations could thus reflect a sleep-protecting response that follows initial

stimulus evaluation during N2 and N3. Besides sleep spindles, previous studies suggested that also the

slope of a SO during stimulus presentation affects stimulus processing. In particular the negative slope

has been found to be associated with decreased responses in studies using somatosensory stimuli and

simple tones as compared to the positive SO slope. Surprisingly, in our study stimulus delivery during

negative and positive slopes revealed similar responses with responses in both conditions being tuned to

stimulus salience. Specifically, as during all other sleep stages UFV stimuli elicited stronger (delta to

sigma) ERS than FV stimuli. These results were supported by ERP analyses indicating that UFV stimuli

induced a more pronounced down-state that was preceded by an up-state. The findings thereby contrast

earlier findings and suggest that also the negative slope of a SO does at least not uniformly inhibit

information processing and allows continued evaluation of stimulus characteristics. Likewise, the findings

also suggest that during a positive SO slope the brain is not uniformly open to external stimulation.

In conclusion, this study shows that stimulus characteristics and especially the familiarity of a

voice continue to be evaluated during all stages of NREM sleep and thus even in the complete absence of

behavioural consciousness. Surprisingly, this is the case even during REM sleep with processing of

external seeming to be slowed and decreased though. Our findings thereby provide support for the idea of

a 'sentinel processing mode' of the brain during sleep, i.e. the continued processing of environmental

stimuli even in the absence of consciousness that may then be followed by either an inhibitory sleep-

protective response or awakening depending on the result of stimulus evaluation. Beyond this, it appears

that even 'transient oscillatory activity', i.e. sleep spindles and slow oscillations are sensitive to

paralinguistic emotional stimulus characteristics. Furthermore, we provide novel evidence that even

during spindles and the negative slope of a SO the brain reacts differentially to incoming information. In a

wider context, our findings also suggest that using emotional stimuli such as familiar voices, or favourite

sounds and music may be helpful in the medical and therapeutic context when patients are in states of

reduced or altered awareness e.g. following severe brain injury. Besides, the results may open up new

perspectives for insomnia research, where a relative deficit in processing of environmental stimuli during

sleep may be related to problems of "letting go of consciousness" and thus falling asleep.

Methods and Materials

Participants

We recruited 20 healthy individuals for the study. Three participants were excluded from the data

analysis, one dropped out after the adaptation night and two had to be excluded due to technical problems

during the acquisition. The remaining sample comprised 17 participants (three males) and had a median

age of 22.6 years ($SD = 2.3$ years). Prior to the study, participants gave written informed consent. Ethical

consent had been obtained from the ethics committee of the University of Salzburg and the study was in

accordance with the Declaration of Helsinki (40). For more details on the study sample please see

supplementary material.

Experimental procedure

Participants were advised to keep a regular sleep/wake rhythm with eight hours time in bed (TIB) for

at least four days prior to their first visit at our sleep laboratory, which was verified with wrist actigraphy

(Cambridge Neurotechnology Actiwatch ©). Participants slept in the sleep laboratory of the University of

Salzburg for two nights, one adaptation night and one experimental night.

The adaptation and experimental nights were comparable except for no auditory stimulation during

sleep taking place during the adaptation night. On both nights and the following mornings participants

were tested during wakefulness resulting in four wakefulness recordings per participant. The wakefulness

part comprised a passive listening as well as an active counting condition, during which participants

listened to the stimuli presented via in-ear headphones at a volume of approximately 65 dB. For the

passive condition participants were instructed to listen attentively to the stimuli while in the active

condition they were to count the number of presentations of one specific name (i.e. the target). The

passive condition always preceded the active one. In this publication, we only present the results from the

passive listening condition, in which participants were presented with their own name (SON) as well as

two unfamiliar names (UNs) as it is the only condition that can be analysed meaningfully across 'longer

lasting brain' or vigilance stages (i.e. wakefulness, NREM and REM sleep). Moreover, each name was

uttered by a familiar (mother, father) and by an unfamiliar voice (lab member unknown to participant).

The stimulus set was specific for each participant and all names of one stimulus set were matched

regarding the number of syllables and the occurrence in the general population. During the wakefulness

recording, each stimulus was presented 40 times and the interstimulus interval (ISI) was 2000ms.

Following the wakefulness recordings in the evenings, participants went to bed for eight hours of

sleep. During the experimental night, stimulation was continued and the volume was adjusted individually

so stimuli were clearly audible, but participants felt they could sleep despite the stimulation. The auditory

stimulation protocol was akin to the passive condition of the wake part, although during the night, the

stimulus onset asynchrony (SOA) was jittered between 2.8 and 7.8 s in 500ms steps. SOA was jittered

specifically in the sleep protocol as this was necessary to allow for an investigation of stimulus processing

in relation to various EEG sleep phenomena (i.e. sleep spindles and slow oscillations) independent of

expectation effects. SOA was not jittered during wakefulness as this would have rendered the tasks

lengthy and probably too fatiguing. During the night each stimulus was presented 690 times and had the

same probability of occurrence as had each SOA. For more details on the experimental procedure please

see the supplementary material.

Electrophysiological data collection and reduction

475   For EEG acquisition we used a 256 electrode GSN HydroCel Geodesic Sensor Net (Electrical

476   Geodesics Inc., Eugene, Oregon, USA) and a Net Amps 400 amplifier.

Wakefulness data

EEG data were processed using the Fieldtrip toolbox (41) in Matlab (Mathworks, Natick, USA).

First, the number of electrodes was reduced to 183 as the others contained a lot of 'non-neural' artefacts

and high-pass filtered at 0.5Hz. Subsequently, eye movement artefacts were corrected using independent

component analysis (ICA), data were segmented into 4s epochs (symmetrically to stimulus onset) and bad

intervals were removed manually during visual data inspection. In the next step, the number of electrodes

was further reduced to a final number of 173 electrodes now excluding 10 more electrodes that had

initially been kept for the identification of eye and muscular artefacts. Bad channels identified during

visual data inspection were interpolated and data were re-referenced to average reference. Subsequently,

we randomly selected the same number of trials for each stimulus to account for imbalances in the

stimulus set (only one SON, but two UNs were presented). We then applied a Morlet wavelet

transformation (cycles = 3, 1-16Hz, 1Hz frequency steps) to each of the segments, which was followed by

a baseline correction (baseline interval: -600 to 0ms relative to stimulus onset) and averaging across trials.

Sleep data

Sleep was scored semi-automatically by The Siesta Group© (Somnolyzer 24×7; cf. 42, 43)

according to standard criteria (32). Spindles were detected automatically during NREM sleep stages N2

and N3 at central leads using the algorithm by Anderer, Gruber (42). Slow oscillations (SOs) were also

detected automatically on frontal electrodes using lab-internal Matlab routines (cf. 44) based on the

criteria by Massimini, Huber (5). For more details on the detection of spindles and SOs please see

supplementary material. Pre-processing for the sleep data was essentially the same as for the wakefulness

data; but we refrained from an automatic eye movement correction in order to not remove REMs. Beyond

investigating processing of different stimuli across 'longer lasting brain states', that is in each sleep stage,

we also investigated stimulus processing with regard to 'transient brain states', that is sleep spindles and

SOs. To this end, we compared evoked oscillatory responses elicited by different stimuli when a spindle

was present during stimulus onset (i.e. spindle offset min. 200ms after stimulus onset) or when there was

a substantial overlap between a spindle and stimulus presentation (spindle onset 0-400ms after stimulus

onset, i.e. spindle overlapping with at least half of the stimulus on average, cf. Suppl.Fig.1, A). Moreover,

we were interested in stimulus-specific differences in the evoked slow oscillatory responses ("SO

evoked"). More precisely, a SO was defined as "evoked" when the negative peak occurred between 300

and 600ms after stimulus onset (cf. Suppl.Fig.1, B1), that is the time range when the negative components

of evoked K-complexes (i.e. N350 and N550) have been found to occur (45). Beyond this, we compared

stimulus processing when stimulus onset was during the positive going slope of a SO (cf. Suppl.Fig.1,

B2) to when stimulus onset coincided with the down-state (cf. Suppl.Fig.1, B3). For more details on data

collection and analysis please refer to the supplementary material.

Event-Related Potentials

Although we focus on oscillatory activity in different frequency bands in the present manuscript,

we provide results from event-related potential (ERP) analyses in the supplementary material (and Fig.1).

Statistical Analyses

Statistical analyses were performed using the cluster-based permutation approach implemented in

Fieldtrip to correct for multiple comparisons that uses a Monte Carlo method for calculating significance

probabilities (46). This approach has originally been introduced by (47) and is referred to as the 'cluster

mass test' in the fMRI literature (for more details please see suppl. material). Three tests were run for the

main effects of *name* (SON vs. UNs), *voice* (FV vs. UFV) and the *name* × *voice* interaction with

significant (or marginally significant) interaction clusters being followed by post-hoc tests. Thus, we

report three *p*-values per condition (i.e. sleep stage and interaction with sleep spindle or SO). We ran a

first set of tests for the delta range that included the dimensions electrode and frequency (1-3Hz in 1Hz frequency steps). In the delta range, values were averaged across time (0-1000ms after stimulus onset for the WAKE condition, 0-1200ms during SLEEP) as time resolution obtained with these low frequencies was considered insufficient for an analysis in the time dimension. A second test was then run for the theta, alpha and sigma ranges including the dimensions electrode, frequency (4-15Hz, 1Hz frequency steps) and time (0-1000ms, five time windows à 200ms each in the WAKE condition, 6 time windows from 0-1200ms during SLEEP). For the "spindle vs. no spindle" and "negative vs. positive SO slope" contrasts we calculated averaged values for FV/UVF for each condition, which we then compared. For all permutation tests the critical *p*-value for the *T*-statistic for dependent samples was set to 0.05 and 1000 randomisations were used. Spatial clusters were formed only if electrodes had a minimum of two neighbouring electrodes that were also significant. We report the Monte Carlo approximation for the estimate of *p*-values. Effects with (one-sided) *Monte Carlo p* < .05 are denoted significant, effects with *Monte Carlo p* < .1 are denoted marginally significant. Critical *p*-values for post-hoc tests were adjusted for multiple comparisons using Bonferroni-corrected *p*-values. We report Cohen's *d* (*d* = mean difference in a significant cluster/standard deviation of the differences) as a measure of the effect size for all analyses. For more details on the statistical analysis please see the supplementary material.

Conflict of interest

The authors declare no competing financial interests.

Acknowledgements

We thank Daniel Koerner and Adriana Michalak for their help with the data collection and pre-processing and Vincenzo Muto for his advice on the sleep staging. We also thank Wolfgang Klimesch for the valuable discussion about the results.

References

1.   Santhi N, Lazar AS, McCabe PJ, Lo JC, Groeger JA, Dijk D-J. Sex differences in the circadian

regulation of sleep and waking cognition in humans. Proceedings of the National Academy of Sciences of

the United States of America. 2016;113(19):E2730-E9.

2.   Dijk D-J, Duffy JF, Czeisler CA. Circadian and sleep/wake dependent aspects of subjective

alertness and cognitive performance. Journal of Sleep Research. 1992;1(2):112-7.

3.   Wyatt JK, Cecco AR-D, Czeisler CA, Dijk D-J. Circadian temperature and melatonin rhythms,

sleep, and neurobehavioral function in humans living on a 20-h day. American Journal of Physiology -

Regulatory, Integrative and Comparative Physiology. 1999;277(4):R1152-R63.

4.   Blume C, del Giudice R, Lechinger J, Wislowska M, Heib DPJ, Hoedlmoser K, et al. Preferential

processing of emotionally and self-relevant stimuli persists in unconscious N2 sleep. Brain and language.

2016.

5.   Massimini M, Huber R, Ferrarelli F, Hill S, Tononi G. The sleep slow oscillation as a traveling

wave. The Journal of Neuroscience. 2004;24(31):6862-70.

6.   Perrin F, Garcia-Larrea L, Mauguiere F, Bastuji H. A differential brain response to the subject's

own name persists during sleep. Clin Neurophysiol. 1999;110(12):2153-64.

7.   Bastuji H, García-Larrea L. Evoked potentials as a tool for the investigation of human sleep.

Sleep Medicine Reviews. 1999;3(1):23-45.

8.   Strauss M, Sitt JD, King J-R, Elbaz M, Azizi L, Buiatti M, et al. Disruption of hierarchical

predictive coding during sleep. Proceedings of the National Academy of Sciences. 2015;112(11):E1353-

E62.

9.   Steriade M. Normal and Altered States of Function. Cerebral Cortex. 1991;9:279e357.

10.   Massimini M, Amzica F. Extracellular calcium fluctuations and intracellular potentials in the

cortex during the slow sleep oscillation. Journal of Neurophysiology. 2001;85(3):1346-50.

11. Schabus M, Dang-Vu TT, Heib DPJ, Boly M, Desseilles M, Vandewalle G, et al. The fate of

incoming stimuli during NREM sleep is determined by spindles and the phase of the slow oscillation.

Frontiers in Neurology. 2012;3.

12. Dang-Vu TT, Bonjean M, Schabus M, Boly M, Darsaud A, Desseilles M, et al. Interplay between

spontaneous and induced brain activity during human non-rapid eye movement sleep. Proceedings of the

National Academy of Sciences. 2011;108(37):15438-43.

13. De Gennaro L, Ferrara M. Sleep spindles: an overview. Sleep Medicine Reviews. 2003;7(5):423-

40.

14. Massimini M, Rosanova M, Mariotti M. EEG slow ($\sim$ 1 Hz) waves are associated with

nonstationarity of thalamo-cortical sensory processing in the sleeping human. Journal of

Neurophysiology. 2003;89(3):1205-13.

15. Elton M, Winter O, Heslenfeld D, Loewy D, Campbell K, Kok A. Event-related potentials to

tones in the absence and presence of sleep spindles. Journal of Sleep Research. 1997;6(2):78-83.

16. Cote KA, Epps TM, Campbell KB. The role of the spindle in human information processing of

high-intensity stimuli during sleep. Journal of Sleep Research. 2000;9(1):19-26.

17. Oswald I, Taylor AM, Treisman M. Discriminative responses to stimulation during human sleep.

Brain : a journal of neurology. 1960.

18. Knyazev GG. EEG delta oscillations as a correlate of basic homeostatic and motivational

processes. Neuroscience & Biobehavioral Reviews. 2012;36(1):677-95.

19. Knyazev GG. Motivation, emotion, and their inhibitory control mirrored in brain oscillations.

Neuroscience & Biobehavioral Reviews. 2007;31(3):377-95.

20. Klimesch W. EEG alpha and theta oscillations reflect cognitive and memory performance: a

review and analysis. Brain Research Reviews. 1999;29(2-3):169-95.

21. Klimesch W, Schack B, Sauseng P. The functional significance of theta and upper alpha

oscillations. Experimental Psychology. 2005;52(2):99-108.

22.   Klimesch W, Doppelmayr M, Russegger H, Pachinger T, Schwaiger J. Induced alpha band power

changes in the human EEG and attention. Neurosci Lett. 1998;244(2):73-6.

23.   Portas CM, Krakow K, Allen P, Josephs O, Armony JL, Frith CD. Auditory processing across the

sleep-wake cycle: simultaneous EEG and fMRI monitoring in humans. Neuron. 2000;28(3):991-9.

24.   Berlad I, Pratt H. P300 in response to the subject's own name. Electroencephalography and

clinical neurophysiology. 1995;96(5):472-4.

25.   del Giudice R, Lechinger J, Wislowska M, Heib DP, Hoedlmoser K, Schabus M. Oscillatory

brain responses to own names uttered by unfamiliar and familiar voices. Brain research. 2014;1591:63-73.

26.   Wood N, Cowan N. The cocktail party phenomenon revisited: how frequent are attention shifts to

one's name in an irrelevant auditory channel? Journal of Experimental Psychology: Learning, Memory,

and Cognition. 1995;21(1):255-60.

27.   Amzica F, Steriade M. The K-complex: Its slow (<1-Hz) rhythmicity and relation to delta waves.

Neurology. 1997;49(4):952-9.

28.   Cash SS, Halgren E, Dehghani N, Rossetti AO, Thesen T, Wang C, et al. The human K-complex

represents an isolated cortical down-state. Science. 2009;324(5930):1084-7.

29.   Bastien C, Campbell K. The evoked K-complex: all-or-none phenomenon? Sleep.

1992;15(3):236-45.

30.   Laurino M, Menicucci D, Piarulli A, Mastorci F, Bedini R, Allegrini P, et al. Disentangling

different functional roles of evoked K-complex components: Mapping the sleeping brain while quenching

sensory processing. NeuroImage. 2014;86:433-45.

31.   Vallat R, Lajnef T, Eichenlaub J-B, Berthomier C, Jerbi K, Morlet D, et al. Increased Evoked

Potentials to Arousing Auditory Stimuli during Sleep: Implication for the Understanding of Dream

Recall. Frontiers in human neuroscience. 2017;11(132).

32.   American Academy of Sleep Medicine, Iber C. The AASM manual for the scoring of sleep and

associated events: rules, terminology and technical specifications: American Academy of Sleep Medicine;

2007.

33.   Nofzinger EA, Mintun MA, Wiseman M, Kupfer DJ, Moore RY. Forebrain activation in REM

sleep: an FDG PET study. Brain research. 1997;770(1):192-201.

34.   Maquet P. Functional neuroimaging of normal human sleep by positron emission tomography.

Journal of sleep research. 2000;9(3):207-32.

35.   Uchida S, Maloney T, Feinberg I. Beta (20-28 Hz) and delta (0.3-3 Hz) EEGs oscillate

reciprocally across NREM and REM sleep. Sleep. 1992;15(4):352-8.

36.   Wehrle R, Kaufmann C, Wetter TC, Holsboer F, Auer DP, Pollmächer T, et al. Functional

microstates within human REM sleep: first evidence from fMRI of a thalamocortical network specific for

phasic REM periods. European Journal of Neuroscience. 2007;25(3):863-71.

37.   Llinás RR, Paré D. Of dreaming and wakefulness. Neuroscience. 1991;44(3):521-35.

38.   Andrillon T, Poulsen AT, Hansen LK, Léger D, Kouider S. Neural Markers of Responsiveness to

the Environment in Human Sleep. The Journal of Neuroscience. 2016;36(24):6583-96.

39.   Sela Y, Vyazovskiy VV, Cirelli C, Tononi G, Nir Y. Responses in rat core auditory cortex are

preserved during sleep spindle oscillations. Sleep. 2016;39(5):1069-82.

40.   World Medical Association (WMA). Ethical Principles for Medical Research Involving Human

Subjects 1964 [10/13:[Available from: http://www.wma.net/en/30publications/10policies/b3/.

41.   Oostenveld R, Fries P, Maris E, Schoffelen J-M. FieldTrip: open source software for advanced

analysis of MEG, EEG, and invasive electrophysiological data. Computational Intelligence and

Neuroscience. 2010;2011.

42.   Anderer P, Gruber G, Parapatics S, Woertz M, Miazhynskaia T, Klösch G, et al. An E-health

solution for automatic sleep classification according to Rechtschaffen and Kales: validation study of the

Somnolyzer 24× 7 utilizing the Siesta database. Neuropsychobiology. 2005;51(3):115-33.

43.   Anderer P, Saletu B, Saletu-Zyhlarz GM, Gruber G, Parapatics S, Miazhynskaia T, et al. Recent

advances in the electrophysiological evaluation of sleep. Essentials and applications of EEG research in

preclinical and clinical pharmacology Berlin: Unipublish Verlag für Studium & Praxis OHG. 2004:307-

39.

44.   Heib DPJ, Hoedlmoser K, Anderer P, Zeitlhofer J, Gruber G, Klimesch W, et al. Slow Oscillation

Amplitudes and Up-State Lengths Relate to Memory Improvement. PloS one. 2013;8(12):e82049.

45.   Cote KA, De Lugt DR, Langley SD, Campbell KB. Scalp topography of the auditory evoked K-

complex in stage 2 and slow wave sleep. Journal of Sleep Research. 1999;8(4):263-72.

46.   Maris E, Oostenveld R. Nonparametric statistical testing of EEG-and MEG-data. Journal of

Neuroscience Methods. 2007;164(1):177-90.

47.   Bullmore ET, Suckling J, Overmeyer S, Rabe-Hesketh S, Taylor E, Brammer MJ. Global, voxel,

and cluster tests, by theory and permutation, for a difference between two groups of structural MR images

of the brain. IEEE Transactions on Medical Imaging. 1999;18(1):32-42.

Supporting Information

S1: Supplementary Material (Methods, Results)