The lectin-specific activity of *Toxoplasma gondii* microneme proteins 1

and 4 binds Toll-like receptor 2 and 4 N-glycans to regulate innate

immune priming

Aline Sardinha-Silva[1,2], Flávia C. Mendonça-Natividade[1], Camila F. Pinzan[1], Carla D.

Lopes[1], Diego L. Costa[2], Damien Jacot[3], Fabricio F. Fernandes[1], André L. V. Zorzetto-

Fernandes[1], Nicholas J. Gay[4], Alan Sher[2], , Dragana Jankovic[2], Dominique Soldati-

Favre[3], Michael E. Grigg[2*], Maria Cristina Roque-Barreira[1*]

[1] Department of Cell and Molecular Biology and Pathogenic Bioagents, Ribeirão Preto

Medical School, University of São Paulo- USP (FMRP/USP), Ribeirão Preto, São

Paulo, 14049-900, Brazil

[2] Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases,

National Institutes of Health, Bethesda, MD 20892, USA

[3] Department of Microbiology and Molecular Medicine, CMU, University of Geneva, 1

rue Michel-Servet, 1211 Geneva 4, Switzerland

[4] Department of Biochemistry, Cambridge University, 80 Tennis Court Road

Cambridge CB2 1GA, United Kingdom

*Corresponding authors:

Michael E. Grigg, Phone: +1 (301) 402-1609, e-mail: griggm@niaid.nih.gov; and Maria

Cristina Roque-Barreira, Phone: +55-16-3315-3062, e-mail: mcrbarre@fmrp.usp.br

ABSTRACT

Infection of host cells by *Toxoplasma gondii* is an active process, which is regulated by secretion of microneme (MICs) and rhoptry proteins (ROPs and RONs) from specialized organelles in the apical pole of the parasite. MIC1, MIC4 and MIC6 assemble into an adhesin complex, secreted on the parasite surface and function to promote infection competency. MIC1 and MIC4 are known to bind terminal sialic acid residues and galactose residues, respectively and to induce IL-12 production from splenocytes. Here we show that rMIC1- and rMIC4-stimulated dendritic cells and macrophages to produce proinflammatory cytokines, and they do so by engaging TLR2 and TLR4. This process depends on sugar recognition, since point mutations in the carbohydrate-recognition domains (CRD) of rMIC1 and rMIC4 inhibit innate immune cells activation. HEK cells transfected with TLR2 glycomutants were selectively unresponsive to MICs. Following *in vitro* infection, parasites lacking MIC1 or MIC4, as well as expressing MIC proteins with point mutations in their CRD, failed to induce wild-type (WT) levels of IL-12 secretion by innate immune cells. However, only MIC1 was shown to impact systemic levels of IL-12 and IFN-γ *in vivo*. Together, our data show that MIC1 and MIC4 interact physically with TLR2 and TLR4 N-glycans to trigger IL-12 responses, and MIC1 is playing a significant role *in vivo* by altering *T. gondii* infection competency and murine pathogenesis.

AUTHOR SUMMARY

Toxoplasmosis is caused by the protozoan *Toxoplasma gondii*, belonging to the Apicomplexa phylum. This phylum comprises important parasites able to infect a broad diversity of animals, including humans. A particularity of *T. gondii* is its ability to invade virtually any nucleated cell of all warm-blooded animals through an active process, which depends on the secretion of adhesin proteins. These proteins are discharged by specialized organelles localized in the parasite apical region, and termed micronemes and rhoptries. We show in this study that two microneme proteins from *T. gondii* utilize their adhesion activity to stimulate innate immunity. These microneme proteins, denoted MIC1 and MIC4, recognize specific sugars on receptors expressed on the surface of mammalian immune cells. This binding activates these innate immune cells to secrete cytokines, which promotes efficient host defense mechanisms against the

parasite and regulate their pathogenesis. This activity promotes a chronic infection by

controlling parasite replication during acute infection.

INTRODUCTION

Toxoplasma gondii is a coccidian parasite belonging to the phylum Apicomplexa

and is the causative agent of toxoplasmosis. This protozoan parasite infects a variety of

vertebrate hosts, including humans with about one-third of the global population being

chronically infected [1]. Toxoplasmosis can be fatal in immunocompromised

individuals or when contracted congenitally [1], and is considered the second leading

cause of death from foodborne illnesses in the United States [2].

T. gondii invades host cells through an active process that relies on the parasite

actinomyosin system, concomitantly with the release of microneme proteins (MICs) and

rhoptry neck proteins (RONs) from specialized organelles in the apical pole of the

parasite [3]. These proteins are secreted by tachyzoites [4, 5] and form complexes

composed of soluble and transmembrane proteins. Some of the MICs act as adhesins,

interacting tightly with host cell-membrane glycoproteins and receptors, and are

involved in the formation of the moving junction [6]. This sequence of events ensures

tachyzoite gliding motility, migration through host cells, invasion and egress from

infected cells [4, 7]. Among the released proteins, MIC1, MIC4, and MIC6 form a

complex that, together with other *T. gondii* proteins, plays a role in the adhesion and

invasion of host cells [8, 9], contributing to the virulence of the parasite [10, 11].

Several studies have shown that host-cell invasion by apicomplexan parasites

such as *T. gondii* involves carbohydrate recognition [12-15]. Interestingly, MIC1 and

MIC4 have lectin domains [11, 16-18] that recognize oligosaccharides with sialic acid

and D-galactose in the terminal position, respectively. Importantly, the parasite's Lac$^+$

subcomplex, consisting of MIC1 and MIC4, induces adherent spleen cells to release IL-

12 [17], a cytokine critical for the protective response of the host to *T. gondii* infection

[19]. In addition, immunization with this native subcomplex, or with recombinant MIC1

(rMIC1) and MIC4 (rMIC4), protects mice against experimental toxoplasmosis [20, 21].

The induction of IL-12 is typically due to detection of the pathogen by innate immunity

receptors, including members of the Toll-like receptor (TLR) family, whose stimulation

involves MyD88 activation and priming of Th1 responses, which protects the host

against *T. gondii* [19, 22]. It is also known that dysregulated expression of IL-12 and

IFN-γ during acute toxoplasmosis can drive a lethal immune response, in which mice

succumb to infection by severe immunopathology, the result of insufficient levels of IL-

10 and/or a collapse in the regulatory CD4+Foxp3+ T cell population [23, 24].

Interestingly, regarding the innate immune receptors associated with IL-12

response during several infections, the extracellular leucine-rich repeat domains of

TLR2 and TLR4 contain four and nine N-glycans, respectively [25]. Therefore, we

hypothesized that MIC1 and MIC4 bind TLR2 and TLR4 N-glycans on antigen-

presenting cells (APCs) and, through this interaction, trigger immune cell activation and

IL-12 production. To investigate this possibility, we assayed the ability of rMIC1 and

rMIC4 to bind and activate TLR2 and TLR4. Using several strategies, we demonstrated

that TLR2 and TLR4 are indeed critical targets for both MIC1 and MIC4. These

parasite and host cell structures establish lectin-carbohydrate interactions that contribute

to the induction of IL-12 production by innate immune cells, and we show here that the

MIC1 lectin promotes *T. gondii* infection competency and regulates parasite virulence

during *in vivo* infection.

RESULTS

Lectin properties of recombinant MIC1 and MIC4 are consistent with those of the

native Lac⁺ subcomplex

The native MIC1/4 subcomplex purified from soluble *T. gondii* antigens has

lectin properties, so we investigated whether their recombinant counterparts retained the

sugar-binding specificity. The glycoarray analysis revealed the interactions of: i) the

Lac⁺ subcomplex with glycans containing terminal α(2-3)-sialyl and β(1-4)- or β(1-3)-

galactose; ii), rMIC1 with α(2-3)-sialyl residues linked to β-galactosides; and iii) of

rMIC4 with oligosaccharides with terminal β(1-4)- or β(1-3)-galactose (Fig 1A). The

combined specificities of the individual recombinant proteins correspond to the dual

sugar specificity of the Lac⁺ fraction, demonstrating that the sugar-recognition

properties of the recombinant proteins are consistent with those of the native ones.

Based on the sugar recognition selectivity of rMIC1 and rMIC4, we tested two

oligosaccharides (α(2-3)-sialyllactose and lacto-N-biose) for their ability to inhibit the

interaction of the MICs with the glycoproteins fetuin and asialofetuin [26]. Sialyllactose

inhibited the binding of rMIC1 to fetuin, and lacto-N-biose inhibited the binding of

rMIC4 to asialofetuin (Fig 1B). To ratify the carbohydrate recognition activity of rMIC1

and rMIC4, we generated point mutations into the carbohydrate recognition domains

(CRDs) of the rMICs to abolish their lectin properties [11, 18, 27]. These mutated

forms, i.e. rMIC1-T126A/T220A and rMIC4-K469M, lost their capacity to bind to

fetuin and asialofetuin, respectively (Fig 1B), having absorbance as low as that in the

presence of the specific sugars. Thus, our results indicate that rMIC1 and rMIC4

maintained their lectin properties, and that the CRD function can be blocked either by

competition with specific sugars or by targeted mutations.

rMIC1 and rMIC4 trigger the activation of DCs and macrophages

We have previously demonstrated that the native Lac[+] subcomplex stimulates

murine adherent spleen cells to produce proinflammatory cytokines [20]. We evaluated

whether recombinant MIC1 and MIC4 retained this property and exerted it on BMDCs

and BMDMs. BMDCs (Fig 2A-2D) and BMDMs (Fig 2E-2H) produced high levels of

the proinflammatory cytokines IL-12 (Fig 2A and 2E), TNF-α (Fig 2B and 2F), and IL-

6 (Fig 2C and 2G). This was not attributable to residual LPS contamination as the

recombinant protein assays were done in the presence of polymyxin B, and LPS levels

were less than 0.5ng/ml [see Materials and Methods section]. Although conventional

CD4[+] Th1 cells are known to be the major producers of IL-10 during murine *T. gondii*

infection [28], we also found that rMIC1 and rMIC4 induced the production of this

cytokine by BMDCs (Fig 2D) and BMDMs (Fig 2H). We verified that the two

recombinant proteins induced the production of similar levels of IL-12, TNF-α, and IL-

6 by both BMDCs (Fig 2A-2C) and BMDMs (Fig 2E-2G). Both MICs induced the

production of similar levels of IL-10 in BMDCs (Fig 2D); however, BMDMs produced

significantly higher levels of IL-10 when stimulated with rMIC1 than when stimulated

with rMIC4 (Fig 2H). These cytokine levels were similar to those induced by the TLR4

agonist LPS. Thus, recombinant MIC1 and MIC4 induce a proinflammatory response in

innate immune cells, which is consistent with the results obtained for the native Lac[+]

subcomplex [20].

The activation of macrophages by rMIC1 and rMIC4 depends on TLR2 and TLR4

To investigate the mechanisms through which *T. gondii* MIC1 and MIC4

stimulate innate immune cells to produce cytokines, we assessed whether these MICs

can activate specific TLRs. To this end, BMDMs from WT, MyD88[-/-], TRIF[-/-], TLR2[-/-],

TLR4[-/-], or TLR2/4 DKO mice, as well as HEK293T cells transfected with TLR2 or

TLR4, were cultured in the presence or absence of rMIC1 and rMIC4 for 48 hours. The

production of IL-12 by BMDMs (Fig 3A-3I) and IL-8 by HEK cells (Fig 3J and 3K)

were used as an indicator of cell activation. IL-12 production by BMDMs from MyD88[-]

[/-], TRIF[-/-], TLR2[-/-], and TLR4[-/-] mice was lower than that of BMDMs from WT mice

(Fig 3A-3D); no IL-12 was detected in cultures of TLR2/4 DKO mice cells stimulated

with either rMIC1 or rMIC4 (Fig 3E). These results show that TLR2 and TLR4 are both

relevant for the activation of macrophages induced by rMIC1 and rMIC4. The residual

cytokine production observed in macrophages from TLR2[-/-] or MyD88[-/-] mice may be

the result of activation of TLR4 (Fig 3A and 3C), and vice versa; e.g., the residual IL-12

levels produced by macrophages from TLR4[-/-] mice may be the result of TLR2

activation. The finding that MICs fail to induce IL-12 production in DKO mice

BMDMs suggests that cell activation triggered by *T. gondii* MIC1 or MIC4 does not

require the participation of other innate immunity receptors beyond TLR2 and TLR4.

Nevertheless, because parasite components such as DNA or profilin engage TLR9,

TLR11, and TLR12 to produce IL-12 in macrophages [19, 22, 29], we investigated the

involvement of these receptors, as well as TLR3 and TLR5, in the response to rMIC1 or

rMIC4. BMDMs from TLR3[-/-], TLR5[-/-], TLR9[-/-], and TLR11/12 DKO mice stimulated

with rMIC1 or rMIC4 produced similar levels of IL-12 as cells from WT (Fig 3F-3I),

indicating that the activation triggered by rMIC1 or rMIC4 does not depend on these

receptors. Additionally, stimulation of HEK cells transfected with human TLR2 (Fig 3J)

or TLR4 (Fig 3K) with optimal concentrations of rMIC1 (Fig S1A and S1C) and rMIC4

(Fig S1B and S1D) induced IL-8 production at levels that were higher than those

detected in the absence of stimuli (medium), and similar to those induced by the

positive controls. Finally, by means of a pull-down experiment, we demonstrated a

physical interaction between rMIC1 and TLR2 or TLR4 and between rMIC4 and TLR2

or TLR4 (Fig 3L).

Cell activation induced by rMIC1 and rMIC4 results from the interaction of their

CRDs with TLR2 and TLR4 N-glycans

We hypothesized that in order to trigger cell activation, rMIC1 and rMIC4 CRDs

target oligosaccharides of the ectodomains of TLR2 (four N-linked glycans) [25] and

TLR4 (nine N-linked glycans) [30]. This hypothesis was tested by stimulating BMDCs

(Fig 4A) and BMDMs (Fig 4B) from WT mice with intact rMIC1 and rMIC4 or with

the mutated forms of these microneme proteins, namely rMIC1-T126A/T220A and

rMIC4-K469M, which lack carbohydrate binding activity [11, 18, 27]. IL-12 levels in

culture supernatants were lower upon stimulation with rMIC1-T126A/T220A or rMIC4-K469M, showing that WT induction of cell activation requires intact rMIC1 and rMIC4 CRDs. The same microneme proteins were used to stimulate TLR2-transfected HEK293T cells (Fig 4C), and similarly, lower IL-8 production was obtained in response to mutated rMIC1 or rMIC4 compared to that seen in response to intact proteins. These observations demonstrated that rMIC1 and rMIC4 CRDs are also necessary for inducing HEK cell activation.

We used an additional strategy to examine the ability of rMIC1 and rMIC4 to bind to TLR2 N-glycans. In this approach, HEK cells transfected with the fully N-glycosylated TLR2 ectodomain or with the TLR2 glycomutants [25] were stimulated with a control agonist (FSL-1) or with rMIC1 or rMIC4. HEK cells transfected with any TLR2 form, except those expressing totally unglycosylated TLR2 (mutant Δ1,2,3,4), were able to respond to FSL-1 (Fig 4D), a finding that is consistent with the previous report that the Δ1,2,3,4 mutant is not secreted by HEK293T cells [25]. Cells transfected with TLR2 lacking only the first or the third N-glycan (mutant Δ1; Δ3) responded to all stimuli. The response to the rMIC1 stimulus was significantly reduced in cells transfected with five different TLR2 mutants, lacking some combination of the second, third, and fourth N-glycans (Fig 4D). Moreover, rMIC4 stimulated IL-8 production was significantly reduced in cells transfected with the mutants lacking some combination of the third and fourth N-glycans (Fig 4D).

These results indicate that *T. gondii* MIC1 and MIC4 use their CRDs to induce TLR2- and TLR4-mediated cell activation. Among the TLR2 N-glycans, the rMIC1 CRD likely targets the second, third, and fourth glycan, whereas the rMIC4 CRD targets only the third and fourth. Additionally, our findings suggested that TLR2 and TLR4 activation is required to enhance the production of IL-12 by APCs following rMIC stimulation.

The IL-12 production during *T. gondii in vitro* infection depends partially on MIC1 and MIC4 proteins and their ability to recognize carbohydrates on APCs surface.

Because IL-12 production is induced by rMICs that engage TLR2 and TLR4 N-glycans expressed on innate immune cells, we investigated whether such production is impaired when APCs are infected with *T. gondii* lacking MIC1 and/or MIC4 proteins, as well as complemented strains expressing mutant versions of these proteins that fail to bind TLR2 or TLR4 carbohydrates. We generated Δ*mic1* and Δ*mic4* strains in an RH

strain expressing GFP and Luciferase using CRISPR/Cas9 to replace the endogenous

MIC gene with the drug-selectable marker HPT (HXGPRT – hypoxanthine-xanthine-

guanine phosphoribosyl transferase) (Fig 5A and 5B). We then complemented MIC

deficient parasites with mutated versions expressing an HA-tag, thus generating the

Δ*mic1*::MIC1-T126A/T220A[HA] or Δ*mic4*::MIC4-K469M[HA] strains (Fig 5A) that

expressed endogenous levels of MIC1 and MIC4 as confirmed by Western Blotting (Fig

5C).

IL-12 secretion by BMDCs and BMDMs infected with WT, Δ*mic1*,

Δ*mic1*::MIC1-T126A/T220A, Δ*mic4* and Δ*mic4*::K469M parasites was assessed at 24

hours post infection. All mutant strains (Δ*mic1*, Δ*mic1*::MIC1-T126A/T220A, Δ*mic4*

and Δ*mic4*::K469M) induced lower IL-12 secretion by BMDCs (Fig 5D) and BMDMs

(Fig 5E) compared to that induced by WT parasites, indicating that engagement of

TLR2 and TLR4 cell surface receptors by the MIC lectin-specific activity led to an early

release of IL-12.

Using flow cytometry, we confirmed that parasites deficient in MIC1or MIC4,

or mutated in their carbohydrate recognition domain resulted in lower intracellular IL-

12 production than WT infected BMDCs (Fig 5F-5H). Interestingly, the Toxo[+] BMDCs

presented the same level of intracellular IL-12, independent of the *T. gondii* strain

infected (Fig 5F and 5H). Whereas the Toxo[-] BMDCs produced less IL-12 when they

were infected with knockout or CRD-mutated *T. gondii* compared to WT-infected cells

(Fig 5G and 5H). Taken altogether, these results indicate that MIC1 and MIC4 induce

IL-12 production in innate immune cells during *in vitro T. gondii* infection. It is known

that other parasite factors act as IL-12 inducers, such as profilin, which is a TLR11 and

TLR12 agonist [29, 31], or GRA7 [32], GRA15 [33], and GRA24 [34], which directly

trigger intracellular signalling pathways in a TLR-independent manner, and these likely

account for the majority of IL-12 released after 24 hours of intracellular infection.

MIC1, but not MIC4, contributes to the cytokine storm and acute death during *in*

***vivo* murine infection with *T. gondii*.**

Given the importance of MIC1 and MIC4 as lectins that engage TLR2 and TLR4

N-glycans to induce increased levels of IL-12 release during *T. gondii in vitro* infection,

we investigated the biological relevance of these proteins during *in vivo* infection. Mice

were injected with 50 tachyzoites of RH WT, Δ*mic1*, Δ*mic1*::MIC1-T126A/T220A,

Δ*mic4* and Δ*mic4*::MIC4-K469M strains into the peritoneum of CD-1 outbred mice, a

lethal dose that causes acute mortality. The survival curve showed that parasites

deficient in MIC1 (Δ*mic1* group) or mutated to remove MIC1 lectin binding activity

(Δ*mic1*::MIC1-T126A/T220A group) were less virulent, resulting in a slight, but

significant (p=0.0017) increase in mouse survival (12 days post-infection) compared to

WT infected mice that all succumbed to infection by day 10 (Fig 6A). This was not the

result of a difference in parasite load, which was equivalent across all *T. gondii*-infected

mice at Day 5 (Fig 6D and 6I). Whereas, the absence of the MIC4 gene or MIC4 lectin

activity did not change the survival curve (Fig 6E) indicating that MIC4 is less relevant

than MIC1 during *in vivo* infection.

Acute mortality in CD-1 mice infected with Type I *T. gondii* is related to the

induction of a cytokine storm, mediated by high levels of IFN-γ production. Thus, we

measured systemic levels of IFN-γ and IL-12 in mice infected with WT, Δ*mic1*,

Δ*mic1*::MIC1-T126A/T220A, Δ*mic4* and Δ*mic4*::MIC4-K469M strains. According to

Kugler et al. (2013), the peak of systemic IL-12p40 and IFN-γ during ME49-*T. gondii*

infection is between days 5-6 post-infection, therefore, we measured these cytokines in

the serum of CD-1-infected mice at day 5. Mice infected with Δ*mic1* or Δ*mic1*::MIC1-

T126A/T220A strains had 3-5 fold lower systemic levels of IL-12 (Fig 6B; p=0.016)

and IFN-γ (Fig 6C; p≤0.0002) than WT infected mice. In contrast, mice infected with

parasites lacking the MIC4 gene, or those expressing the mutant version of MIC4

showed no difference in IL-12 (Fig 6F) or IFN-γ (Fig 6G) compared to WT infected

mice. Hence, only MIC1 altered systemic levels of key cytokines induced during *T.*

gondii in vivo infection, and mice survived longer with lower systemic levels of

cytokines typically associated with acute mortality.

MIC1 wild type complemented strain restores the cytokine storm and acute

mortality kinetics during *in vivo* infection with *T. gondii*.

To formally show that MIC1 alters systemic levels of pro-inflammatory

cytokines associated with acute mortality, we complemented Δ*mic1* parasites at the

endogenous locus with a Type I allele of MIC1 expressing an HA tag (MIC1[HA]).

Western blotting for either MIC1 or HA expression showed WT levels of MIC1

expression in the complemented parasites Δ*mic1*::MIC1[HA] (Fig 7A). The complemented

strain restored WT virulence kinetics during *in vivo* infection and all mice died acutely,

in contrast to Δ*mic1* or Δ*mic1*::MIC1-T126A/T220A parasites, that had a slight, but

significant delay in their acute mortality kinetics (Fig 7B; p=0.0082). Systemic levels of

IFN-γ (Fig 7C) and parasite load (Fig 7D and 7E) from mice infected with the

complemented strain were indistinguishable from WT. To better resolve the apparent

difference in acute mortality, parasites were injected into the right footpad to monitor

mouse weight loss and survival kinetics [35]. Mice infected locally in the footpad with

Δ*mic1* survived significantly longer, or did not die (Fig 7G; p=0.0031), and lost less

weight during acute infection (Fig 7F) than those infected with WT or Δ*mic1*::MIC1

complemented parasites. Further, mice infected with Δ*mic1*::MIC1-T126A/T220A

parasites that fail to bind TLR2 and TLR4 N-glycans *in vivo* also lost less weight and

survived significantly longer than WT or Δ*mic1*::MIC1 complemented parasites (Fig 7F

and G). In conclusion, our results suggest that MIC1 operates in two distinct ways; as an

adhesin protein that promotes parasite infection competency, and as a lectin that

engages TLR N-glycans to induce a stronger proinflammatory immune response, one

that is unregulated and results in acute mortality upon RH infection of CD-1 mice.

DISCUSSION

In this study, we report a new function for MIC1 and MIC4, two *T. gondii*

microneme proteins involved in the host-parasite relationship. We show that rMIC1 and

rMIC4, by interacting directly with N-glycans of TLR2 and TLR4, trigger a

noncanonical carbohydrate recognition-dependent activation of innate immune cells.

This results in IL-12 secretion and the production of IFN-γ, a pivotal cytokine that

mediates parasite clearance and the development of a protective T cell response [19,

22], but in some cases, as seen during RH infection of CD-1 mice, promotes a

dysregulated cytokine storm and acute mortality, as seen during RH infection of CD-1

mice [36]. This MIC-TLR activation event explains, at least in part, the resistance

conferred by rMIC1 and rMIC4 administration against experimental toxoplasmosis [20,

21].

T. gondii tachyzoites express microneme proteins either on their surface or

secrete them in their soluble form. These proteins may form complexes, such as those of

MIC1, MIC4, and MIC6 (MIC1/4/6), in which MIC6 is a transmembrane protein that

anchors the two soluble molecules MIC1 and MIC4 [8]. Genetic disruption of each one

of these three genes does not interfere with parasite survival [8] nor its interaction with,

and attachment to, host cells [10]; however, MIC1 has been shown to play a role in

invasion and contributes to virulence in mice [10]. We previously isolated soluble

MIC1/4, a lactose-binding complex from soluble *T. gondii* antigens (STAg) [17], and its

lectin activity was confirmed by the ability of MIC1 to bind sialic acid [9] and MIC4 to

β-galactose [18]. We also reported that MIC1/4 stimulates adherent splenic murine cells

to produce IL-12 at levels as high as those induced by STAg [20]. Recently, it was also

demonstrated that MIC1, MIC4 and MIC6 are capable of inducing IFN-γ production

from memory T cells in mice chronically infected with *T. gondii* [37]. Our data herein

shows that MIC1/4 binds to and activates TLRs via a novel lectin-carbohydrate

interaction, rather than by its cognate receptor-ligand binding groove, establishing

precisely how the interactions of microneme protein(s) with defined glycosylated

receptor(s) expressed on the host cell surface are capable of altering innate priming of

the immune system.

To formally demonstrate the MIC1/MIC4 binding to glycosylated TLR cell

surface receptors we generated recombinant forms of MIC1 and MIC4, which retained

their specific sialic acid- and β-galactose-binding properties as indicated by the results

of their binding to fetuin and asialofetuin as well as the glycoarray assay. Both

recombinant MIC1 and MIC4 triggered the production of proinflammatory and anti-

inflammatory cytokines in DCs and macrophages via their specific recognition of TLR2

and TLR4 N-glycans, as well as by signaling through MyD88 and, partially, TRIF.

Importantly, our results establish how binding of rMIC1 and rMIC4 to specific N-

glycans present on TLR2 and TLR4 induces cell activation through this novel lectin-

carbohydrate interaction. The ligands for MIC1 and MIC4, α2-3-sialyllactosamine and

β1-3- or β1-4-galactosamine, respectively, are terminal N-glycan residues found on a

wide-spectrum of mammalian cell surface-associated glycoconjugates. Thus, it is

possible that additional lectin-carbohydrate interactions may exist between MIC1/4 and

other cell surface receptors beyond TLR2 and TLR4. Such interactions likely evolved to

facilitate adhesion and promote the infection competency of a wide-variety of host cells

infected by *T. gondii*, further underscoring how these proteins exist as important

virulence factors [10] beyond immune priming. However, it is the immunostimulatory

capacity of rMIC1 and rMIC4 to target N-glycans on the ectodomains of TLR2 and

TLR4 that likely rationalizes how these microneme proteins function as a double-edged

sword during *T. gondii* infection. Mice infected by Type I strains die acutely due to a

failure to regulate the cytokine storm induced by high levels of IL-12 and IFN-γ [38,

39]. In this study, *T. gondii* Type I strains engineered to be deficient in MIC1 or

defective in binding TLR2/4 N-glycans lost less weight, survived significantly longer,

and produced less IL-12 and IFN-γ. Future studies that test whether the

immunostimulatory effect of MIC1/4 alters the pathogenesis and cyst burden of Type II

strains of *T. gondii* should be pursued to formally demonstrate that Type II parasites

rely on MIC1/4 induction of Th1-biased cytokines in order to limit tachyzoite

proliferation and induce a life-long persistent bradyzoite infection.

Several pathogens are known to synthesize lectins, which are most frequently

reported to interact with glycoconjugates on host cells to promote adherence, invasion,

and colonization of tissues [40-43]. Nonetheless, there are currently only a few

examples of lectins from pathogens that recognize sugar moieties present in TLRs and

induce IL-12 production by innate immune cells. Paracoccin, a GlcNAc-binding lectin

from the human pathogen *Paracoccidioides brasiliensis,* induces macrophage

polarization towards the M1 phenotype [44] and the production of inflammatory

cytokines through its interaction with TLR2 N-glycans [45]. Furthermore, the galactose-

adherence lectin from *Entamoeba histolytica* activates TLR2 and induces IL-12

production [46]. In addition, the mammalian soluble lectin SP-A, found in lung alveoli,

interacts with the TLR2 ectodomain [47]. The occurrence of cell activation and IL-12

production as a consequence of the recognition of TLR N-glycans has also been

demonstrated using plant lectins with different sugar-binding specificities [48, 49].

The binding of MIC1 and MIC4, as well as the lectins above, to TLR2 and

TLR4 may be associated with the position of the specific sugar residue present on the

receptor's N-glycan structure. Since the N-glycan structures of TLR2 and TLR4 are still

unknown, we assume that the targeted MIC1 and MIC4 residues, e.g. sialic acid α2-3-

linked to galactose β1-3- and β1-4-galactosamines, are appropriately placed in the

receptors' oligosaccharides to allow the recognition phenomenon and trigger the

activation of innate immune responses.

Several *T. gondii* proteins have previously been shown to activate innate

immune cells in a TLR-dependent manner, but independent of sugar recognition. This is

the case for profilin (TgPRF), which is essential for the parasite's gliding motility based

on actin polymerization; it is recognized by TLR11 [29] and TLR12 [31, 50]. In

addition, *T. gondii*-derived glycosylphosphatidylinositol anchors activate TLR2 and

TLR4 [51], and parasite RNA and DNA are ligands for TLR7 and TLR9, respectively

[19, 22, 50]. The stimulation of all of these TLRs culminate in MyD88 activation which

results in IL-12 production [19, 22]. Several other *T. gondii* secreted effector proteins

regulate the production of proinflammatory cytokines such as IL-12, independent of

TLRs. For example, the dense granule protein 7 (GRA7) induces MyD88-dependent

NF-kB activation, which facilitates IL-12, TNF-α, and IL-6 production [32]. MIC3 is

reported to induce TNF-α secretion and macrophage M1 polarization [52], whereas

GRA15 expressed by Type II strains activates NF-kB, promoting the release of IL-12

[33], and GRA24 triggers the autophosphorylation of p38 MAP kinase and

proinflammatory cytokine and chemokine secretion [34]. In contrast, TgIST interferes

with IFN-γ induction by actively inhibiting STAT1-dependent proinflammatory gene

expression indicating that the parasite is capable of both activating as well as inhibiting

effector arms of the host immune response to impact its pathogenesis *in vivo* [53]. Thus,

multiple secretory effector proteins of *T. gondii*, including MIC1 and MIC4, appear to

work in tandem to ultimately promote protective immunity by either inducing or

dampening the production of proinflammatory cytokines, the timing of which is central

to controlling both the parasite's proliferation during the acute phase of infection and

the induction of an effective immune response capable of establishing a chronic

infection [19].

Our results regarding soluble MIC1 and MIC4 confirmed our hypothesis that

these two effector proteins induce the innate immune response against *T. gondii* through

TLR2- and TLR4-dependent pathways. This is consistent with previous studies that

highlight the importance of TLR signaling, as well as the MyD88 adapter molecule, as

essential for conferring resistance to *T. gondii* infection [29, 51, 54, 55]. In addition, we

show that both MIC1 and MIC4 on the parasite surface contribute to the secretion of IL-

12 by macrophages and DCs during *in vitro* infection, but only MIC1 plays a significant

role during *in vivo* infection, demonstrated by its ability to promote a dysregulated

induction of systemic levels of IFN-γ and a proinflammatory cytokine storm that leads

to acute mortality during murine infection.

METHODS

Ethics statement

All experiments were conducted in accordance to the Brazilian Federal Law

11,794/2008 establishing procedures for the scientific use of animals, and State Law

establishing the Animal Protection Code of the State of Sao Paulo. All efforts were

made to minimize suffering, and the animal experiments were approved by the Ethics

Committee on Animal Experimentation (*Comissão de Ética em Experimentação Animal*

- CETEA) of the Ribeirao Preto Medical School, University of Sao Paulo (protocol

number 065/2012), following the guidelines of the National Council for Control of

Animal Experimentation (*Conselho Nacional de Controle de Experimentação Animal* -

CONCEA).

Lac+ fraction and recombinant MIC1 and MIC4

The lactose-eluted (Lac+) fraction was obtained as previously reported [17, 21].

Briefly, the total soluble tachyzoite antigen (STAg) fraction was loaded into a lactose

column (Sigma-Aldrich, St. Louis, MO) and equilibrated with PBS containing 0.5 M

NaCl. The material adsorbed to the resin was eluted with 0.1 M lactose in equilibrating

buffer and dialyzed against ultrapure water. The obtained fraction was denoted as Lac+

and confirmed to contain MIC1 and MIC4. For the recombinant proteins, rMIC1 and

rMIC4 sequences were amplified from cDNA of the *T. gondii* strain ME49 with a 6-

histidine tag added on the N-terminal, cloned into pDEST17 vector (Gateway Cloning,

Thermo Fisher Scientific Inc., Grand Island, NY), and used to transform DH5α *E. coli*

chemically competent cells for ampicillin expression selection, as described before [21].

The plasmids with rMIC1-T126A/T220A and rMIC4-K469M were synthesized by

GenScript (New Jersey, US) using a pET28a vector, and the MIC sequences carrying

the mutations were cloned between the *Nde*I and *Bam*H I sites. All plasmids extracted

from DH5α *E. coli* were transformed in *E. coli* BL21-DE3 chemically competent cells

to produce recombinant proteins that were then purified from inclusion bodies and

refolded by gradient dialysis, as described previously for rMIC1 and rMIC4 wild type

forms [21]. Endotoxin concentrations were measured in all protein samples using the

Limulus Amebocyte Lysate Kit – QCL-1000 (Lonza, Basel, Switzerland). The rMIC1,

rMIC1-T126A/T220A, rMIC4 and rMIC4-K469M contained 7.2, 3.2, 3.5 and 1.1 EU

endotoxin/μg of protein, respectively. Endotoxin was removed by passing over two

polymyxin-B columns (Affi-Prep Polymyxin Resin; Bio-Rad, Hercules, CA).

Additionally, prior to all *in vitro* cell-stimulation assays, the proteins samples were

incubated with 50 μg/mL of polymyxin B sulphate salt (Sigma-Aldrich, St. Louis, MO)

for 30 min at 37 °C to remove possible residual LPS.

Glycan array

The carbohydrate-binding profile of microneme proteins was determined by Core H (Consortium for Functional Glycomics, Emory University, Atlanta, GA), using a printed glycan microarray, as described previously [56]. Briefly, rMIC1-Fc, rMIC4-Fc, and Lac$^+$-Fc in binding buffer (1% BSA, 150 mM NaCl, 2 mM CaCl$_2$, 2 mM MgCl$_2$, 0.05% (w/v) Tween 20, and 20 mM Tris-HCl, pH 7.4) were applied onto a covalently printed glycan array and incubated for 1 hour at 25 °C, followed by incubation with Alexa Fluor 488-conjugate (Invitrogen, Thermo Fisher Scientific Inc., Grand Island, NY). Slides were scanned, and the average signal intensity was calculated. The common features of glycans with stronger binding are depicted in Fig. 1a. The average signal intensity detected for all of the glycans was calculated and set as the baseline.

Sugar-inhibition assay

Ninety-six-well microplates were coated with 1 μg/well of fetuin or asialofetuin, glycoproteins diluted in 50 μL of carbonate buffer (pH 9.6) per well, followed by overnight incubation at 4 °C. Recombinant MIC1 or MIC4 proteins (both wild type (WT) and mutated forms), previously incubated or not with their corresponding sugars, i.e. α(2-3)-sialyllactose for MIC1 and lacto-N-biose for MIC4 (V-lab, Dextra, LA, UK), were added into coated wells and incubated for 2 h at 25 °C. After washing with PBS, *T. gondii*-infected mouse serum (1:50) was used as the source of the primary antibody. The assay was then developed with anti-mouse peroxidase-conjugated secondary antibody, and the absorbance was measured at 450 nm in a microplate-scanning spectrophotometer (Power Wave-X; BioTek Instruments, Inc., Winooski, VT).

Mice and parasites

Female C57BL/6 (WT), MyD88$^{-/-}$, TRIF$^{-/-}$, TLR2$^{-/-}$, TLR3$^{-/-}$, TLR4$^{-/-}$, double knockout (DKO) TLR2$^{-/-}$/TLR4$^{-/-}$, TLR5$^{-/-}$, and TLR9$^{-/-}$ mice (all from the C57BL/6 background), 8 to 12 weeks of age, were acquired from the University of São Paulo - Ribeirão Preto campus animal facility, Ribeirão Preto, São Paulo, Brazil, and housed in the animal facility of the Department of Cell and Molecular Biology - Ribeirão Preto Medical School, under specific pathogen-free conditions. The TLR11$^{-/-}$/TLR12$^{-/-}$ DKO mice were maintained at American Association of Laboratory Animal Care-accredited animal facilities at NIAID/NIH. For the *in vivo* infections, female CD-1 outbred mice, 6 weeks of age were acquired from Charles River Laboratories, Germantown, MD, USA.

A clonal isolate of the *T. gondii* RH-Δ*ku80*/Δ*hpt* strain was used to generate the

GFP/Luciferase strain, which was the recipient strain to generate the single-knockout

parasites. The GFP/Luc sequence was inserted into the UPRT locus of *Toxoplasma* by

double crossover homologous recombination using CRISPR/Cas-based genome editing

and selected for FUDR resistance to facilitate the targeted GFP/Luc gene cassette

knock-in. The MIC1 and MIC4 genes were replaced by the drug-selectable marker *hpt*

(*hxgprt* - hypoxanthine-xanthine-guanine phosphoribosyl transferase) flanked by LoxP

sites. For all gene deletions, 30 μg of guide RNA was transfected along with 15 μg of a

repair oligo. Parasites were transfected and selected as previously described [57, 58].

For the MIC gene complementation, the sequence was amplified from RH genomic

DNA with the addition of one copy of HA-tag sequence

(TACCCATACGATGTTCCAGATTACGCT) before the stop codon, and cloned into

pCR2.1-TOPO vector, followed by site-directed mutagenesis using the Q-5 kit (New

England Biolabs) in order to generate point mutations into MIC1 (MIC1-

T126A/T220A) and MIC4 (MIC4-K469M) sequences. For transfections, 30 μg of guide

RNA was transfected along with 20 μg of linearized pTOPO vector containing the MIC

mutated sequences.

Strains were maintained in human foreskin fibroblast (HFF) cells grown in

Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% heat-

inactivated foetal bovine serum (FBS), 0.25 mM gentamicin, 10 U/mL penicillin, and

10 μg/mL streptomycin (Gibco, Thermo Fisher Scientific Inc., Grand Island, NY).

Bone marrow-derived dendritic cells and macrophages

Bone marrows of WT, MyD88[-/-], TRIF[-/-], TLR2[-/-], TLR3[-/-], TLR4[-/-], DKO

TLR2[-/-]/TLR4[-/-], TLR5[-/-], TLR9[-/-], and DKO TLR11[-/-]/TLR12[-/-] mice were harvested

from femurs and hind leg bones. Cells were washed with RPMI medium and

resuspended in RPMI medium with 10% FBS, 10 U/mL penicillin, and 10 μg/mL

streptomycin (Gibco). For dendritic cell (DC) differentiation, we added 10 ng/mL of

recombinant murine GM-CSF (Prepotech, Rocky Hill, NJ), and 10 ng/mL murine

recombinant IL-4 (eBioscience, San Diego, CA); for macrophage differentiation, 30%

of L929 conditioned medium was added to RPMI medium with 10% FBS. The cells

were cultured in 100 × 20 mm dish plates (Costar; Corning Inc., Corning, NY),

supplemented with respective conditioned media at days 3 and 6 for DCs, and at day 4

for macrophages. DCs were incubated for 8–9 days and macrophages for 7 days; the

cells were then harvested and plated into 24-well plates at 5×10^5 cells/well for protein

stimulations or *T. gondii* infections, followed by ELISA. Cell purity was analyzed by

flow cytometry. Eighty-five percent of differentiated dentritic cells were

CD11b$^+$/CD11c$^+$, while 94% of differentiated macrophages were CD11b$^+$.

HEK293T cells transfection

Human embryonic kidney 293T (HEK293T) cells, originally acquired from

American Tissue Culture Collection (ATCC, Rockville, MD), were used as an

expression tool [59] for TLR2 and TLR4 [45, 60]. The cells grown in DMEM

supplemented with 10% FBS (Gibco), and were seeded at 3.5×10^5 cells/mL in 96-well

plates (3.5×10^4 cells/well) 24 h before transfection. Then, HEK293T cells were

transiently transfected (70-80% confluence) with human TLR2 plasmids as described

previously [25] or with CD14, CD36, MD-2 and TLR4 [61] using Lipofectamine 2000

(Invitrogen) with 60 ng of NF-κB Luc, an NF-κB reporter plasmid, and 0.5 ng of

Renilla luciferase plasmid, together with 60 ng of each gene of single and multiple

glycosylation mutants and of TLR2 WT genes [25]. After 24 h of transfection, the cells

were stimulated overnight with positive controls: P3C (Pam3CSK4; EMC

Microcollections, Tübingen, Germany), fibroblast stimulating ligand-1 (FSL-1; EMC

Microcollections), or LPS Ultrapure (standard LPS, *E. coli* 0111:B4; Sigma-Aldrich); or

with the negative control for cell stimulation (the medium). Cells transfected with

empty vectors, incubated either with the medium or with agonists (FSL-1 or P3C), were

also assayed; negative results were required for each system included in the study. IL-8

was detected in the culture supernatants. The absence of Mycoplasma contamination in

the cell culture was certified by indirect fluorescence staining as described previously

[62].

Cytokine measurement

The quantification of human IL-8 and mouse IL-12p40, IL-6, TNF-α, and IL-10

in the supernatant of the cultures was performed by ELISA, following the

manufacturer's instructions (OptEIA set; BD Biosciences, San Jose, CA). Human and

murine recombinant cytokines were used to generate standard curves and determine

cytokine concentrations. The absorbance was read at 450 nm using the Power Wave-X

spectrophotometer (BioTek Instruments).

TLR2-FLAG and TLR4-FLAG plasmids

The pcDNA4/TO-FLAG plasmid was kindly provided by Dr. Dario Simões

Zamboni. The pcDNA4-FLAG-TLR2 and pcDNA4-FLAG-TLR4 plasmids   were

constructed as follows. RNA from a P388D1 cell line (ATCC, Rockville, MD) was

extracted and converted to cDNA   with Maxima H Minus Reverse Transcriptase

(Thermo-Fisher Scientific, Waltham, MA USA)   and oligo(dT). TLR2 and TLR4 were

amplified from total cDNA from murine macrophages by using   Phusion High-Fidelity

DNA Polymerase and the phosphorylated primers TLR2_F:

ATGCTACGAGCTCTTTGGCTCTTCTGG, TLR2_R:

CTAGGACTTTATTGCAGTTCTCAGATTTACCCAAAAC, TLR4_F:

TGCTTAGGATCCATGATGCCTCCCTGGCTCCTG and TLR4_R:

TGCTTAGCGGCCGCTCAGGTCCAAGTTGCCGTTTCTTG. The fragments were

isolated from 1% agarose/Tris-acetate-ethylenediaminetetraacetic acid gel, purified

with   GeneJET Gel Extraction Kit (Thermo-Fisher Scientific), and inserted into the

pcDNA4/TO-FLAG vector by using the restriction enzymes sites for NotI and XbaI

(Thermo-Fisher Scientific) for TLR2, and BamHI and NotI (Thermo-Fisher Scientific)

for TLR4. Ligation reactions were performed by using a 3:1 insert/vector   ratio with T4

DNA Ligase (Thermo-Fisher Scientific) and transformed into chemically   competent

Escherichia coli DH5α cells. Proper transformants were isolated from LB agar

medium plates under ampicillin selection (100 μg/mL) and analyzed by PCR,

restriction fragment   analysis, and DNA sequencing. All reactions were performed

according to the manufacturer's   instructions.

Pull-down assay and Western Blot

We used the lysate of HEK293T cells transfected (70-80% confluence) with

plasmids containing TLR2-FLAG or TLR4-FLAG. After 24 h of transfection, the HEK

cells were lysed with a non-denaturing lysis buffer (20 mM Tris, pH 8.0, 137 mM NaCl,

and 2 mM EDTA) supplemented with a protease inhibitor (Roche, Basel, Switzerland).

After 10 min of incubation on ice, the lysate was subjected to centrifugation (16,000 *g*,

at 4 °C for 5 min). The protein content in the supernatant was quantified by the BCA

method, aliquoted, and stored at -80 °C. For the pull-down assay, 100 μg of the lysate

from TLR2-FLAG- or TLR4-FLAG-transfected HEK cells were incubated with 10 μg

of TgMIC1 or TgMIC4 overnight at 4 °C. Since these proteins had a histidine tag, the

samples were purified on nickel-affinity resin (Ni Sepharose High Performance; GE

Healthcare, Little Chalfont, UK) after incubation for 30 min at 25 °C and centrifugation

of the fraction bound to nickel to pull down the TgMIC-His that physically interacted

with TLR-FLAG (16,000 g, 4 °C, 5 min). After washing with PBS, the samples were

resuspended in 100 μL of SDS loading dye with 5 μL of 2-mercaptoethanol, heated for

5 min at 95 ºC, and 25 μL of total volume was run on 10% SDS-PAGE. After

transferring to a nitrocellulose membrane (Millipore, Billerica, MA), immunoblotting

was performed by following the manufacturer's protocol. First, the membrane was

incubated with anti-FLAG monoclonal antibodies (1:2,000) (Clone G10, ab45766,

Sigma-Aldrich) to detect the presence of TLR2 or TLR4. The same membrane was then

subjected to secondary probing and was developed with anti-TgMIC1 (IgY; 1:20,000)

or anti-TgMIC4 (IgY; 1:8,000) polyclonal antibodies and followed by incubation with

secondary polyclonal anti-chicken IgY-HRP (1:4,000) (A9046, Sigma-Aldrich) to

confirm the presence of TgMIC1 and TgMIC4.

***In vitro* infections**

Bone marrow-derived dendritic cells (BMDCs) and bone marrow-derived

macrophages (BMDMs) were infected with WT (Δ*ku80*/Δ*hpt*), Δ*mic1*, Δ*mic1*::MIC1-

T126A/T220A, Δ*mic4* or Δ*mic4*::MIC4-K469M (Type I, RH background) strains

recovered from T25 flasks with HFF cell cultures. The T25 flasks were washed with

RPMI medium to completely remove parasites, and the collected material was

centrifuged for 5 min at 50 g to remove HFF cell debris. The resulting pellet was

discarded, and the supernatant containing the parasites was centrifuged for 10 min at

1,000 g and resuspended in RPMI medium for counting and concentration adjustments.

BMDCs and BMDMs were dispensed in 24-well plates at 5×10^5 cells/well (in RPMI

medium supplemented with 10% FBS), followed by infection with 3 parasites per cell

(multiplicity of infection, MOI 3). Then, the plate was centrifuged for 3 min at 200 g to

synchronize the contact between cells and parasites and incubated at 37 °C. The

supernatants were collected at 6, 12, 24, and 48 h after infection for quantification of

IL-12p40.

***In vivo* infections and Luciferase assay**

Six-week-old female CD-1 outbred mice were infected by intraperitoneal

injection with 50 tachyzoites of RH engineered strains diluted in 500 μl of phosphate-

buffered saline. The mice were weighed daily and survival was evaluated

Bioluminescent detection of firefly luciferase activity was performed at day 5

post-infection using an IVIS BLI system from Xenogen to monitor parasite burden.

Mice were injected with 3 milligrams (200 µl) of D-luciferin (PerkinElmer) substrate,

and after 5 minutes the mice were imaged for 300 seconds to detect the photons emitted.

Statistical analysis

The data were plotted and analysed using GraphPad Prism 7.0 software

(GraphPad, La Jolla, CA). Statistical significance of the obtained results was calculated

using analysis of variance (One-way ANOVA) followed by Bonferroni's multiple

comparisons test. Differences were considered significant when the P value was <0.05.

ACKNOWLEDGEMENTS

We are grateful to Dr. Tiago Wilson Patriarca Mineo (Universidade Federal de

Uberlândia - MG) for kindly provided us the wild type *Toxoplasma gondii* (RH); to Dr.

Larissa Dias Cunha and Dr. Dario Simões Zamboni (Universidade de São Paulo - SP)

for help with double knockout TLR2/TLR4 mice generation and for kindly provide the

pcDNA4/TO-FLAG plasmid; to Izaíra Tincani Brandão and Ana Paula Masson for

technical assistance with endotoxin measurements; to Patricia Vendrusculo, Sandra

Thomaz and Sara Hieny for all technical support essential for this study. We wish to

acknowledge the grants from Consortium for Functional Glycomics (#GM62116), for

doing the glycoarray assays; UK Medical Research Council (#G1000133 to N.J.G.), and

Wellcome Investigator Award (#WT100321/z/12/Z to N.J.G.). This work was

supported in part by the National Institute of Allergy and Infectious Diseases, National

Institutes of Health (A.S.S. and M.E.G.).

REFERENCES

1.   Dubey JP. Toxoplasmosis of animals and humans. 2nd ed. Boca Raton, FL,

USA: CRC Press; 2009. 336 p.

2.   Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, et al.

Foodborne illness acquired in the United States--major pathogens. Emerging infectious

diseases. 2011;17(1):7-15. Epub 2011/01/05. doi: 10.3201/eid1701.P11101

10.3201/eid1701.091101p1. PubMed PMID: 21192848; PubMed Central PMCID:

PMC3375761.

3.   Carruthers VB, Sibley LD. Sequential protein secretion from three distinct

organelles of Toxoplasma gondii accompanies invasion of human fibroblasts. European

journal of cell biology. 1997;73(2):114-23. Epub 1997/06/01. PubMed PMID: 9208224.

4.   Carruthers VB, Giddings OK, Sibley LD. Secretion of micronemal proteins is

associated with toxoplasma invasion of host cells. Cellular microbiology.

1999;1(3):225-35. Epub 2001/02/24. PubMed PMID: 11207555.

5.   Lovett JL, Sibley LD. Intracellular calcium stores in Toxoplasma gondii govern

invasion of host cells. Journal of cell science. 2003;116(Pt 14):3009-16. Epub

2003/06/05. doi: 10.1242/jcs.00596. PubMed PMID: 12783987.

6.   Tomley FM, Soldati DS. Mix and match modules: structure and function of

microneme proteins in apicomplexan parasites. Trends in parasitology. 2001;17(2):81-8.

Epub 2001/03/03. PubMed PMID: 11228014.

7.   Carruthers VB, Tomley FM. Microneme proteins in apicomplexans. Sub-cellular

biochemistry. 2008;47:33-45. Epub 2008/06/03. PubMed PMID: 18512339; PubMed

Central PMCID: PMC2847500.

8.   Reiss M, Viebig N, Brecht S, Fourmaux MN, Soete M, Di Cristina M, et al.

Identification and characterization of an escorter for two secretory adhesins in

Toxoplasma gondii. The Journal of cell biology. 2001;152(3):563-78. Epub 2001/02/07.

PubMed PMID: 11157983; PubMed Central PMCID: PMC2196004.

9.   Friedrich N, Santos JM, Liu Y, Palma AS, Leon E, Saouros S, et al. Members of

a novel protein family containing microneme adhesive repeat domains act as sialic acid-

binding lectins during host cell invasion by apicomplexan parasites. The Journal of

biological chemistry. 2010;285(3):2064-76. Epub 2009/11/11. doi:

10.1074/jbc.M109.060988. PubMed PMID: 19901027; PubMed Central PMCID:

PMC2804363.

10.   Cerede O, Dubremetz JF, Soete M, Deslee D, Vial H, Bout D, et al. Synergistic

role of micronemal proteins in Toxoplasma gondii virulence. The Journal of

experimental medicine. 2005;201(3):453-63. Epub 2005/02/03. doi:

10.1084/jem.20041672. PubMed PMID: 15684324; PubMed Central PMCID:

PMC2213027.

11.   Blumenschein TM, Friedrich N, Childs RA, Saouros S, Carpenter EP,

Campanero-Rhodes MA, et al. Atomic resolution insight into host cell recognition by

Toxoplasma gondii. The EMBO journal. 2007;26(11):2808-20. Epub 2007/05/12. doi:

10.1038/sj.emboj.7601704. PubMed PMID: 17491595; PubMed Central PMCID:

PMC1888667.

12.   Monteiro VG, Soares CP, de Souza W. Host cell surface sialic acid residues are

involved on the process of penetration of Toxoplasma gondii into mammalian cells.

FEMS microbiology letters. 1998;164(2):323-7. Epub 1998/07/31. PubMed PMID:

9682481.

13.   Ortega-Barria E, Boothroyd JC. A Toxoplasma lectin-like activity specific for

sulfated polysaccharides is involved in host cell infection. The Journal of biological

chemistry. 1999;274(3):1267-76. Epub 1999/01/09. PubMed PMID: 9880495.

14.   Carruthers VB, Hakansson S, Giddings OK, Sibley LD. Toxoplasma gondii uses

sulfated proteoglycans for substrate and host cell attachment. Infection and immunity.

2000;68(7):4005-11. Epub 2000/06/17. PubMed PMID: 10858215; PubMed Central

PMCID: PMC101681.

15.   Paing MM, Tolia NH. Multimeric assembly of host-pathogen adhesion

complexes involved in apicomplexan invasion. PLoS pathogens. 2014;10(6):e1004120.

Epub 2014/06/20. doi: 10.1371/journal.ppat.1004120. PubMed PMID: 24945143;

PubMed Central PMCID: PMC4055764.

16.   Brecht S, Carruthers VB, Ferguson DJ, Giddings OK, Wang G, Jakle U, et al.

The toxoplasma micronemal protein MIC4 is an adhesin composed of six conserved

apple domains. The Journal of biological chemistry. 2001;276(6):4119-27. Epub

2000/10/29. doi: 10.1074/jbc.M008294200. PubMed PMID: 11053441.

17.   Lourenco EV, Pereira SR, Faca VM, Coelho-Castelo AA, Mineo JR, Roque-

Barreira MC, et al. Toxoplasma gondii micronemal protein MIC1 is a lactose-binding

lectin. Glycobiology. 2001;11(7):541-7. Epub 2001/07/12. PubMed PMID: 11447133.

18.   Marchant J, Cowper B, Liu Y, Lai L, Pinzan C, Marq JB, et al. Galactose

recognition by the apicomplexan parasite Toxoplasma gondii. The Journal of biological

chemistry. 2012;287(20):16720-33. Epub 2012/03/09. doi: 10.1074/jbc.M111.325928.

PubMed PMID: 22399295; PubMed Central PMCID: PMC3351351.

19.   Yarovinsky F. Toll-like receptors and their role in host resistance to Toxoplasma

gondii. Immunology letters. 2008;119(1-2):17-21. Epub 2008/07/12. doi:

10.1016/j.imlet.2008.05.007. PubMed PMID: 18617274.

20.   Lourenco EV, Bernardes ES, Silva NM, Mineo JR, Panunto-Castelo A, Roque-

Barreira MC. Immunization with MIC1 and MIC4 induces protective immunity against

Toxoplasma gondii. Microbes and infection / Institut Pasteur. 2006;8(5):1244-51. Epub

2006/04/18. doi: 10.1016/j.micinf.2005.11.013. PubMed PMID: 16616574.

21.   Pinzan CF, Sardinha-Silva A, Almeida F, Lai L, Lopes CD, Lourenco EV, et al.

Vaccination with Recombinant Microneme Proteins Confers Protection against

Experimental Toxoplasmosis in Mice. PloS one. 2015;10(11):e0143087. Epub

2015/11/18. doi: 10.1371/journal.pone.0143087. PubMed PMID: 26575028; PubMed

Central PMCID: PMC4648487.

22.   Yarovinsky F. Innate immunity to Toxoplasma gondii infection. Nature reviews

Immunology. 2014;14(2):109-21. Epub 2014/01/25. doi: 10.1038/nri3598. PubMed

PMID: 24457485.

23.   Gazzinelli RT, Wysocka M, Hieny S, Scharton-Kersten T, Cheever A, Kuhn R,

et al. In the absence of endogenous IL-10, mice acutely infected with Toxoplasma

gondii succumb to a lethal immune response dependent on CD4+ T cells and

accompanied by overproduction of IL-12, IFN-gamma and TNF-alpha. J Immunol.

1996;157(2):798-805. Epub 1996/07/15. PubMed PMID: 8752931.

24.   Oldenhove G, Bouladoux N, Wohlfert EA, Hall JA, Chou D, Dos Santos L, et al.

Decrease of Foxp3+ Treg cell number and acquisition of effector cell phenotype during

lethal infection. Immunity. 2009;31(5):772-86. doi: 10.1016/j.immuni.2009.10.001.

PubMed PMID: 19896394; PubMed Central PMCID: PMC2814877.

25.   Weber AN, Morse MA, Gay NJ. Four N-linked glycosylation sites in human

toll-like receptor 2 cooperate to direct efficient biosynthesis and secretion. The Journal

of biological chemistry. 2004;279(33):34589-94. Epub 2004/06/03. doi:

10.1074/jbc.M403830200. PubMed PMID: 15173186.

26.   Santos A, Carvalho FC, Roque-Barreira MC, Zorzetto-Fernandes AL, Gimenez-

Romero D, Monzo I, et al. Evidence for Conformational Mechanism on the Binding of

TgMIC4 with beta-Galactose-Containing Carbohydrate Ligand. Langmuir : the ACS

journal of surfaces and colloids. 2015;31(44):12111-9. Epub 2015/10/22. doi:

10.1021/acs.langmuir.5b03141. PubMed PMID: 26488670.

27.   Hager KM, Carruthers VB. MARveling at parasite invasion. Trends in

parasitology. 2008;24(2):51-4. Epub 2008/01/22. doi: 10.1016/j.pt.2007.10.008.

PubMed PMID: 18203663; PubMed Central PMCID: PMC2662992.

28.   Jankovic D, Kullberg MC, Feng CG, Goldszmid RS, Collazo CM, Wilson M, et

al. Conventional T-bet(+)Foxp3(-) Th1 cells are the major source of host-protective

regulatory IL-10 during intracellular protozoan infection. The Journal of experimental

medicine. 2007;204(2):273-83. Epub 2007/02/07. doi: 10.1084/jem.20062175. PubMed

PMID: 17283209; PubMed Central PMCID: PMC2118735.

29.   Yarovinsky F, Zhang D, Andersen JF, Bannenberg GL, Serhan CN, Hayden MS,

et al. TLR11 activation of dendritic cells by a protozoan profilin-like protein. Science.

2005;308(5728):1626-9. Epub 2005/04/30. doi: 10.1126/science.1109893. PubMed

PMID: 15860593.

30.   da Silva Correia J, Ulevitch RJ. MD-2 and TLR4 N-linked glycosylations are

important for a functional lipopolysaccharide receptor. The Journal of biological

chemistry. 2002;277(3):1845-54. Epub 2001/11/14. doi: 10.1074/jbc.M109910200.

PubMed PMID: 11706042.

31.   Koblansky AA, Jankovic D, Oh H, Hieny S, Sungnak W, Mathur R, et al.

Recognition of profilin by Toll-like receptor 12 is critical for host resistance to

Toxoplasma gondii. Immunity. 2013;38(1):119-30. Epub 2012/12/19. doi:

10.1016/j.immuni.2012.09.016. PubMed PMID: 23246311; PubMed Central PMCID:

PMC3601573.

32.   Yang CS, Yuk JM, Lee YH, Jo EK. Toxoplasma gondii GRA7-Induced TRAF6

Activation Contributes to Host Protective Immunity. Infection and immunity.

2015;84(1):339-50. Epub 2015/11/11. doi: 10.1128/IAI.00734-15. PubMed PMID:

26553469; PubMed Central PMCID: PMC4693986.

33.   Rosowski EE, Lu D, Julien L, Rodda L, Gaiser RA, Jensen KD, et al. Strain-

specific activation of the NF-kappaB pathway by GRA15, a novel Toxoplasma gondii

dense granule protein. The Journal of experimental medicine. 2011;208(1):195-212.

Epub 2011/01/05. doi: 10.1084/jem.20100717. PubMed PMID: 21199955; PubMed

Central PMCID: PMC3023140.

34.   Braun L, Brenier-Pinchart MP, Yogavel M, Curt-Varesano A, Curt-Bertini RL,

Hussain T, et al. A Toxoplasma dense granule protein, GRA24, modulates the early

immune response to infection by promoting a direct and sustained host p38 MAPK

activation. The Journal of experimental medicine. 2013;210(10):2071-86. Epub

2013/09/18. doi: 10.1084/jem.20130103. PubMed PMID: 24043761; PubMed Central

PMCID: PMC3782045.

35.   Ma JS, Sasai M, Ohshima J, Lee Y, Bando H, Takeda K, et al. Selective and

strain-specific NFAT4 activation by the Toxoplasma gondii polymorphic dense granule

protein GRA6. The Journal of experimental medicine. 2014;211(10):2013-32. Epub

2014/09/17. doi: 10.1084/jem.20131272. PubMed PMID: 25225460; PubMed Central

PMCID: PMC4172224.

36.   Hunter CA, Sibley LD. Modulation of innate immunity by Toxoplasma gondii

virulence effectors. Nature reviews Microbiology. 2012;10(11):766-78. Epub

2012/10/17. doi: 10.1038/nrmicro2858. PubMed PMID: 23070557; PubMed Central

PMCID: PMC3689224.

37.   Saraav I, Wang Q, Brown KM, Sibley LD. Secretory Microneme Proteins

Induce T-Cell Recall Responses in Mice Chronically Infected with Toxoplasma gondii.

mSphere. 2019;4(1). Epub 2019/03/01. doi: 10.1128/mSphere.00711-18. PubMed

PMID: 30814319; PubMed Central PMCID: PMC6393730.

38.   Gavrilescu LC, Denkers EY. IFN-gamma overproduction and high level

apoptosis are associated with high but not low virulence Toxoplasma gondii infection. J

Immunol. 2001;167(2):902-9. Epub 2001/07/07. PubMed PMID: 11441097.

39.   Mordue DG, Monroy F, La Regina M, Dinarello CA, Sibley LD. Acute

toxoplasmosis leads to lethal overproduction of Th1 cytokines. J Immunol.

2001;167(8):4574-84. Epub 2001/10/10. PubMed PMID: 11591786.

40.   Takabatake N, Okamura M, Yokoyama N, Ikehara Y, Akimitsu N, Arimitsu N,

et al. Glycophorin A-knockout mice, which lost sialoglycoproteins from the red blood

cell membrane, are resistant to lethal infection of Babesia rodhaini. Veterinary

parasitology. 2007;148(2):93-101. Epub 2007/07/27. doi: 10.1016/j.vetpar.2007.06.011.

PubMed PMID: 17651898.

41.   Persson KE, McCallum FJ, Reiling L, Lister NA, Stubbs J, Cowman AF, et al.

Variation in use of erythrocyte invasion pathways by Plasmodium falciparum mediates

evasion of human inhibitory antibodies. The Journal of clinical investigation.

2008;118(1):342-51. Epub 2007/12/08. doi: 10.1172/JCI32138. PubMed PMID:

18064303; PubMed Central PMCID: PMC2117763.

42.   Favila MA, Geraci NS, Jayakumar A, Hickerson S, Mostrom J, Turco SJ, et al.

Differential Impact of LPG-and PG-Deficient Leishmania major Mutants on the

Immune Response of Human Dendritic Cells. PLoS neglected tropical diseases.

2015;9(12):e0004238. Epub 2015/12/03. doi: 10.1371/journal.pntd.0004238. PubMed

PMID: 26630499; PubMed Central PMCID: PMC4667916.

43.   Nogueira PM, Assis RR, Torrecilhas AC, Saraiva EM, Pessoa NL, Campos MA,

et al. Lipophosphoglycans from Leishmania amazonensis Strains Display

Immunomodulatory Properties via TLR4 and Do Not Affect Sand Fly Infection. PLoS

neglected tropical diseases. 2016;10(8):e0004848. Epub 2016/08/11. doi:

10.1371/journal.pntd.0004848. PubMed PMID: 27508930; PubMed Central PMCID:

PMC4980043.

44.   Freitas MS, Oliveira AF, da Silva TA, Fernandes FF, Goncales RA, Almeida F,

et al. Paracoccin Induces M1 Polarization of Macrophages via Interaction with TLR4.

Frontiers in microbiology. 2016;7:1003. Epub 2016/07/28. doi:

10.3389/fmicb.2016.01003. PubMed PMID: 27458431; PubMed Central PMCID:

PMC4932198.

45.   Alegre-Maller AC, Mendonca FC, da Silva TA, Oliveira AF, Freitas MS, Hanna

ES, et al. Therapeutic administration of recombinant Paracoccin confers protection

against paracoccidioides brasiliensis infection: involvement of TLRs. PLoS neglected

tropical diseases. 2014;8(12):e3317. Epub 2014/12/05. doi:

10.1371/journal.pntd.0003317. PubMed PMID: 25474158; PubMed Central PMCID:

PMC4256291.

46.   Campbell D, Mann BJ, Chadee K. A subunit vaccine candidate region of the

Entamoeba histolytica galactose-adherence lectin promotes interleukin-12 gene

transcription and protein production in human macrophages. European journal of

immunology. 2000;30(2):423-30. Epub 2000/02/12. doi: 10.1002/1521-

4141(200002)30:2<423::AID-IMMU423>3.0.CO;2-0. PubMed PMID: 10671197.

47.   Murakami S, Iwaki D, Mitsuzawa H, Sano H, Takahashi H, Voelker DR, et al.

Surfactant protein A inhibits peptidoglycan-induced tumor necrosis factor-alpha

secretion in U937 cells and alveolar macrophages by direct interaction with toll-like

receptor 2. The Journal of biological chemistry. 2002;277(9):6830-7. Epub 2001/11/29.

doi: 10.1074/jbc.M106671200. PubMed PMID: 11724772.

48.   Unitt J, Hornigold D. Plant lectins are novel Toll-like receptor agonists.

Biochemical pharmacology. 2011;81(11):1324-8. Epub 2011/03/23. doi:

10.1016/j.bcp.2011.03.010. PubMed PMID: 21420389.

49.   Souza MA, Carvalho FC, Ruas LP, Ricci-Azevedo R, Roque-Barreira MC. The

immunomodulatory effect of plant lectins: a review with emphasis on ArtinM

properties. Glycoconjugate journal. 2013;30(7):641-57. Epub 2013/01/10. doi:

10.1007/s10719-012-9464-4. PubMed PMID: 23299509; PubMed Central PMCID:

PMC3769584.

50.   Andrade WA, Souza Mdo C, Ramos-Martinez E, Nagpal K, Dutra MS, Melo

MB, et al. Combined action of nucleic acid-sensing Toll-like receptors and

TLR11/TLR12 heterodimers imparts resistance to Toxoplasma gondii in mice. Cell host

& microbe. 2013;13(1):42-53. Epub 2013/01/08. doi: 10.1016/j.chom.2012.12.003.

PubMed PMID: 23290966; PubMed Central PMCID: PMC3552114.

51.   Debierre-Grockiego F, Campos MA, Azzouz N, Schmidt J, Bieker U, Resende

MG, et al. Activation of TLR2 and TLR4 by glycosylphosphatidylinositols derived

from Toxoplasma gondii. J Immunol. 2007;179(2):1129-37. Epub 2007/07/10. PubMed

PMID: 17617606.

52.   Qiu J, Wang L, Zhang R, Ge K, Guo H, Liu X, et al. Identification of a TNF-

alpha inducer MIC3 originating from the microneme of non-cystogenic, virulent

Toxoplasma gondii. Scientific reports. 2016;6:39407. Epub 2016/12/22. doi:

10.1038/srep39407. PubMed PMID: 28000706; PubMed Central PMCID:

PMC5175157.

53.   Gay G, Braun L, Brenier-Pinchart MP, Vollaire J, Josserand V, Bertini RL, et al.

Toxoplasma gondii TgIST co-opts host chromatin repressors dampening STAT1-

dependent gene regulation and IFN-gamma-mediated host defenses. The Journal of

experimental medicine. 2016;213(9):1779-98. Epub 2016/08/10. doi:

10.1084/jem.20160340. PubMed PMID: 27503074; PubMed Central PMCID:

PMC4995087.

54.   Scanga CA, Aliberti J, Jankovic D, Tilloy F, Bennouna S, Denkers EY, et al.

Cutting edge: MyD88 is required for resistance to Toxoplasma gondii infection and

regulates parasite-induced IL-12 production by dendritic cells. J Immunol.

2002;168(12):5997-6001. Epub 2002/06/11. PubMed PMID: 12055206.

55.   Plattner F, Yarovinsky F, Romero S, Didry D, Carlier MF, Sher A, et al.

Toxoplasma profilin is essential for host cell invasion and TLR11-dependent induction

of an interleukin-12 response. Cell host & microbe. 2008;3(2):77-87. Epub 2008/03/04.

doi: 10.1016/j.chom.2008.01.001. PubMed PMID: 18312842.

56.   von Gunten S, Smith DF, Cummings RD, Riedel S, Miescher S, Schaub A, et al.

Intravenous immunoglobulin contains a broad repertoire of anticarbohydrate antibodies

that is not restricted to the IgG2 subclass. The Journal of allergy and clinical

immunology. 2009;123(6):1268-76 e15. Epub 2009/05/16. doi:

10.1016/j.jaci.2009.03.013. PubMed PMID: 19443021; PubMed Central PMCID:

PMC2777748.

57. Soldati D, Boothroyd JC. Transient transfection and expression in the obligate

intracellular parasite Toxoplasma gondii. Science. 1993;260(5106):349-52. Epub

1993/04/16. PubMed PMID: 8469986.

58. Kim K, Soldati D, Boothroyd JC. Gene replacement in Toxoplasma gondii with

chloramphenicol acetyltransferase as selectable marker. Science. 1993;262(5135):911-

4. Epub 1993/11/05. PubMed PMID: 8235614.

59. Aricescu AR, Owens RJ. Expression of recombinant glycoproteins in

mammalian cells: towards an integrative approach to structural biology. Curr Opin

Struct Biol. 2013;23(3):345-56. doi: 10.1016/j.sbi.2013.04.003. PubMed PMID:

23623336; PubMed Central PMCID: PMC4757734.

60. Mariano VS, Zorzetto-Fernandes AL, da Silva TA, Ruas LP, Nohara LL,

Almeida IC, et al. Recognition of TLR2 N-glycans: critical role in ArtinM

immunomodulatory activity. PLoS One. 2014;9(6):e98512. doi:

10.1371/journal.pone.0098512. PubMed PMID: 24892697; PubMed Central PMCID:

PMC4043963.

61. Carneiro AB, Iaciura BM, Nohara LL, Lopes CD, Veas EM, Mariano VS, et al.

Lysophosphatidylcholine triggers TLR2- and TLR4-mediated signaling pathways but

counteracts LPS-induced NO synthesis in peritoneal macrophages by inhibiting NF-

kappaB translocation and MAPK/ERK phosphorylation. PLoS One. 2013;8(9):e76233.

doi: 10.1371/journal.pone.0076233. PubMed PMID: 24312681; PubMed Central

PMCID: PMC3848743.

62. Young L, Sung J, Stacey G, Masters JR. Detection of Mycoplasma in cell

cultures. Nat Protoc. 2010;5(5):929-34. doi: 10.1038/nprot.2010.43. PubMed PMID:

20431538.

FIGURES

Fig 1. Lectin activity of rMIC1 and rMIC4. (A) Glycoarray of the native MIC1/MIC4

subcomplex (Lac+) and of the recombinant forms of MIC1 and MIC4. In total, 320

oligosaccharide probes were analysed by reading their fluorescence intensities, and the 20

best recognized glycans are shown. The results were represented as previously reported

[18]. **(B)** The activity and inhibition assays were performed in microplates coated with

glycoproteins with or without sialic acid, fetuin (black bars), or asialofetuin (white bars),

separately. After coating, wild type or mutated rMIC1 and rMIC4, pre-incubated with

PBS or their corresponding sugars, were added to the wells. Later, bound proteins were

detected through the addition of serum from *T. gondii*-infected mice. Data in **(B)** are

expressed as mean ±S.D. of triplicate wells and significance was calculated with

ANOVA followed by Bonferroni's multiple comparisons test. *p<0.05. Data are

representative of two **(B)** independent experiments. Gal: galactose; GalNAc: *N*-

acetylgalactosamine; Glc: glucose; Man: mannose; Fuc: fucose; Neu5Ac: *N*-

Fig 2. Microneme proteins stimulate cytokine production by dendritic cells and macrophages. (A-D) Bone marrow-derived dendritic cells (BMDCs) and **(E-H)** bone marrow-derived macrophages (BMDMs) from C57BL/6 mice were stimulated with rMIC1 (5 μg/mL) and rMIC4 (5 μg/mL) for 48 h. LPS (100 ng/mL) was used as positive control. The levels of IL-12p40, TNF-α, and IL-6 were measured by ELISA. For this assay, rMIC1 and rMIC4 were passed through polymyxin B column, followed by incubation with polymyxin B sulphate salt media preparation that was added to the culture (see Material and Methods). Data are expressed as mean ±S.D. of triplicate wells and significance was calculated with ANOVA followed by Bonferroni's multiple comparisons test. *$p<0.05$. Data are representative of three independent experiments.

Fig 3. The IL-12 production induced by rMICs is dependent on binding to TLR2

and TLR4. (A-I) Bone marrow-derived macrophages from WT, TLR2[-/-], TLR4[-/-],

double knockout TLR2[-/-]/TLR4[-/-], TLR3[-/-], TLR5[-/-], TLR9[-/-], and double knockout

TLR11[-/-]/TLR12[-/-] mice, all of the C57BL/6 background, were stimulated with rMIC1

or rMIC4 (5 μg/mL) for 48 h. Pam3CSK4 (P3C) (1 μg/mL), LPS (100 ng/mL), Poly I:C

(10 μg/mL), Flagellin (FLA) (1 μg/mL) and CpG (25 μg/mL) were used as positive

controls. IL-12p40 levels were measured by ELISA. **(J and K)** Transfected HEK293T

cells expressing TLR2 were stimulated with rMIC1 (750 nM) or rMIC4 (500 nM), and

rMIC1 (200 nM) or rMIC4 (160 nM) for HEK cells expressing TLR4, for 24 h. FSL-1

(100 ng/mL) and LPS (100 ng/mL) were used as positive controls. IL-8 levels were

measured by ELISA. **(L)** The interaction between rMICs and TLRs was evaluated by western blot. HEK293T cells transiently expressing TLR2-FLAG and TLR4-FLAG were lysed and incubated with His-rMIC1 (rMIC1^{His}) or His-rMIC4 (rMIC4^{His}). His-rMICs were subjected to Ni^{2+}-affinity resin pull-down (lanes 6 to 9) and analysed for TLR2 and TLR4 binding by protein blotting with antibodies specific for FLAG-tag and then for rMIC (IgY, polyclonal). For these assays, rMIC1 and rMIC4 were passed through polymyxin B column, followed by incubation with polymyxin B sulphate salt media preparation that was added to the culture (see Material and Methods). Data in **(A-K)** are expressed as mean ±S.D. of triplicate wells and significance was calculated with ANOVA followed by Bonferroni's multiple comparisons test. *p<0.05. Data are representative of three **(A-K)** and two **(L)** independent experiments.

Fig 4. The cellular activation induced by rMICs via TLRs depends on

carbohydrate recognition. (A) Bone marrow-derived macrophages and **(B)** bone

marrow-derived dendritic cells from C57BL/6 mice and **(C)** transfected HEK293T cells

expressing fully glycosylated TLR2 were stimulated with rMIC1 (WT) and rMIC4

(WT) or with their mutated forms, rMIC1-T126A/T220A and rMIC4-K469M, 5 μg/mL

of each, for 48 h. LPS (100 ng/mL) and FSL-1 (100 ng/mL) were used as positive

controls. IL-12p40 and IL-8 levels were measured by ELISA. **(D)** HEK293T cells

expressing fully glycosylated TLR2 (with 4 N-glycans, WT) or glycosylation mutants

of TLR2 (Δ-1; Δ-4; Δ-1,2; Δ-3,4; Δ-2,4; Δ-1,2,3; Δ-1,2,3,4) were stimulated with rMIC1

or rMIC4. FSL-1 (100 ng/mL) was used as positive control. IL-8 levels were measured

by ELISA. The statistical analysis compared fully glycosylated TLR2 (WT) and TLR2

mutants for the N-glycosylation sites for the same stimuli. For these assays, rMIC1,

rMIC1-T126A/T220A, rMIC4 and rMIC4-K469M were passed through polymyxin B

column, followed by incubation with polymyxin B sulphate salt media preparation that

was added to the culture (see Material and Methods). Data are expressed as mean ±S.D.

of triplicate wells and significance was calculated with ANOVA followed by

Bonferroni's multiple comparisons test. *p<0.05. Data are representative of three

independent experiments.

Fig 5. The IL-12 production during *T. gondii in vitro* infection partially depends on MICs and their ability to recognize carbohydrates on APCs surface. (A) Schematic representation of knockout and complementation constructs for MIC1 and MIC4 loci. The endogenous loci were disrupted using the hypoxanthine-xanthine-guanine phosphoribosyl transferase (HPT)-selectable marker and CRISPR methodology. **(B)** PCR analysis for MIC1 and MIC4 loci of gDNA from parental (WT RH-Δ*ku80*/Δ*hpt*-GFP/Luc) and knockout (RH-Δ*ku80*/Δ*mic1*-GFP/Luc and RH-Δ*ku80*/Δ*mic4*-GFP/Luc) strains. **(C)** Western blot analysis of an equal loading of whole cell lysates corresponding to 3×10^6 tachyzoites (1×10^8/mL) from WT, Δ*mic1*, Δ*mic1*::MIC1-T126A/T220A, Δ*mic4* and Δ*mic4*::K469M parasites. The membrane was probed with

anti-MIC1 (IgY, 1:20,000) and anti-MIC4 (IgY, 1:8,000). **(D)** Bone marrow-derived

dendritic cells (BMDCs) and **(E)** Bone marrow-derived macrophages (BMDMs) from

C57BL/6 were infected with WT, Δ*mic1*, Δ*mic1*::MIC1-T126A/T220A, Δ*mic4* and

Δ*mic4*::K469M strains (MOI 3). LPS (100 ng/mL) was used as positive control. Cell-

culture supernatants were collected 24 hours post-infection and the IL-12p40 production

was analyzed by ELISA. **(F and G)** Frequency of IL-12p40$^+$ BMDCs (CD11b$^+$IL-

12p40$^+$) after 20-24 hours of *in vitro* infection with WT, Δ*mic1*, Δ*mic1*::MIC1-

T126A/T220A, Δ*mic4* and Δ*mic4*::K469M strains (MOI 1). Brefeldin A was added to

the culture for 8 hours. LPS (100 ng/mL) was used as positive control. **(H)**

Representative dot plots showing IL-12p40 staining in *T. gondii* infected or non-

infected (SAG1$^+$ or SAG1$^-$) CD11b$^+$ cells after 20-24 hours of *in vitro* infection with

WT, Δ*mic1*, Δ*mic1*::MIC1-T126A/T220A, Δ*mic4* and Δ*mic4*::K469M strains (MOI 1).

Brefeldin A was added to the culture for 8 hours. LPS (100 ng/mL) was used as positive

control. Data are expressed as mean ±S.D. of triplicate wells and significance was

calculated with ANOVA followed by Bonferroni's multiple comparisons test. *p<0.05.

Data are representative of three **(D and E)** and two **(F-H)** independent experiments.

1004

1005

Fig 6. MIC1 lectin activity, but not MIC4, contributes to virulence in mice during

in vivo **infection with *T. gondii*.** CD-1 mice were infected intraperitoneally with RH

engineered strains of *T. gondii* at an infectious dose of 50 tachyzoites/mouse (n=6).

Mortality kinetics of mice infected with **(A)** WT, Δ*mic1* and Δ*mic1*::MIC1-

T126A/T220A strains or **(E)** WT, Δ*mic4* and Δ*mic4*::MIC4-K469M parasites. At day 5

post-infection the sera were collected for measuring systemic **(B and F)** IL-12p40 and

(C and G) IFN-γ. **(D, H and I)** Bioluminescent detection in photons/sec/cm^2 shows

parasite burden 5 days post-infection. Data are expressed as mean ±S.D. and

significance was calculated with ANOVA followed by Bonferroni's multiple

comparisons test. *p<0.05. Data are representative of three independent experiments,

with total n=16.

1017

1018

Fig 7. MIC1 wild type complemented strain restores virulence in mice during *in*

***vivo* infection with *T. gondii*. (A)** Western blot analysis of an equal loading of whole

cell lysates corresponding to 3×10^6 tachyzoites (1×10^8/mL) from WT, Δ*mic1*,

Δ*mic1*::MIC1-T126A/T220A and Δ*mic1*::MIC1 parasites. The membrane was probed

with anti-MIC1 (IgY, 1:20,000), anti-HA (rabbit, 1:5,000) and anti-SAG1 (rabbit,

1:10,000). **(B)** Mortality kinetics of CD-1 mice infected intraperitoneally with WT,

Δ*mic1*, Δ*mic1*::MIC1-T126A/T220A and Δ*mic1*::MIC1 at an infectious dose of 50

tachyzoites/mouse (n=5). **(C)** At day 5 post-infection the sera were collected for

measuring systemic IFN-γ. **(D and E)** Bioluminescent detection in photons/sec/cm^2

shows parasite burden 5 days post-infection. **(F and G)** Body mass and mortality

kinetics of CD-1 mice infected subcutaneously with WT, Δ*mic1*, Δ*mic1*::MIC1-

T126A/T220A and Δ*mic1*::MIC1 using an infectious dose of 10^4 tachyzoites/mouse.

Data are expressed as mean ±S.D. and significance were calculated with ANOVA

followed by Bonferroni's multiple comparisons test. *p<0.05. Data are representative of

two independent experiments, total n=10.

1034

SUPPLEMENTARY INFORMATION

S1 Fig. Effect of different concentrations of rMIC1 and rMIC4 on the transfected

HEK cells. HEK293T cells expressing **(A and B)** TLR2 or **(C and D)** TLR4 were

stimulated with increasing concentrations of **(A and C)** rMIC1 and **(B and D)** rMIC4

for 24 h. FSL-1 (100 ng/mL) LPS (100 ng/mL) were used as positive controls. IL-8

levels were measured by ELISA. Data are expressed as mean ±S.D. of triplicate wells

and significance was calculated with ANOVA. *p<0.05. Data are representative of two

independent experiments.