1 **Cortical Connectivity In A Macaque Model Of Congenital Blindness**
2

3 Loïc Magrou[1], Pascal Barone[2,3], Nikola T. Markov[4], Herbert Killackey[5], Pascale Giroud[1],
4 Michel Berland[6], Kenneth Knoblauch[1*], Colette Dehay[1*], Henry Kennedy[1*]
5

1. Univ Lyon, Université Claude Bernard Lyon 1, Inserm, Stem Cell and Brain Research
Institute U1208, 69500 Bron, France.
2. Université De Toulouse Paul Sabatier, 31062 Toulouse, France
3. Centre De Recherche Cerveau & Cognition, CNRS, UMR 5549, 31059 Toulouse, France
4. Princeton Neuroscience Institute and Department of Psychology, Princeton
University, Princeton, 08544, USA.
5. Department of Neurobiology and Behavior, University of California, Irvine, Irvine,
California 92717, USA
6. Faculté de Médecine Lyon-Sud, Service de Gynécologie Obstétrique, Centre
Hospitalier Lyon-Sud, 69495 Pierre Bénite Cedex, France
16

17 * Corresponding authors

18
19
20 **Abstract**
21

Brain-mapping of the congenitally blind human reveals extensive plasticity[1]. The
visual cortex of the blind has been observed to support higher cognitive functions
including language and numerical processing[2,3]. This functional shift is
hypothesized to reflect a metamodal cortical function, where computations are
defined by the local network. In the case of developmental deafferentation, local
circuits are considered to implement higher cognitive functions by
accommodating diverse long-distance inputs[4-7]. However, the extent to which
visual deprivation triggers a reorganization of the large-scale network in the
cortex is still controversial[8]. Here we show that early prenatal ablation of the
retina, an experimental model of anophthalmia in macaque, leads to a major
reduction of area V1 and the creation of a default extrastriate cortex (DEC)[9,10].
Anophthalmic and normal macaques received retrograde tracer injections in DEC,
as well as areas V2 and V4 post-natally. This revealed a six-fold expansion of the
spatial extent of local connectivity in the DEC and a surprisingly high location of
the DEC derived from a computational model of the cortical hierarchy[11]. In the
anophthalmic the set of areas projecting to the DEC, area V2 and V4 does not differ
from that of normal adult controls, but there is a highly significant increase in the
relative cumulative weight of the ventral stream areas input to the early visual
areas. These findings show that although occupying the territory that would have
become primary visual cortex the DEC exhibits features of a higher order area,
thus reflecting a combination of intrinsic and extrinsic factors on cortical
specification. Understanding the interaction of these contributing factors will
shed light on cortical plasticity during primate development and the neurobiology
of blindness.

46
47

Main text

Early visual cortex deafferentation via bilateral removal of the eyes at early stages of prenatal development in the macaque provides a non-human primate (NHP) model of anophthalmia. In NHP anophthalmics, there is an in-depth modification of the development of the visual system accompanied by characteristic sulci malformations; cortex destined to become striate cortex (area V1) reverts to a default phenotype (Default Extrastriate Cortex DEC)[9,10,12]. The three anophthalmics used in this study each showed important shifts in the border of striate cortex (area V1) leading to an important reduction in its dimensions (**Figure 1**). In these animals instead of the typical border between areas V1 and V2 one can detect a large region of interceding cortex where the stria of Genari is absent and its cytoarchitecture can be broadly defined as extrastriate. This stretch of cortex exhibits small islands of striate cortex and the hybrid expression of histochemical phenotypes of striate and extrastriate cortex both during *in utero* development and postnatally[12,13]. The cortex between area V2 and the reduced striate cortex area V1 corresponds to the DEC (lower panels of **Figure 1A,B**). The proportion of area V1 with respect to the total cortex is considerably reduced in anophthalmic brains compared to the normal (**Figure 1C**). This contrasts with the proportions of total visual cortex (including DEC) with respect to neocortex, which is similar in anophthalmics and normals, therefore coherent with deafferentation causing a border shift rather than merely a shrinkage of area V1. Hence the DEC plus the remaining area V1 in the anophthalmic matches the extent of area V1 in the normal animal[14].

We used retrograde tracers that allow exploring the intrinsic labelling of a cortical area, which corresponds to the local connectivity[15]. In the normal brain intrinsic connectivity corresponds to 80-90% of the total connectivity and exhibits an exponential decline with distance[15,16]. In the anophthalmic brain the space constant of the exponential decline is significantly larger and intrinsic projections extend over considerable distances (**Figure 2A**). Hence in the normal V1 the 75% threshold is at 0.25mm, the 80% at 0.35mm and the 95% at 0.80mm. In the DEC these distances are increased 4 to 6 fold (**Figure 2B**), so that local connectivity extends across a large extent of the DEC on the operculum (**Figure S1**).

The topography of connectivity in the anophthalmic was overall similar to that in the normal cortex. Following retrograde tracer injection in a target area, the numbers of labelled neurons in a given source area with respect to the total number of labelled neurons in the brain defines the Fraction of Labelled Neurons FLN, a weight index reflecting the strength of the particular pathway[15]. High frequency sampling of labelled neurons in the cortex allows characterization with a single injection of the weighted connectivity of a pathway linking any two cortical areas[17]. Injections limited to the grey matter were performed in DEC, V2 and V4 in three anophtalmic brains (**Figures S2 Table 1**). Inspection of cortical labelling suggested that early visual deafferentation leads to an increase of numbers of labelled neurons in ventral stream areas (**Figures S2, Fig S3**). Injections in DEC and area V2 show that deafferentation profoundly affects the relative strengths of the dorsal and ventral pathways, as seen after summing the FLN values across all ventral versus dorsal stream areas (**Figure 2C, D**). Differences between normal and anophthalmic cumulative FLN values were not found to be significant following injections in area V4, suggesting that the effect of deafferentation are restricted to early cortical areas.

The laminar distribution of retrogradely labelled parent neurones of a pathway is defined by its proportion of supragranular labelled neurons or SLN index, which has been shown to be highly consistent across individuals[11]. The SLN values of a pathway define it as feedforward or feedback and specify a hierarchical distance[18]. In the absence of the retina there is an increase in numbers of labelled supragranular layer neurons (**Figures S4**). The SLN value for area V2 projections to DEC is significantly higher than for the V2 projection to

V1 and likewise the projection of V3, FST and PIP to DEC have significantly increased SLN
values (**Fig S4, panels A**). Following deafferentation, projections to area V2 showed
significant increases in the SLN in areas V1 and V3 as well as the dorsal stream areas MT,
V3A, LIP, PIP, STP and PGa as well as an increase in the ventral stream area TEO (**Figures
S4, panel B**). By contrast, deafferentation had little or no effect on the SLN values for any of
the projections to area V4 (**Figure S4D**) with the marked exception of V1 where only
infragranular neurons were observed. However, given the very low FLN value, this cannot be
considered significant. The most marked change in the SLN is the projection of area V1 to the
DEC (Figure 3A). Labelled neurons in area V1 projecting to the DEC are entirely located in
the supragranular layers, which makes this projection very different from any projection from
area V1 in the normal brain (**Figure 3A**). Area V1 also projects strongly to area V2, but the
V1->V2 pathway originates from both infra- and supragranular layers. A projection of V1
which is entirely from the supragranular layers would be to area V4, but the V1->V4 pathway
is considerably weaker than the DEC->V4 pathway.
112
SLN has been shown to be a robust indicator of hierarchical distance[11,19] (see Materials
and Methods). When the SLN values extracted from injections at different levels are mapped
on to hierarchical space by means of a sigmoid function they display surprisingly good
agreement (Figure 3B). This is shown by the probit transformed values of SLN plotted in a
pairwise fashion; if the transformation leads to a coherent measure of hierarchical distance the
points will cluster around lines of unit slope. Importantly, both the normal and anophthalmic
brains display this consistency in laminar organization (**Figure, 3B, C, D**). All the common
projections to areas V1 and V2 in the normal cortex are feedback and, as expected, are
observed in the lower left quadrant. This contrasts with the anophthalmic where V1 is in the
top right quadrant, indicating it to be feedforward to DEC and V2. Consideration of an
ensemble of SLN values following injections in areas V1, V2 and V4 allows fitting a
hierarchical model to both the normal and anophthalmic data sets (**Figure 4A**)[11]. This shows
that the overall layout of the ventral and dorsal stream areas remain approximately similar to
that observed in the normal. However, in the anophthalmic brain, DEC and area V2 are
considerably higher in the hierarchy than expected. The goodness of fit of the model is
shown by close agreement between the empirical and estimated SLN values by source and
target areas (**Figure 4B**, normal r^2=0.72; enucleate r^2=0.67).
The present results show that in the anophthalmic the topography of connectivity and
global organization of the ventral and dorsal streams are largely conserved as has been
suggested by imaging studies in the human congenitally blind[20-26], in line with the evidence of
early developmental specification of the functional streams[27,28]. Further, in the anophthalmic
brain we observe en expansion of the ventral pathway that could reflect cross modal
plasticity[24]..
The present findings show that early primate visual cortex in anophthalmia is
profoundly modified both in its cytoarchitecture and local connectivity. While the global
hierarchy remains largely conserved, there are important local changes in the hierarchical
organization. These changes would seem to reflect a persistence of immature features making
the congenitally blind 'visual' cortex neotenic. Indeed, interareal connectivity during *in utero*
development in the primate undergoes extensive remodelling characterised by a greatly
expanded population of supragranular projecting neurons, a global hierarchical organization
similar to that found in the adult, an absence of ectopic pathways and finally a markedly
extensive local connectivity[11,29-31]. The relatively high position in the cortical hierarchy and the
conservation of an extensive local connectivity in the DEC could ensure the long time
constants which would be required for the observed higher cognitive functions of the
deafferentated cortex of the blind[19,32].

The present findings need to be considered in view of current understanding of the
developmental specification of the cortex. Developmental patterning of the neocortex is
consequent to an interplay between intrinsic genetic mechanisms based on morphogens and
secreted signalling molecules and extrinsic inputs relayed to the cortex by thalamocortical
axons[33-35]. The role of thalamic axons in arealization is a multistep hierarchical process
involving events at progenitor and neuronal levels[36]. A recent spatiotemporal transcriptome
analysis of the pre- and postnatal macaque forebrain revealed a small number of genes that
have persistent expression across cortical development, suggesting a large potential for
extrinsic shaping of the cortex[37]. Interestingly, this study shows that areal and laminar
molecular phenotypes are acquired late postnatally indicating a wide and potentially
important role of contextual shaping of the structure and function of the cortex, suggesting
that particular attention should be paid to the care of the young congenitally blind.
160
161 **MATERIALS & METHODS**
162
We examined the connectivity of the cortex in two 25-day old and one 10 month-old
macaques that had been enucleated between 92 and 107 days before birth i.e. between
embryonic day 58 (E58) and E73 (**Table 1**). In these three experimental animals we made six
tracer injections, a fast blue (FB) and a dyamidino (DY) injection in each and we compared
the results to 10 injections made in 8 adult controls.
Anaesthesia and Surgery. The present study is based on observations following
bilateral enucleation performed in three monkey foetuses and contrasted to eight normal
controls. The enucleated foetuses were carried to term and after birth injected with retrograde
tracers (Diamidino Yellow, DY; and Fast Blue, FB) at different postnatal ages (**Table 1).**
Pregnant cynomolgus monkeys (Macaca fascicularis) received atropine (1.25 mg, i.m.),
dexamethasone (4 mg, i.m.), isoxsuprine (2.5 mg, i.m.), and chlorpromazine (2 mg/kg, i.m.)
surgical premedication. They were prepared for surgery under ketamine hydrochloride (20
mg/kg, i.m) anaesthesia. Following intubation, anaesthesia was continued with 1-2%
halothane in a N_2O/O_2 mixture (70/30). The heart rate was monitored, and the expired CO_2
maintained between 4.5% and 6%. Body temperature was maintained using a thermostatically
controlled heating blanket. Between embryonic day 58 (E58) and E73 and using sterile
procedures a midline abdominal incision was made, and uterotomy was performed. The foetal
head was exposed, bilateral eye removal performed, and the foetus replaced in the uterus after
closing the incisions. The mother was returned to her cage and given an analgesic
(visceralgine, 1.25 mg, i.m.) twice daily for 2 days. All foetuses were allowed normal
development until term (E165).
Injections of Retrograde Tracers Identical medication, anaesthesia and monitoring
procedures were used as described above. Tracer injections were placed in the DEC, area V2
and area V4. Injections were made by means of Hamilton syringes in a stereotypic fashion.
Following injections, artificial dura mater was applied, the bone flaps were closed, cemented
and the scalp stitched back into position.
The tracer injection sites are shown in **Figure 2**. Three injections are located in the
DEC (top three rows in Figure 2), one in area V2 (fourth row **Figure 2**) and two in area V4
(last row **Figure 2**). All injections in the enucleate brain were confined to the cortical grey
matter, and except for case BB122 LH DY (third row injection sites). BB122 LH DY
injection in the DEC was very small and restricted to upper layers and is only considered for
the examination of topography. Side-by-side injections in target areas of retrograde tracers
reveal the topology of connectivity in source areas. Such side-by-side injections were made in
the DEC in the lower part of the medial operculum in case BB181 corresponding in normal
cortex to area V1subserving parafoveal visual field[38]. In case BB122 a single injection was

made in V2 near the lip of the lunate sulcus where foveal visual field is represented in the
normal cortex[39]. Finally, a pair of very large injections were made on the dorsal part of the
prelunate gyrus spanning the central and peripheral representation of area V4 in the normal
brain[40].

The full extent of labelled neurons were charted across the cortex, which was
parcellated according to a 91 area atlas[17]. Injection of retrograde tracer in an area leads to a
region of labelling in each afferent area. This region is referred to as the projection zone. So
as to obtain reliable counts of labelled neurons it is necessary to chart labelled neurons
throughout the full extent of the projection zone in each area[11,29]. This makes it possible to
estimate the fraction of labelled neuron (FLN) and the ratio of supragranular layer neurons
(SLN) in each area (see Materials and Methods). FLN is a weight index which allows
construction of a weighted and directed matrix of the cortical network [15], while SLN value of
interareal pathways constitutes an index of hierarchical distance, allowing areas to be
organized in a determinate hierarchy[11].

Animal euthanasia After 10 to 12 days of recovery that allows optimal retrograde
labelling of neurons projecting to the pick-up zone, animals were anesthetised with ketamine
(20 mg/kg, i.m.) followed by a lethal dose of Nembutal (60 mg/kg, i.p.) and perfused through
the heart with a 1.25% paraformaldehyde and 1.5% glutaraldehyde solution. After fixation,
perfusion was continued with a 10-30% sucrose solution to provide cryoprotection of the
brain.

Data Acquisition Depending in the enucleation case, parasagital (BB181) or horizontal
(BB122 and BB169) sections (40-μm thick) were cut on a freezing microtome and at least 1
in 3 sections were stained for Nissl substance. Normal controls were cut in horizontal and
coronal planes. Sections were observed in UV light with oil-immersion objectives using a
Leitz fluorescence microscope equipped with a D-filter set (355-425 nm). High precision
maps were made using Mercator software running on Exploranova technology, coupled to the
microscope stage. Controlled high frequency sampling gives stable neuron counts despite
curvature of the cortex and heterogeneity of neuron distribution in the projection zones of
individual areas[11,41] Characteristics of neurons labelled with FB or DY are described by Keizer
and colleagues[42]. Area limits and layer 4 were marked on the charts of labelled neurons. These
neurons were then attributed to areas of our atlas based on landmarks and histology, and
counted according to that parcellation[17].

Statistical analysis All statistical analyses were performed in the R statistical
environment[43] with additional tools from the MASS, aod, and Betareg packages[44-46]. Each
injection gave rise to retrogradely labelled neurons, which were plotted and compared against
those of normal animals, injected at anatomically equivalent locations. As previously shown[15],
the FLN (Fraction of Labelled Neurons), corresponds to the proportion of cells located in a
given source area with respect to the total number of labelled neurons in the cortex. The
connectivity profile is defined by the FLN values for each of the structures labelled from the
injected target area. The SLN measurement of the proportion of supragranular labelled
neurons in an area, has been shown to be a stable anatomical assessment of areal hierarchical
relationships[11,18].

FLN. The distribution of FLN values has been successfully modelled previously by a
negative binomial distribution[15,17]. This can be performed using a Generalized Linear Model
(GLM) when the dispersion parameter is fixed. We initially estimated the dispersion
parameter for individual areas obtaining values between 10.43 and 11.7, and for subsequent
tests used an average value of 4. We then used this model to compare connection strengths
(i.e. FLN values) between normal (i.e. non-enucleated) and enucleated animals. As
explanatory variables, we used a 2 level factor, Group (Normal/Enclueated) and an 2 level
factor, Area for the labelled areas projecting on the target injection. The linear predictor in

the GLM includes main effect of both factors and their interaction. In the model, the raw
neuron counts enter as the response variable, but a log link is used with the natural log (i.e.
base e) of the total number of labelled neurons in each case included as a fixed offset. In this
way, the model coefficients estimate FLN values. Confidence intervals were computed to
assess the significance of the difference, based on the model fitted estimates. To test the effect
of enucleation in either stream, the linear predictor was modified to include the factor Group
and the two level factor, Stream (Ventral/Dorsal).

SLN. A similar approach was used to analyse SLNs (Supragranular Labelled Neurons,
the proportion of labelled neurons situated above the layer 4 (granular layer) vs. below it, in
each area) but using beta regression[44] to model the proportions. Since the beta distribution is
defined on values in the interval (0, 1), it is useful to analyse proportions. Like the binomial,
the parameters of interest are linked to the explanatory values via a linear predictor. In this
case a logit link was used. Like the negative binomial distribution on counts, the model
includes a dispersion parameter that can account for overdispersion beyond that expected
from a purely binomial process. As before, confidence intervals were evaluated and
significance was assessed using likelihood ratio tests.

Hierarchy. SLN values were used to estimate hierarchical distances between areas in
normal and enucleated data sets with a previously proposed model first[11,19]. A probit model is
used to transform SLN values to a linear predictor

Acknowledgements
This work was supported LABEX CORTEX (ANR-11-LABX-0042) of Université de Lyon
(ANR-11-IDEX-0007) operated by the French National Research Agency (ANR) (H.K.).

Author contribution: CD, HK proposed the project; PG, MB, HK surgical intervention; PB,
PG, CD histological processing; LM, PB, NTM, HK, PG data acquisition; LM, PB, NMT,
KK, HK data analysis; KK computational modeling; LM, KK, CD, HK wrote the paper. All
authors contributed to editing the final document.

References

1 Sadato, N. *et al.* Activation of the primary visual cortex by Braille reading in blind
subjects. *Nature* **380**, 526-528., (1996).
2 Amedi, A., Floel, A., Knecht, S., Zohary, E. & Cohen, L. G. Transcranial magnetic
stimulation of the occipital pole interferes with verbal processing in blind subjects.
Nat Neurosci **7**, 1266-1270, (2004).
3 Kanjlia, S., Lane, C., Feigenson, L. & Bedny, M. Absence of visual experience modifies
the neural basis of numerical thinking. *Proc Natl Acad Sci U S A* **113**, 11172-11177,
(2016).
4 Pascual-Leone, A. & Hamilton, R. The metamodal organization of the brain. *Prog Brain*
Res **134**, 427-445, (2001).
5 Bedny, M. Evidence from Blindness for a Cognitively Pluripotent Cortex. *Trends Cogn*
Sci, (2017).
6 Sur, M. & Leamey, C. A. Development and plasticity of cortical areas and networks. *Nat*
Rev Neurosci **2**, 251-262, (2001).
7 Merabet, L. B. & Pascual-Leone, A. Neural reorganization following sensory loss: the
opportunity of change. *Nat Rev Neurosci* **11**, 44-52, (2010).

8   Hasson, U., Andric, M., Atilgan, H. & Collignon, O. Congenital blindness is associated
with large-scale reorganization of anatomical networks. *Neuroimage* **128**, 362-372,
(2016).
9   Dehay, C., Horsburgh, G., Berland, M., Killackey, H. & Kennedy, H. Maturation and
connectivity of the visual cortex in monkey is altered by prenatal removal of retinal
input. *Nature* **337**, 265-267, (1989).
10 Rakic, P. Specification of cerebral cortical areas. *Science* **241**, 170-176, (1988).
11 Markov, N. T. *et al.* The Anatomy of Hierarchy: Feedforward and feedback pathways
in macaque visual cortex. *J Comp Neurol* **522**, 225-259, (2014).
12 Dehay, C., Giroud, P., Berland, M., Killackey, H. P. & Kennedy, H. Phenotypic
characterisation of respecified visual cortex subsequent to prenatal enucleation in the
monkey: development of acetylcholinesterase and cytochrome oxidase patterns. *J
Comp Neurol* **376**, 386-402, (1996).
13 Dehay, C., Giroud, P., Berland, M., Killackey, H. & Kennedy, H. Contribution of thalamic
input to the specification of cytoarchitectonic cortical fields in the primate: effects of
bilateral enucleation in the fetal monkey on the boundaries, dimensions, and
gyrification of striate and extrastriate cortex. *J Comp Neurol* **367**, 70-89, (1996).
14 Dehay, C., Horsburgh, G., Berland, M., Killackey, H. & Kennedy, H. The effects of
bilateral enucleation in the primate fetus on the parcellation of visual cortex. *Dev
Brain Res* **62**, 137-141, (1991).
15 Markov, N. T. *et al.* Weight Consistency Specifies Regularities of Macaque Cortical
Networks. *Cereb Cortex* **21**, 1254-1272, (2011).
16 Horvat, S. *et al.* Spatial Embedding and Wiring Cost Constrain the Functional Layout
of the Cortical Network of Rodents and Primates. *PLoS Biol* **14**, e1002512, (2016).
17 Markov, N. T. *et al.* A weighted and directed interareal connectivity matrix for
macaque cerebral cortex. *Cereb Cortex* **24**, 17-36, (2014).
18 Barone, P., Batardiere, A., Knoblauch, K. & Kennedy, H. Laminar distribution of
neurons in extrastriate areas projecting to visual areas V1 and V4 correlates with the
hierarchical rank and indicates the operation of a distance rule. *J Neurosci* **20**, 3263-
3281, (2000).
19 Chaudhuri, R., Knoblauch, K., Gariel, M. A., Kennedy, H. & Wang, X. J. A Large-Scale
Circuit Mechanism for Hierarchical Dynamical Processing in the Primate Cortex.
Neuron **88**, 419-431, (2015).
20 Striem-Amit, E., Cohen, L., Dehaene, S. & Amedi, A. Reading with sounds: sensory
substitution selectively activates the visual word form area in the blind. *Neuron* **76**,
640-652, (2012).
21 Striem-Amit, E., Dakwar, O., Reich, L. & Amedi, A. The large-scale organization of
"visual" streams emerges without visual experience. *Cereb Cortex* **22**, 1698-1709,
(2012).
22 Striem-Amit, E. *et al.* Functional connectivity of visual cortex in the blind follows
retinotopic organization principles. *Brain* **138**, 1679-1695, (2015).
23 Matteau, I., Kupers, R., Ricciardi, E., Pietrini, P. & Ptito, M. Beyond visual, aural and
haptic movement perception: hMT+ is activated by electrotactile motion stimulation
of the tongue in sighted and in congenitally blind individuals. *Brain Res Bull* **82**, 264-
270, (2010).
24 Ptito, M., Matteau, I., Gjedde, A. & Kupers, R. Recruitment of the middle temporal area
by tactile motion in congenital blindness. *Neuroreport* **20**, 543-547, (2009).
25 Pietrini, P. *et al.* Beyond sensory images: Object-based representation in the human
ventral pathway. *Proc Natl Acad Sci U S A* **101**, 5658-5663, (2004).

26 van den Hurk, J., Van Baelen, M. & Op de Beeck, H. P. Development of visual category
selectivity in ventral visual cortex does not require visual experience. *Proc Natl Acad*
Sci U S A **114**, E4501-E4510, (2017).
27 Deen, B. *et al.* Organization of high-level visual cortex in human infants. *Nat Commun*
8, 13995, (2017).
28 Livingstone, M. S. *et al.* Development of the macaque face-patch system. *Nat Commun*
8, 14897, (2017).
29 Barone, P., Dehay, C., Berland, M., Bullier, J. & Kennedy, H. Developmental remodeling
of primate visual cortical pathways. *Cereb Cortex* **5**, 22-38, (1995).
30 Luhmann, H. J., Martinez Millan, L. & Singer, W. Development of horizontal intrinsic
connections in cat striate cortex. *Exp Brain Res* **63**, 443-448, (1986).
31 Khazipov, R. & Luhmann, H. J. Early patterns of electrical activity in the developing
cerebral cortex of humans and rodents. *Trends Neurosci* **29**, 414-418, (2006).
32 Honey, C. J. *et al.* Slow cortical dynamics and the accumulation of information over
long timescales. *Neuron* **76**, 423-434, (2012).
33 O'Leary, D. D., Chou, S. J. & Sahara, S. Area patterning of the mammalian cortex.
Neuron **56**, 252-269, (2007).
34 De la Rossa, A. *et al.* In vivo reprogramming of circuit connectivity in postmitotic
neocortical neurons. *Nat Neurosci* **16**, 193-200, (2013).
35 Geschwind, D. H. & Rakic, P. Cortical evolution: judge the brain by its cover. *Neuron*
80, 633-647, (2013).
36 Pouchelon, G. & Jabaudon, D. Nurturing the cortex's thalamic nature. *Curr Opin Neurol*
27, 142-148, (2014).
37 Bakken, T. E. *et al.* A comprehensive transcriptional map of primate brain
development. *Nature* **535**, 367-375, (2016).
38 Gattass, R., Sousa, A. P. & Rosa, M. G. Visual topography of V1 in the Cebus monkey. *J*
Comp Neurol **259**, 529-548, (1987).
39 Gattass, R., Gross, C. G. & Sandell, J. H. Visual topography of V2 in the macaque. *J Comp*
Neurol **201**, 519-539, (1981).
40 Li, C. Y., Tanaka, M. & Creutzfeldt, O. D. Attention and eye movement related activation
of neurons in the dorsal prelunate gyrus (area DP). *Brain Res* **496**, 307-313, (1989).
41 Vezoli, J. *et al.* Quantitative analysis of connectivity in the visual cortex: extracting
function from structure. *Neuroscientist* **10**, 476-482, (2004).
42 Keizer, K., Kuypers, H. G. J. M., Huisman, A. M. & Dann, O. Diamidino Yellow
dihydrochloride (DY 2HCl): a new fluorescent retrograde neuronal tracer, which
migrates only very slowly out of the cell. *Exp Brain Res* **51**, 179-191, (1983).
43 R Development Core Team. R: A language and environment for statistical computing
(R Foundation for statistical computing. http://www.r-project.org/, Vienna, Austria.,
2016).
44 Cribari-Neto, F. & Zeileis, A. Beta Regression in R. *J Stat Soft* **34**, 1-24, (2010).
45 Lesnoff, M. & Lancelot, R. aod: Analysis of Overdispersed Data. *R package version 1.3*
http://cran.r-project.org/package=aod, (2012).
46 Venables, W. N. & Ripley, B. D. *Modern Applied Statistics with S*. 4th edn, (Springer,
2002).
389
390

Figure 1. Effects of early enucleation on cortical parcellation. (**A**) Upper panel parasaggital Nissl stained sections showing cytoarchtecture; lower panel schematic showing the limits of striate cortex and area V2; (**B**) upper panel, parasaggital Nissl stained section in the neonate following prenatal enucleation at 68 days after conception (E68); lower panel, limits of areas V1, V2 and default extrastriate cortex. Sections in A and B taken from equivalent levels, arrow heads indicate limits of striate cortex. Note, large reduction of striate cortex on operculum and more modest reduction in the calcarine, scale bars, 2 mm. (**C**) Quantitative effects of enucleation on proportions of visual cortex; left-hand panel, surface area of striate cortex (p = 4.04e-05, 7 enucleates, 6 normals); middle-panel, proportion of striate cortex with respect to total visual cortex (p = 3.04e-06, 7 enucleates, 6 normals); right-hand panel, proportion of visual cortex with respect to total cortex (p = 0.63, 6 enucleates, 6 normals).

Figure 2. Intrinsic connectivity following enucleation and effects of ventralization. (A) Exponential decay of density of intrinsic neurons with distance following injection in area V1 of a normal (yellow dots, dashed line fit) and in the DEC (enucleation at E73) (blue dots, dotted line fit). The dashed and dotted lines represent exponential fits. (B) Distances within which the 3 thresholds (75%, 80%, and 95%) of intrinsic labelling are attained in normal V1 and in the DEC (enucleation at E73). (C) mean cumulative sum of Fraction of Labelled Neurons (FLN) in ventral stream areas; far-left panel, injections in area V1 and default extrastriate cortex (DEC; p = 0.0155); middle panel, injections in area V2 (p < 2e-16); right-hand panel, injections in area V4 (p = 0.301). Enucleate vs. normal across all injection: p = 2.52e-04. All tests were performed assuming that the proportions followed a beta distribution[44]. (D) Effect of enucleation on connexion strength in ventral stream areas. Log scale dot plot of FLN. Enucleates, blue dots; normal controls, black dots; upper-panel, injections in normal striate cortex (V1) and Default Extrastriate Cortex (DEC) (1 enucleate, 5 normals); middle-panel, injections in area V2 (1 enucleate, 3 normals); bottom panel, injections in area V4 (2 enucleates, 3 normal). For abbreviations of area names see glossary.

418
Figure 3. (A) High power plots comparing laminar distributions of V1->DEC to projections in the normal cortex.
This comparison shows that the V1->DEC has no counter part in the normal cortex, either in terms of strength of
projection nor laminar distribution. **(B)** Schematic illustration of distance relations motivating the pairs plots of
SLN values. Suppose that SLN is related to the hierarchical distance between an injected area A and the areas
projecting to it B_A. In particular, suppose that some function, h, of the SLN provides this distance so that
$h(\mathrm{SLN}_{B \to A}) = h_{BA}$ is the hierarchical distance between A and B. If we assume that this distance measure does
not depend on the area injected or the pairs of areas compared, then we should expect that for injections in areas
A and C that have a common projection B, the distance $h_{BA} = h_{BC} + h_{CA}$. As the relation does not depend on the
particular common area B, it should be true for all of the common projections to areas A and C, i.e., that they are
related by the fixed distance between areas A and C. This implies in turn that if there is a transformation of the
SLN values that maps onto a common distance scale across areas, then when we plot the transformed SLN
values of the common areas for two injection sites against each other, the values will fall along a line of unit
slope (blue dashed line in (B) and (C)) whose intercept is the distance between the two injection sites, h_{CA}. **(C)**
and (D) Pairs plots between probit-transformed SLN values of common areas from injections in V1/DEC, V2
and V4 (as indicated in the boxes along the diagonal). Each point represents the average pair of SLN values
obtained in a single source area; the blue dashed lines indicate the best fitting lines of unit slope. **(C)** Data from
normal controls (yellow background). **(D)** Enucleated cases (blue background). Area labels for points of
potential interest and outliers are indicated to the right of the point.

437
438
Figure 4. Developmental plasticity of the visual hierarchy. (A) Hierarchical relationship between areas in the
visual system based on injections in V1, V2 and V4 in normal animals (A), and following enucleation and
injections in Default Extrastriate Cortex (DEC), V2 and V4. The model fitted was probit(E(SLN)) = Xβ, where
probit is the inverse of a Gaussian cumulative distribution function applied to the expected value of the SLN and
X is an $n \times p$ incidence matrix for the connectivity with a column for each of p areas and a row for each of n pairs
of connected areas (repeated injections may appear as multiple rows). All elements of X are 0 except in the two
columns corresponding to the connecting pairs for that row, taking on the values -1 and 1 for the target and
source, respectively. One column is dropped for model identifiability. β is a vector of hierarchical values to be
estimated. The hierarchical levels are estimated by maximum likelihood assuming that the SLN values follow a
beta-binomial distribution (see Markov et al., 2014). The model is only determined up to a linear combination
so that the values have been scaled to the range 1-10. (B) Detailed comparison of the model with the data for
each injection by labelled area; black dots, empirical values; blue squares, predicted values. Error bars are
empirical SEs **(or 95% confidence intervals?)** for the data and model based SEs for the predictions.S