Inferring the selection window in antimicrobial resistance using deep mutational scanning data and biophysics-based fitness models

Pouria Dasmeh[1,2], Anh-Tien Ton[1,2], Caroline Quach[3,4], Adrian W.R. Serohijos[1,2]

[1]Departement de Biochimie, [2]Centre Robert Cedergren en Bioinformatique et Génomique, Université de Montréal, 2900 Edouard-Montpetit, Montreal, Quebec H3T 1J4, Canada.[3]Département de microbiologie, infectiologie et immunologie, Université de Montréal, [4]Infection Prevention and Control Unit, Department of Medical Microbiology and Infectious Diseases, CHU Sainte-Justine

Correspondence: adrian.serohijos@umontreal.ca

Abstract

Mutant-selection window (MSW) hypothesis in antimicrobial resistance implies a range for antimicrobial concentration that promotes selection of single-step resistant mutants. Since the inception and experimental verification, MSW has been at the forefront of strategies to minimize development of antimicrobial resistance (AR). Setting the upper and lower limits of MSW requires an understanding of the dependence of selection coefficient of arising mutations to antimicrobial concentration. In this work, we employed a biophysics-based and experimentally calibrated fitness model to estimate MSW in the case of Ampicillin and Cefotaxime resistance in E.coli TEM-1 beta lactamase. In line with experimental observations, we show that selection is active at very low levels of antimicrobials. Furthermore, we elucidate the dependence of MSW to catalytic efficiency of mutants, fraction of mutants in the population and discuss the role of population genetic parameters such as population size and mutation rate. Altogether, our analysis and formalism provide a predictive model of MSW with direct implications in the design of dosage strategies.

Introduction

The gain of resistance to antimicrobials is indeed one of the most challenging threats to public health. An effective approach to contain antimicrobial resistance (AMR) is multi-layered, employing knowledge of the mechanisms for gain of resistance, ecological dynamics of bacterial species and eventually treatment regimens and strategies[1].

Since their first medical use, antimicrobials have been administered at a minimum dosage sufficient to inhibit bacterial growth while keeping hazards minimal. This concentration, termed as minimum inhibitory concentration (MIC)[2,3], measures susceptibility of bacteria to drugs and is the basis of clinical protocols that compare effectiveness of antimicrobials[4]. The emergence of AMR requires that at least two sub-populations of bacterial species exist, with distinct MICs. The dosage response of bacterial growth to antimicrobials thus defines a concentration window that is above the MIC of wild-type and below the MIC of resistant mutants, i.e., mutant-selection window (MSW) hypothesis[5-7]. The upper limit of this window (i.e., MIC of resistant mutants) is termed as mutant-prevention concentration (MPC)[7]. An effective treatment strategy for AMR sets drug dosage above MPC to inhibit the selection of resistant mutant[6].

Setting the upper and lower limits of MSW is critically important. Obviously, MPC should be high enough to prevent emergence of AMR, but should be the lowest efficacious concentration to minimize side effects. The lower limit, on the other hand, is the minimum selective concentration (MSC)[8] below which resistance and efficacy are hardly achieved and can be regarded as a safety level for antimicrobial concentration in natural environments[9]. MSC was traditionally assumed to be the MIC of WT strains, but

numerous findings found MSC to be 10- to 100-folds lower and concluded that the

selection of resistant mutants is present at sub-MIC concentrations[10].

Emergence of AMR by drug selection pressure is an example of an evolutionary

process whose fate is determined by the competition between mutational supply,

selection and immigration[11]. Therefore, the rate of acquired resistance depends on the

competition between WT and resistant mutants, which in turn depends on several

factors such as the mutation rate, selection regime, population size, etc. In the limit of

large population size and low mutation rate, selective advantage of the arising mutation

becomes the key factor in determining the evolutionary dynamics. In the simplest

definition, the selection coefficient (s) of a mutant is the normalized differential fitness

between WT and mutant strains, i.e., F_{WT} and F_{mut}:

$$s = \frac{F_{mut} - F_{WT}}{F_{WT}}$$ (Eq. 1)

In principle, deleterious, neutral and beneficial nature of mutations is determined by the

product of effective population size and selection coefficient, i.e., $N_{eff} \times s$ so that $N_{eff} \times s <$

-1, = 1 and > 1 corresponds to deleterious, neutral and beneficial mutations,

respectively.

Over the last decade, considerable efforts have been made to predict protein

phenotypes that derive adaptation[12,13], in particular to drug concentration[14,15] with the

aim of providing an *ab initio* molecular model for AMR. When the gain of resistance is

through *de novo* mutations in target proteins[16], the knowledge of genotype to phenotype

mapping is crucial in predicting fitness and eventually the selection regime. In these

studies, fitness is proportional to bacterial growth, which depends on the metabolic flux

of enzymes targeted by antimicrobials such as *dihydrofolate reductase* (DHFR) in folate

pathway[14] or competitive inhibition in the case of TEM-1 beta lactamase[15,17].

Given such predictive models of AMR and the resulting fitness landscape from

exhaustive mutagenesis experiments, an interesting question is whether these models

can be employed to investigate the limits of MSW in AMR. If proved successful, this

"bottom-up" approach would result in a predictive model for selective window that will

eventually aid in optimizing dosage regimens. This is the major goal of the current

work. We will start with a general summary of the predictive model of AR in TEM-1 beta

lactamase which relates fitness to the rate of peptidoglycan synthesis and cell wall

formation. This rate in turn depends on variety of molecular and cellular factors from

permeability of antimicrobials to the catalytic efficiency, k_{cat}/K_M, of both penicillin binding

proteins (PBPs) and TEM-1 beta lactamase[17,18]. MSW is then derived from the fitness

equation and the landscape for selection coefficients of arising AMR is explored. We

then employ this method to investigate the lower and upper theoretical limits of selective

window in AMR and comment on dependence of this limit to factors such as enzyme

kinetics and the fraction of resistant mutants in the population.

95
96
97   **Results**
98
99   **Fitness model for TEM-1 beta lactamase**
100
TEM-1 beta lactamase or in general beta lactamases are a family of enzymes that give

resistance to beta lactam antibiotics[19-22]. This group of antibiotics which share the

common functional group of beta lactam, inhibits formation of the peptidoglycan cross-

links in the bacterial cell wall. The specific mode of action of beta lactam antibiotics is

the competitive binding with DD-transpeptidases, which are collectively known as

penicillin binding proteins (PBP)[23]. PBPs are involved in the final step of peptidoglycan

synthesis and thus cell wall biogenesis[24]. Resistance to beta lactam antibiotics, by beta

lactamases, is achieved through attacking the beta lactam ring of these antibiotics.

Therefore, bacterial growth under the concentration of beta lactam antibiotics is

proportional to the activity of PBPs.

Figure 1 depicts the mechanistic model for the function of TEM-1 beta

lactamases within the cell. As in Figure 1A, beta lactam antibiotics diffuse through

membrane barrier and once in periplasm, they face either hydrolyses by TEM-1 beta

lactamases or diffusion towards their target, PBPs. The fitness of a bacterial cell, i.e.,

growth rate, is modeled by the PBPs activity, v^{PBP} (Figure 1B-C) which is expressed as:

116
$$Fitness \propto v^{PBP} = \frac{k_{cat}^{PBP}.E^{PBP}.C}{\left(1+\frac{[\beta lac]p}{K_{\beta lac}^{PBP}}\right)K_m^{PBP}+C}$$
(Eq. 2)

Here, C and [ßlac]$_p$ are the periplasmic concentration of peptidyglycan and beta lactam

antibiotic, k_{cat}^{PBP}, K_m^{PBP}, E^{PBP} and $K_{\beta lac}^{PBP}$ are turnover rate, Michaelis constant,

concentration of active PBPs in the periplasm and, dissociation constant of beta lactam

antibiotic from PBPs. As assumed in previous works[17,18], the rate of beta lactam

hydrolysis is equal to that of diffusion across periplasm. Hence, one can find the

following expression for the concentration of beta lactam antibiotic in the periplasm:

123
$$[\beta lac]_p = 0.5\left([\beta lac]_m - K_m^{TEM} - \frac{k_{cat}^{TEM}.E^{TEM}}{D} + \sqrt{\left(-[\beta lac]_m + K_m^{TEM} + \frac{k_{cat}^{TEM}.E^{TEM}}{D}\right)^2 + 4[\beta lac]_m K_m^{TEM}}\right)$$
(Eq.

3)

In Eq. 3, $[\beta lac]_m$ and D are the concentration and the permeability constant for beta

lactam antibiotic and k_{cat}^{PBP}, K_m^{PBP} and E^{PBP} are turnover rate, Michaelis constant and the

concentration of active beta lactamases in the periplasm.

Substituting $[\beta lac]_p$ from Eq. 3 into Eq. 2, selection coefficient (Eq. 1) of an arising

mutation can then be calculated as:

130
$$s = \frac{\left(\frac{K_m^{PBP}}{K_{\beta lac}^{PBP}}\right)\left([\beta lac]_p^{WT} - [\beta lac]_p^{mutant}\right)}{\left(1 + \frac{[\beta lac]_p^{mutant}}{K_{\beta lac}^{PBP}}\right)K_m^{PBP} + c} \qquad \text{(Eq. 4)}$$

We define the lower and upper limits of MSW, namely MSC and MPC, as the two roots

of Eq. 4. We then systematically explore MSC, MPC and MSW as a function of

biophysical properties and population genetic parameters. For these analyses, we used

three datasets. First, a set of k_{cat} and K_M of 4997 TEM-1 mutants against ampicillin that

was obtained by fitting the described fitness model to the relative fitness data from a

recent mutational scan[17]. Second, two sets of known kinetic data for TEM-1 variants as

tabulated in Table S1-2 for reactions with ampicillin and cefotaxime.

We first considered the dependence of selection coefficient to the antimicrobial

concentration. Figure 2 shows the average and standard deviation of s for all 4997

TEM-1 mutants against Ampicillin. From the figure, mutants become more deleterious

i.e., having larger negative selection coefficients, at higher ampicillin concentrations.

Figure 2B shows the same plot with a finer range of s = -0.02 to 0.02. The area depicted

in gray shows the nearly neutral range (hereafter called nearly-neutral zone)

corresponding to $|N_{eff} \times s| \sim 1$, where the effective population size, N_{eff}, is assumed to be

10^7. Among 4997 TEM-1 mutants, 32 mutants possess positive selection coefficients at

different ranges of antibiotic concentration. As shown in the case of I173H mutant,

selection coefficient of beneficial mutations has a quadratic dependence on the

antimicrobial concentration and crosses the nearly-neutral zone at a lower and a higher

concentration of antibiotic. The lower and higher concentrations correspond to the

minimum selective concentration, MSC and, mutant prevention concentration, MPC,

respectively (see Methods).

We calculated MSC and MPC for TEM-1 beneficial mutations against ampicillin

in Figure 3A-B. As shown in Figure 3A, MSC is ~ two-three orders of magnitude lower

than MIC of TEM-1 against ampicillin (~1000-4000 μg/ml [25-27]). This shows that

selection is active at these very low concentrations in line with previous observations for

the prevalence of mutant selection at sub-MIC regimes[8,10,28,29]. There is an inverse

exponential relationship between MPC and MSC as shown in Figure 3C. Beneficial

mutations with higher selective advantages at lower antimicrobial concentrations would

indeed require higher antimicrobial concentrations for prevention. The maximum MPC

belonged to P27E mutant (~10^6 μg/ml). Notably, mutations in residue 173 that are

observed in laboratory evolution of TEM-1[30,31] and in clinical isolates[32] are among

beneficial mutations.

Next, we looked at the individual s curves for TEM-1 mutants against ampicillin.

As shown in Figure 4A, selection coefficients are sigmoidal in antimicrobial

concentrations with two interesting features: I) selection coefficients are clustered and

non-randomly distributed at different concentrations and II) mutations do not preserve

their selective rank at all antimicrobial concentrations. Simply put a high fit mutant to

lower concentrations of ampicillin (compared to all other mutants) would not retain the

same fitness rank at higher concentrations. Below, we discuss these two features in

detail.

The clustering of selection coefficients of several mutations implies that s values

are non-uniformly distributed. To see the pattern of clustering, we compared the

distribution of distances between selection coefficients with that of an equidistance

distribution using Kolmogorov-smirnov test. Selection coefficients at all antimicrobial

concentrations were significantly non-equidistant (p-value $< 10^{-16}$). Then, we looked at

the optimum number of clusters and the centroids as shown in Figure 4C-D for each

ampicillin concentration up to 10000 µg/ml (~two folds higher than MIC). Several

features of these plots are interesting. First, at low antimicrobial concentrations most

selection coefficients are within the 8^{th} and 9^{th} clusters being highly deleterious

(compare upper left dashed squares in Figures 4C and 4C). This trend is reversed at

high concentrations. As shown in bottom-right dashed squares in Figures 4C-D, at high

concentrations of antimicrobials, almost half of mutations have neutral or negligible

fitness costs. This behavior is in contrast with the average behavior shown in Figure 2A

showing a rich heterogeneity in selective advantage/disadvantage of mutant clusters at

different antibiotic concentrations. Interestingly, a sudden overpopulation of mutants

(~2/5 of all mutants) in lower more neutral clusters occurs around MIC, shown with a

black arrow in Figure 4C.

Another interesting feature of Figure 4A is that s-curves cross each other at different

antimicrobial concentrations. This implies that the rank of different mutants is not

preserved at all antimicrobial concentrations. To illustrate this observation further, we

plotted selection coefficients of two mutants, L152C and G251R, in black and gray in

Figure 5A. The corresponding values for V_{max} and K_M of both mutants are shown accordingly. From the figure, at low antimicrobial concentrations, the mutant with lower half-saturation limit (L152C) has a higher selective advantage despite having ~16% less turnover rate. The more efficient G251R mutant gains a higher fitness once antimicrobial concentration reaches the corresponding half-saturation limit.

The lack of rank-preservation at different antibiotic concentrations means that a mutant that confers resistance at one antibiotic concentration would not necessarily do so at a different concentration. This adds another level of impairment for predictability of antimicrobial resistance in addition to stochastic rise and fixation of resistant mutations. To check the rank of mutants at different antimicrobial concentrations systematically, we compared the rank of all TEM-1 mutants at 9 antimicrobial concentrations 1000 to 9000 µg/ml to the ranks at 1 µg/ml. As shown in Figure 5B, the original rank is lost at all higher concentrations. In fact, the correlation between the rank of mutants among all concentrations although significant is very weak (R~0.01, p-value<10^{-3}; see Figure S1-2). We also checked the preservation of ranks at ampicillin concentrations below 1000 µg/ml. As shown in Figure 5C, the maximum correlation is ~0.35 which decays to ~0.05 as the antimicrobial concentration increases.

To explore the dependence of MSW to enzyme kinetics, we plotted V_{max} and K_M of TEM-1 beneficial mutations. Deleterious and neutral mutations are shown in gray and beneficial mutations in black. From the figure, beneficial mutations have maximized V_{max} and minimized K_M leading to maximized V_{max}/K_M ratio (see the fitted dashed line to all beneficial mutations in Figure 6A). We then plotted MSC and MPC versus V_{max}/K_M of all beneficial mutants of TEM-1 (Figure 6B-C). From the figure, mutants with a higher

catalytic efficiency have lower MSC and higher MPC giving rise to an increased MSW.

We thus conclude that V_{max}/K_M ratio is a strong determinant of MSC and MPC in the

case of TEM-1 and ampicillin resistance.

To what extend the selection regime and the concentration-dependence of

selection coefficient discussed in Figure 2 are common to other beta-lactam

antimicrobials? To answer this question, we sought to analyze fitness effects of arising

mutations in TEM-1 against cefotaxime. Resistance to cefotaxime is widely reported for

TEM-1 and the extended-spectrum beta lactamases (ESBLs). Figure S2 shows the

selection coefficient curves for TEM-1 mutants and natural variants where kinetic data

for reaction with cefotaxime were available (Table S2). Evidently, a major difference

with the case of ampicillin is the degree to which mutants are beneficial. While almost all

TEM-1 mutants (~99% in the DMS data set) and all natural variants (Figure S3 and

Table S2) were deleterious at all ampicillin concentrations, kinetic data against

cefotaxime shows a substantially higher fraction of beneficial mutants (12 out of 16

mutants, Figure S4). From the figure, all ESBLs and TEM-1 mutants share the initial

rise in selection coefficient within the few hundred µg/ml of cefotaxime. Note that MIC of

cefotaxime is ~6 µM. The strongly beneficial nature of mutants in TEM-1 against

cefotaxime causes MSC to be ~1 µg/ml for all mutants. MPC is never reached within the

concentration range studied here, i.e., up to 10^6 µg/ml. As shown in Figure S5, the

maximum *s* in cefotaxime dataset reaches is almost two orders of magnitude higher

than the selection coefficient of the most beneficial TEM-1 mutant against ampicillin.

Discussion

The knowledge of antimicrobial concentration at which resistance is prevented is crucial for designing dosage strategies. As we showed in this work, and for the specific case of TEM-1 beta lactamase against ampicillin and cefotaxime, a direct determination of MSC and MPC is feasible if the fitness landscape of enzymes conferring antimicrobial resistance, such as TEM-1 beta lactamase is known. As we showed in this work, both MSC and MPC depend on catalytic efficiency of mutants, i.e., the V_{max}/K_M ratio in the case of ampicillin. However, for strongly beneficial mutations, as observed in the case of cefotaxime, MSC is essentially zero and MPC fall beyond the range of biologically relevant concentrations. One would thus expect that resistance to cefotaxime is much more prevalent than to ampicillin. Therefore, it is essential to estimate not only the fraction of beneficial mutations but also the magnitude of fitness effects, i.e., the full distribution of fitness effects (DFE). We thus propose that DFE obtained from deep mutational scanning combined with fitness models, either biophysics-based as in this work or phenomenological[33], has a great potential to estimate the spectrum of resistance.

In all calculations in this work, we assumed that both WT and mutants have equal initial finesses. This assumption is biologically unrealistic as resistant mutants are shown to be enriched in the population from very low fractions of the order of 10^{-4} [29]. To check for this effect, we used the fraction of mutants in the population as a scaling factor for fitness in Equation 2 (see Methods) and plotted selection coefficients versus ampicillin for the case of I173H mutant in Figure 7B. From the figure, MSW is essentially zero for the same beneficial mutation but with negligible fractions in the population.

Therefore, figures 7B entails a window of opportunity for containment of AMR if

antibiotics are administered at the very onset of emergence of resistance. An exact

profiling of both type and fraction of resistant mutants is necessary for this purpose.

Two factors, namely bacterial population size and immune response, influence

the near-neutrality zone and potentially influence MSW and the fraction of beneficial

mutations. First, from classical population genetics, population size scales up fixation of

beneficial mutations. Since selection is stringer at higher population sizes, slightly

beneficial mutations at a smaller population (e.g., in the regime of $N_{eff}.s \sim 1$) could be

fixated with a higher rate at the larger population. Therefore, higher population sizes

decrease the width of near-neutrality zone and gives rise to lower MSC, higher MPC

and a wider MSW.

Second, in the absence of immune defense as in laboratory settings, the near-

neutrality zone defines the lowest limit of selection. However, a major difference

between these settings and the clinical settings as in the patient body is the presence of

immune defense. In fact, the first response to bacterial infections is through the immune

system. Therefore, one expects that the real selection limit is extended above the near-

neutrality zone and to the limit below which infection is contained by the immune

defense. This factor narrows down the MSW and thus the number of resistant

conferring mutations is further decreased.

We summarize the role of confounding factors on MSW as studied in this work in

Figure 7B. First, any factor that increases selective advantage of mutants such as a

higher catalytic efficiency widens MSW. We provided an equation for the relationship

between MSC and MPC to the V_{max}/K_M ratio and expect that such relationships exist for

other antimicrobials and target enzymes. Second, MSW is substantially narrowed when

the fraction of resistant mutations is negligible in the population. Third, lower population

sizes or higher immune defense extend the near-neutrality zone and thus narrow down

MSW. We anticipate that optimization of MSW with respect to each of these factors

would provide a powerful strategy for rational design of dosage strategy in the treatment

of antimicrobial resistance.

291
292
293   **Materials and method:**
294

Fitness function

Determination of both upper and lower limits to MSW in our approach requires proper

definition of fitness of an organism. Following the original work of Zimmerman and

Rosselet[18] and the following theoretical works[17,34,35], we assumed that beta lactam

antimicrobials passively diffuse through the outer membranes[36,37]. Once in the

periplasm, the antimicrobial either get hydrolyzed by beta lactamase or diffuses to the

target PBP where competitively inhibits formation of peptidoglycan by interacting with

penicillin binding proteins (Figure 1A). The rate equations for beta lactam antibiotics and

peptidoglycan synthesis can be written as

$\frac{d}{dt}[\beta lac]_p = D\big([\beta lac]_m - [\beta lac]_p\big) - \frac{k_{cat}^{\beta lactamase}[\beta lactamase][\beta lac]_p}{K_M^{\beta lactamase}+[\beta lac]_p} = 0$ (Eq. 5)

$\frac{d}{dt}[PepGly] = \frac{k_{cat}^{PBP}[PBP][PepGly]}{\left(1+\frac{[\beta lac]_p}{K_{\beta lac}^{PBP}}\right)K_M^{PBP}+[PepGly]} \equiv Fitness$ (Eq. 6)

As described earlier, the rate of beta lactam diffusion in the periplasm is assumed to be

equal to that of hydrolysis by beta lactamase[17,18]. Using fitness from Eq. 6, the relative

fitness ($\hat{F}$) of each mutant to WT TEM-1 is then expressed as:

$$\hat{F} = \frac{A}{1+B[\beta lac]_p^{mut}} - \frac{A}{1+B[\beta lac]_p^{WT}} \qquad \text{(Eq.7)}$$

where $A = k_{cat}^{PBP}[PBP][PepGly]$ and $B = \frac{K_M^{PBP}}{K_{\beta lac}^{PBP}(K_M^{PBP}+[PepGly])}$. Stiffler et al.[17] fitted equation

7 to relative fitness obtained from deep mutational scanning data and estimated both A

and B constants and K_{cat} and K_M of 4997 TEM-1 mutants using a monte carlo simulated

annealing approach.

314
315
316
317
318   **Table 1.** Parameters used in the fitness function with the corresponding values.

Parameter	Definition	Value (unit)	References
K_{cat}^{PBP}	Enzyme turnover rate of PBPs	~0.05 (s^{-1})^a	38,39
K_M^{PBP}	Michaelis constant of PBPs	~ 83 mM	38,39
E_{PBP}	Concentration of folded and active PBPs	0.12 - 4 µM	*Calculated ^{b}*
E_{PG}	Periplasmic concentration of peptidoglycan	~11 µM	40
K_{blac}^{PBP}	Dissociation constant of beta-lactam from PBPs	200 – 1600 µM^{-1}	41
k_{cat}^{TEM-1}	Enzyme turnover rate of TEM-1	~1- 5000 (1/s)	25,26,42-50
K_M^{TEM-1}	Michaelis constant of TEM-1	~4 - 3500 (1/s)	25,26,42-50
E_{TEM-1}	Concentration of folded and active TEM-1	~3-4 µM	17
D	permeability parameter for beta-lactam	~0.002 - 0.005	17

319   a: For values in ranges, the average value is used in the model. b: See supplementary information for
320   details.

321
322
323   **References**
324

325   1.   Lee, C.-R., Cho, I.H., Jeong, B.C. & Lee, S.H. Strategies to minimize antibiotic resistance.
326   *International journal of environmental research and public health* **10**, 4274-4305 (2013).
327   2.   Birkhaug, K.E. Phenyl-mercuric-nitrate. *The Journal of Infectious Diseases*, 250-261
328   (1933).

3.   Garrod, L.P. The effect of bacterial numbers on minimum bacteriostatic concentrations.
Journal of Infectious Diseases **57**, 247-251 (1935).

4.   Schwalbe, R., Steele-Moore, L. & Goodwin, A.C. *Antimicrobial susceptibility testing*
protocols, (Crc Press, 2007).

5.   Drlica, K. The mutant selection window and antimicrobial resistance. *Journal of*
Antimicrobial Chemotherapy **52**, 11-17 (2003).

6.   Drlica, K. & Zhao, X. Mutant selection window hypothesis updated. *Clinical infectious*
diseases **44**, 681-688 (2007).

7.   Dong, Y., Zhao, X., Domagala, J. & Drlica, K. Effect of fluoroquinolone concentration on
selection of resistant mutants of Mycobacterium bovis BCG andStaphylococcus aureus.
Antimicrobial agents and chemotherapy **43**, 1756-1758 (1999).

8.   Liu, A. *et al.* Selective advantage of resistant strains at trace levels of antibiotics: a
simple and ultrasensitive color test for detection of antibiotics and genotoxic agents.
Antimicrobial agents and chemotherapy **55**, 1204-1210 (2011).

9.   Baquero, F., Martínez, J.-L. & Cantón, R. Antibiotics and antibiotic resistance in water
environments. *Current opinion in biotechnology* **19**, 260-265 (2008).

10.   Andersson, D.I. & Hughes, D. Microbiological effects of sublethal levels of antibiotics.
Nature Reviews Microbiology **12**, 465-478 (2014).

11.   zur Wiesch, P.A., Kouyos, R., Engelstädter, J., Regoes, R.R. & Bonhoeffer, S. Population
biological principles of drug-resistance evolution in infectious diseases. *The Lancet*
infectious diseases **11**, 236-247 (2011).

12.   Dean, A.M. & Thornton, J.W. Mechanistic approaches to the study of evolution: the
functional synthesis. *Nature Reviews Genetics* **8**, 675-688 (2007).

13.   Harms, M.J. & Thornton, J.W. Evolutionary biochemistry: revealing the historical and
physical causes of protein properties. *Nature Reviews Genetics* **14**, 559-571 (2013).

14.   Rodrigues, J.V. *et al.* Biophysical principles predict fitness landscapes of drug resistance.
Proceedings of the National Academy of Sciences, 201601441 (2016).

15.   Knies, J.L., Cai, F. & Weinreich, D.M. Enzyme efficiency but not thermostability drives
cefotaxime resistance evolution in TEM-1 β-lactamase. *Molecular biology and evolution*
34, 1040-1054 (2017).

16.   Woodford, N. & Ellington, M.J. The emergence of antibiotic resistance by mutation.
Clinical Microbiology and Infection **13**, 5-18 (2007).

17.   Stiffler, M.A., Hekstra, D.R. & Ranganathan, R. Evolvability as a function of purifying
selection in TEM-1 β-lactamase. *Cell* **160**, 882-892 (2015).

18.   Zimmermann, W. & Rosselet, A. Function of the outer membrane of Escherichia coli as a
permeability barrier to beta-lactam antibiotics. *Antimicrobial Agents and Chemotherapy*
12, 368-372 (1977).

19.   Livermore, D.M. beta-Lactamases in laboratory and clinical resistance. *Clinical*
microbiology reviews **8**, 557-584 (1995).

20.   Philippon, A., Labia, R. & Jacoby, G. Extended-spectrum beta-lactamases. *Antimicrobial*
agents and chemotherapy **33**, 1131 (1989).

21.   Jacoby, G.A. & Medeiros, A.A. More extended-spectrum beta-lactamases. *Antimicrobial*
Agents and Chemotherapy **35**, 1697 (1991).

22. Bradford, P. Extended-spectrum betalactamases in the 21st century: characterization,
epidemiology, and detection of this important resistance threat. ClinMicrobiol Rev 2001;
14: 933-51. *B, III* (1999).
23. Hartman, B.J. & Tomasz, A. Low-affinity penicillin-binding protein associated with beta-
lactam resistance in Staphylococcus aureus. *Journal of Bacteriology* **158**, 513-516 (1984).
24. Sauvage, E., Kerff, F., Terrak, M., Ayala, J.A. & Charlier, P. The penicillin-binding proteins:
structure and role in peptidoglycan biosynthesis. *FEMS microbiology reviews* **32**, 234-
258 (2008).
25. Huang, W., Le, Q.-Q., LaRocco, M. & Palzkill, T. Effect of threonine-to-methionine
substitution at position 265 on structure and function of TEM-1 beta-lactamase.
Antimicrobial agents and chemotherapy **38**, 2266-2269 (1994).
26. Marciano, D.C., Brown, N.G. & Palzkill, T. Analysis of the plasticity of location of the
Arg244 positive charge within the active site of the TEM-1 β-lactamase. *Protein Science*
18, 2080-2089 (2009).
27. Salverda, M.L. *et al.* Initial mutations direct alternative pathways of protein evolution.
PLoS genetics **7**, e1001321 (2011).
28. Zhanel, G.G., Mayer, M., Laing, N. & Adam, H.J. Mutant prevention concentrations of
levofloxacin alone and in combination with azithromycin, ceftazidime, colistin
(Polymyxin E), meropenem, piperacillin-tazobactam, and tobramycin against
Pseudomonas aeruginosa. *Antimicrobial agents and chemotherapy* **50**, 2228-2230
(2006).
29. Gullberg, E. *et al.* Selection of resistant bacteria at very low antibiotic concentrations.
PLoS pathogens **7**, e1002158 (2011).
30. Barlow, M. & Hall, B.G. Predicting evolutionary potential: in vitro evolution accurately
reproduces natural evolution of the TEM β-lactamase. *Genetics* **160**, 823-832 (2002).
31. Barlow, M. & Hall, B.G. Experimental prediction of the natural evolution of antibiotic
resistance. *Genetics* **163**, 1237-1241 (2003).
32. Zarnayová, M. *et al.* Survey of Enterobacteriaceae producing extended-spectrum β-
lactamases in a Slovak hospital: dominance of SHV-2a and characterization of TEM-132.
Antimicrobial agents and chemotherapy **49**, 3066-3069 (2005).
33. Levison, M.E. & Levison, J.H. Pharmacokinetics and pharmacodynamics of antibacterial
agents. *Infectious disease clinics of North America* **23**, 791-815 (2009).
34. Nikaido, H. & Normark, S. Sensitivity of Escherichia coli to various β-lactams is
determined by the interplay of outer membrane permeability and degradation by
periplasmic β-lactamases: a quantitative predictive treatment. *Molecular microbiology*
1, 29-36 (1987).
35. Lakaye, B., Dubus, A., Joris, B. & Frère, J.-M. Method for estimation of low outer
membrane permeability to β-lactam antibiotics. *Antimicrobial agents and chemotherapy*
46, 2901-2907 (2002).
36. Richmond, M. & Sykes, R. The β-lactamases of gram-negative bacteria and their possible
physiological role. *Advances in microbial physiology* **9**, 31-88 (1973).
37. Nikaido, H. Outer membrane of Salmonella typhimurium: transmembrane diffusion of
some hydrophobic substances. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **433**,
118-132 (1976).

38. Zhang, W., Shi, Q., Meroueh, S.O., Vakulenko, S.B. & Mobashery, S. Catalytic mechanism
of penicillin-binding protein 5 of Escherichia coli. *Biochemistry* **46**, 10113-10121 (2007).
39. Van der Linden, M., De Haan, L., Dideberg, O. & Keck, W. Site-directed mutagenesis of
proposed active-site residues of penicillin-binding protein 5 from Escherichia coli.
Biochemical Journal **303**, 357-362 (1994).
40. Wientjes, F., Woldringh, C. & Nanninga, N. Amount of peptidoglycan in cell walls of
gram-negative bacteria. *Journal of bacteriology* **173**, 7684-7691 (1991).
41. Fuda, C., Suvorov, M., Vakulenko, S.B. & Mobashery, S. The basis for resistance to β-
lactam antibiotics by penicillin-binding protein 2a of methicillin-resistant Staphylococcus
aureus. *Journal of Biological Chemistry* **279**, 40802-40806 (2004).
42. Cantu, C., Huang, W. & Palzkill, T. Cephalosporin substrate specificity determinants of
TEM-1 β-lactamase. *Journal of Biological Chemistry* **272**, 29144-29150 (1997).
43. LENFANT, F. *et al.* Site-directed mutagenesis of β-lactamase TEM-1. *The FEBS Journal*
217, 939-946 (1993).
44. Imtiaz, U., Manavathu, E.K., Lerner, S. & Mobashery, S. Critical hydrogen bonding by
serine 235 for cephalosporinase activity of TEM-1 beta-lactamase. *Antimicrobial agents*
and chemotherapy **37**, 2438-2442 (1993).
45. Gibson, R., Christensen, H. & Waley, S. Site-directed mutagenesis of β-lactamase I.
Single and double mutants of Glu-166 and Lys-73. *Biochemical journal* **272**, 613-619
(1990).
46. Majiduddin, F.K. & Palzkill, T. An analysis of why highly similar enzymes evolve
differently. *Genetics* **163**, 457-466 (2003).
47. Raquet, X. *et al.* TEM β-lactamase mutants hydrolysing third-generation cephalosporins:
a kinetic and molecular modelling analysis. *Journal of molecular biology* **244**, 625-639
(1994).
48. Salverda, M.L., De Visser, J.A.G. & Barlow, M. Natural evolution of TEM-1 β-lactamase:
experimental reconstruction and clinical relevance. *FEMS microbiology reviews* **34**,
1015-1036 (2010).
49. Sideraki, V., Huang, W., Palzkill, T. & Gilbert, H.F. A secondary drug resistance mutation
of TEM-1 β-lactamase that suppresses misfolding and aggregation. *Proceedings of the*
National Academy of Sciences **98**, 283-288 (2001).
50. Thomas, V.L. *et al.* Structural consequences of the inhibitor-resistant Ser130Gly
substitution in TEM β-lactamase. *Biochemistry* **44**, 9330-9338 (2005).
449

450
Figure 1. Scheme of the fitness model and the corresponding dosage dependence profile for
selection coefficient. A) Beta lactam antimicrobials diffuses through the outer membrane and faces
either diffusion to their targets, penicillin binding proteins, PBPs or hydrolysis by beta lactamases. B)
Fitness is proportional to the rate of peptidoglycan formation and the activity of penicillin binding proteins.
Beta lactams involve in competitive inhibition of PBPs. The resulting fitness function is a function of I)
catalytic efficiency of beta lactamases, catalytic efficiency of PBPs, concentration of beta lactams in the
periplasm and cellular constants such a permeability of beta lactam antimicrobials. D) Using the
functionality of growth rate to different molecular and cellular constants, fitness of different mutants can
be expressed as a function of antimicrobial concentration. E) Selection coefficient of an arising mutation
can be expressed as a function of antimicrobial concentration depending on the fitness curves of WT and
resistant mutants. If mutant is more fit than WT (e.g., has a higher kinetic efficiency-the upper curve),
selection coefficient is positive, i.e., red arrow, otherwise negative, i.e., the blue arrow.

472
Figure 2. Selection coefficient of TEM-1 mutants versus ampicillin concentration. A) Average (black
line) and standard deviation (gray bars) of selection coefficient of 4997 mutants of TEM-1 versus
ampicillin concentration. B) Beneficial mutations show a quadratic concave dependence to ampicillin
concentrations. The gray area is the zone at which the mutations have nearly-neutral effects on fitness.
Above and below this zone, mutations are either beneficial or deleterious. Minimum selective
concentration (MSC) and mutant prevention concentration (MPC) are the concentrations at which
selection coefficients fall in the nearly-neutral zone.

480
481

482
483 **Figure 3. Lower bound, minimum selective concentration, MSC, and upper bound, mutant**
484 **prevention concentration, MPC of mutant selection window.** A) MSC B) MPC and C) MPC versus
485 MSC of beneficial mutants in TEM-1 with ampicillin.   The dashed line shows minimum inhibitory
486 concentration, MIC, of WT TEM-1.

487
488

Figure 4. Significant heterogeneity of selection coefficients at different ampicillin concentrations.
A) The selection coefficient curves (s-curves) for a set of 100 mutants. B) Clustering of s-curves at different ampicillin concentrations. Up to 9 unique clusters are detected with different sizes. C) Size and D) centroid of each cluster at different antimicrobial concentrations.

506
507 **Figure 5. TEM-1 mutants do not preserve their ranks in selective advantage at all concentrations.**
508 A) S-curves of two TEM-1 mutants, G251R and L152C, which crosses at antimicrobial concentration of
509 ~4000 µg/ml. B) Ranks of TEM-1 mutants in selective advantage/disadvantage compared to the rank at 1
510 µg/ml as the reference set. C) Spearman correlation coefficients between the ranks at each concentration
511 below 1000 µg/ml and the reference at 1 µg/ml.

512
513

514
515 **Figure 6. MSC and MPC of TEM-1 beneficial mutations as a function of catalytic efficiency.** A) K_M
516 versus V_{max} for all (shown in gray) and beneficial (shown in black) TEM-1 mutants. B) MSC and C) MPC
517 of beneficial mutants versus catalytic efficiency.

532
533 **Figure 7.** A) Selection coefficient of a beneficial TEM-1 mutant (I173H) versus ampicillin concentration as
534 a function of mutant fraction in the population. B) The schematic influence of confounding factors such as
535 enzymatic efficiency and fraction in the population on MSW.

536
537