1 **Invariant errors reveal limitations in motor correction rather than constraints on error sensitivity**
2

3 Hyosub E. Kim[1,2], J. Ryan Morehead[3], Darius E. Parvin[1,2], Reza Moazzezi[4], Richard B. Ivry[1,2]

4 [1]Department of Psychology and [2]Helen Wills Neuroscience Institute, University of California, Berkeley
5 [3]John A. Paulson School of Engineering and Applied Sciences, Harvard University
6 [4]Berkeley, CA
7

Implicit sensorimotor adaptation is traditionally described as a process of error reduction,

whereby a fraction of the error is corrected for with each movement. Here, in our study of healthy

human participants, we characterize two constraints on this learning process: the size of adaptive

corrections is only related to error size when errors are smaller than 6°, and learning functions

converge to a similar level of asymptotic learning over a wide range of error sizes. These findings

are problematic for current models of sensorimotor adaptation, and point to a new theoretical

perspective in which learning is constrained by the size of the error correction, rather than

sensitivity to error.

Movement errors are ubiquitous, arising from numerous sources such as motor noise, fatigue, or changes

in the environment. A large body of evidence has revealed that the motor system compensates for errors

via sensorimotor adaptation (Shadmehr, Smith, & Krakauer, 2010). This implicit learning process is

thought to be driven by sensory prediction error, the discrepancy between the actual and predicted

sensory outcome of a motor command (Burge, Ernst, & Banks, 2008; Chaisanguanthum, Shen, & Sabes,

2014; Kording, Tenenbaum, & Shadmehr, 2007; Mazzoni & Krakauer, 2006; Wolpert, Ghahramani, &

Jordan, 1995). A core issue for models of adaptation has centered on how this error signal is used to

modify motor output (Atkeson, 1989; Fine & Thoroughman, 2006; Thoroughman & Shadmehr, 2000; Wei

& Körding, 2009).

In classic models of sensorimotor adaptation, the response to error is assumed to be linear, with trial-by-

trial corrections a constant fraction of error size (Jordan & Rumelhart, 1992; Thoroughman & Shadmehr,

2000). The theoretical foundation for this relationship centers on the delta learning rule, where the weights

between putative sensorimotor neurons are updated as a function of the magnitude of the difference

between the actual and predicted output (Cheng & Sabes, 2006; Tanaka, Krakauer, & Sejnowski, 2012;

Thoroughman & Shadmehr, 2000). This error term is multiplied by a scalar 'learning rate' to determine

how much the state of the system will change from trial-to-trial. However, empirical studies have shown

limitations with the assumption of a constant learning rate. When operationalized as the ratio of the

change in behavior relative to the error, sensitivity appears to be higher for small errors, with the system

correcting for a relatively large fraction of the error, and then rapidly decreases as errors become large

(Fine & Thoroughman, 2006; Kasuga, Hirashima, & Nozaki, 2013; Marko, Haith, Harran, & Shadmehr,

2012; Wei & Körding, 2009).

However, due to inherent confounds in standard sensorimotor adaptation tasks, estimates of the error

sensitivity function in many of these studies may be contaminated by other learning processes (e.g.,

explicit strategies) (McDougle, Ivry, & Taylor, 2016). Here, we use task-irrelevant clamped visual

feedback, a method thought to induce implicit adaptation from sensory prediction errors without

interference from explicit learning or performance-driven corrections (Morehead, Taylor, Parvin, & Ivry,

2017) (Fig. 1b). We employ this method to measure the rate of adaptive corrections to small and large

perturbations, building on previous studies that have examined this issue using errors that vary randomly

from trial to trial (Kasuga et al., 2013; Marko et al., 2012; Wei & Körding, 2009). Notably, the error signal

remains invariant with the clamp method; thus, we can examine the full accumulated adaptive response

to errors of a given size, something not possible when the size and direction of the error is randomized.

In a between-subject design, participants (n=96, 12/group) were presented with visual feedback that was

clamped to a fixed path which was angularly offset from the target by 0°, 1°, 1.75°, 3.5°, 6°, 10°, 15°, or

45°. This manipulation was explicitly described to the participants and they were instructed to ignore the

feedback and simply move directly to the target (Fig. 1a). With the exception of the 0° control group, all

groups implicitly adapted to the clamp (t_{11}=-.04, p=.97 for 0° group; t_{11}>5.9, p<.0001 for all other groups;

Fig. 1c,d).

Within all adapting groups, there was an effect of clamp size on the average per-trial rate of learning over

the first five cycles (ANOVA: $F_{6,77}$=6.45, p<.0001, η^2=.33). Although there was a modest linear

relationship between clamp size and early adaptation rate (r_{82}=.29, p=0.01), the adaptation rate appeared

to be composed of two zones, one where the rate scaled in proportion to error size, and another where

rates were virtually invariant. To formally assess this hypothesis, we performed segmented linear

regressions. Taking model complexity into account, a two-region segmented regression yielded the best

model (see Supplementary Fig. 1). This model predicted that the breakpoint between the proportional

and saturated zones was at the remarkably low value of 4.4° (95% CI [3.9°, 4.9°], Fig. 1e).

These results, in combination with previous work (Fine & Thoroughman, 2006; Kasuga et al., 2013; Marko

et al., 2012; Morehead et al., 2017; Wei & Körding, 2009), are clearly at odds with models entailing a

fixed learning rate (i.e., adaptation scaling linearly with error size). Prior observations of a nonlinear

response to error have inspired models in which the learning rate saturates for large errors (Tanaka et

al., 2012), or large errors are discounted before the update step of the learning process (Körding &

Wolpert, 2004; Marko et al., 2012; Wei & Körding, 2009). If exposed to a constant perturbation, these

models can generate similar adaptation rates and asymptotes in response to large errors. However, these

models would also predict a lower asymptote in response to small errors in the linearly proportional zone.

Contrary to this prediction, with the exception of the 1° group, the magnitude of adaptation at the end of

training was similar for a wide range of clamp offsets (Fig. 1f and Supplement).

78   **<<< FIGURE 1 >>>**

Figure 1 Initial adaptation rates scale with error size, yet saturate to an invariant response magnitude. (**a**) Illustration of experimental apparatus and task structure for Exp. 1. (**b**) Schematic view of clamped visual feedback paradigm, in which the angular path of the cursor is independent of hand movement direction. (**c-d**) Behavior for all groups, divided into two panels for visualization purposes. The small clamp groups (**c**) demonstrate adaptation rates which scale with error size, whereas the large clamp groups (**d**) show saturated responses. (**e**) Segmented regression indicates that the initial adaptation rate scales between 0° and 4.4° before saturating for all errors above this breakpoint (dashed vertical lines represent 95% C.I.). (**f**) Sensorimotor aftereffects, measured during the first cycle following the clamp block. Dots are individuals; shading and error bars denote SEM. Gray shading denotes cycles without visual feedback. Data used for analyses are provided in *Figure 1-source file1*.

The preceding observation is tempered by the fact that we did not have a sufficient number of trials to

ensure that learning had become asymptotic. To address this issue, we conducted a second experiment

in which the number of cycles was increased from 40 to 160. Participants (n=30, 10/group) were exposed

to clamped feedback with an angular offset of 1.75°, 3.5°, or 15°. These offsets were chosen because

they span the range of early adaptation rates observed in Experiment 1. Consistent with the results of

Experiment 1, there was a clear scaling of the rates across the proportional zone (ANOVA: $F_{2,27} = 18.6$;

$p<.0001$; $\eta^2=.58$), with Tukey Kramer post hoc tests revealing significant differences between all pairwise

comparisons (Fig. 2a). Strikingly, the three groups reached similar asymptotes, with all groups

demonstrating final aftereffects of ~25° (ANOVA: $F_{2,27}=0.39$, $p=.68$; $\eta^2=0.03$) (Fig. 2b; see Supplement,

"Comparisons between Experiment 1 and Experiment 2"). This dissociation between size-dependent

early adaptation rates and invariant asymptotic adaptation is at odds with models that correct for a

constant fraction of error size as well as error discounting models (See Supplement, "Computational

models…"). Even if the learning rate varies as a function of error size, these models predict that

asymptotic behavior will diverge for error sizes eliciting different adaptation rates since the asymptote is

determined by the equilibrium between learning and forgetting.

107   **<<< FIGURE 2 >>>**

Figure 2 Implicit adaptation converges on a common asymptote. (**a**) The 1.75°, 3.5°, and 15° clamp groups in Exp. 2 adapted at markedly different rates (bar graphs depict mean of cycles 3-7). However, there was convergence of all three learning functions by the end of 160 cycles, and (**b**) no difference between groups in the size of the final aftereffects. Asterisk in (**a**) denotes significant differences between groups early in the clamp phase. Data used for analyses are provided in *Figure 2-source file 2*.

In addition to identifying a fundamental limitation with current models of sensorimotor adaptation, our results draw attention to a more general issue. Behavioral responses to error are usually interpreted through the lens of error sensitivity. In this framework, the sensitivity function is generated by dividing the magnitude of the system's adaptive correction by the error size. When applied to the behavioral data that we and others have observed, this divisive operation generates a function in which sensitivity is high for small errors and gradually decreases to near zero for large errors (Fig. 3a,c). However, this transformation of behavioral responses obscures what may be the true limiting factor in adaptation: Adjustments in motor output only scale for small errors before quickly reaching a saturation point for larger errors up to 95° (Fig. 3b).

<<< FIGURE 3 >>>

Figure 3 Two ways of assessing the adaptive response: nonlinearly discounted error sensitivity (left) or
a saturating error correction function (right). Here we plot data from several different studies (Bond &
Taylor, 2015; Morehead et al., 2017; Wei & Körding, 2009 - Experiments 1 and 2), including the present
one. The top panel (**a** and **b**) shows two different ways of evaluating trial-by-trial changes in hand angle
(see Methods). Note how the error correction function (**b**) which starts small then saturates, is obscured
when transformed into a sensitivity function (**a**) that starts large and quickly decays. Plotting the aftereffect
data in terms of a sensitivity function (**c**) also fails to capture the relative invariance of these data within
a given experimental context (**d**).

Focusing on the actual behavioral change from sensory prediction error highlights the limited dependency

of the system on error size, as well as the common asymptotic level of learning in response to small and

large errors (Fig. 3d). We can envision two ways in which current models of adaptation could be modified

to capture these effects. First, the learning rate and retention parameters could be coupled, scaling

together with error size, while the asymptotic response may reflect the limit of proprioceptive recalibration,

which is independent of visual error size. Alternatively, the adaptation system may normalize its

responses to sensory prediction errors with repeated exposure, akin to normalization processes observed

in response to reward prediction errors (Tobler, Fiorillo, & Schultz, 2005). For example, the system may

increase its responses to small, yet persistent errors. Regardless of the appropriate reformulation of

models of sensorimotor adaptation, we expect it will be fruitful to shift the focus away from the error

sensitivity of the learning system, and instead, address the constraints on the behavioral change that

arises in response to the error.

Acknowledgments

We thank Wendy Shwe for assistance with the data collection. We are also grateful to Maurice Smith and

Matthew Boggess for helpful discussions. This work was supported by grant NS092079 from the National

Institutes of Health.

Competing Interests

No competing interests, financial or otherwise, are declared by the authors.

REFERENCES

Atkeson, C. G. (1989). Learning ARM Kinematics and Dynamics. *Annual Review of Neuroscience*,
12(1), 157–183. https://doi.org/10.1146/annurev.ne.12.030189.001105
Bond, K. M., & Taylor, J. A. (2015). Flexible explicit but rigid implicit learning in a visuomotor adaptation
task. *Journal of Neurophysiology*, *113*(10), 3836–3849. https://doi.org/10.1152/jn.00009.2015
Burge, J., Ernst, M. O., & Banks, M. S. (2008). The statistical determinants of adaptation rate in human
reaching. *Journal of Vision*, *8*(4), 20.1-2019. https://doi.org/10.1167/8.4.20
Chaisanguanthum, K. S., Shen, H. H., & Sabes, P. N. (2014). Motor Variability Arises from a Slow
Random Walk in Neural State. *Journal of Neuroscience*, *34*(36), 12071–12080.
https://doi.org/10.1523/JNEUROSCI.3001-13.2014
Cheng, S., & Sabes, P. N. (2006). Modeling sensorimotor learning with linear dynamical systems.
Neuronal Computation, *18*(4), 760–793. https://doi.org/10.1162/089976606775774651.Modeling
Fine, M. S., & Thoroughman, K. (2006). Motor Adaptation to Single Force Pulses: Sensitive to Direction
but Insensitive to Within-Movement Pulse Placement and Magnitude. *Journal of*
Neurophysiology, *96*(2), 710–720. https://doi.org/10.1152/jn.00215.2006
Jordan, M. I., & Rumelhart, D. E. (1992). Forward Models: Supervised Learning with a Distal Teacher,
307354.
Kasuga, S., Hirashima, M., & Nozaki, D. (2013). Simultaneous Processing of Information on Multiple
Errors in Visuomotor Learning. *PLoS ONE*, *8*(8). https://doi.org/10.1371/journal.pone.0072741
Kording, K. P., Tenenbaum, J. B., & Shadmehr, R. (2007). The dynamics of memory as a consequence
of optimal adaptation to a changing body. *Nature Neuroscience*, *10*(6), 779–786.
https://doi.org/10.1038/nn1901
Körding, K. P., & Wolpert, D. M. (2004). The loss function of sensorimotor learning. *Proceedings of the*
National Academy of Sciences of the United States of America, *101*(26), 9839–9842.
https://doi.org/10.1073/pnas.0308394101
Marko, M. K., Haith, A. M., Harran, M. D., & Shadmehr, R. (2012). Sensitivity to prediction error in
reach adaptation. *Journal of Neurophysiology*, *108*(6), 1752–1763.
https://doi.org/10.1152/jn.00177.2012
Mazzoni, P., & Krakauer, J. W. (2006). An implicit plan overrides an explicit strategy during visuomotor
adaptation. *J Neurosci*, *26*(14), 3642–3645. https://doi.org/10.1523/jneurosci.5317-05.2006
McDougle, S. D., Ivry, R. B., & Taylor, J. A. (2016). Taking Aim at the Cognitive Side of Learning in
Sensorimotor Adaptation Tasks. *Trends in Cognitive Sciences*, *20*(7), 535–544.
https://doi.org/10.1016/j.tics.2016.05.002
Morehead, J. R., Taylor, J. A., Parvin, D. E., & Ivry, R. B. (2017). Characteristics of Implicit
Sensorimotor Adaptation Revealed by Task-irrelevant Clamped Feedback. *Journal of Cognitive*
Neuroscience, *29*(6), 1061–1074. https://doi.org/10.1162/jocn_a_01108
Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error correction, sensory prediction, and
adaptation in motor control. *Annu Rev Neurosci*, *33*, 89–108. https://doi.org/10.1146/annurev-
neuro-060909-153135
Tanaka, H., Krakauer, J. W., & Sejnowski, T. J. (2012). Generalization and Multirate Models of Motor
Adaptation. *Neural Computation*, *24*(4), 939–966. https://doi.org/10.1162/NECO_a_00262
Thoroughman, K. A., & Shadmehr, R. (2000). Learning of action through adaptive combination of motor
primitives. *Nature*, *407*(6805), 742–7. https://doi.org/10.1038/35037588
Tobler, P. N., Fiorillo, C. D., & Schultz, W. (2005). Adaptive Coding of Reward Value by Dopamine
Neurons. *Science*, *307*(5715), 1642–1645. https://doi.org/10.1126/science.1105370
Wei, K., & Körding, K. (2009). Relevance of Error: What Drives Motor Adaptation? *Journal of*
Neurophysiology, *101*(2), 655–664. https://doi.org/10.1152/jn.90545.2008
Wolpert, D. M., Ghahramani, Z., & Jordan, M. I. (1995). An internal model for sensorimotor integration.
Science (New York, N.Y.), *269*(5232), 1880–1882.
213

METHODS

Participants: Healthy, young adults (N=126, 89 females, age = 21±2 years old) were recruited from the University of California, Berkeley, community. Each participant was tested in only one experiment. All participants were right-handed, as verified with the Edinburgh Handedness Inventory (Oldfield, 1971). Participants received course credit or financial compensation for their participation. The Institutional Review Board at UC Berkeley approved all experimental procedures.

Experimental Apparatus: The participant was seated at a custom-made tabletop housing an LCD screen (53.2 cm by 30 cm, ASUS), mounted 27 cm above a digitizing tablet (49.3 cm by 32.7 cm, Intuos 4XL; Wacom, Vancouver, WA). The participant made reaching movements by sliding a modified air hockey "paddle" containing an embedded stylus. The position of the stylus was recorded by the tablet at 200 Hz. The experimental software was custom written in Matlab, using the Psychtoolbox extensions (Pelli, 1997).

Reaching Task: Center-out planar reaching movements were performed from the center of the workspace to targets positioned at a radial distance of 8 cm. Direct vision of the hand was occluded by the monitor, and the lights were extinguished in the room to minimize peripheral vision of the arm. The start location and target location were indicated by white and blue circles, respectively (both 6 mm in diameter).

To initiate each trial, the participant moved the digitizing stylus into the start location. The position of the stylus was indicated by a white feedback cursor (3.5 mm diameter). Once the start location was maintained for 500 ms, the target appeared at one of 8 locations, placed in 45° increments around a virtual circle. Participants were instructed to accurately and rapidly "slice" through the target, without needing to stop at the target location. Visual feedback, when presented, was provided during the reach until the movement amplitude exceeded 8 cm. As described below, the feedback either matched the position of the stylus (veridical) or followed a fixed path (clamped). If the movement was not completed within 300 ms, the words "too slow" were generated by the sound system of the computer.

In Experiment 1, the position of the cursor was frozen for 1 s once the movement amplitude reached 8

cm. The participant was free to begin moving back to the start location during this time. After the spatial

feedback period, the cursor disappeared. Once the participant's hand was back within 2 cm of the start

circle, a white ring appeared, indicating the radial distance between the hand and center start position.

The ring was displayed to aid the participant in returning to the start location, without providing angular

information about hand position. Two changes were made in Experiment 2: First, the cursor was turned

off 50 ms after the hand crossed the virtual target ring. Second, during the return movement, the feedback

cursor reappeared when the participant's hand was within 1 cm of the start. These changes reduced the

time required for each trial and allowed the participants to complete the extended number of trials required

in Experiment 2 within our time constraints. Average total trial time in Experiment 1 was $4.45 \pm .62$ s

versus $2.51 \pm .26$ s in Experiment 2.

Experimental Feedback Conditions: Across the experimental session, there were three types of visual

feedback. On no-feedback trials, the cursor disappeared when the participant's hand left the start circle

and only reappeared at the end of the return movement. On veridical feedback trials, the cursor matched

the position of the stylus during the 8 cm outbound segment of the reach. On clamped feedback trials,

the feedback followed a path that was fixed along a specific heading angle (Morehead et al., 2017;

Shmuelof et al., 2012; Vaswani et al., 2015). The radial distance of the cursor from the start location was

still based on the radial extent of the participant's hand during the 8 cm outbound segment, but the angular

position was fixed relative to the target (i.e., independent of the angular position of the hand).

The primary instructions to the participant remained the same across the experimental session:

Specifically, that they were to reach directly towards the visual target. Prior to the introduction of task-

irrelevant clamped feedback trials, participants were briefed about the feedback manipulation. They were

informed that the position of the cursor would now follow a fixed trajectory and that the angular position

would be independent of their movement. They were explicitly instructed to ignore the cursor and continue

to reach directly to the target. The same instructions in abbreviated form ("Ignore the cursor and move

your hand directly to the target location") were repeated verbally and with onscreen text after 20

movement cycles in Experiment 1 (exact mid-point) and every 40 movement cycles during Experiment

2.

Experiment 1

In a previous experiment, adaptation to task-irrelevant clamped visual feedback was statistically uniform

to offsets between 7.5° - 95°. The main goal of Experiment 1 was to investigate if there was a dependency

on error size for angles smaller than 7.5°. Participants (n=96, 12/group) were randomly assigned to one

of eight groups that differed in terms of the size of the clamped visual feedback: 1°, 1.75°, 3.5°, 6°, 10°,

15°, and 45° (with a 0° group included as a control). Given that the target diameter was 6 mm and the

feedback cursor diameter was 3.5 mm, a substantial portion of the cursor overlapped with the target for

the 1° and 1.75° clamps, and was fully embedded in the case of the 0° clamp. Half of the participants

trained with a clockwise clamp offset, and the other half with a counterclockwise clamp offset.

The session began with two baseline blocks, the first comprised of 5 movement cycles (40 reaches to 8

targets) without visual feedback and the second comprised of 5 cycles with a veridical cursor displaying

hand position. The experimenter then informed the participant that the visual feedback would no longer

be veridical and would now be clamped at a fixed angle from the target location. The clamp block had 40

cycles. A short break (<30 s), as well as a reminder of the task instructions, was provided at the mid-way

point of this block. Immediately following the perturbation block, there were two washout blocks, first a 5

cycle block in which there was no visual feedback, followed by 5 cycles with veridical visual feedback.

Participants were debriefed at the end of the experiment and asked whether they ever intentionally tried

to reach to locations other than the target. All subjects reported aiming to the target throughout the

experiment.

Experiment 2

In Experiment 2 we assessed adaptation over an extended number of task-irrelevant clamped visual

feedback trials. The purpose of extending the perturbation block was to ensure that participants reached

asymptotic levels of learning. We were particularly interested in whether asymptotic adaptation would

converge in response to small and large clamps.

Participants (n=30, 10/group) were assigned to either a 1.75°, 3.5°, or a 15° clamped visual feedback

group. Clockwise and counterclockwise perturbations were counterbalanced within each group. As in

Experiment 1, the session started with two baseline blocks, 5 cycles without visual feedback and then 5

cycles with veridical feedback. However, the number of trials in the clamped visual feedback block was

quadrupled to 160 cycles. We included 1 cycle with no visual feedback after every 40 movement cycles.

The purpose of these interspersed no-feedback trials was to gauge adaptation magnitudes in the

absence of the learning stimulus (i.e., clamped visual feedback) at different time points within the

extended clamp block (see Supplement, *"All no-feedback probes"*). Immediately prior to the no feedback

block, the participant was informed that there would be a few trials without feedback and reminded to

always reach directly to the target. The experiment ended with a final block of 5 cycles with veridical

visual feedback of the participant's hand position.

Comparison of error sensitivity and error correction:

For the comparison of error sensitivity and error correction functions in Figure 3, we used the early

adaptation rate (panels a and b) and aftereffect data (panels c and d) from the present study, as well as

data sets from three other studies that have compared adaptive responses to a range of error sizes (Bond

& Taylor, 2015; Morehead et al., 2017; Wei & Körding, 2009). The data from Experiments 1 and 2 in Wei

and Kording were transformed from Cartesian coordinates (as presented in their paper) to polar

coordinates. The data from Bond and Taylor were restricted to the aftereffect data from their Experiment

3 (comparison of adaptation to different rotation sizes). We used the data from the initial adaptation cycles

(mean change in hand angle over first ten movement cycles) and aftereffect phase from Experiment 4 of

Morehead et al. (comparison of different clamp offsets). To obtain measures of error sensitivity, the raw

response magnitudes were divided by their corresponding error size. We did not include the data from

the 1° group in our Experiment 1 since the group function was clearly not at (or likely approaching)

asymptote.

Data Analysis:

All statistical analyses and modeling were performed using Matlab 2015b and the Statistics Toolbox. The

primary dependent variable in all experiments was endpoint hand angle, defined by the angle of the hand

position relative to the target at the time the radial distance of the hand reached 8 cm from the start

position (i.e., angle between lines connecting start position to target and start position to hand). Additional

analyses were performed using hand angle at peak radial velocity rather than endpoint hand angle. The

results were essentially identical for the two dependent variables; as such, we only report the results of

the analyses using endpoint hand angle.

Outlier responses were removed from the analyses. To identify these, the Matlab "smooth" function was

used to calculate a moving average (using a 5-trial window) of the hand angle data for each target

location. Outliers were trials in which the observed hand angle deviated by more than 3 standard

deviations from the moving average function. This procedure resulted in the elimination of ~1% of trials

involved in our statistical analyses of early adaptation rates and aftereffects; our findings are the same

whether tests were performed with or without outlier removal (values reported in main text are with outlier

removal). In total, less than 1% of trials overall, with a maximum of 2% for an individual, were removed.

Movement cycles consisted of 8 consecutive reaches (1 reach/target). Early adaptation rate was

quantified by averaging the endpoint hand angle values over cycles 3-7 of the clamp, and dividing by the

number of cycles (i.e., 5) to get an estimate of the per trial rate of change in hand angle. (As a check, we

performed a secondary analysis using cycles 2-10 and obtained nearly identical results.) The aftereffect

was quantified by using the data from the first no-feedback cycle following the last clamp cycle. Details

for all four no-feedback cycles in Experiment 2 are provided in the Supplemental section.

All t-tests were two-tailed. In order to confirm that there was a robust adaptive response in Experiment 1,

a paired t-test was performed comparing baseline hand angle during the last cycle of the veridical

feedback baseline to the first no feedback cycle (i.e., aftereffect) immediately following the perturbation

block. Posthoc tests following significant ANOVAs were performed using Tukey-Kramer's Honest

Significant Difference in order to determine specific differences in group means. Partial eta squared (η^2)

values are provided as a measure of effect size.

For the segmented linear regression (SLR) performed in Experiment 1, contiguous regression lines were

fit to the data, with each line having an independent intercept and slope. Parameters for the regression

lines were identified by a least-squares fitting procedure. Boundaries for the adjoining segments were

estimated by finding the break point(s), an additional parameter defining where two separate regression

lines meet, that minimized the residual sum of squares. Relative fits were compared using corrected

Akaike Information Criterion (AICc) values, a procedure that adjusts for the number of data points and

assigns penalties for extra parameters.

No statistical methods were used to predetermine sample sizes. The chosen sample sizes were based

on our previous study using the clamp method (Morehead et al., 2017), as well as prior psychophysical

studies of human sensorimotor learning (Galea, Mallia, Rothwell, & Diedrichsen, 2015; Gallivan, Logan,

Wolpert, & Flanagan, 2016; Huang, Haith, Mazzoni, & Krakauer, 2011; Vaswani et al., 2015).

Reaction and Movement Times:

Average movement times were quite fast, averaging 119 $\pm$ 29 ms in Experiment 1 and 136 $\pm$ 25 ms in

Experiment 2. The instructions did not impose any constraints on reaction time. On average, in

Experiment 1 participants initiated their reaches in 429 $\pm$ 72 ms, while in Experiment 2 reaction times

were 364 $\pm$ 54 ms. No significant correlations were found between these temporal variables and our

primary measures of adaptation (rate and aftereffect magnitude).

SUPPLEMENTAL INFORMATION
379
The present study revealed core features of implicit adaptation to sensory prediction errors in the absence

of task performance error. In contrast to previous studies using clamped visual feedback (Scheidt,

Conditt, Secco, & Mussa-Ivaldi, 2005; Vaswani et al., 2015), our method includes full explanation of the

nature of the manipulation, including explicit instructions to ignore the cursor feedback while always

aiming directly to the target. As verified in post-session surveys, participants do not volitionally alter their

aiming behavior since they are aware that this will not impact the visual feedback.

Experiment 1 analyses

a. Segmented regression analysis

389
390
Supplementary Figure 1
Segmented regression analysis of early adaptation rates. In our previous study using the clamped
feedback method, we observed similar adaptation rates for clamp offsets of 7.5° - 95°. In the current
study, we focus on clamps < 7.5°. As described in the text, there was a clear scaling of adaptation rates
for the smallest clamps (Figs. 1e, 2a). To determine the error size at which adaptation rates saturate (i.e.,
the breakpoint), we performed a segmented regression analysis, a widely used method for fitting data
that are hypothesized to have at least two different linear regions (Hudson, 1966). This figure shows the
results for a simple linear regression (1 region) and two segmented regression models, one with 2 regions
and the other with 3 regions, along with the residual sum of squares (RSS) and corrected Akaike
Information Criterion (AICc) scores. The best model, as determined by the AICc scores, was the two-
region model. Models with more than three regions scored progressively worse and are not shown.
Consistent with our prediction of a breakpoint < 7.5°, the predicted breakpoint was 4.4°.

b. Aftereffect data

In Experiment 1, there was an effect of error size on the aftereffect data obtained after 40 cycles of

exposure to the clamped feedback ($F_{6,77}$=3.25, p=.007, η^2=.20, Fig. 1g). However, this effect was driven

by the 1° group, and likely due to the trivial reason that the design did not include a sufficient number of

training cycles for this group to reach asymptote. A Tukey-Kramer posthoc test revealed the only

significant differences were between the 1° group and larger clamp sizes. Furthermore, excluding the 1°

group, there were no reliable differences among all of the other pairwise comparisons (all t_{22}<2.07, all

p>.05). Indeed, the magnitude of the aftereffect was not correlated with error size (r_{82}=0.04, p=0.71).

These results suggested that asymptotic adaptation may be independent of error size, a hypothesis which

was more rigorously tested in Experiment 2.

Experiment 2 analyses
417

c. All no-feedback probes

Supplementary Figure 2
Data for all of the no-feedback probes (x-axis represents number of clamp cycles preceding the no-
feedback probe) from Experiment 2. The results of a two-way repeated measures ANOVA revealed no
main effect for group (between subjects factor: $F_{2,27}$=.62, p=.54, η^2=.04), a significant effect for probe
(within subjects factor: $F_{3.81}$=19.2, p<10^{-7}, η^2=.07), and a significant interaction effect ($F_{6,81}$=2.64, p=.02,
η^2=.02). Due to the interaction, separate between subjects one-way ANOVAs were conducted for each
of the four no feedback cycles. There was a marginal effect of group (i.e., clamp size) at the first probe
only ($F_{2,27}$ = 2.77; p = .08; η^2=.21). Over subsequent blocks learning functions for each group began to
converge, resulting in the corresponding values for no-feedback probes 2-4: $F_{2,27}$ = .92, .30, .39; p= .41,
.75, .68; η^2= .07, .02, .03).
430

433   *d. Computational models of sensorimotor adaptation predict divergence of asymptotic adaptation when*
434   *early adaptation rates are different.*
435

436   A single rate state-space model can generally provide a reasonable account of the performance changes

437   observed in standard adaptation studies in which the feedback is contingent on the movement (Galea,

438   Mallia, Rothwell, & Diedrichsen, 2015; Huang, Haith, Mazzoni, & Krakauer, 2011; Thoroughman &

439   Shadmehr, 2000). This model takes the following form:

440                                 $z_{n+1} = Az_n + Be_n$

441                                 $\hat{y}_n = -z_n$

z_n represents the state estimate of the perturbation on trial n. The learning rate, B, corresponds to the

proportion of the error, e_n, that is corrected for, and A is a retention factor which represents the proportion

of the state retained from one trial to the next. The reach direction relative to the target on trial n, $\hat{y}_{n,,}$ is

opposite in sign to the state estimate. Note that unlike standard adaptation experiments in which the error

changes over trials, the error remains constant with the clamp method.

We evaluated different state-space models, focusing on the data from Experiment 2 (reproduced in the

top panel of Fig. S3). For our model fitting and simulation procedures we applied standard bootstrapping

techniques, constructing group-averaged hand angle data 1000 times by randomly resampling with

replacement from the participant pool. Using Matlab's *fmincon* function, we estimated the retention and

learning parameters which minimized the least squared error between the bootstrapped data and model

output ($\hat{y}_n$). All values in brackets represent the 95% bootstrapped C.I. of the resampled means.

We first verified that a single rate state-space model, which in its basic form assumes single, constant A

and B values for all error sizes, cannot account for the key features of the current experiments. We

estimated A and B for each of the 1000 bootstrap simulations ([.86, .95] and [.11, .29] for A and B,

respectively). In contrast to the invariant asymptotes observed in Experiment 2, the simulated functions

generated with these parameter estimates showed wildly different asymptotic values: [3.2°, 4.8°], [6.5°,

9.6°], and [27.9°, 41.0°] in response to 1.75°, 3.5°, and 15° clamps, respectively. As the error term is

constant during a clamp (i.e., e_n is equal to clamp size on every trial), and the asymptote is equal to

(Be)/(1-A), different error terms (i.e., clamp sizes) will always generate different asymptotic values if *A*

and *B* are both fixed.

As noted in the main text, various studies have suggested that the learning rate, *B*, may vary with error

size (Fine & Thoroughman, 2006; Kasuga et al., 2013; Körding & Wolpert, 2004; Marko et al., 2012;

Tanaka et al., 2012; Wei & Körding, 2009). To examine this class of models, we used a state space

model with a common retention factor and three learning rates, one for each error (clamp) size. We again

obtained 1000 sets of parameter estimates from the bootstrapped data in order to generate the

simulations seen in the bottom panel of Fig. S3. This 4-parameter model (a common *A*, and separate *B*

values for each clamp size) also failed to capture the key features of the behavior. Although the model

generated more similar asymptotes across clamp sizes than the single rate state-space model with

constant *A* and *B* ([18.4°, 28.8°], [25.1°, 32.6°] and [21.6°, 34.5°] for 1.75°, 3.5°, and 15° clamps), it failed

to capture the clear separation of early adaptation rates observed in the behavioral data (Fig. S3, top

panel). In fact, the model predicted nearly identical early adaptation rates (i.e., mean change in hand

angle per movement cycle over first 5 cycles) for the 3.5° and 15° clamps ([.9°, 1.4°] and [.7°, 1.5°]), and

only a modestly slower rate for the 1.75° clamp [.7°, 1.1°]. Moreover, the estimates of the mean error

size-dependent *B* values varied from .10 to .70, a range that seems highly unlikely and discrepant with

estimates in the adaptation literature for visuomotor rotations (Galea et al., 2015; Huang et al., 2011;

Thoroughman & Shadmehr, 2000).

We also considered a variant of this model in which we recapitulate the different adaptation rates by

using the best fit *A* value of .95 and approximating the mean *B* values directly from the early cycles of

the bootstrapped data, using the mean change in hand angle over the first 5 cycles divided by clamp

size. This procedure yielded the following *B* values: [.36, .60], [.31, .59], and [.13, .21] for 1.75°, 3.5°, and

15° clamps, respectively. Here the predicted asymptotes in response to the three clamp sizes again

strayed far from the observed behavior: [12.2°, 20.1°], [20.4°, 39.3°], and [36.4°, 60.4°].

Lastly, we tested a model (Tanaka et al., 2012) motivated by observations suggesting that the perceived

error is linear for small errors before saturating at an upper bound for larger errors (Fine & Thoroughman,

2006). To model this nonlinear effect, the authors replaced the error term, *e*, with a function which

saturates for larger errors (*s*tanh(e_n/s)*). *s* is a scaling parameter for the hyperbolic tangent function:

$$z_{n+1} = Az_n + B{*}s{*}tanh(e_n/s)$$

The 95% confidence intervals for *A* and *s* were [.93, .96] and [1.0, 2.5], respectively. This model produced

learning functions which suffered from the same shortcomings as the state-space model with error-

dependent learning rates, again predicting virtually identical early adaptation rates for the 3.5° and 15°

clamp ([.8°, 1.3°] and [.8°, 1.5°]), a *B* value [.70, 1.0] which far exceeds normal estimates of this value,

and a slightly lower asymptote for the 1.75° clamp ([18.7°, 27.4°] versus [24.8°, 31.6°] and [25.2°, 33.7°]

for the 3.5° and 15° clamps, respectively).

In summary, even models that account for variation in sensitivity to error fail to capture the combined

effects of divergent early adaptation rates and invariant asymptotic performance.

504
Supplementary Figure 3
Model simulations fail to capture key features of adaptation to different error sizes. (**a**) Behavioral data
from Experiment 2 (same as Fig. 2 of main text, but with no-feedback cycles removed). (**b**) Simulations
using a common retention factor and error size dependent learning rates (best fit A and learning rates,
B_e, for each clamp size; 95% CI in brackets). The simulated functions predict identical adaptation rates
for the 3.5° and 15° clamp conditions over the initial portion of the perturbation block, in contrast to the
markedly different rates observed in the actual behavior (top panel). Lines and shading denote mean and
SEM, respectively.

e. Comparisons between Experiment 1 and Experiment 2

Overall, the magnitudes of final aftereffects in Experiment 2 were considerably larger than the aftereffect

of matched groups in Experiment 1. One reason for this is, of course, the increase in the number of clamp

cycles in Experiment 2, which had 160 compared to the 40 cycles used in Experiment 1. As can be seen

in Supplementary Figure 2, the magnitude of adaptation is still rising after 40 cycles of clamped feedback.

As an exploratory analysis, we performed a between-experiment 2-way ANOVA, with one factor being

experiment (1 or 2) and the other error size, limiting this to the three conditions in Experiment 1 that were

included in Experiment 2 (1.75°, 3.5° and 15°). We compared the aftereffect data after 40 clamp cycles

(i.e., final aftereffects from Experiment 1 compared to first no feedback probe in Experiment 2). There

was a significant effect of experiment ($F_{1,60}$=5.27, p=.03, η^2=.07), a marginal effect of clamp size ($F_{2,60}$ =

2.8, p=.07, η^2=.08), and no interaction ($F_{2,60}$=.57, p=.57, η^2=.02). These results suggest that some

performance differences may have arisen from the methodological changes introduced in Experiment 2,

namely, eliminating frozen endpoint feedback and utilizing a quicker method for finding the start position

before each trial (see Methods). This resulted in a shorter total trial duration in Experiment 2, likely

reducing the decay of learning associated with delay (Kim, Oh, & Schweighofer, 2015). This point

highlights that asymptotic magnitude is context specific, as adaptation is highly sensitive to variables

such as temporal delay between hand and cursor movement (Held, Efstathiou, & Greene, 1966;

Kitazawa, Kohno, & Uka, 1995), viewing angle (Jalali, Miall, & Galea, 2017), and the nature of visual

feedback (e.g., online versus endpoint feedback) (J A Taylor, Krakauer, & Ivry, 2014; Jordan A. Taylor &

Ivry, 2011). As such, our claim that the asymptote is independent of error size holds for a given context;

the value will shift in other contexts, although the shift will be uniform for all error sizes.

539   **References for Methods and Supplemental Information**

540   Bond, K. M., & Taylor, J. A. (2015). Flexible explicit but rigid implicit learning in a visuomotor adaptation
541   task. *Journal of Neurophysiology*, *113*(10), 3836–3849. https://doi.org/10.1152/jn.00009.2015
542   Galea, J. M., Mallia, E., Rothwell, J., & Diedrichsen, J. (2015). The dissociable effects of punishment
543   and reward on motor learning. *Nature Neuroscience*, *18*(4), 597–602.
544   https://doi.org/10.1038/nn.3956
545   Gallivan, J. P., Logan, L., Wolpert, D. M., & Flanagan, J. R. (2016). Parallel specification of competing
546   sensorimotor control policies for alternative action options. *Nature Neuroscience*, *19*(2), 320–
547   326. https://doi.org/10.1038/nn.4214
548   Held, R., Efstathiou, A., & Greene, M. (1966). Adaptation to displaced and delayed visual feedback
549   from the hand. *Journal of Experimental Psychology*, *72*(6), 887.
550   Huang, V. S., Haith, A., Mazzoni, P., & Krakauer, J. W. (2011). Rethinking motor learning and savings
551   in adaptation paradigms: model-free memory for successful actions combines with internal
552   models. *Neuron*, *70*(4), 787–801. https://doi.org/10.1016/j.neuron.2011.04.012
553   Hudson, D. J. (1966). Fitting Segmented Curves Whose Join Points Have to Be Estimated. *Journal of*
554   *the American Statistical Association*, *61*(316), 1097–1129.
555   https://doi.org/10.1080/01621459.1966.10482198
556   Jalali, R., Miall, R. C., & Galea, J. M. (2017). No consistent effect of cerebellar transcranial direct
557   current stimulation (tDCS) on visuomotor adaptation. *Journal of Neurophysiology*,
558   jn.00896.2016. https://doi.org/10.1152/jn.00896.2016
559   Kim, S., Oh, Y., & Schweighofer, N. (2015). Between-Trial Forgetting Due to Interference and Time in
560   Motor Adaptation. *PLoS ONE*, *10*(11). https://doi.org/10.1371/journal.pone.0142963
561   Kitazawa, S., Kohno, T., & Uka, T. (1995). Effects of delayed visual information on the rate and amount
562   of prism adaptation in the human. *Journal of Neuroscience*, *15*(11), 7644–7652.
563   Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory.
564   *Neuropsychologia*, *9*(1), 97–113.
565   Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: transforming numbers into
566   movies. *Spatial Vision*, *10*(4), 437–442. https://doi.org/10.1163/156856897X00366
567   Scheidt, R. A., Conditt, M. A., Secco, E. L., & Mussa-Ivaldi, F. A. (2005). Interaction of Visual and
568   Proprioceptive Feedback During Adaptation of Human Reaching Movements. *Journal of*
569   *Neurophysiology*, *93*(6), 3200–3213. https://doi.org/10.1152/jn.00947.2004
570   Shmuelof, L., Huang, V. S., Haith, A. M., Delnicki, R. J., Mazzoni, P., & Krakauer, J. W. (2012).
571   Overcoming Motor "Forgetting" Through Reinforcement Of Learned Actions. *Journal of*
572   *Neuroscience*, *32*(42), 14617–14621a. https://doi.org/10.1523/JNEUROSCI.2184-12.2012
573   Taylor, J. A., & Ivry, R. B. (2011). Flexible Cognitive Strategies during Motor Learning. *PLoS*
574   *Computational Biology*, *7*(3), e1001096. https://doi.org/10.1371/journal.pcbi.1001096
575   Taylor, J. A., Krakauer, J. W., & Ivry, R. B. (2014). Explicit and implicit contributions to learning in a
576   sensorimotor adaptation task. *J Neurosci*, *34*(8), 3023–3032.
577   https://doi.org/10.1523/jneurosci.3619-13.2014
578   Vaswani, P. A., Shmuelof, L., Haith, A. M., Delnicki, R. J., Huang, V. S., Mazzoni, P., … Krakauer, J.
579   W. (2015). Persistent Residual Errors in Motor Adaptation Tasks: Reversion to Baseline and
580   Exploratory Escape. *Journal of Neuroscience*, *35*(17), 6969–6977.
581   https://doi.org/10.1523/JNEUROSCI.2656-14.2015
582