Insights into Origin and Evolution of α-proteobacterial Gene Transfer Agents

Migun Shakya[1,2], Shannon M. Soucy[1], and Olga Zhaxybayeva[1,3,*]

[1]Department of Biological Sciences, Dartmouth College, Hanover NH 03755, USA

[2]Present address: Bioscience Division, Los Alamos National Laboratory, Los Alamos, NM 87544

[3]Department of Computer Science, Dartmouth College, Hanover NH 03755, USA

*Corresponding author, E-mail: olgazh@dartmouth.edu; Tel: (603) 646-8616

Research Paper

Keywords: exaptation, domestication, horizontal gene transfer, bacterium-virus co-evolution, bacteriophage

1 Abstract

Several bacterial and archaeal lineages produce nanostructures that morphologically resemble

small tailed viruses, but, unlike most viruses, contain apparently random pieces of the host

genome. Since these elements can deliver the packaged DNA to other cells, they were dubbed

Gene Transfer Agents (GTAs). Because many genes involved in GTA production have viral

homologs, it has been hypothesized that the GTA ancestor was a virus. Whether GTAs

represent an atypical virus, a defective virus, or a virus co-opted by the prokaryotes for some

function, remains to be elucidated. To evaluate these possibilities, we examined the distribution

and evolutionary histories of genes that encode a GTA in the α-proteobacterium *Rhodobacter*

capsulatus (RcGTA). We report that although homologs of many individual RcGTA genes are

abundant across bacteria and their viruses, RcGTA-like genomes are mainly found in one

subclade of α-proteobacteria. When compared to the viral homologs, genes of the RcGTA-like

genomes evolve significantly slower, and do not have higher %A+T nucleotides than their host

chromosomes. Moreover, they appear to reside in stable regions of the bacterial chromosomes

that are generally conserved across taxonomic orders. These findings argue against RcGTA

being an atypical or a defective virus. Our phylogenetic analyses suggest that RcGTA ancestor

likely originated in the lineage that gave rise to contemporary α-proteobacterial orders

Rhizobiales, Rhodobacterales, Caulobacterales, Parvularculales, and Sphingomonadales, and

since that time the RcGTA-like element has co-evolved with its host chromosomes. Such

evolutionary history is compatible with maintenance of these elements by bacteria due to some

selective advantage. As for many other prokaryotic traits, horizontal gene transfer played a

substantial role in the evolution of RcGTA-like elements, not only in shaping its genome

components within the orders, but also in occasional dissemination of RcGTA-like regions

across the orders and even to different bacterial phyla.

1. Introduction

Prokaryotes are hosts not only of their own genetic material, but also of mobile genetic elements, a broad class of entities that includes integrated viruses (prophages) (Frost et al. 2005). Traditionally, viruses are viewed as selfish genetic elements, but in some instances they can serve as vectors of horizontal gene transfer (HGT) (Touchon et al. 2017), an important driver of evolutionary success of many cellular lifeforms (Koonin 2016; Zhaxybayeva & Doolittle 2011). In other instances, prophage-like elements can provide immunity against other viruses (Canchaya et al. 2003), and therefore are beneficial to their host. While the evolutionary relationship between viruses and cellular lifeforms is still intensely debated, co-option of genes by both empires is likely frequent (Krupovic & Koonin 2017). For example, a few prokaryotic cellular functions that are associated with adaptations, such as cell-cell warfare, microbe-animal interactions, HGT, and response to environmental stress, are carried out using structures that resemble viral "heads" (e.g. McHugh et al. 2014; Sutter et al. 2008) or "tails" (e.g. Borgeaud et al. 2015; Shikuma et al. 2014). These structures appear to be widespread across archaea and bacteria (e.g. Böck et al. 2017; Sarris et al. 2014). In other instances, viral components appear to have originated from the cellular proteins (Krupovic & Koonin 2017). Such intertwined history of genes from cellular and non-cellular lifeforms resulted in a spectrum of genetic elements that, from the cellular host "point of view", span from parasitic to benign to beneficial.

One class of such genetic elements is yet to be placed in this spectrum. Several unrelated bacterial and archaeal lineages are observed to produce particles that morphologically resemble viruses, yet appear to carry random pieces of host DNA instead of their own viral genome (Berglund et al. 2009; Bertani 1999; Humphrey et al. 1997; Marrs 1974; Rapp & Wall 1987). These particles were shown to transfer DNA between cells through a transduction-like mechanism (Lang et al. 2012; McDaniel et al. 2010), and that's why they were

dubbed "Gene Transfer Agents" (GTAs) (Marrs 1974). Since the transferred DNA may benefit

the recipient cells, GTAs were postulated to be viruses that were "domesticated" by prokaryotes

(Bobay et al. 2014) and maintained by them as a mechanism of HGT (Lang et al. 2012). Yet,

the details of such co-option event as well as impact of the GTA-mediated HGT remain to be

deciphered, and alternatives, such as GTA being instead a defective or atypical virus, need to

be evaluated.

The laboratory studies of *Rhodobacter capsulatus*' GTA (or RcGTA for short) show that

this element neither has a typical genome of a bacterial virus nor acts like a typical lysogenic

virus. Its genome is distributed among five loci dispersed across the *R. capsulatus* chromosome

(Hynes et al. 2016 and **Figure 1A**). Seventeen of the genes are found in a single locus that has

a genomic architecture typical of a siphovirus (Lang & Beatty 2007). Most of the genes in the

locus encode proteins involved in head and tail morphogenesis of the RcGTA particle and are

thus referred to as the 'head-tail' cluster (after Lang et al. 2017). The remaining four loci contain

seven genes shown to be important for RcGTA production, release, and DNA transfer (Fogg et

al. 2012; Hynes et al. 2012, 2016; Westbye et al. 2015). Even if RcGTA would preferentially

package the regions of DNA that correspond to its own genome, the small head size of RcGTA

particle can accommodate only ~1/5 of the genome (Lang et al. 2017), and therefore RcGTA

can propagate itself only with the division of the host cell. Expression of the RcGTA genes, as

well as production and release of RcGTA particles can be triggered by phosphate concentration

(Leung et al. 2010; Westbye et al. 2013), salinity (McDaniel et al. 2012), and quorum sensing

(Brimacombe et al. 2013; Schaefer et al. 2002). The latter is regulated via CckA-ChpT-CtrA

phosphorelay (Leung et al. 2012; Mercer & Lang 2014), a widely used bacterial signaling

network that also controls cell cycle (Chen & Stephens 2007; Mann et al. 2016) and flagellar

motility (Zan et al. 2013). These observations suggest that RcGTA is under control of the

bacterial host and is well-integrated with the host's cellular systems. When did such integration

occur? Could RcGTA still represent a selfish genetic element? Through phylogenomic analyses of a much more extensive genomic data set, we show that this α-proteobacterial GTA is a virus-related element whose genome evolves differently from what is expected of a typical bacterial virus. We also infer that RcGTA-like element likely originated in an ancestor of an α-proteobacterial subclade that gave rise to at least five contemporary taxonomic orders. Since that time, the genomes of these elements were shaped by both co-evolution with the bacterial hosts and HGT.

2. Methods

Detection of RcGTA homologs in viruses and bacteria

Twenty-four genes from the RcGTA genome and their homologs from *Rhodobacterales* were used as queries in BLASTP (E-value < 0.001; query and subject overlap by at least 60% of their length) and PSI-BLAST searches (E-value < 0.001; query and subject overlap by at least 40% of their length; maximum of 6 iterations) against viral and bacterial databases using BLAST v. 2.2.30+ (Altschul et al. 1997). The viral database consisted of 1,783 genomes of dsDNA bacterial viruses extracted from the RefSeq database (release 76; last accessed on July 1, 2016) using "taxid=35237 AND host=bacteria" as a query. Bacterial RefSeq database (release 76; last accessed on June 29, 2016) and 255 completely sequenced α-proteobacterial genomes (**Supplementary Table S1**; genomes downloaded from ftp://ftp.ncbi.nlm.nih.gov/genomes/Bacteria/ on November 1, 2015) served as two bacterial databases. The detected RefSeq homologs were mapped to their source genomes using "release76.AutonomousProtein2Genomic.gz" file (ftp.ncbi.nlm.nih.gov/refseq/release/release-catalog/, last accessed on June 29, 2016).

Assignment of genes to viral gene families

All annotated protein-coding genes from 255 α-proteobacterial genomes were used as queries in BLASTP searches (E-value < 0.001; query and subject overlap by at least 40% of their length) against the POG (Phage Orhologous Group) database (Kristensen et al. 2013), which was downloaded from ftp://ftp.ncbi.nlm.nih.gov/pub/kristensen/thousandgenomespogs/ blastdb/ in February 2016. The genes were assigned to a POG family based on the POG affiliation of the top-scoring BLASTP match.

Identification of RcGTA homologs adjacency in the bacterial genomes

Relative positions of RcGTA homologs in each genome were calculated from genome annotations in the GenBank feature tables downloaded from ftp.ncbi.nlm.nih.gov/genomes/all/ on June 29, 2016. Initially, a region of the genome was classified as putative RcGTA-like region if it had at least one RcGTA homolog. The identified regions were merged if two adjacent regions were separated by < 15 open reading frames (ORFs). If the resulting merged region contained at least nine RcGTA homologs, it was designated as a large cluster (LC). The remaining regions were labeled as small clusters (SC).

Examination of viral characteristics of RcGTA-like genes and regions

Prophage-like regions in 255 α-proteobacterial genomes were detected using PhiSpy v. 2.3 (Akhter et al. 2012) with "-n=5" (at least 5 consecutive prophage genes), "-w=30" (a window size of 30 ORFs), and "-t=0" ("Generic Test Set") settings. An RcGTA-like region was classified

as a putative prophage if it overlapped with a PhiSpy-detected prophage region.

Presence of viral integrases was assessed within all prophage-like and RcGTA-like

regions, as well as among 5 ORFs upstream and downstream of the regions. The integrase

homologs were identified using the hmmsearch program of the HMMER package v. 3.1b2 (Eddy

1998; E-value < 10^{-5}), with HMM profiles of integrase genes from the pVOG database 2016

update (Grazziotin et al. 2017) as queries (VOG# 221, 275, 375, 944, 2142, 2405, 2773, 3344,

3995, 4650, 5717, 6225, 6237, 6282, 6466, 7017, 7518, 8218, 8244, and 10948) and 255 α-

proteobacterial genomes as the database.

For both large and small clusters, %G+C of the corresponding nucleotide sequence

(defined as DNA sequence between the first and last nucleotide of the first and last RcGTA

homolog of the region, respectively) was calculated using in-house Python scripts. The relative

difference in GC content of the RcGTA-like region and the whole host genome was calculated

by taking the difference in their %G+C and dividing it by the %G+C of the host genome. This

correction allowed us to compare changes in the GC content of the RcGTA-like regions among

genomes with variable GC contents. To account for possible heterogeneity of GC content within

each genome, 100 genomic regions of the same size were randomly selected from the genome,

their relative GC content calculated as described above and compared to that of the RcGTA-like

region(s) using pairwise t-test.

Pairwise phylogenetic distances (PPD) among RcGTA homologs in viruses, and large

and small clusters were calculated in RAxML v. 8.1.3 (Stamatakis 2014) using '-f x' and '-m

PROTGAMMAAUTO' options. The latter choice selects the best amino acid substitution model

and uses Γ distribution with four discrete rate categories to correct for among site rate variation

(Yang 1994).

Identification of gene families

Sequences of protein-coding genes in 255 α-proteobacterial genomes were subjected to all-against-all BLASTP searches (v. 2.2.30+, E-value < 0.0001), in which reciprocal top-scoring BLASTP matches were retained. The bit scores of the matches were converted to pairwise distances, which were used to form gene families via Markov clustering with the inflation parameter set to 1.2 (van Dongen 2000), as implemented in OrthoMCL v. 2.0.9 (Li 2003). The gene presence/absence in the resulting 33,048 gene families was used to as a proxy of gene family conservation within and across α-proteobacteria.

Characterization of immediate gene neighborhoods of small and large clusters

ORFs without detectable similarity to RcGTA genes, but found immediately upstream, downstream, or within 'head-tail' cluster, were classified based on (a) conservation within α-proteobacterial genomes and (b) potential viral origin. The proportion of α-proteobacterial genomes that have an ORF homolog was used as a proxy for conservation. The ORF was designated as viral if it was assigned to a POG.

Reference phylogeny of α-proteobacteria

From the list of 104 genes families used to reconstruct α-proteobacterial phylogeny (Williams et al. 2007), 99 gene families that are present in 80% of the 255 α-proteobacterial genomes were selected (**Supplementary Table S2**). For each gene family, homologs from *Geobacter sulfurreducens* PCA, *Ralstonia solanacearum* GMI1000, *Chromobacterium violaceum* ATCC 12472, *Xanthomonas axonopodis* pv. citri str. 306*, Pseudomonas aeruginosa*

PAO1*, Virbrio vulnificus* YJ016*, Pasteurella multocida* subsp *multocida* str Pm70, *Escherichia*

coli str. K12 substr. DH10B were added as an outgroup. The amino acid sequences of the

resulting gene set were aligned using MUSCLE v. 3.8.31 (Edgar 2004). The best substitution

model (listed in **Supplementary Table S2**) was selected using *ProteinModelSelection.pl* script

downloaded from http://sco.h-

its.org/exelixis/resource/download/software/ProteinModelSelection.pl in August 2016. Among

site rate variation was modeled using Γ distribution with four rate categories (Yang 1994). All

gene sets were combined into a single dataset with 99 data partitions (one per gene.) The

maximum likelihood phylogenetic tree was reconstructed in RAxML v. 8.1.3 (Stamatakis 2014),

using the best substitution model for each partition and 20 independent tree space searches.

One hundred bootstrap samples were analyzed as described above, and the bootstrap support

values > 60% were mapped to the maximum likelihood tree. Reference phylogeny of only LC-

containing taxa was obtained by pruning taxa that only have SCs.

Phylogenetic analysis of the large-cluster gene families

Since phylogenetic histories of many RcGTA homologs are poorly resolved in

Rhodobacterales (Hynes et al. 2016), amino acid sequences of LC gene families were

combined into one "LC-locus" dataset in order to increase the number of phylogenetically

informative sites. To ensure that such concatenation does not result in a mixture of genes with

different evolutionary histories, phylogenetic trees reconstructed from the alignments of

individual LC genes and of the concatenated LC-locus were compared. Amino acid sequences

of gene families consisting only of LC homologs from α-proteobacterial genomes were aligned

using MUSCLE v. 3.8.31 (Edgar 2004). The individual gene family alignments were

concatenated into the "LC-locus" alignment. Since many LCs do not have homologs of all

RcGTA 'head-tail' genes, in the concatenated alignment the absences were designated as

missing data (Stamatakis 2014). Additionally, "taxa-matched" LC-locus alignments (*i.e.*

concatenated alignments pruned to contain taxa found only in a specific LC gene family) were

created. Maximum likelihood trees were reconstructed from the concatenated and individual LC

gene alignments, as well as from the taxa-matched LC-locus alignments, using RAxML v. 8.2.9

(Stamatakis 2014) under the LG+Γ substitution model (Le & Gascuel 2008; Yang 1994). The LG

model was determined as the best substitution model for each individual gene set using the

PROTGAMMAAUTO option of RAxML v. 8.2.9 (Stamatakis 2014). For each LC gene,

consensus trees were reconstructed from 10 bootstrap samples of taxa-matched LC-locus

alignments. These consensus trees were compared to 100 bootstrap sample trees for the

corresponding LC gene. The congruence between the phylogenies was measured using relative

"Tree Certainty All" (TCA) values (Salichos et al. 2014), as implemented in RAxML v. 8.2.9

(Stamatakis 2014). The TCA values were classified into "strongly conflicting" ($-1.0 \leq$ TCA $\leq -$

0.7), "moderately conflicting" ($-0.7 <$ TCA < 0.7) and "not conflicting" ($0.7 \leq$ TCA ≤ 1.0)

categories. None of the individual LC gene phylogenies had strongly conflicting TCA values,

although 6 out of 17 LC gene phylogenies had moderately conflicting values (**Supplementary**

Table S3). Since an uncertain position of just one taxon can affect support values of multiple

bipartitions (Aberer et al. 2013) and TCA values calculated from them, a Site Specific

Placement Bias (SSPB) analysis (Berger et al. 2011) was carried out, as implemented in

RAxML v. 8.2.9 (Stamatakis 2014). The LC-locus phylogeny and the LC-locus alignment were

used as the reference tree and the input sequence alignment, respectively. The sliding window

size was set to 100 amino acids. The SSPB analysis revealed that for 15 of the 17 genes only

three nodes, on average, separate the optimal positions of each taxon in the gene and LC-locus

phylogenies, indicating largely compatible evolutionary histories of individual LC genes and of

their combination (**Supplementary Figure S1**). Consequently, we decided to use the combined,

taxa-matched LC-locus alignment for further phylogenetic analyses.

To assess the similarity of the evolutionary histories of LC loci and of their hosts, the

topologies of the reference α-proteobacterial tree were compared to 100 bootstrap sample trees

reconstructed from the taxa-matched LC-locus alignment assembled as described above. The

congruence was quantified using Internode Certainty (IC) values (Salichos et al. 2014), as

implemented in RAxML v. 8.2.9 (Stamatakis 2014). The IC values were classified into "strongly

conflicting" ($-1.0 \leq IC \leq -0.7$), "moderately conflicting" ($-0.7 < IC < 0.7$) and "not conflicting" (0.7

$\leq IC \leq 1.0$) categories.

Conservation of gene neighborhoods across phylogenetic distance

Forty ORFs upstream and downstream of the ATP synthase operon, the ribosomal

protein operon, and the LCs were extracted from 87 α-proteobacterial genomes with at least

one LC. The same procedure was performed for a transposase from IS3/IS911 family, which

was detected in 18 of the 87 genomes. Each ORF was assigned to a gene family, and the

pairwise conservation of the corresponding flanking regions was calculated as a proportion of

gene families shared between a pair of genomes. PPDs of 16S rRNA genes were used as a

proxy for time t. 16S rRNA genes were aligned against the GreenGenes database (Desantis et

al. 2006; last accessed in August 2016) using mothur v. 1.35.1 (Schloss et al. 2009). The PPDs

were calculated from the alignment using RAxML v. 8.1.3 (Stamatakis 2014) under GTR+Γ

substitution model. Based on *model 3* from Rocha (2006), the gene neighborhood decay was

modeled as p^t, where p is the probability of the 40 ORFs to remain in the same genomic region

and t is time. This model assumes that every gene has equal probability of separating from the

region. The parameter p was estimated, and the fit between the model and data was assessed

using non-linear least squares method (Bates & Watts 1988), as implemented in the *nls* R

function.

Examination of genes that flank RcGTA-like clusters in non-α-proteobacterial genomes

Amino acid sequences of five ORFs upstream and downstream of the RcGTA-like

clusters in the genomes of actinobacteria *Streptomyces purpurogeneiscleroticus* NRRL B-2952

(GenBank accession number LGEI00000000.1;Ju et al. 2015) and *Asanoa ferruginea* NRRL B-

16430 (LGEJ00000000.1; Ju et al. 2015), a γ-proteobacterium *Pseudomonas bauzanensis*

W13Z2 (JFHS00000000.1; Wang et al. 2014), and a cyanobacterium *Scytonema millei*

VB511283 (JTJC00000000.1; Sen et al. 2015) were searched against the database of protein-

coding genes of 255 α-proteobacterial genomes using BLASTP (E-value < 0.001; query and

subject overlap by at least 60% of their length). Three genes that flank LCs in *P. bauzanensis*,

S. millei and a few α-proteobacteria (*kpdE*, *pbpC*, *dnaJ*) were used as queries to detect their

homologs in other cyanobacteria and γ-proteobacteria in BLASTP searches of γ-proteobacteria

and cyanobacteria in the *nr* database (https://blast.ncbi.nlm.nih.gov/Blast.cgi, last accessed on

October 31, 2016; E-value < 0.001; query and subject overlap by at least 60% of their length).

For each of the three gene sets, four homologs from each identified taxonomic group were

selected and combined with all available homologs from LC-containing α-proteobacteria. The

amino acid sequences of the combined data sets were aligned using MUSCLE v. 3.8.31 (Edgar

2004). Maximum likelihood trees were reconstructed using FastTree v. 2.1.8 (Price et al. 2010)

under JTT+Γ substitution model. The trees were visualized and annotated using EvolView (He

et al. 2016; http://evolgenius.info/evolview, last accessed on November 13, 2016).

Identification of CRISPR/*Cas* defense systems

Spacers of the Clustered Regularly Interspaced Short Palindromic Repeats (CRISPRs) were identified by scanning 255 α-proteobacterial genomes for both CRISPR leader sequences and repeat structures using CRISPRleader v. 1.0.2 (Alkhnbashi et al. 2016) and PILER-CR v. 1.06 (repeat length between 16 and 64; spacer length between 8 and 64; minimum array length of 3 repeats; minimum conservation of repeats 0.9; minimum repeat length ratio 0.9; minimum spacer length ratio 0.75) (Edgar 2007), respectively. CRISPR-associated proteins were identified using the hmmsearch program of the HMMER package v. 3.1b2 (Eddy 1998; E-value < 10^{-5}), with TIGRFAM families of the CRISPR-associated proteins as queries (TIGRFAM # 287, 372, 1573, 1587, 1596, 1863, 1865, 1868, 1869, 1876, 1907, 2562, 2589, 2590, 2593, 2621, 3158, 3637, 3638, 3639, 3640, 3641, 3983; Selengut et al. 2007), and protein-coding genes of 255 α-proteobacterial genomes as database. Genomes were designated to have putative functional CRISPR/*Cas* systems if both CRISPR spacer arrays and at least three CRISPR-associated proteins from the same CRISPR class (Makarova et al. 2015) were present. Spacer sequences from these putatively functional CRISPR systems were used as BLASTN (v. 2.4.0+; E-value < 10; task = tblastn-short) queries against all gene sequences in LCs.

Identification of decaying RcGTA 'head-tail' cluster homologs

The nucleotide sequences of LC and SC regions were extracted from each genome and translated in their entirety into all six reading frames. RcGTA 'head-tail' cluster genes were used as BLASTP (v. 2.4.0+, E-value ≤ 0.001) queries in searches against the database of translated LC and SC regions. Homologous sequences within LC and SC regions that did not match the annotated ORFs were designated as putatively decaying RcGTA genes.

1 3. Results

2 Many RcGTA genes share evolutionary history with viruses

Consistent with the earlier proposed hypothesis that RcGTA is an element originated

from a virus (Lang & Beatty 2000), 14 out of 24 RcGTA genes can be assigned to a POG

(**Figure 1A** and **Supplementary Table S4**), and 18 out of the 24 genes have at least one

readily detectable homolog in available genomes of *bona fide* bacterial viruses (**Figure 1B**).

However, none of the bacterial viruses in RefSeq database contained a substantial number of

the RcGTA homologs (at most eight of them in *Rhizobium* phage 16-3 [GenBank Acc. No.

DQ500118.1]), and no α-proteobacterial genomes with CRISPR systems contained significant

matches between CRISPR spacers and RcGTA genome, suggesting that the presumed

progenitor virus is either extinct or remains unsampled. Collectively, RcGTA-like genes are

unevenly represented across viral genomes (**Figure 1B**). Among the five most abundant are

homologs of *rcc001683* (*g2*), *rcc001684* (*g3*), *rcc001686* (*g4*) and *rcc001687* (*g5*), which

encode terminase, portal protein, prohead protease, and major capsid protein, respectively.

Their widespread occurrence is not surprising. First, these genes correspond to the so-called

"viral hallmark genes", defined as genome replication and virion formation genes with a wide

distribution among diverse viral genomes (Iranzo et al. 2016; Koonin et al. 2006). Second, *g3*,

g4 and *g5* belong to HK97 family, and HK97-like genes are common in Caudovirales (Iranzo et

al. 2017) - one of the most abundant viral groups (Cobián Güemes et al. 2016). However, 13 of

the RcGTA genes are either sparsely represented (found in 10 or fewer viral genomes; 7 genes)

or not detected at all (6 genes) in viruses. These genes may be fast-evolving viral genes,

auxiliary (non-hallmark) viral genes, which could be specific to the viral lineage that gave rise to

RcGTA, non-viral (cellular) genes, or simply be a result of undetected homology due to the

paucity of viral genomes in GenBank. The latter possibility cannot be ignored, especially given

that (i) three of the 13 genes are assigned to a POG (**Figure 1A**), and therefore are likely of viral

origin, and (ii) only 94 of the 1,783 screened bacterial viruses (5%) have α-proteobacteria as

their known host. Since about half of bacterial genomes are estimated to contain at least one

prophage in their genome (Touchon et al. 2016), inclusion of RcGTA homologs from bacterial

genomes could reduce the possibility of undetected homology due to lack of available viral

genomes. Therefore, we looked for RcGTA homologs in all bacterial RefSeq records, and in 255

completely sequenced α-proteobacterial genomes in particular (**Supplementary Table S1**).

Homologs of RcGTA genes are abundant in bacteria, but RcGTA-

like genomes are mainly found in class α-proteobacteria

Every RcGTA gene has at least 198 homologs in RefSeq database (**Figure 1C**), and

therefore even the 13 "rare" genes are relatively abundant in bacterial genomes. Moreover, six

of the 13 genes (*rcc00171*, *rcc00556*, *rcc01688* [*g6*], *rcc01692* [*g10*], *rcc001695* [*g12*]*, and

rcc01866), including three assigned to POGs, are found in more than 1000 bacterial genomes

(**Figure 1C**). Therefore, it is unlikely that the 13 rare genes are auxiliary viral genes specific to

the viral lineage that gave rise to RcGTA. However, a larger number of genomes of α-

proteobacterial viruses is needed to identify if these genes are of viral or cellular origin.

Collectively, homologs of RcGTA genes are found within 13 bacterial phyla (**Figure 1C**

and **Supplementary Figure S2a**). Interestingly, 84% and 35% of the identified homologs

belong to the phylum Proteobacteria and to its class α-proteobacteria, respectively.

Furthermore, many α-proteobacterial genomes contain more 'head-tail' RcGTA homologs than

any available viral genome, and these homologs are often clustered (**Supplementary Figure**

S2b). In contrast, most other taxa have only few RcGTA homologs that are scattered across the

genome (**Supplementary Figures S2b** and **S2c**). Since in the RcGTA genome the 'head-tail'

genes are in one locus (**Figure 1A**) and are presumed to be co-transcribed (Kuchinski et al.

2016), functional GTAs of this family might be restricted mainly to α-proteobacteria.

Within α-proteobacteria, homologs of RcGTA genes are detectable in 197 out of 255

examined genomes (77%). These homologs are distributed across eight α-proteobacterial

orders with at least one available genome and two unclassified α-proteobacterial genera

(**Figure 2**). While 13 out of 17 'head-tail' genes are abundant across α-proteobacteria (and

across Bacteria in general), only two of the genes from the remaining four loci (*rcc00555* and

rcc01866) are frequently detected in bacterial genomes (**Figure 1C** and **Figure 2**). Since within

Rhodobacterales genes outside of 'head-tail' cluster evolve faster than their 'head-tail'

counterparts (Hynes et al. 2016), we conjecture that many α-proteobacterial homologs of the

former genes were not detected in our searches. Therefore, for the subsequent investigations

we focused mostly on the analyses of the 'head-tail' locus genes.

Two distinct classes of RcGTA 'head-tail' homologs are present in

α-proteobacteria

Within 187 α-proteobacterial genomes with at least one 'head-tail' cluster homolog, the

RcGTA-like genes are dispersed across 474 genomic regions, 245 of which contain only 1-2

RcGTA homologs (**Figure 3**). Due to significant similarity between RcGTA and genuine viral

genes, some of these regions may belong to prophages unrelated to RcGTA. Indeed, 261 out of

474 regions (55%) overlap with putative prophages identified using PhiSpy (Akhter et al. 2012)

(**Supplementary Figure S3**). However, PhiSpy also classified the RcGTA as a prophage,

suggesting that the predictions may include other false positives. Of 889 PhiSpy-predicted

prophage regions within 255 α-proteobacterial genomes, only 535 (60%) are associated with an

integrase gene - a gene expected to be present in a functional prophage but not in a GTA.

RcGTA-like regions are overrepresented among the predicted prophages without an integrase

gene (234 out of 354 [66%] vs. 27 out of 535 [5%]), hinting that PhiSpy might not be able to

distinguish GTAs from genuine prophages. Also, PhiSpy did not detect small RcGTA-like

regions due to the program's requirement of at least five consecutive genes with "viral"

functional annotations in a window of 30 genes. Therefore, we examined the RcGTA-like

regions, and genes within them, for additional characteristics associated with viral genes –

skewed nucleotide composition (Rocha & Danchin 2002) and faster substitution rates (Drake

1999; Paterson et al. 2010) – as well as the presence of neighboring viral genes.

Genes of viral origin generally have lower fraction of Gs and Cs (GC content) than their

host genomes (Rocha & Danchin 2002). However, we do not observe this trend for the RcGTA-

like regions. GC content of the regions with less than nine RcGTA homologs (hereafter referred

as small clusters, or SC) is, on average, equivalent to the GC content in the host (median of

percent of relative change = -0.1%; pairwise Wilcoxon test; p-value=0.60), albeit with substantial

variability across examined genomes (**Figure 4**). On the other hand, GC content of the regions

with at least nine RcGTA homologs (hereafter referred as large clusters, or LC) is consistently,

and in most cases significantly, higher than the GC content of their host (median of percent of

relative change = 6.57%; pairwise t test; p-value < 0.001) (**Figure 4**). The elevated GC content

of LCs also persists when compared to randomly sampled regions of the host genome of

equivalent size (pairwise t-test; *p-value* <0.001). These observations prompted us to examine

the features of LCs and SCs separately.

Using the pairwise phylogenetic distances (PPD) as a proxy for substitution rates, we

found that LC genes evolve significantly slower than both their SC (all of the 13 possible

pairwise comparisons; Wilcoxon tests; p-values <0.0001) and viral homologs (14 out of 15

possible comparisons; Wilcoxon tests; p-values <0.005) (**Supplementary Figure S4** and

Supplementary Table S5). While many SC genes also evolve significantly slower than their

viral homologs (six out of 10 possible comparisons; Wilcoxon tests; *P*-values <0.005), the 25-75

percentile range for PPDs of SC genes often overlapped with that of the viral homologs

(**Supplementary Figure S4**). Hence, SCs are more virus-like than LCs. This conjecture is

supported by two additional observations. First, SCs are more frequently associated with

unrelated viral genes than LCs (76 vs. 56%; **Supplementary Figure S3**). In particular, SCs are

more likely to reside in a vicinity of an integrase gene than LCs (45 out of 383 SCs vs. 1 out of

91 LCs). In some cases, SCs likely belong to viral elements that are clearly not GTAs: for

example, in all analyzed genomes of the *Rickettsiales'* genera *Anaplasma* and *Ehrlichia* a

singleton homolog of *rcc001695* (g12) is found within 1 kb of *vir*B homologs, which in

Anaplasma phagocytophilum belong to a virus-derived type IV secretion system (Al-Khedery et

al. 2012). Second, SCs have very little conservation of flanking genes (**Supplementary Figure

S5)**, while in LCs flanking genes are conserved within orders and sometimes even across larger

phylogenetic distances (**Supplementary Figure S6)**.

Taken together, we hypothesize that large and small clusters represent genomic regions

under different selective pressures in the host chromosome and may have separate

evolutionary histories. We further hypothesize that most SCs are viral elements unrelated to

RcGTA and only LCs represent RcGTA-like elements.

RcGTA-like element was likely absent in the last common

ancestor of α-proteobacteria

In what lineage did the RcGTA-like element originate? How did it propagate across α-

proteobacteria? To address these questions, we examined the phylogenetic history of LCs.

Within α-proteobacteria, LCs are found only within a monophyletic clade defined by node 1 in

Figure 5 (hereafter referred as Clade 1), which consists of all analyzed representatives from the

orders *Rhizobiales, Caulobacterales*, *Rhodobacterales, Parvularculales* and

Sphingomonadales. Two deeper branching α-proteobacterial clades, which include

Rickettsiales, *Rhodospirillales,* and yet unclassified *Micavibrio* spp. and SAR 116 cluster

bacteria, contain only SCs (**Figure 5**). The largest of these SCs is made up of five genes, but

most are singletons (**Figures 2** and **3**). RcGTA homologs are completely absent from

Pelagibacterales (formerly SAR11) and the basal α-proteobacterial order *Magnetococcales*

(**Figure 5**), although only five genomes from these two groups were available for the analyses.

It is possible that SCs represent decaying remnants of the previously functional GTAs.

Bacterial genomes do not generally contain many pseudogenes due to genome streamlining

(Lawrence et al. 2001). However, if pieces of RcGTA homologs that no longer form an ORF

remain within the regions, they may still be detectable by similarity searches. Indeed, putative

pseudogenes of RcGTA-like genes were detected in both large and small clusters, but only 35%

of them reside in SCs (**Supplementary Figures S2, S5** and **S6**). Furthermore, only 3 of 199

SCs found in taxa outside of Clade 1 contain detectable putative pseudogenes of the RcGTA

'head-tail' genes (**Supplementary Figure 3**), suggesting that SCs outside of Clade 1 are

probably unrelated to RcGTA.

Taking into account the observation that the fraction of genomes with no detectable

RcGTA homologs is higher in taxa outside than inside of Clade 1 - 30% versus 15% of the

genomes, respectively, - we hypothesize that the RcGTA-like element originate at the earliest

on the branch leading to the last common ancestor of Clade 1 and not in the last common

ancestor of α-proteobacteria (Clade 2 in **Figure 5**), as was previously suggested (Lang et al.

2002; Lang & Beatty 2007).

1 Evolution of the RcGTA-like genome: Vertically Inherited or

2 Horizontally Exchanged?

Did the RcGTA-like element appear on the branch leading to the last common ancestor

of Clade 1, or did it spread across Clade 1 via HGT? Lang & Beatty (2007) inferred that RcGTA-

like element had evolved vertically within α-proteobacteria, although only few genomes were

available at the time. Within *Rhodobacterales* RcGTA-like 'head-tail' clusters are also predicted

to be vertically inherited, but phylogenetic trees of many individual genes were poorly resolved

(Hynes et al. 2016). In contrast, homologs of *rcc00171* were likely horizontally exchanged within

Rhodobacterales (Hynes et al. 2016). To investigate the extent of HGT in the evolution of the

RcGTA-like 'head-tail' genes across Clade 1, we reconstructed the phylogeny of LC genes and

compared it to the reference tree of α-proteobacteria represented by a set of conserved α-

proteobacterial genes.

Of 83 internal branches of the reference phylogeny, 42 are in disagreement with the

branching order of the LC-locus phylogeny (**Figure 6**). Thirty-one of the 42 conflicts (74%) are

within taxonomic orders, with 14 out of the 21 strongly supported conflicts limited to genera

(**Figure 6** and **Supplementary Figure S7**). Among the nine conflicts at deep branches, eight

are due to an alternative position of *Methylobacterium nodulans* ORS 2060 (**Figure 6** and

Supplementary Figure S7), which, in addition to grouping outside of its order *Rhizobiales,* has

five non-identical LCs (**Supplementary Figure S6**). Therefore, evolution of 'head-tail' locus is

likely impacted by HGT events, but most of them have occurred between closely related taxa.

Intriguingly, at least occasionally LCs have also been transferred across very large

evolutionary distances. While LCs are predominantly found in α-proteobacteria, the genomes of

the actinobacteria *Streptomyces purpurogeneiscleroticus* NRRL B-2952 and *Asanoa ferruginea*

NRRL B-16430, the cyanobacterium *Scytonema meliloti* VB511283, and the γ-proteobacterium

Pseudomonas bauzanensis W13Z2 also contain one LC each (**Supplementary Figures S2** and

S6). These four non- α-proteobacterial taxa branch within the Clade 1 in three separate

locations (**Figure 6**), indicating that these taxa likely acquired the LCs from the α-proteobacteria

at least three times independently. Moreover, genes immediately upstream and downstream of

S. millei and *P. bauzanensis'* LCs are homologous to the genes that flank LCs in their

respective α-proteobacterial sister taxa (**Supplementary Figure S6**). These flanking genes are

also of α-proteobacterial origin (**Supplementary Figure S8**), and hence were likely acquired in

the same HGT events.

The duration of RcGTA-like elements' association with bacterial genomes can also be

gauged from the stability of the host chromosome regions that surround LCs. For example,

highly mobile genes, such as transposable elements and some prophages, are often found in

genomic islands (Juhas et al. 2009) - dynamic genomic regions that exhibit poor gene synteny

even across short phylogenetic distances (Rodriguez-Valera et al. 2016). On the other hand,

gene synteny of stable regions is preserved across larger evolutionary distances, although it is

still subject to decay due to gene rearrangements (Brilli et al. 2013; Rocha 2006). LCs found in

the same taxonomic order are generally flanked by the same, conserved α-proteobacterial

genes (**Supplementary Figure S6**), hinting that they might be part of stable regions of bacterial

chromosomes and not within genomic islands. We modeled gene order decay upstream and

downstream of LC loci as a function of phylogenetic distance (Rocha 2006) and compared the

conservation of genes flanking LCs to that of a transposable element from IS911/IS3 family and

two operons of conserved housekeeping genes, the ribosomal protein operon and the ATP

synthase operon. The IS911/IS3 element can undergo non-targeted integration (Chandler et al.

2015), and thus, genes adjoining these transposons are not expected to be homologous even

across closely related taxa. In contrast, gene order near presumably non-mobile housekeeping

operons are expected to be much more conserved across time. Indeed, we found that the

regions surrounding the transposable element are too variable to even fit the model, while the

decay of gene synteny near LCs is similar to that of the two operons (**Supplementary Figure**

S9). Therefore, it is unlikely that RcGTA-like elements reside in dynamic regions of their host

genomes.

Although sequence divergence of the remaining four loci of the RcGTA genome makes it

difficult to track their evolutionary history, in 66 α-proteobacterial genomes with detected

homologs of *rcc00171* and *rcc00555*, 22 LCs from Rhizobiales and Rhodobacterales have

rcc00171 homologs at their 3' end and 6 of the 22 LCs additionally have *rcc00555* immediately

downstream of an *rcc00171* homolog (**Supplementary Figure S6**). This anecdotal evidence

suggests that in the past RcGTA-like elements may have carried at least some of the genes in

one locus with the 'head-tail' genes, and that the division of the GTA genome into multiple loci

may have happened after the bacterium-GTA association was already established.

Taken together, our analyses suggest a complex evolutionary history of the RcGTA-like

elements. The congruence of the reference and the LC-locus phylogenies at taxonomic order

and family levels and conservation of LC-flanking genes within orders suggest that RcGTA-like

element was likely acquired before the contemporary orders within Clade 1 have diversified, and

since that time the element co-existed with the clade members. However, the congruence does

not automatically equate to vertical inheritance, since even genes highly conserved across α-

proteobacteria are not immune to HGT and thus the reference phylogeny may not represent the

strictly vertical history of α-proteobacteria (Gogarten et al. 2002; Andam et al. 2010). Indeed,

strong phylogenetic conflicts between the reference and LC locus phylogenies of Clade 1 at

both deep and shallow branches, and presence of LCs in several taxa outside of the α-

proteobacteria indicate that HGT has been shaping the evolution of RcGTA-like element to a

non-negligible degree. Since the majority of the strongly supported conflicts are found within

genera, HGT probably played a larger role in the recent evolution of RcGTA-like elements than

in their dissemination across Clade 1.

4. Discussion

Our survey of RcGTA homologs in the genomes of bacteria and bacterial viruses provide new insights into fascinatingly intertwined evolutionary histories of virus-like elements and their bacterial hosts. Although we infer that the RcGTA-like element is not as ancient as the class α-proteobacteria itself, RcGTA homologs are ubiquitous across a clade that spans multiple α-proteobacterial orders (Clade 1 in **Figure 5**) and, according to molecular clock estimates, has originated between 777 Mya (Luo et al. 2013) and 1710 Mya (with a 95% CI of 1977-1402 Ma; Gregory Fournier, pers. comm.) Therefore, either RcGTA-like element has been associated with a group of α-proteobacteria for hundreds of millions of years, or it is a vestige of a very successful virus capable of inter-species and inter-order infections. The latter scenario is supported by experimentally observed long-range HGT via GTAs produced by *Roseovirus nubinhibens* ISM (Mcdaniel et al. 2010). However, lack of CRISPR repeats that match RcGTA genome, absence of closely-related contemporary viruses in GenBank, and order-level congruence of the evolutionary histories of RcGTA and conserved α-proteobacterial genes point against the GTA being derived from a broad host-range virus that recently invaded this clade. Instead, our data are more compatible with a long-term association between a GTA and its bacterial host.

Over this time, the GTA genome has gone through extensive changes, as hinted by the putative HGT events within taxonomic orders, presence of likely pseudogenized RcGTA homologs within a few putative GTAs, and division of the RcGTA genome into multiple loci. Could these modifications simply mean that this RcGTA-like element is just a defective prophage (Redfield 2001; Solioz & Marrs 1977) not yet purged from the genomes? Since bacterial genomes are generally under deletion bias of unnecessary DNA (Mira et al. 2001),

conservation of "junk" regions across over millions of years is unexpected. Moreover, given the

universal mutation bias towards AT-rich DNA (Hershberg & Petrov 2010), over such prolonged

time we would expect the defective prophage region to become more AT-rich than the host

DNA, and to accumulate mutations evenly across non-synonymous and synonymous sites of

ORFs (Andersson & Andersson 2001). Yet, genomic regions that encode LCs are not AT-richer

than the rest of the host chromosomes, and, at least across *Rhodobacterales*, RcGTA

homologs do not exhibit increased rates of non-synonymous substitutions expected of

pseudogenes (Lang et al. 2012), suggesting that LCs are unlikely to be decaying prophages.

Surprisingly, RcGTA-like gene clusters have a significantly elevated % G+C than

chromosomes of their hosts. Evolution of GC content and its variation across and within the

prokaryotic genomes is an unsolved puzzle, with many possibilities invoked to explain it

(Hildebrand et al. 2010; Lassalle et al. 2015; Rocha & Feil 2010). In one hypothesis, AT-

richness of highly expressed genes is explained by accumulation of additional mutations during

the single-stranded DNA state required for transcription (Hildebrand et al. 2010). Since RcGTA-

producing cells lyse and thus leave no progeny, the GTA genes are never expressed by cells

that reproduce. Hence, it is tempting to speculate that selection for reduced gene expression

may have played a role in driving the GC content of the 'head-tail' locus higher than that of the

expressed host genes. Future testing of this hypothesis and exploring other possible

explanations are necessary to account for the aberrant GC content of the RcGTA-like regions

within Clade 1 genomes.

But if RcGTA-like elements are indeed functional, why do we observe putatively

pseudogenized and apparently incomplete gene repertoires of many LCs when compared to the

RcGTA genome? Perhaps, genomes of these elements are also split into different pieces, have

some components replaced from sources unrelated to the RcGTA, or have evolved to a

different functionality than RcGTA. For example, GTA gene expression and GTA particle

release was demonstrated for *Ruegeria pomeroyi* DSS-3 (Biers et al. 2008) and *Rhodovulum*

sulfidophilum (Nagao et al. 2015), suggesting that GTA is functional in these lineages. However,

the *R. pomeroyi* and *R. sulfidophilum* genome contain no detectable homologs of *g1*, and this

gene is essential for RcGTA production in *R. capsulatus* (Hynes et al. 2016). Therefore, even

presumably "incomplete" LCs may encode functional GTAs – a conjecture that awaits

experimental demonstrations.

With the evidence pointing against the *decaying* or a *selfish* nature of the RcGTA-like

elements, and given that cells that produce RcGTA lyse (Westbye et al. 2013) and, therefore,

the selection for the maintenance of RcGTA cannot occur at a level of an organism, earlier

suggestions that GTA maintenance is due to some *population*-level benefits (Lang et al. 2012)

remain most plausible explanation of persistence of RcGTA-like elements in the Clade 1

genomes. However, specific advantage(s) associated with acquisition of ~4kb pieces of random

DNA via GTA remain to be elucidated. Among the proposed drivers of GTA maintenance are

effective exchange of useful genes among the population of heterogeneous cells (Smillie et al.

2011), improved response to environmental stressors, such as DNA damage (Marrs et al. 1977;

Brimacombe et al. 2015) and nutritional deficits (Lang et al. 2012). Possibilities unrelated to

GTA-mediated DNA delivery, such as protection of the host population against infection by other

viruses via the superinfection immunity mechanism (Díaz-Muñoz 2017), and increase of the

host population resilience against various environmental stressors such as antibiotics (Wang et

al. 2010), should also be considered. Perhaps, there is no single benefit associated with

RcGTA-like elements across all Clade 1 taxa. Instead, different lineages may experience

selection pressures unique to their ecological niches.

Interestingly, GTA-associated benefits have been proposed to drive evolutionary

innovations of cellular lifeforms that go beyond population level. For example, it has been

hypothesized that abundance and mixed ancestry of α-proteobacterial genes in eukaryotic

genomes is due to the delivery of such genes by RcGTA-like elements into proto-eukaryote

genome, and that such "seeding" facilitated the later integration of mitochondrial progenitor and

the proto-eukaryotic host (Richards & Archibald 2011). Our study, however, does not support

this hypothesis. First, based on the estimated age of the Clade 1 (777-1710 Mya) and the

appearance of eukaryotic, and hence mitochondria-containing, organisms in the fossil record

(1600 and possibly 1800 Mya; Knoll 2014), the RcGTA-like system likely originated after the

eukaryogenesis took place. Second, α-proteobacterial lineage(s) suggested to have given rise

to mitochondrial progenitor (Brindefalk et al. 2011; Thrash et al. 2011; Wang & Wu 2015)

appear to lack LCs and therefore are unlikely to have had RcGTA-like elements. In another

intriguing hypothesis, an unrelated GTA in *Bartonella* has been proposed to have facilitated

adaptive radiation within the genus (Guy et al. 2013). Perhaps, acquisition of the RcGTA-like

element by the last common ancestor of Clade 1 taxa represent a similar innovation that led to

diversification of this clade and success of its members in a variety of ecological settings that

span soil, freshwater, marine, waste water, wetland, and eukaryotic intracellular and

extracellular environments.

14   Despite the appearance that bacteria maintain GTA genes for the potential contribution

to their own evolutionary success, there is an undeniable shared evolutionary history between

RcGTA-like elements and *bona fide* viruses. Presence of homologs for most RcGTA-like genes

in viral genomes, as well as similarity in the organization of the 'head-tail' cluster and

corresponding region of a typical siphovirus genome (Huang et al. 2011), led to a prevailing

hypothesis that RcGTA originated in an initial co-option of a virus and evolved in subsequent

modification of the progenitor prophage genome via vertical descent and HGT (Lang et al.

2017). Our data are compatible with this scenario, and further suggest an even larger role of

HGT in the evolution of RcGTA-like elements than currently acknowledged, shaping GTAs both

within α-proteobacterial orders, but also at least occasionally transferring RcGTA-like regions

across the higher-order α-proteobacterial taxa and even to different phyla. Yet, there are hints of

even more entangled connection between RcGTA-like genes found in cellular and viral

genomes: a few viruses that infect *Paracoccus*, *Roseobacter*, *Caulobacter*, and *Rhodobacter*

spp., and a yet unclassified marine *Rhizobiales* str. JL001 bacterium, carry RcGTA-like genes

that are phylogenetically placed within their cellular homologs (Zhan et al. 2016), suggesting

that some viruses acquired genes from the cellular RcGTA-like regions. Given recent

propositions of (*i*) cellular origin of even such "hallmark" viral genes like those encoding capsid

proteins (Krupovic & Koonin 2017), (*ii*) the possibly virus-independent origin of cellular

microcompartments that resemble viral structures (Bobik et al. 2015), and (*iii*) the potentially

primordial origin of type VI secretion system that resembles a bacteriophage tail (Böck et al.

2017), we should not exclude a possibility that genomic regions encoding RcGTA-like

nanostructures did not originate by a co-option of a prophage and its subsequent modification,

but instead are a mosaic originally assembled within bacterial genomes from the available

cellular and viral parts.

13 Acknowledgements

14 We would like to thank Andrew S. Lang, J. Thomas Beatty and Rosemary J. Redfield for

15 numerous stimulating discussions regarding GTA evolution.

17 **Conflict of Interest:** None declared

18 Funding

19 This work was supported by the National Science Foundation (NSF-DEB 1551674 to O.Z.); the

20 Simons Foundation (Investigator in Mathematical Modeling of Living Systems award 327936 to

21 O.Z.); the Neukom Institute CompX award to O.Z.; and Dartmouth start-up funds to O.Z.

Author Contributions

MS and OZ designed the study. MS and SMS performed the analyses. MS, SMS, and OZ wrote the manuscript. All authors have seen and approved the final manuscript.

Data Availability

Amino acid sequences of RcGTA homologs in bacterial and viral genomes; amino acid sequence alignments of RcGTA homologs from the LCs, SCs, and viruses; concatenated alignment of 99 conserved α-proteobacterial genes and RcGTA homologs from LCs; and discussed phylogenetic trees in Newick format are available via FigShare at DOI 10.6084/m9.figshare.5406733.

Figure Legends

Figure 1

RcGTA genome and distribution of its homologs in viral and bacterial genomes. Panel A. The RcGTA genome architecture. The RcGTA genome consists of 24 genes scattered across five loci in the *R. capsulatus* SB1003 genome. The genes are represented by arrows. The majority of the 'head-tail' cluster genes (*R. capsulatus* SB1003 locus tags *rcc01682-rcc01698*; also known as *g1-g15*) encode genes involved in head and tail morphogenesis, *rcc00171* (*tsp*) - a tail spike protein (Hynes et al. 2016), *rcc00555* and *rcc00556* - endolysin and holin (Fogg et al. 2012; Hynes et al. 2012; Westbye et al. 2013), *rcc01079* (*ghsA*) and *rcc01080* (*ghsB*) - a

head spike (Westbye et al. 2015), and *rcc01865* and *rcc01866* - putative regulatory elements

(Hynes et al. 2016). The GenBank functional annotations of the genes are listed in

Supplementary Table S4. Genes with similarity to gene families in the POG database

(Kristensen et al. 2013) are shaded and their assigned POG categories are listed in

Supplementary Table S4. Panel B. Presence of RcGTA gene homologs in 1,783 viral

genomes. Panel C. Presence of RcGTA gene homologs in 30,503 bacterial genomes. The

numbers above arrows represent the total number of detected homologs for each RcGTA gene,

while the percent of viral (**panel B**) bacterial (**panel C**) genomes in which these homologs are

found is depicted in shades of gray (see figure inset for scale). Genomic regions were visualized

using R package genoplotR (Guy et al. 2010).

Figure 2

Presence of RcGTA-like genes in α-proteobacterial genomes. Each line represents the

distribution of genes in the RcGTA genome within a taxonomic group. The number of surveyed

genomes in a taxonomic group is listed in parentheses. The shades of gray represent the

proportion of these genomes in which the RcGTA homologs are found (see figure inset).

Genomic regions were visualized using R package genoplotR (Guy et al. 2010).

Figure 3

Size variation of the RcGTA-like 'head-tail' clusters among α-proteobacterial genomes.

The number of homologs per genome (1-17) is shown on the X axis. The total number of

genomes in each size category is shown in parentheses on the X-axis labels. The genomes are

binned into taxonomic groups (arranged on Y axis). The diameter of each circle is scaled with

respect to the number of genomes (see inset for scale). Only 91 of 474 clusters (19%) have at

least nine RcGTA homologs. Sizes of RcGTA-like regions across several other bacterial

taxonomic groups are shown in **Supplementary Figure S2**.

Figure 4

Difference in GC content between clusters of RcGTA 'head-tail' gene homologs and their

host chromosome. The relative difference in GC content is calculated as $(GC_{GTA}-GC_{host})/GC_{host}$

(Y axis), and the data is presented by the number of RcGTA 'head-tail' homologs found in a

cluster (X axis). In the box-and-whisker plots the horizontal line marks the median; the boxes

extend from the 25th to 75th percentile; the whiskers encompass values at most 1.5 times away

from the 25th and 75th percentile range, and any value outside of the range is shown as a dot.

The vertical dashed line separates two groups of clusters: $\leq$ 8 genes, or small clusters (SC, in

blue) and $\geq$9 genes, or large clusters (LC, in orange). The inset depicts the combined data from

all small and large clusters.

Figure 5

Distribution of RcGTA homologs in α-proteobacteria. The phylogenetic tree is a schematic

version of the α-proteobacterial reference phylogeny, in which branches are collapsed at the

taxonomic order level. The number of taxa in each collapsed clade is shown in parentheses.

The tree is rooted using taxa from β-, γ- and δ-proteobacteria ("outgroup"). The non-collapsed

version of this tree is provided in **Supplementary Figure S3**. The nodes with at least 60%

bootstrap support are labeled with a black circle. Bars associated with each clade show the

relative proportion of genomes that contain at least one LC (orange), only SC(s) (blue), or no

detectable RcGTA homologs (gray). Nodes 1 and 2 define Clades 1 and 2 discussed in the text.

Twenty-three of 150 genomes within Clade 1 and 32 out of 105 genomes outside of Clade 1

have no detectable RcGTA homologs. While relationships among the α-proteobacterial orders

remain debated (Ferla et al. 2013; Gupta & Mok 2007; Lee et al. 2005; Luo 2014; Thrash et al.

2011; Williams et al. 2007), the alternative histories do not affect the inferences of this study

(data not shown). Scale bar, substitutions per site.

Figure 6

Comparison of the phylogenetic history of large clusters (right) to the reference α-

proteobacterial phylogeny (left). The branches of the reference tree are color-coded

according to the strength of conflict, as measured by an Internode Certainty (IC) value (Salichos

et al. 2014). IC values for all branches are shown in **Supplementary Figure S7.** The overall

number of branches with strongly conflicting IC values are shown as numbers (in pink) for

"deep" and "shallow" branches, as demarcated by a vertical dashed line. Fourteen of the 15

strong conflicts at shallow branches occur within genera, and five of the six strong conflicts at

deep branches are due to the different position of *Methylobacterium nodulans* ORS 2060. The

phylogenetic tree on the left is rooted with the same outgroup as in **Supplementary Figure S3**,

while the phylogenetic tree on the right should be considered unrooted. Underlined taxa names

represent four non- α-proteobacterial genomes that contain an LC (see also **Supplementary**

Figures S6 and **S9**). Taxa names in brackets represent lineages that branch within a taxonomic

order different from the one designated by the NCBI Taxonomy database (NCBI Resource

Coordinators 2017). The tree heights (*i.e.* the sum of all branch lengths of a phylogenetic tree)

of the LC-locus and reference phylogenies are 31.19 and 19.39 substitutions per site,

respectively, suggesting that LC locus evolves ~1.6 times faster than the conserved α-

proteobacterial genes. Scale bar, substitutions per site.

Supplementary Data

SupplementaryFigures.pdf: Supplementary Figures 1-9

SupplementaryTables.pdf: Supplementary Tables 1-5

References

Aberer, A. J., Krompass, D., & Stamatakis, A. (2013). 'Pruning rogue taxa improves phylogenetic accuracy: an efficient algorithm and web-service', *Systematic Biology*, 62/1: 162–6.

Akhter, S., Aziz, R. K., & Edwards, R. A. (2012). 'PhiSpy: a novel algorithm for finding prophages in bacterial genomes that combines similarity- and composition-based strategies', *Nucleic Acids Research*, 40/16: e126.

Al-Khedery, B., Lundgren, A. M., Stuen, S., et al. (2012). 'Structure of the type IV secretion system in different strains of *Anaplasma phagocytophilum*', *BMC Genomics*, 13: 678.

Alkhnbashi, O. S., Shah, S. A., Garrett, R. A., et al. (2016). 'Characterizing leader sequences of CRISPR loci', *Bioinformatics* , 32/17: i576–85.

Altschul, S. F., Madden, T. L., Schäffer, A. A., et al. (1997). 'Gapped BLAST and PSI-BLAST: a new generation of protein database search programs', *Nucleic Acids Research*, 25/17: 3389–402.

Andersson, J. O., & Andersson, S. G. (2001). 'Pseudogenes, junk DNA, and the dynamics of *Rickettsia* genomes', *Molecular Biology and Evolution*, 18/5: 829–39.

Bates, D. M., & Watts, D. G. (1988). 'Nonlinear regression analysis and its applications', Wiley&Sons, Inc.

Berger, S. A., Krompass, D., & Stamatakis, A. (2011). 'Performance, accuracy, and Web server for evolutionary placement of short sequence reads under maximum likelihood', *Systematic Biology*, 60/3: 291–302.

Berglund, E., Frank, A., Calteau, A., et al. (2009). 'Run-off replication of host-adaptability genes is associated with gene transfer agents in the genome of mouse-infecting *Bartonella grahamii*', *PLoS Genetics*, 5/7: e1000546.

Bertani, G. (1999). 'Transduction-like gene transfer in the methanogen *Methanococcus voltae*', *Journal of Bacteriology*, 181/10: 2992–3002.

Biers, E. J., Wang, K., Pennington, C., et al. (2008). 'Occurrence and expression of gene transfer agent genes in marine bacterioplankton', *Applied and Environmental Microbiology*,

74/10: 2933–9.

Bobay, L. M., Touchon, M., & Rocha, E. P. (2014). 'Pervasive domestication of defective prophages by bacteria', *Proceedings of the National Academy of Sciences of the United States of America*.111/33: 12127-32.

Bobik, T. A., Lehman, B. P., & Yeates, T. O. (2015). 'Bacterial microcompartments: widespread prokaryotic organelles for isolation and optimization of metabolic pathways', *Molecular Microbiology*, 98/2: 193–207.

Böck, D., Medeiros, J. M., Tsao, H.-F., et al. (2017). 'In situ architecture, function, and evolution of a contractile injection system', *Science*, 357/6352: 713–7.

Borgeaud, S., Metzger, L. C., Scrignari, T., et al. (2015). 'Bacterial evolution. The type VI secretion system of *Vibrio cholerae* fosters horizontal gene transfer', *Science*, 347/6217: 63–7.

Brilli, M., Liò, P., Lacroix, V., et al. (2013). 'Short and long-term genome stability analysis of prokaryotic genomes', *BMC Genomics*, 14: 309.

Brimacombe, C. A., Ding, H., Johnson, J. A., et al. (2015). 'Homologues of genetic transformation DNA import genes are required for *Rhodobacter capsulatus* gene transfer agent recipient capability regulated by the response regulator CtrA', *Journal of Bacteriology*, 197/16: 2653–63.

Brimacombe, C., Stevens, A., Jun, D., et al. (2013). 'Quorum-sensing regulation of a capsular polysaccharide receptor for the *Rhodobacter capsulatus* gene transfer agent (RcGTA)', *Molecular Microbiology*, 87/4: 802–17.

Brindefalk, B., Ettema, T. J. G., Viklund, J., et al. (2011). 'A phylometagenomic exploration of oceanic alphaproteobacteria reveals mitochondrial relatives unrelated to the SAR11 clade', *PLoS One*, 6/9: e24457.

Canchaya, C., Proux, C., Fournous, G., et al. (2003). 'Prophage genomics', *Microbiology and Molecular Biology Reviews*, 67/2: 238–76.

Chandler, M., Fayet, O., Rousseau, P., et al. (2015). 'Copy-out-Paste-in transposition of IS911: A major transposition pathway', *Microbiology Spectrum*, 3/4.

Chen, J. C., & Stephens, C. (2007). 'Bacterial cell cycle: completing the circuit', *Current Biology,* 17/6: R203–6.

Cobián Güemes, A. G., Youle, M., Cantú, V. A., et al. (2016). 'Viruses as winners in the game of life', *Annual Review of Virology*, 3/1: 197–214.

Desantis, T., Hugenholtz, P., Larsen, N., et al. (2006). 'GreenGenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB', *Applied and Environmental Microbiology*, 72/7: 5069–72.

Díaz-Muñoz, S. L. (2017). 'Viral coinfection is shaped by host ecology and virus-virus interactions across diverse microbial taxa and environments', *Virus Evolution*, 3/1: vex011.

van Dongen, S. M. (2000). *Graph clustering by flow simulation*, Ph.D.thesis, University of

Utrecht.

Drake, J. W. (1999). 'The distribution of rates of spontaneous mutation over viruses,
prokaryotes, and eukaryotes', *Annals of the New York Academy of Sciences*, 870: 100–7.

Eddy, S. R. (1998). 'Profile hidden Markov models', *Bioinformatics* , 14/9: 755–63.

Edgar, R. (2004). 'MUSCLE: a multiple sequence alignment method with reduced time and
space complexity', *BMC Bioinformatics*, 5: 113.

Edgar, R. (2007). 'PILER-CR: fast and accurate identification of CRISPR repeats', *BMC
Bioinformatics*, 8: 18.

Ferla, M., Thrash, J. C., Giovannoni, S., et al. (2013). 'New rRNA gene-based phylogenies of
the alphaproteobacteria provide perspective on major groups, mitochondrial ancestry and
phylogenetic instability', *PLoS One*, 8/12: e83383.

Fogg, P. C., Westbye, A., & Beatty, J. T. (2012). 'One for all or all for one: heterogeneous
expression and host cell lysis are key to gene transfer agent activity in *Rhodobacter
capsulatus*', *PLoS One*, 7/8: e43772.

Frost, L., Leplae, R., Summers, A., et al. (2005). 'Mobile genetic elements: the agents of open
source evolution', *Nature Reviews of Microbiology*, 3/9: 722–32.

Grazziotin, A. L., Koonin, E. V., & Kristensen, D. M. (2017). 'Prokaryotic Virus Orthologous
Groups (pVOGs): a resource for comparative genomics and protein family annotation',
Nucleic Acids Research, 45/D1: D491–8.

Gupta, R. S., & Mok, A. (2007). 'Phylogenomics and signature proteins for the alpha
proteobacteria and its main groups', *BMC Microbiology*, 7: 106.

Guy, L., Kultima, J. R., & Andersson, S. G. E. (2010). 'genoPlotR: comparative gene and
genome visualization in R', *Bioinformatics* , 26/18: 2334–5.

Guy, L., Nystedt, B., Toft, C., et al. (2013). 'A gene transfer agent and a dynamic repertoire of
secretion systems hold the keys to the explosive radiation of the emerging pathogen
Bartonella', *PLoS Genetics*, 9/3: e1003393.

Hershberg, R., & Petrov, D. A. (2010). 'Evidence that mutation is universally biased towards AT
in bacteria', *PLoS Genetics*, 6/9: e1001115.

He, Z., Zhang, H., Gao, S., et al. (2016). 'Evolview v2: an online visualization and management
tool for customized and annotated phylogenetic trees', *Nucleic Acids Research*, 44/W1:
W236–41.

Hildebrand, F., Meyer, A., & Eyre-Walker, A. (2010). 'Evidence of selection upon genomic GC-
content in bacteria', *PLoS Genetics*, 6/9: e1001107.

Huang, S., Zhang, Y., Chen, F., et al. (2011). 'Complete genome sequence of a marine
roseophage provides evidence into the evolution of gene transfer agents in
alphaproteobacteria', *Virology Journal*, 8: 124.

Humphrey, S. B., Stanton, T. B., Jensen, N. S., et al. (1997). 'Purification and characterization

of VSH-1, a generalized transducing bacteriophage of *Serpulina hyodysenteriae*', *Journal of Bacteriology*, 179/2: 323–9.

Hynes, A. P., Mercer, R. G., Watton, D., et al. (2012). 'DNA packaging bias and differential expression of gene transfer agent genes within a population during production and release of the *Rhodobacter capsulatus* gene transfer agent, RcGTA', *Molecular Microbiology*, 85/2: 314–25.

Hynes, A. P., Shakya, M., Mercer, R. G., et al. (2016). 'Functional and evolutionary characterization of a gene transfer agent's multilocus "genome"', *Molecular Biology and Evolution*, 33/10: 2530–43.

Iranzo, J., Krupovic, M., & Koonin, E. V. (2016). 'The double-stranded DNA virosphere as a modular hierarchical network of gene sharing', *mBio*, 7/4: e00978-16

Iranzo, J., Krupovic, M., & Koonin, E. V. (2017). 'A network perspective on the virus world', *Communicative & Integrative Biology*, 10/2: e1296614.

Juhas, M., van der Meer, J. R., Gaillard, M., et al. (2009). 'Genomic islands: tools of bacterial horizontal gene transfer and evolution', *FEMS Microbiology Reviews*, 33/2: 376–93.

Ju, K.-S., Gao, J., Doroghazi, J. R., et al. (2015). 'Discovery of phosphonic acid natural products by mining the genomes of 10,000 actinomycetes', *Proceedings of the National Academy of Sciences of the United States of America*, 112/39: 12175–80.

Knoll, A. H. (2014). 'Paleobiological perspectives on early eukaryotic evolution', *Cold Spring Harbor perspectives in biology*, 6/1: a016121

Koonin, E., Senkevich, T., & Dolja, V. V. (2006). 'The ancient Virus World and evolution of cells', *Biology Direct*, 1: 29.

Koonin, E. V. (2016). 'Horizontal gene transfer: essentiality and evolvability in prokaryotes, and roles in evolutionary transitions', *F1000Research*, 5: 1805.

Kristensen, D. M., Waller, A. S., Yamada, T., et al. (2013). 'Orthologous gene clusters and taxon signature genes for viruses of prokaryotes', *Journal of Bacteriology*, 195/5: 941–50.

Krupovic, M., & Koonin, E. V. (2017). 'Multiple origins of viral capsid proteins from cellular ancestors', *Proceedings of the National Academy of Sciences of the United States of America*, 114/12: E2401–10.

Kuchinski, K. S., Brimacombe, C. A., Westbye, A. B., et al. (2016). 'The SOS response master regulator LexA regulates the Gene Transfer Agent of *Rhodobacter capsulatus* and represses transcription of the signal transduction protein CckA', *Journal of Bacteriology*, 198/7: 1137–48.

Lang, A. S., & Beatty, J. T. (2000). 'Genetic analysis of a bacterial genetic exchange element: the gene transfer agent of Rhodobacter capsulatus', *Proceedings of the National Academy of Sciences of the United States of America*, 97/2: 859–64.

Lang, A. S., & Beatty, J. T. (2007). 'Importance of widespread gene transfer agent genes in α-proteobacteria', *Trends in Microbiology*, 15/2: 54–62.

Lang, A. S., Taylor, T., & Beatty, J. T. (2002). 'Evolutionary implications of phylogenetic analyses of the gene transfer agent (GTA) of *Rhodobacter capsulatus*', *Journal of Molecular Evolution*, 55/5: 534–43.

Lang, A. S., Westbye, A. B., & Beatty, J. T. (2017). 'The distribution, evolution, and roles of gene transfer agents in prokaryotic genetic exchange', *Annual Review of Virology*, 4/1: 87-104.

Lang, A. S., Zhaxybayeva, O., & Beatty, J. T. (2012). 'Gene transfer agents: phage-like elements of genetic exchange', *Nature Reviews. Microbiology*, 10/7: 472–82.

Lassalle, F., Périan, S., Bataillon, T., et al. (2015). 'GC-content evolution in bacterial genomes: the biased gene conversion hypothesis expands', *PLoS Genetics*, 11/2: e1004941.

Lawrence, J. G., Hendrix, R. W., & Casjens, S. (2001). 'Where are the pseudogenes in bacterial genomes?', *Trends in Microbiology*, 9/11: 535–40.

Lee, K. B., Liu, C. T., Anzai, Y., et al. (2005). 'The hierarchical system of the "Alphaproteobacteria": description of Hyphomonadaceae fam. nov., Xanthobacteraceae fam. nov. and Erythrobacteraceae fam. nov', *International Journal of Systematic and Evolutionary Microbiology*, 55/Pt 5: 1907–19.

Le, S. Q., & Gascuel, O. (2008). 'An improved general amino acid replacement matrix', *Molecular Biology and Evolution*, 25/7: 1307–20.

Leung, M., Brimacombe, C., Spiegelman, G. B., et al. (2012). 'The GtaR protein negatively regulates transcription of the gtaRI operon and modulates gene transfer agent (RcGTA) expression in *Rhodobacter capsulatus*', *Molecular Microbiology*, 83/4: 759–74.

Leung, M., Florizone, S., Taylor, T., et al. (2010). 'The gene transfer agent of Rhodobacter capsulatus', *Advances in Experimental Medicine and Biology*, 675: 253–64.

Li, L., Stoeckert, C. J., Jr, & Roos, D. S. (2003). 'OrthoMCL: Identification of ortholog groups for eukaryotic genomes', *Genome Research*, 13/9: 2178–89.

Luo, H. (2014). 'Evolutionary origin of a streamlined marine bacterioplankton lineage', *The ISME Journal*, 9/6: 1423–33.

Luo, H., Csuros, M., Hughes, A., et al. (2013). 'Evolution of divergent life history strategies in marine alphaproteobacteria', *mBio*, 4/4: e00373–13.

Makarova, K. S., Wolf, Y. I., Alkhnbashi, O. S., et al. (2015). 'An updated evolutionary classification of CRISPR-Cas systems', *Nature Reviews Microbiology*, 13/11: 722–36.

Mann, T. H., Seth Childers, W., Blair, J. A., et al. (2016). 'A cell cycle kinase with tandem sensory PAS domains integrates cell fate cues', *Nature Communications*, 7: 11454.

Marrs, B. (1974). 'Genetic recombination in *Rhodopseudomonas capsulata*', *Proceedings of the National Academy of Sciences of the United States of America*, 71/3: 971–3.

Mcdaniel, L., Young, E., Delaney, J., et al. (2010). 'High frequency of horizontal gene transfer in the oceans', *Science*, 330/6000: 50.

McDaniel, L., Young, E., Ritchie, K., et al. (2012). 'Environmental factors influencing gene
transfer agent (GTA) mediated transduction in the subtropical ocean', *PLoS One*, 7/8:
e43506.

McHugh, C. A., Fontana, J., Nemecek, D., et al. (2014). 'A virus capsid-like nanocompartment
that stores iron and protects bacteria from oxidative stress', *The EMBO journal*, 33/17:
1896–911.

Mercer, R. G., & Lang, A. S. (2014). 'Identification of a predicted partner-switching system that
affects production of the gene transfer agent RcGTA and stationary phase viability in
Rhodobacter capsulatus', *BMC Microbiology*, 14: 71.

Mira, A., Ochman, H., & Moran, N. A. (2001). 'Deletional bias and the evolution of bacterial
genomes', *Trends in Genetics*, 17/10: 589–96.

Nagao, N., Yamamoto, J., Komatsu, H., et al. (2015). 'The gene transfer agent-like particle of
the marine phototrophic bacterium *Rhodovulum sulfidophilum*', *Biochemistry and
Biophysics Reports*, 4: 369–74.

NCBI Resource Coordinators. (2017). 'Database resources of the National Center for
Biotechnology Information', *Nucleic Acids Research*, 45/D1: D12–7.

Paterson, S., Vogwill, T., Buckling, A., et al. (2010). 'Antagonistic coevolution accelerates
molecular evolution', *Nature*, 464/7286: 275–8.

Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). 'FastTree 2--approximately maximum-likelihood
trees for large alignments', *PLoS One*, 5/3: e9490.

Rapp, B. J., & Wall, J. D. (1987). 'Genetic transfer in *Desulfovibrio desulfuricans*', *Proceedings
of the National Academy of Sciences of the United States of America*, 84/24: 9128–30.

Redfield, R. J. (2001). 'Do bacteria have sex?', *Nature Reviews Genetics*, 2/8: 634–9.

Richards, T. A., & Archibald, J. M. (2011). 'Cell evolution: gene transfer agents and the origin of
mitochondria', *Current biology*, 21/3: R112–4.

Rocha, E. P. C. (2006). 'Inference and analysis of the relative stability of bacterial
chromosomes', *Molecular biology and evolution*, 23/3: 513–22.

Rocha, E. P. C., & Danchin, A. (2002). 'Base composition bias might result from competition for
metabolic resources', *Trends in Genetics*, 18/6: 291–4.

Rocha, E. P. C., & Feil, E. J. (2010). 'Mutational patterns cannot explain genome composition:
Are there any neutral sites in the genomes of bacteria?', *PLoS Genetics*, 6/9: e1001104.

Rodriguez-Valera, F., Martin-Cuadrado, A.-B., & López-Pérez, M. (2016). 'Flexible genomic
islands as drivers of genome evolution', *Current Opinion in Microbiology*, 31: 154–60.

Salichos, L., Stamatakis, A., & Rokas, A. (2014). 'Novel information theory-based measures for
quantifying incongruence among phylogenetic trees', *Molecular Biology and Evolution*,
31/5: 1261–71.

Sarris, P. F., Ladoukakis, E. D., Panopoulos, N. J., et al. (2014). 'A phage tail-derived element

with wide distribution among both prokaryotic domains: a comparative genomic and phylogenetic study', *Genome Biology and Evolution*, 6/7: 1739–47.

Schaefer, A. L., Taylor, T., Beatty, J. T., et al. (2002). 'Long-chain acyl-homoserine lactone quorum-sensing regulation of *Rhodobacter capsulatus* gene transfer agent production', *Journal of Bacteriology*, 184/23: 6515–21.

Schloss, P., Westcott, S., Ryabin, T., et al. (2009). 'Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities', *Applied and Environmental Microbiology*, 75/23: 7537–41.

Selengut, J. D., Haft, D. H., Davidsen, T., et al. (2007). 'TIGRFAMs and Genome Properties: tools for the assignment of molecular function and biological process in prokaryotic genomes', *Nucleic Acids Research*, 35/Database issue: D260–4.

Sen, D., Chandrababunaidu, M. M., Singh, D., et al. (2015). 'Draft genome sequence of the terrestrial cyanobacterium *Scytonema millei* VB511283, isolated from Eastern India', *Genome announcements*, 3/2: e00009-15

Shikuma, N. J., Pilhofer, M., Weiss, G. L., et al. (2014). 'Marine tubeworm metamorphosis induced by arrays of bacterial phage tail-like structures', *Science*, 343/6170: 529–33.

Smillie, C. S., Smith, M. B., Friedman, J., et al. (2011). 'Ecology drives a global network of gene exchange connecting the human microbiome', *Nature*, 480/7376: 241–4.

Solioz, M., & Marrs, B. (1977). 'The gene transfer agent of *Rhodopseudomonas capsulata*. Purification and characterization of its nucleic acid', *Archives of Biochemistry and Biophysics*, 181/1: 300–7.

Stamatakis, A. (2014). 'RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies', *Bioinformatics* , 30/9: 1312–3.

Sutter, M., Boehringer, D., Gutmann, S., et al. (2008). 'Structural basis of enzyme encapsulation into a bacterial nanocompartment', *Nature Structural & Molecular Biology*, 15/9: 939–47.

Thrash, J., Boyd, A., Huggett, M., et al. (2011). 'Phylogenomic evidence for a common ancestor of mitochondria and the SAR11 clade', *Scientific Reports*, 1: 1–9.

Touchon, M., Bernheim, A., & Rocha, E. P. (2016). 'Genetic and life-history traits associated with the distribution of prophages in bacteria', *The ISME journal,* 10: 2744-2754.

Touchon, M., Moura de Sousa, J. A., & Rocha, E. P. (2017). 'Embracing the enemy: the diversification of microbial gene repertoires by phage-mediated horizontal gene transfer', *Current Opinion in Microbiology*, 38: 66–73.

Wang, X., Jin, D., Zhou, L., et al. (2014). 'Draft genome sequence of halotolerant polycyclic aromatic hydrocarbon-degrading *Pseudomonas bauzanensis* Strain W13Z2', *Genome Announcements*, 2/5: e01049-14

Wang, X., Kim, Y., Ma, Q., et al. (2010). 'Cryptic prophages help bacteria cope with adverse environments', *Nature Communications*, 1: 147.

Wang, Z., & Wu, M. (2015). 'An integrated phylogenomic approach toward pinpointing the origin

of mitochondria', *Scientific Reports*, 5: 7949.

Westbye, A. B., Kuchinski, K., Yip, C. K., et al. (2015). 'The gene transfer agent RcGTA contains head spikes needed for binding to the *Rhodobacter capsulatus* polysaccharide cell capsule', *Journal of Molecular Biology,* 428: 477-91.

Westbye, A. B., Leung, M. M., Florizone, S. M., et al. (2013). 'Phosphate concentration and the putative sensor kinase protein CckA modulate cell lysis and release of the *Rhodobacter capsulatus* gene transfer agent', *Journal of Bacteriology*, 195/22: 5025–40.

Williams, K. P., Sobral, B. W., & Dickerman, A. W. (2007). 'A robust species tree for the alphaproteobacteria', *Journal of Bacteriology*, 189/13: 4578–86.

Yang, Z. (1994). 'Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: approximate methods', *Journal of Molecular Evolution*, 39/3: 306– 14.

Zan, J., Heindl, J., Liu, Y., et al. (2013). 'The CckA-ChpT-CtrA phosphorelay system is regulated by quorum sensing and controls flagellar motility in the marine sponge symbiont *Ruegeria* sp. KLH11', *PLoS One*, 8/6: e66346.

Zhan, Y., Huang, S., Voget, S., et al. (2016). 'A novel roseobacter phage possesses features of podoviruses, siphoviruses, prophages and gene transfer agents', *Scientific Reports*, 6: 30372.

Zhaxybayeva, O., & Doolittle, W. F. (2011). 'Lateral gene transfer', *Current Biology* , 21/7: R242–6.

A. RcGTA Genome

POGs   No detected POGs

B. Viruses (1,783)

0%   5.6%   11.2%   16.8%   22.4%

C. Bacteria (30,503)

0%   3.0%   12.0%   18.0%   24.0%

2 kb

Proportion of genomes with an RcGTA homolog:

Number of *rcc01682–rcc01698* homologs in a cluster

Rhizobiales (102)

Caulobacterales (7)

Rhodobacterales II (3)

Parvularculales (1)

Rhodobacterales I (22)

Sphingomonadales (15)

Rhodospirillales & SAR116 (31)

Rickettsiales (69)

Pelagibacterales (4)

Magnetococcales (1)

Outgroup

① ②

Large Cluster
Small Cluster
No RcGTA homologs

All α-proteobacteria (255)