Unique ATP-cone-driven allosteric regulation of ribonucleotide

reductase via the radical-generating subunit

Inna Rozman Grinberg[1], Daniel Lundin[1], Mahmudul Hasan[1,2,#], Mikael Crona[3,#],

Venkateswara Rao Jonna[4,#], Christoph Loderer[1], Margareta Sahlin[1], Natalia Markova[5],

Ilya Borovok[6], Gustav Berggren[7], Anders Hofer[4], Derek T Logan[2,*] and Britt-Marie

Sjöberg[1,*]

[1]Department of Biochemistry and Biophysics, Stockholm University, Sweden, [2]Department of

Biochemistry and Structural Biology, Lund University, Sweden, [3]Swedish Orphan Biovitrum

AB, Stockholm, Sweden, [4]Department of Medical Biochemistry and Biophysics, Umeå

University, Sweden, [5]Malvern Instruments Inc., Sweden, [6]Department of Molecular

Microbiology and Biotechnology, Tel-Aviv University, Israel, and [7]Department of Chemistry,

Uppsala University. Sweden.

[#]These authors contributed equally

*Corresponding authors

Abstract

Ribonucleotide reductases (RNRs) are key enzymes in DNA synthesis and repair, with

sophisticated allosteric mechanisms controlling both substrate specificity and overall activity.

In RNRs, the activity master-switch, the ATP-cone, has been found exclusively in the

catalytic subunit. In two class I RNR subclasses whose catalytic subunit lacks the ATP-cone,

we discovered ATP-cones in the radical-generating subunit. The ATP-cone in the

Leewenhoekiella blandensis radical-generating subunit regulates activity via modifications of

quaternary structure induced by binding of nucleotides. ATP induces enzymatically

competent dimers, whereas dATP induces non-productive tetramers, resulting in different

holoenzyme complexes. The tetramer forms solely by interactions between ATP-cones, as

evidenced by a 2.45 Å crystal structure. We also present evidence for an $Mn^{III}Mn^{IV}$ metal

center. In summary, lack of an ATP-cone domain in the catalytic subunit was compensated

by evolutionary capture of the domain by the radical-generating subunit. Our findings present

a novel opportunity for dATP-regulation of engineered proteins.

Introduction

Allosteric regulation of an enzyme is defined as regulation of activity by binding of an effector
molecule to a different location of the enzyme than the active site. The effector influences
the distribution of tertiary or quaternary conformations of an enzyme, alone or in
combination, modulating its activity (Swain & Gierasch, 2006). Allostery is an intrinsic
property of many, if not all, dynamic proteins (Gunasekaran, Ma, & Nussinov, 2004) and an
important factor in disease (Nussinov & Tsai, 2013), and has hence attracted considerable
scientific interest. A substantial part of this interest has been focused on ribonucleotide
reductase (RNR), which has been termed a "paradigm for allosteric regulation of enzymes"
(Aravind, Wolf, & Koonin, 2000).

RNRs are essential enzymes in all free-living cells, providing the only known *de novo*
pathway for the biosynthesis of deoxyribonucleotides (dNTPs), the immediate precursors for
DNA synthesis and repair (Hofer, Crona, Logan, & Sjöberg, 2012; Nordlund & Reichard,
2006). To avoid misbalanced levels of dNTPs and the increased mutation rates that are the
inevitable consequences of this (Kumar et al., 2011; Mathews, 2006; Watt, Buckland, Lujan,
Kunkel, & Chabes, 2016), RNRs are tightly controlled through transcriptional and allosteric
regulation, subcellular compartmentalization and small protein inhibitors (Pai & Kearsey,
2017; Torrents, 2014). Allosteric regulation of RNRs affects both substrate specificity and
overall activity. The specificity regulation has been intensively studied and described for all
three classes of RNRs (Andersson, Westman, Hofer, & Sjöberg, 2000; Hofer et al., 2012;
Larsson et al., 2004; Reichard, 2010; Torrents et al., 2000; Zimanyi, Chen, Kang, Funk, &
Drennan, 2016). The s-site binds dNTPs and determines which nucleotide will be reduced at
the active site to ensure balanced levels of the four dNTPs in the cell. Additionally, many
RNRs possess an overall activity regulation site (a-site) (Brown & Reichard, 1969; Thelander
& Reichard, 1979) positioned in an N-terminal domain of ~100 amino acid residues called
the ATP-cone (Aravind et al., 2000; Eriksson et al., 1997). Acting as a regulatory master
switch, the a-site senses intracellular nucleotide concentrations by competitive binding of
ATP and dATP. In presence of ATP the enzyme is active, and when concentrations of
dNTPs rise, binding of dATP switches the enzyme off. This mechanism ensures sufficient
but not excessive amounts of nucleotides that may also cause increased mutation rates
(Mathews, 2006).

To date, ATP-cone domains in RNRs have been observed exclusively at the N-terminus of
the catalytic subunit NrdA (class I), NrdJ (class II) and NrdD (class III). Class I RNRs consist
of a large, catalytic subunit (α or NrdA), and a smaller, radical-generating subunit (β or NrdB)
(Huang, Parker, & Stubbe, 2014; Nordlund & Reichard, 2006). NrdA and NrdB interact to
form the active complex, in which the two proteins need to be precisely positioned such that
the radical can be transferred from NrdB, where it is generated and stored, to NrdA where it
starts the substrate reduction. In class I, it has long been noted that ATP-cones are absent
from subclass Ib (NrdE) but present in several, but not all, NrdAs. A recent phylogenetic
subclassification of RNRs reveals that three phylogenetically well-supported subclasses of
class I never have ATP-cones (Jonna et al., 2015) (http://rnrdb.pfitmap.org): NrdE, NrdAi
and NrdAk. In two of these subclasses we instead discovered ATP-cones attached to their
corresponding radical-generating subunit: NrdF (the Ib subclass) and NrdBi. It hence
appears as if the lack of activity regulation through an ATP-cone in the catalytic subunit is

compensated by the presence of this domain in the non-homologous radical-generating
subunit of some RNRs.

Three distinct types of class I complexes have been mechanistically characterized and found
to operate via nucleotide-induced regulation of quaternary structure (Johansson et al., 2016;
Jonna et al., 2015; Kashlan, Scott, Lear, & Cooperman, 2002; Rofougaran, Crona, Vodnala,
Sjöberg, & Hofer, 2008; Rofougaran, Vodnala, & Hofer, 2006; Torrents, Westman, Sahlin, &
Sjöberg, 2006). Crystal structures, cryo-electron microscopy reconstructions and small-angle
X-ray scattering studies of inhibited complexes have revealed that when dATP is bound at
the a-site, high molecular mass oligomers are formed, in which the radical transfer pathway
is distorted (Ando et al., 2011; Ando et al., 2016; Fairman et al., 2011; Johansson et al.,
2016). Conversely, when ATP is bound, an active enzyme complex is formed. Interestingly,
the structure and organization of subunits in active and inactive complexes varies
considerably between species (Ahmad & Dealwis, 2013; Hofer et al., 2012). In *Escherichia*
coli RNR, the active NrdAB complex is $\alpha_2\beta_2$, whereas the inactive form is an $\alpha_4\beta_4$ ring-
shaped octamer where the ATP-cones in the α subunits interact with the β subunits (Ando et
al., 2011). In the eukaryotic class I RNR, the inactive complex differs from *E. coli* in that it
has an α_6 stoichiometry. This hexamer can only bind one β_2 subunit in an unproductive
manner without a properly aligned electron transport chain (Fairman et al., 2011). Activation
by ATP creates a different type of α_6 complex that binds one or more β_2 complexes (Ando et
al., 2016; Aye & Stubbe, 2011; Crona et al., 2013; Fairman et al., 2011; Rofougaran et al.,
2006). The different complexes are formed by subtle changes at the a-site induced by
binding of the different nucleotides (Fairman et al., 2011). Another oligomerization
mechanism has been recently found in *Pseudomonas aeruginosa* class I RNR, which
possesses two consecutive ATP-cones, of which only the N-terminal one binds nucleotides.
The active complex is once again $\alpha_2\beta_2$, but the inactive *P. aeruginosa* RNR complex is a
dATP-induced α_4 complex consisting of a ring of four α subunits interacting via their outer
ATP-cones (Johansson et al., 2016; Jonna et al., 2015). A single β_2 can bind to this ring, but
the complex is inactive.

The unexpected finding of an ATP-cone fused to the radical-generating subunits poses
questions regarding how it might regulate activity. Here we describe the mechanism of
activity regulation by the ATP-cone N-terminally fused to the radical-generating NrdBi from
Leeuwenhoekiella blandensis strain MED217. *L. blandensis* was isolated from
Mediterranean surface seawater and belongs to Flavobacteriaceae, the major family of
marine Bacteroidetes, with important roles in carbon flow and nutrient turnover in the sea
during and following algal blooms (Fernandez-Gomez et al., 2013; Pinhassi et al., 2006). *L.*
blandensis possesses two RNRs: a class II without ATP-cone, and the class I NrdAi/NrdBi,
which lacks an ATP-cone in NrdA and instead contains an ATP-cone positioned at the N-
terminus of NrdB. Superficially, the allosteric mechanism of *L. blandensis* NrdAi/NrdBi
holoenzyme is similar to when the ATP-cone is contained in the catalytic subunit. At high
dATP concentrations, inhibited higher oligomeric complexes of the holoenzyme are
favoured. However, in the *L. blandensis* class I RNR, the oligomerization is driven by a shift
towards tetramers of the radical-generating subunit. This illustrates how allosteric regulation
controlled by ATP-cones can evolve in a highly dynamic way, requiring few adaptations to
the core of the enzyme. The relative ease by which ATP-cone-driven activity regulation
appears to evolve, provides a potential route to regulate engineered enzymes by dATP-
inhibition.

Results

The activity allosteric regulatory ATP-cone is linked to the radical-containing subunit in some class I RNRs

We detected ATP-cones in the radical-generating subunits of RNRs from two distinct
phylogenetic RNR subclasses: NrdBi and NrdF (Fig. 1a). In the NrdBi sequences, the ATP-
cone was found at the N-terminus of the protein, whereas it was found at the C-terminus of
the NrdF proteins (Fig. 1b). Interestingly, the corresponding catalytic subunit subclasses –
NrdAi and NrdE respectively – have been found to lack ATP-cones. Ninety-three sequences
in NCBI's RefSeq database are NrdBi proteins with N-terminally positioned ATP-cones. They
are encoded by viruses and bacteria from several phyla, although the Flavobacteriales order
in the Bacteroidetes phylum predominate (70 sequences, http://rnrdb.pfitmap.org). NrdF
proteins with a C-terminally positioned ATP-cone are only encoded by four species of the
Meiothermus genus of the Deinococcus-Thermus phylum (http://rnrdb.pfitmap.org). All
species encoding NrdB proteins with ATP-cones in their genomes also encode other RNRs.

Glutaredoxin ATP-cone domain NrdA, class I catalytic component NrdB, class I radical-generator

Figure 1. Unrooted phylogenetic tree of NrdB sequences and genome arrangements
of NrdB sequences with ATP-cones. a) Maximum likelihood phylogeny of class I RNR
radical generating subunit with subclasses in color, names to the left and organism

distributions to the right (see inset legend; sizes of circles are proportional to the number of
sequences found in each taxon). Bootstrap support values greater than 0.7 are shown with
colored diamonds. NrdBs with ATP-cones were discovered in two subclasses: NrdBi and
NrdF (formerly class Ib). Neither of the two subclasses have corresponding catalytic
subunits, NrdAi and NrdE respectively, with ATP-cones. In both subclasses, NrdB
sequences with ATP-cones were rare and phylogenetically limited, see inset arrows. b)
Arrangement of class I RNR genes in three genomes encoding NrdB proteins with ATP-
cones (*green borders*, NrdAi/NrdBi; *purple borders*, NrdE/NrdF). Genes are shown 5' to 3',
so that ATP-cones in the N-terminus are to the left in the gene.

Substrate specificity regulation of *L. blandensis* RNR via the s-site

Initially we cloned, expressed and purified the *L. blandensis* NrdBi and NrdAi proteins. Using
a four-substrate activity assay in the presence of saturating concentrations of the s-site
effectors dTTP, dGTP or ATP, we found that *L. blandensis* RNR has a similar specificity
regulation pattern to most characterized RNRs (Hofer et al., 2012). ATP stimulated the
reduction of CDP and UDP, whereas dTTP stimulated the reduction of GDP, and dGTP
stimulated the reduction of ADP and GDP (Fig. 2). The enzyme was completely inactive in
the absence of allosteric effectors. Using mixtures of allosteric effectors, we observed that
dTTP-induced GDP reduction dramatically increased in the presence of ATP (Fig. 2).

Figure 2: Allosteric specificity regulation of the *L. blandensis* class I RNR. Enzyme
activity was measured after 10 and 30 min in the presence of all four substrates (0.5 mM
each) and with the indicated allosteric effectors (2 mM of each). Error bars indicate the
extremes of two measurements. Protein concentrations were 1 µM NrdB and 4 µM NrdA.

Overall activity of *L. blandensis* RNR is regulated via the NrdB-linked ATP-cone

We performed a series of activity assays with CDP as substrate to elucidate the potential roles of ATP and dATP in activating and inhibiting the enzyme (Fig. 3). The presence of ATP clearly activated the enzyme (Fig. 3a), while dATP activated the enzyme when used at low concentrations and was inhibitory at 30 µM and higher (Fig. 3b). An ATP-cone deletion mutant NrdBΔ99, lacking the N-terminal 99 residues, reached a lower maximum activity compared to the wild type enzyme, suggesting that it was not activated by ATP beyond saturation of the s-site in the NrdA, nor was it inhibited by dATP (Fig. 3a-b). From the NrdBΔ99 effector titrations, we could calculate K_L values – the concentrations of allosteric effectors that give half maximal enzyme activity – for binding of ATP and dATP to the s-site in NrdA to 30 and 1.4 µM respectively. Titration of ATP into wild type NrdB in the presence of an excess of NrdA saturated with dTTP showed that it activates the enzyme through the a-site with a K_L of 96 µM (Fig. 3c). For the corresponding inhibition by dATP binding to the a-site, we calculated the K_i value - the binding constant of a non-competitive inhibitor - to be 20 µM in the presence of an s-site saturating dTTP concentration (Fig. 3d). We also tested if only the triphosphate form of (deoxy)adenosine nucleotides would interact with the a-site in presence of s-site saturating dTTP concentrations. In addition to ATP and dATP, dADP was also found to interact, whereas ADP, AMP and dAMP had no effect (Fig. 3e). Titration with dADP inhibited the enzyme activity, although less strongly than dATP (Fig. 3f).

Figure 3: Activity of *L. blandensis* RNR with wild type NrdB or NrdBΔ99 in the presence of allosteric effectors. NrdB was used in excess over NrdA when studying the s-site and NrdA was used in excess when studying the a-site. a-b) CDP reduction assayed in mixtures of 0.5 µM NrdA and 2 µM of either wild type NrdB (black circles) or NrdBΔ99 (white circles), titrated with ATP (a) or dATP (b). c) GDP reduction assay mixtures with 2 µM NrdA and 0.5 µM wild type NrdB titrated with ATP in the presence of an s-site saturating concentration of dTTP (2 mM). d) Reduction of GDP assayed with 2 µM NrdA and 0.5 µM wild type NrdB, titrated with dATP in the presence of an s-site saturating concentration of dTTP (2 mM). e) GDP reduction in presence of s-site saturating dTTP (2 mM) and 2 mM of the indicated adenosine nucleotides. Assay mixtures contained 4 µM NrdA and 1 µM NrdB. 100% activity corresponded to 639 nmol x mg^{-1} x min^{-1}. f) CDP reduction assays titrated with dADP. Assay mixtures contained 2 µM NrdA and 0.5 µM NrdB. Error bars in panels E and F indicate the standard deviation of three measurements.

dATP binding to NrdB induces formation of higher oligomeric complexes

To elucidate the mechanism of allosteric overall activity regulation governed by the NrdB-linked ATP-cone, activity-independent oligomer-distribution experiments were performed by gas-phase electrophoretic macromolecule analysis (GEMMA). In the absence of allosteric effectors, NrdB (β) is mainly monomeric (theoretically 51.8 kDa) and in contrast to most studied NrdB proteins it does not readily form dimers at the low protein concentration range suitable for GEMMA analyses (Fig. 4a). Addition of dATP (50 µM) dramatically shifted the equilibrium towards tetramers β_4, which became the major form under these conditions (Fig. 4a). Titration of increasing concentrations of dATP to NrdB showed that the tetramers reached their half-maximum mass concentration at around 10 µM dATP (Supplementary Fig. S1). Addition of 50 µM dADP also induced formation of NrdB tetramers (Fig. 4a). The NrdBΔ99 mutant, lacking the ATP-cone (Fig. 4b), was in contrast mainly monomeric regardless if dATP was present or not (Fig. 4b), demonstrating that the NrdB-linked ATP-cone is required for dATP-induced tetramer formation. NrdA was a monomer (theoretically 70.6 kDa) in the absence of allosteric effectors, while addition of dATP prompted formation of dimers (Fig. 4c) To assess the oligomeric state of the complete enzymatic complex of the inactive *L. blandensis* RNR, a mixture of NrdA (α) and NrdB (β) in the presence of 100 µM dATP was analyzed with GEMMA (Fig. 4c). The two subunits formed a large complex of 340 kDa with the expected mass of an $\alpha_2\beta_4$ complex (theoretically 348 kDa), and at higher protein concentration a complex of 465 kDa with the expected mass of an $\alpha_4\beta_4$ complex (theoretically 488 kDa).

Figure 4: GEMMA analysis of the *L. blandensis* RNR subunits α, β and their combinations in the presence and the absence of allosteric effectors. a) 0.1 mg/ml NrdB (~2 µM) analyzed in the absence and presence of 50 µM dATP or dADP. b) The NrdBΔ99 mutant analyzed in the absence and the presence of 50 µM dATP. c) In the top two traces 0.1 mg/ml NrdA (~1.4 µM) is analyzed in the absence or presence of 100 µM dATP. In the third trace, 0.1 mg/ml NrdB is added to the mixture. The last trace is similar to the third trace, but the concentration of NrdA and NrdB is increased to 0.4 mg/ml NrdA and 0.3 mg/ml NrdB (~6 µM of each). The analyses of NrdA-NrdB complexes were performed at a very low pressure (1.4 Psi) to avoid the influence of magnesium-nucleotide clusters on the measurement and to minimize the risk of false protein-protein interactions. The baselines of the individual experiments are distributed in the vertical direction to be able to fit many traces in each panel.

To complement the GEMMA analyses of oligomer formation, we performed analytical size
exclusion chromatography (SEC) using higher protein concentrations and physiologically
reasonable concentrations of effectors (3 mM ATP and 0.1 to 0.5 mM dATP) (Bochner &
Ames, 1982; Buckstein, He, & Rubin, 2008). At the higher protein concentrations used in
SEC (as compared to GEMMA) NrdB eluted predominantly as dimers rather than monomers
in the absence of effectors and similar results were also obtained in the presence of ATP. In
agreement with the GEMMA results, the distribution was shifted towards tetramers in the
presence of dATP (Fig. 5a). Without effectors, NrdA was mainly in a monomeric state, while
it was dominated by dimers in the presence of either of the two effectors (Fig. 5b). When
NrdA and NrdB were mixed in the absence of allosteric effectors, no complex was visible.
After addition of ATP, a new complex of ~200 kDa appeared (Fig. 5c). To verify its
composition and stoichiometry, we analyzed fractions eluted from SEC on SDS PAGE. Both
NrdA and NrdB were visible on the gel in a 1:1 molar ratio (Fig. 5c insert). This complex,
formed in conditions promoting active RNR, is conceivably an $\alpha_2\beta_2$ complex. After addition of
the inhibiting effector dATP, a complex with a molecular mass consistent with $\alpha_4\beta_4$ eluted.
The few differences between GEMMA and SEC are due to the different protein
concentrations used and a complementary SEC study at lower protein concentration
showed that the two methods are in good agreement with each other (Supplementary Fig.
S2). A summary of the protein complexes formed in the presence and absence of effectors
is presented in figure 5d.

Figure 5: Size exclusion chromatography of *L. blandensis* RNR in the absence and
presence of 3 mM ATP or 0.5 mM dATP. Proteins at concentrations of 20 μM each were
incubated separately or mixed together for 10 minutes in the absence (black dotted line) or
presence of ATP (blue dashed line) or dATP (red solid line), centrifuged and applied to the
column equilibrated with SEC buffer. Panels show NrdB (a), NrdA (b), and an equimolar
mixture of NrdA and NrdB (c). C Insert: SDS PAGE of the eluted proteins run in the
presence and absence of the indicated effectors. Elution positions of fractions applied to gel
are indicated by *red* (in the presence of dATP) and *blue* (in the presence of ATP) arrows,
respectively. d) Summary of multiple SEC experiments varying protein concentrations of
NrdA (10-113 μM), NrdB (5-150 μM) and mixtures of NrdA and NrdB at ratios of 1:1 or 1:2.
Molecular masses and standard deviations are calculated for 3-5 experiments, and closest
estimated complex stoichiometry are shown in parenthesis, with major species underlined
when appropriate (see Supplementary Fig. S2 for details).

274 Three-dimensional structure of *L. blandensis* NrdB

The crystal structure of the dATP-inhibited complex of *L. blandensis* NrdB at 2.45 Å
resolution (PDB 5OLK) revealed a novel tetrameric arrangement hitherto not observed in the
RNR family, with approximate 222 point symmetry (Fig. 6a, Supplementary Table S1). Each
monomer consists of an ATP-cone domain (residues 1-103) joined by a short linker (104-
106) to a metal-binding α-helical core domain (residues 107-398) and a disordered C-
terminus (399-427). The latter two features are typical of the NrdB/F family. This domain
arrangement gives the NrdB monomer and dimer extended conformations that are
presumably flexible in solution (Fig. 6b). The dimer buries about 1100 Å^2 of solvent
accessible area on each monomer. The tetrameric arrangement is completely dependent on
interactions between the ATP cone domains, as no contacts are made between the core
domains in the two dimers that make up the tetramer (Fig. 6a).

The ATP-cone domain in *L. blandensis* NrdB is structurally very similar to the one recently
identified in the NrdA protein of *P. aeruginosa* (Johansson et al., 2016). The root-mean-
square deviation for 92 equivalent Cα atoms is 1.2 Å. The electron density unambiguously
confirms the *L. blandensis* ATP-cone's ability to bind two molecules of dATP, which it shares
with *P. aeruginosa* NrdA (Fig. 6c). Despite a local sequence identity of only 31% to *P.*
aeruginosa NrdA, all amino acids involved in binding both dATP molecules are conserved
(Fig. 6c). The two dATP molecules bind in a "tail-to-tail" arrangement that orients the base of
the "non-canonical" dATP towards the fourth, most C-terminal helix, an arrangement made
possible by the binding of a Mg^{2+} ion between the triphosphate moieties.

Remarkably, the interactions between the ATP-cones in *L. blandensis* NrdB are also very
similar to those seen in *P. aeruginosa* NrdA (Johansson et al., 2016), despite the fact that
the ATP-cone is attached to a structurally completely different core domain. The main
interactions occur between the last two helices α3 and α4 in respective ATP-cones (Fig. 6d).
A hydrophobic core in *L. blandensis* NrdB involving residues Met80, Ile92 and Ile93 in both
monomers is reinforced by salt bridges between residues Asp73, Asp76 in one monomer
and Lys89 in the other. The two domains bury ~510 Å^2 of solvent accessible area. This is
slightly less than the ~640 Å^2 buried in the equivalent interaction involving the ATP-cones of
P. aeruginosa NrdA. Within each ATP cone, helices α3 and α4 have the same relative

orientation, partly determined by an internal salt bridge between the conserved Asp73 and
Arg95, but the helix pair in *L. blandensis* NrdB is rotated relative to its counterpart in the
other ATP cone by about 15° compared to *P. aeruginosa* NrdA, which reduces the number
of possible interactions between them. Interestingly, the dATP-induced tetramer leaves free
the surfaces of both dimers of *L. blandensis* NrdB that are thought to interact with the NrdA
subunit in productive RNR complexes, which implies that one or two dimers of *L. blandensis*
NrdA could attach to these surfaces in a near-productive fashion in $\alpha_2\beta_4$ or $\alpha_4\beta_4$ complexes
(Fig. 6a).

Two metal ions are found to bind to each of the monomers of *L. blandensis* NrdB.
Comparison of their B-factors with those of surrounding atoms suggest that they are fourth
row elements with close to full occupancy, but does not enable us to distinguish between
Mn, Fe or Ca. No metal ions were added to the protein preparation, but crystals were
obtained in 0.2 M $CaCl_2$. The coordination distances are long for an RNR metal center, with
the smallest distances being 2.3–2.4 Å. The distance between metal ions varies between
3.4–3.8 Å in the four monomers (Supplementary Fig. S3), but the metal coordination is very
similar. Interestingly, Tyr207 is found near the metal site at the position expected for a
radical-carrying Tyr, but it is not hydrogen-bonded to the metal site, its hydroxyl group being
at around 6 Å from the side chain of Glu170. Tyr207 is H-bonded to the side chain of
Thr294, but the latter is not H-bonded to a metal center ligand. On the other side of the metal
site, Trp177 is H-bonded to the side chain of Glu263. This tryptophan is completely
conserved in the NrdBi subclass (Supplementary Fig. S4). It makes the same interaction as
Trp111 from *E. coli* NrdB, despite the fact that it is projected from the first helix of the 4-helix
bundle containing the metal center ligands rather than the second. The first helix of the
bundle has a very unusual distortion in the middle (Supplementary Fig. S5). Normally this is
an undistorted α-helix, but in *L. blandensis* NrdB, residues 171–175 form a loop that bulges
away from the metal site, with the exception of Lys174, whose side chain is projected
towards the metal site and is H-bonded to Glu170 (Fig. 6e). The significance of this distortion
is at present not clear.

The nature of the metallo-cofactor was addressed by X-band EPR spectroscopy and
catalytically active samples were analyzed at 5 - 32 K. The spectra revealed a mixture of
signals from low and high-valent manganese species (Fig. 7a). In particular, at 5 K a 6-line
signal attributable to low valent Mn (Mn^{II}) was clearly visible, overlaid with a complex
multiline signal with a width of approximately 1250G. Increasing the temperature to 32 K
resulted in a significant decrease of the latter signal, while the 6-line feature remained
relatively intense (Fig. 7a, top and middle). The shape, width and temperature dependence
of the multiline signal are all highly reminiscent of the signal reported for super-oxidized
manganese catalase as well as the 16-line signal observed during the assembly of the
dimanganese/tyrosyl radical cofactor in NrdF RNR, and is attributable to a strongly coupled
$Mn^{III}Mn^{IV}$ dimer ($S_{Total} = \frac{1}{2}$) (Fig. 7a, bottom) (Cotruvo, Stich, Britt, & Stubbe, 2013; Zheng,
Khangulov, Dismukes, & Barynin, 1994). The low valent species appeared to be weakly
bound to the protein, while the $Mn^{III}Mn^{IV}$ dimer was retained in the protein following desalting
(Supplementary Fig. S6). The observation of a high-valent Mn cofactor is consistent with the
Mn-dependent increase in catalytic activity of the enzyme and its inhibition by the addition of
Fe^{2+} (Fig. 7b-c), and is suggestive of a novel high-valent homodimeric Mn-cofactor. Indeed,
earlier calculations have shown that high-valent Mn-dimers have reduction potentials similar
to that of the tyrosyl radical in standard class I RNRs, but are hitherto unobserved in RNRs

(Roos & Siegbahn, 2011). A more detailed biophysical characterization of this cofactor is
currently ongoing.

Figure 6. Structure of tetrameric *L. blandensis* NrdB in complex with dATP. a) Overall structure of the NrdB tetramer. The two monomers of each dimer are colored in different shades of gray, while the N-terminal ATP-cones are colored orange and light blue respectively. The dATP molecules are shown in CPK representation. The curved lines at the top and bottom of each dimer core indicate the proposed binding area in the active $\alpha_2\beta_2$ complex of RNRs. b) Surface representation of the NrdB dimer, showing the presumably flexible nature of the ATP-cones in solution. The color scheme is as in panel a). c) Binding of two dATP molecules to the ATP-cone. The dATP molecules are shown as sticks. Residues involved in binding the two dATP molecules are labeled in blue and red respectively. Polar contacts are shown as dotted yellow lines. d) The interface between ATP-cones in the NrdB tetramer. Chains A and D are shown in light blue and orange respectively. Side chains of residues involved in the interface are labeled. The two dATP molecules closest to the interface are shown as sticks. Polar contacts are shown as yellow dotted lines. e) Structure of the metal site in chain C, which is representative of the others. Metal ions are shown as purple spheres. 2m|Fo|-D|Fc| electron density is shown as a grey mesh, contoured at 1.4 σ. The density for Lys174 is shown in red for clarity.

Figure 7. Type of dinuclear metal center of *L. blandensis* NrdB and metal-dependency of enzyme activity. (a) X-band EPR spectra of catalytically active, non-reconstituted, samples recorded at 5 K (*black*, top); 32 K (*red*, middle) (signal intensity multiplied by 3.7 for clarity; multiline spectrum obtained by subtraction of a scaled 32K spectrum from the 5K spectrum (*blue*, bottom) (signal intensity multiplied by 3 for clarity). Instrument settings: microwave frequency = 9.28 GHz; power = 1 mW; modulation amplitude = 10G; modulation frequency = 100 kHz. (b) Enzyme activity of NrdBΔ99 purified from heterologously expressed cultures grown with addition of different divalent metal ions as indicated; the Mn-sample was used for the EPR analysis. (c) Enzyme activity was measured after addition of a total concentration of 20 μM divalent metal ions to 10 μM of wild-type or NrdBΔ99 protein as

382 indicated. Enzyme activity without addition of metals was set as 100% and corresponded to
383 592 and 217 nmol mg^{-1} min^{-1} for wild type and NrdBΔ99 enzymes respectively. Error bars
384 indicate the standard deviation of three measurements.

L. blandensis NrdB binds two nucleotide molecules per ATP-cone

The finding that two dATP molecules were bound to the ATP-cone in the 3D structure
prompted us to investigate nucleotide binding using isothermal titration calorimetry (ITC).
Binding curves for dATP, ATP and dADP to NrdB, including a reverse titration of NrdB to
dATP, were all consistent with a single set of binding sites except for the ATP-cone deletion
mutant (NrdBΔ99), which did not bind nucleotides at all (Fig. 8). Using information available
from the 3D structure, a fixed stoichiometry of 2 was used to fit dATP binding to NrdB. The fit
indicated 59% active (i.e. nucleotide binding) protein. Fitting dADP and ATP with a 59%
proportion of active protein taken from the dATP experiment, we could calculate
stoichiometries of 2.2 and 2.1 for ATP and dADP respectively, indicating that each NrdB
monomer can bind two molecules of adenosine nucleotides (Fig. 8f). K_d for the three
different nucleotides (Fig. 8f) indicated a 38-fold and 15-fold lower affinity for ATP and dADP
compared to dATP. Thermodynamic parameters (Fig. 8f) indicated that the interactions are
enthalpy-driven, with negative ΔH values of -79, -44 and -103 for dATP, ATP and dADP
respectively.

protein	ligand	N	K_d (µM)	ΔH (kJ mol^{-1})	ΔG (kJ mol^{-1})	-TΔS (kJ mol^{-1})
NrdB	dATP	2*	1 ± 0.2	-79 ± 10	-33	47 ± 10
NrdB	ATP	2.2 ± 0.4	38 ± 2	-44 ± 4	-24	20 ± 4
NrdB	dADP	2.1 ± 0.2	15 ± 1.5	-103 ± 9	-26	77 ± 8
NrdBΔ99	dATP	ND	ND	ND	ND	ND

Figure 8: Representative ITC thermograms obtained by titration of dATP (a), ATP (c) and
dADP (d) into NrdB. Isothermal calorimetric enthalpy changes (upper panel) and resulting
binding isotherms (lower panel) are shown. Reverse titration of NrdB to dATP (b). Titration of
dATP to NrdBΔ99 (e). Thermodynamic parameters of ligand binding to NrdB (f). Binding
isotherms were fitted using a one-set-of sites binding model. Values are reported as the
mean ± SD of three titrations (and additional two reverse titrations for dATP). All titrations

were performed at 10°C as described in Materials and Methods. *=binding stoichiometry was
kept constant N=2, considering monomeric protein concentrations. ND=not detected.

410 Discussion

It is critically important for an organism to control the supply of dNTPs to allow fidelity in DNA
replication and repair (Mathews, 2006). Specificity regulation of RNR makes sure relative
concentrations of dNTPs fit the organism's DNA composition. On the other hand, activity
regulation assures absolute concentrations of dNTPs follow the different requirements
through the cell cycle (Hofer et al., 2012). Specificity regulation of RNRs is ubiquitous,
integrated in the catalytic subunit of the enzyme, and works via the classical homooligomeric
model of allosteric regulation (Andersson et al., 2000; Hofer et al., 2012; Larsson et al.,
2004; Reichard, 2010; Swain & Gierasch, 2006; Torrents et al., 2000; Zimanyi et al., 2016).
In contrast, the activity regulation is controlled by an accessory domain, the ATP-cone, and
works by affecting the distribution of the holoenzyme heteromeric complexes. Moreover, the
ATP-cone is only found in some RNRs, and appears to be gained by domain shuffling when
evolutionary selection favours it and lost when selection decreases (Lundin, Berggren,
Logan, & Sjöberg, 2015). These dynamics are further evidenced by the differences in
mechanisms recently discovered in class I RNRs (Ando et al., 2011; Ando et al., 2016;
Fairman et al., 2011; Johansson et al., 2016; Jonna et al., 2015). The current study was
prompted by the interesting observation that several radical-generating subunits of the NrdBi
subclass possess an N-terminal ATP-cone and that a few radical-generating subunits of the
NrdF subclass possess a C-terminal ATP-cone. Our discovery evokes several pertinent
questions: does the ATP-cone fused to a radical-generating subunit function as an allosteric
on/off switch; how does it affect the distribution of holoenzyme complexes under active and
inhibited conditions; is its structural mode of action similar to that of ATP-cones fused to the
catalytic subunit?

We have delineated the function of the ATP-cone that is N-terminally fused to the *L.*
blandensis NrdBi protein. In the presence of the positive effector ATP, *L. blandensis* NrdBi
was a dimer, which by interaction with the *L. blandensis* catalytic subunit NrdAi formed the
common active $\alpha_2\beta_2$ complex, also found in e.g. *E. coli*, *P. aeruginosa* and eukaryotic class I
RNRs. Binding of dATP to the ATP-cone instead promoted oligomerization of *L. blandensis*
NrdBi to an inactive β_4 complex, with a novel tetrameric structure revealed by
crystallography. This oligomerization is reminiscent of the "ring-shaped" α_4 and α_6 complexes
formed by dATP binding to the NrdA-linked ATP-cones in *P. aeruginosa* and eukaryotic
RNRs. When *L. blandensis* NrdAi was added to the dATP-loaded NrdBi tetramer, higher
molecular mass complexes of $\alpha_2\beta_4$ and $\alpha_4\beta_4$ appeared. The NrdA dimers presumably bind to
the NrdB tetramers in a 'nonproductive' orientation, which does not allow efficient electron
transfer between NrdA and NrdB. The crystal structure shows that the tetramerization of *L.*
blandensis NrdBi leaves the putative interaction surface for NrdAi free, which is consistent
with the possibility to form both $\alpha_2\beta_4$ and $\alpha_4\beta_4$ oligomers. However the structure does not
suggest the structural basis for a disruption of the cysteinyl radical generation pathway in
these non-productive complexes.

The structure of the dATP-loaded *L. blandensis* NrdB shows that it binds two dATP
molecules per ATP-cone. Both molecules bind to the same site and interact with each other
through a Mg^{2+} ion. Most allosterically regulated RNRs characterized so far bind only one

dATP molecule per ATP-cone. However, a novel class of ATP-cones that binds two dATP
molecules was recently characterized in *P. aeruginosa* NrdA (Johansson et al., 2016; Jonna
et al., 2015). The NrdB-linked ATP-cone of *L. blandensis* has sequence motifs characteristic
of this kind of ATP-cone. The structure confirms that both dATP molecules bind essentially
identically as in *P. aeruginosa* NrdA and that the ATP-cones make similar interactions to
each other, distinct from those made in the eukaryotic α_6 complexes. It has also been shown
in the RNR transcriptional regulator NrdR, that ATP and dATP bind with positive
cooperativity to its ATP-cone (McKethan & Spiro, 2013), implying binding of more than one
nucleotide molecule.

The ATP-cone in *L. blandensis* NrdBi offers a unique possibility to measure its binding
capacity for (deoxy)adenosine nucleotides. This has not been possible in earlier studied
RNRs, where the ATP-cone is bound to the α subunit that also possesses a binding site for
allosteric regulation of substrate specificity. ITC ligand binding studies confirmed that the *L.
blandensis* ATP-cone bound two dATP molecules, and showed that it also can bind two
molecules of ATP or two molecules of dADP. Structurally, binding of dADP is not surprising,
since the γ-phosphates of the dATP molecules make only one interaction with the protein,
through Arg102, and they only contribute 2 of the 6 coordinating atoms of the intervening
Mg^{2+} ion. Nonetheless, this is the first observation of dADP binding to the ATP-cone of an
RNR enzyme. dADP inhibits enzyme activity in a similar manner to dATP, but higher
concentrations are required. *In vivo*, deoxyribonucleoside diphosphates are rapidly
converted to triphosphates and cellular dADP concentrations are very low (Mathews, 2014;
Traut, 1994). Nevertheless, dADP is one of the products of *L. blandensis* RNR, and perhaps
local concentrations are higher. The ability of dADP to regulate the activity of *L. blandensis*
RNR may enable it to react more rapidly to changes in (deoxy)adenosine nucleotide
concentrations and provide the cell with a fitness advantage. Binding of deoxyadenosine di-
and monophosphates has been described for the ATP-cone in NrdR (Grinberg et al., 2006;
McKethan & Spiro, 2013).

Based on the variable occurrence of the ATP-cone in RNR catalytic subunits, we have
earlier hypothesized that its presence in RNRs is part of a dynamic process of gains and
losses on a relatively short evolutionary time scale (Lundin et al., 2015). This suggests that
the ATP-cone, contrary to what might be expected for a domain involved in allostery, does
not require a long evolutionary period of integration with a protein to contribute to regulation
and that it hence lends itself to a highly dynamic evolution of allosteric activity control
(Lundin et al., 2015). This hypothesis is nicely supported by the N-terminally positioned ATP-
cone, in the radical-generating subunit, NrdB, of *L. blandensis* RNR described here. No RNR
in the phylogenetic subclass NrdAi/Bi contains an ATP-cone in the catalytic subunit, and only
a minority – mostly encoded by Flavobacteria, a marine class of Bacteroidetes – have an
NrdB with an ATP-cone, suggesting the relatively recent acquisition of the ATP-cone to an
RNR subclass that is otherwise not activity regulated (Fig. 1). Moreover, in the Ib subclass,
which also lacks ATP-cones in the catalytic subunit, we detected a C-terminally positioned
ATP-cone in the radical-generating subunit. This was found in only two sequences from
closely related organisms and might hence be an example of an even more recent
evolutionary event.

The evidence we have presented here for an ATP/dATP-sensing master switch of the *L.
blandensis* NrdAi/NrdBi class I RNR, suggests a potential for the use of ATP-cones to
control the activity of engineered proteins. Multimeric enzymes could be inactivated through

sequestration of one member of an active complex, by control of dATP concentrations in the
reaction mixture.

The surprises did not end with the discovery of a functional master switch in the *L.*
blandensis NrdAi/NrdBi RNR. The active center of the radical generating subunit of class I
RNRs have earlier been found to consist either of an $Fe^{III}Fe^{III}$ or $Mn^{III}Mn^{III}$ pair coupled to a
tyrosine residue acting as long-term storage for the catalytically essential radical (Berggren,
Lundin, & Sjöberg, 2017; Cotruvo & Stubbe, 2012), or a $Fe^{III}Mn^{IV}$ center not coupled to an
amino acid radical (Griese, Srinivas, & Högbom, 2014). Although further analyses are
required to fully characterize the *L. blandensis* NrdBi active center, it appears to present a
fourth type, a high-valent $Mn^{III}Mn^{IV}$ center that lacks a suitably positioned radical storage
amino acid. The evolutionary flexibility displayed by the ATP-cone appears hence all but
equaled by the evolutionary tuning possibilities of the metal centers in the radical generating
subunit of class I RNR.

Materials and Methods

Cloning

DNA fragments encoding NrdAi (WP_009781766) and NrdBi (EAQ51288) were amplified by
PCR from *Leeuwenhoekiella blandensis* MED217 genomic DNA, isolated as described
previously (Pinhassi et al., 2006), using specific primers: NrdA: LBR1_For 5'-
cgagCATATGAGAGAAACACTACCAAAC-3' and LBR1_Rev 5'-
gcaaGGATCCTTAAGCTTCACAGCTTACA-3'. NrdB: LBR2_For 5'-
cgagCATATGAGTTCACAAGAGATCAAA-3', LBR2_REV 5'-
gcaaGGATCCTTAAAATAAGTCGTCGCTG-3', The PCR products were purified, cleaved
with NdeI and BamHI restriction enzymes and inserted into a pET-28a(+) expression vector
(Novagen, Madison, Wisconsin, USA). The obtained constructs pET-*nrdA* and pET-*nrdB*
contained an N-terminal hexahistidine (His) tag and thrombin cleavage site. To construct a
truncated NrdB mutant, lacking the entire ATP-cone domain, new forward primer
LBR2Δ99_For 5'-cgatCATATGCTGGAGCGTAAACAAAT-3' was used with LBR2_REV to
yield a pET-*nrdB*Δ99. The cloning process and the resulting construct was similar to that of
the wild type protein, except that it lacked sequence coding for the N-terminal 99 amino
acids.

Protein expression

Overnight cultures of *E. coli* BL21(DE3)/pET28a(+) bearing pET-*nrdA*, pET-*nrdB* and pET-
*nrdB*Δ99 were diluted to an absorbance at 600 nm of 0.1 in LB (Luria-Bertani) liquid medium,
containing kanamycin (50 µg/ml) and shaken vigorously at 37°C. At an absorbance at 600
nm of 0.8, isopropyl-β-D-thiogalactopyranoside (Sigma) was added to a final concentration
of 0.01 mM for NrdA expression and 0.5 mM for NrdB and NrdBΔ99 expression. For
particular experiments, 0.5 mM $MnSO_4$ or 0.5 mM $FeNH_4(SO4)_2$ or the combination of both
metals (0.4 mM and 0.25 mM respectively) were added to NrdBΔ99 cultures. The cells were
grown overnight at 14°C for NrdA expression and 20°C for NrdB and NrdBΔ99 expression
and harvested by centrifugation.

Protein purification

The cell pellet was resuspended in lysis buffer: 50 mM Tris-HCl pH 7.6 containing 300 mM NaCl, 20% glycerol, 10 mM imidazole, 1 mM PMSF. Cells were disrupted by high pressure homogenization and the lysate was centrifuged at 18,000 × g for 45 min at 4°C. The recombinant His-tagged protein was first isolated by metal-chelate affinity chromatography using ÄKTA prime system (GE Healthcare): the supernatant was loaded on a HisTrap FF Ni Sepharose column (GE Healthcare), equilibrated with lysis buffer (w/o PMSF), washed thoroughly with buffer and eluted with buffer containing 500 mM imidazole. Further purification was accomplished by fast protein liquid chromatography (FPLC) on a 125 ml column packed with Superose 12 Prep Grade or HiLoad 16/600 Superdex 200 pg column (GE Healthcare) using ÄKTA prime system, equilibrated with buffer containing 50 mM Tris-HCl pH 7.6, 300 mM NaCl, 10-20% glycerol. Eluted protein was collected. In the case of NrdA, all purification steps were performed in the presence of 2 mM DTT. Protein concentration was determined by measuring the UV absorbance at 280 nm based on protein theoretical extinction coefficients 91,135, 46,870 and 39,420 $M^{-1} cm^{-1}$ for NrdA, NrdB and NrdBΔ99 respectively. Proteins were concentrated using Amicon Ultra-15 centrifugal filter units (Millipore), frozen in liquid nitrogen and stored at -80°C until used.

For crystallization, NrdB was subsequently cleaved by thrombin (Novagen) to remove the hexahistidine tag. 41 mg NrdB was incubated with 25 U thrombin at 6°C for 3 hours, in a buffer containing 40 mM Tris-HCl pH 8.4, 150 mM NaCl, and 2.5 mM $CaCl_2$ in a total volume of 50 ml. Subsequently, imidazole to a final concentration of 20 mM was added and the reaction mixture was applied to a HisTrap FF Ni Sepharose column in a buffer containing 20 mM imidazole. Unbound protein was collected, concentrated and further purified by FPLC (see above) in a buffer containing Tris-HCl pH 7.6, 300 mM NaCl, and 10% glycerol. The thrombin-cleaved NrdB contained three additional residues (GlySerHis) that originated from the cleavage site of the enzyme at its N-terminus. Protein purity was evaluated by SDS–PAGE (12%) stained with Coomassie Brilliant Blue.

For EPR measurements, NrdBΔ99 was frozen in liquid nitrogen in EPR tubes directly after imidazole elution from the HisTrap column. Additional EPR samples were frozen after desalting the protein using PD-10 desalting columns (GE Healthcare).

Enzyme activity assays

Enzyme assays were performed at room temperature in 50 mM Tris-HCl at pH 8 in volumes of 50 µl. Reaction conditions, giving maximal activity were determined experimentally. In a standard reaction the constituents were; 10 mM DTT, 10 mM MgAc, 10 mM KCl, 0.8 mM (or when indicated 3 mM) CDP, and various concentrations of allosteric effectors ATP, dATP or dADP. Mixtures of 0.5 µM NrdA and 2 µM wild type NrdB or NrdBΔ99 (for determination of s-site K_L) (Fig. 3a-b) or 0.5 µM of NrdB and 2 uM NrdA (for determination of a-site K_L), (Fig. 3 c-f) were used. Some components were explicitly varied in specific experiments. Generally, 0.8 mM CDP was used as substrate. High CDP concentration (3 mM) was used for dADP titrations, to exclude potential product inhibition by binding of dADP to the active site. When dTTP was used as an s-site effector, 0.5 mM or 0.8 mM GDP was used as substrate. In four substrate assays, the four substrates CDP, ADP, GDP and UDP were simultaneously present in the mixture at concentrations of 0.5 mM each. The substrate mixture was added

last to start the reactions. Certain assays were performed in the presence of $Mn(CH_3COO)_2$
or $FeNH_4(SO4)$: 20 µM of the indicated metal was added to 10 µM NrdB or NrdBΔ99 protein,
incubated for 10 minutes, mixed with NrdA and added to the reaction mixture.

Enzyme reactions were incubated for 10-30 minutes and then stopped by the addition of
methanol. The chosen incubation time gave a maximum substrate turnover of 30%.
Substrate conversion was analyzed by HPLC using a Waters Symmetry C18 column (150 ×
4.6 mm, 3.5 µm pore size) equilibrated with buffer A. 25 µl samples were injected and eluted
at 1 ml/min with a linear gradient of 0–100% buffer B (buffer A: 10% methanol in 50 mM
potassium phosphate buffer, pH 7.0, supplemented with 10 mM tributylammonium
hydroxide; buffer B: 30% methanol in 50 mM potassium phosphate buffer, pH 7.0,
supplemented with 10 mM tributylammonium hydroxide). Compound identification was
achieved by comparison with injected standards. Relative quantification was obtained by
peak height measurements in the chromatogram (UV absorbance at 271 nm) in relation to
standards. Specific activities of either NrdA (Fig. 2 a-b) or NrdB (Fig. 2 c-f) were determined.
Specific activities varied between protein preparations. In some cases the data was
standardized to activity percent, where 100% was determined as maximum enzyme activity
in a specific condition.

From a direct plot of activity versus concentration of effector, the K_L values for binding of
effectors to the s-site and the a-site, were calculated in SigmaPlot using the equation:

$v = V_{max} \times [dNTP] / (K_L + [dNTP])$

and K_i for non-competitive dATP inhibition at NrdB was calculated in Sigmaplot using the
equation:

$v = V_{max} / (1 + ([dNTP]/K_i)$

604 GEMMA analysis

In GEMMA, biomolecules are electrosprayed into gas phase, neutralized to singly charged
particles, and the gas phase electrophoretic mobility is measured with a differential mobility
analyzer. The mobility of an analyzed particle is proportional to its diameter, which therefore
allows for quantitative analysis of the different particle sizes contained in a sample
(Kaufman, Skogen, Dorman, Zarrin, & Lewis, 1996). The GEMMA instrumental setup and
general procedures were as described previously (Rofougaran et al., 2008). NrdA, NrdB and
NrdBΔ99 proteins were equilibrated by Sephadex G-25 chromatography into a buffer
containing 100 mM ammonium acetate, pH 7.8. In addition, 2 mM DTT was added to the
NrdA protein solutions to increase protein stability. Prior to GEMMA analysis, the protein
samples were diluted to a concentration of 0.025–0.1 mg/ml in a buffer containing 20 mM
ammonium acetate, pH 7.5, 0.005% (v/v) Tween 20, nucleotides (when indicated), and
magnesium acetate (equimolar to the total nucleotide concentration), incubated for 5 min at
room temperature, centrifuged and applied to the GEMMA instrument. Protein
concentrations higher than normally recommended for GEMMA were needed to see the
larger oligomeric complexes and the experiments to measure NrdA-NrdB interactions were
run at as low flow rate as possible (driven by 1.4 Psi pressure) to minimize false interactions
that may appear with elevated protein concentration if the flow-rate recommended by the

manufacturer is used (3.7 Psi). The lower flow-rates give less sensitivity, though, and a flow
rate driven by 2 Psi was sufficient to prevent in most experiments.

Analytical size exclusion chromatography

Fast protein liquid chromatography on a Superdex 200 PC 3.2/30 column (with a total
volume of 2.4 ml) and ÄKTA prime system (GE Healthcare) was performed. The column was
equilibrated with SEC buffer containing 50 mM Tris-HCl pH 8, 50 mM KCl, 10% glycerol, 10
mM magnesium acetate, 2 mM DTT and when applicable either 3 mM ATP or 0.1-0.5 mM
dATP. 50 µL samples containing NrdA, NrdB or both subunits in the presence or the
absence of indicated amounts of nucleotides, were incubated for 10 minutes in room
temperature, centrifuged and applied to the column with a flow rate of 0.07 ml/min. When
nucleotides were added to proteins, they were also included in the buffer at the same
concentration to avoid dissociation of nucleotide-induced protein complexes during the run.
Varying concentrations of proteins were used in the range of 10-113 µM and 5-150 µM for
NrdA and NrdB respectively. For complex formation, 10-20 µM NrdA and 10-40 µM NrdB
were used in ratios of 1:1 or 1:2. Representative SEC chromatograms in which 20 µM NrdA,
20 µM NrdB or a mixture of 25 µM and 50 µM NrdA and NrdB respectively are shown in Fig.
5. Molecular weight was estimated based on a calibration curve, derived from globular
protein standards using high molecular weight SEC marker kit (GE Healthcare). Standard
deviations were calculated from at least 3 SEC experiments.

EPR measurements

642 Measurements were performed on a Bruker ELEXYS E500 spectrometer using an ER049X
643 SuperX microwave bridge in a Bruker SHQ0601 cavity equipped with an Oxford Instruments
644 continuous flow cryostat and using an ITC 503 temperature controller (Oxford Instruments).
645 Measurement temperatures ranged from 5 to 32 K, using liquid helium as coolant. The
646 spectrometer was controlled by the Xepr software package (Bruker).

Bioinformatics

RNR protein sequences were collected and scored using HMMER (Eddy, 2011) HMM
profiles in the RNRdb (http://rnrdb.pfitmap.org). ATP-cones were identified with the Pfam
ATP-cone HMM profile: PF03477. Sequences representing the diversity of NrdB were
selected by clustering all NrdB sequences from RefSeq at an identity threshold of 70% using
VSEARCH (Rognes, Flouri, Nichols, Quince, & Mahe, 2016). Sequences were aligned with
ProbCons (Do, Mahabhashyam, Brudno, & Batzoglou, 2005) and reliable positions for
phylogenetic reconstruction were manually selected. A maximum likelihood phylogeny was
estimated with FastTree 2 (Price, Dehal, & Arkin, 2010).

Isothermal titration calorimetry (ITC) measurements

Isothermal titration calorimetry (ITC) experiments were carried out on a MicroCal PEAQ-ITC
system (Malvern Instruments Ltd) in a buffer containing 25 mM HEPES (pH 7.65), 150 mM
NaCl, and 10% glycerol, 2 mM tris(2-carboxyethyl)phosphine, and 5 mM $MgCl_2$.
Measurements were done at 10°C. The initial injection volume was 0.4 µl over a duration of
0.8 s. All subsequent injection volumes were 2-2.5 µl over 4-5 s with a spacing of 150 s
between the injections. Data for the initial injection were not considered. For dATP binding

analysis, the concentration of NrdB in the cell was 12 µM and dATP in syringe 120 or 140
µM. Reverse titrations were performed with 113 µM NrdB in the syringe and 12 or 30 µM
dATP in the cell. For titration of dATP into NrdBΔ99, protein concentration in the cell and
dATP concentration in the syringe were 50 µM and 900 µM respectively. For dADP binding
analysis, the NrdB concentration in the cell was 20-50 µM and ligand concentrations in the
syringe were 500-750 µM. For titration of ATP into NrdB, cell and syringe concentrations
were 50 and 1600 µM respectively. The data were analyzed using the built-in one set of
sites model of the MicroCal PEAQ-ITC Analysis Software (Malvern Instruments Ltd). A fixed
ligand/protein stoichiometry of 2 was used for dATP to NrdB titrations. Standard deviations in
thermodynamic parameters, N and K_d were estimated from the fits of three different
titrations.

674 Crystallization and data collection

The purified NrdB, digested by thrombin to remove the hexahistidine tag (see above), was
used for crystallization. The protein at a concentration of 9.6 mg/ml was mixed with 20 mM
MgCl$_2$, 2 mM TCEP and 5 mM dATP, incubated for 30 minutes and used for setting up drops
using commercially available screens. An initial crystal hit was obtained by the sitting drop
vapor diffusion method with a protein to reservoir volume ratio of 200:200 nL and incubated
with a 45 µl reservoir in a Triple Drop UV Polymer Plate (Molecular Dimensions, UK). A
Mosquito nanoliter pipetting robot (TTP Labtech, UK) was used to set up drops, which were
imaged by the Minstrel HT UV imaging system (Rigaku Corporation, USA) available at the
Lund Protein Production Platform (LP3). Crystals were obtained with a reservoir containing
0.2 M CaCl$_2$, 0.1 M Tris pH 8.0 and 20% w/v PEG 600 (condition #57 of the PACT screen).
The crystals were then further optimized using an additive screen (Hampton Research, USA)
and diffraction quality crystals were obtained within 1 week from a crystallization solution
containing an additional 3% 6-aminohexanoic acid. Crystals were picked up directly from the
drop without cryoprotectant and data were collected at 100 K using the ID23-1 beamline at
the ESRF, Grenoble, France.

690 Structure determination and model building

The diffraction images were integrated using the program XDS (Kabsch, 2010) and scaled
using the program Aimless (Evans & Murshudov, 2013) from the CCP4 package (Winn et
al., 2011). The structure was solved by molecular replacement (MR) in two steps, using
Phaser (McCoy et al., 2007). First the structure of NrdBΔ99 was solved to 1.7 Å resolution
using the most homologous structure in the PDB, that of NrdF from *Chlamydia trachomatis*
(1SYY) (Högbom et al., 2004). This structure was rebuilt manually in Coot (Emsley,
Lohkamp, Scott, & Cowtan, 2010) and using Buccaneer (Cowtan, 2006) then refined to
convergence using Refmac5 (Murshudov, Vagin, & Dodson, 1997) and Buster (Bricogne,
2016). Full details of this structure will be presented elsewhere. In the second step, a multi-
body molecular replacement search was carried out using four copies of NrdBΔ99 and four
copies of the ATP-cone from *P. aeruginosa* NrdA (Johansson et al., 2016) prepared by side
chain truncation using Chainsaw (Stein, 2008). A single solution was found in which all 8
bodies were placed, with a translation function Z score (TFZ) of 12.2. After rearrangement of
the ATP-cones from the MR solution to the N-termini of their respective core domains, the

structure was refined using Refmac5. Some autobuilding was performed using Buccaneer
and manual rebuilding in Coot. Final refinement was done using Buster (Bricogne, 2016).

Automatically-generated non-crystallographic symmetry restraints were used. Geometry was
validated using the MolProbity server (Chen et al., 2010).

Acknowledgements

We want to thank Jarone Pinhassi and Sabina Arnautovic, Linneaus University, Kalmar,
Sweden, for the kind gift of *Leeuwenhoekiella blandensis* DNA, Yuliya Leontyeva, Jan-Olov
Persson and Rolf Sundberg, Department of Mathematics, Stockholm University, Sweden, for
excellent statistical analysis of ligand binding, Eva Muñoz and Angel Piñeiro, AFFINImeter,
Santiago de Compostela, Spain for valuable assistance with analysis of ITC titrations, Hanna
Jankevics Jones and John Stenson, Malvern Instruments Ltd, Malvern, UK for highly
appreciated help with protein biophysical characterization and Fredrik Tholander, Karolinska
Institute, Stockholm, Sweden, for generously providing his HPLC instrument for the enzyme
assays. We thank the crystallization facility at Lund Protein Production Platform for
crystallization trials and staff at EMBL Hamburg and ESRF beamlines for assistance with
data collection. This study was supported by grants from the Swedish Cancer Society (CAN
2016/670 to BMS), the Swedish Research Council (2016-01920 to BMS, 2016-04855 to
DTL), the Wenner-Gren Foundations (to BMS) and the Carl Trygger Foundation (to AH).
Work in the laboratory of GB is supported by the Swedish Research Council (621-2014-
5670), the Swedish Research Council for Environment, Agricultural Sciences and Spatial
Planning (213-2014-880) and the European Research Council (714102).

Accession codes

727 References accessions from Protein Data Bank: 5OLK

Author contributions

729 IRG, DL, IB, NM, GB, AH, DTL and BMS designed and analyzed experiments. IRG, DL, MH,
730 VRJ, GB, AH and DTL performed experiments. MC, CL and MS helped with setting up
731 activity assays. All authors contributed to the final manuscript.

References

Ahmad, M. F., & Dealwis, C. G. (2013). The structural basis for the allosteric regulation of
ribonucleotide reductase. *Prog Mol Biol Transl Sci, 117*, 389-410. doi:10.1016/b978-
0-12-386931-9.00014-3
Andersson, J., Westman, M., Hofer, A., & Sjöberg, B. M. (2000). Allosteric regulation of the
class III anaerobic ribonucleotide reductase from bacteriophage T4. *J Biol Chem,
275*(26), 19443-19448. doi:10.1074/jbc.M001490200
Ando, N., Brignole, E. J., Zimanyi, C. M., Funk, M. A., Yokoyama, K., Asturias, F. J., . . .
Drennan, C. L. (2011). Structural interconversions modulate activity of Escherichia
coli ribonucleotide reductase. *Proc Natl Acad Sci U S A, 108*(52), 21046-21051.
doi:10.1073/pnas.1112715108

Ando, N., Li, H., Brignole, E. J., Thompson, S., McLaughlin, M. I., Page, J. E., . . . Drennan,
C. L. (2016). Allosteric inhibition of human ribonucleotide reductase by dATP entails
the stabilization of a hexamer. *Biochemistry, 55*(2), 373-381.
doi:10.1021/acs.biochem.5b01207
Aravind, L., Wolf, Y. I., & Koonin, E. V. (2000). The ATP-cone: an evolutionarily mobile, ATP-
binding regulatory domain. *J Mol Microbiol Biotechnol, 2*(2), 191-194.
Aye, Y., & Stubbe, J. (2011). Clofarabine 5'-di and -triphosphates inhibit human
ribonucleotide reductase by altering the quaternary structure of its large subunit. *Proc*
Natl Acad Sci U S A, 108(24), 9815-9820. doi:10.1073/pnas.1013274108
Berggren, G., Lundin, D., & Sjöberg, B.-M. (2017). Homo- and heterometallic dinuclear
manganese proteins: active site assembly. In M. K. Johnson & R. A. Scott (Eds.),
Metalloprotein Active Site Assembly: John Wiley & Sons, Ltd.
Bochner, B. R., & Ames, B. N. (1982). Complete analysis of cellular nucleotides by two-
dimensional thin layer chromatography. *J Biol Chem, 257*(16), 9759-9769.
Bricogne, G. B., E.; Brandl, M.; Flensburg, C.; Keller, P.; Paciorek, W.; Roversi, P.; Sharff,
A.; Smart, O.S.; Vonrhein, C.; Womack, T.O. (2016). BUSTER version 2.10.2.
Cambridge, United Kingdom: Global Phasing Ltd.
Brown, N. C., & Reichard, P. (1969). Role of effector binding in allosteric control of
ribonucleoside diphosphate reductase. *J Mol Biol, 46*(1), 39-55. doi:0022-
2836(69)90056-4 [pii]
Buckstein, M. H., He, J., & Rubin, H. (2008). Characterization of nucleotide pools as a
function of physiological state in Escherichia coli. *J Bacteriol, 190*(2), 718-726.
doi:10.1128/jb.01020-07
Chen, V. B., Arendall, W. B., 3rd, Headd, J. J., Keedy, D. A., Immormino, R. M., Kapral, G.
J., . . . Richardson, D. C. (2010). MolProbity: all-atom structure validation for
macromolecular crystallography. *Acta Crystallogr D Biol Crystallogr, 66*(Pt 1), 12-21.
doi:10.1107/s0907444909042073
Cotruvo, J. A., Jr., Stich, T. A., Britt, R. D., & Stubbe, J. (2013). Mechanism of assembly of
the dimanganese-tyrosyl radical cofactor of class Ib ribonucleotide reductase:
enzymatic generation of superoxide is required for tyrosine oxidation via a
Mn(III)Mn(IV) intermediate. *J Am Chem Soc, 135*(10), 4027-4039.
doi:10.1021/ja312457t
Cotruvo, J. A., Jr., & Stubbe, J. (2012). Metallation and mismetallation of iron and
manganese proteins in vitro and in vivo: the class I ribonucleotide reductases as a
case study. *Metallomics, 4*(10), 1020-1036. doi:10.1039/c2mt20142a
Cowtan, K. (2006). The Buccaneer software for automated model building. 1. Tracing protein
chains. *Acta Crystallogr D Biol Crystallogr, 62*(Pt 9), 1002-1011.
doi:10.1107/s0907444906022116
Crona, M., Avesson, L., Sahlin, M., Lundin, D., Hinas, A., Klose, R., . . . Sjöberg, B. M.
(2013). A rare combination of ribonucleotide reductases in the social amoeba
Dictyostelium discoideum. *J Biol Chem, 288*(12), 8198-8208.
doi:10.1074/jbc.M112.442434
Do, C. B., Mahabhashyam, M. S. P., Brudno, M., & Batzoglou, S. (2005). ProbCons:
Probabilistic consistency-based multiple sequence alignment. *Genome Research,*
15(2), 330-340. doi:10.1101/gr.2821705
Eddy, S. R. (2011). Accelerated profile HMM searches. *PLoS Comput Biol, 7*(10), e1002195.
doi:10.1371/journal.pcbi.1002195
Emsley, P., Lohkamp, B., Scott, W. G., & Cowtan, K. (2010). Features and development of
Coot. *Acta Crystallogr D Biol Crystallogr, 66*(Pt 4), 486-501.
doi:10.1107/s0907444910007493
Eriksson, M., Uhlin, U., Ramaswamy, S., Ekberg, M., Regnström, K., Sjöberg, B. M., &
Eklund, H. (1997). Binding of allosteric effectors to ribonucleotide reductase protein
R1: reduction of active-site cysteines promotes substrate binding. *Structure, 5*(8),
1077-1092.

Evans, P. R., & Murshudov, G. N. (2013). How good are my data and what is the resolution?
Acta Crystallogr D Biol Crystallogr, 69(Pt 7), 1204-1214.
doi:10.1107/s0907444913000061
Fairman, J. W., Wijerathna, S. R., Ahmad, M. F., Xu, H., Nakano, R., Jha, S., . . . Dealwis, C.
G. (2011). Structural basis for allosteric regulation of human ribonucleotide reductase
by nucleotide-induced oligomerization. *Nat Struct Mol Biol, 18*(3), 316-322.
doi:nsmb.2007 [pii]10.1038/nsmb.2007
Fernandez-Gomez, B., Richter, M., Schuler, M., Pinhassi, J., Acinas, S. G., Gonzalez, J. M.,
& Pedros-Alio, C. (2013). Ecology of marine Bacteroidetes: a comparative genomics
approach. *ISME J, 7*(5), 1026-1037. doi:10.1038/ismej.2012.169
Griese, J. J., Srinivas, V., & Högbom, M. (2014). Assembly of nonheme Mn/Fe active sites in
heterodinuclear metalloproteins. *J Biol Inorg Chem, 19*(6), 759-774.
doi:10.1007/s00775-014-1140-7
Grinberg, I., Shteinberg, T., Gorovitz, B., Aharonowitz, Y., Cohen, G., & Borovok, I. (2006).
The Streptomyces NrdR transcriptional regulator is a Zn ribbon/ATP cone protein that
binds to the promoter regions of class Ia and class II ribonucleotide reductase
operons. *J Bacteriol, 188*(21), 7635-7644. doi:JB.00903-06 [pii]10.1128/JB.00903-06
Gunasekaran, K., Ma, B., & Nussinov, R. (2004). Is allostery an intrinsic property of all
dynamic proteins? *Proteins, 57*(3), 433-443. doi:10.1002/prot.20232
Hofer, A., Crona, M., Logan, D. T., & Sjöberg, B. M. (2012). DNA building blocks: keeping
control of manufacture. *Crit Rev Biochem Mol Biol, 47*(1), 50-63.
doi:10.3109/10409238.2011.630372
Huang, M., Parker, M. J., & Stubbe, J. (2014). Choosing the right metal: case studies of
class I ribonucleotide reductases. *J Biol Chem, 289*(41), 28104-28111.
doi:10.1074/jbc.R114.596684
Högbom, M., Stenmark, P., Voevodskaya, N., McClarty, G., Gräslund, A., & Nordlund, P.
(2004). The radical site in chlamydial ribonucleotide reductase defines a new R2
subclass. *Science, 305*(5681), 245-248. doi:10.1126/science.1098419
Johansson, R., Jonna, V. R., Kumar, R., Nayeri, N., Lundin, D., Sjöberg, B. M., . . . Logan,
D. T. (2016). Structural mechanism of allosteric activity regulation in a ribonucleotide
reductase with double ATP cones. *Structure, 24*(6), 906-917.
doi:10.1016/j.str.2016.03.025
Jonna, V. R., Crona, M., Rofougaran, R., Lundin, D., Johansson, S., Brännström, K., . . .
Hofer, A. (2015). Diversity in overall activity regulation of ribonucleotide reductase. *J*
Biol Chem, 290(28), 17339-17348. doi:10.1074/jbc.M115.649624
Kabsch, W. (2010). XDS. *Acta Crystallogr D Biol Crystallogr, 66*(Pt 2), 125-132.
doi:10.1107/s0907444909047337
Kashlan, O. B., Scott, C. P., Lear, J. D., & Cooperman, B. S. (2002). A comprehensive
model for the allosteric regulation of mammalian ribonucleotide reductase. Functional
consequences of ATP- and dATP-induced oligomerization of the large subunit.
Biochemistry, 41(2), 462-474. doi:10.1021/bi011653a
Kaufman, S. L., Skogen, J. W., Dorman, F. D., Zarrin, F., & Lewis, K. C. (1996).
Macromolecule analysis based on electrophoretic mobility in air: globular proteins.
Anal Chem, 68(11), 1895-1904. doi:10.1021/ac951128f
Kumar, D., Abdulovic, A. L., Viberg, J., Nilsson, A. K., Kunkel, T. A., & Chabes, A. (2011).
Mechanisms of mutagenesis in vivo due to imbalanced dNTP pools. *Nucleic Acids*
Res, 39(4), 1360-1371. doi:10.1093/nar/gkq829
Larsson, K. M., Jordan, A., Eliasson, R., Reichard, P., Logan, D. T., & Nordlund, P. (2004).
Structural mechanism of allosteric substrate specificity regulation in a ribonucleotide
reductase. *Nat Struct Mol Biol, 11*(11), 1142-1149. doi:10.1038/nsmb838
Lundin, D., Berggren, G., Logan, D. T., & Sjöberg, B. M. (2015). The origin and evolution of
ribonucleotide reduction. *Life (Basel), 5*(1), 604-636. doi:10.3390/life5010604
Mathews, C. K. (2006). DNA precursor metabolism and genomic stability. *FASEB J, 20*(9),
1300-1314. doi:20/9/1300 [pii]10.1096/fj.06-5730rev

Mathews, C. K. (2014). Deoxyribonucleotides as genetic and metabolic regulators. *FASEB J,*
28(9), 3832-3840. doi:10.1096/fj.14-251249
McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C., & Read,
R. J. (2007). Phaser crystallographic software. *J Appl Crystallogr, 40*(Pt 4), 658-674.
doi:10.1107/s0021889807021206
McKethan, B. L., & Spiro, S. (2013). Cooperative and allosterically controlled nucleotide
binding regulates the DNA binding activity of NrdR. *Mol Microbiol, 90*(2), 278-289.
doi:10.1111/mmi.12364
Murshudov, G. N., Vagin, A. A., & Dodson, E. J. (1997). Refinement of macromolecular
structures by the maximum-likelihood method. *Acta Crystallogr D Biol Crystallogr,*
53(Pt 3), 240-255. doi:10.1107/s0907444996012255
Nordlund, P., & Reichard, P. (2006). Ribonucleotide reductases. *Annu Rev Biochem, 75*,
681-706. doi:10.1146/annurev.biochem.75.103004.142443
Nussinov, R., & Tsai, C. J. (2013). Allostery in disease and in drug discovery. *Cell, 153*(2),
293-305. doi:10.1016/j.cell.2013.03.034
Pai, C. C., & Kearsey, S. E. (2017). A critical balance: dNTPs and the maintenance of
genome stability. *Genes (Basel), 8*(2). doi:10.3390/genes8020057
Pinhassi, J., Bowman, J. P., Nedashkovskaya, O. I., Lekunberri, I., Gomez-Consarnau, L., &
Pedros-Alio, C. (2006). Leeuwenhoekiella blandensis sp. nov., a genome-sequenced
marine member of the family Flavobacteriaceae. *Int J Syst Evol Microbiol, 56*(Pt 7),
1489-1493. doi:10.1099/ijs.0.64232-0
Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2 – approximately maximum-
likelihood trees for large alignments. *PLoS One, 5*(3), e9490.
doi:10.1371/journal.pone.0009490
Reichard, P. (2010). Ribonucleotide reductases: substrate specificity by allostery. *Biochem*
Biophys Res Commun, 396(1), 19-23. doi:10.1016/j.bbrc.2010.02.108
Rofougaran, R., Crona, M., Vodnala, M., Sjöberg, B.-M., & Hofer, A. (2008). Oligomerization
status directs overall activity regulation of the Escherichia coli class Ia ribonucleotide
reductase. *J Biol Chem, 283*(51), 35310-35318. doi:10.1074/jbc.M806738200
Rofougaran, R., Vodnala, M., & Hofer, A. (2006). Enzymatically active mammalian
ribonucleotide reductase exists primarily as an alpha6beta2 octamer. *J Biol Chem,*
281(38), 27705-27711. doi:M605573200 [pii]10.1074/jbc.M605573200
Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahe, F. (2016). VSEARCH: a versatile
open source tool for metagenomics. *PeerJ, 4*, e2584. doi:10.7717/peerj.2584
Roos, K., & Siegbahn, P. E. (2011). Oxygen cleavage with manganese and iron in
ribonucleotide reductase from Chlamydia trachomatis. *J Biol Inorg Chem, 16*(4), 553-
565. doi:10.1007/s00775-011-0755-1
Stein, N. (2008). CHAINSAW: a program for mutating pdb files used as templates in
molecular replacement. *J Appl Cryst, 41*, 641-643.
Swain, J. F., & Gierasch, L. M. (2006). The changing landscape of protein allostery. *Curr*
Opin Struct Biol, 16(1), 102-108. doi:10.1016/j.sbi.2006.01.003
Thelander, L., & Reichard, P. (1979). Reduction of ribonucleotides. *Annu Rev Biochem, 48*,
133-158. doi:10.1146/annurev.bi.48.070179.001025
Torrents, E. (2014). Ribonucleotide reductases: essential enzymes for bacterial life. *Front*
Cell Infect Microbiol, 4, 52. doi:10.3389/fcimb.2014.00052
Torrents, E., Buist, G., Liu, A., Eliasson, R., Kok, J., Gibert, I., . . . Reichard, P. (2000). The
anaerobic (class III) ribonucleotide reductase from Lactococcus lactis. Catalytic
properties and allosteric regulation of the pure enzyme system. *J Biol Chem, 275*(4),
2463-2471.
Torrents, E., Westman, M., Sahlin, M., & Sjöberg, B. M. (2006). Ribonucleotide reductase
modularity: atypical duplication of the ATP-cone domain in Pseudomonas
aeruginosa. *J Biol Chem, 281*(35), 25287-25296. doi:M601794200
[pii]10.1074/jbc.M601794200
Traut, T. W. (1994). Physiological concentrations of purines and pyrimidines. *Mol Cell*
Biochem, 140(1), 1-22.

Watt, D. L., Buckland, R. J., Lujan, S. A., Kunkel, T. A., & Chabes, A. (2016). Genome-wide
analysis of the specificity and mechanisms of replication infidelity driven by
imbalanced dNTP pools. *Nucleic Acids Res, 44*(4), 1669-1680.
doi:10.1093/nar/gkv1298
Winn, M. D., Ballard, C. C., Cowtan, K. D., Dodson, E. J., Emsley, P., Evans, P. R., . . .
Wilson, K. S. (2011). Overview of the CCP4 suite and current developments. *Acta*
Crystallogr D Biol Crystallogr, 67(Pt 4), 235-242. doi:10.1107/s0907444910045749
Zheng, M., Khangulov, S. V., Dismukes, G. C., & Barynin, V. V. (1994). Electronic structure
of dimanganese(II,III) and dimanganese(III,IV) complexes and dimanganese catalase
enzyme - a general EPR spectral simulation approach. *Inorg Chem, 33*(2), 382-387.
doi:10.1021/ic00080a030
Zimanyi, C. M., Chen, P. Y., Kang, G., Funk, M. A., & Drennan, C. L. (2016). Molecular basis
for allosteric specificity regulation in class Ia ribonucleotide reductase from
Escherichia coli. *Elife, 5*, e07141. doi:10.7554/eLife.07141