Genetic interactions between the chromosome axis-associated protein Hop1 and

homologous recombination determinants in *Schizosaccharomyces pombe*

Simon David Brown[1,2], Olga Dorota Jarosinska[1], and Alexander Lorenz[1,]*

[1]Institute of Medical Sciences (IMS), University of Aberdeen, Aberdeen AB25 2ZD, United

Kingdom

[2]Present address: MRC Institute of Genetics & Molecular Medicine, University of Edinburgh,

Edinburgh EH4 2XU, United Kingdom

Key words: *Schizosaccharomyces pombe*, meiosis, homologous recombination, chromosome

axis, Hop1

*Correspondence should be addressed to:

Alexander Lorenz

Institute of Medical Sciences (IMS)

University of Aberdeen

Foresterhill

Aberdeen AB25 2ZD

United Kingdom

Phone: +44 1224 437323

E-mail: a.lorenz@abdn.ac.uk

Alexander Lorenz ORCID 0000-0003-1925-3713

Abstract

Hop1 is a component of the meiosis-specific chromosome axis and belongs to the evolutionarily conserved family of HORMA domain proteins. Hop1 and its orthologs in higher eukaryotes are a major factor in promoting double-strand DNA break formation and inter-homolog recombination. In budding yeast and mammals they are also involved in a meiotic checkpoint kinase cascade monitoring the completion of double-strand DNA break repair. We used the fission yeast, *Schizosaccharomyces pombe*, which lacks a canonical synaptonemal complex to test whether Hop1 has a role beyond supporting the generation of double-strand DNA breaks and facilitating inter-homolog recombination events. We determined how mutants of homologous recombination factors genetically interact with *hop1*, studied the role(s) of the HORMA domain of Hop1, and characterized a bio-informatically predicted interactor of Hop1, Aho1 (SPAC688.03c). Our observations indicate that in fission yeast, Hop1 does require its HORMA domain to support wild-type levels of meiotic recombination and localization to meiotic chromatin. Furthermore, we show that *hop1Δ* only weakly interacts genetically with mutants of homologous recombination factors, and in fission yeast likely has no major role beyond break formation and promoting inter-homolog events. We speculate that after the evolutionary loss of the synaptonemal complex, Hop1 likely has become less important for modulating recombination outcome during meiosis in fission yeast, and that this led to a concurrent rewiring of genetic pathways controlling meiotic recombination.

Introduction

Chromosomes of diploid organisms are present in two sets, each derived from one of the parents. To reproduce sexually, this genetic material is halved exactly to produce haploid cells that contain only one chromosome set, the gametes. During fertilization, two gametes from different sexes fuse to form a zygote, re-establishing a double chromosome set. A specialized kind of cell division, called meiosis, is employed to halve the genetic material. In meiosis, one round of chromosome duplication is followed by two rounds of chromosome segregation. Remarkably, correct segregation of chromosomes in meiosis requires formation of double-strand DNA breaks (DSBs) by the conserved transesterase Spo11 (Lam and Keeney 2015), followed by repair through homologous recombination. Homologous recombination, in combination with sister chromatid cohesion, establishes physical connections (chiasmata) between maternal and paternal chromosomes that guide their proper segregation, and promotes reciprocal exchange (crossovers) of maternal and paternal genetic information (Hunter 2015). Careful regulation of this complex process is imperative to ensure the required number of physical links between the correct partner chromosomes, and that unrepaired DNA breaks are mended (Lam and Keeney 2015; Hunter 2015).

The repair of the Spo11-induced DSBs is started by DNA resection. During the initial step of DNA resection Spo11, which is covalently linked to the DNA ends, is endonucleolytically removed by the Mre11-Rad50-Nbs1 (Mre11-Rad50-Xrs2 in *Saccharomyces cerevisiae*) complex in cooperation with Ctp1 (Sae2 in *S. cerevisiae*) (Neale et al. 2005; Hartsuiker et al. 2009). Long-range resection then exposes substantial 3' tails which are used to invade homologous template DNA on the homologous chromosome or the sister chromatid (strand exchange) to mend the DSB by recombination (reviewed in Mimitou and

Symington 2011; Hunter 2015; Lorenz 2017). Key factors for the strand exchange process in

meiosis are the RecA-type recombinases Rad51 and Dmc1 (Shinohara et al. 1997; Grishchuk

and Kohli 2003) supported by their mediators Rad55-Rad57, Swi5-Sfr1, and Rlp1-Rdl1-Sws1

(SHU complex in *S. cerevisiae*) (Tsutsui et al. 2001; Grishchuk and Kohli 2003; Akamatsu et al.

2003; Martín et al. 2006; Sasanuma et al. 2013; Hong et al. 2013; Lorenz et al. 2014). These

Rad51/Dmc1-mediators are important for generating and stabilizing Rad51/Dmc1 coated

nucleoprotein filaments which then invade homologous template DNA (Sung 1997; Haruta et

al. 2006). The DNA on the homologous chromosome (or sister chromatid) being invaded is

opened as a displacement loop (D-loop) which can then be extended by DNA synthesis using

the invading sequence as primer (reviewed in Brown and Bishop 2014). The way in which D-

loops or recombination intermediates derived from it are processed, determines whether

repair will occur as a crossover (CO) or a non-CO, and Rad51/Dmc1-mediators play an

important role in this (reviewed in Brown and Bishop 2014; Hunter 2015; Lorenz 2017).

In fission yeast processing of most meiotic recombination intermediates into COs and

some non-COs depends on the structure-selective endonuclease Mus81-Eme1, the lack of

which results in meiotic catastrophe and strongly reduced spore viability (Osman et al. 2003;

Smith et al. 2003). Additionally, a sizeable fraction of non-COs is produced by the action of

the FANCM-ortholog Fml1, a DEAD/DEAH-box DNA helicase, and it has been shown that

Mus81-Eme1 and Fml1 work in parallel to process meiotic recombination intermediates

(Lorenz et al. 2012). In a wild-type situation Fml1 action seems to be curtailed by

Rad51/Dmc1-mediators, because weakening the strand exchange reaction by removing any

Rad51/Dmc1-mediator in a *mus81Δ* background ameliorates the strong spore viability defect

of *mus81Δ* without restoring CO formation; this rescue depends on the presence of Fml1

(Lorenz et al. 2012, 2014).

To facilitate the generation of DSBs and their correct repair during meiosis,

chromosomes undergo a progression of changes starting with the establishment of meiosis-

specific chromosome axes and, in most organisms, culminating in the formation of the

synaptonemal complex – an elaborate proteinaceous structure (reviewed in Cahoon and

Hawley 2016). Components of the synaptonemal complex seem to be critical for controlling

several aspects of meiotic chromosome metabolism, from pairing of homologous

chromosomes, to initiation of recombination and maturation of the physical interactions

between the homologous chromosomes (reviewed in Cahoon and Hawley 2016; Gray and

Cohen 2016). However, some organisms have overcome the necessity for a fully-fledged

synaptonemal complex and perform meiosis with a simplified version of it (reviewed in Loidl

2016). The fission yeast *Schizosaccharomyces pombe* is the best-studied organism executing

meiosis without a canonical synaptonemal complex (reviewed in Loidl 2006).

Despite lacking a synaptonemal complex, fission yeast meiotic chromosomes

assemble axes, called linear elements (Olson et al. 1978; Bähler et al. 1993; Molnar 2003),

consisting of proteins which are evolutionarily related to axial/lateral elements of the

synaptonemal complex (Lorenz et al. 2004), and building upon the existing mitotic

chromosome axis organization (reviewed in Ding et al. 2016; Iwasaki and Noma 2016). One

of these components is Hop1 which, in contrast to other meiotic chromosome axis factors, is

conserved on the protein sequence level with homologs from yeast to man (Lorenz et al.

2004; Rosenberg and Corbett 2015). In both *S. cerevisiae* and *Sz. pombe*, Hop1 localizes to

the meiosis-specific chromosome axis (Hollingsworth et al. 1990; Smith and Roeder 1997;

Lorenz et al. 2004), and deletion of *HOP1/hop1*[+] causes a decrease in meiotic recombination

and homologous chromosome pairing (Hollingsworth and Byers 1989; Loidl et al. 1994;

Latypov et al. 2010). Hop1 is characterized by a conserved N-terminal HORMA (Hop1-Rev7-

Mad2) domain (Aravind and Koonin 1998) (Fig. 1a) and is the founding member of the

meiotic HORMA domain (HORMAD) family proteins. In contrast to yeasts which contain only

one meiotic HORMAD (Hop1), higher eukaryotes contain multiple paralogs of meiotic

HORMADs: These include HIM-3, HTP-1, HTP-2, and HTP-3 in *C. elegans*; ASY1 and ASY2 in

Arabidopsis; and HORMAD1 and HORMAD2 in mammals (Zetka et al. 1999; Caryl et al. 2000;

Couteau and Zetka 2005; Martinez-Perez and Villeneuve 2005; Goodyer et al. 2008; Fukuda

et al. 2010). In fungi, plants, and mammals the HORMA domain is followed by several [S/T]Q

residues representing potential Tel1/ATM-Rad3/Mec1/ATR phosphorylation sites, which

enable meiotic HORMADs to act as structural adaptors for a meiotic checkpoint kinase

cascade (Sanchez-Moran et al. 2007; Carballo et al. 2008; Daniel et al. 2011; Kogo et al.

2012; Osman et al. 2018) (Fig. 1a). Fungal Hop1 proteins also contain a centrally located CxxC

Zn-finger motif (Hollingsworth et al. 1990) (Fig. 1a) which has been demonstrated to be

important for *in vitro* DNA-binding of *S. cerevisiae* Hop1 (Tripathi et al. 2007). Despite its *in*

vitro DNA-binding capabilities, Hop1 requires the presence of Rec10 (Red1 in *S. cerevisiae*) to

localize to chromosome axes *in vivo* (Smith and Roeder 1997; Lorenz et al. 2004); indeed,

Hop1 and Rec10/Red1 have been shown to physically interact with each other (de los Santos

and Hollingsworth 1999; Spirek et al. 2010). Rec10 in turn is recruited to chromosome axes

by the meiosis-specific cohesin subunit Rec11 upon phosphorylation by casein kinase 1

(Phadnis et al. 2015; Sakuno and Watanabe 2015).

Here, we investigate genetic interactions of *hop1* with meiotic recombination

determinants in fission yeast to test whether *hop1* has (a) role(s) beyond its established

function in promoting DSB formation (Latypov et al. 2010). Genetic interactions with factors

important for recombination intermediate processing (*mus81* and *fml1*) suggested a

potential, likely indirect, role for Hop1 in the strand exchange process. However, subsequent

epistasis analysis with the strand exchange factor *dmc1* and Rad51/Dmc1-mediators (*rad55,*

sfr1, and *rlp1*) did not support the hypothesized separate function of Hop1 in strand

exchange. Furthermore, we also show that the HORMA domain is required for wild-type

levels of meiotic recombination, and for localization to meiotic chromatin in *Sz. pombe*.

Implications of the evolutionary differences regarding meiotic HORMAD proteins and their

interactions are discussed.

Material and methods

***Sz. pombe* and *E. coli* culture conditions**

E. coli strains were grown on LB and SOC media – where appropriate containing 100 µg/ml

Ampicillin (Sambrook and Russell 2000). Competent cells of *E. coli* strains NEB10®-beta (New

England BioLabs Inc., Ipswich, MA, USA), and XL1-blue (Agilent Technologies, Santa Clara, CA,

USA) were transformed following the protocols provided by the manufacturers.

Sz. pombe strains used for this study are listed in Supplementary Table S1. Yeast cells

were cultured on yeast extract (YE), yeast nitrogen base glutamate (YNG) and Edinburgh

minimal media with glutamate as nitrogen source (EMMG) agar plates containing the

required supplements (concentration 250 mg/l on YE, 75 mg/l on YNG and EMMG). Crosses

were performed on malt extract (ME) agar containing the required supplements at a final

concentration of 50 mg/l, unless strains to be crossed contained plasmids, in which case they

were sporulated on sporulation agar with supplements (SPAS) plates with the necessary

supplements at 50 mg/l to maintain selection for the plasmids (Smith 2009; Sabatinos and

Forsburg 2010). Transformation of yeast strains was done using a standard lithium-acetate

procedure (Keeney and Boeke 1994) with the modifications described in Brown and Lorenz

(2016).

To obtain meiotic *Sz. pombe* cells for chromatin spreading and cDNA preparation

meiotic time-courses were executed according to a published protocol (Loidl and Lorenz

2009).

Yeast strain and plasmid construction

For specifics of strain and plasmid construction, please refer to Supplementary Materials.

Construction of the *kanMX6*-marked *hop1Δ-1* allele (Latypov et al. 2010) and the

kanMX6-marked *mug20$^+$-GFP* (Estreicher et al. 2012) have been described previously. The

natMX6-marked *hop1Δ-25* was derived from *hop1Δ-1* using an established marker swap

protocol (Sato et al. 2005; Lorenz 2015).

To produce *hop1*-alleles expressing a particular splice variant or the *hop1-ΔHORMAD*

mutant, we first partially deleted *hop1* retaining the coding sequence directly upstream of

the first and directly downstream of the last intron using a *ura4* marker (Grimm et al. 1988).

The resulting strain carrying the *hop1-3* allele (UoA725) was subsequently transformed with

the cDNA of one of the two splice variants or a *hop1-ΔHORMAD* construct; candidate strains

were selected by resistance to FOA (5-fluoroorotic acid) and then subjected to further

testing.

Epitope-tagging of Hop1 with 13myc was achieved by cloning a transformation

cassette into pFA6a-*13myc-kanMX6* (Bähler et al. 1998) with sequences from the 3' end and

downstream flanking region of the *hop1* coding sequence. This transformation cassette was

amplified by PCR and transformed into strains UoA746, UoA868, and UoA938.

A deletion cassette for *aho1* was constructed by cloning flanking sequences of

SPAC688.03c into pAG25 up- and downstream of the *natMX4* marker (Goldstein and

McCusker 1999). The resulting plasmid (pALo116) was linearized by a restriction digest and

transformed into the standard lab strain FO652.

To over-express *aho1*[+], the coding sequence of *SPAC688.03c* was amplified by PCR

and cloned into pREP41-*eGFPC* (Craven et al. 1998).

To construct plasmids for over-expression of *hop1*[+] in *Sz. pombe* we cloned PCR

products representing genomic or cDNA versions of the *hop1* coding sequence into pJR-41XL

(Moreno et al. 2000). During DNA sequencing we found discrepancies with the *hop1* coding

sequence prediction on PomBase (https://www.pombase.org/, last accessed 03/06/2017)

(see Results section for details). To corroborate this initial finding we also cloned meiotic

cDNA clones from two independently constructed strains (UoA399 and UoA722) into pUC8.

All plasmid constructs (for sequences see supporting online material, Lorenz 2018),

as well as epitope-tagged, splice variant, and *hop1-ΔHORMAD* strains were verified by DNA

sequencing (Source BioScience plc, Nottingham, UK).

All DNA modifying enzymes (high-fidelity DNA polymerase Q5, restriction

endonucleases, T4 DNA ligase) and NEBuilder HiFi DNA Assembly Master Mix were obtained

from New England BioLabs. Oligonucleotides were supplied by Sigma-Aldrich Co. (St. Louis,

MO, USA).

Genetic and molecular biology techniques

Spot assays to monitor growth and genotoxin sensitivities were essentially performed as

described previously (Doe et al. 2000; Lorenz et al. 2009). Spot assays of deletion strains

were done on YE plates, and of strains harboring plasmids on EMMG plates containing the

required supplements. To repress gene expression from pREP41-type plasmids in strains

grown on EMMG, thiamine was added to a final concentration of 4 μM (Sabatinos and

Forsburg 2010).

Genetic recombination assays and spore viability testing by random spore analysis

have been described in detail (Osman et al. 2003; Lorenz et al. 2010, 2012, 2014; Sabatinos

and Forsburg 2010).

Meiotic cDNA was obtained from cells undergoing synchronized azygotic wild-type

meiosis. Aliquots of cell culture were drawn at several time-points during a meiotic time-

course by first extracting RNA using the NucleoSpin TriPrep kit (Macherey-Nagel GmbH & Co.

KG, Düren, Germany) and then reverse-transcribing the RNA into cDNA using the ProtoScript

II First Strand cDNA Synthesis kit (New England BioLabs) following the instructions of the

respective manufacturer.

Chromatin spreads, immunocytochemistry, and microscopy

Nuclear spreads of meiotic *Sz. pombe* cells were prepared as described previously (Loidl and

Lorenz 2009), with the only modification that Lallzyme MMX (Lallemand Inc., Montréal,

Canada) at a final concentration of 100 mg/ml was used as the sole enzyme in the

spheroplasting solution (Flor-Parra et al. 2014).

Immunostaining was performed as previously described (Loidl and Lorenz 2009) using

polyclonal rabbit α-myc (ab9106; Abcam PLC, Cambridge, UK) at a 1:500 dilution and

monoclonal rat α-GFP [3H9] (ChromoTek GmbH, Planegg-Martinsried, Germany) at a 1:100

dilution as primary antibodies. Antibody-bound protein was visualized using donkey α-rabbit

IgG AlexaFluor-555 (ab150062; Abcam) and donkey α-rat IgG AlexaFluor-488 (ab150153;

Abcam), both at a 1:500 dilution, as secondary antibodies conjugated to fluorophores. DNA

was stained by Hoechst 33342 (Molecular Probes, Eugene, OR, USA) at a final concentration

of 1 µg/ml.

Analysis was done using a Zeiss Axio Imager.M2 (Carl Zeiss AG, Oberkochen,

Germany) epifluorescence microscope equipped with the appropriate filter sets to detect

red, green, and blue fluorescence. Black-and-white images were taken with a Zeiss AxioCam

MRm CCD camera controlled by AxioVision 40 software v4.8.2.0. Images were pseudo-

coloured and overlayed using Adobe Photoshop CC (Adobe Systems Inc., San José, CA, USA).

Data presentation and statistics

Data presented as box-and-whisker plots were created in RStudio 1.0.136 (R version 3.3.1)

(http://www.r-project.org/) using the boxplot() function with its standard settings. The lower

and upper 'hinges' of the box represent the first and third quartile, respectively, and the

black bar within the box indicates the median (=second quartile). The 'whiskers' represent

the minimum and maximum of the range, unless they differ more than 1.5-times the

interquartile distance from the median. In the latter case, the borders of the 1.5-times

interquartile distance around the median are indicated by the 'whiskers' and values outside

this range ('outliers') are shown as open circles. Raw data and R scripts used for the

boxplot() function are available online as supporting material (Lorenz 2018).

Initially, all data involving multiple comparisons underwent an overall statistical

assessment using a Kruskal-Wallis one-way analysis of variance to verify that there indeed

are statistically significant differences ($\alpha < 0.05$) between data sets within one experiment.

All direct one-on-one statistical comparisons between control and experimental data were

done using a two-tailed Mann-Whitney U test. Both Kruskal-Wallis and Mann-Whitney U

tests are non-parametric and therefore do not assume normal distribution of data points

within a data set. Kruskal-Wallis analyses were performed in IBM SPSS Statistics 24.0.0.0

(International Business Machines Corporation, Armonk, NY, USA), and Mann-Whitney U tests

in G*Power 3.1.9.2 (Faul et al. 2007, 2009). P values listed in Supplementary Tables S2 and

S3 were determined with a statistical power $(1 - \beta)$ set to 0.8.

Results

Correction of the *Sz. pombe hop1* coding sequence

When constructing plasmids to over-express *hop1*[+] in fission yeast cells, sequencing of our

cDNA clones revealed major discrepancies compared to the annotated sequence on

PomBase (https://www.pombase.org/, last accessed 03/06/2017). These differences

resulted from the incorrect annotation of the position of the third intron (Figs. 1b, S1, S2).

Additionally, we detected two splice variants (sv) of *hop1*

(https://www.ebi.ac.uk/ena/data/view/LT963776;

https://www.ebi.ac.uk/ena/data/view/LT907816). The main cDNA version (*hop1-sv1*)

translated into a protein with 527 amino acids (AAs) in length which is one AA shorter than

the currently annotated one (528 AAs). The discrepancy affects AAs 138-156/157 in the core

region of the HORMA domain (Fig. 1). A second, shorter splice variant (*hop1-sv2*) seems to

use a downstream AG as an alternative splice acceptor site at the third intron, thereby

expanding it by another 21 bps (Figs. 1b, S1, S2). In a total of 25 cDNA clones from meiotic

cDNA preparations of two independently constructed strains drawn from different time-

points of a meiotic time-course, 23 were of the *hop1-sv1* type, one conformed to the *hop1-*

sv2 type, and one was aberrant representing a rather short form of *hop1* with a frame-shift

(Fig. 1c). Considering the scarcity of *hop1-sv2* in our samples, it could well represent a

splicing accident or even a PCR artefact.

Using ClustalΩ (Sievers et al. 2014) the translation of the experimentally determined

Hop1-sv1 HORMA domain sequence also produces a slightly better alignments with Hop1

HORMA domains of other *Schizosaccharomyces* (Fig. S3) species than the predicted one. The

region of discrepancy in the experimentally derived *Sz. pombe* Hop1 HORMA domain

(VCKFFNRLHESFIKLINVQKE) aligns without much gapping (only one AA position in *Sz.*

octosporus and *Sz. cryophilus* shows a gap) and 7 fully conserved or strongly similar positions

to its counterpart in the other *Schizosaccharomyces* species. The same region from the

predicted *Sz. pombe* Hop1 (VSYQRSERFVIKLFLSGNVKTE) produces a large gap in the

alignment with the *Sz. japonicus* sequence, receives a three AAs gap compared to *Sz.*

octosporus and *Sz. cryophilus*, and aligns with only 4 fully conserved or strongly similar

positions. Over the whole length of the HORMA domain in this multi-species comparison,

this results in 10 fully conserved and 34 strongly similar AA residues in the alignment for the

predicted *Sz. pombe* Hop1 sequence, and in 11 fully conserved and 35 strongly similar AA

residues with the experimentally derived one (Fig. S3). This demonstrates that the current

annotation of the *hop1* open reading frame on PomBase (https://www.pombase.org/) is

erroneous,   and   that   the   long,   predominant   splice   variant   (*hop1-sv1*;

https://www.ebi.ac.uk/ena/data/view/LT963776) we identified in our cDNA clones is the

correct transcript of *hop1*.

The long *hop1* splice variant *hop1-sv1* is sufficient for meiotic recombination

We were interested in establishing whether the long, predominant version of Hop1 (Hop1-

sv1) is fully functional, and whether solely expressing the rare short version Hop1-sv2

(lacking 7 AAs in the HORMA domain, see Fig. 1) causes a phenotype. Additionally, we

constructed a Hop1 version which lacks the HORMA domain altogether, Hop1-ΔHORMAD. In

S. cerevisiae removing the N-terminal 268 AAs of Hop1 – which includes the HORMA domain

– has previously been shown to abolish spore viability completely, however the C-terminal

fragment remaining is still capable of binding DNA *in vitro* (Khan et al. 2013). In contrast, the

M127K point mutation in the center of the HORMA domain of *C. elegans HTP-1*, one of four

meiotic HORMA-paralogs in the nematode, results in only a minor meiotic phenotype

despite HTP-1[M127K] not localizing to meiotic chromosome axes (Silva et al. 2014). The *Sz.*

pombe short form *hop1-sv2* is apparently not strongly expressed (if not a splicing accident or

PCR artefact) in a wild-type background and thus unlikely to play a (major) role. However, in

light of the distinct phenotypes of HORMA-domain mutants in budding yeast and *C. elegans*,

we wanted to use the *hop1-sv2* variant and the *hop1-ΔHORMAD* mutant to probe the role of

the HORMA domain for meiotic recombination and Hop1 localization to meiotic chromatin in

fission yeast. The 7 AAs (positions 157-163) missing in Hop1-sv2 are close to the boundary

between the HORMA core domain and the HORMA "safety-belt" region (Fig. 1a, b; Fig. S2)

(Rosenberg and Corbett 2015; West et al. 2017), and their loss might affect interaction of

Hop1 with the meiotic chromosome axis similar to nematode HTP-1[M127K] and also interfere

with Hop1's self-oligomerization. The *hop1-ΔHORMAD* strains serve as controls, to monitor

the effect removal of the complete HORMA domain has on meiotic recombination and Hop1

localization.

Previously, it has been shown that the full deletion of *hop1* causes a reduction in

both intragenic (gene conversion) and intergenic (crossover) recombination at several

genetic intervals in fission yeast with a concomitant increase in intrachromosomal

recombination (Latypov et al. 2010). The *hop1* deletion also leads to defects in homologous

pairing along chromosome I during meiosis (Latypov et al. 2010). These phenotypes can at

least be partially explained by a reduction in DSB formation and a preponderance of DSBs

being repaired from the sister chromatid rather than the homolog (Rothenberg et al. 2009;

Latypov et al. 2010). Using our meiotic recombination assay system (Lorenz et al. 2010)

which allows us to concomitantly measure spore viability, and the frequency of gene

conversion (GC), crossovers (COs) and COs with GC events we found a statistically significant

decrease in all recombination measures in a *hop1Δ* compared to a wild-type cross (Fig. 2).

Indeed, GC decreased 4.7-fold ($p = 4.53 \times 10^{-13}$) (Fig. 2b), COs 2.8-fold ($p = 3.61 \times 10^{-6}$) (Fig.

2c), and COs associated with GC events were reduced by 8.7 percentage points ($p = 0.01$)

(Fig. 2d) (see also Table S2). Overall, this reduction in recombination outcome did not

negatively affect spore viability (Table S2). Recombination of a strain expressing only the

long, predominant version *hop1-sv1* (from the *hop1*-promoter at its original locus without

any marker genes inserted) was indistinguishable from wild type, whereas replacing the

wild-type copy of *hop1* with *hop1-sv2* led to a moderate reduction in GC (1.4-fold, $p = 1.51 \times$

10^{-3}) and COs (1.7-fold, $p = 2.27 \times 10^{-3}$), but not in COs associated with GC events ($p = 0.76$)

compared to wild type (Fig. 2, Table S2). The *hop1-ΔHORMAD* in turn was indistinguishable

from a full deletion (Fig. 2, Table S2). These results indicate that in line with *hop1-sv1* being

the predominant, if not the exclusive, splice variant behaves like wild-type *hop1* containing

all its introns, that *hop1-sv2* lacking 21 nucleotides in the HORMA domain is not fully

functional producing a hypomorphic phenotype, and that removal of the complete HORMA

domain (*hop1-ΔHORMAD*) produces a null phenotype.

Hop1 localizes to meiotic chromatin and forms linear elements independent of a full-

length HORMA domain

As with organisms possessing a fully-fledged synaptonemal complex, Hop1 localizes to

meiotic chromosome axes, called linear elements, in fission yeast (Lorenz et al. 2004).

However, the deletion of *hop1* has little impact on linear element formation *per se* (Lorenz

et al. 2006). We were keen to understand whether the presence of the HORMA domain is

required for localization of Hop1 to linear elements. We used immunostaining on chromatin

spreads from meiotic fission yeast cells to detect heterozygously 13myc-tagged Hop1-sv1,

Hop-sv2, and Hop1-ΔHORMAD in homozygous *hop1-sv1*, *hop1-sv2*, and *hop1-ΔHORMAD*

strains, respectively. Hop1-sv1-13myc formed linear elements, and all four morphological

classes of linear elements (dots, threads, networks, and bundles) were observed (Fig. 3a-d)

(Bähler et al. 1993; Lorenz et al. 2004, 2006). This result is consistent with the wild-type

phenotype for meiotic recombination of *hop1-sv1* crosses. Intriguingly, Hop1-sv2-13myc was

indistinguishable from Hop1-sv1-13myc (Fig. 3e-h). Therefore, the loss of the 7 AAs close to

the boundary between the HORMA core domain and the HORMA "safety-belt" region in the

hop1-sv2 allele does not conspicuously influence recruitment of Hop1 to linear elements,

despite affecting meiotic recombination (Fig. 2b-d). Hop1-ΔHORMAD-13myc could not be

detected on meiotic chromatin (Fig. S4) (a GFP-tagged version of the linear element protein

Mug20 served as control to identify meiotic nuclei containing linear elements; (Estreicher et

al. 2012)), this is consistent with *hop1-ΔHORMAD* displaying a meiotic recombination

phenotype similar to *hop1Δ* (Fig. 2b-d).

Searching for Hop1 interactors

A bio-informatics screen using the *Pombe* Interactome (PInt) (Pancaldi et al. 2012) for

potential Hop1 interactors retrieved *SPAC688.03c* as the highest hit from an uncharacterized

open reading frame (10[th] highest overall). The gene product of *SPAC688.03c* localizes

preferentially to the nucleus (Matsuyama et al. 2006) (Fig. S5) and is upregulated at the

onset of meiosis (Mata et al. 2002), which is consistent with a potential role in meiotic

chromatin metabolism. Notably, a point mutation in the human ortholog of *SPAC688.03c*,

AMMECR1, is associated with Alport syndrome (A), mental retardation (M), midface

hypoplasia (M), and elliptocytosis (E) (Vitelli et al. 1999; Andreoletti et al. 2017). Therefore,

we investigated the phenotype of a *SPAC688.03c* (from here on *aho1*, for *AMMECR1-*

homolog) deletion in vegetative and generative cells.

First, *aho1*Δ was assayed for changes in genotoxin sensitivity and meiotic

recombination outcome. The wild-type and *aho1*Δ strains exhibited similar levels of

resistance to the topoisomerase I-poison camptothecin (CPT), the alkylating agent methyl

methanesulfonate (MMS), the ribonucleotide reductase blocker hydroxyurea (HU), and

ultraviolet (UV) radiation (Fig. 4a). The deletion of *aho1* in a wild-type or *hop1*Δ background

did not affect spore viability (Fig. 4b). The meiotic recombination outcome of an *aho1*Δ

single mutant was similar to wild type, and an *aho1*Δ *hop1*Δ double mutant was

indistinguishable from a *hop1*Δ single mutant, except for a moderate 1.4-fold reduction in

gene conversion (p = 0.034) (Fig. 4c-e, Table S2). These results indicate that under standard

laboratory conditions *aho1*$^+$ is not required for DNA damage repair or meiotic

recombination. Therefore, Aho1 is not an essential interactor of Hop1 because it does not

mirror the phenotype of *hop1*Δ and we did not observe strong genetic interactions with

*hop1*Δ.

We tested whether over-expression of *aho1*$^+$ might uncover a DNA repair and/or

recombination role. The *aho1*$^+$ gene was put under the control of the thiamine-repressible

nmt1-promoter at the medium pREP41-level and tagged with GFP at its C-terminus. Spot

assays and recombination assays were performed after Aho1-GFP over-expression was

induced for 24 hours (see also Fig. S5). Cells over-expressing Aho1-GFP did not show

increased genotoxin sensitivities compared to cells containing an empty plasmid, cells over-

expressing GFP alone, or cells grown in the presence of thiamine (in which expression is not

induced) (Fig. 5a, b). Over-expression of Aho1-GFP also did not alter spore viability and

meiotic recombination outcome in comparison to GFP over-expression (Fig. 5c-f, Table S3),

indicating that a massive surplus of Aho1 protein has no negative impact on reproductive

success.

hop1 displays genetic interactions with the structure-selective endonuclease mus81 and

the DNA helicase fml1

Considering that the absence of *hop1* reduces recombination outcome in a similar fashion to

Rad51/Dmc1-mediator (*rad55-rad57*, *swi5-sfr1*, and *rlp1-rdl1-sws1*) mutants (Lorenz et al.

2012, 2014) (Fig. 2) we were wondering how *hop1* genetically interacts with the key meiotic

determinants of CO and non-CO formation, *mus81* and *fml1*.

The deletion of *hop1Δ* in a *mus81Δ* background improves spore viability from 1.85%

to 18.7% (Fig. 6, Table S2) similar to what has been seen in *mus81Δ* Rad51/Dmc1-mediator

double mutants (Lorenz et al. 2010, 2012, 2014). This rescue in spore viability is not

accompanied by a restoration of CO frequency. A *hop1Δ mus81Δ* mutant shows only 0.2% of

COs which is significantly different both from wild type ($p = 1.9 \times 10^{-7}$) and the *hop1Δ* single

mutant ($p = 2.7 \times 10^{-3}$) (Fig. 6c, Table S2). Also, the rate of COs associated with GC events at

0.71% is significantly lower in the *hop1Δ mus81Δ* double mutant than in the wild type

(63.0%, $p = 5.3 \times 10^{-12}$) or the *hop1Δ* single mutant (54.3%, $p = 3.4 \times 10^{-15}$) (Fig. 6d, Table S2).

Removal of *fml1+* increases COs associated with GC events to 72.2% (~10 percentage

points more than wild type) (Lorenz et al. 2012), but has little effect on other recombination

outcomes (Fig. 6). In *fml1Δ hop1Δ* double mutants, there are 72.6% of COs associated with

GC events which is significantly different from both wild type ($p = 3.4 \times 10^{-3}$) and a *hop1Δ*

single mutant (p = 4.9 × 10^{-9}), but indistinguishable from a *fml1Δ* single mutant (p = 0.78)

(Fig. 6d, Table S2). Overall CO formation is not changed compared to *hop1Δ* (p = 0.78), but

significantly different from wild type (p = 7 × 10^{-7}) and *fml1Δ* (p = 6.3 × 10^{-8}) (Fig. 6c, Table

S2). Also GC frequency is unchanged compared to *hop1Δ* (p = 0.49) (Fig. 6b, Table S2)

suggesting that in the absence of *hop1*$^{+}$ the DNA helicase Fml1 does not gain any additional

functionality.

hop1 does not show strong genetic interactions with Rad51/Dmc1-mediator mutants

Rad51/Dmc1-mediators are highly conserved, positive determinants of meiotic

recombination (Schwacha and Kleckner 1997; Ellermeier et al. 2004; Hayase et al. 2004;

Bleuyard et al. 2005; Hyppa and Smith 2010; Lorenz et al. 2012, 2014; Sasanuma et al. 2013).

In fission yeast the meiotic recombination phenotype of the *hop1Δ* crosses strongly

resembles that of Rad51/Dmc1-mediator mutants in that it reduces all meiotic

recombination outcomes and alleviates the effects of a *mus81Δ* mutants (see above).

Therefore, we were interested in how *hop1* genetically interacts with the mediators *sfr1*,

rad55, and *rlp1*, as well as *dmc1* itself. The deletion of *hop1*$^{+}$ leads to a notable reduction in

DSB formation (Rothenberg et al. 2009; Latypov et al. 2010), and some of the decrease in

recombination frequency in *hop1Δ* is likely due to low DSB formation. This reduction may be

unrelated to Rad51/Dmc1-mediator function, because absence of the Sfr1-cofactor *swi5*$^{+}$ or

of *dmc1*$^{+}$ does not appreciably reduce DSB formation (Young et al. 2004). We pose that in

the absence of *hop1*$^{+}$, which contributes to the establishment of a meiosis-specific

chromatin environment, successful sexual reproduction will become more reliant on the

mitotic repair factors Rad55 and Rlp1. Therefore, we expected a substantial negative genetic

interaction of *hop1* with *sfr1*, *rad55*, *rlp1*, and *dmc1*.

Indeed, *hop1Δ rad55Δ* and *hop1Δ rlp1Δ* double mutants displayed significantly lower

GC (p = 5.3 × 10^{-5} and p = 0.02, respectively) than a *hop1Δ* single mutant; the GC level is also

notably lower than in *rad55Δ* (p = 2.9 × 10^{-5}) and *rlp1Δ* (p = 1.3 × 10^{-4}) (Fig. 7a , Table S2)

(Lorenz et al. 2014). However, overall CO frequency and COs associated with GC events were

not significantly different from *hop1Δ* or *rad55Δ* in *hop1Δ rad55Δ*, and from *hop1Δ* or *rlp1Δ*

in *hop1Δ rlp1Δ* (Fig. 7b, c; Table S2). Deletion of *sfr1*$^{+}$ decreased GC rate by 7.5-fold in a

hop1Δ background to 0.036% (p = 1.4 × 10^{-8}) which is also significantly lower than the 0.13%

(p = 1 × 10^{-4}) found in an *sfr1Δ* single mutant (Fig. 7d, Table S2) (Lorenz et al. 2014). Other

recombination outcomes were indistinguishable from *hop1Δ* (Fig. 7e, f; Table S2). The

interaction of *hop1Δ* with *dmc1Δ* is completely epistatic (Fig. 7d-f, Table S2). Importantly, no

noteworthy decreases of spore viability have been observed in any of the double mutants

(Table S2). This stands in contrast to double mutant combinations of Rad51/Dmc1-mediators

with each other or with *dmc1Δ*; *dmc1Δ rad55Δ*, *dmc1Δ rlp1Δ*, *rad55Δ sfr1Δ* and *rlp1Δ sfr1Δ*

display strong synergistic reductions in spore viability (Lorenz et al. 2014). Overall, this

suggests that in fission yeast meiosis the absence of *hop1*$^{+}$ does not generate a situation

which necessitates the presence of a fully functional recombination machinery to produce

viable progeny.

Discussion

Here we characterize the genetic interactions of *hop1*, which encodes a meiotic HORMA

domain protein, with homologous recombination factors in fission yeast. Meiotic HORMADs

are components of the meiosis-specific chromosome axis which promote DSB formation to

allow for the efficient pairing of homologous chromosomes. At the same time HORMADs

attenuate undesirable recombination between sister chromatids (reviewed in Rosenberg

and Corbett 2015). In organisms possessing a synaptonemal complex, HORMADs support its

formation, and depending on the organism may mount a checkpoint arrest when DSB repair

is not completed or homologous chromosomes are not synapsed (Woltering et al. 2000;

Armstrong et al. 2002; Couteau and Zetka 2005; Martinez-Perez and Villeneuve 2005;

Carballo et al. 2008; Daniel et al. 2011; Rinaldi et al. 2017). *Sz. pombe* meiosis progresses

without the formation of a synaptonemal complex, therefore any roles related to the

synaptonemal complex will likely not apply. Indeed, DSB formation is reduced and inter-

sister recombination is increased in fission yeast *hop1*Δ (Rothenberg et al. 2009; Latypov et

al. 2010). Consistent with DSB reduction, decreased numbers of Rec7-foci (an essential co-

factor of Spo11) have been observed when *hop1* was absent (Lorenz et al. 2006). As a result

of DSB reduction, meiotic inter-homolog recombination frequencies are lower in *hop1*Δ than

in wild type (Latypov et al. 2010) (Fig. 2). Thus, the Hop1 function in promoting DSB

formation seems to be conserved in fission yeast. In budding yeast Hop1 also acts

downstream of DSB formation in promoting Dmc1-dependent inter-homolog recombination

by preventing Dmc1-independent Rad51-driven recombination (Niu et al. 2005; Carballo et

al. 2008). The suppression of inter-sister recombination is achieved by Hop1-dependent

activation of Mek1 which in turn phosphorylates and deactivates the Rad51-mediator Rad54,

and also phosphorylates and activates the meiosis-specific Rad51-inhibitor Hed1 (Tsubouchi

and Roeder 2006; Niu et al. 2009; Callender et al. 2016). Together, this regulatory network

has specific genetic consequences: (I) meiotic recombination cannot proceed without Dmc1

which causes a meiotic arrest resulting in an inability to sporulate, and (II) deleting *hop1* in a

*dmc1*Δ background restores spore formation by allowing Rad51-dependent DSB repair (Niu

et al. 2005; Carballo et al. 2008). In fission yeast the genetic interaction between *hop1* and

dmc1 looks quite different (Fig. 7) in that it is epistatic, however neither single mutant has as

strong a phenotype as the corresponding mutants in *S. cerevisiae* (Hollingsworth and Byers

1989; Bishop et al. 1992; Grishchuk and Kohli 2003; Latypov et al. 2010). Two features may

account for this apparent difference between the two yeasts: The budding yeast is probably

the only organism which shuts down Rad51 strand exchange activity via Hed1 (Tsubouchi

and Roeder 2006), and fission yeast employs both Rad51 and Dmc1 in their direct strand

exchange capacity (Grishchuk and Kohli 2003). Interestingly, the genetic interaction of

meiotic *Hormad* and *dmc1* genes generally does not show much conservation. In *Arabidopsis*

mutation of *asy1* or *dmc1* causes a strong reduction or absence of chiasmata, respectively,

leading to fertility defects. However, the *asy1 dmc1* double mutant looks like a *dmc1* single

mutant (Sanchez-Moran et al. 2007). In the mouse both $Dmc1^{-/-}$ and $Hormad1^{-/-}$ single

mutants undergo meiotic arrest at mid-pachytene, and so does the $Dmc1^{-/-}$ $Hormad1^{-/-}$

double mutant (Daniel et al. 2011), i.e. there is no reciprocal rescue as in *S. cerevisiae hop1Δ*

dmc1Δ. These disparate observations in various model organisms suggest that the genetic

networks controlling DSB repair and CO formation, and how these processes are integrated

with chromosome axis organization, have undergone considerable rewiring during evolution.

Because fission yeast relies on both Rad51 and Dmc1 to invade a repair template during

meiotic DSB repair, we also looked at the genetic interaction of *hop1* with Rad51/Dmc1-

mediator mutants. Apart from an additive effect on GC frequency in *hop1Δ rad55Δ*, *hop1Δ*

rlp1Δ, and *hop1Δ sfr1Δ* double mutants, which can easily be explained by the early functions

of Hop1 (reduced DSB formation and increased inter-sister recombination), other

recombination outcomes were not different from the *hop1Δ* single mutant (Fig. 7). This

indicates that in a fission yeast *hop1Δ* mutant gamete production does not become more

reliant on strand exchange than in a wild-type background.

Genetic interaction of *hop1Δ* with *mus81Δ* resembled that of *mus81Δ* combined with

mutants affecting strand exchange (Ellermeier et al. 2004; Lorenz et al. 2012, 2014), rescuing

mus81Δ's spore viability defect without rescuing its CO defect (Fig. 6). This interaction can

be explained by Hop1 promoting DSB formation and inter-homolog recombination: Fewer

DSBs and fewer inter-homolog events in *hop1Δ* decrease the probability of the production of

recombination intermediates requiring the action of Mus81. Additionally, the absence of

hop1 might allow DNA helicases to process recombination intermediates as non-COs more

efficiently than in a wild-type background.

It is well established that fungal Hop1 localizes to the chromosome axis during meiotic

prophase by physically interacting with the linear/lateral element protein Rec10/Red1 in

fission and budding yeast, respectively (de los Santos and Hollingsworth 1999; Spirek et al.

2010). Considering Hop1's central role in coordinating meiotic progression, it is not unlikely

that it would physically interact with previously uncharacterized meiotic players; indeed with

IHO1, a novel HORMAD1-interactor has recently been described in mice (Stanzione et al.

2016). A bio-informatical protein interaction-predictor PInt (Pancaldi et al. 2012) suggested

that the uncharacterized AMMECR1-homolog Aho1 might be a potential Hop1-interactor in

fission yeast. However, neither deletion nor over-expression of Aho1 indicates a clear

meiotic function, independent of the presence or absence of Hop1 (Figs. 4, 5).

One of the determining features of Hop1 is its HORMA domain whose function was

enigmatic until recently, when it was implicated in the recruitment of Hop1 to the meiotic

chromosome axis and in its self-assembly into larger homomeric complexes (Kim et al. 2014;

Rosenberg and Corbett 2015). We exploited a cDNA clone of *hop1* (*hop1-sv2*) which

produces a protein that lacks 7 AAs in the HORMA domain and constructed a version of hop1

lacking the complete HORMA domain (*hop1-ΔHORMAD*) to examine whether the HORMA

domain in fission yeast Hop1 is essential for its function and localization. Interestingly,

expressing Hop1-sv2 in place of wild-type Hop1 did not result in a notable reduction in

localization to chromatin, however it did cause a hypomorphic phenotype for GC and CO

frequency outcome, whereas the complete removal of the HORMA domain from Hop1

mimicked a deletion phenotype (Figs. 2, 3, S4). This suggests that the shortened HORMA

domain in Hop1-sv2 is still partially functional, because absence of Hop1's HORMA domain in

fission yeast completely abolishes localization to meiotic chromatin causing a *hop1Δ*

phenotype in recombination outcome.

It is very interesting how genetic pathways driving meiotic chromosome organization and

meiotic recombination are rewired during evolution. The fission yeast, which is lacking a

synaptonemal complex, but retains remnants of a meiosis-specific chromosome axis, is a

great model for probing the roles of conserved meiotic factors such as Hop1. Clearly, Hop1

fulfills an important meiotic function in promoting DSB formation and inter-homolog

recombination in this organism, but lacks regulatory roles in lieu of the synaptonemal

complex, which appears crucial in the majority of organisms performing meiosis with a

synaptonemal complex.

Acknowledgments

We are grateful to Shin-ichiro Hiraga, Franz Klein, Jürg Kohli, Josef Loidl, Walter W. Steiner,

Matthew C. Whitby, and the National BioResource Project (NBRP) Japan for providing strains

and/or plasmids, and to K.C. Chan, M. Roca, and D. Whyte for technical assistance. Kevin

Corbett, Nancy Hollingsworth, Takashi Kubota, and Josef Loidl are thanked for critically

reading a previous version of this manuscript. Microscopy was performed at the University

of Aberdeen Microscopy & Histology facility (Kevin Mackenzie). This work was supported by

the Biotechnology and Biological Sciences Research Council UK (BBSRC, Doctoral Training

Grant BB/F016964/1), and the University of Aberdeen (College of Life Sciences and Medicine

Start-up grant to AL).

Author contributions

578 SDB: unpublished essential reagents (yeast strains, plasmids), acquisition of data, analysis

579 and interpretation of data, revising the manuscript; ODJ: acquisition of data, revising the

580 manuscript; AL: conception and design, unpublished essential reagents (yeast strains,

581 plasmids), acquisition of data, analysis and interpretation of data, drafting and revising the

582 manuscript.

583
Figure 1. *Schizosaccharomyces pombe* **Hop1 domain structure and alignment between**
annotated and experimentally determined versions. (**a**) Schematic of the domain structure
of the experimentally derived sv1 (splice variant 1) version of Hop1 with the HORMA domain
at the N terminus (AAs 11-211), four potential [S/T]Q phosphorylation sites (red dots)
distributed between AAs 247 to 283, and residues 334-385 comprising the CxxC Zn-finger
domain (dark blue). The HORMA domain can be subdivided into a core domain (dark green)
represented by the most C-terminal ~150 AAs and a "safety-belt" region (light green;
residues 161-211) (Rosenberg and Corbett 2015; West et al. 2017). Positions of key AA
residues are indicated below the schematic. (**b**) ClustalΩ alignment (Sievers et al. 2014) of
part of the HORMA domain (starting at residue 125) of the current PomBase-annotation of
Hop1, and the two observed splice variants Hop1-sv1 and Hop1-sv2. Colouring of AAs
follows the standard ClustalΩ scheme according to physicochemical properties: red for small
and small & hydrophobic AAs (AVFPMILW); blue for acidic AAs (DE); magenta for basic AAs
(RK); and green for AAs containing hydroxyl-, sulfhydryl-, or amine-groups. Consensus
symbols below the alignment indicate full conservation (*) and conservation of groups with
strongly similar (:) or weakly similar (.) properties, for details see
http://www.ebi.ac.uk/Tools/msa/clustalo/help/faq.html. (**c**) Relative abundance of different
hop1 splice variants found in cDNA of the indicated diploid strains at given time-points of a
meiotic time-course.
603

604
Figure 2. Wild-type and *hop1* mutant levels of meiotic recombination at the *ade6-3083*

hotspot. (**a**) Schematic of the recombination assay at *ade6* (yellow) and its prevalent

outcomes: positions of *ade6* and the artificially introduced markers *ura4⁺-aim2* (green) and

his3⁺-aim (blue) on chromosome III are given in bp; positions of the point mutations in *ade6*

in the two parental strains are indicated in red (*ade6-3083*) and light blue (*ade6-469*). (**b**)

Frequency of gene conversion (GC) at *ade6-3083*, (**c**) frequency of crossovers (COs) between

ura4⁺-aim2 and *his3⁺-aim*, and (**d**) percentage of COs between *ura4⁺-aim2* and *his3⁺-aim*

associated with a GC event at *ade6-3083* in crosses of wild type (WT) ALP733×ALP731 (n =

12), *hop1Δ* UoA200×UoA199 (n = 17), *hop1-sv1* UoA766×UoA767 (n = 10), *hop1-sv2*

UoA853×UoA854 (n = 12), and *hop1-ΔHORMAD* UoA945×UoA944 (n = 12); n indicates the

number of independent crosses (see also Table S2).

617
Figure 3. Linear element morphology delineated by immunostaining of Hop1-13myc on
chromatin spreads from meiotic fission yeast cells. Linear elements as seen by
immunostaining against Hop1-13myc develop from dots via threads to networks and
bundles in h^{+N}/h^{-smt0} *hop1-sv1/hop1-sv1::13myc-kanMX6 ade6-M210/ade6-M216* (UoA878)
(**a-d**) or h^{+N}/h^{-smt0} *hop1-sv2/hop1-sv2::13myc-kanMX6 ade6-M210/ade6-M216* (UoA930) (**e-**
h). Hop1-13myc is shown in red and Hoechst 33342-stained DNA in blue in the merge panels.
Scale bar represents 2 μm.
625

626
Figure 4. The role of *aho1⁺* in genotoxic stress resistance and meiotic recombination. (a)

Spot assays comparing wild type (WT, FO652) and *aho1Δ* (UoA423) for sensitivity to a range

of genotoxins; number of cells plated in each spot is indicated at the bottom of the figure.

(**b**) Percent spore viability, (**c**) frequency of gene conversion (GC) at *ade6-3083*, (**d**) frequency

of crossovers (COs) between *ura4⁺-aim2* and *his3⁺-aim*, and (**e**) percentage of COs between

ura4⁺-aim2 and *his3⁺-aim* associated with a GC event at *ade6-3083* in crosses of wild type

(WT) ALP733×ALP731 (n = 12), *hop1Δ* UoA200×UoA199 (n = 16), *aho1Δ* UoA424×UoA427 (n

= 6), and *aho1Δ hop1Δ* UoA785×UoA786 (n = 13); n indicates the number of independent

crosses (see also Table S2).

637
Figure 5. Over-expression of the *aho1*[+] in wild-type fission yeast cells and its consequences

for genotoxic stress resistance and meiotic recombination. (**a**, **b**) Effect of GFP and Aho1-

GFP over-expression on the genotoxin sensitivity of wild type (WT, FO652); number of cells

plated in each spot is indicated at the bottom of the figure. (**c**) Percent spore viability, (**d**)

frequency of gene conversion (GC) at *ade6-3083*, (**e**) frequency of crossovers (COs) between

ura4[+]*-aim2* and *his3*[+]*-aim*, and (**f**) percentage of COs between *ura4*[+]*-aim2* and *his3*[+]*-aim*

associated with a GC event at *ade6-3083* in crosses of wild type (WT) ALP733×UoA841 over-

expressing GFP (n = 6), or Aho1-GFP (n = 6); n indicates the number of independent crosses

(see also Table S3).

648
Figure 6. Genetic interaction of *hop1Δ* with deletions of factors required for meiotic

recombination intermediate processing. (**a**) Percent spore viability, (**b**) frequency of gene

conversion (GC) at *ade6-3083*, (**c**) frequency of crossovers (COs) between $ura4^+$-*aim2* and

$his3^+$-*aim*, and (**d**) percentage of COs between $ura4^+$-*aim2* and $his3^+$-*aim* associated with a

GC event at *ade6-3083* in crosses of wild type (WT) ALP733×ALP731 (n = 12), *hop1Δ*

UoA200×UoA199 (n = 16), *mus81Δ* ALP812×ALP813 (n = 10; in a) and ALP802×ALP822 (n =

10; in c-d), *hop1Δ mus81Δ* UoA204×UoA203 (n = 6), *fml1Δ* ALP1133×FO2608 (n = 15), and

fml1Δ hop1Δ UoA182×UoA181 (n = 14); n indicates the number of independent crosses (see

also Table S2).

659
Figure 7. Genetic interaction of *hop1*Δ with deletions of Rad51/Dmc1-mediator genes. (a,
d) frequency of gene conversion (GC) at *ade6-3083*, (**b**, **e**) frequency of crossovers (COs)
between *ura4+-aim2* and *his3+-aim*, and (**c**,**f**) percentage of COs between *ura4+-aim2* and
his3+-aim associated with a GC event at *ade6-3083* in crosses of wild type (WT)
ALP733×ALP731 (n = 12), *hop1*Δ UoA200×UoA199 (n = 16), *rad55*Δ ALP1649×ALP1648 (n =
12), *hop1*Δ *rad55*Δ UoA846×UoA847 (n = 10), *rlp1*Δ ALP1623×ALP1620 (n = 18), *hop1*Δ *rlp1*Δ
UoA848×UoA849 (n = 10), *sfr1*Δ ALP800×ALP782 (n = 22), *hop1*Δ *sfr1*Δ UoA844×UoA845 (n =
12), *dmc1*Δ ALP1545×ALP1544 (n = 12), and *dmc1*Δ *hop1*Δ UoA851×UoA852 (n = 10); n
indicates the number of independent crosses (see also Table S2).

References

Akamatsu Y, Dziadkowiec D, Ikeguchi M, et al (2003) Two different Swi5-containing protein
complexes are involved in mating-type switching and recombination repair in fission
yeast. Proc Natl Acad Sci U S A 100:15770–5. doi: 10.1073/pnas.2632890100

Andreoletti G, Seaby EG, Dewing JM, et al (2017) *AMMECR1*: a single point mutation causes
developmental delay, midface hypoplasia and elliptocytosis. J Med Genet 54:269–277.
doi: 10.1136/jmedgenet-2016-104100

Aravind L, Koonin E V (1998) The HORMA domain: a common structural denominator in
mitotic checkpoints, chromosome synapsis and DNA repair. Trends Biochem Sci 23:284–
6. doi: 10.1016/S0968-0004(98)01257-2

Armstrong SJ, Caryl AP, Jones GH, Franklin FCH (2002) Asy1, a protein required for meiotic
chromosome synapsis, localizes to axis-associated chromatin in *Arabidopsis* and
Brassica. J Cell Sci 115:3645–55. doi: 10.1242/jcs.00048

Bähler J, Wu JQ, Longtine MS, et al (1998) Heterologous modules for efficient and versatile
PCR-based gene targeting in *Schizosaccharomyces pombe*. Yeast 14:943–51. doi:
10.1002/(SICI)1097-0061(199807)14:10<943::AID-YEA292>3.0.CO;2-Y

Bähler J, Wyler T, Loidl J, Kohli J (1993) Unusual nuclear structures in meiotic prophase of
fission yeast: a cytological analysis. J Cell Biol 121:241–56. doi: 10.1083/jcb.121.2.241

Bishop DK, Park D, Xu L, Kleckner N (1992) *DMC1*: a meiosis-specific yeast homolog of E. coli
recA required for recombination, synaptonemal complex formation, and cell cycle
progression. Cell 69:439–56. doi: 10.1016/0092-8674(92)90446-J

Bleuyard J-Y, Gallego ME, Savigny F, White CI (2005) Differing requirements for the
Arabidopsis Rad51 paralogs in meiosis and DNA repair. Plant J 41:533–45. doi:
10.1111/j.1365-313X.2004.02318.x

Brown MS, Bishop DK (2014) DNA strand exchange and RecA homologs. Cold Spring Harb
Perspect Biol 7:a016659. doi: 10.1101/cshperspect.a016659

Brown SD, Lorenz A (2016) Single-step marker switching in *Schizosaccharomyces pombe*
using a lithium acetate transformation protocol. Bio-protocol 6:e2075. doi:
10.21769/BioProtoc.2075

Cahoon CK, Hawley RS (2016) Regulating the construction and demolition of the
synaptonemal complex. Nat Struct Mol Biol 23:369–377. doi: 10.1038/nsmb.3208

Callender TL, Laureau R, Wan L, et al (2016) Mek1 down regulates Rad51 activity during
yeast meiosis by phosphorylation of Hed1. PLoS Genet 12:e1006226. doi:
10.1371/journal.pgen.1006226

Carballo JA, Johnson AL, Sedgwick SG, Cha RS (2008) Phosphorylation of the axial element
protein Hop1 by Mec1/Tel1 ensures meiotic interhomolog recombination. Cell
132:758–70. doi: 10.1016/j.cell.2008.01.035

Caryl AP, Armstrong SJ, Jones GH, Franklin FC (2000) A homologue of the yeast *HOP1* gene is
inactivated in the *Arabidopsis* meiotic mutant *asy1*. Chromosoma 109:62–71. doi:
10.1007/s004120050413

Couteau F, Zetka M (2005) HTP-1 coordinates synaptonemal complex assembly with
homolog alignment during meiosis in *C. elegans*. Genes Dev 19:2744–56. doi:

10.1101/gad.1348205

Craven RA, Griffiths DJ, Sheldrick KS, et al (1998) Vectors for the expression of tagged
proteins in *Schizosaccharomyces pombe*. Gene 221:59–68. doi: 10.1016/S0378-
1119(98)00434-X

Daniel K, Lange J, Hached K, et al (2011) Meiotic homologue alignment and its quality
surveillance are controlled by mouse HORMAD1. Nat Cell Biol 13:599–610. doi:
10.1038/ncb2213

de los Santos T, Hollingsworth NM (1999) Red1p, a *MEK1*-dependent phosphoprotein that
physically interacts with Hop1p during meiosis in yeast. J Biol Chem 274:1783–90. doi:
10.1074/jbc.274.3.1783

Ding DQ, Haraguchi T, Hiraoka Y (2016) A cohesin-based structural platform supporting
homologous chromosome pairing in meiosis. Curr Genet 62:499–502. doi:
10.1007/s00294-016-0570-x

Doe CL, Dixon J, Osman F, Whitby MC (2000) Partial suppression of the fission yeast *rqh1*(-)
phenotype by expression of a bacterial Holliday junction resolvase. EMBO J 19:2751–62.
doi: 10.1093/emboj/19.11.2751

Ellermeier C, Schmidt H, Smith GR (2004) Swi5 acts in meiotic DNA joint molecule formation
in *Schizosaccharomyces pombe*. Genetics 168:1891–8. doi:
10.1534/genetics.104.034280

Estreicher A, Lorenz A, Loidl J (2012) Mug20, a novel protein associated with linear elements
in fission yeast meiosis. Curr Genet 58:119–27. doi: 10.1007/s00294-012-0369-3

Faul F, Erdfelder E, Buchner A, Lang A-G (2009) Statistical power analyses using G*Power 3.1:
tests for correlation and regression analyses. Behav Res Methods 41:1149–60. doi:
10.3758/BRM.41.4.1149

Faul F, Erdfelder E, Lang A-G, Buchner A (2007) G*Power: A flexible statistical power analysis
program for the social, behavioral, and biomedical sciences. Behav Res Methods
39:175–191. doi: 10.3758/BF03193146

Flor-Parra I, Zhurinsky J, Bernal M, et al (2014) A Lallzyme MMX-based rapid method for
fission yeast protoplast preparation. Yeast 31:61–6. doi: 10.1002/yea.2994

Fukuda T, Daniel K, Wojtasz L, et al (2010) A novel mammalian HORMA domain-containing
protein, HORMAD1, preferentially associates with unsynapsed meiotic chromosomes.
Exp Cell Res 316:158–71. doi: 10.1016/j.yexcr.2009.08.007

Goldstein AL, McCusker JH (1999) Three new dominant drug resistance cassettes for gene
disruption in *Saccharomyces cerevisiae*. Yeast 15:1541–53. doi: 10.1002/(SICI)1097-
0061(199910)15:14<1541::AID-YEA476>3.0.CO;2-K

Goodyer W, Kaitna S, Couteau F, et al (2008) HTP-3 links DSB formation with homolog
pairing and crossing over during *C. elegans* meiosis. Dev Cell 14:263–74. doi:
10.1016/j.devcel.2007.11.016

Gray S, Cohen PE (2016) Control of meiotic crossovers: From double-strand break formation
to designation. Annu Rev Genet 50:175–210. doi: 10.1146/annurev-genet-120215-
035111

Grimm C, Kohli J, Murray J, Maundrell K (1988) Genetic engineering of *Schizosaccharomyces*

pombe: a system for gene disruption and replacement using the *ura4* gene as a
selectable marker. Mol Gen Genet 215:81–6. doi: doi.org/10.1007/BF00331307

Grishchuk AL, Kohli J (2003) Five RecA-like proteins of *Schizosaccharomyces pombe* are
involved in meiotic recombination. Genetics 165:1031–43.

Hartsuiker E, Neale MJ, Carr AM (2009) Distinct requirements for the Rad32(Mre11)
nuclease and Ctp1(CtIP) in the removal of covalently bound topoisomerase I and II from
DNA. Mol Cell 33:117–23. doi: 10.1016/j.molcel.2008.11.021

Haruta N, Kurokawa Y, Murayama Y, et al (2006) The Swi5-Sfr1 complex stimulates
Rhp51/Rad51- and Dmc1-mediated DNA strand exchange in vitro. Nat Struct Mol Biol
13:823–30. doi: 10.1038/nsmb1136

Hayase A, Takagi M, Miyazaki T, et al (2004) A protein complex containing Mei5 and Sae3
promotes the assembly of the meiosis-specific RecA homolog Dmc1. Cell 119:927–40.
doi: 10.1016/j.cell.2004.10.031

Hollingsworth NM, Byers B (1989) *HOP1*: a yeast meiotic pairing gene. Genetics 121:445–62.

Hollingsworth NM, Goetsch L, Byers B (1990) The *HOP1* gene encodes a meiosis-specific
component of yeast chromosomes. Cell 61:73–84. doi: 10.1016/0092-8674(90)90216-2

Hong S, Sung Y, Yu M, et al (2013) The logic and mechanism of homologous recombination
partner choice. Mol Cell 51:440–453. doi: 10.1016/j.molcel.2013.08.008

Hunter N (2015) Meiotic recombination: the essence of heredity. Cold Spring Harb Perspect
Biol 7:a016618. doi: 10.1101/cshperspect.a016618

Hyppa RW, Smith GR (2010) Crossover invariance determined by partner choice for meiotic
DNA break repair. Cell 142:243–55. doi: 10.1016/j.cell.2010.05.041

Iwasaki O, Noma K (2016) Condensin-mediated chromosome organization in fission yeast.
Curr Genet 62:739–743. doi: 10.1007/s00294-016-0601-7

Keeney JB, Boeke JD (1994) Efficient targeted integration at *leu1-32* and *ura4-294* in
Schizosaccharomyces pombe. Genetics 136:849–56.

Khan K, Madhavan TPV, Kshirsagar R, et al (2013) N-terminal disordered domain of
Saccharomyces cerevisiae Hop1 protein is dispensable for DNA binding, bridging, and
synapsis of double-stranded DNA molecules but is necessary for spore formation.
Biochemistry 52:5265–79. doi: 10.1021/bi4005528

Kim Y, Rosenberg SC, Kugel CL, et al (2014) The chromosome axis controls meiotic events
through a hierarchical assembly of HORMA domain proteins. Dev Cell 31:487–502. doi:
10.1016/j.devcel.2014.09.013

Kogo H, Tsutsumi M, Ohye T, et al (2012) HORMAD1-dependent checkpoint/surveillance
mechanism eliminates asynaptic oocytes. Genes to Cells 17:439–54. doi:
10.1111/j.1365-2443.2012.01600.x

Lam I, Keeney S (2015) Mechanism and regulation of meiotic recombination initiation. Cold
Spring Harb Perspect Biol 7:a016634. doi: 10.1101/cshperspect.a016634

Latypov V, Rothenberg M, Lorenz A, et al (2010) Roles of Hop1 and Mek1 in meiotic
chromosome pairing and recombination partner choice in *Schizosaccharomyces pombe*.
Mol Cell Biol 30:1570–81. doi: 10.1128/MCB.00919-09

Loidl J (2016) Conservation and variability of meiosis across the eukaryotes. Annu Rev Genet
50:293–316. doi: 10.1146/annurev-genet-120215-035100

Loidl J (2006) *S. pombe* linear elements: the modest cousins of synaptonemal complexes.
Chromosoma 115:260–71. doi: 10.1007/s00412-006-0047-7

Loidl J, Klein F, Scherthan H (1994) Homologous pairing is reduced but not abolished in
asynaptic mutants of yeast. J Cell Biol 125:1191–200. doi: 10.1083/jcb.125.6.1191

Loidl J, Lorenz A (2009) Analysis of *Schizosaccharomyces pombe* meiosis by nuclear
spreading. Methods Mol Biol 558:15–36. doi: 10.1007/978-1-60761-103-5_2

Lorenz A (2017) Modulation of meiotic homologous recombination by DNA helicases. Yeast
34:195–203. doi: 10.1002/yea.3227

Lorenz A (2015) New cassettes for single-step drug resistance and prototrophic marker
switching in fission yeast. Yeast 32:703–710. doi: 10.1002/yea.3097

Lorenz A (2018) Supporting Online Material - Genetic interactions between the chromosome
axis-associated protein Hop1 and homologous recombination determinants in
Schizosaccharomyces pombe. doi: 10.6084/m9.figshare.5373163.v2

Lorenz A, Estreicher A, Kohli J, Loidl J (2006) Meiotic recombination proteins localize to linear
elements in *Schizosaccharomyces pombe*. Chromosoma 115:330–40. doi:
10.1007/s00412-006-0053-9

Lorenz A, Mehats A, Osman F, Whitby MC (2014) Rad51/Dmc1 paralogs and mediators
oppose DNA helicases to limit hybrid DNA formation and promote crossovers during
meiotic recombination. Nucleic Acids Res 42:13723–13735. doi: 10.1093/nar/gku1219

Lorenz A, Osman F, Folkyte V, et al (2009) Fbh1 limits Rad51-dependent recombination at
blocked replication forks. Mol Cell Biol 29:4742–56. doi: 10.1128/MCB.00471-09

Lorenz A, Osman F, Sun W, et al (2012) The fission yeast FANCM ortholog directs non-
crossover recombination during meiosis. Science 336:1585–8. doi:
10.1126/science.1220111

Lorenz A, Wells JL, Pryce DW, et al (2004) *S. pombe* meiotic linear elements contain proteins
related to synaptonemal complex components. J Cell Sci 117:3343–51. doi:
10.1242/jcs.01203

Lorenz A, West SC, Whitby MC (2010) The human Holliday junction resolvase GEN1 rescues
the meiotic phenotype of a *Schizosaccharomyces pombe mus81* mutant. Nucleic Acids
Res 38:1866–73. doi: 10.1093/nar/gkp1179

Martín V, Chahwan C, Gao H, et al (2006) Sws1 is a conserved regulator of homologous
recombination in eukaryotic cells. EMBO J 25:2564–74. doi: 10.1038/sj.emboj.7601141

Martinez-Perez E, Villeneuve AM (2005) HTP-1-dependent constraints coordinate homolog
pairing and synapsis and promote chiasma formation during *C. elegans* meiosis. Genes
Dev 19:2727–43. doi: 10.1101/gad.1338505

Mata J, Lyne R, Burns G, Bähler J (2002) The transcriptional program of meiosis and
sporulation in fission yeast. Nat Genet 32:143–7. doi: 10.1038/ng951

Matsuyama A, Arai R, Yashiroda Y, et al (2006) ORFeome cloning and global analysis of
protein localization in the fission yeast *Schizosaccharomyces pombe*. Nat Biotechnol
24:841–7. doi: 10.1038/nbt1222

Mimitou EP, Symington LS (2011) DNA end resection-Unraveling the tail. DNA Repair (Amst)
10:344–348. doi: 10.1016/j.dnarep.2010.12.004

Molnar M (2003) Linear element formation and their role in meiotic sister chromatid
cohesion and chromosome pairing. J Cell Sci 116:1719–1731. doi: 10.1242/jcs.00387

Moreno MB, Durán A, Ribas JC (2000) A family of multifunctional thiamine-repressible
expression vectors for fission yeast. Yeast 16:861–72. doi: 10.1002/1097-
0061(20000630)16:9<861::AID-YEA577>3.0.CO;2-9

Neale MJ, Pan J, Keeney S (2005) Endonucleolytic processing of covalent protein-linked DNA
double-strand breaks. Nature 436:1053–7. doi: 10.1038/nature03872

Niu H, Wan L, Baumgartner B, et al (2005) Partner choice during meiosis is regulated by
Hop1-promoted dimerization of Mek1. Mol Biol Cell 16:5804–18. doi:
10.1091/mbc.E05-05-0465

Niu H, Wan L, Busygina V, et al (2009) Regulation of meiotic recombination via Mek1-
mediated Rad54 phosphorylation. Mol Cell 36:393–404. doi:
10.1016/j.molcel.2009.09.029

Olson LW, Eden U, Egel-Mitani M, Egel R (1978) Asynaptic meiosis in fission yeast? Hereditas
89:189–199. doi: 10.1111/j.1601-5223.1978.tb01275.x

Osman F, Dixon J, Doe CL, Whitby MC (2003) Generating crossovers by resolution of nicked
Holliday junctions: a role for Mus81-Eme1 in meiosis. Mol Cell 12:761–74. doi:
10.1016/S1097-2765(03)00343-5

Osman K, Yang J, Roitinger E, et al (2018) Affinity proteomics reveals extensive
phosphorylation of the Brassica chromosome axis protein ASY1 and a network of
associated proteins at prophase I of meiosis. Plant J 93:17–33. doi: 10.1111/tpj.13752

Pancaldi V, Saraç OS, Rallis C, et al (2012) Predicting the fission yeast protein interaction
network. G3 (Bethesda) 2:453–67. doi: 10.1534/g3.111.001560

Phadnis N, Cipak L, Polakova S, et al (2015) Casein kinase 1 and phosphorylation of cohesin
subunit Rec11 (SA3) promote meiotic recombination through linear element formation.
PLoS Genet 11:e1005225. doi: 10.1371/journal.pgen.1005225

Rinaldi VD, Bolcun-Filas E, Kogo H, et al (2017) The DNA damage checkpoint eliminates
mouse oocytes with chromosome synapsis failure. Mol Cell 67:1026–1036. doi:
10.1016/j.molcel.2017.07.027

Rosenberg SC, Corbett KD (2015) The multifaceted roles of the HORMA domain in cellular
signaling. J Cell Biol 211:745–55. doi: 10.1083/jcb.201509076

Rothenberg M, Kohli J, Ludin K (2009) Ctp1 and the MRN-complex are required for
endonucleolytic Rec12 removal with release of a single class of oligonucleotides in
fission yeast. PLoS Genet 5:e1000722. doi: 10.1371/journal.pgen.1000722

Sabatinos SA, Forsburg SL (2010) Molecular genetics of *Schizosaccharomyces pombe*.
Methods Enzymol 470:759–95. doi: 10.1016/S0076-6879(10)70032-X

Sakuno T, Watanabe Y (2015) Phosphorylation of cohesin Rec11/SA3 by Casein Kinase 1
promotes homologous recombination by assembling the meiotic chromosome axis. Dev
Cell 32:220–230. doi: 10.1016/j.devcel.2014.11.033

Sambrook JF, Russell DW (2000) Molecular Cloning: A Laboratory Manual, 3rd edn. Cold

Spring Harbor Laboratory Press, Cold Spring Harbor

Sanchez-Moran E, Santos J-L, Jones GH, Franklin FCH (2007) ASY1 mediates AtDMC1-
dependent interhomolog recombination during meiosis in *Arabidopsis*. Genes Dev
21:2220–33. doi: 10.1101/gad.439007

Sasanuma H, Tawaramoto MS, Lao JP, et al (2013) A new protein complex promoting the
assembly of Rad51 filaments. Nat Commun 4:1676. doi: 10.1038/ncomms2678

Sato M, Dhut S, Toda T (2005) New drug-resistant cassettes for gene disruption and epitope
tagging in *Schizosaccharomyces pombe*. Yeast 22:583–91. doi: 10.1002/yea.1233

Schwacha A, Kleckner N (1997) Interhomolog bias during meiotic recombination: meiotic
functions promote a highly differentiated interhomolog-only pathway. Cell 90:1123–35.
doi: 10.1016/S0092-8674(00)80378-5

Shinohara A, Gasior S, Ogawa T, et al (1997) *Saccharomyces cerevisiae recA* homologues
RAD51 and *DMC1* have both distinct and overlapping roles in meiotic recombination.
Genes to Cells 2:615–29. doi: 10.1046/j.1365-2443.1997.1480347.x

Sievers F, Wilm A, Dineen D, et al (2014) Fast, scalable generation of high-quality protein
multiple sequence alignments using Clustal Omega. Mol Syst Biol 7:539–539. doi:
10.1038/msb.2011.75

Silva N, Ferrandiz N, Barroso C, et al (2014) The fidelity of synaptonemal complex assembly is
regulated by a signaling mechanism that controls early meiotic progression. Dev Cell
31:503–511. doi: 10.1016/j.devcel.2014.10.001

Smith GR (2009) Genetic analysis of meiotic recombination in *Schizosaccharomyces pombe*.
Methods Mol Biol 557:65–76. doi: 10.1007/978-1-59745-527-5_6

Smith GR, Boddy MN, Shanahan P, Russell P (2003) Fission yeast Mus81.Eme1 Holliday
junction resolvase is required for meiotic crossing over but not for gene conversion.
Genetics 165:2289–93.

Smith A V, Roeder GS (1997) The yeast Red1 protein localizes to the cores of meiotic
chromosomes. J Cell Biol 136:957–67. doi: 10.1083/jcb.136.5.957

Spirek M, Estreicher A, Csaszar E, et al (2010) SUMOylation is required for normal
development of linear elements and wild-type meiotic recombination in
Schizosaccharomyces pombe. Chromosoma 119:59–72. doi: 10.1007/s00412-009-0241-
5

Stanzione M, Baumann M, Papanikos F, et al (2016) Meiotic DNA break formation requires
the unsynapsed chromosome axis-binding protein IHO1 (CCDC36) in mice. Nat Cell Biol
18:1208–1220. doi: 10.1038/ncb3417

Sung P (1997) Yeast Rad55 and Rad57 proteins form a heterodimer that functions with
replication protein A to promote DNA strand exchange by Rad51 recombinase. Genes
Dev 11:1111–21. doi: 10.1101/gad.11.9.1111

Tripathi P, Pal D, Muniyappa K (2007) *Saccharomyces cerevisiae* Hop1 protein zinc finger
motif binds to the Holliday junction and distorts the DNA structure: implications for
Holliday junction migration. Biochemistry 46:12530–42. doi: 10.1021/bi701078v

Tsubouchi H, Roeder GS (2006) Budding yeast Hed1 down-regulates the mitotic
recombination machinery when meiotic recombination is impaired. Genes Dev

922   20:1766–75. doi: 10.1101/gad.1422506

923 Tsutsui Y, Khasanov FK, Shinagawa H, et al (2001) Multiple interactions among the
924   components of the recombinational DNA repair system in *Schizosaccharomyces pombe*.
925   Genetics 159:91–105.

926 Vitelli F, Piccini M, Caroli F, et al (1999) Identification and characterization of a highly
927   conserved protein absent in the Alport syndrome (A), mental retardation (M), midface
928   hypoplasia (M), and elliptocytosis (E) contiguous gene deletion syndrome (AMME).
929   Genomics 55:335–40. doi: 10.1006/geno.1998.5666

930 West AMV, Komives EA, Corbett KD (2017) Conformational dynamics of the Hop1 HORMA
931   domain reveal a common mechanism with the spindle checkpoint protein Mad2.
932   Nucleic Acids Res 1–14. doi: 10.1093/nar/gkx1196

933 Woltering D, Baumgartner B, Bagchi S, et al (2000) Meiotic segregation, synapsis, and
934   recombination checkpoint functions require physical interaction between the
935   chromosomal proteins Red1p and Hop1p. Mol Cell Biol 20:6646–58. doi:
936   10.1128/MCB.20.18.6646-6658.2000

937 Young JA, Hyppa RW, Smith GR (2004) Conserved and nonconserved proteins for meiotic
938   DNA breakage and repair in yeasts. Genetics 167:593–605. doi:
939   10.1534/genetics.103.023762

940 Zetka MC, Kawasaki I, Strome S, Müller F (1999) Synapsis and chiasma formation in
941   *Caenorhabditis elegans* require HIM-3, a meiotic chromosome core component that
942   functions in chromosome segregation. Genes Dev 13:2258–2270. doi:
943   10.1101/gad.13.17.2258