A novel combination of chemotherapy and imunotherapy controls tumor growth in mice with a human immune system

Running title: anti-ICOS and cyclophosphamide for breast cancer immunotherapy

Aude Burlion [1], Rodrigo N. Ramos [2], Kélhia Sendeyo [1], Aurélien Corneau [3], Christine Ménétrier-Caux [4], Eliane Piaggio [2], Daniel Olive [5], Christophe Caux [4] and Gilles Marodon [1*]

1 Sorbonne Université, CNRS, INSERM U1135, Centre d'Immunologie et des Maladies Infectieuses, CIMI, Paris, France.

2 Paris Sciences et Lettres Research University, INSERM U932, SiRIC Translational Immunotherapy Team, Institut Curie, Paris, France.

3 Sorbonne Université, UMS 30, Plateforme de Cytométrie, CyPS, Paris, France.

4 Université Claude Bernard Lyon 1, INSERM 1052, CNRS 5286, Department of Translational Research and Innovation, Centre Léon Bérard, Centre de Recherche en Cancérologie de Lyon, CRCL, Lyon, France.

5 Aix Marseille Université, INSERM U1068, CNRS, Institut Paoli - Calmettes, Centre de recherche en Cancérologie de Marseille, CRCM, Marseille, France.

* To whom correspondence should be addressed: gilles.marodon@upmc.fr

The authors declare no potential conflict of interests.

Abbreviations: ICOS, Inducible Costimulatory; NSG, NOD.SCID.gc-null; Treg, regulatory T cells; CTX, cyclophosphamide; HuMice, humanized mice; CyTOF, cytometry time-of-flight; tSNE, t-distributed stochastic neighbor embedding; pDCs, plasmacytoid dendritic cells; DC, dendritic cells; ICD, immunogenic cell death

Abstract

Mice reconstituted with a human immune system and bearing human tumors represent a promising model for developing novel cancer immunotherapies. A critical parameter will be to determine whether the quality of the immune infiltrate in tumors bears any similarities in humanized mice and patients. Here, we used mass cytometry and multi-parametric flow cytometry to characterize human leukocytes

35 infiltrating a human breast cancer tumor model in immunocompromised NOD.SCID.γc-null mice reconstituted with human hematopoietic progenitors and compared it to breast cancer samples. Among many similarities, we observed that $ICOS^+$ $CD4^+$ regulatory T cells (Tregs) were enriched in the tumor relative to the periphery both in humanized mice and in breast cancer patients. The presence of Tregs in the tumor has been associated with poor survival in breast cancer patients. Thus, targeting Tregs with

40 an anti-ICOS mAb could improve anti-tumor immunity and allow better tumor control. Indeed, a single administration of a neutralizing mAb to human ICOS affected Tregs in humanized mice. Moreover, a combination of the anti-ICOS mAb and chemotherapy reduced tumor growth compared to single treatments, associated to a marked decrease in Treg in the tumor. Similar results were observed with a fully humanized anti-ICOS mAb currently in clinical trial. Exploiting the similarities between

45 humanized mice and cancer patients to design and validate novel therapeutic strategies, as illustrated here, should lead to faster and more efficient treatments.

One sentence summary: single cell analysis of breast tumor infiltrates in patients and functional studies

50 *in vivo* in humanized mice reveal that ICOS expressed on Tregs is a promising target to improve tumor control in combination with chemotherapy.

Introduction

Preventing immune suppression in tumors is the "next frontier" in immuno-oncology research. Despite recent success of anti-PD-1 mAbs for metastatic melanomas and other solid tumors, it seems reasonable to propose that combination therapy will be key to the success of cancer immunotherapy, as suggested in patients receiving a combination of anti-PD-1 and anti-CTLA-4 *(1)*. Likely, the most efficient combinations would emerge from the vast list of mechanisms that prevent an efficient immune response. Recently, a murine study combining a tumor-targeting mAb, a recombinant cytokine, an anti-PD-1 mAb and a T cell vaccine achieved a remarkable efficacy at clearing large established syngeneic tumors *(2)*, illustrating the power of combinations for T-cell and innate cell mediated tumor control. However, studies purely performed in syngeneic models are only useful to gain biological insights but are not relevant to validate mAbs or drugs selectively targeting human cells. Therefore, a mouse model in which the impact on tumor control of mAbs targeting human T cells could be tested would be most invaluable.

It has been shown that human breast tumor morphology and metastatic properties were conserved in xenograft immunodeficient mice *(3,4)*. This was also showed for ovarian cancers *(5)* and latter confirmed for melanomas *(6)*, suggesting that a large panel of human tumors engrafts efficiently in immunodeficient mice and reproduces features of clinical progression. Xenograft models might be useful to target human tumors but these immunodeficient models are not relevant to test drugs such as mAbs that target the human immune system. Mice carrying a human immune system (HuMice) and human tumors could therefore represent a most relevant model for cancer immunotherapy research, at the interface of mice and humans *(7)*. Most of the literature regarding the use of HuMice for immuno-oncology relies on direct transfer of total PBMC into immunodeficient mice of various genetic backgrounds. Therapeutic efficacy of a combination of an anti-PD-1 and an anti-4-1-BB mAbs has been demonstrated in a model in which the tumor and the PBMC were from the same patient *(8)*. However, PBMC-implanted mice invariably and rapidly suffer from Graft vs Host Disease (GVHD), which complicate the interpretation of the results and limits the duration of the experiments. To preclude this problem, NOD.SCID.gamma-c-null (NSG) immunodeficient mice can be reconstituted with human $CD34^+$ hematopoietic progenitors *(9)*, most often isolated from cord blood. Because we demonstrated that a pool of human T cells with a diverse repertoire was then generated *(10)*, it becomes feasible to assess combination therapy targeting human tumors and/or human T cells *in vivo*,

generating results in a closer setting to human physiology than syngeneic murine models. However, there is still a lack of information on the composition and the function of human leukocytes infiltrating human tumors in CD34-reconstituted HuMice. Indeed, very few studies have evaluated the nature of human leukocytes in these models *(11–15)*□. Activated T cells in the tumor were observed in a peculiar model of tumor implantation at birth in NSG HuMice that did not lead to tumor rejection *(11)*□. A more advanced model of HuMice (MISTRG) was used to decipher the contribution of the innate immune system to tumor control, but the presence of T cells in the tumor was not reported *(13)*□. Very few T cells were detected in a Patient-Derived Xenograft (PDX) HuMice model of head and neck squamous cell carcinomas *(14)*□. Likewise, very few T cells were detected in breast cancer-bearing NSG HuMice with myeloid cells being the subset the most represented in the tumor *(12)*□. Finally, a recent study described the presence of Tregs in the tumor of NSG HuMice but a precise quantification of T cells was not reported *(15)*□. Thus, the presence of human T cells in tumors of HuMice is still open to debate. This is a crucial question to be answered if one wants to use the model in a reliable and relevant way.

Without tumors, the expected effects of the mAbs Teplizumab (anti-CD3) and Ipilimumab (anti-CTLA-4) were successfully reproduced in CD34-HuMice *(16,17)*□. Moreover, Avastin (anti-VEGF) had the expected effect on tumor growth *(13)*□. Altogether, these results show that human-specific mAbs are effective in HuMice. A recent study found that a humanized IgG1 mAb targeting the TNFR family member GITR, currently in clinical trial, was effective at preventing growth of a melanoma cell line in NSG HuMice *(15)*□. Thus, novel mAbs that would be effective on tumor control in HuMice might represent promising candidates for cancer immunotherapy in humans.

Cytometry Time-of-flight (CyTOF) or mass cytometry is a recent technology allowing quantification of the expression of 30 to 40 proteins at the same time at the single cell level *(18)*□, a very useful feature when analyzing samples containing a limited number of cells. Here, we used a combination of single-cell mass cytometry and high dimension flow cytometry to compare the immune landscape of human breast cancer in CD34-reconstituted NSG and in patients, as a reference. From this analysis, the Inducible COStimulatory molecule (ICOS) emerged in both human and HuMice as an eligible marker to target Tregs, a crucial T cell subset dampening the anti-tumoral immune response *(19)*□. We provide the first proof-of-principle that a combination of chemotherapy and neutralizing anti-ICOS mAb affected Tregs and reduced tumor growth in HuMice, opening the possibility to translate this combination into the clinics.

Results

T cell infiltrate in the tumors of HuMice

The general design of our study is represented on Fig. 1A. Irradiated NSG newborns were injected in the liver with cord blood CD34$^+$ cells. After 16 to 20 weeks, mice were grafted s.c with MDA-MB-231 human breast cancer cells. After 30 to 40 days, mice were euthanized and the spleen and tumors were analyzed by mass cytometry. The CyTOF panel was designed to include the most common lineage markers for murine and human leukocytes, but also several activation and proliferation markers of human T lymphocytes (Table S1).

We first compared the frequencies of human and murine cells of the hematopoietic lineage (CD45$^+$) in the spleen and in the tumor in 4 independent experiments, totalizing 15 CD34-reconstituted NSG mice. Within human cells, a classical 2D analysis of the CyTOF data revealed the presence of CD20$^+$ B cells and CD3$^+$ T cells (CD4$^+$ and CD8$^+$) in the spleen and the tumor, although B cells tended to disappear from the tumor (Fig. S1A). To get a deeper insight on the nature of immune cells present in the tumor, we performed unsupervised clustering of the data using the tSNE algorithm. This algorithm reduces the multidimensional nature of mass cytometry data to a 2D representation, in which each dot is a cell, clustered according to the level of expression of chosen markers *(20)*□. We first focused our analysis on manually gated hCD45$^+$ and ran the tSNE algorithm based on lineage and activation markers. Clusters of CD3$^+$, CD4$^+$ and CD8$^+$ cells in the spleen and the tumor were readily detected with that method (Fig. 1B). Thus, these results demonstrate that the MDA-MB-231 cell line is infiltrated by human T cells in CD34-reconstituted HuMice.

To compare the distribution of human leukocytes in HuMice to human samples, we determined the distribution of human leukocytes in the blood and tumors of 9 breast cancer patients by regular flow cytometry (Fig. 1C). The mean frequencies of CD3$^+$ T cells were similar in HuMice and patients in the tumor, representing around 50-60% of total CD45$^+$ cells. In contrast, the frequencies of CD4+ T cells was higher in the tumor of HuMice than in patients. Conversely, the frequencies of CD8$^+$ T cells were lower in the tumor of HuMice than in patients. Collectively, NSG HuMice and breast cancer patients harbored similar frequencies of human T cells in the tumor but the ratio of CD4$^+$ to CD8$^+$ T cells was higher in HuMice tumors than in patients.

Innate immune cells in the tumors of HuMice

We next investigated the nature of non-CD19, non-CD3-expressing cells in the same t-SNE analysis than before. A distinct cluster of $CD33^+$ $HLA-DR^+$ cells was observed in the spleen and tumor of HuMice (Fig. 2A). Zooming on in $CD33^+$, two distinct populations of $CD14^+HLA-DR^{lo}$ and $CD123^+HLA-DR^{hi}$ cells further segregate, most likely representing monocytes and plasmacytoid dendritic cells (pDCs) precursors, since pDC precursors and mature pDCs differ by CD33 expression *(21)* (Fig. 2B). A subset of $CD33^+$ cells not expressing CD14 nor CD123, and heterogeneous for HLA-DR were enriched in the tumor, likely including classical DCs. In addition, two clusters of Granzyme-B-expressing cells ($Grz-B^+$) were observed in the spleen and the tumor (Fig. 2C): one expressed CD8 and CD3 (Fig. 1D) and thus represented CTL, while the others weakly expressed CD335 (NKp46) (Fig. 2C) but not CD3 (Fig. 1D) and thus represented NK cells. In this later cluster, NKp46 expression was even lower in the tumor than in the spleen. Thus, in addition to $CD4^+$ and $CD8^+$ T cells, which represented the vast majority of human cells, the immune landscape of tumors in CD34-HuMice also included monocytes, pDC and NK cells, albeit in small proportions.

The same unsupervised representation of the data was performed on gated $mCD45^+$ cells based on myeloid-specific lineage markers (Fig. S1B). As expected from NSG mice, which are deprived of all lymphoid lineages, most $mCD45^+$ cells expressed CD11b in the spleen and the tumor (Fig. S1C). Clusters of $Ly-6C^+$ monocytes, $Ly6-G^+$ neutrophils and $CD11c^+$ DC were readily observed in the spleen and the tumor (Fig. S1C). Overall, myeloid cells of NSG mice were similarly represented in the spleen and the tumor, with the exception of $Ly6G^+$ neutrophils which were less abundant in the tumor (Fig. S1D).

Activated/memory T cells in tumors of HuMice

Having established unambiguously that tumors in HuMice contained human T cells, we next investigated the activation status of those human T cells. We performed a t-SNE analysis on human $CD3^+$ T cells, allowing visualization of two main clusters of $CD4^+$ and $CD8^+$ T cells (Fig. 3A). When activation/memory and functional markers such as CD45RO, HLA-DR, PD-1, CD25 or Grz-B were considered, we noticed a general increase in high expressing cells in the tumor relative to the spleen for both $CD4^+$ and $CD8^+$ cells, a first indication that T cells were activated in the tumor environment. Up

regulation of CD45RO and HLA-DR expression was observed in both CD4$^+$ and CD8$^+$ subsets whereas higher PD-1 and CD25 expression was noted among CD4$^+$ T cells. As expected, up regulation of Grz-B expression was restricted to CD8$^+$ T cells. These main features were reproduced across 3 independent experiments (Fig. S2A). To determine the activation status of human T cells more precisely, we performed a boolean analysis calculating the frequencies of cells expressing 0, 1, any combination of 2 or all 3 above mentioned activation markers in CD4$^+$ and CD8$^+$ T cells. Results depicted in Fig. 3B show that the frequencies of CD4$^+$ or CD8$^+$ cells co expressing 2 or 3 activation markers were vastly increased in the tumor, while frequencies of T cells expressing none of the activation markers were drastically reduced. Despite this massive activation, the frequencies of proliferating Ki-67$^+$ cells co-expressing 3 activation markers were lower in tumors relative to the spleen (Fig. S2B). Importantly, a similar enrichment for activated/memory T cells was observed in the tumor relative to the blood in breast cancer patients, assessed by the co expression of 2 or 3 activation/memory markers (Fig. 3C). Thus, tumor infiltrating T cells were of an activated/memory phenotype in both HuMice and breast cancer patients. Of note, PD-1 expression was correlated with higher proliferation of CD4$^+$ and CD8$^+$ T cells in the spleen. In contrast, it was inversely correlated with proliferation of CD4$^+$ but surprisingly not of CD8$^+$ T cells in the tumor (Fig S2C-E), appearing as an exhaustion marker only for CD4$^+$ T cells of the tumor.

Increased expression of ICOS on regulatory T cells in tumors

Effector Treg are defined by high expression of the transcription factor FOXP3 and the IL-2R-alpha chain CD25, and by the lack of CD45RA expression *(22)*. They are important actors of immunosuppression in the tumors of human patients. It has also been shown that ICOS/ICOS-L interaction can regulate Treg functions in humans *(23)* and that ICOS expression is linked to Treg-mediated immunosuppression in breast cancer patients *(24,25)*. Furthermore, ICOS expression was associated with poor prognosis in breast cancer patients due to its promoting action on Treg *(25,26)*. A distinct cluster of CD4$^+$ cells expressing FOXP3, high levels of ICOS and negative for CD45RA, representing effector Treg was detected at higher frequencies in breast tumors compared to blood (Fig. 4A-B). It has been shown in humans that CD4$^+$FOXP3$^+$CD45RA^{neg} cells might include activated effector T cells with lower FOXP3 expression than effector Treg *(22)*. We observed that the frequencies of ICOS$^+$ cells were higher in FOXP3^{hi} cells than in FOXP3^{lo} cells in the tumor (Fig. 4C),

suggesting that CD4$^+$CD45RA^{neg}FOXP3^{hi}ICOS$^+$ cells represented *bona fide* effector Treg. The higher expression of ICOS on Treg of the tumor was also documented by an increase in the MFI of ICOS in

210 FOXP3$^+$ vs. FOXP3^{neg} cells (Fig. 4D). Importantly, a similar increase was observed in the tumors of HuMice (Fig. 4D), showing that over expression of ICOS by Treg in human tumors was recapitulated in HuMice. We thus surmise that ICOS could represent a suitable target to affect Treg, with a possible impact on tumor growth, a testable hypothesis in HuMice.

215 **Impact of anti-ICOS mAb on Treg and Teff**

To test the hypothesis that ICOS/ICOS-L neutralization might affect Treg *in vivo*, we injected once an anti-hICOS mAb, reported as neutralizing in vitro *(25)*□, in tumor-bearing HuMice and determined the frequencies of Treg in the spleen and the tumor 30 days after (Fig. 5A). Technical issues

220 prevented CyTOF analysis of these experiments. Thus, regular flow cytometry was performed to determine frequencies of FOXP3$^+$ICOS$^+$ and FOXP3$^+$CD25$^+$ cells following treatment. There was a statistically significant effect of the treatment on the proportions of FOXP3$^+$ICOS$^+$ cells (p=0.0027, 2-way ANOVA) and of FOXP3$^+$CD25$^+$ cells (p=0.0039, 2-way ANOVA). By multiple comparisons with corrected p-values, the treatment led to a statistically significant reduction in the frequencies of

225 FOXP3$^+$ cells expressing CD25 or ICOS in the spleen of HuMice and similar tendencies were observed in the tumor (Fig. 5A). Moreover, the anti-ICOS mAb treatment was associated with increased proliferation of total CD4$^+$ but not CD8$^+$ T cells in the spleen (Fig. 5B). The treatment also affected the absolute numbers of total T cells in the spleen, roughly by a factor of 6, but that did not reach statistical significance. In contrast, Treg counts were significantly reduced by the treatment with numbers

230 dropping almost 20-fold (Fig. 5C). Thus, the anti-ICOS mAb led to a significant reduction of FOXP3$^+$ cells in treated animals but also improved proliferation of CD4$^+$ T cells.

Combination of chemotherapy and anti-ICOS to control tumor growth in HuMice

To evaluate the impact of the mAb on tumor growth, we dispatched HuMice with similar T cell reconstitution prior the initiation of the experiments (Fig. S3) into a pre-clinical-like trial with random assignments into groups, blinded evaluation, mixed sex ratio and sufficient number of experiments and

animals to detect the observed effect with enough statistical power (the experimental scheme is
summarized on Fig. 6A). Despite the clear effects on Treg and Teff reported above, the anti-ICOS mAb
injected alone at day 7 post-tumor implantation had no effect on tumor growth (Fig. 6B). We thus
reasoned that combining the anti-ICOS mAb with a known inducer of immunogenic cell death (ICD)
might be more efficient, as shown in syngeneic murine models with anti-CTLA-4 or anti-PD-1 mAbs
(27)□. Cyclophosphamide (CTX) is widely used as a chemotherapy for treatment of breast cancer for
its cytostatic properties, has also been described as a potent inducer of ICD and may affect Treg as well
(28)□. Indeed, CTX at a dose of 3 mg per mouse completely abolished tumor growth in non-
humanized NSG mice, whereas a dose of 1.5 mg only moderately affected tumor growth in non-
humanized (Fig. S4A) and HuMice alike (Fig. 6B), and moderately affected the proportions of Treg in
treated animals (Fig. S4B). The combination of the anti-ICOS mAb and CTX injected at day 7 post
tumor implantation profoundly reduced tumor growth in HuMice compared to single treatments (Fig.
6B). Thus, a combination of CTX with neutralizing anti-ICOS mAb efficiently controlled tumor growth
in CD34-reconstituted HuMice. As expected from results reported above, the proportion of Treg was
lower and the CD8/Treg ratio was higher in the tumor of the combo group relative to CTX alone (Fig.
6C), showing that the combined effect of CTX and anti-ICOS on Treg was associated to better tumor
control. Furthermore, frequencies of human CD45$^+$CD3$^+$ T cells were similar in all groups, indicating
that better tumor control in the combo group was not associated to a quantitative increase of human T
cells in the tumor (Fig. S5).

To test the therapeutic efficacy of the combination further, we used a fully humanized IgG4
anti-ICOS mAb, currently entered into a clinical trial (NCT02723955) in a similar unbiased pre-
clinical-like trial in 4 different experiments with repeated injections once a week. At day 30, the results
were already significant and reproduced the observations made with a single injection of the murine
mAb (Fig. 6D). Thus, better tumor control during combination of chemotherapy and anti-ICOS mAb
did not depend on the murine or human nature of the Fc fragment of the mAb.

Discussion

        In the present study, we provide the first demonstration that CD34-reconstituted HuMice can be used as a platform for the discovery and the validation of novel combinations of chemotherapy and immunotherapy, potentially applicable to patients. This demonstration originates from several similarities between HuMice and patients that we reveal here using multi parametric flow and mass
270   cytometry. A first similarity that we uncover here is the composition of the tumor immune landscape. Within total human hematopoietic cells, it is remarkable to note that a similar proportion of CD3$^+$ T cells infiltrated the tumors in patients and in HuMice. Thus, HuMice models might be useful to decipher the molecules and mechanisms at play that attracts T cells within the tumor environment. We also observed similar infiltration of memory/activated CD4$^+$ and CD8$^+$ T cells that expressed a
275   combination of activation markers. However, this activation status did not translate into increased proliferation in the tumor of HuMice, suggesting that the tumor microenvironment of this triple negative (TN) PR^{neg}ER^{neg}HER-2^{neg} cell line might be immunosuppressive *in vivo*. The immunosuppressive status of the tumor microenvironment in breast cancer patients vary according to the nature of the tumor (ER+, HER2+, TN) that may condition T cell infiltrate and T cell activation
280   status *(29)*□. Among breast tumor entities, TN are the most infiltrated by T cells (30)□, CD8$^+$ and FOXP3$^+$ cells alike, suggesting that TN might be an immunosuppressive-prone microenvironment. Here, we did not compare the proliferative status of T cells from patients with HuMice since none were carrying TN tumors. Further studies should confirm whether the immunosuppressive environment of the TN cell line observed here in HuMice is also observed in patients.

285   The enrichment for ICOS-expressing Treg is a remarkably conserved feature in HuMice and patients and led us to investigate the therapeutic potential of targeting ICOS for cancer immunotherapy. ICOS, member of the CD28 superfamily is an essential co-stimulatory molecule for T cell activation and function. Although originally thought of as a co-stimulatory molecule for effective T helper cell response, the biology of ICOS now extends well beyond this restricted view *(31)*□. Relevant to the
290   present study, we and others have shown that the survival, the expansion, and the suppressive functions of human Treg were dependent on ICOS signaling *(23,25,32)*□. Thus, we evaluated whether anti-ICOS mAbs could affect human Treg *in vivo*, with possible impact on tumor growth. The reduction in the absolute numbers of total T cells and Treg after anti-ICOS mAb treatment that we report here is in line with an important role for ICOS/ICOS-L on T cells survival. However, an alternative possibility is that

295   Treg would be physically depleted by the mAb, even though the murine IgG1 isotype used herein is generally not recognized as a strong inducer of ADCC. Moreover, NSG mice are deficient for complement-dependent ADCC, owing to a mutation in the C5 gene (33)□. Thus, the anti-ICOS mAb might have engaged FcRs expressed by myeloid cells, leading to active killing of ICOS-expressing cells, including Treg. Actually, Treg depletion mediated by myeloid cells has been proposed as a

300   mechanism to explain slower tumor growth following administration of a murine IgG2a isotype to GITR in complement-sufficient mice (34)□. Nevertheless, the fact that we observed a similar tumor control with murine or human mAbs in the context of chemotherapy is not in favor of the intervention of murine Fc receptors in the observed outcome, although this remains to be formally proven. As put forward in the study of Bulliard et al (34)□, much remains to be done to decipher the mechanisms

305   responsible for the therapeutic effects of mAbs used for immunotherapy. Whatever the mechanism, Treg depletion is now considered a major correlate of therapeutic efficacy for anti-CTLA-4 or anti-OX40 mAbs in mice (35–37)□ and possibly of Ipilimumab in humans (38)□. Additionally, the engagement of ICOS/ICOS-L pathway is required for anti-CTLA-4 efficacy in some models (39)□. Several mAbs targeting Treg are in development or already approved, including but not limited to anti-

310   CCR4 (40)□, anti-OX-40 (41)□, anti-GITR (42)□ and anti-CD25 mAb (43)□. In addition, the chemokine receptor CCR8 was recently proposed as an attractive target to affect Treg in breast cancer and alleviate immunosuppression, although this was not directly demonstrated (44)□. Our results indicate that ICOS might be a useful addition to this growing list of Treg targets.

        In line with published results in murine models (45)□, we confirm here that blocking ICOS

315   alone was not sufficient to impact tumor growth. However, a combination of chemotherapy and anti-ICOS mAb significantly impacted tumor growth in CD34-HuMice. This result supports the notion that the reduction in Treg is associated to better tumor control if ICD is simultaneously induced by chemotherapy. We did not find an association between frequencies of TILs and better control, suggesting that qualitative rather than quantitative aspects of the response are crucial to improve anti-

320   tumor immunity. We observed that CD4+ T cell proliferation was improved upon anti-ICOS treatment, suggesting that CD4+ T cells might possess improved functional properties. However, we are unable to discriminate the origin of this improved proliferation since it might be due to Treg decrease or an agonist effect of the anti-ICOS mAb. Furthermore, our CyTOF analysis showed that the tumors of HuMice also contained rare populations of immune cells known to play important roles on the anti-

325   tumor immune response, such as monocytes or pDCs. The pDC subset that we describe here for the

first time in the tumor of HuMice might have played a role on the amplification of ICOS[+] Treg, as described in breast cancer patients *(25)*□. It will be important in the future to monitor the human myeloid compartment more closely in HuMice models optimized for human myeloid development, since those cells can positively *(46)*□ and/or negatively affect the local anti-tumor immune response

330 *(47)*□. We also revealed the presence of NK cells with low expression of NKp46 in the tumor, suggesting impaired function, in agreement with clinical observations *(48)*□. Finally, a role for myeloid cells of the NSG mice cannot be excluded.

To our knowledge, our study represents the first demonstration that ICOS represents a target to impact tumor growth in a context of chemotherapy in a non-syngeneic humanized pre-clinical model.

335 The presence of ICOS[+] Treg has been described in breast *(24)*□, ovarian *(49)*□, and gastric *(50)*□ carcinomas, melanomas *(51,52)*□ and more recently follicular lymphomas *(53)*□, suggesting that ICOS-based cancer immunotherapy might be applicable to a wide range of cancers. There are currently at least 3 open clinical trials that evaluate anti-ICOS mAb alone or in combination with checkpoint blockade inhibitors in advanced solid tumors or lymphomas (NCT02520791, NCT02904226, and

340 NCT02723955 at clinicaltrials.gov). From our results, we predict that combination of anti-ICOS mAb with chemotherapy, not tested on those trials, might be more efficient at controlling tumor growth in patients than single treatments.

Methods

345 Mice and humanization

NOD.SCID.gc-null mice (stock ≠005557) were originally purchased from the Jackson Laboratory and were bred in our own animal facility under a 14hrs-10hrs light cycle with ad libitum food and water. Mice were given Baytril in their water every other week. Newborns NSG mice were
350 reconstituted by intra hepatic injection of $5.\,10^4$ to 10^5 magnetically-purified $CD34^+$ cord blood cells according to the manufacturer's instructions (Miltenyi, Paris, France) or were purchased from ABCell Bio (Paris, France). Reconstitution was followed over time in the blood by multicolor flow cytometry using the following markers in various combinations of fluorescent dyes: mCD45, hCD45, hCD3, hCD20, hCD4, and hCD8. Validated males and females humanized mice of 16 to 20 weeks old were
355 grafted s.c in the right flank with $1.5.10^6$ MDA-MB-231 breast cancer cells. Mice were euthanized when the tumor reached 3000 mm3 in control groups or 4 to 5 weeks after tumor implantation. All protocols were approved by the French National Ethical Committee on Animal Experimentation (Ce5/2012/025). To assess the effects of the various treatments on tumor growth, a total of 4 experiments are presented. Not every experiment included all conditions. Males and females NSG were
360 randomly dispatched into the various experimental groups to avoid sex-linked effects. Experimental groups were dispatched in different cages to avoid cage-related effects. Tumor growth was monitored in a blinded fashion. The number of mice used in each condition is indicated in the figure legend.

Cell line

The triple-negative ($PR^{neg}ER^{neg}HER-2^{neg}$) MDA-MB-231 cell line was grown in DMEM media supplemented with 10% FCS, L-glutamine and antibiotics (Penicillin/Streptomycin) (all from Thermo) in tissue culture flasks. Cells were transduced with a lentiviral vector co-expressing GFP and Luciferase to follow efficient engraftment *in vivo* using luciferase and to allow exclusion of tumor cells
370 from the analysis based on GFP expression. Cells were confirmed of being mycoplasma-free by a standard Hoechst-dye coloration on indicator Vero cells before injection into mice. A genetic profiling was established and confirmed the identity of the cell line (Eurofins Forensic Department, Ebersberg, Germany).

375 Reagents preparation and injection

The 314.8 mAb (mouse IgG1 anti-human ICOS) has been described before *(25)*□. The antibody was expressed *in vitro* using serum free medium and purified by protein-A affinity chromatography. Endotoxin was <1EU/mg. The 422 mAb from GSK is a fully humanized IgG4 anti-

380 ICOS mAb. Cyclophosphamide (CTX, Sigma Aldrich) was prepared extemporaneously according to supplier technical data sheet, i.e to 20 mg/ml of injectable water. All reagents were injected intra peritoneally.

Phenotypic analysis of leukocytes in the spleens and tumors of humanized mice

Splenocytes and tumors were digested with 1.6 mg/ml of collagenase IV and 10 µg/ml of DNAse I (Sigma Aldrich) for 2 hrs at 37° with regular flushing of the tissue. Cells were passed through a 40µm-cell strainer and resuspended in PBS 3% SVF. To eliminate dead cells and debris, tumor cell suspensions were isolated on a Ficoll gradient. Rings were collected, washed and cell pellets were

390 resuspended in PBS 3%SVF before counting with Trypan blue exclusion. Subsequently, 3 to 6. 10^6 cells live cells were stained with corresponding antibodies (lanthanide labeled mAbs for CyTOF or fluorochrome-labeled mAbs for FACS analysis). The details of each panel (each one corresponding to one experiment) can be found in Table S1. For CyTOF, 1 to 3µl of each lanthanide-labelled mAbs was added in 1.5 ml Eppendorf tubes in a final volume of 50µl of Max Par Staining Buffer (Fluidigm, San

395 Francisco, USA), according to manufacturer protocol. Intracellular staining was performed with FOXP3 staining kit (eBioscience, Courtaboeuf, France). Cell events were acquired on the CyTOF-2 mass cytometer and CyTOF software version 6.0.626 (Fluidigm) at the Cytometry Pitié-Salpétrière core (CyPS). Dual count calibration, noise reduction, cell length threshold between 10 and 150 pushes, and a lower convolution threshold equal to 10 were applied during acquisition. Data files produced by

400 the CyTOF-2 were normalized with the MatLab Compiler software normalizer using the signal from the 4-Element EQ beads (Fluidigm) as recommended by the software developers. Except for the analysis shown in Fig. 1, GFP exclusion was performed to remove tumor cells from the analysis. To normalize the variability between mice for supervised (i.e 2D plots) and unsupervised (i.e tSNE)

analysis, samples from tumors and spleen individually acquired on the CyTOF were aggregated in individual file in each organ and in each experiment.

Clinical samples

The main clinical characteristics of the patients are summarized in Table 1. Luminal breast tumors were collected from 9 untreated cancer patients undergoing standard surgery at Institut Curie Hospital, in accordance with institutional ethical guidelines and approved by the ethical (CPP ref: 99-15) and medical (ANSM ref: 2015-A00824-45) committees. Flow cytometry data in those patients were collected prior the initiation of the HuMice study, hence no CyTOF data were collected from human patients.

Tumor samples from patients and cell isolation

Samples were cut into small fragments, digested with 0.1 mg/ml Liberase TL in the presence of 0.1 mg/ml DNase (Roche, Meylan, France) and incubated for 30 min at 37°C in 5% of CO2 incubator. Subsequently, cells were filtered on a 40-µm cell strainer (BD Biosciences, Le Pont-de-Claix, France), washed and submitted to staining with specific antibodies. Peripheral blood from breast cancer patients was collected in tubes containing EDTA, washed in PBS and submitted to staining with specific fluorescent-labelled antibodies.

Phenotypic analysis of leukocytes in whole blood and tumors in human breast cancer patients

Tumor cell suspension and whole blood were stained with LIVE/DEAD Fixable Aqua (Life Technologies, Courtaboeuf, France) for 10 min at RT. Then, cells were washed and stained with Aqua BV500 dead cells exclusion dye (Life Technologies); anti-CD3 (clone OKT3, BV650), anti-CD4 (clone OKT4, BV785), anti-PD-1 (EH122.2H7, BV711), anti-CD27 (O323, BV605) from Biolegend (London, UK); anti-CD45 (clone 2D1, APC Cy7), anti-CD8 (clone RPA-T8, BUV395), anti-CD19 (clone HIB 19, Alexa 700) anti-CD56 (NCAM16.2, BUV737), anti-HLA-DR (G46.6, PECy5), anti-PD-L1 (MIHI, PE-CF594) from BD Biosciences; anti-EpCAM (1B7, eFluor660), anti-CD14 (2G5, FITC), anti-CD45RA (HI100, PECy7), and anti-ICOS (ISA3, PERCPe710) from eBioscience for 20 min at 4°C.

After incubation, cells were washed and permeabilized for 16 hours according to manufacturer's instructions for staining with anti-FOXP3 (236A/E7, PE, eBioscience) and anti-Granzyme-B (GB11, BV421, BD Biosciences). Cells were then fixed for subsequent analysis on a Fortessa flow cytometer (BD Biosciences). Data were analyzed with FlowJo Version v10 (FlowJO LCC, Ashland, USA), or Cytobank (www.cytobank.org).

Mass cytometry data analysis

For CyTOF data analysis, 10 healthy cord blood donors were used in 4 independent experiments, each experiment including mice reconstituted with different donors. A total of 15 mice were analyzed at the steady state by mass cytometry for human and murine cell content in the spleen and the tumor. Due to the paucity of cells in tissues of some HuMice, and to increase resolution of the analysis, tissue samples from 2 to 3 of those mice were pooled before staining and CyTOF analysis. The frequency of hCD45$^+$ determined by flow cytometry was used to normalize the representation of each mouse within the pool. In case where sufficient number of cells were collected, tissue samples from individual mice were run into CyTOF. To normalize the analysis from these various conditions, concatenation of individual mice from the same tissue in the same experiment was performed using FlowJo v10. Therefore, this analysis method does not allow comparisons of individual mice but of individual experiments. To document reproducibility of our observations, we provide frequencies of cells in each of the 4 individual experiments, when applicable. Concatenated samples were exported in Cytobank for unsupervised viSNE analysis or were analyzed using FlowJo v10 for supervised analysis. The default settings were applied: 1000 iterations, perplexity of 30 and a theta factor of 0.5.

Statistical analysis

Statistical analyses were performed using Prism v6.0h for Mac (GraphPad) with statistical models indicated in the figure legends. Outliers detection method is reported in the figure legends when applicable. All tests were performed with an α of 0.05 (probability of detecting a difference in the means by chance below 5%). No a priori sample size estimation to fit statistical power with the observed effect on tumor growth was used. However, a reverse analysis of our data (G-Power; gpower.hhu.de) showed that given the number of mice included in the study, the observed difference in

the means at the end of the experiment and the standard deviations in both groups, the β power was >95%, hence validating the rejection of the null hypothesis by statistical modeling of the data. For statistical analysis of tumor growth, the null hypothesis stating that one curve fits all the data in the compared groups was rejected based if the p-value was inferior to 0.05, determined by nonlinear

470   regression modeling of the data using the exponential growth equation.

Author's contributions (according to (54): Conceptualization, AB, CC, and GM; Methodology, AB, RR, EP and GM; Formal Analysis, AB, RR, KS, AC and GM; Investigation, AB, RR, KS and AC; Resources, EP, DO and CMC; Writing-Original Draft, AB and GM; Writing-Review and editing, AB, RR, EP, CMC, CC and GM; Visualization, AC, GM; Supervision, GM; Funding acquisition, CMC, CC, DO and GM.

Acknowledgements: the authors would like to thank Ms S Just-Landi (Institut Paoli-Calmettes, Marseille, France) for providing the 314.8 mAb, Dr P Mayes (GSK) for providing the 422 humanized mAb, Dr S Brunel and Mr Pukar KC for technical help, Dr H. Yssel (CIMI-PARIS, Paris, France) for monitoring mycoplasma infection in cell culture, Dr V. Vieillard (CIMI-PARIS) for advice on NK cells analysis, Dr M. Miyara (CIMI-PARIS) and Dr C. Combadière (CIMI-PARIS) for access to Cytobank, Dr B Salomon (CIMI-PARIS) for critical reading of the manuscript, C. Enond, O. Brégerie and B. Kane (Centre d'Exploration Fonctionnelle, Pitié Salpêtrière, Paris, France) for animal husbandry, and all the mothers for cord blood donations and patients for blood samples. **Funding:** This study was supported by the Ligue Nationale contre le cancer (LNC), the Agence Nationale de la Recherche (ANR-11-EMMA-0045 VICIT) and Glaxo Smith Kline (GSK). Dr A. Burlion was supported by a doctoral fellowship from the French Ministère de l'Education Supérieure et de la Recherche and from LNC. Dr K. Sendeyo was supported by a postdoctoral fellowship from GSK. **Competing interests:** The authors declare no competing interests. GSK had neither implication in the design of the experiments nor in the interpretation of the results.

References

1. Wolchok JD, Kluger H, Callahan MK, Postow M a, Rizvi N a, Lesokhin AM, et al. Nivolumab plus ipilimumab in advanced melanoma. N Engl J Med. 2013;369:122–33.

2. Moynihan KD, Opel CF, Szeto GL, Tzeng A, Zhu EF, Engreitz JM, et al. Eradication of large established tumors in mice by combination immunotherapy that engages innate and adaptive immune responses. Nat Med. 2016;22:1402–10.

3. DeRose YS, Wang G, Lin Y-C, Bernard PS, Buys SS, Ebbert MTW, et al. Tumor grafts derived from women with breast cancer authentically reflect tumor pathology, growth, metastasis and disease outcomes. Nat Med. Nature Publishing Group, a division of Macmillan Publishers Limited. All Rights Reserved.; 2011;17:1514–20.

4. Valdez KE, Fan F, Smith W, Allred DC, Medina D, Behbod F. Human primary ductal carcinoma in situ (DCIS) subtype-specific pathology is preserved in a mouse intraductal (MIND) xenograft model. J Pathol. 2011;225:565–73.

5. Bankert RB, Balu-Iyer S V., Odunsi K, Shultz LD, Kelleher RJ, Barnas JL, et al. Humanized mouse model of ovarian cancer recapitulates patient solid tumor progression, ascites formation, and metastasis. Orsulic S, editor. PLoS One. Public Library of Science; 2011;6:e24420.

6. Quintana E, Piskounova E, Shackleton M, Weinberg D, Eskiocak U, Fullen DR, et al. Human melanoma metastasis in NSG mice correlates with clinical outcome in patients. Sci Transl Med. 2012;4:159ra149.

7. Morton JJ, Bird G, Refaeli Y, Jimeno A. Humanized Mouse Xenograft Models: Narrowing the Tumor-Microenvironment Gap. Cancer Res. 2016;76:6153–8.

8. Sanmamed MF, Rodriguez I, Schalper KA, Onate C, Azpilikueta A, Rodriguez-Ruiz ME, et al. Nivolumab and Urelumab Enhance Antitumor Activity of Human T Lymphocytes Engrafted in Rag2-/-IL2R null Immunodeficient Mice. Cancer Res. 2015;75:3466–78.

9. Shultz LD, Lyons BL, Burzenski LM, Gott B, Chen X, Chaleff S, et al. Human Lymphoid and Myeloid Cell Development in NOD/LtSz-scid IL2R null Mice Engrafted with Mobilized Human Hemopoietic Stem Cells. J Immunol. 2005;174:6477–89.

10. Marodon G, Desjardins D, Mercey L, Baillou C, Parent P, Manuel M, et al. High diversity of the immune repertoire in humanized NOD.SCID.gamma c-/- mice. Eur J Immunol. 2009;39:2136–45.

11. Wege AK, Ernst W, Eckl J, Frankenberger B, Vollmann-Zwerenz A, Männel DN, et al. Humanized tumor mice-A new model to study and manipulate the immune response in advanced cancer therapy. Int J Cancer. 2011;129:2194–206.

12. Escobar G, Moi D, Ranghetti A, Ozkal-Baydin P, Squadrito ML, Kajaste-Rudnitski A, et al. Genetic engineering of hematopoiesis for targeted IFN-α delivery inhibits breast cancer progression. Sci Transl Med. 2014;6:217ra3.

13. Rongvaux A, Willinger T, Martinek J, Strowig T, Gearty S V, Teichmann LL, et al. Development and function of human innate immune cells in a humanized mouse model. Nat Biotechnol. 2014;32:364–72.

14. Morton JJ, Bird G, Keysar SB, Astling DP, Lyons TR, Anderson RT, et al. XactMice: humanizing mouse bone marrow enables microenvironment reconstitution in a patient-derived xenograft model of head and neck cancer. Oncogene. 2016;35:290–300.

15. Mahne A, Mauze S, Joyce-Shaikh B, Xia J, Bowman E, Beebe A, et al. Dual roles for regulatory T cell depletion and co-stimulatory signaling in agonistic GITR targeting for tumor immunotherapy. Cancer Res. 2016;77:1108–18.

16. Waldron-Lynch F, Henegariu O, Deng S, Preston-Hurlburt P, Tooley J, Flavell R, et al. Teplizumab Induces Human Gut-Tropic Regulatory Cells in Humanized Mice and Patients. Sci Transl Med. 2012;4:118ra12.

17. Vudattu NK, Waldron-Lynch F, Truman LA, Deng S, Preston-Hurlburt P, Torres R, et al. Humanized Mice as a Model for Aberrant Responses in Human T Cell Immunotherapy. J Immunol. 2014;193:587–96.

18. Spitzer MH, Nolan GP. Mass Cytometry: Single Cells, Many Features. Cell. Elsevier Inc.; 2016;165:780–91.

19. Tanaka A, Sakaguchi S. Regulatory T cells in cancer immunotherapy. Cell Res. Nature Publishing Group; 2017;27:109–18.

20. Amir ED, Davis KL, Tadmor MD, Simonds EF, Levine JH, Bendall SC, et al. viSNE enables visualization of high dimensional single-cell data and reveals phenotypic heterogeneity of leukemia. Nat Biotechnol. Nature Publishing Group; 2013;31:545–52.

21. See P, Dutertre C-A, Chen J, Günther P, McGovern N, Irac SE, et al. Mapping the human DC lineage through the integration of high-dimensional techniques. Science. 2017;356:eaag3009.

22. Miyara M, Yoshioka Y, Kitoh A, Shima T, Wing K, Niwa A, et al. Functional Delineation and Differentiation Dynamics of Human CD4+ T Cells Expressing the FoxP3 Transcription Factor. Immunity. 2009;30:899–911.

23. Ito T, Hanabuchi S, Wang YH, Park WR, Arima K, Bover L, et al. Two functional subsets of FOXP3+ regulatory T cells in human thymus and periphery. Immunity. 2008/06/03. 2008;28:870–80.

24. Gobert M, Treilleux I, Bendriss-Vermare N, Bachelot T, Goddard-Leon S, Arfi V, et al. Regulatory T Cells Recruited through CCL22/CCR4 Are Selectively Activated in Lymphoid Infiltrates Surrounding Primary Breast Tumors and Lead to an Adverse Clinical Outcome. Cancer Res. 2009;69:2000–9.

25. Faget J, Bendriss-Vermare N, Gobert M, Durand I, Olive D, Biota C, et al. ICOS-ligand expression on plasmacytoid dendritic cells supports breast cancer progression by promoting the accumulation of immunosuppressive CD4+ T cells. Cancer Res. 2012/10/03. 2012;72:6130–41.

26. Faget J, Sisirak V, Blay J-Y, Caux C, Bendriss-Vermare N, Ménétrier-Caux C. ICOS is associated with poor prognosis in breast cancer as it promotes the amplification of immunosuppressive CD4(+) T cells by plasmacytoid dendritic cells. Oncoimmunology. 2013;2:e23185.

27. Pfirschke C, Engblom C, Rickelt S, Cortez-Retamozo V, Garris C, Pucci F, et al. Immunogenic Chemotherapy Sensitizes Tumors to Checkpoint Blockade Therapy. Immunity. Elsevier Inc.; 2016;44:343–54.

28. Sistigu A, Viaud S, Chaput N, Bracci L, Proietti E, Zitvogel L. Immunomodulatory effects of cyclophosphamide and implementations for vaccine design. Semin Immunopathol. 2011;1–15.

29. Stanton SE, Disis ML. Clinical significance of tumor-infiltrating lymphocytes in breast cancer. J Immunother cancer. BioMed Central; 2016;4:59.

30. Stanton SE, Adams S, Disis ML. Variation in the Incidence and Magnitude of Tumor-Infiltrating Lymphocytes in Breast Cancer Subtypes A Systematic Review. JAMA Oncol. 2016;2:1354–60.

31. Wikenheiser DJ, Stumhofer JS. ICOS Co-Stimulation: Friend or Foe? Front Immunol. Frontiers; 2016;7:304.

32. Tuettenberg A, Huter E, Hubo M, Horn J, Knop J, Grimbacher B, et al. The Role of ICOS in Directing T Cell Responses: ICOS-Dependent Induction of T Cell Anergy by Tolerogenic Dendritic Cells. J Immunol. 2009;182:3349–56.

33. Baxter AG, Cooke A. Complement lytic activity has no role in the pathogenesis of autoimmune diabetes in NOD mice. Diabetes. 1993;42:1574–8.

34. Bulliard Y, Jolicoeur R, Windman M, Rue SM, Ettenberg S, Knee D a, et al. Activating Fc γ receptors contribute to the antitumor activities of immunoregulatory receptor-targeting antibodies. J Exp Med. 2013;210:1685–93.

35. Simpson TR, Li F, Montalvo-Ortiz W, Sepulveda MA, Bergerhoff K, Arce F, et al. Fc-dependent depletion of tumor-infiltrating regulatory T cells co-defines the efficacy of anti-CTLA-4 therapy against melanoma. J Exp Med. 2013;210:1695–710.

590   36.   Selby MJ, Engelhardt JJ, Quigley M, Henning K a., Chen T, Srinivasan M, et al. Anti-CTLA-4 Antibodies of IgG2a Isotype Enhance Antitumor Activity through Reduction of Intratumoral Regulatory T Cells. Cancer Immunol Res. 2013;1:32–42.

37.   Bulliard Y, Jolicoeur R, Zhang J, Dranoff G, Wilson NS, Brogdon JL. OX40 engagement depletes intratumoral Tregs via activating FcγRs, leading to antitumor efficacy. Immunol Cell
595   Biol. 2014;92:475–80.

38.   Romano E, Kusio-Kobialka M, Foukas PG, Baumgaertner P, Meyer C, Ballabeni P, et al. Ipilimumab-dependent cell-mediated cytotoxicity of regulatory T cells ex vivo by nonclassical monocytes in melanoma patients. Proc Natl Acad Sci. 2015;112:201417320.

39.   Fan X, Quezada S a, Sepulveda M a, Sharma P, Allison JP. Engagement of the ICOS pathway
600   markedly enhances efficacy of CTLA-4 blockade in cancer immunotherapy. J Exp Med. 2014;211:715–25.

40.   Chang D-K, Peterson E, Sun J, Goudie C, Drapkin RI, Liu JF, et al. Anti-CCR4 monoclonal antibody enhances antitumor immunity by modulating tumor-infiltrating Tregs in an ovarian cancer xenograft humanized mouse model. Oncoimmunology. Taylor & Francis;
605   2016;5:e1090075.

41.   Marabelle A, Kohrt H, Sagiv-Barfi I, Ajami B, Axtell RC, Zhou G, et al. Depleting tumor-specific Tregs at a single site eradicates disseminated tumors. J Clin Invest. 2013;123:2447–63.

42.   Sukumar S, Wilson DC, Yu Y, Wong J, Naravula S, Ermakov G, et al. Characterization of MK-4166, a clinical agonistic antibody that targets human GITR and inhibits the generation and
610   suppressive effects of T regulatory cells. Cancer Res. 2017;77:canres.1439.2016.

43.   Arce Vargas F, Furness AJS, Solomon I, Joshi K, Mekkaoui L, Lesko MH, et al. Fc-Optimized Anti-CD25 Depletes Tumor-Infiltrating Regulatory T Cells and Synergizes with PD-1 Blockade to Eradicate Established Tumors. Immunity. 2017;46:577–86.

44.   Plitas G, Konopacki C, Wu K, Bos PD, Morrow M, Putintseva E V., et al. Regulatory T Cells
615   Exhibit Distinct Features in Human Breast Cancer. Immunity. Elsevier Inc.; 2016;45:1122–34.

45.   Metzger TC, Long H, Potluri S, Pertel T, Bailey-Bucktrout SL, Lin JC, et al. ICOS promotes the function of CD4+ effector T cells during anti-OX40-mediated tumor rejection. Cancer Res. American Association for Cancer Research; 2016;76:3684–9.

46.   Broz ML, Binnewies M, Boldajipour B, Nelson AE, Pollack JL, Erle DJ, et al. Dissecting the
620   Tumor Myeloid Compartment Reveals Rare Activating Antigen-Presenting Cells Critical for T Cell Immunity. Cancer Cell. Elsevier Inc.; 2014;26:638–52.

47. Gabrilovich DI, Ostrand-Rosenberg S, Bronte V. Coordinated regulation of myeloid cells by tumours. Nat Rev Immunol. Nature Publishing Group, a division of Macmillan Publishers Limited. All Rights Reserved.; 2012;12:253–68.

48. Mamessier E, Sylvain A, Thibult M, Houvenaeghel G, Jacquemier J, Castellano R, et al. Human breast cancer cells enhance self tolerance by promoting evasion from NK cell antitumor immunity. J Clin Invest. 2011;121:3609–22.

49. Conrad C, Gregorio J, Wang YH, Ito T, Meller S, Hanabuchi S, et al. Plasmacytoid dendritic cells promote immunosuppression in ovarian cancer via ICOS costimulation of Foxp3+ T-regulatory cells. Cancer Res. 2012;72:5240–9.

50. Nagase H, Takeoka T, Urakawa S, Morimoto-Okazawa A, Kawashima A, Iwahori K, et al. ICOS [+] Foxp3 [+] TILs in gastric cancer are prognostic markers and effector regulatory T cells associated with *H elicobacter pylori*. Int J Cancer. 2016;1–10.

51. Martin-Orozco N, Li Y, Wang Y, Liu S, Hwu P, Liu Y-J, et al. Melanoma cells express ICOS ligand to promote the activation and expansion of T-regulatory cells. Cancer Res. 2010/11/26. 2010;70:9581–90.

52. Strauss L, Bergmann C, Szczepanski MJ, Lang S, Kirkwood JM, Whiteside TL. Expression of ICOS on human melanoma-infiltrating CD4+CD25highFoxp3+ T regulatory cells: implications and impact on tumor-mediated immune suppression. J Immunol. 2008/02/23. 2008;180:2967–80.

53. Le K-S, Thibult M-L, Just-Landi S, Pastor S, Gondois-Rey F, Granjeaud S, et al. Follicular B Lymphomas Generate Regulatory T Cells via the ICOS/ICOSL Pathway and Are Susceptible to Treatment by Anti-ICOS/ICOSL Therapy. Cancer Res. 2016;76:4648–60.

54. Brand A, Allen L, Altman M, Hlava M, Scott J. Beyond authorship: Attribution, contribution, collaboration, and credit. Learn Publ. 2015;28:151–5.

Cod. Pat.	Age (years)	Meno status	Histo.	Grad.	Size (mm)	pTNM	ER (%)	PR (%)	HER2 status	Ki67 (%)	TIL (%)
808220	86	yes	Lobular	2	21	pT2N3M0	100	0	neg	20	15
808243	55	yes	Ductal	2	30	pT2N1M0	100	15	neg	20	5
808254	65	yes	Ductal	2	25	pT2N2M0	95	95	neg	15	NA
808364	52	yes	Ductal	2	18	pT1N2M0	90	80	neg	15	<10
808430	32	no	Ductal	2	26	pT2N2M0	100	95	neg	40	20
809090	40	no	Ductal	3	25	pT2N2M0	90	90	neg	40	30
809113	60	yes	Ductal	2	28	pT2N1M0	100	30	pos	17	10

Table 1. Clinical characteristics of the primary breast tumors samples. Cod. Pat.= Patient code/ Age= age of the patient at the time of surgery/Meno. status = Menopausal status/ Histo.= Histologic 655 type/ Grade= SBR Grade (histo-prognostic grade based on "Scarff-Bloom-Richardson")/ pTNM= tumor classification based on Tumor-Nodes-Metastasis score/ ER= % of estrogen receptor positivity/ PR= % of progesterone receptor positivity/ Ki67%= cellular marker of proliferation/ TILs=% of Tumor-infiltrated Lymphocytes.

Fig. 1. Distribution of human and murine leukocytes in HuMice and in breast cancer patients.
(**A**) Experimental scheme of the study, as detailed in the text. (**B-C**) Frequencies of human and murine hematopoietic cells in the spleen (**B**) and the tumor (**C**) of HuMice were determined based on hCD45$^+$ and mCD45$^+$ expression as depicted. Each dot is an experiment. Negative cells are split in two on each axes due to the arc sin transformation of CyTOF data and arbitrary settings for best 2D representation in FlowJo. (**D**) viSNE representation of gated human CD45$^+$ cells in the spleen and the tumor of HuMice from a representative experiment. The viSNE plot was generated according to NKp46, CD38, CD33, CD45RO, PD-1, CD4, CD8, CD20, CD25, Grz-B, and HLA-DR expression using proportional sampling with 54422 events in the spleen and 10274 events in the tumor. The level of expression of each of the indicated markers is represented by a color scale on the right. (**E**) Frequencies of the indicated subsets were determined in HuMice by supervised 2D-gating from CyTOF data and in patients from multi-parametric flow cytometry. Each dot represents an experiment or a patient. Horizontal line represents the mean value. Four independent experiments totalizing 15 mice were performed. The p-values in B, C and E are from non-parametric two-tailed Mann-Whitney t-test.

Fig. 2. Detection of human myeloid cells and NK cells in HuMice. (A) The viSNE plot was generated as in Fig. 1E and represents the expression of CD33 and HLA-DR in hCD45$^+$ cells of the spleen and tumor. **(B)** The viSNE plots were generated in gated hCD45$^+$CD33$^+$ cells according to CD14, CD45RO, CD123, CD4, CD11b and HLA-DR expression using proportional sampling with 535 cells in the spleen and 476 events in the tumors. Indicated on the plots are the frequencies of CD14-expressing cells in the CD33$^+$ cluster. **(C)** The visNE plots was generated as in Fig. 1E and represents the expression of Granzyme-B (Grz-B) and NKp46 in the spleen and the tumor. Gates highlight the localization of putative NK cells.

Fig. 3. Activation status of Tumor-infiltrating T cells of HuMice and breast cancer patients. (A) hCD45$^+$CD3$^+$ from HuMice were gated in Cytobank and a viSNE plot was generated according to CD4, CD8, CD45RO, HLA-DR, PD-1, CD25, Ki-67 and GrzB expression using equal sampling with 8574 events in the spleen and tumor. **(B)** Mean frequencies of cells expressing 0, 1, any combination of 2 or all 3 activation/memory markers are shown in the indicated tissue. Activation/memory markers used for boolean analysis were CD25, HLA-DR and PD-1 for CD4$^+$ T cells and HLA-DR, Grz-B and CD45RO for CD8$^+$ T cells. Results are from one experiment out of 3. **(C)** Same analysis for T cells from breast cancer patients. Results are mean frequencies in 9 breast cancer patients in 9 independent experiments in the blood and the tumor determined by flow cytometry. The activation markers used for boolean analysis were HLA-DR, ICOS and PD-1 for CD4$^+$ T cells and CD45RA, Grz-B and HLA-DR for CD8$^+$ T cells (negativity for CD45RA was considered as an activated/memory phenotype).

Fig. 4. ICOS expression by Treg in HuMice and breast cancer patients. (**A**) viSNE plot of hCD45$^+$ from a representative breast cancer patient showing FOXP3, ICOS and CD45RA expression in the blood and the tumor, determined by flow cytometry. (**B**) Frequencies of FOXP3$^+$CD45RA^{neg} cells in the indicated tissue from breast cancer patients among CD4$^+$CD3$^+$ cells. (**C**) Frequencies of ICOS$^+$ cells in FOXP3^{lo} or FOXP3^{hi} CD4$^+$CD45RA^{neg} in the tumor of breast cancer patients. (**D**) MFI of ICOS in CD3$^+$CD4$^+$Foxp3$^-$ or CD3$^+$CD4$^+$Foxp3$^+$ in the tumor of breast cancer patients or in the tumor of HuMice. Horizontal line represents the mean value. Each dot represents a patient or a mouse. The p values indicated on the graphs are from non-parametric two-tailed Mann-Whitney t-test.

Fig 5. Impact of anti-ICOS mAb on Treg and Teff in HuMice. (**A**) Frequencies of FOXP3[+]ICOS[+] or FOXP3[+]CD25[+] cells in CD4[+]CD3[+] T cells in the spleen and tumor of HuMice injected with isotype control (Iso) or anti-ICOS mAb (50µg/mouse). Results shown are cumulative of at least 3 independent experiments. Only high expressing cells were gated. The p-values reported on the graph are from a 2-way ANOVA with multiple comparisons test corrected by the Sidak method. (**B**) Shown are the frequencies of Ki-67[+] cells in the indicated subsets in the spleen of HuMice treated with the isotype control (Iso) or the anti-ICOS mAb (50µg/mouse) from 2 independent experiments. Each dot is a mouse. The p-values indicated on the graphs are from a 2-way ANOVA with multiples comparisons using the Sidak correction. **C**) Absolute numbers of total T cells (CD3[+]) or Treg (FOXP3[+]CD25[+]) were based on absolute counts of the spleen of HuMice treated with isotype control (Iso) or anti-ICOS mAb. Each dot represents a mouse. Fold change in mean numbers are indicated on the right. The p-values indicated on the graphs are from an unpaired two-tailed non-parametric Kolgomorov-Smirnov t-test. Data are cumulative of at least 3 independent experiments. Horizontal line represents the mean value.

Figure 6. Therapeutic efficacy of anti-ICOS mAbs in humanized mice (A) Experimental design of the study, as detailed in the text. (B) Tumor growth was determined in 4 independent experiments in the indicated number of mice treated by PBS and isotype control (Vehicles), the murine anti-ICOS 314.8 and PBS (anti-ICOS; 50 µg/mouse), cyclophosphamide and isotype control (CTX; 1.5 mg/mouse) or a combination of CTX and anti-ICOS (Combo). Error bars are SEM. The arrow indicates the day the anti-ICOS and the CTX treatment was performed. (C) Frequencies of Foxp3$^+$CD25$^+$ cells in CD4$^+$CD3$^+$ cells of the tumor in the indicated conditions. The CD8 to Treg ratio was obtained by dividing the frequencies of CD8$^+$ T cells by the frequencies of Foxp3$^+$CD25$^+$ cells in CD3+ cells of the tumor in the indicated conditions. Each dot is a mouse from two experiments. (D) Tumor growth was determined in 4 independent experiments in the indicated number of mice treated with isotype controls (iso), the humanized IgG4 anti-ICOS mAb (422) (0.5mg/kg once a week), cyclophosphamide (CTX; 1.5mg/mouse) or the combination (Combo). Error bars are SEM.